



A PRACTITIONER'S GUIDE FOR
ADVANCING HEALTH EQUITY
Community Strategies for Preventing Chronic Disease



TOBACCO-FREE LIVING STRATEGIES



National Center for Chronic Disease Prevention and Health Promotion
Division of Community Health



This document was developed by the Centers for Disease Control and Prevention, U.S. Department of Health and Human Services and Prevention Institute, with funding support under contract 200-2007-20014 Task 009. The design and published components of this document were completed by Ogilvy Public Relations with funding from the Centers for Disease Control and Prevention under contract 200-2010-F-33546.

WWW.CDC.GOV/HEALTHEQUITYGUIDE

**Centers for Disease Control and Prevention
Division of Community Health**

**4770 Buford Highway, NE, Mailstop F-81
Atlanta, Georgia 30341**

Phone: 770-488-1170

Fax: 770-488-5964

Email: cdcinfo@cdc.gov

Suggested Citation: Centers for Disease Control and Prevention – Division of Community Health. *A Practitioner’s Guide for Advancing Health Equity: Community Strategies for Preventing Chronic Disease*. Atlanta, GA: US Department of Health and Human Services; 2013.

Disclaimer: *A Practitioner’s Guide for Advancing Health Equity* is intended as a resource for public health practitioners working to advance health equity through chronic disease-focused community health efforts. However, this guide is not intended to reflect the Centers for Disease Control and Prevention (CDC) activities or promote strategies that may not adhere to restrictions regarding the use of federal investments.

Additionally, *A Practitioner’s Guide for Advancing Health Equity* is not intended to serve as step-by-step instructions, as there is no one-size-fits all approach to advancing health equity. Although this document discusses a variety of evidence- and practice-based strategies, it is not exhaustive. Strategies included may not be appropriate for every organization’s situation. Communities must decide what is appropriate for their local context. Therefore, strategies and examples in this guide should be considered in accordance with an organization’s and, where applicable, its funder’s established protocols and regulations.

In accordance with applicable laws, policies, and regulations, we note that, no federal funds are permitted to be used for impermissible lobbying in support of or opposition to proposed or pending legislative matters. Any organization using this guide should be aware of restrictions related to their organization’s funding sources when considering the strategies included in this document. For additional information on federal restrictions on lobbying for CDC funding recipients using CDC funds, see Anti-Lobbying Restrictions for CDC Grantees¹ and Additional Requirement #12.²

Information in this guide does not constitute legal advice. Use of any strategy outlined should only be considered within the context of guidance from legal counsel, as appropriate, to ensure compliance with applicable laws and policies.

Finally, links to non-federal government organizations found in this document are provided solely as a service to the reader. These links do not constitute an endorsement of these organizations or their programs by CDC or the Federal Government, and none should be inferred. CDC is not responsible for the content of any other Web sites contained in this document.



LETTER FROM THE DIVISION OF COMMUNITY HEALTH

PUBLIC HEALTH PRACTITIONER,

There is a growing body of literature exploring how environments in this nation shape our health. To address this issue, public health practitioners are implementing chronic disease policy, systems, and environmental improvements where people live, learn, work, and play. Practitioners are also considering how to ensure such improvements are designed to reverse the negative trends of chronic health conditions among vulnerable population groups. In response to the mounting needs of practitioners seeking reliable tools to advance health equity, the Centers for Disease Control and Prevention (CDC) developed *A Practitioner's Guide for Advancing Health Equity: Community Strategies for Preventing Chronic Disease (Health Equity Guide)*.

The purpose of the *Health Equity Guide* is to assist practitioners with addressing the well-documented disparities in chronic disease health outcomes. This resource offers lessons learned from practitioners on the front lines of local, state, and tribal organizations that are working to promote health and prevent chronic disease health disparities. It provides a collection of health equity considerations for several policy, systems, and environmental improvement strategies focused on tobacco-free living, healthy food and beverages, and active living. Additionally, the *Health Equity Guide* will assist practitioners with integrating the concept of health equity into local practices such as building organizational capacity, engaging the community, developing partnerships, identifying health inequities, and conducting evaluations. The *Health Equity Guide* is designed for the novice interested in the concept of health equity, as well as the skillful practitioner tackling health inequities.

We encourage you to visit WWW.CDC.GOV/HEALTHEQUITYGUIDE for additional tools and resources that promote health and the integration of health equity into everyday practice. We hope you find the information and examples provided to be useful and an impetus in your efforts to reduce health disparities and advance health equity.

Sincerely,

Leonard Jack, Jr, PhD, MSc
Director, Division of Community Health (DCH)
National Center for Chronic Disease
Prevention and Health Promotion
(NCCDPHP)

Pattie Tucker, DrPH, RN
Acting Associate Director for Health Equity, DCH
National Center for Chronic Disease
Prevention and Health Promotion
(NCCDPHP)



TABLE OF CONTENTS

ACKNOWLEDGEMENTS v

INTRODUCTION 1

SECTION 2: MAXIMIZING TOBACCO-FREE LIVING STRATEGIES TO ADVANCE HEALTH EQUITY 5

Comprehensive Smoke-Free Policies 6

 Native American Tribes Adopt Tobacco Protections for Tribal Members and Future Generations—Montana 8

 Partnerships and Educational Initiatives Lead to Smoke-Free Air Protections—Birmingham, AL 9

Smoke-Free Multi-Unit Housing Policies 10

 Creating Healthy Environments through Smoke-Free Multi-Unit Housing Policies—San Antonio, TX 12

 Housing Authority and Public Health Commission Partner on Smoke-Free Housing—Boston, MA 13

Tobacco Cessation Services 14

 Expanding Cessation Services in Marginalized Communities—St. Louis, MO 16

 Using Partnerships to Increase Access to Cessation Services—Santa Clara County, CA 17

Point-of-Sale Strategies to Address Access and Exposure to Tobacco Products 18

 Citywide Restrictions Tackle Flavored Tobacco Products—New York, NY 20

 Tobacco-Free Pharmacies Promote Health for All—San Francisco, CA 21

APPENDICES 22

REFERENCES 30



ACKNOWLEDGEMENTS

This resource was developed with the input and feedback of practitioners and researchers from across the United States. We would like to thank all who have contributed to the development and design of *A Practitioner's Guide for Advancing Health Equity*. The affiliations listed below are those of the contributors at the time of their participation.

Core Project Team

CDC expresses sincere appreciation to the Core Project Team for the development of this resource. The Core Project Team's time, expertise, and dedication contributed significantly to the vision and content of this publication.

Carolyn Brooks
Centers for Disease Control and Prevention

Rebecca Bunnell
Centers for Disease Control and Prevention

Rebecca Payne
Centers for Disease Control and Prevention

Manal Aboelata
Prevention Institute

Dalila Butler
Prevention Institute

Virginia Lee
Prevention Institute

Shayla Spilker
Prevention Institute

Special thanks to Natalie Stahl, Centers for Disease Control and Prevention; Lesley Guyot, SciMetrika; and Menaka Mohan, Prevention Institute for their project support and coordination.

Contributing Writers and Researchers

CDC thanks the contributing writers and researchers for their diligent and thoughtful analyses and reflections of the literature and input from community practitioners.

Centers for Disease Control and Prevention

Carolyn Brooks	Jennifer Kohr
Belsie Gonzalez	Terry Njoroge
Joi Hudson	Natalie Stahl

Consultants

Kim Hodgson,
Cultivating Healthy Places

Elva Yañez,
Colibri Strategies Inc.

Emory University Rollins School of Public Health

Jessica Pittman

Prevention Institute

Manal Aboelata	Menaka Mohan
Dalila Butler	Janet Pan
Jeremy Cantor	Nicole Schneider
Sana Chehimi	Linda Shak
Tony Dang	Ben Simons
Rachel Davis	Juliet Sims
Phebe Gibson	Shayla Spilker
Carolina Guzman	Sandra Viera
Virginia Lee	Ann Whidden
Leslie Mikkelsen	

SciMetrika

Lesley Guyot

Advisors and Reviewers

CDC gratefully acknowledges the time, dedication, and expertise of the many individuals and organizations involved in the development and review of *A Practitioner's Guide for Advancing Health Equity*. The insight of these individuals has been invaluable in shaping the content and ensuring it reflects both the practice- and evidence-base.

Given the range of featured topics, a Technical Review Team (TRT) of more than 80 individuals representing **local health departments, community-based organizations, national organizations, and CDC subject matter experts** was developed. TRT members contributed expertise in health equity, particular strategies, specific population groups, and/or settings. These individuals were engaged at various stages of development and participated in various ways. Specific TRT members also participated on Strategy Review Teams for each strategy within the Tobacco-Free Living, Healthy Food and Beverage, and Active Living sections of the guide. These individuals served as the main consultative body for the development of each respective strategy. The individuals listed below contributed to sections for which they had expertise. However, they are not responsible for the final content of the guide.



ACKNOWLEDGEMENTS (Continued)

Centers for Disease Control and Prevention

National Center for Chronic Disease Prevention and Health Promotion—

Division of Community Health

Anna Berkowitz
Joyce Buckner-
Brown
Rebecca Bunnell
Shannon Griffin-
Blake
John Francis
Shalon Irving
Leonard Jack, Jr.

Jennifer Kohr
Rashid Njai
Rebecca Payne
Mark Rivera
Angela Ryan
Robin Soler
Pattie Tucker

Division of Diabetes Translation

Larry Alonso Keri Norris

Division of Nutrition, Physical Activity, and Obesity

Latetia Moore
Freeman
Laurence
Grummer-Strawn
Sonia Kim
Joel Kimmons
Beverly Kingsley

Carol MacGowan
Stephen Onufrak
Terry O'Toole
Tom Schmid
Katherine Shealy
Demia Wright

Division of Population Health

Lynda Anderson Holly Hunt
Sherry Everett-
Jones Sarah Lee
Susie McCarthy

Office on Smoking and Health

Steven Babb Brian King
Bridgette Garrett Michael Tynan

National Center on Birth Defects and Developmental Disabilities— Division of Human Development and Disabilities

Jacqui Butler

National Center for Environmental Health

Daneen Farrow-
Collier Dee Merriam
Chris Kochtitzky Margie Walling
Arthur Wendel

National Center for Injury Prevention and Control— Division of Violence Prevention

Sarah Bacon Greta Massetti
Linda Dahlberg Neil Rainford
Marci Hertz

Oak Ridge Institute for Science and Education Research Participation Program at CDC

W. Brad Jones

SciMetrika

Tarisha Cockrell Martha Rider
Lesley Guyot John Wingfield

Representatives from National and Community Organizations

Linda Aragon
*Los Angeles County Tobacco Control
and Prevention Program*

Sabrina Baronberg
*New York City Department of Health
and Mental Hygiene*

Adam Becker
*Consortium to Lower Obesity in
Chicago Children*

Jim Bergman
*Smoke-Free Environments Law
Project and The Center for Social
Gerontology, Inc.*

Elan Bobay
*Monterey County Health Department,
Steps to Healthier Salinas*

Lorraine Boyd
*New York City Department of Health
and Mental Hygiene*

Marie Bresnahan
*New York City Department of Health
and Mental Hygiene*

Isaiah Brokenleg
*Great Lakes Tribal
Epidemiology Center*

Karen Brawley
*Public Health - Seattle and
King County*

Hannah Burton
ChangeLab Solutions

Vicki Carll
Pinellas County Health Department

Caroline Chappell
*North Carolina Division of
Public Health*



ACKNOWLEDGEMENTS (Continued)

Serena Chen
*American Lung Association in
California, Bay Area Smoke-Free
Housing Project*

Kristin Cipriani
*National Association for Sports
and Physical Education*

Ann Cody
BlazeSports America

Keith Cooper
*American Lung Association,
Southwest*

Kerry Cork
Public Health Law Center

Shannon Cosgrove
YMCA of the USA

Dave Cowan
*Safe Routes to School
National Partnership*

Quang Dang
ChangeLab Solutions

Char Day
Americans for Nonsmokers' Rights

Daisy DeLaRosa
Boston Public Health Commission

Dr. Scout
*Network for LGBT Health Equity
at the Fenway Institute*

Delbert Elliott
University of Colorado

Karen Farley
California WIC Association

Mark Fenton
Mark Fenton Associates

Catherine Fields
American Lung Association

Chione Flegal
PolicyLink

Rebecca Flournoy
PolicyLink

Barry Freedman
*St. Louis County Department
of Health*

Roberta Friedman
*Rudd Center for Food Policy
and Obesity, Yale University*

Christine Fry
ChangeLab Solutions

Lark Galloway-Gilliam
Community Health Councils, Inc.

Justin Garrett
*American Lung Association in
California, Center for Tobacco Policy
and Organizing*

Tony Gomez
*Public Health - Seattle and
King County*

Priscilla Gonzalez
Berkeley Media Studies Group

Bob Gordon
*California LGBT Tobacco Education
Partnership (LGBT Partnership)*

Christine Green
National Complete Streets Coalition

Erin Hagen
PolicyLink

Cynthia Hallett
Americans for Nonsmokers' Rights

Muriel Harris
*University of Louisville
School of Public Health and
Information Sciences*

Alisa Haushalter
*Bureau of Population
Health Programs*

Kenneth Hecht
C & K Hecht Consulting

Seung Hee Lee
Johns Hopkins University

Stacy Ignoffo
*Respiratory Health Association
of Metropolitan Chicago*

Delmonte Jefferson
*National African American
Tobacco Control Network*

Allison Karpyn
The Food Trust

Noelle Kleszynski
*Association of American
Indian Physicians*

Ingrid Krasnow
Berkeley Media Studies Group

Hannah Laurison
ChangeLab Solutions

Kathy Lawrence
School Food Focus

Mary Lee
PolicyLink

Christy Lefall
Urban Habitat

Whitney Maegher
NASBE

Shireen Malekafzali
PolicyLink

Jennifer Moore
*Multnomah County
Health Department*



ACKNOWLEDGEMENTS (Continued)

Tammy Morales
Urban Foodlink

Amanda Navarro
PolicyLink

Vu-Bang Nguyen
Urban Habitat

Jeannette Noltenius
Indiana Latino Institute, Inc.

Faryle Nothwehr
*University of Iowa, College
of Public Health*

Ana Novais
Rhode Island Department of Health

Odessa Ortiz
Boston Public Health Commission

Sara Padilla
Community Food Security Coalition

Megan Patterson
Boston Public Health Commission

Katharine Dupont Phillips
Nemours

Lisa Pivec
*Community Health Promotion -
Cherokee Nation*

Janet Porter
Break Free Alliance

Stephanie Ramirez
*National Association of Latino
Elected Officials*

Kurt Ribisl
*University of North Carolina Gillings
School of Global Public Health*

Cheryl Richardson
*National Association for Sports
and Physical Education*

Cynthia Roberts
Rhode Island Department of Health

Catherine Saucedo
*Smoking Cessation Leadership
Center at UCSF*

Sheila Savannah
Houston Public Health Department

Regina Schaefer
American Academy of Pediatrics

Susan Schoenmarklin
*Attorney and former consultant
to the Smoke-Free Environmental
Law Project of the Tobacco Control
Legal Consortium*

Kevin Schroth
*New York City Department of
Mental Health and Hygiene*

Dave Shaibley
Tobacco Control Law Consortium

Brian Smedley
*Health Policy Institute, Joint Center
for Political and Economic Studies*

Seth Strongin
City Project

Lynnette Swain
Pinellas County Health Department

Erika Terl
*National Recreation and
Park Association*

Mary Thomas
*San Antonio Metropolitan
Health District*

Abby Thorne-Lyman
Reconnecting America

Laurie True
California WIC Association

Adrienne Udarbe
*Arizona Department of
Health Services*

Amanda Wagner
*Philadelphia Department of
Public Health*

Elizabeth Walker
*National Association of State Boards
of Education - Center for Safe and
Healthy Schools*

Y. Claire Wang
*Columbia University, Mailman
School of Public Health*

Billie Weiss
Advisor to Safe States Alliance, SAVIR

Stephanie Weiss
The Food Trust

Alzen Whitten
*New York City Department of Health
and Mental Hygiene*

Missy Wilson
Mobile County Health Department

Heather Wooten
ChangeLab Solutions

Ellen Wu
*California Pan-Ethnic
Health Network*

Marc Zimmerman
University of Michigan

Additionally, special thanks to the many unnamed members of local communities presented as examples throughout *A Practitioner's Guide for Advancing Health Equity*.

CDC would also like to thank ICF International, Prevention Institute, and Ogilvy Public Relations for their support in the development and production of the resource.



INTRODUCTION



Heart disease, cancer, diabetes, and stroke are the most common causes of illness, disability, and death affecting a growing number of Americans.⁴ Many of these chronic conditions tend to be more common, diagnosed later, and result in worse outcomes for particular individuals,⁵⁻⁷ such as people of color, people in low-income neighborhoods, and others whose life conditions place them at risk for poor health.

(See Appendix A for list of population groups experiencing chronic disease disparities.)

Despite decades of efforts to reduce and eliminate health disparities, they persist—and in some cases, they are widening among some population groups.⁸⁻¹¹ Such disparities do not have a single cause. They are created and maintained through multiple, interconnected, and complex pathways. Some of the factors influencing health and contributing to health disparities include the following:^{12,13}

- **Root causes or social determinants of health** such as poverty, lack of education, racism, discrimination, and stigma.
- **Environment and community conditions** such as how a community looks (e.g., property neglect), what residents are exposed to (e.g., advertising, violence), and what resources are available there (e.g., transportation, grocery stores).
- **Behavioral factors** such as diet, tobacco use, and engagement in physical activity.
- **Medical services** such as the availability and quality of medical services.

INTRODUCTION (Continued)

HEALTH EQUITY MEANS THAT EVERY PERSON HAS AN OPPORTUNITY TO ACHIEVE OPTIMAL HEALTH REGARDLESS OF:

- THE COLOR OF THEIR SKIN
- LEVEL OF EDUCATION
- GENDER IDENTITY
- SEXUAL ORIENTATION
- THE JOB THEY HAVE
- THE NEIGHBORHOOD THEY LIVE IN
- WHETHER OR NOT THEY HAVE A DISABILITY³

While health disparities can be addressed at multiple levels, this resource focuses on **policy, systems, and environmental improvement strategies** designed to improve the places where people live, learn, work, and play. Many of the 20th and 21st century's greatest public health achievements (e.g., water fluoridation, motor vehicle safety, food safety) have relied on the use of laws, regulations, and environmental improvement strategies.^{14,15} Health practitioners play an important role in these improvements by engaging the community, identifying needs, conducting analyses, developing partnerships, as well as implementing and evaluating evidence-based interventions.

These intervention approaches are briefly described below:

- **Policy improvements** may include “a law, regulation, procedure, administrative action, incentive, or voluntary practice of governments and other institutions.”¹⁶
Example: A voluntary school wellness policy that ensures food and beverage offerings meet certain standards.
- **Systems improvements** may include a “change that impacts all elements, including social norms of an organization, institution, or system.”¹⁷
Example: The integration of tobacco screening and referral protocols into a hospital system.
- **Environmental improvements** may include changes to the physical, social, or economic environment.¹⁷
Example: A change to street infrastructure that enhances connectivity and promotes physical activity.



INTRODUCTION (Continued)

Such interventions have great potential to prevent and reduce health inequities, affect a large portion of a population, and can also be leveraged to address root causes, ensuring the greatest possible health impact is achieved over time. **However, without careful design and implementation, such interventions may inadvertently widen health inequities.** To maximize the health effects for all and reduce health inequities, it is important to consider the following:

- Different strategies require varying levels of individual or community effort and resources, which may affect who benefits and at what rate.
- Certain population groups may face barriers to or negative unintended consequences from certain strategies (see Appendix B for a list of common barriers). Such barriers can limit the strategy's effect and worsen the disparity.
- Population groups experiencing health disparities have further to go to attain their full health potential, so even with equitable implementation, health effects may vary.
- Health equity should not only be considered when designing interventions. To help advance the goal, health equity should be considered in other aspects of public health practice (e.g., organizational capacity, partnerships, evaluation).

A Practitioner's Guide to Advancing Health Equity provides lessons learned and practices from the field, as well as from the existing evidence-base. This resource offers ideas on how to maximize the effects of several policy, systems, and environmental improvement strategies with a goal to reduce health inequities and advance health equity. Additionally, the resource will help communities incorporate the concept of health equity into core components of public health practice such as organizational capacity, partnerships, community engagement, identifying health inequities, and evaluation.

This resource has four major sections:

- Incorporating Health Equity into **Foundational Skills** of Public Health
- Maximizing **Tobacco-Free Living** Strategies to Advance Health Equity
- Maximizing **Healthy Food and Beverage** Strategies to Advance Health Equity
- Maximizing **Active Living** Strategies to Advance Health Equity



TERMINOLOGY



A clear understanding of definitions is important. The following definitions are offered as a starting place as you review this resource:

Health equity: Health equity is attainment of the highest level of health for all people. Achieving health equity requires valuing everyone equally with focused and ongoing societal efforts to address avoidable inequalities, historical and contemporary injustices, and the elimination of health and health care disparities.¹²

Health disparities: Health disparities are differences in health outcomes and their determinants between segments of the population, as defined by social, demographic, environmental, and geographic attributes.⁷

Health inequalities: Health inequalities is a term sometimes used interchangeably with the term health disparities. It is more often used in the scientific and economic literature to refer to summary measures of population health associated with individual- or group-specific attributes (e.g., income, education, or race/ethnicity).⁷

Health inequities: Health inequities are a subset of health inequalities that are modifiable, associated with social disadvantage, and considered ethically unfair.^{7,18,19}

Social determinants of health: Social determinants of health are conditions in the environments in which people are born, live, learn, work, play, worship, and age that affect a wide range of health, functioning, and quality-of-life outcomes and risks.²⁰



SECTION 2 Maximizing Tobacco-Free Living Strategies to Advance Health Equity



USE THE CONTENT TO:

- FOSTER DIALOGUE ON HEALTH EQUITY CONCERNS WITHIN A COMMUNITY.
- TRAIN STAFF AND PARTNERS ON EQUITY ISSUES SURROUNDING TOBACCO-FREE LIVING STRATEGIES.
- IDENTIFY WAYS TO ADDRESS HEALTH EQUITY IN THE DESIGN AND IMPLEMENTATION OF STRATEGIES.
- DEVELOP YOUR OWN APPROACH FOR ENSURING EFFORTS ARE ADDRESSING HEALTH INEQUITIES.

Despite overall declines in cigarette smoking, some population groups have disproportionately higher rates of smoking. These groups include certain racial/ethnic minority groups, particularly American Indians/Alaska Natives; those with low socioeconomic status; those with mental health and substance abuse conditions; those in the lesbian, gay, bisexual, and transgender communities;³⁰ and those with disabilities.³¹ Identifying and eliminating tobacco-related health inequities among population groups is an important component of tobacco control efforts.³²

The Tobacco-Free Living section of *A Practitioner's Guide for Advancing Health Equity* provides equity-oriented considerations, key partners, and community examples related to the design and implementation of the following strategies:

- Comprehensive Smoke-Free Policies
- Smoke-Free Multi-Unit Housing Policies
- Tobacco Cessation Services
- Point-of-Sale Strategies to Address Access and Exposure to Tobacco Products

The content presented is not exhaustive and is not intended to act as a “how-to” guide. Rather, this section is devised to stimulate ideas for ensuring tobacco-free living strategies are designed to address the needs of populations experiencing health inequities. Please refer to the disclaimer on page ii when using this Section.



COMPREHENSIVE SMOKE-FREE POLICIES


COMPREHENSIVE SMOKE-FREE POLICIES MAY INCLUDE STATE OR LOCAL LAWS OR REGULATIONS THAT PROHIBIT SMOKING IN ALL INDOOR AREAS OF WORKSITES AND PUBLIC PLACES, INCLUDING RESTAURANTS AND BARS.^{33,34}

MAKE THE CASE:

Why Is This A Health Equity Issue?

The issues below highlight the need for comprehensive smoke-free policies that advance health equity:

- **Differential Policy Coverage in Workplaces Employing Vulnerable Populations:** Comprehensive smoke-free policies are the most effective means to fully protect all workers from secondhand smoke exposure in workplaces.³³ In contrast, policies that exempt venues, such as restaurants, bars, hotels, casinos, and factories, may exclude many blue-collar and service sector workers from smoke-free protections and create disparities in secondhand smoke exposure.^{35,36} These workers—many of whom are racial or ethnic minorities, immigrants, and individuals with limited education and low incomes—may have disproportionate exposure to secondhand smoke in the workplace.^{37,38}
- **Lack of Enforcement and Compliance with Existing Smoke-Free Policies in Some Communities:** Even when a comprehensive smoke-free policy exists, some groups may not fully benefit from the policy due to inconsistent education and enforcement regarding the policy.^{35,39,40} Lack of community engagement or culturally appropriate efforts to inform these groups about policies and failure to provide these populations with cessation services may also influence who benefits from the policy.
- **Challenges with Adopting Comprehensive Smoke-free Policies in Rural Areas and Tribes:** Some rural areas or tribes may be resistant to smoke-free policies as indicated by higher smoking rates in these areas.^{30,41} Others may be resistant because the economy may rely on tobacco production or use.⁴²⁻⁴⁴ Additionally, in many American Indian and Alaska Native tribes, barriers to such policies may arise if cultural and historical norms regarding ceremonial or traditional tobacco practices are not considered when adopting and implementing smoke-free policies.⁴⁴ Considering the cultural and social norms in communities is critical for the development of successful, smoke-free strategies.



Design and Implement with Health Equity in Mind

To maximize health impact and advance health equity, consider these factors and others when designing, implementing, and evaluating comprehensive smoke-free policies:

KEY FACTORS	BARRIERS OR UNINTENDED CONSEQUENCES	OPPORTUNITIES TO MAXIMIZE IMPACT
<p>COMMUNITY AWARENESS & INVOLVEMENT</p> <p>Engage communities to understand and shift social norms around smoking and secondhand smoke</p>	<p>Some communities, particularly those with high rates of smoking, may be reluctant to implement comprehensive smoke-free policies. In other communities, tobacco control interventions may not be a priority.</p>	<ul style="list-style-type: none"> • Understand the social norms around smoking and secondhand smoke in underserved communities. • Work with organizations that serve population groups experiencing inequities to engage the community. • Use culturally appropriate media and education efforts to build awareness of the health effects of smoking and secondhand smoke exposure in the underserved communities.
<p>CAPACITY & INFRASTRUCTURE</p> <p>Build community capacity and infrastructure to support implementation of comprehensive smoke-free policies</p>	<p>Limited capacity and infrastructure among agencies serving populations experiencing inequities may be a challenge to implementing comprehensive smoke-free policies.⁴⁴ Additionally, some of these organizations may receive financial and other supports from the tobacco industry.⁴⁵⁻⁴⁷</p>	<ul style="list-style-type: none"> • Prioritize inclusion of organizations serving or working with populations experiencing inequities in tobacco control coalitions. • Identify community leaders and train them to educate stakeholders about the disparities that result when policies are not prioritized in underserved communities. • Use partnerships to leverage resources. Explore funding opportunities to support organizations that want to join smoke-free implementation efforts.
<p>ACCESS TO CESSATION SERVICES</p> <p>Integrate cessation support as part of a comprehensive approach</p>	<p>Given existing inequities in access to and quality of health care,⁴⁸ access to cessation supports and services may vary.^{49,50}</p>	<ul style="list-style-type: none"> • Incorporate free or low-cost cessation services before and during policy implementation to help motivated individuals quit. • See strategy on Tobacco Cessation Services for more information.
<p>LACK OF SUPPORTIVE DATA</p> <p>Identify and track health inequities</p>	<p>Lack of timely and comprehensive data that fully explore health inequities may be a barrier to tobacco control efforts⁴⁹ (e.g., data examining inequities in secondhand smoke exposure among different groups).</p>	<ul style="list-style-type: none"> • Improve collection and use of standardized data across population groups (e.g., geography, occupation, sexual orientation) to assess inequities in secondhand smoke exposure and policy coverage. • Use findings to identify where interventions are needed, monitor effects of an intervention, and track progress in addressing health inequities.
<p>VARIABILITY IN IMPLEMENTATION & ENFORCEMENT</p> <p>Expand smoke-free policies and institutional practices</p>	<p>State and local smoking restrictions may not cover all settings, including certain worksites (e.g., bars and casinos), outdoor public spaces (e.g., dining areas, construction sites), and institutions (e.g., mental health and substance abuse treatment facilities).</p>	<ul style="list-style-type: none"> • Eliminate exemptions in existing smoking restrictions. • Prioritize efforts in institutions with high rates of secondhand smoke exposure when local or state policies do not cover these settings. • Use media to address the health benefits of smoke-free policies and any misperceptions about these policies. • Develop appropriate enforcement mechanisms to support policy implementation.

Build the Team: Partnership for Success

Successful efforts to implement comprehensive smoke-free policies depend on bringing a diverse set of partners to the table early, consistently, and authentically. These partners may include the following:

- Area Agencies on Aging
- Cessation support services
- Community members (of diverse abilities, ages, cultures, gender, income levels, race/ethnicity, and sexual orientation)
- Employee rights groups
- Health care systems, hospitals, community clinics, and health care providers
- Leaders and community champions from multiple sectors
- Local businesses
- Mental health and substance abuse treatment facilities
- Organizations serving populations experiencing health inequities
- Parks and recreation department
- Public health agencies
- School districts, universities, and community colleges
- State tobacco control programs
- Tobacco control groups
- Youth, the elderly, and people with disabilities



Native American Tribes Adopt Tobacco Protections for Tribal Members and Future Generations

Montana

In Montana, 43% of Native American adults self-report cigarette smoking.⁵¹ These high rates of commercial tobacco products use contributes to high rates of disease and premature death among Montana's Native Americans.⁵² To address the commercial use of tobacco in their communities, the Blackfeet and Fort Peck tribes worked together to implement comprehensive smoke-free indoor protections. These protections also safeguard casino visitors and employees from secondhand smoke.

Respect for cultural traditions of tobacco use was instrumental in the development and implementation of strategies to create smoke-free environments. Several years earlier, the Native American Tobacco Coalition of Montana approached tribal elders to ask if they would support the creation of smoke-free environments. Initially, the elders were not supportive, because they believed this could potentially hinder traditional uses of tobacco, which are rooted in spiritual beliefs and medicinal

practices. The elders engaged in a four-year process of teaching the historical and ceremonial practices of traditional tobacco use, including spiritual offerings. In turn, the coalition educated elders about the impact of commercial tobacco use and secondhand smoke exposure on tribal youth and future generations.

With support from the elders, the coalition educated the tribal members about the distinction between the sacred use of tobacco and the use of commercial tobacco. Community engagement activities included commercial tobacco-free celebrations, health fairs, youth-focused events, and trainings. By conducting extensive educational initiatives for tribal members and elders, the Blackfeet and Fort Peck Tribal Nations were able to create smoke-free indoor environments that included casinos. As a result, other tribes have created smoke-free environments in most tribal facilities. The coalition learned a valuable lesson: to be successful, smoke-free strategies need to be true to the people and rooted in cultural tradition.



Smokefree table tent at Birmingham, AL restaurant. Photo courtesy of the JCDH and the HAP.

Partnerships and Educational Initiatives Lead to Smoke-Free Air Protections

Birmingham, AL

Jefferson County Department of Health (JCDH) and the Health Action Partnership (HAP) are helping to implement smoke-free protections in Birmingham—impacting approximately 356,000 Jefferson County residents and commuters. With support from CDC's *Communities Putting Prevention to Work* program, the health department conducted community needs assessments and used geographic information systems (GIS) mapping to track rates of smoking, heart attack, and cancer to identify communities with the highest need for smoke-free protections. Then they overlaid those maps with maps of low-income areas.

After identifying high-need communities, JCDH and HAP conducted evaluation interviews in these areas to assess the key organizations and community champions that could become a conduit for educating residents on secondhand smoke issues. Working with a variety of local organizations, faith-based leaders, and the media, the community was able to successfully

educate and increase community awareness about the benefits of smoke-free environments. The Friends of West End, a local organization with strong ties to the targeted communities, educated nearly 100 neighborhood association presidents. The presidents then educated their respective communities while local pastors did the same among their congregations. JCDH's understanding of culturally appropriate educational media led to a well-received radio soap opera, *Live Well Camberwell*. The educational radio program and health expert interviews were aired on stations with largely African American audiences.

All of these educational initiatives contributed to increasing awareness around the health effects of secondhand smoke exposure in indoor places of employment including restaurants, bars, and hotels. When smoke-free protections were put in place, the HAP provided technical assistance to ensure proper implementation of and compliance with the smoke-free protections.



SMOKE-FREE MULTI-UNIT HOUSING POLICIES


SMOKE-FREE MULTI-UNIT HOUSING POLICIES AIM TO PROTECT NONSMOKERS WHO LIVE IN, WORK, AND VISIT MULTI-UNIT RESIDENCES SUCH AS APARTMENTS, CONDOMINIUMS, TOWNHOUSES, DUPLEXES, AND AFFORDABLE HOUSING COMPLEXES FROM SECONDHAND SMOKE.

MAKE THE CASE:

Why Is This A Health Equity Issue?

The issues below highlight the need for smoke-free multi-unit housing policies that advance health equity:

- **Increased Secondhand Smoke Risks among Vulnerable Populations:** Low-income individuals generally have higher smoking rates,³⁰ which may result in increased exposure to secondhand smoke in affordable and public housing. Given that many residents living in these settings are vulnerable population groups (e.g., children, older adults, racial/ethnic minorities, and those with a disability),⁵³ secondhand smoke exposure is critical to address. For instance, many racial/ethnic minorities and low-income populations suffer higher rates of asthma and other tobacco-related health issues,⁵⁴ making them particularly vulnerable to the effects of secondhand smoke exposure. Children are also vulnerable to developing health effects from secondhand smoke exposure.⁵⁵
- **Residents Who are Being Exposed to Secondhand Smoke May Have Limited Alternative Housing Options:** Even when residents of multi-unit housing do not allow smoking in their unit, secondhand smoke can enter their unit from other units or common areas through shared ventilation systems, air spaces, windows, and hallways, putting residents at risk.^{54,56,57} Low-income residents in affordable or public housing complexes who are being involuntarily exposed to secondhand smoke in their homes in this manner may have limited alternative housing options or be unable to move.



Design and Implement with Health Equity in Mind

To maximize health impact and advance health equity, consider these factors and others when designing, implementing, and evaluating smoke-free multi-unit housing policies:

KEY FACTORS	BARRIERS OR UNINTENDED CONSEQUENCES	OPPORTUNITIES TO MAXIMIZE IMPACT
<p>MISPERCEPTIONS</p> <p>Clarify intent and address misperceptions about smoke-free multi-unit housing strategies</p>	<p>Organizations that work on behalf of low-income residents (e.g., residents' rights organizations, affordable housing groups) may have misconceptions about the intent or effects of smoke-free multi-unit housing strategies.</p>	<ul style="list-style-type: none"> • Ensure residents and owners understand that the smoke-free policy is designed to promote a healthy home environment and reduce secondhand smoke exposure—not to remove smokers or prevent new smokers from moving in, as long as they comply with the policy. • Consider working across different types of multi-unit housing (e.g., public, affordable, and market-rate) to show everyone deserves clean air and prevent concerns about discrimination.
<p>STAKEHOLDER SUPPORT</p> <p>Address concerns and build support among housing providers</p>	<p>If landlords are unaware of resident support for smoke-free policies, they may have concerns that such a policy will lead to reduced occupancy or will be difficult to enforce.</p>	<ul style="list-style-type: none"> • Alleviate concerns of stakeholders (e.g., landlords, apartment owners) by educating them on the business, health, and safety benefits of smoke-free policies. • Provide tools and training on how to gather resident feedback, navigate the implementation process, develop monitoring mechanisms, and address noncompliance.
<p>COMMUNITY INVOLVEMENT & AWARENESS</p> <p>Engage residents in strategy development and implementation</p>	<p>Strategies developed without resident input may negatively affect strategy implementation and enforcement.</p>	<ul style="list-style-type: none"> • Identify residents to serve as champions and involve them throughout the development and implementation process. • Gather input through resident surveys and forums, and collaborate with resident associations to develop the strategy. • Establish mechanisms to ensure that residents are aware of the policy and its benefits (e.g., culturally appropriate education initiatives, resident forums).
<p>ACCESS</p> <p>Increase access to free or low-cost cessation services</p>	<p>Residents without access to cessation supports may not comply with smoke-free policies, placing other residents at risk for secondhand smoke exposure.</p>	<ul style="list-style-type: none"> • Provide smokers with free access to evidence-based cessation treatments to ease the transition to a smoke-free environment, increase compliance, prevent smokers from feeling stigmatized, and help them quit smoking, thus maximizing the policy's health benefits. • Offer and promote cessation services in or near the complex at convenient times before and during policy implementation.
<p>EQUITABLE IMPLEMENTATION</p> <p>Anticipate additional challenges to policy implementation</p>	<p>After policy implementation, smokers may tend to congregate in outdoor areas near buildings, exposing residents in outdoor or adjacent indoor areas to secondhand smoke, which may enter the building through doors, windows, or vents.</p>	<ul style="list-style-type: none"> • Make the entire property smoke-free or restrict smoking to a few designated outdoor areas located far enough away from entrances and exits to prevent secondhand smoke from infiltrating indoor areas. • Improve compliance by conducting tailored resident engagement, education, and cessation efforts. • In supportive housing settings, (e.g., homeless shelters, mental health and substance abuse treatment facilities), work with staff to find tailored, context-specific approaches.

Build the Team: Partnership for Success

Successful efforts to implement smoke-free multi-unit housing policies depend on bringing a diverse set of partners to the table early, consistently, and authentically. These partners may include the following:

- Area Agencies on Aging
- Cessation support services
- Community members (of diverse abilities, ages, cultures, gender, income levels, race/ethnicity, and sexual orientation)
- Condominium owners
- Faith-based organizations
- Housing industry organizations, local housing authorities, and nonprofit housing associations
- Immigrants and refugees
- Low-income residents
- Organizations serving populations experiencing health inequities
- Real estate developers
- Residential property owners, management companies
- State tobacco control programs
- Residents' rights/fair housing organizations
- Youth, the elderly, and people with disabilities



HEALTH EQUITY IN ACTION

Creating Healthy Environments through Smoke-Free Multi-Unit Housing Policies

San Antonio, TX

Everyone has the right to breathe clean air, especially in their own home. Yet in many San Antonio multi-unit housing complexes, residents were being exposed to secondhand smoke infiltrating their units from neighboring units. To address this issue, the San Antonio Metropolitan Health District (SAMHD) worked with the San Antonio Housing Authority (SAHA) to implement a smoke-free multi-unit housing policy for all 70 of its housing sites. These efforts were made possible with support from a combination of state and local funds.

SAMHD focused on making multi-unit housing smoke-free with a strategy that covered all indoor areas, since a comprehensive policy would have a greater impact on health equity for all SAHA residents and staff. Both organizations recognized the importance of community engagement to ensure policy success. They worked closely with residents and staff in mini-community centers associated with each housing campus. Engaging

the residents through the community centers—a resource that residents were already turning to for information—was key to the success of this approach.

SAMHD bolstered its outreach and education efforts through a partnership with the American Cancer Society (ACS). ACS educated community center staff on how to answer questions about the new policy, to refer residents to the state quitline, and to discuss options for obtaining nicotine replacement therapy (e.g. nicotine patches). All materials were printed in both English and Spanish, and staff in many centers were able to communicate the benefit of smoke-free air protections to residents in their own language.

ACS and SAHA also helped organize town hall meetings to educate residents and discuss the benefits of smoke-free protections in an open forum. As a result of the policy, nearly 16,000 residents (many of them low-income immigrants and racial/ethnic minorities) now have access to cleaner air in their homes.



Housing Authority and Public Health Commission Partner on Smoke-Free Housing

Boston, MA

Building relationships with the Boston Housing Authority, other city agencies, and community-based organizations, the Boston Public Health Commission (BPHC), with support from the CDC's *Communities Putting Prevention to Work* program, educated housing providers about the voluntary adoption of smoke-free multi-unit housing policies. The Commission has also successfully leveraged support for smoke-free housing among residents in all market sectors (e.g., market rate, public housing), as well as used data to identify disparities in health outcomes across residents of different types of housing. The goal is to ensure that residents, particularly those most vulnerable to secondhand smoke exposure, have clean air to breathe in their homes. Smoke-free protections in public housing, affordable housing, and market-rate housing have the potential to provide protection for populations with high exposure to secondhand smoke, including low-income families, children with asthma, immigrants, elderly residents, and persons with chronic diseases or disabilities.

The BPHC has taken a series of steps to increase smoke-free protections in and around multi-unit housing complexes. One key to the success of the BPHC's policy was its approach of working closely with stakeholders such as the Boston Housing Authority and landlords, property management companies, and community development

corporations. Input from residents in policy development was critical, and mini-grants to community development corporations supported community engagement. The BPHC also worked strategically with a variety of community partners to make the case to landlords and housing agencies that a majority of residents wanted smoke-free housing, and that going smoke-free would benefit all stakeholders. The BPHC provided technical assistance in developing and implementing a smoke-free multi-unit housing policy. Because Massachusetts has relatively good coverage of smoking cessation treatments, many smokers in the state can access affordable cessation services.

The BPHC approach led to some notable early successes. The Boston Housing Authority portfolio became 100% smoke-free in September 2012, assuring a healthier environment for the 23,000 residents living in their 12,000 units. Community development corporations have transitioned over 600 units to smoke-free status, and continue to bring additional units on line as smoke-free. There are now 6,600 non-public smoke-free units and the BPHC sees this as just the beginning. As Margaret Reid, Director of the Division of Healthy Homes and Community Supports at the BPHC, says, "We are supporting smoke-free policies and awareness so that smoke-free becomes the norm in Boston multi-unit housing, whether it is public, subsidized, or market-rate."



TOBACCO CESSATION SERVICES

TOBACCO CESSATION STRATEGIES HELP PEOPLE QUIT SMOKING OR USING OTHER FORMS OF TOBACCO. THESE STRATEGIES MAY INCLUDE CLINICAL SCREENING AND REFERRAL SYSTEMS, QUIT LINES, BEHAVIORAL COUNSELING, AND CESSATION MEDICATIONS.⁵⁸

MAKE THE CASE:

Why Is This A Health Equity Issue?

The issues below highlight the need for tobacco cessation strategies that advance health equity:

- **Varying Rates of Smoking and Cessation among Different Communities:** In the United States, certain population groups stand out as having higher-than-average smoking rates, lower-than-average cessation rates, or higher-than-average rates of tobacco-related diseases.^{49,59,60} For example, population groups with disproportionately high rates of smoking include American Indian and Alaska Natives,^{7,60} the lesbian, gay, bisexual, and transgender (LGBT) communities,⁶¹⁻⁶³ people with mental illness⁶⁴ and substance abuse conditions,⁷ and people with disabilities.⁶⁵ African American adults have been found to be more likely to express interest in quitting and more likely to have tried to quit in the past year than white adults, but are less likely to use proven treatments (e.g., counseling and/or medications) and are less likely to succeed in quitting.⁵⁹ Adults aged 65 or older have also been found to be less likely to attempt smoking cessation compared to younger adults.^{59,66} Additionally, low-income populations are more likely to smoke, less likely to quit, and often lack access to affordable cessation support.^{49,60,67}
- **Barriers to Accessing Cessation Resources:** Differential access and quality of health care may present barriers to quitting.⁵⁰ For example, uninsured smokers may be less likely to receive quitting advice or other forms of cessation treatment from health care providers.⁵⁰ Additionally, certain population groups, including African Americans and Hispanics, are less likely to be screened for tobacco use or receive smoking cessation interventions.^{59,68-70}
- **Challenges to the Widespread Use of Evidence-Based Cessation Interventions:** There is limited research on effective approaches for promoting and increasing utilization of cessation interventions among population groups experiencing health disparities.^{71,72} A lack of sensitivity to social norms and cultural traditions in developing cessation interventions may influence intervention use and ultimate effectiveness.



Design and Implement with Health Equity in Mind

To maximize health impact and advance health equity, consider these factors and others when designing, implementing, and evaluating tobacco cessation strategies:

KEY FACTORS	BARRIERS OR UNINTENDED CONSEQUENCES	OPPORTUNITIES TO MAXIMIZE IMPACT
<p>ACCESS</p> <p>Increase access to cessation services by integrating them into health systems and making them convenient</p>	<p>Certain population groups are less likely to be screened for tobacco use or receive tobacco cessation counseling.^{69,70} Additionally, cessation services may be underused because of limited knowledge, awareness, and cultural beliefs.^{73,74}</p>	<ul style="list-style-type: none"> • Prioritize integration of tobacco screening and provision of/referral for evidence-based cessation treatments into institutions that are already serving vulnerable populations (e.g., community health centers, Federally Qualified Health Centers, rural health clinics, mental health and substance abuse treatment facilities). • Prioritize the promotion of existing cessation services including tobacco quitlines and Web sites to populations experiencing health disparities. • Integrate cessation programs and support into community institutions located in places where people already go (e.g., public housing, faith-based settings, social service agencies). • Train community health outreach workers to provide cessation services during home visits.
<p>COST</p> <p>Remove/reduce cost and insurance barriers</p>	<p>Costs associated with cessation services, which may result from lack of/inadequate health insurance coverage pose particularly significant barriers for low-income populations.⁷⁵⁻⁷⁷</p>	<ul style="list-style-type: none"> • Develop relationships with private and public health insurers and health care systems, including state Medicaid programs, to expand insurance coverage of cessation services. • Consider ways to eliminate or minimize cost and other barriers (e.g., co-pays, prior authorization requirements) to accessing cessation treatments.
<p>DIVERSE NORMS AND CUSTOMS</p> <p>Ensure that cessation services are culturally relevant and appropriate</p>	<p>Limited research on effective approaches to promote cessation services interventions across different groups may hinder the utilization of such interventions.^{49,71,78} For example, tobacco quitlines may be accessed less by groups with cultural norms that avoid seeking counseling from strangers.</p>	<ul style="list-style-type: none"> • Evaluate the effectiveness of cessation interventions across different population groups. • Ensure that underserved populations have access to and are aware of cessation services (e.g., promote services through culturally appropriate communication channels). • If such populations are still not using, or are unsatisfied with, existing cessation services, partner with relevant organizations to increase culturally relevant training of providers or to tailor these services to meet the populations' needs.

Build the Team: Partnership for Success

Successful efforts to implement tobacco cessation strategies depend on bringing a diverse set of partners to the table early, consistently, and authentically. These partners may include the following:

- Academic institutions
- Cessation support services
- Community-based organizations
- Community health centers, including Federally Qualified Health Centers and rural health clinics
- Community members (of diverse abilities, ages, cultures, gender, income levels, race/ethnicity, and sexual orientation)
- Employers
- Health care systems
- Health insurers
- Lay health providers/promotoras
- Leaders and community champions from multiple sectors
- Media
- Mental health and substance abuse treatment facilities
- National culturally specific cessation guidance organizations
- Public health agencies
- State quitline providers
- State tobacco control programs
- Tobacco control groups
- Workplace wellness organizations



HEALTH EQUITY IN ACTION

Expanding Cessation Services in Marginalized Communities

St. Louis, MO

With the goal of reducing tobacco-related health disparities, the St. Louis County Department of Health (DOH), with support from CDC's *Communities Putting Prevention to Work* program, set out to increase access to cessation services among populations with high smoking rates. "Being in their neighborhood and speaking their language" was a critical strategy for helping people most in need of cessation resources, noted Barry Freedman, Project Manager for the *Communities Putting Prevention to Work* program at the DOH. DOH partnered with three trusted organizations that had strong ties in the community to provide free and low-cost services and culturally competent care. Each partnership is briefly described below.

Casa de Salud, a local health clinic, provides cessation services to low-income and limited-English-speaking Hispanic populations, including onsite one-on-one cessation counseling and nicotine replacement therapy (NRT), such as nicotine patches. All of this is done using culturally appropriate materials in an environment where clients can feel safe.

SAGE Metro St. Louis works with the lesbian, gay, bisexual, and transgender (LGBT) communities and

offers cessation counseling services and NRT free of charge. SAGE also provides education on the techniques the tobacco industry has used to target LGBT communities. Having a presence at the city's three major gay pride events proved an effective outreach and education approach.

The St. Louis Christian Chinese Community Service Center worked to provide cessation services, including individual counseling, support, and other resources, to Asian-American restaurant employees with high rates of smoking. Because of limited health literacy in the communities they serve, the Center conducted traditional Chinese puppet shows to encourage cessation and provided health information to over 500 Chinese Americans. The shows are especially powerful because they respect Chinese cultural norms while conveying important health messages to multiple generations.

In addition to these partnerships, the DOH is helping support low-income and uninsured community clinic clients. DOH is training providers in those clinics to facilitate a free cessation program, and is offering a free three-month supply of NRT products to smokers who want to quit.



Using Partnerships to Increase Access to Cessation Services

Santa Clara County, CA

“A one-stop shop [is] more likely to work than having people go to different places for [tobacco cessation] services,” noted Kris Vantornhout, Program Manager for the *Communities Putting Prevention to Work* program at the Santa Clara County Public Health Department (SCCPHD). With this in mind, Santa Clara County’s Tobacco Prevention Program strategically partnered with established community-based organizations (CBOs) throughout the county. These CBOs were poised to implement cessation services in neighborhoods with high numbers of smokers. At times, securing diverse community leadership involvement was challenging, but focused efforts were successful in finding champions to lead the way.

SCCPHD awarded twenty-seven mini-grants to CBOs to expand cessation counseling, referrals, and access to nicotine replacement therapy (NRT). These grants supported organizations working with the Vietnamese, African American, Latino, and Lesbian, Gay, Bisexual, and Transgender (LGBT) communities, and engaged diverse partners. Efforts also focused on improving cessation services and referral systems in mental health facilities, health care clinics, and college campuses.

CBOs integrated cessation services into organizational practice by implementing the “ask, advise, refer” model during intake processes and, as appropriate, referring patients or students to trained staff for cessation assistance. Providing culturally and linguistically relevant messaging around secondhand smoke exposure was also important for increasing the uptake of cessation services. Tobacco-free messaging and cessation information were shared onsite, as well as at outreach events such as the San Jose LGBT Pride Celebration, the annual Martin Luther King Luncheon, and the Holiday Fair.

As a result of the Tobacco Prevention Program’s partnership efforts, cessation services are now available to some of the most vulnerable populations



Smoke-free sign on medical center campus in Santa Clara County, CA. Photo courtesy of Breathe California.

in the county. Thirty health facilities, 8 colleges, and 11 CBOs now have staff or clinicians using the “ask, advise, refer” model to reach over 544,000 residents. Approximately 8,000 units of NRT were distributed through these networks within less than two years, and post-intervention surveys have shown an overall 39% quit success rate, with an amazing 50% quit rate within the Vietnamese community.



POINT-OF-SALE STRATEGIES TO ADDRESS ACCESS AND EXPOSURE TO TOBACCO PRODUCTS

POINT-OF-SALE (POS) STRATEGIES TO REDUCE THE AVAILABILITY AND APPEAL OF TOBACCO PRODUCTS MAY INCLUDE ADDRESSING THE MARKETING AND AFFORDABILITY OF THESE PRODUCTS THROUGH RESTRICTIONS ON THE POINT-OF-SALE TOBACCO ADVERTISING, PROMOTION (INCLUDING PRICE PROMOTIONS), DISPLAYS, AND PLACEMENT; THE SALES OF SPECIFIC TYPES OF TOBACCO PRODUCTS (E.G., FLAVORED PRODUCTS); AND TOBACCO RETAILER LOCATION AND DENSITY.^{79-81,*}


MAKE THE CASE:

Why Is This A Health Equity Issue?

The issues below highlight the need for POS strategies that advance health equity:

- **Different Exposure to POS Advertising and Targeted Marketing:** Studies have consistently shown that low-income communities⁸² and communities of color are more heavily exposed to POS tobacco advertising than other communities.^{83,84} Additionally, such advertising may be targeted to or disproportionately impact certain population groups. For instance, the messaging used in marketing menthol cigarettes has been culturally tailored and targeted toward communities of color, especially African Americans.⁸⁵
- **Placement of and Price Discounts on Tobacco Products:** Tobacco companies have used a variety of point-of-sale strategies to place tobacco products prominently in the retail environment and keep these products affordable.⁸⁶ For example, in 2011, the tobacco industry spent an estimated \$8 billion, or nearly \$23 million per day, on cigarette advertising and promotional expenses in the United States alone. Approximately 84% (or nearly \$7 billion) of this expenditure was spent on price discounts to cigarette retailers or wholesalers to reduce the price of cigarettes to consumers.⁸⁷ A placement strategy may include placing tobacco products (e.g., cigarillos, cigars) next to candy or within the view of children and youth.⁸⁸ Additionally, tobacco companies may deeply discount their products in stores in lower-income communities and require targeted placement of signs advertising lower prices in these stores.⁸⁹ Youth and low-income individuals may be particularly sensitive to prominently placed, inexpensive tobacco products.^{79,90}
- **Greater Density of Tobacco Retailers in Underserved Communities:** Research has shown that tobacco retail outlets are more heavily concentrated in low-income communities and communities of color than in higher income communities.⁹¹⁻⁹⁴ This makes tobacco products more readily accessible, potentially increasing consumption.⁹⁴

Note: The Tobacco Control Act preserves the authority of state, local, and tribal governments to regulate tobacco products in certain specific respects. It also prohibits, with certain exceptions, state and local requirements that are different from, or in addition to, requirements under the provisions of the FDCA relating to specified areas.



Design and Implement with Health Equity in Mind

To maximize health impact and advance health equity, consider these factors and others when designing, implementing, and evaluating POS strategies:

KEY FACTORS	BARRIERS OR UNINTENDED CONSEQUENCES	OPPORTUNITIES TO MAXIMIZE IMPACT
<p>RESOURCE & FUNDING LIMITATIONS</p> <p>Prioritize and prepare resources for communities in greatest need</p>	<p>Underserved communities may have fewer resources to implement POS strategies. Additionally, such communities may have other priorities that make it difficult to implement tobacco control initiatives.</p>	<ul style="list-style-type: none"> • Conduct assessments to examine tobacco retail density, the amount of POS advertising, and tobacco-related health disparities. • Prioritize intervention efforts to address areas with greatest need. • Engage partners who can provide technical assistance to identify viable POS strategies and overcome barriers. • Participate in community coalitions and events in order to understand community priorities, align POS efforts with those priorities, and educate and mobilize the community around these efforts.
<p>ECONOMIC STABILITY</p> <p>Support retailers when implementing POS strategies</p>	<p>Underserved communities, which can have disproportionately high concentrations of tobacco retailers, may oppose POS strategies due to concerns about the potential financial effects on local businesses.</p>	<ul style="list-style-type: none"> • Find creative mechanisms to support retailers that are implementing POS strategies. • Establish programs that may help retailers transition from relying on sales of tobacco product to selling healthier products (e.g., Healthy Food Financing,⁹⁵ healthy corner store initiatives⁹⁶).
<p>COMMUNITY AWARENESS</p> <p>Build community awareness and skills to counter tobacco advertising</p>	<p>Tobacco product displays and POS advertising may distort perceptions of the pervasiveness of tobacco use among adolescents,⁹⁷ increase the likelihood of youth smoking initiation,^{97,98} and may prompt impulse buys (e.g., among smokers who are trying to quit).⁹⁹</p>	<ul style="list-style-type: none"> • Increase community awareness of industry marketing tactics to help people critically assess the advertising they see around them. • Work with media outlets serving specific population groups to reinforce positive messaging and to counter any negative effects of tobacco marketing.
<p>VARIABLE IMPLEMENTATION & ENFORCEMENT</p> <p>Support Implementation and Enforcement of POS Strategies, particularly in communities with tobacco-related inequities.</p>	<p>Disparities may increase if POS strategies are not fully implemented or enforced in communities with high smoking rates or social norms that support tobacco use.</p>	<ul style="list-style-type: none"> • Consider developing implementation plans to support consistent and equitable policy compliance. • Consider establishing processes for accountability and gathering feedback from all communities affected by the policy.

Build the Team: Partnership for Success

Successful efforts to implement POS strategies in communities with tobacco-related disparities depend on bringing a diverse set of partners to the table early, consistently, and authentically. These partners may include the following:

- Community members (of diverse abilities, ages, cultures, gender, income levels, race/ethnicity, and sexual orientation)
- Healthy food retail groups
- Leaders and community champions from multiple sectors
- Local governments
- Local store owners
- Public health agencies
- Public Works Department
- School districts, universities, and community colleges
- Senior centers
- State tobacco control programs
- Tobacco control groups (including groups representing populations experiencing health disparities)
- Youth volunteers/coalitions
- Zoning and planning organizations



Citywide Restrictions Tackle Flavored Tobacco Products

New York, NY

“The price point for entry into the world of tobacco is one dollar.” Kevin Schroth, Senior Legal Counsel for Policy, at New York City’s (NYC) Department of Health and Mental Hygiene was describing sales of single, flavored cigarillos and little cigars. These inexpensive alternatives to cigarettes are heavily marketed in stores in low-income and minority urban neighborhoods. These products are often flavored, come in brightly colored packages, and are prominently placed in stores next to gum and candy—all features that make them appealing to children and youth.

“Kids spend twice as much time in convenience stores as adults,” said Schroth. “It’s not a coincidence that these flavored tobacco products look similar to other products marketed to kids.” The point-of-sale marketing of these products contribute to perceptions among youth that these products are easily accessible and that their use is acceptable and cool.

The health department partnered with the New York City Coalition for a Smoke-Free City and community-based organizations such as Korean Community Services, which serves an Asian population with high smoking rates. Together, with support from state and city funds, they educated community stakeholders about the health risks associated with flavored non-cigarette tobacco products. In 2009, the NYC City Council passed a law that restricted the sale of flavored non-cigarette tobacco products, with the exception of menthol, mint, wintergreen, and tobacco-flavored products, in stores throughout the city, with the exception of tobacco bars. The strategy is part of a comprehensive approach to protect all residents, especially impressionable youth, in all New York City communities from tobacco industry marketing. The example has motivated other local jurisdictions, such as nearby Providence, Rhode Island, to implement similar measures.



Tobacco-Free Pharmacies Promote Health for All

San Francisco, CA

Many people go to pharmacies to purchase medications and other items to improve their health. Why then do many major pharmacy chains and independent pharmacies sell tobacco products which contribute to severe health effects? The California LGBT Tobacco Education Partnership (the Partnership) saw this discrepancy as an opportunity to decrease the widespread availability of tobacco products.

With the high rates of smoking within the LGBT population, the Castro District (historically considered the center of San Francisco's LGBT communities) was the Partnership's priority location in its attempt to eliminate tobacco sales in pharmacies. The Partnership strategically engaged independent pharmacies that were already tobacco-free, acknowledging them with certificates. These pharmacies also educated community stakeholders on the benefits of tobacco-free pharmacies. In addition, a public opinion survey of smokers and nonsmokers assessed the need for tobacco-free pharmacies.

To ensure that every San Francisco resident, not only those living in the Castro District, has access to tobacco-free pharmacies, the Partnership, with

support from the California Department of Public Health, Tobacco Control Program and others, implemented a citywide strategy. Working with diverse partners, including the San Francisco Tobacco-Free Coalition, the University of California at San Francisco School of Pharmacy, and the Board of Supervisors, the Partnership made a compelling research-supported argument that pharmacies should be hubs for health, and that this role was inconsistent with selling tobacco products. Even in the face of opposition from some local media outlets and national retailers, the Partnership remained focused on the message that health for all should be the priority.



On October 1, 2008, San Francisco became the first city in the United States to eliminate the sale of tobacco products in its pharmacies (through changes to the Health Code), affecting an estimated 100 pharmacies and all 805,000 city residents. The policy's success has inspired similar efforts in cities from nearby Richmond, CA to Boston, MA. As Bob Gordon of the Partnership stated, the policy has "changed a social norm around the availability and accessibility of tobacco products in San Francisco. That alone is an amazing outcome."



APPENDICES



APPENDIX A

Health Disparities in Chronic Disease Risk Factors by Population Group

APPENDIX B

Considerations for Health Equity-Oriented Strategy Selection, Design, and Implementation

APPENDIX C

Example Resources for Identifying and Understanding Health Inequities


APPENDIX D

Health Equity Checklist: Considering Health Equity in the Strategy Development Process



HEALTH DISPARITIES IN CHRONIC DISEASE RISK FACTORS BY POPULATION GROUP

Despite decades of effort to reduce and eliminate health disparities, they have largely persisted—and in some cases are widening.⁹⁻¹¹ Specifically related to chronic diseases, there is a concentrated, disproportionate burden of chronic disease in many underserved populations and communities. The table below describes disparities in chronic disease risk factors by various population groups.


**PEOPLE OF COLOR
(RACIAL/ETHNIC
MINORITIES)**

According to the 2010 Census, approximately 16% of Americans identified themselves as Hispanic or Latino, 13% as Black, 5% as Asian, 1% as American Indian and Alaska Native, and 0.2% as Native Hawaiian and other Pacific Islander.²³¹ On a variety of health indicators, significant disparities among these racial and ethnic minorities continue to exist.^{7,232} For example, adult obesity rates in the U.S. are higher among non-Hispanic African Americans (50%) and Mexican Americans (40%) than among non-Hispanic Whites (35%), and they are highest among African American women, at 59%.²³³ In 2011, cigarette smoking among adults was highest among American Indian/Alaska Native populations (32%), compared to other racial/ethnic groups.²³⁴

**PEOPLE WITH
MENTAL OR
SUBSTANCE USE
DISORDERS**

In the United States, adults with mental or substance use disorders comprise approximately 25% of the population. However, this population accounts for an estimated 40% of all cigarettes smoked resulting in a disproportionate burden from the health consequences of smoking.²³⁵

**PEOPLE LIVING
IN RURAL
COMMUNITIES**

Approximately 19%, or 60 million Americans, live in rural areas.²³⁶ Rural residents are more likely to be elderly, in poverty, in fair or poor health, and to have chronic health conditions.⁴⁸ For example, the prevalence of obesity is higher in rural adults (40%) than urban adults (33%).²³⁷ Adults living in non-metropolitan counties also have a higher average annual percentage of smoking (27%) than adults living in large metropolitan counties (18%).²³⁸

**PEOPLE WITH
DISABILITIES**

Approximately 20% of U.S. adults have a disability.²³⁹ Approximately 28% of adults with disabilities smoke, compared to 16% of those without a disability.³¹ Adults with disabilities are more likely to be physically inactive (22%) than are adults without disabilities (10%).²⁴⁰ Obesity is also higher among adults with a disability (38%) compared to those without a disability (24%), according to self-reported data.²⁴¹

**PEOPLE WITH
LOW-INCOME
AND THOSE
EXPERIENCING
POVERTY**

In 2011, an estimated 15% of the U.S. population lived below the federal poverty level.¹⁵² Poverty is correlated with perceived and actual poor health outcomes. People living in poverty are five times more likely to report their health as “poor” compared to high-income individuals.²⁴² People with a household income below the poverty line (29%) have a much higher prevalence of smoking compared to people with household incomes at or above the poverty line (18%).²³⁴ Healthy eating (specifically fruit and vegetable consumption) is also lower among low-income populations compared to higher income populations.²⁴³

**PEOPLE WITH
LESS THAN A
HIGH SCHOOL
EDUCATION**

Approximately 15 % of Americans 25 years old and older have not earned a high school diploma.²⁴⁴ Those with undergraduate degrees have a lower prevalence of smoking (9%), compared to those with less than a high school education (25%) or only a high school diploma (24%).²³⁴ Additionally, those with a GED have the highest prevalence of smoking (45%). Regarding obesity, college graduates or above had the lowest rate of obesity (28%) in 2009-2010, compared to those with less than a high school education (38%).²⁴⁵

OLDER ADULTS

The proportion of our nation’s population aged 65 years and older is expected to increase from approximately 13% of the population in 2010 to an estimated 19% in 2030.²⁴⁶ In 2009–2010, 45% of adults aged 65 and over were diagnosed with two or more chronic conditions.²⁴⁷ Regarding inequities, older adults living in poverty and isolation may be particularly vulnerable.²⁴⁸

**PEOPLE WHO
IDENTIFY AS
LESBIAN, GAY,
BISEXUAL, OR
TRANSGENDER
(LGBT)**

The lesbian, gay, or bisexual population is estimated at 3.5% in the United States, with an additional 0.3% identifying as transgender.²⁴⁹ Regarding sexual orientation, use of any tobacco products have been found to be higher among lesbian, gay, bisexual, and transgender populations (38.5%) compared to the heterosexual/straight population (25.3%).⁶¹ Obesity prevalence has also been noted among the LGBT community, particularly among lesbians who have been shown to have a higher prevalence of being overweight and obese than heterosexual women who are overweight and obese.²⁵⁰

NOTE: This list is not exhaustive and the groups are not mutually exclusive; individuals may belong to more than one population group.



CONSIDERATIONS FOR HEALTH EQUITY-ORIENTED STRATEGY SELECTION, DESIGN, AND IMPLEMENTATION

Policy, systems, and environmental improvement strategies have great potential to prevent and reduce health inequities, affect a large portion of a population, and can also be leveraged to address the underlying social determinants of health. **However, without careful design and implementation, such interventions may inadvertently widen health inequities.** Collaborate with partners and community members, including those experiencing health inequities, to identify possible barriers or negative unintended consequences that may limit a strategy’s effectiveness. Then, account for identified challenges in strategy development to maximize the health effects for all and reduce health inequities. Consider the following barriers, unintended consequences, and questions when selecting, designing, and implementing equity-oriented strategies:

1

LIMITED COMMUNITY CAPACITY AND RESOURCES

Variability in community capacity and resources can influence decisions about which communities and community organizations to partner with, especially if resources are limited. While there are benefits to funding and collaborating with partners that can “hit the ground running,” it is also important to build the capacity of other groups through training and additional support.

- Has lack of capacity or resources kept critical partners away?
- What training opportunities can build the capacity of residents or organizations to make community improvements?
- Are the same organizations repeatedly benefiting from funds distributed in the community? What steps can you take to engage other organizations?

2

VARIABILITY IN HEALTH LITERACY

Addressing health literacy means ensuring that all members of the community have the capacity to access and understand the information they need to engage in health improvement strategies or reap their health benefits.

- Will the improvements be understood by all community members?
- Is training needed to support and sustain the improvements?
- How will language, culture, and other differences be accommodated?

3

LACK OF COMMUNITY ENGAGEMENT, AWARENESS, AND PARTICIPATION

A well-designed effort may fail to reach its full potential if residents are unaware of the improvements or were not invited to participate in the planning and implementation process. Community residents and stakeholders should be consulted and engaged from the very start, and this engagement should be sustained throughout the process.

- How will stakeholders representative of the community’s diversity be engaged?
- What steps will be taken to engage community members in planning, implementation, and evaluation?

4

COST, RESOURCES, AND OTHER FISCAL CONSIDERATIONS

There may be costs related to strategy implementation, either for the institutions making the improvements, or for the people who are the intended beneficiaries of these improvements. Examine how budget constraints may hinder implementation or uptake in underserved communities.

- Will costs prevent underserved populations from fully benefitting from the strategy? How can affordability be ensured for all?
- Which partners might be able to help provide required resources (e.g., funding, materials, staff, other assets) to implement the strategy?

5

TRANSPORTATION CHALLENGES

Lack of personal transportation, unaffordable or unreliable public transportation, or inadequate infrastructure may reduce access to goods, services, or environmental improvements, including tobacco cessation services and other health care services. Explore whether transportation issues such as access, cost, and proximity exist.

- Is lack of transportation a problem for the intended beneficiaries of the strategy?
- Are the locations where services are provided too distant, inconvenient, inaccessible, or unsafe?

6

POTENTIAL DISPLACEMENT EFFECTS

Changing community conditions may contribute to cycles of displacement. It is important to ensure that improvements will benefit residents rather than create conditions that displace them. Identify factors that may drive displacement and protections that can prevent it.

- How might community improvement strategies lead to displacement in the future?
- What protections can be put in place to preserve affordable housing and prevent displacement?
- How might concerns about displacement prevent residents from engaging in community improvements?

7

VARIABILITY IN IMPLEMENTATION

Uneven implementation of a policy or systems improvement may worsen inequities. Explore the factors (including those listed in this table) that might prevent consistent implementation of a strategy and develop solutions early in the planning process.

- Once your strategy is adopted or implemented, what steps will ensure proper implementation?
- How will you ensure implementation occurs where it's needed most?
- Which institutions need additional support to implement the improvements?

8

CRIME/SAFETY INFLUENCES (BOTH REAL AND PERCEIVED)

Even if effective strategies are put in place, fear of crime at locations where the intervention or service is being delivered may keep residents from using the new resources. Assess safety conditions and residents' perceptions of these conditions, and, if necessary, take steps to ensure participants' safety.

- How might concerns about safety prevent the community from benefitting from the strategy?
- Are there visible signs of crime and violence?

9

LACK OF AWARENESS OF DIVERSE NORMS AND CUSTOMS

Understanding the diversity in culture, norms, and customs among population groups can ensure that strategies are designed to be inclusive. Institutions also have their own customs and norms, and these should also be considered, as they might affect decision making.

- How will community members with different norms and customs be engaged in strategy design?
- Are differences in culture and norms understood in ways that result in respectful strategy development?



EXAMPLE RESOURCES FOR IDENTIFYING AND UNDERSTANDING HEALTH INEQUITIES

This table describes several online resources that you may be able to use to identify and understand health inequities in your area. This list is not exhaustive and you should determine what best fits your local needs.

BEHAVIORAL RISK FACTOR SURVEILLANCE SYSTEM (BRFSS)²⁵¹

A state-based system of health surveys that collects information on health risk behaviors, preventive health practices, and health care access primarily related to chronic disease and injury.

CENSUS DATA²⁵²

A database that provides demographic information on income, education, race/ethnicity, housing, and other factors that are viewable at multiple levels: national, state, county, and smaller geographic areas. Interactive features also allow cross tabulation of indicators and population groups.

COMMUNITY COMMONS²⁵³

An online interactive mapping tool that provides free geographic information systems (GIS) data from the state level to the block group level. The Commons is linked to the National Prevention Strategy and provides a peer learning network and other resources.

COMMUNITY HEALTH ASSESSMENT & GROUP EVALUATION (CHANGE): BUILDING A FOUNDATION OF KNOWLEDGE TO PRIORITIZE COMMUNITY NEEDS⁷

A tool to help community teams develop a community action plan. This tool provides steps for community team members to use in an assessment process. It also helps define and prioritize possible areas of improvement to address the root causes of chronic diseases, as well as related risk factors.

COUNTY HEALTH RANKINGS: MOBILIZING ACTION TOWARD COMMUNITY HEALTH²⁵⁴

A ranking of counties in each of the 50 states according to summaries of a variety of health measures. Summary measures include health outcomes (mortality and morbidity) and health factors (health behaviors, clinical care, social and economic aspects, and physical environment).

**COMMUNITY
HEALTH STATUS
INDICATORS
(CHSI)²⁵⁵**

A report that contains over 200 measures for each of the 3,141 U.S. counties. The report presents indicators for deaths due to heart disease and cancer as well as on behavioral factors such as tobacco use, diet, physical activity, alcohol and drug use, sexual behavior, and others that substantially contribute to these deaths.



**DATA SET
DIRECTORY
OF SOCIAL
DETERMINANTS
OF HEALTH AT THE
LOCAL LEVEL²⁵⁶**

A directory that contains an extensive list of existing data sets that can be used to address social determinants of health. The data sets are organized according to 12 dimensions (broad categories) of the social environment.

**HEALTHY
COMMUNITIES
NETWORK (HCN)²⁵⁷**

A network that tracks over 200 health and quality-of-life indicators. It also provides guidance on 1,800-plus community-level interventions. Local information is collected and combined with other data.

**HEALTH
DISPARITIES
CALCULATOR²⁵⁸**

Statistical software from the National Cancer Institute that imports population-based health data and calculates different disparity measurements.

**HEALTH EQUITY
INDEX²⁵⁹**

An online tool created by the Connecticut Association of Directors of Health that outlines and measures the social determinants of health with specific health outcomes. The index produces scores as well as GIS maps.

**HEALTH
INDICATORS
WAREHOUSE²⁶⁰**

A Web site maintained by CDC's National Center for Health Statistics. This resource provides data on communities' health status as well as different determinants. There are over 1,000 indicators that can be categorized by geography, initiative, or topic.

**THE TOOL FOR
HEALTH AND
RESILIENCE IN
VULNERABLE
ENVIRONMENTS
(THRIVE)²⁶¹**

A tool intended to help people understand and prioritize the factors within their own communities in order to improve health and safety. The tool identifies key factors around equitable opportunity, people, and place, and allows users to rate how important each factor might be in their community.





HEALTH EQUITY CHECKLIST: CONSIDERING HEALTH EQUITY IN THE STRATEGY DEVELOPMENT PROCESS

The Health Equity Checklist provides questions for consideration when designing a strategy to ensure health equity remains central to all aspects of an initiative.

STEP 1: IDENTIFY

Clearly identify health inequities and protective factors in both health outcomes and community conditions across population groups and geographic areas through the use of existing data, community input, and environmental assessments.

STEP 2: ENGAGE

Include and meaningfully engage representatives of population(s)/area(s) defined in Step 1 in your partnerships, coalitions, or on leadership teams.

STEP 3: ANALYZE

Ensure the selection, design, and implementation of strategies are linked to the inequities identified in Step 1, and will work to advance health equity. Consider the following:

-  Is the strategy TARGETED to a population group(s)/area(s) experiencing health inequities?
 - Is the outcome written in a way that allows you to measure the effect of efforts?
 - Is it culturally tailored to the unique needs of population group(s)/area(s) experiencing health inequities, and are potential barriers addressed?

-  Does the strategy rely on SITE SELECTION (e.g., selecting X number of sites for smoke-free cessation services, creating X number of farmers' markets)?
 - Do selection criteria for sites reflect populations/areas with the highest burden?
 - If not, are selection criteria logical and justified?
 - Are there additional supports provided for selected sites that might require them to be successful?

-  Is the strategy POPULATION-WIDE?
 - Have population(s)/area(s) experiencing health inequities been engaged in efforts to identify possible barriers and unintended consequences of the proposed strategy?
 - Are identified barriers regarding implementation and enforcement being addressed?
 - Have potential unintended consequences been considered and accounted for in proposed activities?

STEP 4: REVIEW

Review evaluation and monitoring plans to ensure health equity-related efforts will be measured. Additionally, ensure appropriate data will be collected to conduct sub-analyses. These data will help in assessing the differential effects of each strategy across population group(s)/area(s), as well as the overall impact of strategies on reducing health inequities.

REFERENCES

1. Centers for Disease Control and Prevention. Anti-lobbying restrictions for CDC grantees. <http://www.cdc.gov/obesity/downloads/Anti-Lobbying-Restrictions-for-CDC-Grantees-July2012-508.pdf>. Accessed May 22, 2013.
2. Centers for Disease Control and Prevention. Additional requirement #12. http://www.cdc.gov/od/pgo/funding/grants/additional_req.shtm#ar12. Accessed May 22, 2013.
3. Braveman PA, Kumanyika S, Fielding J, et al. Health disparities and health equity: the issue is justice. *Am J Public Health*. 2011;101(suppl 1):S149-S155.
4. Centers for Disease Control and Prevention. Chronic diseases and health promotion. <http://www.cdc.gov/chronicdisease/overview/index.htm>. Accessed June 8, 2012.
5. Smedley BD, Stith AY, Nelson AR. *Unequal Treatment: Confronting Racial and Ethnic Disparities in Healthcare*. Washington, DC: National Academies Press;2003.
6. Williams DR. Race, socioeconomic status, and health: the added effects of racism and discrimination. *Ann N Y Acad Sci*. 1999;896(1):173-188.
7. Centers for Disease Control and Prevention. CDC health disparities and inequalities report - United States, 2011. *MMWR Morb Mortal Wkly Rep*. 2011;60(Supplement):1-113.
8. National Center for Health Statistics. *Healthy People 2010 Final Review*. Hyattsville, MD: National Center for Health Statistics;2012.
9. Keppel KG, Percy JN, Heron MP. Is there progress toward eliminating racial/ethnic disparities in the leading causes of death? *Public Health Rep*. 2010;125(5):689-697.
10. Orsi JM, Margellos-Anast H, Whitman S. Black-white health disparities in the United States and Chicago: a 15-year progress analysis. *Am J Public Health*. 2010;100(2):349-356.
11. National Center for Health Statistics. *Health, United States, 2011: With Special Feature on Socioeconomic Status and Health*. Hyattsville, MD: National Center for Health Statistics;2012.
12. Healthypeople.gov. Determinants of health. <http://www.healthypeople.gov/2020/about/DOHAbout.aspx#socialfactors>. Accessed February 26, 2013.
13. US Department of Health and Human Services. Framework: the vision, mission, and goals of Healthy People 2020. <http://www.healthypeople.gov/2020/Consortium/HP2020Framework.pdf>. Accessed February 26, 2013.
14. Centers for Disease Control and Prevention. Ten great public health achievements - United States, 1900-1999. *MMWR Morb Mortal Wkly Rep*. 1999;48(12):241-243.
15. Centers for Disease Control and Prevention. Ten great public health achievements - United States, 2001-2010. *MMWR Morb Mortal Wkly Rep*. 2011;60(19):619-623.
16. Centers for Disease Control and Prevention. State, tribal, local, and territorial public health professionals gateway: policy at CDC. <http://www.cdc.gov/stltpublichealth/Policy/index.html>. Accessed June 8, 2012.
17. Centers for Disease Control and Prevention. Community Health Assessment and Group Evaluation (CHANGE) action guide: building a foundation of knowledge to prioritize community needs. <http://www.cdc.gov/healthycommunitiesprogram/tools/change.htm>. Accessed May 22, 2013.
18. Braveman P. Health disparities and health equity: concepts and measurement. *Annu Rev Public Health*. 2006;27:167-194.
19. Braveman P, Gruskin S. Defining equity in health. *J Epidemiol Community Health*. 2003;57(4):254-258.
20. US Department of Health and Human Services. Social determinants of health. <http://www.healthypeople.gov/2020/topicsobjectives2020/overview.aspx?topicid=39>. Accessed July 19, 2013.
21. Alameda County Public Health Department. Organizational transformation. <http://www.acphd.org/social-and-health-equity/organizational-transformation.aspx>. Accessed June 11, 2012.

REFERENCES (Continued)

22. Bay Area Regional Health Inequities Initiative. Local health department organizational self-assessment for addressing health inequities: toolkit and guide to implementation. <http://www.barhii.org/resources/toolkit.html>. Accessed August 21, 2012.
23. Mohan M. Personal Communication with Lark Galloway-Gilliam. In: Prevention Institute, ed2012.
24. Satcher D. Include a social determinants of health approach to reduce health inequities. *Public Health Rep.* 2010;125(Supplement 4):6-7.
25. Centers for Disease Control and Prevention. *Health Equity Checklist: Considering Health Equity in the Strategy Development Process*. Atlanta, GA: US Dept of Health and Human Services;2010.
26. Centers for Disease Control and Prevention. Health impact assessment. <http://www.cdc.gov/healthyplaces/hia.htm>. Accessed June 01, 2012.
27. Haber R. Health equity impact assessment: a primer. http://www.threesource.ca/documents/March2011/health_equity.pdf. Accessed May 22, 2013.
28. Mapping Our Voices for Equality (MOVE). Map. <http://www.mappingvoices.org/>. Accessed May 22, 2013.
29. Potvin L, Mantoura P, Ridde V. Evaluating equity in health promotion. In: McQueen D, Jones C, eds. *Global Perspectives on Health Promotion Effectiveness*. Atlanta, GA: Springer Science & Business Media; 2007:367-384.
30. Centers for Disease Control and Prevention. Cigarette smoking - United States 1965-2008. *MMWR Morb Mortal Wkly Rep.* 2011;60(Suppl):109-113.
31. Centers for Disease Control and Prevention. *Cigarette Smoking and People with Disabilities: A Tip Sheet for Public Health Professionals*. Atlanta, GA: National Center on Birth Defects and Development Disabilities;2009.
32. Centers for Disease Control and Prevention. *Best Practices for Comprehensive Tobacco Control Programs-2007*. Atlanta, GA: US Dept of Health and Human Services, Centers for Disease Control and Prevention, National Center for Chronic Disease Prevention and Health Promotion, Office of Smoking and Health;2007.
33. Centers for Disease Control and Prevention. State smoke-free laws for worksites, restaurants, and bars--United States, 2000-2010. *MMWR Morb Mortal Wkly Rep.* 2011;60(15):472-475.
34. American Nonsmokers' Rights Foundation. State, commonwealths, and municipalities with 100% smokefree laws in non-hospitality workplaces, restaurants, or bars. <http://www.no-smoke.org/pdf/100ordlist.pdf>. Accessed August 22, 2012.
35. US Department of Health and Human Services. *The Health Consequences of Involuntary Exposure to Tobacco Smoke: A Report of the Surgeon General*. Atlanta, GA: US Dept of Health and Human Services, Centers for Disease Control and Prevention, Coordinating Center for Health Promotion, National Center for Chronic Disease Prevention and Health Promotion, Office on Smoking and Health;2006.
36. Berman M, Post C, Tobacco Control Legal Consortium. Secondhand smoke and casinos. http://publichealthlawcenter.org/sites/default/files/resources/tclc-syn-casinos-2007_0.pdf. Accessed May 2, 2013.
37. US Bureau of Labor Statistics. *Labor Force Characteristics by Race and Ethnicity, 2010*. Washington, D.C.: US Dept of Labor and US Bureau of Labor Statistics;2011.
38. Osypuk TL, Subramanian SV, Kawachi I, Acevedo-Garcia D. Is workplace smoking policy equally prevalent and equally effective among immigrants? *J Epidemiol Community Health.* 2009;63:784-791.
39. Gerlach KK, Shopland DR, Hartman AM, Gibson JT, Pechacek TF. Workplace smoking policies in the United States: results from a national survey of more than 100,000 workers. *Tob Control.* 1997;6(3):199-206.

REFERENCES (Continued)

40. Moore RS, Annechino RM, Lee JP. Unintended consequences of smoke-free bar policies for low-SES women in three California counties. *Am J Prev Med*. 2009;37(suppl 2):S138-S143.
41. Substance Abuse and Mental Health Services Administration. *Results from the 2008 National Survey on Drug Use and Health: National Findings*. Rockville, MD: Substance Abuse and Mental Health Services Administration - Office of Applied Studies;2009.
42. Stevens S, Colwell B, Hutchison L. Tobacco use in rural areas. In: Gamm L, Hutchison L, Dabney B, Dorsey A, eds. *Rural Healthy People 2010: A Companion Document to Healthy People 2010*. Vol 2. College Station, TX: The Texas A&M University System Health Science Center, School of Rural Public Health, Southwest Rural Health Research Center; 2003.
43. Wood LE. *The Economic Impact of Tobacco Production in Appalachia*. Washington, DC: Appalachian Regional Commission; 1998.
44. Báezconde-Garbanati L, Beebe LA, Pérez-Stable EJ. Building capacity to address tobacco-related disparities among American Indian and Hispanic/Latino communities: conceptual and systemic considerations. *Addiction*. 2007;102(suppl 2):112-122.
45. Yerger VB, Malone RE. African American leadership groups: smoking with the enemy. *Tob Control*. 2002;11(4):336-345.
46. Portugal C, Cruz TB, Espinoza L, Romero M, Baezconde-Garbanati L. Countering tobacco industry sponsorship of Hispanic/Latino organizations through policy adoption: a case study. *Health Promot Pract*. 2004;5(Suppl 3):143S-156S.
47. Raebeck A, Campbell R, Balbach E. Unhealthy partnerships: the tobacco industry and African American and Latino labor organizations. *J Immigr Minor Health*. 2010;12(2):228-233.
48. Agency for Healthcare Research and Quality. *National Healthcare Disparities Report 2011*. Rockville, MD: Agency for Healthcare Research and Quality;2012.
49. Fagan P, Moolchan ET, Lawrence D, Fernander A, Ponder PK. Identifying health disparities across the tobacco continuum. *Addiction*. 2007;102(suppl 2):5-29.
50. Moolchan ET, Fagan P, Fernander AF, et al. Addressing tobacco-related health disparities. *Addiction*. 2007;102(suppl 2):30-42.
51. Centers for Disease Control and Prevention. Smoking & tobacco use: Montana. http://www.cdc.gov/tobacco/data_statistics/state_data/state_highlights/2010/states/montana/index.htm. Accessed July 17, 2013.
52. Montana Tobacco Prevention Advisory Board. Montana tobacco use prevention plan. http://s3.amazonaws.com/zanran_storage/tobaccofree.mt.gov/ContentPages/43378690.pdf. Accessed July 17, 2013.
53. US Department of Housing and Urban Development. Resident Characteristics Report: public housing. <https://pic.hud.gov/pic/RCRPublic/rcrmain.asp>. Accessed January 16, 2013.
54. Kraev TA, Adamkiewicz G, Hammond SK, Spengler JD. Indoor concentrations of nicotine in low-income, multi-unit housing: associations with smoking behaviours and housing characteristics. *Tob Control*. 2009;18(6):438-444.
55. DiFranza JR, Aligne CA, Weitzman M. Prenatal and postnatal environmental tobacco smoke exposure and children's health. *Pediatrics*. 2004;113(suppl 3):1007-1015.
56. King BA, Travers MJ, Cummings KM, Mahoney MC, Hyland AJ. Secondhand smoke transfer in multiunit housing. *Nicotine Tob Res*. 2010;12(11):1133-1141.
57. Helburn A. A case for smoke free housing. 2007. http://www.hria.org/uploads/reports/HRIA-Smoke_Free_Housing_2007.pdf. Accessed April 19, 2013.
58. Fiore MC, Jaen CR, Baker TB, et al. *Treating Tobacco Use and Dependence: 2008 Update*. Rockville, MD: US Dept of Health and Human Services - Public Health Service;2008.

REFERENCES (Continued)

59. Centers for Disease Control and Prevention. Quitting smoking among adults --- United States, 2001--2010. *MMWR Morb Mortal Wkly Rep.* 2011;60(44):1513-1519.
60. Centers for Disease Control and Prevention. Vital signs: current cigarette smoking among adults aged ≥ 18 years --- United States, 2005--2010. *MMWR Morb Mortal Wkly Rep.* 2011;60(35):1207-1212.
61. King BA, Dube SR, Tynan MA. Current tobacco use among adults in the United States: findings from the National Adult Tobacco Survey. *Am J Public Health.* 2012;102(11):e93-e100.
62. Lee JGL, Griffin GK, Melvin CL. Tobacco use among sexual minorities in the USA, 1987 to May 2007: a systematic review. *Tob Control.* 2009;18(4):275-282.
63. Ryan H, Wortley PM, Easton A, Pederson L, Greenwood G. Smoking among lesbians, gays, and bisexuals: a review of the literature. *Am J Prev Med.* 2001;21(2):142-149.
64. Lasser K, Boyd JW, Woolhandler S, Himmelstein DU, McCormick D, Bor DH. Smoking and mental illness: a population-based prevalence study. *JAMA.* 2000;284(20):2606-2610.
65. Armour BS, Campbell VA, Crews JE, Malarcher A, Maurice E, Richard RA. State-level prevalence of cigarette smoking and treatment advice, by disability status, United States, 2004. *Prev Chronic Dis.* 2007;4(4):1-11.
66. Kleykamp BA, Heishman SJ. The older smoker. *JAMA.* 2011;306(8):876-877.
67. Barbeau EM, Krieger N, Soobader M. Working class matters: socioeconomic disadvantage, race/ethnicity, gender, and smoking in NHIS 2000. *Am J Public Health.* 2004;94(2):269-278.
68. Levinson AH, Pérez-Stable EJ, Espinoza P, Flores ET, Byers TE. Latinos report less use of pharmaceutical aids when trying to quit smoking. *Am J Prev Med.* 2004;26(2):105-111.
69. Cokkinides VE, Halpern MT, Barbeau EM, Ward E, Thun MJ. Racial and ethnic disparities in smoking-cessation interventions: analysis of the 2005 National Health Interview Survey. *Am J Prev Med.* 2008;34(5):404-412.
70. Sonnenfeld N, Schappert SM, Lin X. Racial and ethnic differences in delivery of tobacco-cessation services. *Am J Prev Med.* 2009;36(1):21-28.
71. Doolan DM, Froelicher ES. Efficacy of smoking cessation intervention among special populations: review of the literature from 2000 to 2005. *Nurs Res.* 2006;55 (suppl 4):S29-S37.
72. Fu SS, Burgess DJ, Hatsukami DK, et al. Race and nicotine replacement treatment outcomes among low-income smokers. *Am J Prev Med.* 2008;35(6):S442-S448.
73. Fu SS, Burgess D, van Ryn M, Hatsukami DK, Solomon J, Joseph AM. Views on smoking cessation methods in ethnic minority communities: a qualitative investigation. *Prev Med.* 2007;44:235-240.
74. Cox LS, Okuyemi K, Choi WS, Ahluwalia JS. A review of tobacco use treatments in US ethnic minority populations. *Am J Health Promot.* 2011;25(suppl 5):11-30.
75. Bandi P, Cokkinides VE, Virgo KS, Ward EM. The receipt and utilization of effective clinical smoking cessation services in subgroups of the insured and uninsured populations in the USA. *J Behav Health Serv Res.* 2012;39(2):202-213.
76. Centers for Disease Control and Prevention. State medicaid coverage for tobacco-dependence treatments- United States, 2007. *MMWR Morb Mortal Wkly Rep.* 2009;58(43):1199-1204.
77. Blumenthal DS. Barriers to the provision of smoking cessation services reported by clinicians in underserved communities. *J Am Board Fam Med.* 2007;20(3):272-279.
78. Dickerson DL, Leeman RF, Mazure CM, O'Malley S. The inclusion of women and minorities in smoking cessation clinical trials: a systematic review. *Am J Addict.* 2009;18(1):21-28.

REFERENCES (Continued)

79. US Department of Health and Human Services. *Preventing Tobacco Use Among Youth and Young Adults: A Report of the Surgeon General*. Atlanta, GA: US Department of Health and Human Services, Centers for Disease Control and Prevention, National Center for Chronic Disease Prevention and Health Promotion, Office on Smoking and Health;2012.
80. 111th Congress. Division A - Family Smoking Prevention and Tobacco Control Act Public Law 111-31. <http://www.fda.gov/TobaccoProducts/GuidanceComplianceRegulatoryInformation/ucm261829.htm>. Accessed June 12, 2012.
81. Institute of Medicine. *Ending the Tobacco Problem: A Blueprint for the Nation*. Washington, DC: The National Academies Press; 2007.
82. Siahpush M, Jones PR, Singh GK, Timsina LR, Martin J. The association of tobacco marketing with median income and racial/ethnic characteristics of neighbourhoods in Omaha, Nebraska. *Tob Control*. 2010;19(3):256.
83. John R, Cheney MK, Azad MR. Point-of-sale marketing of tobacco products: taking advantage of the socially disadvantaged? *J Health Care Poor Underserved*. 2009;20(2):489-506.
84. Laws MB, Whitman J, Bowser DM, Krech L. Tobacco availability and point of sale marketing in demographically contrasting districts of Massachusetts. *Tob Control*. 2002;11(suppl 2):ii71-ii73.
85. Sutton CD, Robinson RG. The marketing of menthol cigarettes in the United States: populations, messages, and channels. *Nicotine Tob Res*. 2004;6 (suppl 1):S83-S91.
86. Feighery EC, Ribisl KM, Clark PI, Haladjian HH. How tobacco companies ensure prime placement of their advertising and products in stores: interviews with retailers about tobacco company incentive programmes. *Tob Control*. 2003;12(2):184-188.
87. Federal Trade Commission. *Federal Trade Commission Cigarette Report for 2011*. Washington, DC: Federal Trade Commission;2013.
88. Feighery EC, Ribisl KM, Schleicher N, Lee RE, Halvorson S. Cigarette advertising and promotional strategies in retail outlets: results of a statewide survey in California. *Tob Control*. 2001;10(2):184-188.
89. Seidenberg AB, Caughey RW, Rees VW, Connolly GN. Storefront cigarette advertising differs by community demographic profile. *Am J Health Promot*. 2010;24(6):e26-e31.
90. Farrelly MC, Nonnemaker JM, Watson KA. The consequences of high cigarette excise taxes for low-income smokers. *PLoS ONE*. 2012;7(9):e43838.
91. Hyland A, Travers MJ, Cummings KM, Bauer J, Alford T, Wieczorek WF. Tobacco outlet density and demographics in Erie County, New York. *Am J Public Health*. 2003;93(7):1075-1076.
92. Schneider JE, Reid RJ, Peterson NA, Lowe JB, Hughey J. Tobacco outlet density and demographics at the tract level of analysis in Iowa: implications for environmentally based prevention initiatives. *Prev Sci*. 2005;6(4):319-325.
93. Siahpush M, Jones PR, Singh GK, Timsina LR, Martin J. Association of availability of tobacco products with socio-economic and racial/ethnic characteristics of neighbourhoods. *Public Health*. 2010;124(9):525-529.
94. Peterson NA, Yu D, Morton CM, Reid RJ, Sheffer MA, Schneider JE. Tobacco outlet density and demographics at the tract level of analysis in New Jersey: a statewide analysis. *Drugs Educ Prev Policy*. 2011;18(1):47-52.
95. PolicyLink, The Food Trust, The Reinvestment Fund. A healthy food financing initiative: an innovative approach to improve health and spark economic development. http://www.policylink.org/atf/cf/%7B97c6d565-bb43-406d-a6d5-eca3bbf35af0%7D/HFFI_ADVOCACY3.PDF. Accessed May 22, 2013.
96. The Food Trust. What we do: in corner stores. <http://thefoodtrust.org/what-we-do/corner-store>. Accessed May 22, 2013.

REFERENCES (Continued)

97. National Cancer Institute. Influence of tobacco marketing on smoking behavior. Tobacco Control Monograph No. 19. In: Davis RM, Gilpin EA, Loken B, Viswanath K, Wakefield MA, eds. *The Role of the Media in Promoting and Reducing Tobacco Use*. Bethesda, MD: National Cancer Institute, US Dept of Health and Human Services, National Institutes of Health; 2008:211-291.
98. Henriksen L, Schleicher NC, Feighery EC, Fortmann SP. A longitudinal study of exposure to retail cigarette advertising and smoking initiation. *Pediatrics*. 2010;126(2):232-238.
99. Carter OBJ, Mills BW, Donovan RJ. The effect of retail cigarette pack displays on unplanned purchases: results from immediate postpurchase interviews. *Tob Control*. 2009;18(3):218-221.
100. Treuhaft S, Karpyn A. The grocery gap: who has access to healthy food and why it matters. http://www.policylink.org/site/c.lkIXLbMNJrE/b.5860321/k.A5BD/The_Grocery_Gap.htm. Accessed August 22, 2012.
101. Larson NI, Story MT, Nelson MC. Neighborhood environments: disparities in access to healthy foods in the US. *Am J Prev Med*. 2009;36(1):74-81.
102. Moore LV, Diez Roux AV. Associations of neighborhood characteristics with the location and type of food stores. *Am J Public Health*. 2006;96(2):325-331.
103. Powell LM, Slater S, Mirtcheva D, Bao Y, Chaloupka FJ. Food store availability and neighborhood characteristics in the United States. *Prev Med*. 2007;44(3):189-195.
104. New Mexico Food and Agriculture Policy Council. Closing New Mexico's rural food gap. http://farmtotablenm.org/wp-content/uploads/2013/03/closing_nm_food_gap_4pgs.pdf. Accessed September 9, 2013.
105. Jetter KM, Cassady DL. The availability and cost of healthier food alternatives. *Am J Prev Med*. 2006;30(1):38-44.
106. Liese AD, Weis KE, Pluto D, Smith E, Lawson A. Food store types, availability, and cost of foods in a rural environment. *J Am Diet Assoc*. 2007;107(11):1916-1923.
107. Andreyeva T, Blumenthal DM, Schwartz MB, Long MW, Brownell KD. Availability and prices of foods across stores and neighborhoods: the case of New Haven, Connecticut. *Health Aff (Millwood)*. 2008;27(5):1381-1388.
108. Zenk SN, Schulz AJ, Israel BA, James SA, Bao S, Wilson ML. Fruit and vegetable access differs by community racial composition and socioeconomic position in Detroit, Michigan. *Ethn Dis*. 2006;16(1):275-280.
109. Fair Food Network. Double Up Food Bucks program. 2011; <http://www.doubleupfoodbucks.org/about>. Accessed December 7, 2011.
110. Walker RE, Keane CR, Burke JG. Disparities and access to healthy food in the United States: a review of food deserts literature. *Health Place*. 2010;16(5):876-884.
111. Hendrickson D, Smith C, Eikenberry N. Fruit and vegetable access in four low-income food deserts communities in Minnesota. *Agric and Human Values*. 2006;23(3):371-383.
112. Flournoy R, PolicyLink. Healthy food, healthy communities: promising strategies to improve access to fresh, healthy food and transform communities. http://www.policylink.org/atf/cf/%7B97c6d565-bb43-406d-a6d5-eca3bbf35af0%7D/HFHC_SHORT_FINAL.PDF. Accessed May 22, 2013.
113. ChangeLab Solutions. Healthy corner stores: the state of the movement. <http://changelabsolutions.org/sites/default/files/documents/HCSRreport.pdf>. Accessed May 22, 2013.
114. ChangeLab Solutions. Getting to grocery: tools for attracting healthy food retail to underserved neighborhoods. <http://changelabsolutions.org/publications/getting-grocery>. Accessed August 23, 2012.

REFERENCES (Continued)

115. The Food Trust. Building healthy communities: expanding access to fresh food retail. http://prc.tulane.edu/uploads/REPORT_FINAL-1290013526.pdf. Accessed August 23, 2012.
116. Poti JM, Popkin BM. Trends in energy intake among US children by eating location and food source, 1977-2006. *J Am Diet Assoc*. 2011;111(8):1156-1164.
117. Nielsen SJ, Siega-Riz AM, Popkin BM. Trends in energy intake in US between 1977 and 1996: similar shifts seen across age groups. *Obes Res*. 2002;10(5):370-378.
118. Block JP, Scribner RA, DeSalvo KB. Fast food, race/ethnicity, and income: a geographic analysis. *Am J Prev Med*. 2004;27(3):211-217.
119. Lewis LB, Sloane DC, Nascimento LM, et al. African Americans' access to healthy food options in South Los Angeles restaurants. *Am J Public Health*. Apr 2005;95(4):668-673.
120. Powell LM, Chaloupka FJ, Bao Y. The availability of fast-food and full-service restaurants in the United States: associations with neighborhood characteristics. *Am J Prev Med*. 2007;33(4):S240-S245.
121. Larson N, Story M, Nelson MC. Restaurant realities: inequalities in access to healthy restaurant choices. [http://www.healthyeatingresearch.org/images/stories/her_research_briefs/her_research_briefs/her%20restaurant%20realities_7-2008.pdf](http://www.healthyeatingresearch.org/images/stories/her_research_briefs/her%20restaurant%20realities_7-2008.pdf). Accessed January 16, 2013.
122. Creel JS, Sharkey JR, McIntosh A, Anding J, Huber Jr JC. Availability of healthier options in traditional and nontraditional rural fast-food outlets. *BMC Public Health*. 2008;8(1):395.
123. Cheyne A, Phil C, Dorfman L, Gonzalez P, Mejia P. Food and beverage marketing to children and adolescents: an environment at odds with good health. http://www.healthyeatingresearch.org/images/stories/her_research_briefs/HER_RS_FoodMarketing_FINAL_4-6-11.pdf. Accessed June 12, 2012.
124. Britt JW, Frandsen K, Leng K, Evans D, Pulos E. Feasibility of voluntary menu labeling among locally owned restaurants. *Health Promot Pract*. 2011;12(1):18-24.
125. Elbel B, Kersh R, Brescoll VL, Dixon LB. Calorie labeling and food choices: a first look at the effects on low-income people in New York City. *Health Aff (Millwood)*. 2009;28(6):w1110-w1121.
126. Mayor's Healthy Hometown Movement Food in Neighborhoods Committee. The state of food: a snapshot of food access in Louisville. <http://www.louisvilleky.gov/NR/rdonlyres/E8C0D055-E234-489D-A592-7792E323D106/0/StateofFoodFINAL.pdf>. Accessed July 19, 2013.
127. Gittelsohn J, Sharma S. Physical, consumer, and social aspects of measuring the food environment among diverse low-income populations. *Am J Prev Med*. Apr 2009;36(4 Suppl):S161-165.
128. Gordon A, Fox MK, Clark M, et al. *School Nutrition Dietary Assessment Study-III: Summary of Findings*. Washington, DC: US Department of Agriculture;2007.
129. Gleason P, Suitor C. *Children's Diets in the Mid-1990s: Dietary Intake and Its Relationship with School Meal Participation*. Alexandria, VA: US Dept of Agriculture, Food and Nutrition Service;2001.
130. Khan S, Pinckney RG, Keeney D, Frankowski B, Carney JK. Prevalence of food insecurity and utilization of food assistance program: an exploratory survey of a Vermont middle school. *J Sch Health*. 2011;81(1):15-20.
131. Story M, Kaphingst KM, French S. The role of child care settings in obesity prevention. *Future Child*. 2006;16(1):143-168.
132. US Department of Agriculture Food and Nutrition Service. *National School Lunch Program Fact Sheet*. Washington, DC: US Dept of Agriculture;2012.
133. US Department of Agriculture Food and Nutrition Services. *The School Breakfast Program - Fact Sheet*. Washington, DC: US Dept of Agriculture;2013.
134. US Department of Agriculture Food and Nutrition Services. Child & adult care food program. <http://www.fns.usda.gov/cnd/care/CACFP/aboutcacfp.htm>. Accessed May 9, 2013.

REFERENCES (Continued)

135. US Department of Agriculture - Food and Nutrition Services Child Nutrition Programs. *Eligibility Manual for School Meals: Determining and Verifying Eligibility*. Washington, DC: US Dept of Agriculture;2012.
136. White House Task Force on Childhood Obesity. *Solving the Problem of Childhood Obesity Within a Generation - Section III*. Washington, DC: Executive Office of the President of the United States;2010.
137. National Council of La Raza. Profiles of Latino Health Series 3 - Issue 10: Hispanic participation in school-based nutrition programs. http://www.nclr.org/images/uploads/pages/Jan12_Profiles_Issue_10.pdf. Accessed June 10, 2012.
138. Stein K. Erasing the stigma of subsidized school meals. *J Am Diet Assoc*. 2008;108(12):1980-1983.
139. Mirtcheva DM, Powell LM. Participation in the National School Lunch Program: importance of school level and neighborhood contextual factors. *J Sch Health*. 2009;79(10):485-494.
140. United States Government Accountability Office. *School Meal Programs: Competitive Foods are Widely Available and Generate Substantial Revenue for Schools*. Washington, DC: Government Accountability Office;2005. GAO-05-563.
141. Story M, Kaphingst KM, French S. The role of schools in obesity prevention. *Future Child*. 2006;16(1):109-142.
142. Southern Nevada Health District. Community obesity survey: executive summary. <http://www.southernnevadahealthdistrict.org/download/white-papers/obesity-executive-report.pdf>. Accessed July 19, 2013.
143. Haboush A, Davidson D, Phebus T, Lopez E, Pitts C, Nevada Institute for Children's Research and Policy. Health status of children entering kindergarten in Nevada. http://nic.unlv.edu/files/KHS%20Year%205%20Report_514.13_FinalRevised.pdf. Accessed July 19, 2013.
144. Bibb County School District. Bibb County school nutrition program. <http://schools.bibb.k12.ga.us/Page/399>. Accessed April 6, 2012.
145. Arnold CA. *Fair and Healthy Land Use: Environmental Justice and Planning*. Chicago, IL: American Planning Association; 2007.
146. Ellickson R, Been V. *Land-Use Controls: Cases and Materials*. 3rd ed. New York, NY: Aspen Publishers; 2005.
147. Collin RW. Environmental equity: a law and planning approach to environmental racism. *Environ Law J*. 1992;11(495):496-546.
148. Haar CM, Fessler DW. *The Wrong Side of the Tracks: A Revolutionary Rediscovery of the Common Law Tradition of Fairness in the Struggle Against Inequality*. New York, NY: Simon and Schuster; 1986.
149. Bond KW. Toward equal delivery of municipal services in the central cities. *Fordham Urban Law J*. 1975;4(2):263-287.
150. Garcia R, Flores ES. Anatomy of the urban parks movement: equal justice, democracy, and livability in Los Angeles. In: Bullard RD, ed. *The Quest for Environmental Justice: Human Rights and the Politics of Pollution*. San Francisco, CA: Sierra Club Books; 2005:145-167.
151. Rossen LM, Pollack KM. Making the connection between zoning and health disparities. *Environ Justice*. 2012;5(3):119-127.
152. US Census Bureau. 2010 American Community Survey. <http://www.census.gov/acs/www/>. Accessed June 11, 2012.
153. Neuner K, Raja S. Healthy eating and active living: for children in the city of Buffalo. http://foodsystmsplanning.ap.buffalo.edu/wp-content/uploads/2012/08/HKHC-Policy-Brief-1_whyhealthyliving.pdf. Accessed July 19, 2013.

REFERENCES (Continued)

154. Bassford N, Galloway-Gilliam L, Flynn G, CHC Food Resource Development Workgroup. Food desert to food oasis: promoting grocery store development in South Los Angeles. http://chc-inc.org/downloads/PB_Food_Desert_2010.pdf. Accessed July 1, 2013.
155. Los Angeles County Department of Public Health. Life expectancy in Los Angeles County: how long do we live and why?. http://www.publichealth.lacounty.gov/epi/docs/Life%20Expectancy%20Final_web.pdf. Accessed July 1, 2013.
156. Baby-Friendly USA. Baby-friendly hospital initiative. <http://www.babyfriendlyusa.org/about-us/baby-friendly-hospital-initiative>. Accessed May 22, 2013.
157. US Department of Health and Human Services. *The Surgeon General's Call to Action to Support Breastfeeding*. Washington, DC: US Dept of Health and Human Services, Office of the Surgeon General;2011.
158. Grummer-Strawn LM, Shealy KR. Progress in protecting, promoting, and supporting breastfeeding: 1984-2009. *Breastfeed Med*. 2009;4 (suppl 1):S31-S39.
159. Sparks PJ. Rural-urban differences in breastfeeding initiation in the United States. *J Hum Lact*. 2010;26(2):118-129.
160. Centers for Disease Control and Prevention. Breastfeeding trends and updated national health objectives for exclusive breastfeeding—United States, birth years 2000–2004. *MMWR Morb Mortal Wkly Rep*. 2007;56:760-763.
161. Ringel-Kulka T, Jensen E, McLaurin S, et al. Community-based participatory research of breastfeeding disparities in African American women. *Infant Child Adolesc Nutr*. 2011;3(4):233-239.
162. Flower KB, Willoughby M, Cadigan RJ, Perrin EM, Randolph G. Understanding breastfeeding initiation and continuation in rural communities: a combined qualitative/quantitative approach. *Matern Child Health J*. 2008;12(3):402-414.
163. Brown CA, Poag S, Kasprzycki C. Exploring large employers' and small employers' knowledge, attitudes, and practices on breastfeeding support in the workplace. *J Hum Lact*. 2001;17(1):39-46.
164. Gill SL, Reifsnider E, Mann AR, Villarreal P, Tinkle MB. Assessing infant breastfeeding beliefs among low-income Mexican Americans. *J Perinat Educ*. 2004;13(3):39-50.
165. American Academy of Family Physicians. Breastfeeding, family physicians supporting. <http://www.aafp.org/online/en/home/policy/policies/b/breastfeedingpositionpaper.html>. Accessed February 25, 2013.
166. United States Department of Agriculture - Food and Nutrition Service. Women, infants, and children (WIC). <http://www.fns.usda.gov/wic/>. Accessed May 22, 2013.
167. California WIC Association and UC Davis Human Lactation Center. *Increasing Exclusive Breastfeeding in WIC: The Power of Peer Counseling*. Sacramento, CA: CA WIC Association and UC Davis Human Lactation Center;2009.
168. California WIC Association and the UC Davis Human Lactation Center. Collaboration counts: improving hospital breastfeeding policies - California fact sheet: 2011 data. http://calwic.org/storage/restricted/hospitalfactsheetsdata2011/statefactsheet2012_corrected.pdf. Accessed July 19, 2013.
169. California Department of Public Health. Birth & Beyond California: hospital breastfeeding quality improvement and staff training demonstration project. <http://www.cdph.ca.gov/HealthInfo/healthyliving/childfamily/Pages/BirthandBeyondCaliforniaDescription.aspx>. Accessed May 22, 2013.
170. New York City Department of Health and Mental Hygiene. Breastfeeding in New York City hospitals, 2009. <http://www.nyc.gov/html/doh/downloads/pdf/ms/ms-breastfeeding-nyc.pdf>. Accessed July 19, 2013.
171. Kaufman L, Deenadayalan S, Karpati A. Breastfeeding ambivalence among low-income African American and Puerto Rican women in north and central Brooklyn. *Matern Child Health J*. 2010;14(5):696-704.

REFERENCES (Continued)

172. Powell LM, Slater S, Chaloupka FJ, Harper D. Availability of physical activity-related facilities and neighborhood demographic and socioeconomic characteristics: a national study. *Am J Public Health*. 2006;96(9):1676-1680.
173. Centers for Disease Control and Prevention. Youth violence: national statistics - trends in homicide rates among persons ages 10-24 years, by race/ethnicity. http://www.cdc.gov/ViolencePrevention/youthviolence/stats_at-a_glance/hr_trends_race.html. Accessed February 1, 2012.
174. Kelly CM, Schootman M, Baker EA, Barnidge EK, Lemes A. The association of sidewalk walkability and physical disorder with area-level race and poverty. *J Epidemiol Community Health*. 2007;61(11):978-983.
175. Zhu X, Lee C. Walkability and safety around elementary schools: economic and ethnic disparities. *Am J Prev Med*. 2008;34(4):282-290.
176. Powell LM, Slater S, Chaloupka FJ. The relationship between community physical activity settings and race, ethnicity and socioeconomic status. *Evidence-Based Prev Med* 2004;1(2):135-144.
177. Frank L, Kerr J, Rosenberg D, King A. Healthy aging and where you live: community design relationships with physical activity and body weight in older Americans. *J Phys Act Health*. 2010;7 (suppl 1):S82-S90.
178. Davison KK, Werder JL, Lawson CT. Children's active commuting to school: current knowledge and future directions. *Prev Chronic Dis*. 2008;5(3):1-11.
179. Lynott J, Taylor A, Twaddell H, et al. Planning complete streets for an aging America. <http://assets.aarp.org/rgcenter/ppi/liv-com/2009-12-streets.pdf>. Accessed May 16, 2013.
180. Harrell R, Brooks A, Nedwick T, AARP Public Policy Institute. Preserving affordability and access in liveable communities: subsidized housing opportunities near transit and the 50+ population. <http://assets.aarp.org/rgcenter/ppi/liv-com/2009-15x.pdf>. Accessed May 16, 2013.
181. Evenson KR, McGinn AP. Availability of school physical activity facilities to the public in four U.S. communities. *Am J Health Promot*. 2004;18(3):243-250.
182. Pincetl S, Wolch J, Wilson J, Longcore T. Toward a sustainable Los Angeles: a "nature's services" approach. http://sustainablecommunities.environment.ucla.edu/wp-content/uploads/2012/10/Toward_Sustainable_LA_2003.pdf. Accessed May 16, 2013.
183. Centers for Disease Control and Prevention. Physical activity levels among children aged 9-13 years - 2002. *MMWR Morb Mortal Wkly Rep*. 2003;52(33):785-788.
184. Choy LB, McGurk MD, Tamashiro R, Nett B, Maddock J. Increasing access to places for physical activity through a joint use agreement: a case study in urban Honolulu. *Prev Chronic Dis*. 2008;5(3):1-8.
185. National Policy & Legal Analysis Network to Prevent Childhood Obesity. NPLAN Joint Use Agreement Resources. http://changelabsolutions.org/sites/default/files/Joint%20Use%20Agreement%20Resources_FINAL_090901.pdf. Accessed January 30, 2013.
186. Spengler JO, Young SJ, Linton LS. Schools as a community resource for physical activity: legal considerations for decision makers. *Am J Health Promot*. 2007;21(suppl 4):390-396.
187. Healthy people, healthy places: snapshots of where we live, learn, work, and play. <http://ochealthinfo.com/civicax/filebank/blobdload.aspx?BlobID=14814>. Accessed July 18, 2013.
188. Latino Health Access. Latino health access park and community center: promoting healthy living.
189. Aytur SA, Satinsky SB, Evenson KR, Rodriguez DA. Pedestrian and bicycle planning in rural communities tools for active living. *Fam Community Health*. 2011;34(2):173-181.
190. Tarko A, Azam MS. Pedestrian injury analysis with consideration of the selectivity bias in linked police-hospital data. *Accid Anal Prev*. 2011;43(5):1689-1695.

REFERENCES (Continued)

191. US Department of Transportation Federal Highway Administration. *Guidelines and Recommendations to Accommodate Older Drivers and Pedestrians*. McLean, VA: US Department of Transportation;2001.
192. Black JL, Macinko J. Neighborhoods and obesity. *Nutr Rev*. 2008;66(1):2-20.
193. Wilcox S, Castro C, King AC, Housemann R, Brownson RC. Determinants of leisure time physical activity in rural compared with urban older and ethnically diverse women in the United States. *J Epidemiol Community Health*. 2000;54(9):667-672.
194. Reynolds KD, Wolch J, Byrne J, et al. Trail characteristics as correlates of urban trail use. *Am J Health Promot*. 2007;21(suppl 4):335-345.
195. Rimmer J. *Promoting Inclusive Physical Activity Communities for People with Disabilities*. Washington, DC: President's Council on Physical Fitness;2008. Series 9, No.2.
196. Walker JG, Evenson KR, Davis WJ, Bors P, Rodriguez DA. A tale of two trails: exploring different paths to success. *J Phys Act Health*. May 2011;8(4):523-533.
197. Rails to Trails Conservancy. Acquisition overview. http://www.railstotrails.org/ourwork/trailbuilding/toolbox/informationsummaries/acquisition_overview.html. Accessed June 1, 2012.
198. National Institute of Crime Prevention. Crime prevention through environmental design. <http://www.cptedtraining.net/>. Accessed May 22, 2013.
199. Babey SH, Hastert TA, Brown ER, UCLA Center for Health Policy Research. Teens living in disadvantaged neighborhoods lack access to parks and get less physical activity. <http://healthpolicy.ucla.edu/publications/Documents/PDF/Teens%20Living%20in%20Disadvantaged%20Neighborhoods%20Lack%20Access%20to%20Parks%20and%20Get%20Less%20Physical%20Activity.pdf>. Accessed May 15, 2013.
200. Hammerschmidt P, Tackett W, Golzynski M, Golzynski D. Barriers to and facilitators of healthful eating and physical activity in low-income schools. *J Nutr Educ Behav*. 2011;43(1):63-68.
201. Murphy NA, Carbone PS, and The Council on Children with Disabilities. Promoting the participation of children with disabilities in sports, recreation, and physical activities. *Pediatrics*. 2008;121(5):1057-1061.
202. Cox L, Berends V, Sallis J, et al. Engaging school governance leaders to influence physical activity policies. *J Phys Act Health*. 2011;8(suppl 1):40-48.
203. Pollack S, Bluestone B, Billingham C. Maintaining diversity in America's transit-rich neighborhoods: tools for equitable neighborhood change. <http://www.bos.frb.org/commdev/necd/2010/issue1/diversity-transit-rich-neighborhoods.pdf>. Accessed September 9, 2013.
204. Bailey L. Aging Americans: stranded without options. <http://www.transact.org/report.asp?id=232>. Accessed January 11, 2013.
205. Wilson S, Hutson M, Mujahid M. How planning and zoning contribute to inequitable development, neighborhood health, and environmental injustice. *Environmental Justice*. 2008;1(4):211-216.
206. Stommes E, Brown D, Houston C. Moving rural residents to work: lessons learned from implementation of eight job access and reverse commute projects. <http://www.fta.dot.gov/3630.html>. Accessed February 1, 2013.
207. Barten F, Mitlin D, Mulholland C, Hardoy A, Stern R. Integrated approaches to address the social determinants of health for reducing health inequity. *J Urban Health*. 2007;84(suppl 3):i164-173.
208. Sanoff H. Participation Purposes. In: Sanoff H, ed. *Community Participation Methods in Design and Planning*. New York, NY: John Wiley & Sons, Inc; 2000:1-32.
209. Policylink. Equitable development toolkit. http://www.policylink.org/site/c.lk1XLbMNJrE/b.5136575/k.39A1/Equitable_Development_Toolkit.htm. Accessed June 18 2004.

REFERENCES (Continued)

210. Snyder R, The Labor/Community Strategy Center. The Bus Riders Union transit model: why a bus-centered system will best serve US Cities. [http://www.thestrategycenter.org/sites/www.thestrategycenter.org/files/\[LCSC\]_BRU_Transit_Model_2009-04.pdf](http://www.thestrategycenter.org/sites/www.thestrategycenter.org/files/[LCSC]_BRU_Transit_Model_2009-04.pdf) Accessed August 22, 2012.
211. Rosenbloom S, Transportation Research News. The equity implications of financing the nation's surface transportation system. Accessed May 17, 2013.
212. Maryland Department of Health and Mental Hygiene. *Maryland Vital Statistics Annual Report*. Baltimore, MD: Maryland Department of Health and Mental Hygiene;2010.
213. MacDonald J, Golinelli D, Stokes RJ, Bluthenthal R. The effect of business improvement districts on the incidence of violent crimes. *Inj Prev*. 2010;16(5):327-332.
214. Casteel C, Peek-Asa C. Effectiveness of crime prevention through environmental design (CPTED) in reducing robberies. *Am J Prev Med*. 2000;18(suppl 4):99-115.
215. Bureau of Justice Assistance, Office of Justice Programs. *What Have We Learned From Evaluations of Crime Prevention Through Environmental Design Strategies?* Washington, DC: Center for Program Evaluation and Performance Measurement;2012.
216. Cozens PM, Saville G, Hillier D. Crime prevention through environmental design (CPTED): a review and modern bibliography. *Property Management*. 2005;23(5):328-356.
217. Skogan WG, Hartnett SM, Bump N, Dubois J. *Evaluation of CeaseFire-Chicago*. Rockville, MD: National Institute of Justice, National Criminal Justice Reference Service;2008. Document No. 227181.
218. Cohen L, Iton A, Davis RA, Rodriguez S, Prevention Institute. A time of opportunity: local solutions to reduce inequities in health and safety. <http://www.preventioninstitute.org/component/jlibrary/article/id-81/127.html>. Accessed August 22, 2012.
219. Wilkinson R. Why is violence more common where inequality is greater? *Ann NY Acad Sci*. 2004;1036:1-12.
220. Prevention Institute. Making the case: violence and health equity fact sheet. <http://www.preventioninstitute.org/component/jlibrary/article/id-311/127.html>. Accessed August 22, 2012.
221. Centers for Disease Control and Prevention. Youth violence: risk and protective factors. <http://www.cdc.gov/violenceprevention/youthviolence/riskprotectivefactors.html#3>. Accessed May 21, 2013.
222. Centers for Disease Control and Prevention (CDC). Youth violence national and state statistics at a glance. http://www.cdc.gov/ViolencePrevention/youthviolence/stats_at-a_glance/index.html. Accessed February 1, 2012.
223. Centers for Disease Control Prevention (CDC). Youth risk behavior surveillance United States, 2011. *MMWR Morb Mortal Wkly Rep*. 2012;61(4):1-162.
224. Egerter S, Barclay C, Grossman-Kahn R, Braveman P, Robert Wood Johnson Foundation. Exploring the social determinants of health series: violence, social disadvantage and health. http://www.rwjf.org/content/dam/farm/reports/issue_briefs/2011/rwjf70452. Accessed August 22, 2012.
225. Sampson RJ, Morenoff JD, Gannon-Rowley T. Assessing "neighborhood effects": social processes and new directions in research. *Annu Rev Sociol*. 2002;28:443-478.
226. Sampson RJ, Raudenbush SW, Earls F. Neighborhoods and violent crime: a multilevel study of collective efficacy. *Science*. 1997;277(5328):918-924.
227. Boston Public Health Commission. Selected health indicators. http://www.bphc.org/about/research/hob2010/Forms%20%20Documents/HOB11_Figs_4-13_SelHealthIndic_HOB11_PrintCopy_02Dec11.pdf. Accessed July 24, 2013.

REFERENCES (Continued)

228. Boston Public Health Commission. Boston moves for health: an action plan for healthy weight and healthy community. <http://www.bphc.org/programs/cib/chronicdisease/bostonmovesforhealth/Documents/BMH%20Action%20Plan%20-%20April%202012.pdf>. Accessed July 24, 2013.
229. Cohen L, Davis R, Lee V, Valdovinos E. Addressing the intersection: preventing violence and promoting healthy eating and active living. <http://www.preventioninstitute.org/component/jlibrary/article/id-267/127.html>. Accessed August 22, 2012.
230. Smith P, Pennington M, Crabtree L, Illback R. Louisville Metro health equity report: the social determinants of health in Louisville Metro neighborhoods. <http://www.louisvilleky.gov/NR/rdonlyres/29925903-E77F-46E5-8ACF-B801520B5BD2/O/HERFINALJAN23.pdf>. Accessed July 24, 2013.
231. Humes K, Jones N, Ramirez R. *Overview of Race and Hispanic Origin: 2010*. Washington, DC: US Department of Commerce, Economics and Statistics Administration, US Census Bureau;2011.
232. US Department of Health and Human Services. *HHS Action Plan to Reduce Racial and Ethnic Health Disparities: A Nation Free of Disparities in Health and Healthcare*. Washington, D.C.: US Dept of Health and Human Services;2011.
233. Flegal KM, Carroll MD, Kit BK, Ogden CL. Prevalence of obesity and trends in the distribution of body mass index among US adults, 1999-2010. *JAMA*. 2012;307(5):491-497.
234. Centers for Disease Control and Prevention. Current cigarette smoking among adults -- United States, 2011. *MMWR Morb Mortal Wkly Rep*. 2012;61(44):889-908.
235. Substance Abuse and Mental Health Services Administration, Center for Behavioral Health Statistics and Quality. *The NSDUH Report: Adults with Mental Illness or Substance Use Disorder Account for 40 percent of All Cigarettes Smoked*. Rockville, MD: SAMHSA;2013.
236. US Census Bureau. 2010 Census. Number and percent of population: 2010 - United States -- urban/rural and inside/outside metropolitan and micropolitan Area. Summary File 1, Table P1. http://factfinder2.census.gov/faces/tableservices/jsf/pages/productview.xhtml?pid=DEC_10_SF1_GCTP1.US26&prodType=table. Accessed February 13, 2013.
237. Befort CA, Nazir N, Perri MG. Prevalence of obesity among adults from rural and urban areas of the United States: findings from NHANES (2005-2008). *J Rural Health*. 2012(28):392-397.
238. Centers for Disease Control and Prevention, National Center for Health Statistics. Health indicators warehouse. http://healthindicators.gov/Indicators/Cigarette-smoking-adults-percent_1498/Profile/Data. Accessed February 15, 2013.
239. Centers for Disease Control and Prevention. Prevalence and most common causes of disability among adults -- United States, 2005. *MMWR Morb Mortal Wkly Rep*. 2009;58(16):421-426.
240. Centers for Disease Control and Prevention. Physical inactivity and people with disabilities: a tip sheet for public health professionals. http://www.cdc.gov/ncbddd/documents/physical-inactivity-tip-sheet-_phpa_1.pdf. Accessed July 11, 2012.
241. Centers for Disease Control and Prevention. Obesity and people with disabilities: a tip sheet for public health professionals. http://www.cdc.gov/ncbddd/documents/Obesity%20tip%20sheet%20_%20PHPa_1.pdf. Accessed June 06, 2012.
242. Braveman P, Egerter S, Robert Wood Johnson Foundation. Overcoming obstacles to health: report from the Robert Wood Johnson Foundation to the Commission to Build a Healthier America. <http://www.commissiononhealth.org/PDF/ObstaclesToHealth-Report.pdf> Accessed August 21, 2012.
243. Grimm K, Foltz J, Blanck H, Scanlon K. Household income disparities in fruit and vegetable consumption by state and territory: results of the 2009 Behavioral Risk Factor Surveillance System. *J Acad Nutr Diet*. 2012;112(12):2014-2021.

REFERENCES (Continued)

244. Ryan CL, Siebens J, US Census Bureau. Educational attainment in the United States: 2009. <http://www.census.gov/prod/2012pubs/p20-566.pdf>. Accessed June 11, 2012.
245. US Department of Health and Human Services. Nutrition, physical activity, and obesity. <http://www.healthypeople.gov/2020/lhi/nutrition.aspx?tab=data>. Accessed July 19, 2013.
246. Vincent GK, Velkoff VA, US Census Bureau. The next four decades: the older population in the United States: 2010 to 2050. <http://www.census.gov/prod/2010pubs/p25-1138.pdf>. Accessed January 14, 2012.
247. Freid V, Bernstein A, Bush MA, National Center for Health Statistics. Multiple chronic conditions among adults aged 45 and over: trends over the past 10 years. <http://www.cdc.gov/nchs/data/databriefs/db100.pdf>. Accessed January 25, 2012.
248. Wallace S. Social determinants of health inequities and healthcare in old age. In: Prohaska T, Anderson L, Binstock R, eds. *Public Health for an Aging Society*. Baltimore, MD: Johns Hopkins University Press; 2012:99-118.
249. Gates GJ, Williams Distinguished Scholar; The Williams Institute; UCLA School of Law. How many people are lesbian, gay, bisexual, and transgender? <http://williamsinstitute.law.ucla.edu/research/census-lgbt-demographics-studies/how-many-people-are-lesbian-gay-bisexual-and-transgender/>. Accessed August 21, 2012.
250. Boehmer U, Bowen DJ, Bauer GR. Overweight and obesity in sexual-minority women: evidence from population-based data. *Am J Public Health*. 2007;97(6):1134-1140.
251. Centers for Disease Control and Prevention. Behavioral risk factor surveillance system. <http://www.cdc.gov/brfss/>. Accessed May 22, 2013.
252. United States Census Bureau. Census news. <http://www.census.gov/#>. Accessed May 22, 2012.
253. Community Commons. About Community Commons. <http://initiatives.communitycommons.org/About.aspx>. Accessed May 22, 2013.
254. County Health Rankings and Robert Wood Johnson Foundation. Rankings. <http://www.countyhealthrankings.org/>. Accessed May 22, 2013.
255. United States Department of Health & Human Services. Community health status indicators report. <http://wwwn.cdc.gov/CommunityHealth/homepage.aspx?j=1>. Accessed May 22, 2013.
256. Hillmeier M, Lynch J, Harper S, Casper M, Centers for Disease Control and Prevention. Data set directory of social determinants of health at the local level. http://www.cdc.gov/dhdsp/docs/data_set_directory.pdf. Accessed May 22, 2013.
257. Healthy Communities Institute. Healthy communities network. <http://new.healthycommunitiesinstitute.com/healthy-communities-network-2/>. Accessed May 22, 2013.
258. National Cancer Institute. Health disparities calculator (HD*Calc). <http://seer.cancer.gov/hdcalc/>. Accessed May 22, 2013.
259. Connecticut Association of Directors of Health. Health equity index. <http://www.cadh.org/health-equity/health-equity-index.html>. Accessed May 22, 2013.
260. National Center for Health Statistics and Health Indicators Warehouse. About the HIW. <http://healthindicators.gov/About/AboutTheHIW>. Accessed May 22, 2013.
261. Prevention Institute. THRIVE: Tool for Health and Resilience In Vulnerable Environments. <http://thrive.preventioninstitute.org/thrive/about.php>. Accessed May 22, 2013.



[CDC.GOV/HEALTHEQUITYGUIDE](https://www.cdc.gov/healthequityguide)