

# West Virginia

State Nutrition, Physical Activity, and Obesity Profile

2016

National Center for Chronic Disease Prevention and Health Promotion  
Division of Nutrition, Physical Activity, and Obesity





# West Virginia - State Nutrition, Physical Activity, and Obesity Profile

Many American communities lack environments that could support healthy diets and regular physical activity. Healthy dietary and physical activity practices can lower the risk for heart disease, type 2 diabetes, and some cancers.<sup>1,2</sup>

In addition, excess calorie intake and physical inactivity contribute to obesity, which is associated with some of the leading preventable chronic diseases, including heart disease, stroke, type 2 diabetes, and some cancers. Among adults, the medical costs associated with obesity are an estimated \$147 billion.<sup>1,2,3</sup>

Public health approaches can help make healthy options available, accessible, and affordable. The Centers for Disease Control and Prevention's Division of Nutrition, Physical Activity, and Obesity (DNPAO) supports the nation to establish sustainable programs to improve dietary quality, increase physical activity, and reduce obesity and overweight. DNPAO funds and works with state health departments through a cooperative agreement ([State Public Health Actions to Prevent and Control Diabetes, Heart Disease, Obesity and Associated Risk Factors and Promote School Health program](#)).

This profile highlights data on dietary quality, physical activity, and obesity in West Virginia and select activities funded with this cooperative agreement. For more information about CDC's work to support healthy eating and physical activity, please see the [DNPAO website](#).

## Population Statistics

Estimated Population of West Virginia: 1,844,128<sup>4</sup>

## Adult Statistics

### Dietary Behaviors<sup>5</sup>

- 46.0% of adults reported consuming fruit less than one time daily.
- 24.2% of adults reported consuming vegetables less than one time daily.

### Physical Activity<sup>5</sup>

- 47.6% of adults achieved the equivalent of at least 150 minutes of moderate intensity physical activity per week.

### Overweight and Obesity<sup>5</sup>

- 34.0% of adults were overweight.
- 35.7% of adults had obesity.

## Adolescent Statistics

### Dietary Behaviors<sup>5</sup>

- 39.4% of adolescents reported consuming fruit less than one time daily.

- 37.5% of adolescents reported consuming vegetables less than one time daily.

### Physical Activity<sup>5</sup>

- 31.0% of adolescents were physically active at least 60 minutes per day on all 7 days in the past week.

### Overweight and Obesity<sup>5</sup>

- 15.5% of adolescents were overweight.
- 15.6% of adolescents had obesity.

## Child Statistics

### Breastfeeding<sup>5</sup>

- 62.3% of infants were ever breastfed.
- 34.3% of infants were breastfed for at least 6 months.

### Overweight and Obesity<sup>5</sup>

- 13.7% of children aged 2 to 4 years in the Special Supplemental Nutrition Program for Women, Infants, and Children (WIC) program were overweight.
- 14.1% of children aged 2 to 4 years in WIC had obesity.

# West Virginia - State Nutrition, Physical Activity, and Obesity Profile

## State Activities

West Virginia worked on the following select activities during the first 2 years of the State Public Health Actions program.

- Worked with partners to provide technical assistance (TA) to strengthen obesity prevention licensing and regulations in early care and education (ECE) settings. Partners included the West Virginia Bureau for Children and Families, the West Virginia University Extension Office, and the West Virginia Department of Education
- Collaborated with partner organizations to train and provide TA to ECE providers on Nutrition and Physical Activity Self-Assessment for Child Care (NAP SACC) implementation and follow up.
- Provided a condensed NAP SACC survey to all 361 licensed child care centers in the state. A 45% response rate was achieved. Survey analysis will assess where ECE programs can improve healthy eating and physical activity practices.



---

## References

1. US Department of Health and Human Services. National Institutes of Health. Managing Overweight and Obesity in Adults: Systematic Evidence Review from the Obesity Expert Panel, 2013.
2. World Cancer Research Fund / American Institute for Cancer Research. Food, Nutrition, Physical Activity, and the Prevention of Cancer: a Global Perspective. Washington DC: AICR, 2007
3. Finkelstein, EA, Trogon, JG, Cohen, JW, Dietz, W. Annual medical spending attributable to obesity: Payer- and service-specific estimates. *Health Affairs*. 2009;28(5):w822-w831.
4. US Census Bureau. State and County QuickFacts 2015 website. <http://www.census.gov/quickfacts/chart/PST045215/00>. Accessed February 22, 2016.
5. Centers for Disease Control and Prevention. Nutrition, Physical Activity and Obesity Data, Trends and Maps website. [https://nccd.cdc.gov/NPAO\\_DTM/](https://nccd.cdc.gov/NPAO_DTM/). Accessed February 22, 2016.