

Cover illustration courtesy of the Los Angeles County
Childhood Lead Poisoning Prevention Program

Managing Elevated Blood Lead Levels Among Young Children: Recommendations from the Advisory Committee on Childhood Lead Poisoning Prevention

Edited by
Birt Harvey, MD

**Centers for Disease Control
and Prevention**

Jeffrey P. Koplan, MD, MPH, Director

National Center for Environmental Health

Richard J. Jackson, MD, MPH, Director

Division of Environmental Hazards and Health Effects

Michael A. McGeehin, PhD, MSPH, Director

Lead Poisoning Prevention Branch

Gary P. Noonan, MPA, Acting Chief

**U.S. Department of Health
and Human Services, Public Health Service**

March 2002

Suggested reference:

Centers for Disease Control and Prevention. Managing Elevated Blood Lead Levels Among Young Children: Recommendations from the Advisory Committee on Childhood Lead Poisoning Prevention. Atlanta: CDC; 2002.

Table of Contents

List of Tables	iii
List of Figures	v
Foreword	vii
Preface	ix
Members of the Advisory Committee on Childhood Lead Poisoning Prevention	xi
Authors	xv
Acknowledgments	xvii
Glossary	xix
Chapter 1. Introduction	1
Development of the Case Management Recommendations	3
Overview of Comprehensive Case Management	4
General Considerations	9
Chapter 2. Assessment and Remediation of Residential Lead Exposure	13
Summary of Recommendations	15
Introduction	16
Sources and pathways of residential lead exposure	16
Effectiveness and Safety of Lead Hazard Control Measures	19
Recommendations for Assessment and Remediation	22
Enforcement of Laws and Regulations	25
Financial Resources for Lead Hazard Control	26
Recommendations for Future Research	27
Chapter 3. Medical Assessment and Interventions	39
Summary of Recommendations	41
Introduction	42
General Principles of Medical Case Management	42
Medical History	43
Physical Examination	48
Laboratory and Imaging Evaluation	48
Chelation Therapy	49

Monitoring Blood Lead Levels	50
Monitoring the Child	52
Recommendations for Future Research	53
 Chapter 4. Nutritional Assessment and Interventions	 59
Summary of Recommendations	61
Introduction	62
Nutritional Interventions: Summary of the Evidence	62
General Recommendations	68
Recommendations for Future Research	70
 Chapter 5. Developmental Assessment and Interventions	 77
Summary of Recommendations	79
Introduction	80
Detailed Bases for Recommendations	80
General Recommendations	85
Recommendations for Future Research	88
 Chapter 6. Educational Interventions for Caregivers	 97
Summary of Recommendations	99
Introduction	100
Sources and Pathways of Residential Lead Exposure	100
General Principles	100
Studies of Various Interventions	101
General Recommendations	105
Recommendations for Future Research	110
 Appendixes	 115
Appendix I. Published Reports of Less Common Causes of Elevated Blood Lead Levels (EBLLs) in Children.	117
Appendix II. Sources of Information on Lead Abatement	128

List of Tables

Table 1.1. Possible Elements of a Case Management Plan Based on Individualized Child Assessment	11
Table 2.1. Summary of Recommendations for Assessment and Remediation of Residential Lead Exposure	15
Table 2.2. Time Frames for Environmental Investigation and Other Case Management Activities According to a Child’s Blood Lead Level	36
Table 2.3. Common Sources of Lead Exposure to Consider in an Environmental Investigation	37
Table 3.1. Summary of Recommendations for Children with Confirmed (Venous) Elevated Blood Lead Levels	41
Table 3.2. Guidelines for Questions to Ask Regarding a Child’s Environmental History	45
Table 3.3. Recommended Schedule for Obtaining a Confirmatory Venous Sample	51
Table 3.4. Schedule for Follow-Up Blood Lead Testing	51
Table 4.1. Summary of Recommendations for Nutritional Assessment and Interventions	61
Table 5.1. Summary of Recommendations for Developmental Assessment and Interventions	79
Table 6.1. Summary of Recommendations for Educational Interventions for Caregivers	99

List of Figures

Figure 2.1. Pathways of Lead Exposure in the Residential Environment	33
Figure 2.2. Relationship of Housing Age and Condition to Dust Lead Levels	34
Figure 2.3. Relationship of Dust Lead Levels to Blood Lead Levels in Children	35
Figure 3.1. Lowest Reported Effect Levels of Inorganic Lead in Children	58
Figure 4.1. Median Calcium Intake by Race from NHANES III	76

Foreword

The overall reduction in childhood lead levels over the last three decades has been one of the great environmental health success stories in this country. However, our goal has not yet been reached. There are still far too many lower-income children living in older housing who are being hurt by elevated blood lead levels. The public health, housing, and environmental communities must continue to work together to eliminate the threat of lead poisoning for our future generations.

An important factor in the battle against lead poisoning is the proper management of children who have been identified as having elevated blood lead levels. In this publication, the Advisory Committee on Childhood Lead Poisoning Prevention (ACCLPP) and other public health practitioners have developed guidelines for assessment and interventions in the areas of medicine, nutrition, environmental exposure, childhood development, and education. Implementation of these “Best Practices” will greatly assist case managers, medical care providers, and others in delivering the most effective services to the lead poisoned child and the child’s caregiver.

I congratulate the ACCLPP and all the authors of these guidelines and thank them for their efforts. This report is a critical piece in the nation’s effort to eliminate childhood lead poisoning in America by the year 2010.

Richard Joseph Jackson, MD, MPH
Director, National Center for Environmental Health

Preface

Because case management of children with elevated blood lead levels varies markedly among states, cities, and other jurisdictions, the Advisory Committee on Childhood Lead Poisoning Prevention developed these nationally applicable recommendations. Based on recently published studies and augmented with opinions of experts, this report defines the elements of case management and offers assessment and management guidelines for health departments, case managers, primary care physicians, and other professionals. Not all recommendations are appropriate for any individual child because of variations in age, blood lead level, housing status, and—most important—the ability of caregivers to respond to recommendations without being overwhelmed.

The report contains five chapters in addition to the introduction: home environment investigation and interventions, medical evaluation and treatment, nutritional assessment and dietary modification, developmental surveillance and interventions, and education for caregivers. At the beginning of each chapter is a summary table of specific management recommendations. (The remainder of the tables, the figures, and the references are at the end of each chapter.) The text of the chapters provides the detailed information and references upon which most recommendations are based. Each chapter concludes with suggestions for further research.

This report, in addition to addressing the case management of individual children, also discusses the importance of state laws, regulations, and financing related to lead abatement efforts and the provision of appropriate services for affected children. Finally, the authors of this report recognize that case management is involved with the secondary prevention of elevated blood lead levels and that primary prevention by the removal of ongoing lead exposure sources should be promoted as the ideal and most effective means of preventing elevated blood lead levels.

Members of the Advisory Committee on Childhood Lead Poisoning Prevention

March 2002

ACTING CHAIR

Carla C. Campbell, MD, MS
Children's Hospital of Philadelphia
Philadelphia, Pennsylvania

EXECUTIVE SECRETARY

Gary P. Noonan, MPA
Acting Chief, Lead Poisoning Prevention Branch
National Center for Environmental Health, CDC
Atlanta, Georgia

MEMBERS

Cushing N. Dolbeare
Housing and Public Policy Consultant
Washington, D.C.

Anne M. Guthrie, MPH
Alliance to End Childhood Lead Poisoning
Charlottesville, Virginia

Birt Harvey, MD
Pediatrician
Palo Alto, California

Richard E. Hoffman, MD, MPH
Physician
Denver, Colorado

Amy A. Murphy, MPH
City of Milwaukee Health Department
Milwaukee, Wisconsin

Estelle B. Richman, M.A.
Philadelphia Department of Public Health
Philadelphia, Pennsylvania

Joel D. Schwartz, PhD
Harvard School of Public Health
Boston, Massachusetts

Michael W. Shannon, MD, MPH
Children's Hospital Boston
Boston, Massachusetts

Michael L. Weitzman, MD
University of Rochester
Rochester, New York

EX OFFICIO MEMBERS

Agency for Toxic Substances and Disease Registry
Olivia Harris, MA

Centers for Medicare and Medicaid Services
Jerry Zelinger, MD

Health Resources and Services Administration
Byron P. Bailey, MPH

National Institute for Occupational Safety and Health, CDC
Robert J. Roscoe, MS

National Institute of Environmental Health Sciences
Walter Rogan, MD

U.S. Agency for International Development
John Borrazzo, PhD

U.S. Consumer Product Safety Commission
Lori Saltzman

U.S. Department of Housing and Urban Development
David Jacobs, PhD

U.S. Environmental Protection Agency
William H. Sanders, III, DrPH

U.S. Food and Drug Administration
Michael P. Bolger, PhD

LIAISON REPRESENTATIVES

American Academy of Pediatrics

J. Routt Reigart, II, MD

American Association of Poison Control Centers

George C. Rodgers, Jr., MD, PhD

American Industrial Hygiene Association

Steve M. Hays

American Public Health Association

Rebecca Parkin, PhD, MPH

Association of Public Health Laboratories

Henry Bradford, Jr., PhD

Association of State and Territorial Health Officials

Karen Pearson

Council of State and Territorial Epidemiologists

Ezatollah Keyvan-Larijani, MD, DrPH

National Center for Healthy Housing

Pat McLaine, MPH

Authors

Assessment and Remediation of Residential Lead Exposure

Prepared by:

Thomas D. Matte, MD, MPH

Dennis Kim, MD, MPH

Division of Environmental Hazards and Health Effects

National Center for Environmental Health

Mark R. Farfel, PhD

Kennedy Krieger Institute

Developmental Assessment and Interventions

Prepared by:

David Bellinger, PhD, MSc

Harvard Medical School

Leonard Rappaport, MD

Harvard Medical School

Educational Interventions for Caregivers

Prepared by:

James R. Campbell, MD, MPH

University of Rochester School of Medicine and Dentistry

Michael L. Weitzman, MD

University of Rochester School of Medicine and Dentistry

Medical Assessment and Interventions

Prepared by:

James R. Roberts, MD, MPH

Medical University of South Carolina

J. Routt Reigart, MD

Medical University of South Carolina

Nutritional Assessment and Interventions

Prepared by:

James D. Sargent, MD

Dartmouth Medical School

Acknowledgments

I am indebted to many people for their input into this document. The Working Group decided on the format and appropriate individuals to develop each chapter, reviewed with those authorities the initial drafts, and suggested modifications that the chapter authors invariably felt benefitted their work. The Working Group members were Carla Campbell, Isabella Clemente, Susan Cummins, Patricia McLaine, Tom Matte, Joel Schwartz, and Michael Weitzman.

For each chapter, its authors, CDC experts in the subject, and outside experts attended a Working Group meeting. They provided background information, additional citations, and suggestions for modifications. Among these experts were Susan Adubato, Carol Ballew, Andrea Carlson, Julian Chisholm, Mary Cogswell, Peter Dallman, Mark Farfel, Warren Galke, Scott Grosse, Randy Louchart, Kathryn Mahaffey, Morri Markowitz, Tim Morta, George Rhoads and Walter Rogan.

The document was reviewed at meetings attended by ACCLPP members, ex officio members, liaison representatives, and chapter authors. Changes requested by attendees were incorporated in the final document.

Special thanks should go to CDC staff members Alan Bloch and Jerry Hershovitz, who assisted in the initial stages of reviewing and commenting on the document; Becky Wright, who organized meetings and typed revisions; Philip Jacobs and Nikki Kilpatrick, who assisted with final formatting; Pamela Meyer, who shepherded the document to its present form; and Joey Johnson and Connie Woodall, who provided graphics support.

Sheila Jurik, Kevin Moran, and Pam Gillis Watson provided editorial support.

Last, my personal thanks to Susan Cummins, chair of the ACCLPP at the time this project was initiated, without whose efforts this document would never have been developed or published.

Birt Harvey, Editor
Chair, Working Group

Glossary

ACCLPP—Advisory Committee for Childhood Lead Poisoning Prevention.

Acidosis—a condition resulting from the accumulation of acid or depletion of bicarbonate content in the blood and tissues.

Aminoaciduria—an excess of amino acids in the urine.

Asymptomatic—without signs or symptoms.

Ataxia—failure of muscular coordination; irregularity of muscular action.

Bioavailable—readily absorbed and used by the body.

BLL—blood lead level, usually measured in micrograms per deciliter (Fg/dL).

Caregiver—parent, guardian, or other person involved in a child’s daily care.

CDC—Centers for Disease Control and Prevention; part of the U.S. Department of Health and Human Services, Public Health Service.

Chelation therapy—the use of chelating agents (chemical compounds that bind to metals) to remove toxic metals such as lead from the body.

Clearance standards—maximum allowable lead levels on surfaces (e.g., floors, windowsills, and window wells) after a residence has undergone lead abatement.

Drip line—the area under the edges of a roof.

EBLL—elevated blood lead level, defined as any blood lead level ≥ 10 Fg/dL.

Encephalopathy—extensive swelling of the brain.

Environmental investigation—an investigation by trained personnel at a child’s residence (or any secondary addresses where the child spends significant amounts of time) to identify lead hazards.

Gingival lead lines—darkening of the gums just distal to the insertion of the tooth.

Glucosuria—the presence of glucose in urine.

HUD—U.S. Department of Housing and Urban Development.

Hypophosphatemia—an abnormally low blood phosphate level.

Fg/dL—micrograms per deciliter, the usual unit of measure for blood lead levels.

Fg/ft²—micrograms per square foot, a unit of measure for measuring dust lead loading.

Papilledema—excessive fluid in the optic disk; also called choked disk.

PCP—primary care provider, the health professional who oversees a child’s care, usually a physician, nurse practitioner, or physician’s assistant.

Phosphaturia—an abnormally high urine phosphate level.

Pica—compulsive eating of nonnutritive substances such as dirt or flaking paint.

ppb—parts per billion.

ppm—parts per million.

Primary prevention—preventing a problem before it occurs. Primary prevention of lead poisoning would eliminate lead sources, thus preventing exposure.

Proteinuria—excess protein in the urine.

Radiograph—a film record of internal structures produced by passing x-rays or gamma rays through the body; frequently referred to as an “x-ray.”

Renal—having to do with the kidneys.

Secondary prevention—responding to a problem after it has been detected. Secondary prevention of lead poisoning involves identifying children with EBLs and eliminating or reducing their lead exposure.

WIC—Special Supplemental Nutrition Program for Women, Infants, and Children.

