



# HIV Recommended Queries

# HIV diagnoses (2011)

All races/ethnicities | Male | 2011 | Adults and adolescents | Male-to-male sexual contact




## Cases



Legend classified according to 2011 data.

## National Data By Year

With the current query, additional data and graphics are not available. Performing a new query may allow for additional graphics.



**Disclaimer:** This is a user-generated report. The findings and conclusions are those of the user and do not necessarily represent the views of the CDC.



**Data Source:** Query and graphics are based on HIV surveillance data. Data may include a combination of the following: HIV infection or stage 3 (AIDS) diagnoses, deaths among persons with HIV infection or stage 3 (AIDS), or persons living with HIV infection or stage 3 (AIDS); by state, year, race/ethnicity, sex, age group, and transmission category; shown in numbers and rates.

**Notes:** All displayed diagnoses of HIV infection and diagnoses of infection classified as stage 3 (AIDS) data have been statistically adjusted to account for reporting delays, but not for incomplete reporting. *Data displayed are for individuals 13 years of age or older, and therefore may not reflect the exact numbers and rates in the 2011 HIV Surveillance Report.* The estimated numbers and rates of diagnoses of HIV infection and stage 3 (AIDS) are based on data from 56 areas (50 states, the District of Columbia, American Samoa, Guam, the Northern Mariana Islands, Puerto Rico, and the U.S. Virgin Islands) in which laws or regulations require confidential reporting by name for adults and adolescents with confirmed diagnoses of HIV infection. Data for the Republic of Palau are not currently included in the Atlas. For more info, see: [HIV Surveillance Technical Notes](#). Suggested citation: [NCHHSTP Atlas](#)



# HIV diagnoses (2011)

Black/African American | Both sexes | 2011 | Adults and adolescents | All transmission categories




## Rate per 100,000 among selected population



Legend classified according to 2011 data.

## National Data By Sex



**Disclaimer:** This is a user-generated report. The findings and conclusions are those of the user and do not necessarily represent the views of the CDC.

**Data Source:** Query and graphics are based on HIV surveillance data. Data may include a combination of the following: HIV infection or stage 3 (AIDS) diagnoses, deaths among persons with HIV infection or stage 3 (AIDS), or persons living with HIV infection or stage 3 (AIDS); by state, year, race/ethnicity, sex, age group, and transmission category; shown in numbers and rates.

**Notes:** All displayed diagnoses of HIV infection and diagnoses of infection classified as stage 3 (AIDS) data have been statistically adjusted to account for reporting delays, but not for incomplete reporting. *Data displayed are for individuals 13 years of age or older, and therefore may not reflect the exact numbers and rates in the 2011 HIV Surveillance Report.* The estimated numbers and rates of diagnoses of HIV infection and stage 3 (AIDS) are based on data from 56 areas (50 states, the District of Columbia, American Samoa, Guam, the Northern Mariana Islands, Puerto Rico, and the U.S. Virgin Islands) in which laws or regulations require confidential reporting by name for adults and adolescents with confirmed diagnoses of HIV infection. Data for the Republic of Palau are not currently included in the Atlas. **For more info, see: [HIV Surveillance Technical Notes](#). Suggested citation: [NCHHSTP Atlas](#)**


Centers for Disease Control and Prevention

National Center for HIV/AIDS, Viral Hepatitis, STD, and TB Prevention



# HIV diagnoses (2011)

Hispanic/Latino | Both sexes | 2011 | Adults and adolescents | All transmission categories




## Rate per 100,000 among selected population



Legend classified according to 2011 data.



## National Data By Sex



**Disclaimer:** This is a user-generated report. The findings and conclusions are those of the user and do not necessarily represent the views of the CDC.


**Data Source:** Query and graphics are based on HIV surveillance data. Data may include a combination of the following: HIV infection or stage 3 (AIDS) diagnoses, deaths among persons with HIV infection or stage 3 (AIDS), or persons living with HIV infection or stage 3 (AIDS); by state, year, race/ethnicity, sex, age group, and transmission category; shown in numbers and rates.

**Notes:** All displayed diagnoses of HIV infection and diagnoses of infection classified as stage 3 (AIDS) data have been statistically adjusted to account for reporting delays, but not for incomplete reporting. **Data displayed are for individuals 13 years of age or older, and therefore may not reflect the exact numbers and rates in the 2011 HIV Surveillance Report.** The estimated numbers and rates of diagnoses of HIV infection and stage 3 (AIDS) are based on data from 56 areas (50 states, the District of Columbia, American Samoa, Guam, the Northern Mariana Islands, Puerto Rico, and the U.S. Virgin Islands) in which laws or regulations require confidential reporting by name for adults and adolescents with confirmed diagnoses of HIV infection. Data for the Republic of Palau are not currently included in the Atlas. **For more info, see: [HIV Surveillance Technical Notes](#). Suggested citation: [NCHHSTP Atlas](#)**



# Persons living with HIV, stage 3 (AIDS) (2010)

All races/ethnicities | Female | 2010 | Adults and adolescents | All transmission categories




## Rate per 100,000 among selected population



Legend classified according to 2010 data.

## National Data By Age Group



**Disclaimer:** This is a user-generated report. The findings and conclusions are those of the user and do not necessarily represent the views of the CDC.


**Data Source:** Query and graphics are based on HIV surveillance data. Data may include a combination of the following: HIV infection or stage 3 (AIDS) diagnoses, deaths among persons with HIV infection or stage 3 (AIDS), or persons living with HIV infection or stage 3 (AIDS); by state, year, race/ethnicity, sex, age group, and transmission category; shown in numbers and rates.

**Notes:** All displayed diagnoses of HIV infection and diagnoses of infection classified as stage 3 (AIDS) data have been statistically adjusted to account for reporting delays, but not for incomplete reporting. *Data displayed are for individuals 13 years of age or older, and therefore may not reflect the exact numbers and rates in the 2011 HIV Surveillance Report.* The estimated numbers and rates of diagnoses of HIV infection and stage 3 (AIDS) are based on data from 56 areas (50 states, the District of Columbia, American Samoa, Guam, the Northern Mariana Islands, Puerto Rico, and the U.S. Virgin Islands) in which laws or regulations require confidential reporting by name for adults and adolescents with confirmed diagnoses of HIV infection. Data for the Republic of Palau are not currently included in the Atlas. For more info, see: [HIV Surveillance Technical Notes](#). Suggested citation: [NCHHSTP Atlas](#)



# HIV diagnoses (2011)

All races/ethnicities | Both sexes | 2011 | Adults and adolescents | All transmission categories




Rate per 100,000 among selected population



Legend classified according to 2011 data.

## National Data By Race/ethnicity



**Disclaimer:** This is a user-generated report. The findings and conclusions are those of the user and do not necessarily represent the views of the CDC.

**Data Source:** Query and graphics are based on HIV surveillance data. Data may include a combination of the following: HIV infection or stage 3 (AIDS) diagnoses, deaths among persons with HIV infection or stage 3 (AIDS), or persons living with HIV infection or stage 3 (AIDS); by state, year, race/ethnicity, sex, age group, and transmission category; shown in numbers and rates.

**Notes:** All displayed diagnoses of HIV infection and diagnoses of infection classified as stage 3 (AIDS) data have been statistically adjusted to account for reporting delays, but not for incomplete reporting. **Data displayed are for individuals 13 years of age or older, and therefore may not reflect the exact numbers and rates in the 2011 HIV Surveillance Report.** The estimated numbers and rates of diagnoses of HIV infection and stage 3 (AIDS) are based on data from 56 areas (50 states, the District of Columbia, American Samoa, Guam, the Northern Mariana Islands, Puerto Rico, and the U.S. Virgin Islands) in which laws or regulations require confidential reporting by name for adults and adolescents with confirmed diagnoses of HIV infection. Data for the Republic of Palau are not currently included in the Atlas. For more info, see: [HIV Surveillance Technical Notes](#). Suggested citation: [NCHHSTP Atlas](#)

