

Data Brief 193: Hypertension-related Mortality in the United States, 2000–2013

Data table for Figure 1. Number of deaths and rates for hypertension-related and all other causes of death: United States, 2000–2013

Year	Number of hypertension-related deaths	Number of all other causes of death	Age-adjusted hypertension-related death rates	Age-adjusted death rates for all other causes of death combined	Percent of all deaths for hypertension-related deaths
2000	245,220	1,955,792	255.1	2,032.7	11.1
2001	255,250	1,955,374	260.9	1,996.0	11.5
2002	270,896	1,966,551	272.6	1,974.8	12.1
2003	286,160	1,957,513	282.7	1,930.0	12.8
2004	293,026	1,905,037	285.0	1,847.6	13.3
2005	311,685	1,935,335	296.7	1,837.8	13.9
2006	318,201	1,907,654	296.2	1,772.7	14.3
2007	328,427	1,898,936	299.0	1,727.1	14.7
2008	339,672	1,942,882	302.4	1,729.2	14.9
2009	339,828	1,912,984	296.0	1,666.3	15.1
2010	354,813	1,937,472	303.6	1,659.1	15.5
2011	368,855	1,969,529	306.4	1,639.5	15.8
2012	381,406	1,985,744	309.1	1,613.4	16.1
2013	396,675	2,023,816	314.1	1,605.9	16.4

NOTES: Deaths are per 100,000 standard population for persons aged 45 and over. Hypertension-related deaths are identified using ICD–10 codes I10, I11, I12, I13, and I15 for underlying and contributing causes of death, according to the *International Classification of Diseases, 10th revision* (ICD–10).

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality.

Data Brief 193: Hypertension-related Mortality in the United States, 2000–2013

Data table for Figure 2. Hypertension-related death rates, by sex and selected age group: United States, 2000–2013

Year	Women, 85 and over	Men, 85 and over	Women, 75–84	Men, 75–84	Women, 65–74	Men, 65–74	Women, 45–64	Men, 45–64
2000	2,003.1	1,654.8	634.7	683.8	216.8	291.0	44.6	78.0
2001	2,083.3	1,711.1	641.5	703.9	218.8	295.0	45.1	78.7
2002	2,208.4	1,858.0	664.0	728.4	226.4	304.9	46.3	82.7
2003	2,307.8	1,942.2	679.5	755.0	227.2	314.2	49.1	88.7
2004	2,316.4	2,005.7	680.4	768.1	226.2	313.9	49.0	91.1
2005	2,464.4	2,110.4	704.1	788.1	229.6	320.7	50.7	96.3
2006	2,438.5	2,124.6	692.3	793.8	225.7	320.4	51.8	99.6
2007	2,487.3	2,193.7	693.0	799.8	220.1	320.0	52.5	101.5
2008	2,517.0	2,251.3	694.5	806.7	219.8	320.0	54.4	104.1
2009	2,418.8	2,192.0	675.0	791.5	214.1	318.6	54.0	106.6
2010	2,519.2	2,322.0	680.1	810.7	215.3	321.5	55.1	109.5
2011	2,520.6	2,299.9	685.6	815.9	215.5	329.2	57.7	113.9
2012	2,544.5	2,383.5	688.6	818.6	209.4	329.7	58.4	117.7
2013	2,549.7	2,398.8	689.7	837.3	212.7	338.9	61.0	123.4

NOTES: Deaths are per 100,000 population for persons aged 45 and over. Hypertension-related deaths are identified using ICD–10 codes I10, I11, I12, I13, and I15 for underlying and contributing causes of death, according to the *International Classification of Diseases, 10th revision (ICD–10)*.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality.

Data Brief 193: Hypertension-related Mortality in the United States, 2000–2013

Data table for Figure 3. Age-adjusted hypertension-related death rates, by race and Hispanic origin: United States, 2000–2013

Year	Non-Hispanic black	Non-Hispanic white	Hispanic
2000	523.8	228.5	233.7
2001	527.9	234.0	246.8
2002	539.5	246.2	251.7
2003	552.7	256.2	261.9
2004	550.2	259.2	260.6
2005	559.3	270.9	277.9
2006	548.1	271.6	274.2
2007	542.5	276.0	270.1
2008	535.6	280.8	268.5
2009	518.6	275.1	262.9
2010	519.9	282.8	279.3
2011	510.4	288.4	270.1
2012	506.4	291.6	273.8
2013	509.9	296.5	284.8

NOTES: Deaths are per 100,000 standard population for persons aged 45 and over. Hypertension-related deaths are identified using ICD–10 codes I10, I11, I12, I13, and I15 for underlying and contributing causes of death, according to the *International Classification of Diseases, 10th revision* (ICD–10).

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality.