

Data Brief 274: Utilization of Clinical Preventive Services for Cancer and Heart Disease Among Insured Adults: United States, 2015

Data table for Figure 1. Percentage of insured adults aged 50–75 who had any colorectal cancer screening test within the recommended intervals, by sex and age group: United States, 2015

Sex and age group	Percent	Standard error
Men		
Total	63.7	0.98
50–59	54.0	1.54
60–69	70.1	1.33
70–75	76.1	1.69
Women		
Total	64.4	0.86
50–59	58.1	1.36
60–69	69.5	1.25
70–75	70.1	1.91

NOTE: See "Definitions" for more information about the recommended intervals for this screening test.

SOURCE: NCHS, National Health Interview Survey, 2015.

Data Brief 274: Utilization of Clinical Preventive Services for Cancer and Heart Disease Among Insured Adults: United States, 2015

Data table for Figure 2. Percentage of insured women aged 21–65 who have not had a hysterectomy and had a cervical cancer screening test within the recommended intervals, by age group: United States, 2015

Age group	Percent	Standard error
Total	85.5	0.50
21–29	80.5	1.35
30–39	90.5	0.83
40–49	87.8	0.94
50–59	84.8	1.02
60–65	81.0	1.60

NOTE: See "Definitions" for more information about the recommended intervals for this screening test.

SOURCE: NCHS, National Health Interview Survey, 2015.

Data Brief 274: Utilization of Clinical Preventive Services for Cancer and Heart Disease Among Insured Adults: United States, 2015

Data table for Figure 3. Percentage of insured women aged 50–74 who had a mammogram in the past 2 years, by age group: United States, 2015

Age group	Percent	Standard error
Total	73.5	0.76
50–59	73.3	1.13
60–69	74.2	1.11
70–74	72.1	1.75

SOURCE: NCHS, National Health Interview Survey, 2015.

Data Brief 274: Utilization of Clinical Preventive Services for Cancer and Heart Disease Among Insured Adults: United States, 2015

Data table for Figure 4. Percentage of insured adults aged 18 and over who had their blood pressure checked by a doctor, nurse, or other health professional in the past 12 months, by sex and age group: United States, 2015

Sex and age group	Percent	Standard error
Men		
Total	82.0	0.53
18–39	70.4	1.00
40–64	86.1	0.67
65–79	93.8	0.65
80 and over	95.3	1.00
Women		
Total	90.7	0.33
18–39	86.7	0.69
40–64	91.7	0.43
65–79	95.4	0.47
80 and over	94.8	0.79

SOURCE: NCHS, National Health Interview Survey, 2015.

Data Brief 274: Utilization of Clinical Preventive Services for Cancer and Heart Disease Among Insured Adults: United States, 2015

Data table for Figure 5. Percentage of overweight and obese insured adults aged 40–70 who had a fasting test for high blood sugar or diabetes during the past 12 months, by sex and age group: United States, 2015

Sex and age group	Percent	Standard error
Men		
Total	60.1	0.97
40–49	49.5	1.69
50–59	60.9	1.61
60–70	70.1	1.54
Women		
Total	63.7	0.88
40–49	56.4	1.65
50–59	64.4	1.55
60–70	69.5	1.34

SOURCE: NCHS, National Health Interview Survey, 2015.