

**Data Brief 277: Selected Health Conditions Among Native Hawaiian and Pacific Islander Adults: United States, 2014**

**Data table for Figure 1. Age-sex-adjusted percentage of adults aged 18 and over in fair or poor health, or who had experienced serious psychological distress, by race: United States, 2014**

Race	Fair or poor health	Serious psychological distress in the past 30 days
	Percent (95% CI)	
2014 NHPI NHIS		
All NHPI	15.5 (14.0–17.0)	4.1 (2.9–5.3)
2014 NHIS		
Total U.S. population	12.0 (11.6–12.4)	3.1 (2.9–3.3)
Single-race Asian	9.2 (7.6–10.8)	1.6 (1.0–2.2)

NOTES: CI is confidence interval. NHPI is Native Hawaiian and Pacific Islander. NHIS is National Health Interview Survey. All NHPI refers to persons who were identified as NHPI alone or in combination with other races.

SOURCES: NCHS, Native Hawaiian and Pacific Islander National Health Interview Survey (NHPI NHIS), 2014 and NCHS, National Health Interview Survey (NHIS), 2014.

**Data Brief 277: Selected Health Conditions Among Native Hawaiian and Pacific Islander Adults: United States, 2014**

**Data table for Figure 2. Age-sex-adjusted percentage of adults aged 18 and over with cancer, coronary heart disease, or diabetes, by race: United States, 2014**

Race	Cancer	Coronary heart disease	Diabetes
Percent (95% CI)			
2014 NHPI NHIS			
All NHPI	5.7 (4.4–7.0)	6.0 (4.8–7.2)	15.6 (13.8–17.4)
2014 NHIS			
Total U.S. population	7.8 (7.4–8.2)	5.4 (5.1–5.7)	8.7 (8.3–9.1)
Single-race Asian	3.2 (2.3–4.1)	3.2 (2.3–4.1)	8.0 (6.5–9.5)

NOTES: CI is confidence interval. NHPI is Native Hawaiian and Pacific Islander. NHIS is National Health Interview Survey. All NHPI refers to persons who were identified as NHPI alone or in combination with other races.

SOURCES: NCHS, Native Hawaiian and Pacific Islander National Health Interview Survey (NHPI NHIS), 2014 and NCHS, National Health Interview Survey (NHIS), 2014.

**Data Brief 277: Selected Health Conditions Among Native Hawaiian and Pacific Islander Adults: United States, 2014**

**Data table for Figure 3. Age-sex-adjusted percentage of adults aged 18 and over with pain in lower back, arthritis, or migraines, by race: United States, 2014**

Race	Lower back pain in the past 3 months	Arthritis	Migraines or severe headaches in the past 3 months
Percent (95% CI)			
2014 NHPI NHIS			
All NHPI	28.5 (26.4–30.6)	19.7 (17.4–22.0)	14.1 (11.3–16.9)
2014 NHIS			
Total U.S. population	28.3 (27.6–29.0)	21.1 (20.6–21.6)	15.1 (14.6–15.6)
Single-race Asian	17.2 (15.2–19.2)	11.4 (9.6–13.2)	9.9 (8.3–11.5)

NOTES: CI is confidence interval. NHPI is Native Hawaiian and Pacific Islander. NHIS is National Health Interview Survey. All NHPI refers to persons who were identified as NHPI alone or in combination with other races.

SOURCES: NCHS, Native Hawaiian and Pacific Islander National Health Interview Survey (NHPI NHIS), 2014 and NCHS, National Health Interview Survey (NHIS), 2014.

**Data Brief 277: Selected Health Conditions Among Native Hawaiian and Pacific Islander Adults: United States, 2014**

**Data table for Figure 4. Age-sex-adjusted percentage of adults aged 18 and over who ever had asthma or still have asthma, by race: United States, 2014**

Race	Ever had asthma	Still have asthma
Percent (95% CI)		
2014 NHPI NHIS		
All NHPI	19.2 (17.0–21.4)	9.9 (8.2–11.6)
2014 NHIS		
Total U.S. population	12.8 (12.3–13.3)	7.4 (7.0–7.8)
Single-race Asian	9.2 (7.6–10.8)	4.8 (3.6–6.0)

NOTES: CI is confidence interval. NHPI is Native Hawaiian and Pacific Islander. NHIS is National Health Interview Survey. All NHPI refers to persons who were identified as NHPI alone or in combination with other races.

SOURCES: NCHS, Native Hawaiian and Pacific Islander National Health Interview Survey (NHPI NHIS), 2014 and NCHS, National Health Interview Survey (NHIS), 2014.