## Data Brief 282: Drug Overdose Deaths Among Adolescents Aged 15-19 in the United States: 1999-2015

## Data table for Figure 1. Drug overdose death rates for adolescents aged 15–19, by sex: United States, 1999–2015

[Rates per 100,000 population enumerated as of April 1 for 2000 and 2010 and estimated as of July 1 for all other years]

| Year |       |      |        |
|------|-------|------|--------|
|      | Total | Male | Female |
| 1999 | 1.6   | 2.1  | 1.1    |
| 2000 | 2.1   | 2.9  | 1.2    |
| 2001 | 2.5   | 3.4  | 1.5    |
| 2002 | 2.9   | 4.0  | 1.7    |
| 2003 | 3.1   | 4.3  | 1.8    |
| 2004 | 3.7   | 5.2  | 2.0    |
| 2005 | 3.5   | 5.0  | 1.9    |
| 2006 | 3.9   | 5.6  | 2.0    |
| 007  | 4.2   | 6.2  | 2.1    |
| 2008 | 4.2   | 6.1  | 2.1    |
| 2009 | 3.6   | 5.2  | 1.9    |
| 2010 | 3.8   | 5.3  | 2.2    |
| 2011 | 3.8   | 5.4  | 2.2    |
| 2012 | 3.1   | 4.4  | 1.7    |
| 2013 | 3.2   | 4.3  | 2.0    |
| 014  | 3.1   | 4.0  | 2.2    |
| 2015 | 3.7   | 4.6  | 2.7    |

NOTE: Drug overdose deaths are those with International Classification of Diseases, Tenth Revision underlying cause-of-death codes X40-X44, X60-X64, X85, and Y10-Y14.

## Data Brief 282: Drug Overdose Deaths Among Adolescents Aged 15-19 in the United States: 1999-2015

|               | Percent |       |        |  |
|---------------|---------|-------|--------|--|
| Intent        | Total   | Male  | Female |  |
| Total         | 100.0   | 100.0 | 100.0  |  |
| Unintentional | 80.4    | 86.2  | 70.1   |  |
| Suicide       | 13.5    | 8.7   | 21.9   |  |
| Homicide      | 0.9     | 0.6   | 1.4    |  |
| Undetermined  | 5.2     | 4.5   | 6.5    |  |

Data table for Figure 2. Percent distribution of drug overdose deaths for adolescents aged 15–19, by intent and sex: United States, 2015

NOTE: Drug overdose deaths are those with International Classification of Diseases, Tenth Revision underlying cause-of-death codes X40–X44 (unintentional), X60–X64 (suicide), X85 (homicide), and Y10–Y14 (undetermined).

#### Data Brief 282: Drug Overdose Deaths Among Adolescents Aged 15–19 in the United States: 1999–2015

### Data table for Figure 3. Drug overdose death rates for adolescents aged 15–19, by type of drug involved: United States, 1999–2015

[Rates per 100,000 population enumerated as of April 1 for 2000 and 2010 and estimated as of July 1 for all other years]

| Year | Rate                                   |         |                 |                                       |  |
|------|--|---------|-----------------|---------------------------------------|--|
|      | Drug overdose deaths involving opioids | Cocaine | Benzodiazepines | Psychostimulants with abuse potential |  |
| 1999 | 0.8                                    | 0.3     | *               | 0.1                                   |  |
| 2000 | 0.9                                    | 0.3     | 0.1             | 0.2                                   |  |
| 2001 | 1.3                                    | 0.3     | 0.2             | 0.2                                   |  |
| 2002 | 1.5                                    | 0.5     | 0.2             | 0.2                                   |  |
| 2003 | 1.8                                    | 0.5     | 0.3             | 0.2                                   |  |
| 2004 | 2.2                                    | 0.5     | 0.4             | 0.1                                   |  |
| 2005 | 2.0                                    | 0.5     | 0.4             | 0.2                                   |  |
| 2006 | 2.5                                    | 0.7     | 0.5             | 0.2                                   |  |
| 2007 | 2.7                                    | 0.5     | 0.5             | 0.1                                   |  |
| 2008 | 2.7                                    | 0.4     | 0.6             | 0.2                                   |  |
| 2009 | 2.5                                    | 0.2     | 0.7             | 0.1                                   |  |
| 2010 | 2.5                                    | 0.2     | 0.7             | 0.2                                   |  |
| 2011 | 2.6                                    | 0.3     | 0.6             | 0.2                                   |  |
| 2012 | 2.0                                    | 0.2     | 0.4             | 0.2                                   |  |
| 2013 | 2.0                                    | 0.1     | 0.4             | 0.3                                   |  |
| 2014 | 2.0                                    | 0.2     | 0.5             | 0.2                                   |  |
| 2015 | 2.4                                    | 0.3     | 0.6             | 0.4                                   |  |

\* Figure does not meet standards of reliability or precision.

NOTES: Drug overdose deaths are those with International Classification of Diseases, Tenth Revision underlying cause-of-death codes X40–X44, X60–X64, X85, and Y10–Y14. Drug overdose deaths involving opioid analgesics include drug poisoning deaths with multiple cause-of-death codes, including T40.0, T40.1, T40.2, T40.3, T40.4, or T40.6. Drug overdose deaths involving cocaine include code T40.5; benzodiazepines include code T42.4; and psychostimulants with abuse potential include code T43.6. Deaths might involve more than one drug, so categories are not exclusive.

## Data Brief 282: Drug Overdose Deaths Among Adolescents Aged 15-19 in the United States: 1999-2015

# Data table for Figure 4. Drug overdose death rates for adolescents aged 15–19, by type of opioid drug involved: United States, 1999–2015

[Rates per 100,000 population enumerated as of April 1 for 2000 and 2010 and estimated as of July 1 for all other years]

| Year | Rate   |        |                                      |           |  |
|------|--|--------|--------------------------------------|-----------|--|
|      | Natural and semisynthetic,<br>excluding heroin | Heroin | Synthetic,<br>excluding<br>methadone | Methadone |  |
| 999  | 0.2  | 0.3    | *                                    | 0.1       |  |
| 2000 | 0.3  | 0.2    | *                                    | 0.1       |  |
| 2001 | 0.6  | 0.2    | *                                    | 0.2       |  |
| 002  | 0.5  | 0.3    | 0.1                                  | 0.5       |  |
| 003  | 0.6  | 0.3    | 0.2                                  | 0.7       |  |
| 004  | 0.7  | 0.3    | 0.2                                  | 0.9       |  |
| 005  | 0.7  | 0.3    | 0.1                                  | 0.8       |  |
| 006  | 0.8  | 0.3    | 0.3                                  | 1.0       |  |
| 007  | 0.9  | 0.4    | 0.2                                  | 1.1       |  |
| 008  | 1.1  | 0.5    | 0.3                                  | 0.9       |  |
| 009  | 1.0  | 0.4    | 0.2                                  | 0.7       |  |
| 010  | 1.2  | 0.5    | 0.2                                  | 0.6       |  |
| 011  | 1.1  | 0.6    | 0.3                                  | 0.6       |  |
| 012  | 0.8  | 0.6    | 0.2                                  | 0.4       |  |
| 013  | 0.7  | 0.9    | 0.2                                  | 0.3       |  |
| 2014 | 0.6  | 0.9    | 0.3                                  | 0.2       |  |
| 015  | 0.6  | 1.0    | 0.7                                  | 0.3       |  |

\* Figure does not meet standards of reliability or precision.

NOTES: Drug overdose deaths are those with International Classification of Diseases, Tenth Revision underlying cause-of-death codes X40–X44, X60–X64, X85, and Y10–Y14. Drug overdose deaths involving heroin include code T40.1; natural and semisynthetic opioids, excluding heroin include code T40.2; methadone include code T40.3; and synthetic opioids, excluding methadone include code T40.4. Deaths might involve more than one drug, so categories are not exclusive.