	Total		Men		Women	
Year	Number	Age-adjusted death rate ¹	Number	Age-adjusted death rate ¹	Number	Age-adjusted death rate ¹
2000	12,841	7.2	8,133	10.5	4,708	4.5
2001	13,284	7.3	8,451	10.7	4,833	4.6
2002	13,971	7.5	8,960	11.0	5,011	4.7
2003	14,627	7.7	9,433	11.4	5,194	4.8
2004	15,239	7.9	9,786	11.4	5,453	5.0
2005	16,000	8.1	10,502	11.9	5,498	4.9
2006	16,461	8.1	10,724	11.9	5,737	5.0
2007	17,058	8.3	11,293	12.2	5,765	5.0
2008	18,137	8.6	12,257	12.9	5,880	5.0
2009	19,288	8.9	13,007	13.2	6,281	5.3
2010	20,230	9.2	13,615	13.6	6,615	5.5
2011	21,540	9.5	14,585	14.0	6,955	5.6
2012	22,906	9.8	15,522	14.5	7,384	5.8
2013	23,967	10.0	16,265	14.8	7,702	5.9
2014	24,644	10.1	16,591	14.7	8,053	6.1
2015	25,694	10.2	17,377	15.0	8,317	6.1
2016	26,489	10.3	17,797	15.0	8,692	6.3

Data table for Figure 1. Age-adjusted death rates for liver cancer among adults aged 25 and over, by sex: United States, 2000-2016

¹Deaths per 100,000 U.S. standard population aged 25 and over.

NOTE: Liver cancer deaths are identified with International Classification of Diseases, 10th Revision, using underlying cause-of-death code C22.

Data Brief 314. Trends in Liver Cancer Mortality Among Adults Aged 25 and Over in the United States, 2000–2016

Data table for Figure 2. Age-adjusted death rates for liver cancer among adults aged 25 and over, by race and ethnicity: United States, 2000–2016

	Non-His	Non-Hispanic white		Non-Hispanic black		Non-Hispanic Asian or Pacific Islander		Hispanic	
Year	Number	Age-adjusted death rate ¹	Number	Age-adjusted death rate ¹	Number	Age-adjusted death rate ¹	Number	Age-adjusted death rate ¹	
2000	9,083	6.1	1,570	9.5	896	17.5	1,161	11.5	
2001	9,393	6.3	1,637	9.7	860	15.7	1,247	11.4	
2002	9,826	6.5	1,696	9.8	953	16.3	1,359	11.8	
2003	10,107	6.6	1,838	10.3	1,045	17.3	1,499	12.2	
2004	10,501	6.8	1,967	10.8	1,038	16.0	1,561	12.1	
2005	10,907	7.0	2,074	10.8	1,081	16.2	1,793	13.2	
2006	11,269	7.0	2,150	10.9	1,087	15.2	1,798	12.7	
2007	11,674	7.2	2,234	11.1	1,106	14.6	1,877	12.5	
2008	12,281	7.4	2,436	11.6	1,197	15.4	2,048	13.0	
2009	12,954	7.7	2,681	12.2	1,222	14.9	2,248	13.6	
2010	13,602	8.0	2,776	12.3	1,306	15.5	2,317	13.5	
2011	14,369	8.2	2,991	12.9	1,419	15.6	2,551	13.7	
2012	15,298	8.6	3,123	13.1	1,473	15.4	2,762	14.2	
2013	15,853	8.7	3,332	13.3	1,516	14.9	2,978	14.4	
2014	16,323	8.8	3,422	13.4	1,608	14.9	3,011	13.9	
2015	17,072	9.0	3,507	13.3	1,614	14.0	3,173	14.0	
2016	17,374	9.0	3,687	13.6	1,622	13.6	3,458	14.6	

 $^{1}\mbox{Deaths}$ per 100,000 U.S. standard population aged 25 and over.

NOTE: Liver cancer deaths are identified with International Classification of Diseases, 10th Revision, using underlying cause-of-death code C22.

	25–44 years		45–54 years		55–64 years		65–74 years		75 years and over	
Year	Number	Death rate ¹	Number	Death rate ¹						
2000	515	0.6	1,686	4.5	2,256	9.3	3,439	18.7	4,945	29.8
2001	476	0.6	1,868	4.7	2,351	9.4	3,440	18.7	5,149	30.5
2002	519	0.6	1,976	4.9	2,558	9.6	3,581	19.5	5,337	31.1
2003	484	0.6	2,223	5.4	2,765	9.9	3,630	19.6	5,525	31.8
2004	464	0.6	2,348	5.6	3,123	10.7	3,670	19.7	5,634	32.1
2005	469	0.6	2,515	5.9	3,438	11.2	3,747	19.8	5,831	32.8
2006	448	0.5	2,406	5.6	3,842	12.0	3,830	19.9	5,935	33.0
2007	458	0.6	2,503	5.7	4,181	12.6	3,884	19.7	6,032	33.3
2008	416	0.5	2,480	5.6	4,737	13.9	4,105	20.0	6,399	35.0
2009	427	0.5	2,621	5.8	5,292	14.9	4,470	21.1	6,478	35.2
2010	405	0.5	2,493	5.5	5,819	16.0	4,814	22.2	6,699	36.1
2011	413	0.5	2,547	5.7	6,616	17.4	4,989	22.2	6,975	36.9
2012	432	0.5	2,432	5.5	7,159	18.6	5,723	23.9	7,160	37.4
2013	452	0.5	2,216	5.1	7,632	19.4	6,211	24.6	7,456	38.3
2014	386	0.5	2,163	5.0	7,772	19.4	6,606	25.0	7,717	38.9
2015	435	0.5	2,090	4.8	7,979	19.5	7,289	26.5	7,901	39.1
2016	443	0.5	1,878	4.4	8,075	19.5	7,813	27.3	8,280	40.2

Data table for Figure 3. Death rates for liver cancer among adults aged 25 and over, by selected age groups: United States, 2000–2016

¹Deaths per 100,000 U.S. standard population.

NOTE: Liver cancer deaths are identified with International Classification of Diseases, 10th Revision, using underlying cause-of-death code C22.

Data table for Figure 4. Age-adjusted death rates for liver cancer among adults aged 25
and over, by state: United States, 2016

Area	Number	Age-adjusted death rate ¹ 10.3		
United States	26,489			
Alabama	464	11.6		
Alaska	51	11.7		
Arizona	553	9.7		
Arkansas	272	11.2		
California	3,433	12.2		
Colorado	346	8.7		
Connecticut	249	8.0		
Delaware	88	10.4		
District of Columbia	74	16.8		
Florida	1,937	9.9		
Georgia	714	9.7		
Hawaii	158	12.7		
Idaho	134	10.2		
Illinois	1,006	10.2		
Indiana	476	9.0		
lowa	216	8.2		
Kansas	224	9.5		
Kentucky	367	10.3		
Louisiana	500	13.8		
Maine	96	7.4		
Maryland	451	9.4		
Massachusetts	586	10.4		
Michigan	811	9.7		
Minnesota	399	9.0		
Mississippi	294	12.4		
Missouri	463	9.3		
Montana	73	7.7		
Nebraska	117	7.8		
Nevada	215	9.7		
New Hampshire	117	9.9		
New Jersey	661	9.2		
New Mexico	211	12.4		
New York	1,533	9.7		
North Carolina	861	10.5		
North Dakota	47	8.8		
Ohio	951	9.6		
Oklahoma	320	10.4		
Oregon	382	11.0		
Pennsylvania	1,141	10.0		
Rhode Island	97	10.7		
South Carolina	462	10.8		
South Dakota	72	10.3		
Tennessee	561	10.4		
Texas	2,279	12.3		
Utah	131	7.8		
Vermont	36	6.0		
Virginia	639	9.9		
Washington	577	10.1		
West Virginia	148	8.8		
Wisconsin	440	9.2		
Wyoming	56	11.3		

¹Deaths per 100,000 U.S. standard population aged 25 and over.

NOTE: Liver cancer deaths are identified with International Classification of Diseases, 10th Revision, using underlying cause-of-death code C22.