

A revised mortality data file for 2014 was released on April 3, 2017 to include corrections from Tennessee and Massachusetts affecting 125 records previously coded to Accidental discharge of firearms (W32-W34). After the revision, 50% of the changed records are now classified as Assault (homicide) by discharge of firearms (*U01.4 and X93-X95); 42% as Intentional self-harm (suicide) by discharge of firearms (X72-X74); 4% as Discharge of firearms, undetermined intent (Y22-Y24); and the remaining 4% to various other causes. For more information, please see "[Deaths: Final Data for 2014](#)." This table was updated on May 3, 2017 to correct errors. The shaded areas in this table indicate the changes that were made.

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex:
United States, 2014

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
All races, both sexes, all ages				
...	All causes	2,626,418	100.0	823.7
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	614,348	23.4	192.7
2	Malignant neoplasms (C00-C97)	591,700	22.5	185.6
3	Chronic lower respiratory diseases (J40-J47)	147,101	5.6	46.1
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	135,928	5.2	42.6
5	Cerebrovascular diseases (I60-I69)	133,103	5.1	41.7
6	Alzheimer's disease (G30)	93,541	3.6	29.3
7	Diabetes mellitus (E10-E14)	76,488	2.9	24.0
8	Influenza and pneumonia (J09-J18)	55,227	2.1	17.3
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	48,146	1.8	15.1
10	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	42,826	1.6	13.4
11	Septicemia (A40-A41)	38,940	1.5	12.2
12	Chronic liver disease and cirrhosis (K70,K73-K74)	38,170	1.5	12.0
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	30,221	1.2	9.5
14	Parkinson's disease (G20-G21)	26,150	1.0	8.2
15	Pneumonitis due to solids and liquids (J69)	18,792	0.7	5.9
...	All other causes (residual)	535,737	20.4	168.0
All races, both sexes, 1-4 years				
...	All causes	3,830	100.0	24.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,216	31.7	7.6
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	399	10.4	2.5
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	364	9.5	2.3
4	Malignant neoplasms (C00-C97)	321	8.4	2.0
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	149	3.9	0.9
6	Influenza and pneumonia (J09-J18)	109	2.8	0.7
7	Septicemia (A40-A41)	53	1.4	0.3
7	Chronic lower respiratory diseases (J40-J47)	53	1.4	0.3
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	38	1.0	0.2
9	Certain conditions originating in the perinatal period (P00-P96)	38	1.0	0.2
11	Cerebrovascular diseases (I60-I69)	33	0.9	0.2
12	Complications of medical and surgical care (Y40-Y84,Y88)	17	0.4	*
13	Meningitis (G00,G03)	16	0.4	*
13	Acute bronchitis and bronchiolitis (J20-J21)	16	0.4	*
15	Anemias (D50-D64)	15	0.4	*
...	All other causes (residual)	993	25.9	6.2

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2014.

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex:
United States, 2014

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
All races, both sexes, 5-14 years				
...	All causes	5,250	100.0	12.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,480	28.2	3.6
2	Malignant neoplasms (C00-C97)	852	16.2	2.1
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	428	8.2	1.0
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	348	6.6	0.8
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	279	5.3	0.7
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	191	3.6	0.5
7	Chronic lower respiratory diseases (J40-J47)	139	2.6	0.3
8	Influenza and pneumonia (J09-J18)	98	1.9	0.2
9	Cerebrovascular diseases (I60-I69)	88	1.7	0.2
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	74	1.4	0.2
11	Septicemia (A40-A41)	63	1.2	0.2
12	Anemias (D50-D64)	34	0.6	0.1
13	Certain conditions originating in the perinatal period (P00-P96)	32	0.6	0.1
14	Diabetes mellitus (E10-E14)	26	0.5	0.1
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	13	0.2	*
15	Complications of medical and surgical care (Y40-Y84,Y88)	13	0.2	*
...	All other causes (residual)	1,092	20.8	2.7
All races, both sexes, 15-24 years				
...	All causes	28,791	100.0	65.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	11,797	41.0	26.8
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	5,090	17.7	11.6
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	4,171	14.5	9.5
4	Malignant neoplasms (C00-C97)	1,569	5.4	3.6
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	953	3.3	2.2
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	377	1.3	0.9
7	Influenza and pneumonia (J09-J18)	199	0.7	0.5
8	Diabetes mellitus (E10-E14)	181	0.6	0.4
9	Chronic lower respiratory diseases (J40-J47)	178	0.6	0.4
10	Cerebrovascular diseases (I60-I69)	177	0.6	0.4
11	Pregnancy, childbirth and the puerperium (O00-O99)	163	0.6	0.4
12	Septicemia (A40-A41)	126	0.4	0.3
13	Human immunodeficiency virus (HIV) disease (B20-B24)	117	0.4	0.3
14	Anemias (D50-D64)	85	0.3	0.2
15	Legal intervention (Y35,Y89.0)	82	0.3	0.2
...	All other causes (residual)	3,526	12.2	8.0

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2014.

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex:
United States, 2014

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
All races, both sexes, 25-34 years				
...	All causes	47,177	100.0	108.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	17,335	36.7	39.8
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	6,574	13.9	15.1
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	4,173	8.8	9.6
4	Malignant neoplasms (C00-C97)	3,624	7.7	8.3
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,341	7.1	7.7
6	Chronic liver disease and cirrhosis (K70,K73-K74)	725	1.5	1.7
7	Diabetes mellitus (E10-E14)	709	1.5	1.6
8	Human immunodeficiency virus (HIV) disease (B20-B24)	583	1.2	1.3
9	Cerebrovascular diseases (I60-I69)	579	1.2	1.3
10	Influenza and pneumonia (J09-J18)	549	1.2	1.3
11	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	428	0.9	1.0
12	Pregnancy, childbirth and the puerperium (O00-O99)	405	0.9	0.9
13	Chronic lower respiratory diseases (J40-J47)	367	0.8	0.8
14	Septicemia (A40-A41)	359	0.8	0.8
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	236	0.5	0.5
...	All other causes (residual)	7,190	15.2	16.5
All races, both sexes, 35-44 years				
...	All causes	70,996	100.0	175.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	16,031	22.6	39.6
2	Malignant neoplasms (C00-C97)	11,267	15.9	27.8
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	10,368	14.6	25.6
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	6,715	9.5	16.6
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	2,596	3.7	6.4
6	Chronic liver disease and cirrhosis (K70,K73-K74)	2,582	3.6	6.4
7	Diabetes mellitus (E10-E14)	1,999	2.8	4.9
8	Cerebrovascular diseases (I60-I69)	1,745	2.5	4.3
9	Human immunodeficiency virus (HIV) disease (B20-B24)	1,174	1.7	2.9
10	Influenza and pneumonia (J09-J18)	1,125	1.6	2.8
11	Septicemia (A40-A41)	832	1.2	2.1
12	Chronic lower respiratory diseases (J40-J47)	754	1.1	1.9
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	673	0.9	1.7
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	439	0.6	1.1
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	409	0.6	1.0
...	All other causes (residual)	12,287	17.3	30.3

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2014.

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex:
United States, 2014

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
All races, both sexes, 45-54 years				
...	All causes	175,917	100.0	404.8
1	Malignant neoplasms (C00-C97)	44,834	25.5	103.2
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	34,791	19.8	80.1
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	20,593	11.7	47.4
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	8,774	5.0	20.2
5	Chronic liver disease and cirrhosis (K70,K73-K74)	8,627	4.9	19.9
6	Diabetes mellitus (E10-E14)	6,062	3.4	13.9
7	Cerebrovascular diseases (I60-I69)	5,349	3.0	12.3
8	Chronic lower respiratory diseases (J40-J47)	4,402	2.5	10.1
9	Influenza and pneumonia (J09-J18)	2,731	1.6	6.3
10	Septicemia (A40-A41)	2,514	1.4	5.8
11	Human immunodeficiency virus (HIV) disease (B20-B24)	2,234	1.3	5.1
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2,032	1.2	4.7
13	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,952	1.1	4.5
14	Viral hepatitis (B15-B19)	1,774	1.0	4.1
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,428	0.8	3.3
...	All other causes (residual)	27,820	15.8	64.0
All races, both sexes, 55-64 years				
...	All causes	348,808	100.0	870.3
1	Malignant neoplasms (C00-C97)	115,282	33.1	287.6
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	74,473	21.4	185.8
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	18,014	5.2	44.9
4	Chronic lower respiratory diseases (J40-J47)	16,492	4.7	41.2
5	Diabetes mellitus (E10-E14)	13,342	3.8	33.3
6	Chronic liver disease and cirrhosis (K70,K73-K74)	12,792	3.7	31.9
7	Cerebrovascular diseases (I60-I69)	11,727	3.4	29.3
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	7,539	2.2	18.8
9	Septicemia (A40-A41)	5,709	1.6	14.2
10	Influenza and pneumonia (J09-J18)	5,390	1.5	13.4
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	5,063	1.5	12.6
12	Viral hepatitis (B15-B19)	3,930	1.1	9.8
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	3,362	1.0	8.4
14	Human immunodeficiency virus (HIV) disease (B20-B24)	1,810	0.5	4.5
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1,610	0.5	4.0
...	All other causes (residual)	52,273	15.0	130.4

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2014.

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex:
United States, 2014

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
All races, both sexes, 65-74 years				
...	All causes	471,541	100.0	1,786.3
1	Malignant neoplasms (C00-C97)	159,209	33.8	603.1
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	101,683	21.6	385.2
3	Chronic lower respiratory diseases (J40-J47)	35,617	7.6	134.9
4	Cerebrovascular diseases (I60-I69)	19,663	4.2	74.5
5	Diabetes mellitus (E10-E14)	18,204	3.9	69.0
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	11,901	2.5	45.1
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	9,060	1.9	34.3
8	Septicemia (A40-A41)	8,220	1.7	31.1
9	Influenza and pneumonia (J09-J18)	7,861	1.7	29.8
10	Chronic liver disease and cirrhosis (K70,K73-K74)	7,809	1.7	29.6
11	Alzheimer's disease (G30)	5,170	1.1	19.6
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	4,464	0.9	16.9
13	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	4,115	0.9	15.6
14	Parkinson's disease (G20-G21)	3,431	0.7	13.0
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2,981	0.6	11.3
...	All other causes (residual)	72,153	15.3	273.3
All races, both sexes, 75-84 years				
...	All causes	624,504	100.0	4,564.2
1	Malignant neoplasms (C00-C97)	154,053	24.7	1,125.9
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	146,426	23.4	1,070.2
3	Chronic lower respiratory diseases (J40-J47)	47,758	7.6	349.0
4	Cerebrovascular diseases (I60-I69)	36,353	5.8	265.7
5	Alzheimer's disease (G30)	25,393	4.1	185.6
6	Diabetes mellitus (E10-E14)	19,407	3.1	141.8
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	14,868	2.4	108.7
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	13,497	2.2	98.6
9	Influenza and pneumonia (J09-J18)	13,193	2.1	96.4
10	Parkinson's disease (G20-G21)	10,835	1.7	79.2
11	Septicemia (A40-A41)	10,004	1.6	73.1
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	7,023	1.1	51.3
13	Pneumonitis due to solids and liquids (J69)	5,015	0.8	36.7
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	4,791	0.8	35.0
15	Chronic liver disease and cirrhosis (K70,K73-K74)	4,157	0.7	30.4
...	All other causes (residual)	111,731	17.9	816.6

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2014.

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex:
United States, 2014

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
All races, both sexes, 85 years and over				
...	All causes	826,226	100.0	13,407.9
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	241,613	29.2	3,920.9
2	Malignant neoplasms (C00-C97)	100,624	12.2	1,632.9
3	Alzheimer's disease (G30)	62,041	7.5	1,006.8
4	Cerebrovascular diseases (I60-I69)	57,292	6.9	929.7
5	Chronic lower respiratory diseases (J40-J47)	41,318	5.0	670.5
6	Influenza and pneumonia (J09-J18)	23,782	2.9	385.9
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	21,513	2.6	349.1
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	17,400	2.1	282.4
9	Diabetes mellitus (E10-E14)	16,550	2.0	268.6
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	13,374	1.6	217.0
11	Parkinson's disease (G20-G21)	11,216	1.4	182.0
12	Septicemia (A40-A41)	10,900	1.3	176.9
13	Pneumonitis due to solids and liquids (J69)	9,108	1.1	147.8
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	5,381	0.7	87.3
15	Atherosclerosis (I70)	3,573	0.4	58.0
...	All other causes (residual)	190,541	23.1	3,092.1
All races, male, all ages				
...	All causes	1,328,241	100.0	846.4
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	325,077	24.5	207.1
2	Malignant neoplasms (C00-C97)	311,296	23.4	198.4
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	85,340	6.4	54.4
4	Chronic lower respiratory diseases (J40-J47)	69,456	5.2	44.3
5	Cerebrovascular diseases (I60-I69)	55,471	4.2	35.3
6	Diabetes mellitus (E10-E14)	41,111	3.1	26.2
7	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	33,162	2.5	21.1
8	Alzheimer's disease (G30)	28,362	2.1	18.1
9	Influenza and pneumonia (J09-J18)	26,586	2.0	16.9
10	Chronic liver disease and cirrhosis (K70,K73-K74)	24,584	1.9	15.7
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	24,436	1.8	15.6
12	Septicemia (A40-A41)	18,333	1.4	11.7
13	Parkinson's disease (G20-G21)	15,681	1.2	10.0
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	12,989	1.0	8.3
15	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	12,546	0.9	8.0
...	All other causes (residual)	243,811	18.4	155.4

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2014.

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex:
United States, 2014

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
All races, male, 1-4 years				
...	All causes	2,172	100.0	26.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	744	34.3	9.1
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	219	10.1	2.7
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	196	9.0	2.4
4	Malignant neoplasms (C00-C97)	182	8.4	2.2
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	76	3.5	0.9
6	Influenza and pneumonia (J09-J18)	57	2.6	0.7
7	Chronic lower respiratory diseases (J40-J47)	30	1.4	0.4
8	Certain conditions originating in the perinatal period (P00-P96)	24	1.1	0.3
9	Septicemia (A40-A41)	23	1.1	0.3
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	19	0.9	*
10	Cerebrovascular diseases (I60-I69)	19	0.9	*
12	Anemias (D50-D64)	10	0.5	*
13	Meningitis (G00,G03)	7	0.3	*
13	Acute bronchitis and bronchiolitis (J20-J21)	7	0.3	*
13	Complications of medical and surgical care (Y40-Y84,Y88)	7	0.3	*
...	All other causes (residual)	552	25.4	6.8
All races, male, 5-14 years				
...	All causes	3,128	100.0	14.9
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	968	30.9	4.6
2	Malignant neoplasms (C00-C97)	488	15.6	2.3
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	277	8.9	1.3
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	185	5.9	0.9
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	166	5.3	0.8
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	108	3.5	0.5
7	Chronic lower respiratory diseases (J40-J47)	90	2.9	0.4
8	Influenza and pneumonia (J09-J18)	50	1.6	0.2
9	Cerebrovascular diseases (I60-I69)	47	1.5	0.2
10	Septicemia (A40-A41)	34	1.1	0.2
11	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	33	1.1	0.2
12	Anemias (D50-D64)	17	0.5	*
13	Certain conditions originating in the perinatal period (P00-P96)	16	0.5	*
14	Diabetes mellitus (E10-E14)	15	0.5	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	9	0.3	*
...	All other causes (residual)	625	20.0	3.0

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2014.

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex:
United States, 2014

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
All races, male, 15-24 years				
...	All causes	21,117	100.0	93.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	8,765	41.5	38.9
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	4,100	19.4	18.2
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	3,624	17.2	16.1
4	Malignant neoplasms (C00-C97)	952	4.5	4.2
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	616	2.9	2.7
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	211	1.0	0.9
7	Influenza and pneumonia (J09-J18)	116	0.5	0.5
7	Chronic lower respiratory diseases (J40-J47)	116	0.5	0.5
9	Diabetes mellitus (E10-E14)	115	0.5	0.5
10	Cerebrovascular diseases (I60-I69)	102	0.5	0.5
11	Legal intervention (Y35,Y89.0)	78	0.4	0.3
12	Human immunodeficiency virus (HIV) disease (B20-B24)	76	0.4	0.3
13	Septicemia (A40-A41)	65	0.3	0.3
14	Anemias (D50-D64)	41	0.2	0.2
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	39	0.2	0.2
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	39	0.2	0.2
...	All other causes (residual)	2,062	9.8	9.2
All races, male, 25-34 years				
...	All causes	32,697	100.0	148.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	12,822	39.2	58.4
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	5,226	16.0	23.8
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	3,490	10.7	15.9
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,266	6.9	10.3
5	Malignant neoplasms (C00-C97)	1,794	5.5	8.2
6	Chronic liver disease and cirrhosis (K70,K73-K74)	437	1.3	2.0
7	Diabetes mellitus (E10-E14)	432	1.3	2.0
8	Human immunodeficiency virus (HIV) disease (B20-B24)	413	1.3	1.9
9	Cerebrovascular diseases (I60-I69)	331	1.0	1.5
10	Influenza and pneumonia (J09-J18)	288	0.9	1.3
11	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	254	0.8	1.2
12	Chronic lower respiratory diseases (J40-J47)	213	0.7	1.0
13	Septicemia (A40-A41)	185	0.6	0.8
14	Legal intervention (Y35,Y89.0)	160	0.5	0.7
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	129	0.4	0.6
...	All other causes (residual)	4,257	13.0	19.4

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2014.

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex:
United States, 2014

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
All races, male, 35-44 years				
...	All causes	43,693	100.0	216.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	11,052	25.3	54.8
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	7,021	16.1	34.8
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	5,044	11.5	25.0
4	Malignant neoplasms (C00-C97)	4,829	11.1	24.0
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	2,067	4.7	10.3
6	Chronic liver disease and cirrhosis (K70,K73-K74)	1,617	3.7	8.0
7	Diabetes mellitus (E10-E14)	1,211	2.8	6.0
8	Cerebrovascular diseases (I60-I69)	1,001	2.3	5.0
9	Human immunodeficiency virus (HIV) disease (B20-B24)	774	1.8	3.8
10	Influenza and pneumonia (J09-J18)	616	1.4	3.1
11	Septicemia (A40-A41)	419	1.0	2.1
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	399	0.9	2.0
13	Chronic lower respiratory diseases (J40-J47)	359	0.8	1.8
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	299	0.7	1.5
15	Aortic aneurysm and dissection (I71)	251	0.6	1.2
...	All other causes (residual)	6,734	15.4	33.4
All races, male, 45-54 years				
...	All causes	106,377	100.0	496.5
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	24,316	22.9	113.5
2	Malignant neoplasms (C00-C97)	22,053	20.7	102.9
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	13,826	13.0	64.5
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	6,427	6.0	30.0
5	Chronic liver disease and cirrhosis (K70,K73-K74)	5,727	5.4	26.7
6	Diabetes mellitus (E10-E14)	3,795	3.6	17.7
7	Cerebrovascular diseases (I60-I69)	2,995	2.8	14.0
8	Chronic lower respiratory diseases (J40-J47)	2,002	1.9	9.3
9	Human immunodeficiency virus (HIV) disease (B20-B24)	1,662	1.6	7.8
10	Influenza and pneumonia (J09-J18)	1,488	1.4	6.9
11	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,473	1.4	6.9
12	Septicemia (A40-A41)	1,343	1.3	6.3
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,231	1.2	5.7
14	Viral hepatitis (B15-B19)	1,213	1.1	5.7
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	920	0.9	4.3
...	All other causes (residual)	15,906	15.0	74.2

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2014.

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex:
United States, 2014

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
All races, male, 55-64 years				
...	All causes	212,198	100.0	1,098.2
1	Malignant neoplasms (C00-C97)	63,825	30.1	330.3
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	51,795	24.4	268.1
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	12,162	5.7	62.9
4	Chronic liver disease and cirrhosis (K70,K73-K74)	8,871	4.2	45.9
5	Chronic lower respiratory diseases (J40-J47)	8,549	4.0	44.2
6	Diabetes mellitus (E10-E14)	8,196	3.9	42.4
7	Cerebrovascular diseases (I60-I69)	6,796	3.2	35.2
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	5,689	2.7	29.4
9	Influenza and pneumonia (J09-J18)	3,076	1.4	15.9
10	Septicemia (A40-A41)	3,007	1.4	15.6
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2,918	1.4	15.1
12	Viral hepatitis (B15-B19)	2,770	1.3	14.3
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	2,079	1.0	10.8
14	Human immunodeficiency virus (HIV) disease (B20-B24)	1,389	0.7	7.2
15	Aortic aneurysm and dissection (I71)	992	0.5	5.1
...	All other causes (residual)	30,084	14.2	155.7
All races, male, 65-74 years				
...	All causes	268,648	100.0	2,175.5
1	Malignant neoplasms (C00-C97)	87,914	32.7	711.9
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	64,917	24.2	525.7
3	Chronic lower respiratory diseases (J40-J47)	18,096	6.7	146.5
4	Diabetes mellitus (E10-E14)	10,599	3.9	85.8
5	Cerebrovascular diseases (I60-I69)	10,504	3.9	85.1
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	7,515	2.8	60.9
7	Chronic liver disease and cirrhosis (K70,K73-K74)	4,999	1.9	40.5
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	4,827	1.8	39.1
9	Influenza and pneumonia (J09-J18)	4,368	1.6	35.4
10	Septicemia (A40-A41)	4,264	1.6	34.5
11	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	3,286	1.2	26.6
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	2,475	0.9	20.0
13	Parkinson's disease (G20-G21)	2,271	0.8	18.4
14	Alzheimer's disease (G30)	2,206	0.8	17.9
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1,771	0.7	14.3
...	All other causes (residual)	38,636	14.4	312.9

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2014.

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex:
United States, 2014

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
All races, male, 75-84 years				
...	All causes	316,430	100.0	5,369.2
1	Malignant neoplasms (C00-C97)	81,768	25.8	1,387.5
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	79,844	25.2	1,354.8
3	Chronic lower respiratory diseases (J40-J47)	22,958	7.3	389.6
4	Cerebrovascular diseases (I60-I69)	16,075	5.1	272.8
5	Diabetes mellitus (E10-E14)	9,998	3.2	169.6
6	Alzheimer's disease (G30)	9,425	3.0	159.9
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	8,134	2.6	138.0
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	6,994	2.2	118.7
9	Parkinson's disease (G20-G21)	6,917	2.2	117.4
10	Influenza and pneumonia (J09-J18)	6,756	2.1	114.6
11	Septicemia (A40-A41)	4,704	1.5	79.8
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	3,078	1.0	52.2
13	Pneumonitis due to solids and liquids (J69)	2,911	0.9	49.4
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2,714	0.9	46.1
15	Chronic liver disease and cirrhosis (K70,K73-K74)	2,238	0.7	38.0
...	All other causes (residual)	51,916	16.4	880.9
All races, male, 85 years and over				
...	All causes	308,785	100.0	14,642.2
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	93,917	30.4	4,453.4
2	Malignant neoplasms (C00-C97)	47,458	15.4	2,250.4
3	Cerebrovascular diseases (I60-I69)	17,546	5.7	832.0
4	Chronic lower respiratory diseases (J40-J47)	17,024	5.5	807.3
5	Alzheimer's disease (G30)	16,334	5.3	774.5
6	Influenza and pneumonia (J09-J18)	9,680	3.1	459.0
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	8,674	2.8	411.3
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	7,837	2.5	371.6
9	Diabetes mellitus (E10-E14)	6,744	2.2	319.8
10	Parkinson's disease (G20-G21)	6,052	2.0	287.0
11	Pneumonitis due to solids and liquids (J69)	4,638	1.5	219.9
12	Septicemia (A40-A41)	4,201	1.4	199.2
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	4,048	1.3	192.0
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2,422	0.8	114.8
15	Atherosclerosis (I70)	1,164	0.4	55.2
...	All other causes (residual)	61,046	19.8	2,894.7

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2014.

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex:
United States, 2014

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
All races, female, all ages				
...	All causes	1,298,177	100.0	801.7
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	289,271	22.3	178.6
2	Malignant neoplasms (C00-C97)	280,404	21.6	173.2
3	Chronic lower respiratory diseases (J40-J47)	77,645	6.0	48.0
4	Cerebrovascular diseases (I60-I69)	77,632	6.0	47.9
5	Alzheimer's disease (G30)	65,179	5.0	40.3
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	50,588	3.9	31.2
7	Diabetes mellitus (E10-E14)	35,377	2.7	21.8
8	Influenza and pneumonia (J09-J18)	28,641	2.2	17.7
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	23,710	1.8	14.6
10	Septicemia (A40-A41)	20,607	1.6	12.7
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	17,232	1.3	10.6
12	Chronic liver disease and cirrhosis (K70,K73-K74)	13,586	1.0	8.4
13	Parkinson's disease (G20-G21)	10,469	0.8	6.5
14	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	9,664	0.7	6.0
15	Pneumonitis due to solids and liquids (J69)	8,434	0.6	5.2
...	All other causes (residual)	289,738	22.3	178.9
All races, female, 1-4 years				
...	All causes	1,658	100.0	21.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	472	28.5	6.1
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	180	10.9	2.3
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	168	10.1	2.2
4	Malignant neoplasms (C00-C97)	139	8.4	1.8
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	73	4.4	0.9
6	Influenza and pneumonia (J09-J18)	52	3.1	0.7
7	Septicemia (A40-A41)	30	1.8	0.4
8	Chronic lower respiratory diseases (J40-J47)	23	1.4	0.3
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	19	1.1	*
10	Cerebrovascular diseases (I60-I69)	14	0.8	*
10	Certain conditions originating in the perinatal period (P00-P96)	14	0.8	*
12	Complications of medical and surgical care (Y40-Y84,Y88)	10	0.6	*
13	Meningitis (G00,G03)	9	0.5	*
13	Acute bronchitis and bronchiolitis (J20-J21)	9	0.5	*
15	Anemias (D50-D64)	5	0.3	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	5	0.3	*
...	All other causes (residual)	436	26.3	5.6

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2014.

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex:
United States, 2014

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
All races, female, 5-14 years				
...	All causes	2,122	100.0	10.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	512	24.1	2.5
2	Malignant neoplasms (C00-C97)	364	17.2	1.8
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	163	7.7	0.8
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	151	7.1	0.7
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	113	5.3	0.6
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	83	3.9	0.4
7	Chronic lower respiratory diseases (J40-J47)	49	2.3	0.2
8	Influenza and pneumonia (J09-J18)	48	2.3	0.2
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	41	1.9	0.2
9	Cerebrovascular diseases (I60-I69)	41	1.9	0.2
11	Septicemia (A40-A41)	29	1.4	0.1
12	Anemias (D50-D64)	17	0.8	*
13	Certain conditions originating in the perinatal period (P00-P96)	16	0.8	*
14	Diabetes mellitus (E10-E14)	11	0.5	*
15	Complications of medical and surgical care (Y40-Y84,Y88)	6	0.3	*
...	All other causes (residual)	478	22.5	2.4
All races, female, 15-24 years				
...	All causes	7,674	100.0	35.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,032	39.5	14.1
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	990	12.9	4.6
3	Malignant neoplasms (C00-C97)	617	8.0	2.9
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	547	7.1	2.5
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	337	4.4	1.6
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	166	2.2	0.8
7	Pregnancy, childbirth and the puerperium (O00-O99)	163	2.1	0.8
8	Influenza and pneumonia (J09-J18)	83	1.1	0.4
9	Cerebrovascular diseases (I60-I69)	75	1.0	0.3
10	Diabetes mellitus (E10-E14)	66	0.9	0.3
11	Chronic lower respiratory diseases (J40-J47)	62	0.8	0.3
12	Septicemia (A40-A41)	61	0.8	0.3
13	Anemias (D50-D64)	44	0.6	0.2
14	Human immunodeficiency virus (HIV) disease (B20-B24)	41	0.5	0.2
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	37	0.5	0.2
...	All other causes (residual)	1,353	17.6	6.3

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2014.

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex:
United States, 2014

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
All races, female, 25-34 years				
...	All causes	14,480	100.0	67.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	4,513	31.2	20.9
2	Malignant neoplasms (C00-C97)	1,830	12.6	8.5
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,348	9.3	6.3
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,075	7.4	5.0
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	683	4.7	3.2
6	Pregnancy, childbirth and the puerperium (O00-O99)	405	2.8	1.9
7	Chronic liver disease and cirrhosis (K70,K73-K74)	288	2.0	1.3
8	Diabetes mellitus (E10-E14)	277	1.9	1.3
9	Influenza and pneumonia (J09-J18)	261	1.8	1.2
10	Cerebrovascular diseases (I60-I69)	248	1.7	1.2
11	Septicemia (A40-A41)	174	1.2	0.8
11	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	174	1.2	0.8
13	Human immunodeficiency virus (HIV) disease (B20-B24)	170	1.2	0.8
14	Chronic lower respiratory diseases (J40-J47)	154	1.1	0.7
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	107	0.7	0.5
...	All other causes (residual)	2,773	19.2	12.9
All races, female, 35-44 years				
...	All causes	27,303	100.0	134.1
1	Malignant neoplasms (C00-C97)	6,438	23.6	31.6
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	4,979	18.2	24.5
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,347	12.3	16.4
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,671	6.1	8.2
5	Chronic liver disease and cirrhosis (K70,K73-K74)	965	3.5	4.7
6	Diabetes mellitus (E10-E14)	788	2.9	3.9
7	Cerebrovascular diseases (I60-I69)	744	2.7	3.7
8	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	529	1.9	2.6
9	Influenza and pneumonia (J09-J18)	509	1.9	2.5
10	Septicemia (A40-A41)	413	1.5	2.0
11	Human immunodeficiency virus (HIV) disease (B20-B24)	400	1.5	2.0
12	Chronic lower respiratory diseases (J40-J47)	395	1.4	1.9
13	Pregnancy, childbirth and the puerperium (O00-O99)	350	1.3	1.7
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	274	1.0	1.3
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	191	0.7	0.9
...	All other causes (residual)	5,310	19.4	26.1

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2014.

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex:
United States, 2014

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
All races, female, 45-54 years				
...	All causes	69,540	100.0	315.6
1	Malignant neoplasms (C00-C97)	22,781	32.8	103.4
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	10,475	15.1	47.5
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	6,767	9.7	30.7
4	Chronic liver disease and cirrhosis (K70,K73-K74)	2,900	4.2	13.2
5	Chronic lower respiratory diseases (J40-J47)	2,400	3.5	10.9
6	Cerebrovascular diseases (I60-I69)	2,354	3.4	10.7
7	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,347	3.4	10.7
8	Diabetes mellitus (E10-E14)	2,267	3.3	10.3
9	Influenza and pneumonia (J09-J18)	1,243	1.8	5.6
10	Septicemia (A40-A41)	1,171	1.7	5.3
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	801	1.2	3.6
12	Human immunodeficiency virus (HIV) disease (B20-B24)	572	0.8	2.6
13	Viral hepatitis (B15-B19)	561	0.8	2.5
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	508	0.7	2.3
15	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	479	0.7	2.2
...	All other causes (residual)	11,914	17.1	54.1
All races, female, 55-64 years				
...	All causes	136,610	100.0	658.2
1	Malignant neoplasms (C00-C97)	51,457	37.7	247.9
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	22,678	16.6	109.3
3	Chronic lower respiratory diseases (J40-J47)	7,943	5.8	38.3
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	5,852	4.3	28.2
5	Diabetes mellitus (E10-E14)	5,146	3.8	24.8
6	Cerebrovascular diseases (I60-I69)	4,931	3.6	23.8
7	Chronic liver disease and cirrhosis (K70,K73-K74)	3,921	2.9	18.9
8	Septicemia (A40-A41)	2,702	2.0	13.0
9	Influenza and pneumonia (J09-J18)	2,314	1.7	11.1
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2,145	1.6	10.3
11	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,850	1.4	8.9
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,283	0.9	6.2
13	Viral hepatitis (B15-B19)	1,160	0.8	5.6
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	665	0.5	3.2
15	Pneumonitis due to solids and liquids (J69)	517	0.4	2.5
...	All other causes (residual)	22,046	16.1	106.2

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2014.

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex:
United States, 2014

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
All races, female, 65-74 years				
...	All causes	202,893	100.0	1,444.2
1	Malignant neoplasms (C00-C97)	71,295	35.1	507.5
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	36,766	18.1	261.7
3	Chronic lower respiratory diseases (J40-J47)	17,521	8.6	124.7
4	Cerebrovascular diseases (I60-I69)	9,159	4.5	65.2
5	Diabetes mellitus (E10-E14)	7,605	3.7	54.1
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	4,386	2.2	31.2
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	4,233	2.1	30.1
8	Septicemia (A40-A41)	3,956	1.9	28.2
9	Influenza and pneumonia (J09-J18)	3,493	1.7	24.9
10	Alzheimer's disease (G30)	2,964	1.5	21.1
11	Chronic liver disease and cirrhosis (K70,K73-K74)	2,810	1.4	20.0
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,989	1.0	14.2
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1,210	0.6	8.6
14	Parkinson's disease (G20-G21)	1,160	0.6	8.3
15	Pneumonitis due to solids and liquids (J69)	996	0.5	7.1
...	All other causes (residual)	33,350	16.4	237.4
All races, female, 75-84 years				
...	All causes	308,074	100.0	3,955.1
1	Malignant neoplasms (C00-C97)	72,285	23.5	928.0
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	66,582	21.6	854.8
3	Chronic lower respiratory diseases (J40-J47)	24,800	8.1	318.4
4	Cerebrovascular diseases (I60-I69)	20,278	6.6	260.3
5	Alzheimer's disease (G30)	15,968	5.2	205.0
6	Diabetes mellitus (E10-E14)	9,409	3.1	120.8
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	6,734	2.2	86.5
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	6,503	2.1	83.5
9	Influenza and pneumonia (J09-J18)	6,437	2.1	82.6
10	Septicemia (A40-A41)	5,300	1.7	68.0
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	3,945	1.3	50.6
12	Parkinson's disease (G20-G21)	3,918	1.3	50.3
13	Pneumonitis due to solids and liquids (J69)	2,104	0.7	27.0
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2,077	0.7	26.7
15	Chronic liver disease and cirrhosis (K70,K73-K74)	1,919	0.6	24.6
...	All other causes (residual)	59,815	19.4	767.9

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2014.

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex:
United States, 2014

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
All races, female, 85 years and over				
...	All causes	517,441	100.0	12,765.7
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	147,696	28.5	3,643.8
2	Malignant neoplasms (C00-C97)	53,166	10.3	1,311.7
3	Alzheimer's disease (G30)	45,707	8.8	1,127.6
4	Cerebrovascular diseases (I60-I69)	39,746	7.7	980.6
5	Chronic lower respiratory diseases (J40-J47)	24,294	4.7	599.4
6	Influenza and pneumonia (J09-J18)	14,102	2.7	347.9
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	12,839	2.5	316.7
8	Diabetes mellitus (E10-E14)	9,806	1.9	241.9
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	9,563	1.8	235.9
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	9,326	1.8	230.1
11	Septicemia (A40-A41)	6,699	1.3	165.3
12	Parkinson's disease (G20-G21)	5,164	1.0	127.4
13	Pneumonitis due to solids and liquids (J69)	4,470	0.9	110.3
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2,959	0.6	73.0
15	Atherosclerosis (I70)	2,409	0.5	59.4
...	All other causes (residual)	129,495	25.0	3,194.8
White, both sexes, all ages				
...	All causes	2,237,880	100.0	892.9
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	524,695	23.4	209.4
2	Malignant neoplasms (C00-C97)	502,933	22.5	200.7
3	Chronic lower respiratory diseases (J40-J47)	134,541	6.0	53.7
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	117,151	5.2	46.7
5	Cerebrovascular diseases (I60-I69)	111,035	5.0	44.3
6	Alzheimer's disease (G30)	84,990	3.8	33.9
7	Diabetes mellitus (E10-E14)	59,741	2.7	23.8
8	Influenza and pneumonia (J09-J18)	47,293	2.1	18.9
9	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	38,723	1.7	15.5
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	37,976	1.7	15.2
11	Chronic liver disease and cirrhosis (K70,K73-K74)	33,508	1.5	13.4
12	Septicemia (A40-A41)	31,512	1.4	12.6
13	Parkinson's disease (G20-G21)	24,294	1.1	9.7
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	23,639	1.1	9.4
15	Pneumonitis due to solids and liquids (J69)	16,487	0.7	6.6
...	All other causes (residual)	449,362	20.1	179.3

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2014.

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex:
United States, 2014

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
White, both sexes, 1-4 years				
...	All causes	2,592	100.0	21.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	859	33.1	7.2
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	271	10.5	2.3
3	Malignant neoplasms (C00-C97)	250	9.6	2.1
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	200	7.7	1.7
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	84	3.2	0.7
6	Influenza and pneumonia (J09-J18)	72	2.8	0.6
7	Septicemia (A40-A41)	37	1.4	0.3
8	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	28	1.1	0.2
9	Chronic lower respiratory diseases (J40-J47)	25	1.0	0.2
10	Certain conditions originating in the perinatal period (P00-P96)	24	0.9	0.2
11	Cerebrovascular diseases (I60-I69)	20	0.8	0.2
12	Meningitis (G00,G03)	12	0.5	*
13	Complications of medical and surgical care (Y40-Y84,Y88)	10	0.4	*
14	Acute bronchitis and bronchiolitis (J20-J21)	9	0.3	*
15	Anemias (D50-D64)	8	0.3	*
...	All other causes (residual)	683	26.4	5.7
White, both sexes, 5-14 years				
...	All causes	3,753	100.0	12.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,067	28.4	3.4
2	Malignant neoplasms (C00-C97)	666	17.7	2.1
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	348	9.3	1.1
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	254	6.8	0.8
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	160	4.3	0.5
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	116	3.1	0.4
7	Influenza and pneumonia (J09-J18)	73	1.9	0.2
8	Cerebrovascular diseases (I60-I69)	69	1.8	0.2
9	Chronic lower respiratory diseases (J40-J47)	59	1.6	0.2
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	50	1.3	0.2
11	Septicemia (A40-A41)	41	1.1	0.1
12	Certain conditions originating in the perinatal period (P00-P96)	21	0.6	0.1
13	Diabetes mellitus (E10-E14)	14	0.4	*
14	Anemias (D50-D64)	13	0.3	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	8	0.2	*
...	All other causes (residual)	794	21.2	2.6

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2014.

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex:
United States, 2014

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
White, both sexes, 15-24 years				
...	All causes	20,760	100.0	62.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	9,734	46.9	29.4
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	4,186	20.2	12.6
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,371	6.6	4.1
4	Malignant neoplasms (C00-C97)	1,209	5.8	3.6
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	621	3.0	1.9
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	288	1.4	0.9
7	Influenza and pneumonia (J09-J18)	154	0.7	0.5
8	Cerebrovascular diseases (I60-I69)	120	0.6	0.4
9	Diabetes mellitus (E10-E14)	118	0.6	0.4
10	Pregnancy, childbirth and the puerperium (O00-O99)	99	0.5	0.3
11	Chronic lower respiratory diseases (J40-J47)	95	0.5	0.3
12	Septicemia (A40-A41)	89	0.4	0.3
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	54	0.3	0.2
14	Legal intervention (Y35,Y89.0)	47	0.2	0.1
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	41	0.2	0.1
...	All other causes (residual)	2,534	12.2	7.6
White, both sexes, 25-34 years				
...	All causes	35,408	100.0	106.9
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	14,649	41.4	44.2
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	5,638	15.9	17.0
3	Malignant neoplasms (C00-C97)	2,771	7.8	8.4
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,177	6.1	6.6
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,673	4.7	5.1
6	Chronic liver disease and cirrhosis (K70,K73-K74)	587	1.7	1.8
7	Diabetes mellitus (E10-E14)	435	1.2	1.3
8	Influenza and pneumonia (J09-J18)	424	1.2	1.3
9	Cerebrovascular diseases (I60-I69)	387	1.1	1.2
10	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	349	1.0	1.1
11	Septicemia (A40-A41)	258	0.7	0.8
12	Pregnancy, childbirth and the puerperium (O00-O99)	240	0.7	0.7
13	Chronic lower respiratory diseases (J40-J47)	230	0.6	0.7
14	Human immunodeficiency virus (HIV) disease (B20-B24)	194	0.5	0.6
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	140	0.4	0.4
...	All other causes (residual)	5,256	14.8	15.9

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2014.

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex:
United States, 2014

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
White, both sexes, 35-44 years				
...	All causes	53,674	100.0	172.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	13,573	25.3	43.5
2	Malignant neoplasms (C00-C97)	8,610	16.0	27.6
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	7,135	13.3	22.9
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	5,979	11.1	19.2
5	Chronic liver disease and cirrhosis (K70,K73-K74)	2,172	4.0	7.0
6	Diabetes mellitus (E10-E14)	1,355	2.5	4.3
7	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,274	2.4	4.1
8	Cerebrovascular diseases (I60-I69)	1,129	2.1	3.6
9	Influenza and pneumonia (J09-J18)	863	1.6	2.8
10	Septicemia (A40-A41)	605	1.1	1.9
11	Chronic lower respiratory diseases (J40-J47)	534	1.0	1.7
12	Human immunodeficiency virus (HIV) disease (B20-B24)	511	1.0	1.6
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	376	0.7	1.2
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	327	0.6	1.0
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	220	0.4	0.7
...	All other causes (residual)	9,011	16.8	28.9
White, both sexes, 45-54 years				
...	All causes	137,497	100.0	397.1
1	Malignant neoplasms (C00-C97)	35,341	25.7	102.1
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	25,922	18.9	74.9
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	17,431	12.7	50.3
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	8,123	5.9	23.5
5	Chronic liver disease and cirrhosis (K70,K73-K74)	7,533	5.5	21.8
6	Diabetes mellitus (E10-E14)	4,366	3.2	12.6
7	Chronic lower respiratory diseases (J40-J47)	3,675	2.7	10.6
8	Cerebrovascular diseases (I60-I69)	3,573	2.6	10.3
9	Influenza and pneumonia (J09-J18)	2,126	1.5	6.1
10	Septicemia (A40-A41)	1,843	1.3	5.3
11	Viral hepatitis (B15-B19)	1,512	1.1	4.4
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,285	0.9	3.7
13	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,129	0.8	3.3
14	Human immunodeficiency virus (HIV) disease (B20-B24)	1,090	0.8	3.1
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	886	0.6	2.6
...	All other causes (residual)	21,662	15.8	62.6

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2014.

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex:
United States, 2014

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
White, both sexes, 55-64 years				
...	All causes	277,580	100.0	844.2
1	Malignant neoplasms (C00-C97)	93,131	33.6	283.2
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	57,690	20.8	175.4
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	14,997	5.4	45.6
4	Chronic lower respiratory diseases (J40-J47)	14,400	5.2	43.8
5	Chronic liver disease and cirrhosis (K70,K73-K74)	11,071	4.0	33.7
6	Diabetes mellitus (E10-E14)	9,850	3.5	30.0
7	Cerebrovascular diseases (I60-I69)	8,176	2.9	24.9
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	7,109	2.6	21.6
9	Influenza and pneumonia (J09-J18)	4,253	1.5	12.9
10	Septicemia (A40-A41)	4,237	1.5	12.9
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3,304	1.2	10.0
12	Viral hepatitis (B15-B19)	3,122	1.1	9.5
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	2,221	0.8	6.8
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1,336	0.5	4.1
15	Aortic aneurysm and dissection (I71)	1,076	0.4	3.3
...	All other causes (residual)	41,607	15.0	126.5
White, both sexes, 65-74 years				
...	All causes	396,428	100.0	1,769.2
1	Malignant neoplasms (C00-C97)	135,521	34.2	604.8
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	83,591	21.1	373.0
3	Chronic lower respiratory diseases (J40-J47)	32,527	8.2	145.2
4	Cerebrovascular diseases (I60-I69)	15,304	3.9	68.3
5	Diabetes mellitus (E10-E14)	14,226	3.6	63.5
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	10,317	2.6	46.0
7	Chronic liver disease and cirrhosis (K70,K73-K74)	6,963	1.8	31.1
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	6,663	1.7	29.7
9	Influenza and pneumonia (J09-J18)	6,528	1.6	29.1
10	Septicemia (A40-A41)	6,514	1.6	29.1
11	Alzheimer's disease (G30)	4,598	1.2	20.5
12	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	3,905	1.0	17.4
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	3,258	0.8	14.5
14	Parkinson's disease (G20-G21)	3,133	0.8	14.0
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2,586	0.7	11.5
...	All other causes (residual)	60,794	15.3	271.3

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2014.

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex:
United States, 2014

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
White, both sexes, 75-84 years				
...	All causes	545,948	100.0	4,619.3
1	Malignant neoplasms (C00-C97)	135,028	24.7	1,142.5
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	126,744	23.2	1,072.4
3	Chronic lower respiratory diseases (J40-J47)	44,363	8.1	375.4
4	Cerebrovascular diseases (I60-I69)	30,811	5.6	260.7
5	Alzheimer's disease (G30)	22,814	4.2	193.0
6	Diabetes mellitus (E10-E14)	15,525	2.8	131.4
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	13,551	2.5	114.7
8	Influenza and pneumonia (J09-J18)	11,367	2.1	96.2
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	10,954	2.0	92.7
10	Parkinson's disease (G20-G21)	10,063	1.8	85.1
11	Septicemia (A40-A41)	8,355	1.5	70.7
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	5,525	1.0	46.7
13	Pneumonitis due to solids and liquids (J69)	4,376	0.8	37.0
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	4,334	0.8	36.7
15	Chronic liver disease and cirrhosis (K70,K73-K74)	3,809	0.7	32.2
...	All other causes (residual)	98,329	18.0	832.0
White, both sexes, 85 years and over				
...	All causes	749,222	100.0	13,742.6
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	220,405	29.4	4,042.8
2	Malignant neoplasms (C00-C97)	90,352	12.1	1,657.3
3	Alzheimer's disease (G30)	56,765	7.6	1,041.2
4	Cerebrovascular diseases (I60-I69)	51,379	6.9	942.4
5	Chronic lower respiratory diseases (J40-J47)	38,619	5.2	708.4
6	Influenza and pneumonia (J09-J18)	21,329	2.8	391.2
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	20,257	2.7	371.6
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	15,132	2.0	277.6
9	Diabetes mellitus (E10-E14)	13,846	1.8	254.0
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	11,455	1.5	210.1
11	Parkinson's disease (G20-G21)	10,509	1.4	192.8
12	Septicemia (A40-A41)	9,437	1.3	173.1
13	Pneumonitis due to solids and liquids (J69)	8,312	1.1	152.5
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	4,971	0.7	91.2
15	Atherosclerosis (I70)	3,317	0.4	60.8
...	All other causes (residual)	173,137	23.1	3,175.8

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2014.

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex:
United States, 2014

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
White, male, all ages				
...	All causes	1,128,993	100.0	909.4
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	277,921	24.6	223.9
2	Malignant neoplasms (C00-C97)	266,137	23.6	214.4
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	72,807	6.4	58.6
4	Chronic lower respiratory diseases (J40-J47)	62,989	5.6	50.7
5	Cerebrovascular diseases (I60-I69)	45,505	4.0	36.7
6	Diabetes mellitus (E10-E14)	32,920	2.9	26.5
7	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	30,015	2.7	24.2
8	Alzheimer's disease (G30)	25,937	2.3	20.9
9	Influenza and pneumonia (J09-J18)	22,643	2.0	18.2
10	Chronic liver disease and cirrhosis (K70,K73-K74)	21,781	1.9	17.5
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	19,587	1.7	15.8
12	Septicemia (A40-A41)	14,847	1.3	12.0
13	Parkinson's disease (G20-G21)	14,594	1.3	11.8
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	9,998	0.9	8.1
15	Pneumonitis due to solids and liquids (J69)	9,119	0.8	7.3
...	All other causes (residual)	202,193	17.9	162.9
White, male, 1-4 years				
...	All causes	1,452	100.0	23.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	521	35.9	8.6
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	146	10.1	2.4
3	Malignant neoplasms (C00-C97)	142	9.8	2.3
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	105	7.2	1.7
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	42	2.9	0.7
6	Influenza and pneumonia (J09-J18)	37	2.5	0.6
7	Septicemia (A40-A41)	15	1.0	*
8	Chronic lower respiratory diseases (J40-J47)	14	1.0	*
8	Certain conditions originating in the perinatal period (P00-P96)	14	1.0	*
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	12	0.8	*
10	Cerebrovascular diseases (I60-I69)	12	0.8	*
12	Anemias (D50-D64)	4	0.3	*
12	Meningitis (G00,G03)	4	0.3	*
12	Complications of medical and surgical care (Y40-Y84,Y88)	4	0.3	*
15	Diabetes mellitus (E10-E14)	3	0.2	*
15	Nutritional deficiencies (E40-E64)	3	0.2	*
15	Acute bronchitis and bronchiolitis (J20-J21)	3	0.2	*
...	All other causes (residual)	371	25.6	6.1

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2014.

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex:
United States, 2014

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
White, male, 5-14 years				
...	All causes	2,231	100.0	14.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	695	31.2	4.4
2	Malignant neoplasms (C00-C97)	387	17.3	2.4
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	225	10.1	1.4
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	133	6.0	0.8
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	88	3.9	0.6
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	59	2.6	0.4
7	Cerebrovascular diseases (I60-I69)	36	1.6	0.2
7	Influenza and pneumonia (J09-J18)	36	1.6	0.2
7	Chronic lower respiratory diseases (J40-J47)	36	1.6	0.2
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	26	1.2	0.2
11	Septicemia (A40-A41)	20	0.9	0.1
12	Diabetes mellitus (E10-E14)	10	0.4	*
13	Certain conditions originating in the perinatal period (P00-P96)	9	0.4	*
14	Anemias (D50-D64)	8	0.4	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	7	0.3	*
...	All other causes (residual)	456	20.4	2.9
White, male, 15-24 years				
...	All causes	15,041	100.0	88.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	7,246	48.2	42.5
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	3,382	22.5	19.9
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,102	7.3	6.5
4	Malignant neoplasms (C00-C97)	734	4.9	4.3
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	408	2.7	2.4
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	162	1.1	1.0
7	Influenza and pneumonia (J09-J18)	84	0.6	0.5
8	Diabetes mellitus (E10-E14)	79	0.5	0.5
9	Cerebrovascular diseases (I60-I69)	68	0.5	0.4
10	Chronic lower respiratory diseases (J40-J47)	61	0.4	0.4
11	Septicemia (A40-A41)	51	0.3	0.3
12	Legal intervention (Y35,Y89.0)	43	0.3	0.3
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	25	0.2	0.1
14	Human immunodeficiency virus (HIV) disease (B20-B24)	24	0.2	0.1
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	21	0.1	0.1
...	All other causes (residual)	1,551	10.3	9.1

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2014.

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex:
United States, 2014

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
White, male, 25-34 years				
...	All causes	24,572	100.0	145.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	10,819	44.0	64.0
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	4,478	18.2	26.5
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,518	6.2	9.0
4	Malignant neoplasms (C00-C97)	1,425	5.8	8.4
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,267	5.2	7.5
6	Chronic liver disease and cirrhosis (K70,K73-K74)	365	1.5	2.2
7	Diabetes mellitus (E10-E14)	277	1.1	1.6
8	Influenza and pneumonia (J09-J18)	228	0.9	1.3
9	Cerebrovascular diseases (I60-I69)	218	0.9	1.3
10	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	211	0.9	1.2
11	Human immunodeficiency virus (HIV) disease (B20-B24)	148	0.6	0.9
12	Chronic lower respiratory diseases (J40-J47)	136	0.6	0.8
13	Septicemia (A40-A41)	132	0.5	0.8
14	Legal intervention (Y35,Y89.0)	106	0.4	0.6
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	80	0.3	0.5
...	All other causes (residual)	3,164	12.9	18.7
White, male, 35-44 years				
...	All causes	33,482	100.0	212.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	9,345	27.9	59.3
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	4,956	14.8	31.5
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	4,494	13.4	28.5
4	Malignant neoplasms (C00-C97)	3,787	11.3	24.0
5	Chronic liver disease and cirrhosis (K70,K73-K74)	1,402	4.2	8.9
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	939	2.8	6.0
7	Diabetes mellitus (E10-E14)	852	2.5	5.4
8	Cerebrovascular diseases (I60-I69)	633	1.9	4.0
9	Influenza and pneumonia (J09-J18)	479	1.4	3.0
10	Human immunodeficiency virus (HIV) disease (B20-B24)	390	1.2	2.5
11	Septicemia (A40-A41)	307	0.9	1.9
12	Chronic lower respiratory diseases (J40-J47)	240	0.7	1.5
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	232	0.7	1.5
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	168	0.5	1.1
15	Aortic aneurysm and dissection (I71)	159	0.5	1.0
...	All other causes (residual)	5,099	15.2	32.4

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2014.

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex:
United States, 2014

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
White, male, 45-54 years				
...	All causes	84,414	100.0	488.9
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	18,597	22.0	107.7
2	Malignant neoplasms (C00-C97)	17,662	20.9	102.3
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	11,659	13.8	67.5
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	5,947	7.0	34.4
5	Chronic liver disease and cirrhosis (K70,K73-K74)	5,089	6.0	29.5
6	Diabetes mellitus (E10-E14)	2,811	3.3	16.3
7	Cerebrovascular diseases (I60-I69)	2,046	2.4	11.9
8	Chronic lower respiratory diseases (J40-J47)	1,680	2.0	9.7
9	Influenza and pneumonia (J09-J18)	1,167	1.4	6.8
10	Viral hepatitis (B15-B19)	1,037	1.2	6.0
11	Septicemia (A40-A41)	999	1.2	5.8
12	Human immunodeficiency virus (HIV) disease (B20-B24)	906	1.1	5.2
13	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	817	1.0	4.7
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	808	1.0	4.7
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	586	0.7	3.4
...	All other causes (residual)	12,603	14.9	73.0
White, male, 55-64 years				
...	All causes	170,566	100.0	1,063.8
1	Malignant neoplasms (C00-C97)	51,875	30.4	323.5
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	41,036	24.1	255.9
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	10,062	5.9	62.8
4	Chronic liver disease and cirrhosis (K70,K73-K74)	7,754	4.5	48.4
5	Chronic lower respiratory diseases (J40-J47)	7,464	4.4	46.6
6	Diabetes mellitus (E10-E14)	6,229	3.7	38.8
7	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	5,361	3.1	33.4
8	Cerebrovascular diseases (I60-I69)	4,743	2.8	29.6
9	Influenza and pneumonia (J09-J18)	2,427	1.4	15.1
10	Septicemia (A40-A41)	2,247	1.3	14.0
11	Viral hepatitis (B15-B19)	2,215	1.3	13.8
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,930	1.1	12.0
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,418	0.8	8.8
14	Aortic aneurysm and dissection (I71)	818	0.5	5.1
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	799	0.5	5.0
...	All other causes (residual)	24,188	14.2	150.9

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2014.

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex:
United States, 2014

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
White, male, 65-74 years				
...	All causes	227,249	100.0	2,143.3
1	Malignant neoplasms (C00-C97)	75,028	33.0	707.6
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	54,364	23.9	512.7
3	Chronic lower respiratory diseases (J40-J47)	16,351	7.2	154.2
4	Diabetes mellitus (E10-E14)	8,518	3.7	80.3
5	Cerebrovascular diseases (I60-I69)	8,199	3.6	77.3
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	6,496	2.9	61.3
7	Chronic liver disease and cirrhosis (K70,K73-K74)	4,447	2.0	41.9
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3,656	1.6	34.5
9	Influenza and pneumonia (J09-J18)	3,588	1.6	33.8
10	Septicemia (A40-A41)	3,382	1.5	31.9
11	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	3,130	1.4	29.5
12	Parkinson's disease (G20-G21)	2,082	0.9	19.6
13	Alzheimer's disease (G30)	1,980	0.9	18.7
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,835	0.8	17.3
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1,570	0.7	14.8
...	All other causes (residual)	32,623	14.4	307.7
White, male, 75-84 years				
...	All causes	279,177	100.0	5,419.1
1	Malignant neoplasms (C00-C97)	72,145	25.8	1,400.4
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	70,142	25.1	1,361.5
3	Chronic lower respiratory diseases (J40-J47)	21,222	7.6	411.9
4	Cerebrovascular diseases (I60-I69)	13,670	4.9	265.3
5	Alzheimer's disease (G30)	8,530	3.1	165.6
6	Diabetes mellitus (E10-E14)	8,288	3.0	160.9
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	7,404	2.7	143.7
8	Parkinson's disease (G20-G21)	6,456	2.3	125.3
9	Influenza and pneumonia (J09-J18)	5,847	2.1	113.5
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	5,841	2.1	113.4
11	Septicemia (A40-A41)	3,936	1.4	76.4
12	Pneumonitis due to solids and liquids (J69)	2,568	0.9	49.8
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2,500	0.9	48.5
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	2,443	0.9	47.4
15	Chronic liver disease and cirrhosis (K70,K73-K74)	2,073	0.7	40.2
...	All other causes (residual)	46,112	16.5	895.1

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2014.

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex:
United States, 2014

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
White, male, 85 years and over				
...	All causes	282,424	100.0	15,000.3
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	86,681	30.7	4,603.9
2	Malignant neoplasms (C00-C97)	42,921	15.2	2,279.7
3	Cerebrovascular diseases (I60-I69)	15,841	5.6	841.4
4	Chronic lower respiratory diseases (J40-J47)	15,772	5.6	837.7
5	Alzheimer's disease (G30)	15,079	5.3	800.9
6	Influenza and pneumonia (J09-J18)	8,693	3.1	461.7
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	8,139	2.9	432.3
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	6,977	2.5	370.6
9	Diabetes mellitus (E10-E14)	5,852	2.1	310.8
10	Parkinson's disease (G20-G21)	5,671	2.0	301.2
11	Pneumonitis due to solids and liquids (J69)	4,283	1.5	227.5
12	Septicemia (A40-A41)	3,701	1.3	196.6
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	3,509	1.2	186.4
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2,276	0.8	120.9
15	Atherosclerosis (I70)	1,088	0.4	57.8
...	All other causes (residual)	55,941	19.8	2,971.2
White, female, all ages				
...	All causes	1,108,887	100.0	876.7
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	246,774	22.3	195.1
2	Malignant neoplasms (C00-C97)	236,796	21.4	187.2
3	Chronic lower respiratory diseases (J40-J47)	71,552	6.5	56.6
4	Cerebrovascular diseases (I60-I69)	65,530	5.9	51.8
5	Alzheimer's disease (G30)	59,053	5.3	46.7
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	44,344	4.0	35.1
7	Diabetes mellitus (E10-E14)	26,821	2.4	21.2
8	Influenza and pneumonia (J09-J18)	24,650	2.2	19.5
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	18,389	1.7	14.5
10	Septicemia (A40-A41)	16,665	1.5	13.2
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	13,641	1.2	10.8
12	Chronic liver disease and cirrhosis (K70,K73-K74)	11,727	1.1	9.3
13	Parkinson's disease (G20-G21)	9,700	0.9	7.7
14	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	8,708	0.8	6.9
15	Pneumonitis due to solids and liquids (J69)	7,368	0.7	5.8
...	All other causes (residual)	247,169	22.3	195.4

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2014.

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex:
United States, 2014

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
White, female, 1-4 years				
...	All causes	1,140	100.0	19.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	338	29.6	5.8
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	125	11.0	2.1
3	Malignant neoplasms (C00-C97)	108	9.5	1.9
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	95	8.3	1.6
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	42	3.7	0.7
6	Influenza and pneumonia (J09-J18)	35	3.1	0.6
7	Septicemia (A40-A41)	22	1.9	0.4
8	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	16	1.4	*
9	Chronic lower respiratory diseases (J40-J47)	11	1.0	*
10	Certain conditions originating in the perinatal period (P00-P96)	10	0.9	*
11	Meningitis (G00,G03)	8	0.7	*
11	Cerebrovascular diseases (I60-I69)	8	0.7	*
13	Acute bronchitis and bronchiolitis (J20-J21)	6	0.5	*
13	Complications of medical and surgical care (Y40-Y84,Y88)	6	0.5	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	5	0.4	*
...	All other causes (residual)	305	26.8	5.2
White, female, 5-14 years				
...	All causes	1,522	100.0	10.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	372	24.4	2.4
2	Malignant neoplasms (C00-C97)	279	18.3	1.8
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	123	8.1	0.8
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	121	8.0	0.8
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	72	4.7	0.5
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	57	3.7	0.4
7	Influenza and pneumonia (J09-J18)	37	2.4	0.2
8	Cerebrovascular diseases (I60-I69)	33	2.2	0.2
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	24	1.6	0.2
10	Chronic lower respiratory diseases (J40-J47)	23	1.5	0.2
11	Septicemia (A40-A41)	21	1.4	0.1
12	Certain conditions originating in the perinatal period (P00-P96)	12	0.8	*
13	Anemias (D50-D64)	5	0.3	*
14	Diabetes mellitus (E10-E14)	4	0.3	*
15	Pneumonitis due to solids and liquids (J69)	3	0.2	*
15	Complications of medical and surgical care (Y40-Y84,Y88)	3	0.2	*
...	All other causes (residual)	333	21.9	2.2

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2014.

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex:
United States, 2014

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
White, female, 15-24 years				
...	All causes	5,719	100.0	35.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,488	43.5	15.4
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	804	14.1	5.0
3	Malignant neoplasms (C00-C97)	475	8.3	2.9
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	269	4.7	1.7
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	213	3.7	1.3
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	126	2.2	0.8
7	Pregnancy, childbirth and the puerperium (O00-O99)	99	1.7	0.6
8	Influenza and pneumonia (J09-J18)	70	1.2	0.4
9	Cerebrovascular diseases (I60-I69)	52	0.9	0.3
10	Diabetes mellitus (E10-E14)	39	0.7	0.2
11	Septicemia (A40-A41)	38	0.7	0.2
12	Chronic lower respiratory diseases (J40-J47)	34	0.6	0.2
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	29	0.5	0.2
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	20	0.3	0.1
15	Pneumonitis due to solids and liquids (J69)	15	0.3	*
...	All other causes (residual)	948	16.6	5.9
White, female, 25-34 years				
...	All causes	10,836	100.0	66.9
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,830	35.3	23.6
2	Malignant neoplasms (C00-C97)	1,346	12.4	8.3
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,160	10.7	7.2
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	659	6.1	4.1
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	406	3.7	2.5
6	Pregnancy, childbirth and the puerperium (O00-O99)	240	2.2	1.5
7	Chronic liver disease and cirrhosis (K70,K73-K74)	222	2.0	1.4
8	Influenza and pneumonia (J09-J18)	196	1.8	1.2
9	Cerebrovascular diseases (I60-I69)	169	1.6	1.0
10	Diabetes mellitus (E10-E14)	158	1.5	1.0
11	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	138	1.3	0.9
12	Septicemia (A40-A41)	126	1.2	0.8
13	Chronic lower respiratory diseases (J40-J47)	94	0.9	0.6
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	60	0.6	0.4
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	49	0.5	0.3
...	All other causes (residual)	1,983	18.3	12.2

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2014.

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex:
United States, 2014

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
White, female, 35-44 years				
...	All causes	20,192	100.0	130.7
1	Malignant neoplasms (C00-C97)	4,823	23.9	31.2
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	4,228	20.9	27.4
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,179	10.8	14.1
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,485	7.4	9.6
5	Chronic liver disease and cirrhosis (K70,K73-K74)	770	3.8	5.0
6	Diabetes mellitus (E10-E14)	503	2.5	3.3
7	Cerebrovascular diseases (I60-I69)	496	2.5	3.2
8	Influenza and pneumonia (J09-J18)	384	1.9	2.5
9	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	335	1.7	2.2
10	Septicemia (A40-A41)	298	1.5	1.9
11	Chronic lower respiratory diseases (J40-J47)	294	1.5	1.9
12	Pregnancy, childbirth and the puerperium (O00-O99)	219	1.1	1.4
13	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	159	0.8	1.0
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	144	0.7	0.9
15	Human immunodeficiency virus (HIV) disease (B20-B24)	121	0.6	0.8
...	All other causes (residual)	3,754	18.6	24.3
White, female, 45-54 years				
...	All causes	53,083	100.0	305.8
1	Malignant neoplasms (C00-C97)	17,679	33.3	101.8
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	7,325	13.8	42.2
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	5,772	10.9	33.2
4	Chronic liver disease and cirrhosis (K70,K73-K74)	2,444	4.6	14.1
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,176	4.1	12.5
6	Chronic lower respiratory diseases (J40-J47)	1,995	3.8	11.5
7	Diabetes mellitus (E10-E14)	1,555	2.9	9.0
8	Cerebrovascular diseases (I60-I69)	1,527	2.9	8.8
9	Influenza and pneumonia (J09-J18)	959	1.8	5.5
10	Septicemia (A40-A41)	844	1.6	4.9
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	477	0.9	2.7
12	Viral hepatitis (B15-B19)	475	0.9	2.7
13	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	314	0.6	1.8
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	312	0.6	1.8
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	300	0.6	1.7
...	All other causes (residual)	8,929	16.8	51.4

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2014.

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex:
United States, 2014

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
White, female, 55-64 years				
...	All causes	107,014	100.0	635.1
1	Malignant neoplasms (C00-C97)	41,256	38.6	244.9
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	16,654	15.6	98.8
3	Chronic lower respiratory diseases (J40-J47)	6,936	6.5	41.2
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	4,935	4.6	29.3
5	Diabetes mellitus (E10-E14)	3,621	3.4	21.5
6	Cerebrovascular diseases (I60-I69)	3,433	3.2	20.4
7	Chronic liver disease and cirrhosis (K70,K73-K74)	3,317	3.1	19.7
8	Septicemia (A40-A41)	1,990	1.9	11.8
9	Influenza and pneumonia (J09-J18)	1,826	1.7	10.8
10	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,748	1.6	10.4
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,374	1.3	8.2
12	Viral hepatitis (B15-B19)	907	0.8	5.4
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	803	0.8	4.8
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	537	0.5	3.2
15	Alzheimer's disease (G30)	425	0.4	2.5
...	All other causes (residual)	17,252	16.1	102.4
White, female, 65-74 years				
...	All causes	169,179	100.0	1,433.1
1	Malignant neoplasms (C00-C97)	60,493	35.8	512.4
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	29,227	17.3	247.6
3	Chronic lower respiratory diseases (J40-J47)	16,176	9.6	137.0
4	Cerebrovascular diseases (I60-I69)	7,105	4.2	60.2
5	Diabetes mellitus (E10-E14)	5,708	3.4	48.4
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,821	2.3	32.4
7	Septicemia (A40-A41)	3,132	1.9	26.5
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3,007	1.8	25.5
9	Influenza and pneumonia (J09-J18)	2,940	1.7	24.9
10	Alzheimer's disease (G30)	2,618	1.5	22.2
11	Chronic liver disease and cirrhosis (K70,K73-K74)	2,516	1.5	21.3
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,423	0.8	12.1
13	Parkinson's disease (G20-G21)	1,051	0.6	8.9
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1,016	0.6	8.6
15	Pneumonitis due to solids and liquids (J69)	837	0.5	7.1
...	All other causes (residual)	28,109	16.6	238.1

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2014.

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex:
United States, 2014

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
White, female, 75-84 years				
...	All causes	266,771	100.0	4,001.3
1	Malignant neoplasms (C00-C97)	62,883	23.6	943.2
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	56,602	21.2	849.0
3	Chronic lower respiratory diseases (J40-J47)	23,141	8.7	347.1
4	Cerebrovascular diseases (I60-I69)	17,141	6.4	257.1
5	Alzheimer's disease (G30)	14,284	5.4	214.2
6	Diabetes mellitus (E10-E14)	7,237	2.7	108.5
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	6,147	2.3	92.2
8	Influenza and pneumonia (J09-J18)	5,520	2.1	82.8
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	5,113	1.9	76.7
10	Septicemia (A40-A41)	4,419	1.7	66.3
11	Parkinson's disease (G20-G21)	3,607	1.4	54.1
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	3,082	1.2	46.2
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1,834	0.7	27.5
14	Pneumonitis due to solids and liquids (J69)	1,808	0.7	27.1
15	Chronic liver disease and cirrhosis (K70,K73-K74)	1,736	0.7	26.0
...	All other causes (residual)	52,217	19.6	783.2
White, female, 85 years and over				
...	All causes	466,798	100.0	13,079.1
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	133,724	28.6	3,746.8
2	Malignant neoplasms (C00-C97)	47,431	10.2	1,329.0
3	Alzheimer's disease (G30)	41,686	8.9	1,168.0
4	Cerebrovascular diseases (I60-I69)	35,538	7.6	995.7
5	Chronic lower respiratory diseases (J40-J47)	22,847	4.9	640.1
6	Influenza and pneumonia (J09-J18)	12,636	2.7	354.0
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	12,118	2.6	339.5
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	8,155	1.7	228.5
9	Diabetes mellitus (E10-E14)	7,994	1.7	224.0
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	7,946	1.7	222.6
11	Septicemia (A40-A41)	5,736	1.2	160.7
12	Parkinson's disease (G20-G21)	4,838	1.0	135.6
13	Pneumonitis due to solids and liquids (J69)	4,029	0.9	112.9
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2,695	0.6	75.5
15	Atherosclerosis (I70)	2,229	0.5	62.5
...	All other causes (residual)	117,196	25.1	3,283.7

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2014.

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex:
United States, 2014

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
Black, both sexes, all ages				
...	All causes	308,960	100.0	697.3
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	73,095	23.7	165.0
2	Malignant neoplasms (C00-C97)	69,090	22.4	155.9
3	Cerebrovascular diseases (I60-I69)	17,088	5.5	38.6
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	14,135	4.6	31.9
5	Diabetes mellitus (E10-E14)	13,435	4.3	30.3
6	Chronic lower respiratory diseases (J40-J47)	9,934	3.2	22.4
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	8,586	2.8	19.4
8	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	7,903	2.6	17.8
9	Alzheimer's disease (G30)	6,567	2.1	14.8
10	Septicemia (A40-A41)	6,386	2.1	14.4
11	Influenza and pneumonia (J09-J18)	5,611	1.8	12.7
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	5,399	1.7	12.2
13	Certain conditions originating in the perinatal period (P00-P96)	4,032	1.3	9.1
14	Human immunodeficiency virus (HIV) disease (B20-B24)	3,591	1.2	8.1
15	Chronic liver disease and cirrhosis (K70,K73-K74)	3,103	1.0	7.0
...	All other causes (residual)	61,005	19.7	137.7
Black, both sexes, 1-4 years				
...	All causes	1,009	100.0	37.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	291	28.8	10.7
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	151	15.0	5.6
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	104	10.3	3.8
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	55	5.5	2.0
5	Malignant neoplasms (C00-C97)	54	5.4	2.0
6	Influenza and pneumonia (J09-J18)	29	2.9	1.1
7	Chronic lower respiratory diseases (J40-J47)	23	2.3	0.8
8	Septicemia (A40-A41)	12	1.2	*
8	Certain conditions originating in the perinatal period (P00-P96)	12	1.2	*
10	Cerebrovascular diseases (I60-I69)	9	0.9	*
11	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	7	0.7	*
11	Acute bronchitis and bronchiolitis (J20-J21)	7	0.7	*
11	Complications of medical and surgical care (Y40-Y84,Y88)	7	0.7	*
14	Anemias (D50-D64)	6	0.6	*
15	Meningitis (G00,G03)	4	0.4	*
...	All other causes (residual)	238	23.6	8.8

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2014.

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex:
United States, 2014

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
Black, both sexes, 5-14 years				
...	All causes	1,204	100.0	17.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	336	27.9	4.9
2	Malignant neoplasms (C00-C97)	146	12.1	2.1
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	105	8.7	1.5
4	Chronic lower respiratory diseases (J40-J47)	75	6.2	1.1
5	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	72	6.0	1.1
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	61	5.1	0.9
7	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	58	4.8	0.9
8	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	21	1.7	0.3
9	Anemias (D50-D64)	18	1.5	*
10	Influenza and pneumonia (J09-J18)	17	1.4	*
11	Septicemia (A40-A41)	16	1.3	*
12	Cerebrovascular diseases (I60-I69)	15	1.2	*
13	Diabetes mellitus (E10-E14)	11	0.9	*
14	Certain conditions originating in the perinatal period (P00-P96)	10	0.8	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	5	0.4	*
15	Complications of medical and surgical care (Y40-Y84,Y88)	5	0.4	*
...	All other causes (residual)	233	19.4	3.4
Black, both sexes, 15-24 years				
...	All causes	6,602	100.0	89.6
1	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	2,683	40.6	36.4
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,509	22.9	20.5
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	546	8.3	7.4
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	284	4.3	3.9
5	Malignant neoplasms (C00-C97)	272	4.1	3.7
6	Human immunodeficiency virus (HIV) disease (B20-B24)	86	1.3	1.2
7	Chronic lower respiratory diseases (J40-J47)	72	1.1	1.0
8	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	67	1.0	0.9
9	Anemias (D50-D64)	65	1.0	0.9
10	Diabetes mellitus (E10-E14)	60	0.9	0.8
11	Pregnancy, childbirth and the puerperium (O00-O99)	57	0.9	0.8
12	Cerebrovascular diseases (I60-I69)	49	0.7	0.7
13	Influenza and pneumonia (J09-J18)	38	0.6	0.5
14	Septicemia (A40-A41)	32	0.5	0.4
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	29	0.4	0.4
15	Legal intervention (Y35,Y89.0)	29	0.4	0.4
...	All other causes (residual)	724	11.0	9.8

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2014.

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex:
United States, 2014

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
Black, both sexes, 25-34 years				
...	All causes	9,549	100.0	148.6
1	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	2,368	24.8	36.8
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,956	20.5	30.4
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,010	10.6	15.7
4	Malignant neoplasms (C00-C97)	629	6.6	9.8
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	568	5.9	8.8
6	Human immunodeficiency virus (HIV) disease (B20-B24)	376	3.9	5.9
7	Diabetes mellitus (E10-E14)	239	2.5	3.7
8	Cerebrovascular diseases (I60-I69)	155	1.6	2.4
9	Pregnancy, childbirth and the puerperium (O00-O99)	143	1.5	2.2
10	Anemias (D50-D64)	127	1.3	2.0
11	Chronic lower respiratory diseases (J40-J47)	126	1.3	2.0
12	Influenza and pneumonia (J09-J18)	99	1.0	1.5
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	84	0.9	1.3
14	Septicemia (A40-A41)	81	0.8	1.3
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	63	0.7	1.0
...	All other causes (residual)	1,525	16.0	23.7
Black, both sexes, 35-44 years				
...	All causes	13,997	100.0	247.9
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,775	19.8	49.2
2	Malignant neoplasms (C00-C97)	1,976	14.1	35.0
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,853	13.2	32.8
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,222	8.7	21.6
5	Human immunodeficiency virus (HIV) disease (B20-B24)	640	4.6	11.3
6	Diabetes mellitus (E10-E14)	547	3.9	9.7
7	Cerebrovascular diseases (I60-I69)	516	3.7	9.1
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	442	3.2	7.8
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	265	1.9	4.7
10	Influenza and pneumonia (J09-J18)	201	1.4	3.6
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	197	1.4	3.5
12	Septicemia (A40-A41)	189	1.4	3.3
13	Chronic lower respiratory diseases (J40-J47)	188	1.3	3.3
14	Chronic liver disease and cirrhosis (K70,K73-K74)	184	1.3	3.3
15	Anemias (D50-D64)	143	1.0	2.5
...	All other causes (residual)	2,659	19.0	47.1

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2014.

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex:
United States, 2014

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
Black, both sexes, 45-54 years				
...	All causes	31,815	100.0	557.2
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	7,687	24.2	134.6
2	Malignant neoplasms (C00-C97)	7,492	23.5	131.2
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,528	7.9	44.3
4	Cerebrovascular diseases (I60-I69)	1,473	4.6	25.8
5	Diabetes mellitus (E10-E14)	1,439	4.5	25.2
6	Human immunodeficiency virus (HIV) disease (B20-B24)	1,100	3.5	19.3
7	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	728	2.3	12.8
8	Chronic liver disease and cirrhosis (K70,K73-K74)	724	2.3	12.7
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	660	2.1	11.6
10	Chronic lower respiratory diseases (J40-J47)	651	2.0	11.4
11	Septicemia (A40-A41)	593	1.9	10.4
12	Influenza and pneumonia (J09-J18)	500	1.6	8.8
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	485	1.5	8.5
14	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	376	1.2	6.6
15	Viral hepatitis (B15-B19)	188	0.6	3.3
...	All other causes (residual)	5,191	16.3	90.9
Black, both sexes, 55-64 years				
...	All causes	59,972	100.0	1,265.2
1	Malignant neoplasms (C00-C97)	18,040	30.1	380.6
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	14,638	24.4	308.8
3	Cerebrovascular diseases (I60-I69)	2,963	4.9	62.5
4	Diabetes mellitus (E10-E14)	2,900	4.8	61.2
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,462	4.1	51.9
6	Chronic lower respiratory diseases (J40-J47)	1,832	3.1	38.6
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,583	2.6	33.4
8	Septicemia (A40-A41)	1,323	2.2	27.9
9	Chronic liver disease and cirrhosis (K70,K73-K74)	1,312	2.2	27.7
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,018	1.7	21.5
11	Human immunodeficiency virus (HIV) disease (B20-B24)	996	1.7	21.0
12	Influenza and pneumonia (J09-J18)	936	1.6	19.7
13	Viral hepatitis (B15-B19)	651	1.1	13.7
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	359	0.6	7.6
15	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	252	0.4	5.3
...	All other causes (residual)	8,707	14.5	183.7

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2014.

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex:
United States, 2014

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
Black, both sexes, 65-74 years				
...	All causes	60,955	100.0	2,381.5
1	Malignant neoplasms (C00-C97)	18,700	30.7	730.6
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	15,130	24.8	591.1
3	Cerebrovascular diseases (I60-I69)	3,511	5.8	137.2
4	Diabetes mellitus (E10-E14)	3,213	5.3	125.5
5	Chronic lower respiratory diseases (J40-J47)	2,589	4.2	101.2
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2,082	3.4	81.3
7	Septicemia (A40-A41)	1,481	2.4	57.9
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,182	1.9	46.2
9	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,024	1.7	40.0
10	Influenza and pneumonia (J09-J18)	1,012	1.7	39.5
11	Chronic liver disease and cirrhosis (K70,K73-K74)	597	1.0	23.3
12	Alzheimer's disease (G30)	491	0.8	19.2
13	Pneumonitis due to solids and liquids (J69)	360	0.6	14.1
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	311	0.5	12.2
15	Human immunodeficiency virus (HIV) disease (B20-B24)	303	0.5	11.8
...	All other causes (residual)	8,969	14.7	350.4
Black, both sexes, 75-84 years				
...	All causes	60,428	100.0	5,049.0
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	15,575	25.8	1,301.3
2	Malignant neoplasms (C00-C97)	14,276	23.6	1,192.8
3	Cerebrovascular diseases (I60-I69)	4,183	6.9	349.5
4	Diabetes mellitus (E10-E14)	3,012	5.0	251.7
5	Chronic lower respiratory diseases (J40-J47)	2,558	4.2	213.7
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2,083	3.4	174.0
7	Alzheimer's disease (G30)	2,073	3.4	173.2
8	Septicemia (A40-A41)	1,387	2.3	115.9
9	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,218	2.0	101.8
10	Influenza and pneumonia (J09-J18)	1,216	2.0	101.6
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	861	1.4	71.9
12	Parkinson's disease (G20-G21)	489	0.8	40.9
13	Pneumonitis due to solids and liquids (J69)	486	0.8	40.6
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	314	0.5	26.2
15	Chronic liver disease and cirrhosis (K70,K73-K74)	189	0.3	15.8
...	All other causes (residual)	10,508	17.4	878.0

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2014.

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex:
United States, 2014

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
Black, both sexes, 85 years and over				
...	All causes	56,327	100.0	12,136.5
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	15,752	28.0	3,394.0
2	Malignant neoplasms (C00-C97)	7,494	13.3	1,614.7
3	Cerebrovascular diseases (I60-I69)	4,190	7.4	902.8
4	Alzheimer's disease (G30)	3,896	6.9	839.5
5	Diabetes mellitus (E10-E14)	2,012	3.6	433.5
6	Chronic lower respiratory diseases (J40-J47)	1,812	3.2	390.4
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,771	3.1	381.6
8	Influenza and pneumonia (J09-J18)	1,489	2.6	320.8
9	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,402	2.5	302.1
10	Septicemia (A40-A41)	1,219	2.2	262.7
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	734	1.3	158.2
12	Pneumonitis due to solids and liquids (J69)	521	0.9	112.3
13	Parkinson's disease (G20-G21)	420	0.7	90.5
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	281	0.5	60.5
15	Nutritional deficiencies (E40-E64)	194	0.3	41.8
...	All other causes (residual)	13,140	23.3	2,831.2
Black, male, all ages				
...	All causes	157,733	100.0	742.6
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	37,962	24.1	178.7
2	Malignant neoplasms (C00-C97)	35,061	22.2	165.1
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	9,537	6.0	44.9
4	Cerebrovascular diseases (I60-I69)	7,747	4.9	36.5
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	6,823	4.3	32.1
6	Diabetes mellitus (E10-E14)	6,452	4.1	30.4
7	Chronic lower respiratory diseases (J40-J47)	5,035	3.2	23.7
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	4,034	2.6	19.0
9	Septicemia (A40-A41)	2,969	1.9	14.0
10	Influenza and pneumonia (J09-J18)	2,736	1.7	12.9
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	2,481	1.6	11.7
12	Human immunodeficiency virus (HIV) disease (B20-B24)	2,373	1.5	11.2
13	Certain conditions originating in the perinatal period (P00-P96)	2,266	1.4	10.7
14	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,951	1.2	9.2
15	Chronic liver disease and cirrhosis (K70,K73-K74)	1,940	1.2	9.1
...	All other causes (residual)	28,366	18.0	133.5

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2014.

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex:
United States, 2014

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
Black, male, 1-4 years				
...	All causes	583	100.0	42.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	179	30.7	12.9
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	81	13.9	5.9
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	59	10.1	4.3
4	Malignant neoplasms (C00-C97)	28	4.8	2.0
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	26	4.5	1.9
6	Influenza and pneumonia (J09-J18)	19	3.3	*
7	Chronic lower respiratory diseases (J40-J47)	11	1.9	*
8	Certain conditions originating in the perinatal period (P00-P96)	9	1.5	*
9	Septicemia (A40-A41)	7	1.2	*
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	6	1.0	*
11	Anemias (D50-D64)	5	0.9	*
11	Cerebrovascular diseases (I60-I69)	5	0.9	*
13	Acute bronchitis and bronchiolitis (J20-J21)	4	0.7	*
14	Meningitis (G00,G03)	3	0.5	*
14	Complications of medical and surgical care (Y40-Y84,Y88)	3	0.5	*
...	All other causes (residual)	138	23.7	10.0
Black, male, 5-14 years				
...	All causes	725	100.0	21.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	225	31.0	6.5
2	Malignant neoplasms (C00-C97)	77	10.6	2.2
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	70	9.7	2.0
4	Chronic lower respiratory diseases (J40-J47)	51	7.0	1.5
5	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	42	5.8	1.2
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	42	5.8	1.2
7	Diseases of heart (I00-I09,I11,I13,I20-I51)	40	5.5	1.2
8	Influenza and pneumonia (J09-J18)	10	1.4	*
9	Septicemia (A40-A41)	9	1.2	*
10	Anemias (D50-D64)	8	1.1	*
11	Cerebrovascular diseases (I60-I69)	7	1.0	*
11	Certain conditions originating in the perinatal period (P00-P96)	7	1.0	*
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	5	0.7	*
13	Diabetes mellitus (E10-E14)	5	0.7	*
15	Diseases of appendix (K35-K38)	3	0.4	*
...	All other causes (residual)	124	17.1	3.6

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2014.

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex:
United States, 2014

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
Black, male, 15-24 years				
...	All causes	5,058	100.0	135.4
1	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	2,428	48.0	65.0
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,121	22.2	30.0
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	450	8.9	12.0
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	172	3.4	4.6
5	Malignant neoplasms (C00-C97)	161	3.2	4.3
6	Human immunodeficiency virus (HIV) disease (B20-B24)	52	1.0	1.4
7	Chronic lower respiratory diseases (J40-J47)	47	0.9	1.3
8	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	38	0.8	1.0
9	Anemias (D50-D64)	33	0.7	0.9
9	Diabetes mellitus (E10-E14)	33	0.7	0.9
11	Cerebrovascular diseases (I60-I69)	29	0.6	0.8
11	Legal intervention (Y35,Y89.0)	29	0.6	0.8
13	Influenza and pneumonia (J09-J18)	27	0.5	0.7
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	17	0.3	*
15	Septicemia (A40-A41)	12	0.2	*
...	All other causes (residual)	409	8.1	11.0
Black, male, 25-34 years				
...	All causes	6,622	100.0	212.1
1	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	2,115	31.9	67.7
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,465	22.1	46.9
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	629	9.5	20.1
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	477	7.2	15.3
5	Malignant neoplasms (C00-C97)	258	3.9	8.3
6	Human immunodeficiency virus (HIV) disease (B20-B24)	253	3.8	8.1
7	Diabetes mellitus (E10-E14)	130	2.0	4.2
8	Cerebrovascular diseases (I60-I69)	91	1.4	2.9
9	Anemias (D50-D64)	80	1.2	2.6
10	Chronic lower respiratory diseases (J40-J47)	74	1.1	2.4
11	Influenza and pneumonia (J09-J18)	48	0.7	1.5
12	Legal intervention (Y35,Y89.0)	45	0.7	1.4
13	Septicemia (A40-A41)	43	0.6	1.4
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	43	0.6	1.4
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	33	0.5	1.1
...	All other causes (residual)	838	12.7	26.8

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2014.

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex:
United States, 2014

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
Black, male, 35-44 years				
...	All causes	8,210	100.0	308.5
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,717	20.9	64.5
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,291	15.7	48.5
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,056	12.9	39.7
4	Malignant neoplasms (C00-C97)	766	9.3	28.8
5	Human immunodeficiency virus (HIV) disease (B20-B24)	366	4.5	13.8
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	353	4.3	13.3
7	Cerebrovascular diseases (I60-I69)	304	3.7	11.4
8	Diabetes mellitus (E10-E14)	301	3.7	11.3
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	150	1.8	5.6
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	129	1.6	4.8
11	Influenza and pneumonia (J09-J18)	103	1.3	3.9
12	Chronic lower respiratory diseases (J40-J47)	101	1.2	3.8
13	Septicemia (A40-A41)	93	1.1	3.5
14	Chronic liver disease and cirrhosis (K70,K73-K74)	91	1.1	3.4
15	Aortic aneurysm and dissection (I71)	76	0.9	2.9
...	All other causes (residual)	1,313	16.0	49.3
Black, male, 45-54 years				
...	All causes	18,011	100.0	671.8
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	4,812	26.7	179.5
2	Malignant neoplasms (C00-C97)	3,452	19.2	128.8
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,728	9.6	64.5
4	Diabetes mellitus (E10-E14)	824	4.6	30.7
5	Cerebrovascular diseases (I60-I69)	794	4.4	29.6
6	Human immunodeficiency virus (HIV) disease (B20-B24)	722	4.0	26.9
7	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	594	3.3	22.2
8	Chronic liver disease and cirrhosis (K70,K73-K74)	430	2.4	16.0
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	365	2.0	13.6
10	Septicemia (A40-A41)	303	1.7	11.3
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	295	1.6	11.0
12	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	285	1.6	10.6
13	Chronic lower respiratory diseases (J40-J47)	280	1.6	10.4
14	Influenza and pneumonia (J09-J18)	264	1.5	9.8
15	Viral hepatitis (B15-B19)	120	0.7	4.5
...	All other causes (residual)	2,743	15.2	102.3

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2014.

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex:
United States, 2014

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
Black, male, 55-64 years				
...	All causes	34,843	100.0	1,611.5
1	Malignant neoplasms (C00-C97)	9,741	28.0	450.5
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	9,199	26.4	425.5
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,726	5.0	79.8
4	Cerebrovascular diseases (I60-I69)	1,716	4.9	79.4
5	Diabetes mellitus (E10-E14)	1,601	4.6	74.0
6	Chronic lower respiratory diseases (J40-J47)	939	2.7	43.4
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	889	2.6	41.1
8	Chronic liver disease and cirrhosis (K70,K73-K74)	863	2.5	39.9
9	Human immunodeficiency virus (HIV) disease (B20-B24)	703	2.0	32.5
10	Septicemia (A40-A41)	684	2.0	31.6
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	589	1.7	27.2
12	Influenza and pneumonia (J09-J18)	528	1.5	24.4
13	Viral hepatitis (B15-B19)	444	1.3	20.5
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	288	0.8	13.3
15	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	194	0.6	9.0
...	All other causes (residual)	4,739	13.6	219.2
Black, male, 65-74 years				
...	All causes	33,499	100.0	3,047.4
1	Malignant neoplasms (C00-C97)	10,177	30.4	925.8
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	8,735	26.1	794.6
3	Cerebrovascular diseases (I60-I69)	1,867	5.6	169.8
4	Diabetes mellitus (E10-E14)	1,656	4.9	150.6
5	Chronic lower respiratory diseases (J40-J47)	1,464	4.4	133.2
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,004	3.0	91.3
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	784	2.3	71.3
8	Septicemia (A40-A41)	756	2.3	68.8
9	Influenza and pneumonia (J09-J18)	597	1.8	54.3
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	549	1.6	49.9
11	Chronic liver disease and cirrhosis (K70,K73-K74)	419	1.3	38.1
12	Pneumonitis due to solids and liquids (J69)	230	0.7	20.9
13	Human immunodeficiency virus (HIV) disease (B20-B24)	217	0.6	19.7
14	Alzheimer's disease (G30)	193	0.6	17.6
15	Viral hepatitis (B15-B19)	166	0.5	15.1
...	All other causes (residual)	4,685	14.0	426.2

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2014.

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex:
United States, 2014

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
Black, male, 75-84 years				
...	All causes	28,161	100.0	6,172.6
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	7,534	26.8	1,651.4
2	Malignant neoplasms (C00-C97)	7,160	25.4	1,569.4
3	Cerebrovascular diseases (I60-I69)	1,794	6.4	393.2
4	Diabetes mellitus (E10-E14)	1,291	4.6	283.0
5	Chronic lower respiratory diseases (J40-J47)	1,280	4.5	280.6
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	908	3.2	199.0
7	Alzheimer's disease (G30)	722	2.6	158.3
8	Septicemia (A40-A41)	640	2.3	140.3
9	Influenza and pneumonia (J09-J18)	588	2.1	128.9
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	511	1.8	112.0
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	480	1.7	105.2
12	Parkinson's disease (G20-G21)	293	1.0	64.2
13	Pneumonitis due to solids and liquids (J69)	256	0.9	56.1
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	135	0.5	29.6
15	Chronic liver disease and cirrhosis (K70,K73-K74)	99	0.4	21.7
...	All other causes (residual)	4,470	15.9	979.8
Black, male, 85 years and over				
...	All causes	18,100	100.0	13,291.7
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	5,029	27.8	3,693.0
2	Malignant neoplasms (C00-C97)	3,239	17.9	2,378.6
3	Cerebrovascular diseases (I60-I69)	1,127	6.2	827.6
4	Alzheimer's disease (G30)	877	4.8	644.0
5	Chronic lower respiratory diseases (J40-J47)	782	4.3	574.3
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	640	3.5	470.0
7	Diabetes mellitus (E10-E14)	609	3.4	447.2
8	Influenza and pneumonia (J09-J18)	524	2.9	384.8
9	Septicemia (A40-A41)	395	2.2	290.1
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	372	2.1	273.2
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	302	1.7	221.8
12	Parkinson's disease (G20-G21)	216	1.2	158.6
13	Pneumonitis due to solids and liquids (J69)	208	1.1	152.7
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	89	0.5	65.4
15	Nutritional deficiencies (E40-E64)	70	0.4	51.4
...	All other causes (residual)	3,621	20.0	2,659.1

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2014.

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex:
United States, 2014

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
Black, female, all ages				
...	All causes	151,227	100.0	655.5
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	35,133	23.2	152.3
2	Malignant neoplasms (C00-C97)	34,029	22.5	147.5
3	Cerebrovascular diseases (I60-I69)	9,341	6.2	40.5
4	Diabetes mellitus (E10-E14)	6,983	4.6	30.3
5	Chronic lower respiratory diseases (J40-J47)	4,899	3.2	21.2
6	Alzheimer's disease (G30)	4,730	3.1	20.5
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	4,598	3.0	19.9
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	4,552	3.0	19.7
9	Septicemia (A40-A41)	3,417	2.3	14.8
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	2,918	1.9	12.6
11	Influenza and pneumonia (J09-J18)	2,875	1.9	12.5
12	Certain conditions originating in the perinatal period (P00-P96)	1,766	1.2	7.7
13	Human immunodeficiency virus (HIV) disease (B20-B24)	1,218	0.8	5.3
14	Chronic liver disease and cirrhosis (K70,K73-K74)	1,163	0.8	5.0
15	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,080	0.7	4.7
...	All other causes (residual)	32,525	21.5	141.0
Black, female, 1-4 years				
...	All causes	426	100.0	31.9
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	112	26.3	8.4
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	70	16.4	5.2
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	45	10.6	3.4
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	29	6.8	2.2
5	Malignant neoplasms (C00-C97)	26	6.1	1.9
6	Chronic lower respiratory diseases (J40-J47)	12	2.8	*
7	Influenza and pneumonia (J09-J18)	10	2.3	*
8	Septicemia (A40-A41)	5	1.2	*
9	Cerebrovascular diseases (I60-I69)	4	0.9	*
9	Complications of medical and surgical care (Y40-Y84,Y88)	4	0.9	*
11	Acute bronchitis and bronchiolitis (J20-J21)	3	0.7	*
11	Certain conditions originating in the perinatal period (P00-P96)	3	0.7	*
13	Salmonella infections (A01-A02)	1	0.2	*
13	Meningococcal infection (A39)	1	0.2	*
13	Malaria (B50-B54)	1	0.2	*
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1	0.2	*
13	Anemias (D50-D64)	1	0.2	*
13	Nutritional deficiencies (E40-E64)	1	0.2	*
13	Meningitis (G00,G03)	1	0.2	*
13	Pneumonitis due to solids and liquids (J69)	1	0.2	*
...	All other causes (residual)	95	22.3	7.1

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2014.

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex:
United States, 2014

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
Black, female, 5-14 years				
...	All causes	479	100.0	14.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	111	23.2	3.3
2	Malignant neoplasms (C00-C97)	69	14.4	2.1
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	35	7.3	1.0
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	30	6.3	0.9
5	Chronic lower respiratory diseases (J40-J47)	24	5.0	0.7
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	21	4.4	0.6
7	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	16	3.3	*
7	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	16	3.3	*
9	Anemias (D50-D64)	10	2.1	*
10	Cerebrovascular diseases (I60-I69)	8	1.7	*
11	Septicemia (A40-A41)	7	1.5	*
11	Influenza and pneumonia (J09-J18)	7	1.5	*
13	Diabetes mellitus (E10-E14)	6	1.3	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3	0.6	*
14	Certain conditions originating in the perinatal period (P00-P96)	3	0.6	*
14	Complications of medical and surgical care (Y40-Y84,Y88)	3	0.6	*
...	All other causes (residual)	110	23.0	3.3
Black, female, 15-24 years				
...	All causes	1,544	100.0	42.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	388	25.1	10.7
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	255	16.5	7.0
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	112	7.3	3.1
4	Malignant neoplasms (C00-C97)	111	7.2	3.1
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	96	6.2	2.6
6	Pregnancy, childbirth and the puerperium (O00-O99)	57	3.7	1.6
7	Human immunodeficiency virus (HIV) disease (B20-B24)	34	2.2	0.9
8	Anemias (D50-D64)	32	2.1	0.9
9	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	29	1.9	0.8
10	Diabetes mellitus (E10-E14)	27	1.7	0.7
11	Chronic lower respiratory diseases (J40-J47)	25	1.6	0.7
12	Septicemia (A40-A41)	20	1.3	0.6
12	Cerebrovascular diseases (I60-I69)	20	1.3	0.6
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	12	0.8	*
15	Influenza and pneumonia (J09-J18)	11	0.7	*
...	All other causes (residual)	315	20.4	8.7

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2014.

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex:
United States, 2014

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
Black, female, 25-34 years				
...	All causes	2,927	100.0	88.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	491	16.8	14.9
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	381	13.0	11.5
3	Malignant neoplasms (C00-C97)	371	12.7	11.2
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	253	8.6	7.7
5	Pregnancy, childbirth and the puerperium (O00-O99)	143	4.9	4.3
6	Human immunodeficiency virus (HIV) disease (B20-B24)	123	4.2	3.7
7	Diabetes mellitus (E10-E14)	109	3.7	3.3
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	91	3.1	2.8
9	Cerebrovascular diseases (I60-I69)	64	2.2	1.9
10	Chronic lower respiratory diseases (J40-J47)	52	1.8	1.6
11	Influenza and pneumonia (J09-J18)	51	1.7	1.5
12	Anemias (D50-D64)	47	1.6	1.4
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	41	1.4	1.2
14	Septicemia (A40-A41)	38	1.3	1.1
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	30	1.0	0.9
...	All other causes (residual)	642	21.9	19.4
Black, female, 35-44 years				
...	All causes	5,787	100.0	193.9
1	Malignant neoplasms (C00-C97)	1,210	20.9	40.5
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,058	18.3	35.5
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	562	9.7	18.8
4	Human immunodeficiency virus (HIV) disease (B20-B24)	274	4.7	9.2
5	Diabetes mellitus (E10-E14)	246	4.3	8.2
6	Cerebrovascular diseases (I60-I69)	212	3.7	7.1
7	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	166	2.9	5.6
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	115	2.0	3.9
9	Pregnancy, childbirth and the puerperium (O00-O99)	103	1.8	3.5
10	Influenza and pneumonia (J09-J18)	98	1.7	3.3
11	Septicemia (A40-A41)	96	1.7	3.2
12	Chronic liver disease and cirrhosis (K70,K73-K74)	93	1.6	3.1
13	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	89	1.5	3.0
14	Chronic lower respiratory diseases (J40-J47)	87	1.5	2.9
15	Anemias (D50-D64)	69	1.2	2.3
...	All other causes (residual)	1,309	22.6	43.9

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2014.

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex:
United States, 2014

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
Black, female, 45-54 years				
...	All causes	13,804	100.0	455.8
1	Malignant neoplasms (C00-C97)	4,040	29.3	133.4
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,875	20.8	94.9
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	800	5.8	26.4
4	Cerebrovascular diseases (I60-I69)	679	4.9	22.4
5	Diabetes mellitus (E10-E14)	615	4.5	20.3
6	Human immunodeficiency virus (HIV) disease (B20-B24)	378	2.7	12.5
7	Chronic lower respiratory diseases (J40-J47)	371	2.7	12.2
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	295	2.1	9.7
9	Chronic liver disease and cirrhosis (K70,K73-K74)	294	2.1	9.7
10	Septicemia (A40-A41)	290	2.1	9.6
11	Influenza and pneumonia (J09-J18)	236	1.7	7.8
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	190	1.4	6.3
13	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	134	1.0	4.4
14	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	91	0.7	3.0
15	Anemias (D50-D64)	79	0.6	2.6
...	All other causes (residual)	2,437	17.7	80.5
Black, female, 55-64 years				
...	All causes	25,129	100.0	974.8
1	Malignant neoplasms (C00-C97)	8,299	33.0	321.9
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	5,439	21.6	211.0
3	Diabetes mellitus (E10-E14)	1,299	5.2	50.4
4	Cerebrovascular diseases (I60-I69)	1,247	5.0	48.4
5	Chronic lower respiratory diseases (J40-J47)	893	3.6	34.6
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	736	2.9	28.6
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	694	2.8	26.9
8	Septicemia (A40-A41)	639	2.5	24.8
9	Chronic liver disease and cirrhosis (K70,K73-K74)	449	1.8	17.4
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	429	1.7	16.6
11	Influenza and pneumonia (J09-J18)	408	1.6	15.8
12	Human immunodeficiency virus (HIV) disease (B20-B24)	293	1.2	11.4
13	Viral hepatitis (B15-B19)	207	0.8	8.0
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	101	0.4	3.9
15	Pneumonitis due to solids and liquids (J69)	81	0.3	3.1
...	All other causes (residual)	3915	15.6	151.9

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2014.

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex:
United States, 2014

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
Black, female, 65-74 years				
...	All causes	27,456	100.0	1,880.2
1	Malignant neoplasms (C00-C97)	8,523	31.0	583.7
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	6,395	23.3	437.9
3	Cerebrovascular diseases (I60-I69)	1,644	6.0	112.6
4	Diabetes mellitus (E10-E14)	1,557	5.7	106.6
5	Chronic lower respiratory diseases (J40-J47)	1,125	4.1	77.0
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,078	3.9	73.8
7	Septicemia (A40-A41)	725	2.6	49.6
8	Essential hypertension and hypertensive renal disease (I10,I12,I15)	475	1.7	32.5
9	Influenza and pneumonia (J09-J18)	415	1.5	28.4
10	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	398	1.4	27.3
11	Alzheimer's disease (G30)	298	1.1	20.4
12	Chronic liver disease and cirrhosis (K70,K73-K74)	178	0.6	12.2
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	160	0.6	11.0
14	Pneumonitis due to solids and liquids (J69)	130	0.5	8.9
15	Aortic aneurysm and dissection (I71)	91	0.3	6.2
...	All other causes (residual)	4,264	15.5	292.0
Black, female, 75-84 years				
...	All causes	32,267	100.0	4,356.8
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	8,041	24.9	1,085.7
2	Malignant neoplasms (C00-C97)	7,116	22.1	960.8
3	Cerebrovascular diseases (I60-I69)	2,389	7.4	322.6
4	Diabetes mellitus (E10-E14)	1,721	5.3	232.4
5	Alzheimer's disease (G30)	1,351	4.2	182.4
6	Chronic lower respiratory diseases (J40-J47)	1,278	4.0	172.6
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,175	3.6	158.7
8	Septicemia (A40-A41)	747	2.3	100.9
9	Essential hypertension and hypertensive renal disease (I10,I12,I15)	707	2.2	95.5
10	Influenza and pneumonia (J09-J18)	628	1.9	84.8
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	381	1.2	51.4
12	Pneumonitis due to solids and liquids (J69)	230	0.7	31.1
13	Parkinson's disease (G20-G21)	196	0.6	26.5
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	179	0.6	24.2
15	Aortic aneurysm and dissection (I71)	109	0.3	14.7
...	All other causes (residual)	6,019	18.7	812.7

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2014.

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex:
United States, 2014

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
Black, female, 85 years and over				
...	All causes	38,227	100.0	11,656.8
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	10,723	28.1	3,269.8
2	Malignant neoplasms (C00-C97)	4,255	11.1	1,297.5
3	Cerebrovascular diseases (I60-I69)	3,063	8.0	934.0
4	Alzheimer's disease (G30)	3,019	7.9	920.6
5	Diabetes mellitus (E10-E14)	1,403	3.7	427.8
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,131	3.0	344.9
7	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,030	2.7	314.1
7	Chronic lower respiratory diseases (J40-J47)	1,030	2.7	314.1
9	Influenza and pneumonia (J09-J18)	965	2.5	294.3
10	Septicemia (A40-A41)	824	2.2	251.3
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	432	1.1	131.7
12	Pneumonitis due to solids and liquids (J69)	313	0.8	95.4
13	Parkinson's disease (G20-G21)	204	0.5	62.2
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	192	0.5	58.5
15	Atherosclerosis (I70)	136	0.4	41.5
...	All other causes (residual)	9,507	24.9	2,899.0
American Indian, both sexes, all ages				
...	All causes	18,008	100.0	398.5
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,288	18.3	72.8
2	Malignant neoplasms (C00-C97)	3,153	17.5	69.8
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,996	11.1	44.2
4	Chronic liver disease and cirrhosis (K70,K73-K74)	951	5.3	21.0
5	Diabetes mellitus (E10-E14)	945	5.2	20.9
6	Chronic lower respiratory diseases (J40-J47)	788	4.4	17.4
7	Cerebrovascular diseases (I60-I69)	649	3.6	14.4
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	489	2.7	10.8
9	Influenza and pneumonia (J09-J18)	412	2.3	9.1
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	338	1.9	7.5
11	Alzheimer's disease (G30)	304	1.7	6.7
12	Septicemia (A40-A41)	289	1.6	6.4
13	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	264	1.5	5.8
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	176	1.0	3.9
15	Certain conditions originating in the perinatal period (P00-P96)	142	0.8	3.1
...	All other causes (residual)	3,824	21.2	84.6

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2014.

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex:
United States, 2014

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, both sexes, 1-4 years				
...	All causes	95	100.0	30.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	36	37.9	11.7
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	8	8.4	*
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	6	6.3	*
4	Malignant neoplasms (C00-C97)	5	5.3	*
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	4	4.2	*
6	Septicemia (A40-A41)	2	2.1	*
6	Chronic lower respiratory diseases (J40-J47)	2	2.1	*
6	Certain conditions originating in the perinatal period (P00-P96)	2	2.1	*
9	Meningococcal infection (A39)	1	1.1	*
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1	1.1	*
9	Cerebrovascular diseases (I60-I69)	1	1.1	*
9	Influenza and pneumonia (J09-J18)	1	1.1	*
...	All other causes (residual)	26	27.4	8.4
American Indian, both sexes, 5-14 years				
...	All causes	89	100.0	11.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	37	41.6	4.8
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	12	13.5	*
3	Malignant neoplasms (C00-C97)	7	7.9	*
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	7	7.9	*
5	Influenza and pneumonia (J09-J18)	4	4.5	*
5	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	4	4.5	*
7	Diabetes mellitus (E10-E14)	1	1.1	*
7	Diseases of heart (I00-I09,I11,I13,I20-I51)	1	1.1	*
7	Cerebrovascular diseases (I60-I69)	1	1.1	*
7	Chronic lower respiratory diseases (J40-J47)	1	1.1	*
7	Diseases of appendix (K35-K38)	1	1.1	*
7	Complications of medical and surgical care (Y40-Y84,Y88)	1	1.1	*
...	All other causes (residual)	12	13.5	*

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2014.

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex:
United States, 2014

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, both sexes, 15-24 years				
...	All causes	586	100.0	76.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	282	48.1	36.6
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	129	22.0	16.7
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	56	9.6	7.3
4	Malignant neoplasms (C00-C97)	14	2.4	*
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	13	2.2	*
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	8	1.4	*
7	Chronic lower respiratory diseases (J40-J47)	5	0.9	*
8	Influenza and pneumonia (J09-J18)	4	0.7	*
8	Pregnancy, childbirth and the puerperium (O00-O99)	4	0.7	*
10	Chronic liver disease and cirrhosis (K70,K73-K74)	3	0.5	*
10	Legal intervention (Y35,Y89.0)	3	0.5	*
12	Septicemia (A40-A41)	2	0.3	*
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1	0.2	*
13	Diabetes mellitus (E10-E14)	1	0.2	*
13	Meningitis (G00,G03)	1	0.2	*
13	Cerebrovascular diseases (I60-I69)	1	0.2	*
13	Aortic aneurysm and dissection (I71)	1	0.2	*
13	Pneumonitis due to solids and liquids (J69)	1	0.2	*
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1	0.2	*
...	All other causes (residual)	56	9.6	7.3
American Indian, both sexes, 25-34 years				
...	All causes	943	100.0	135.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	380	40.3	54.5
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	133	14.1	19.1
3	Chronic liver disease and cirrhosis (K70,K73-K74)	73	7.7	10.5
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	72	7.6	10.3
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	50	5.3	7.2
6	Malignant neoplasms (C00-C97)	24	2.5	3.4
7	Diabetes mellitus (E10-E14)	16	1.7	*
8	Influenza and pneumonia (J09-J18)	11	1.2	*
9	Cerebrovascular diseases (I60-I69)	10	1.1	*
10	Septicemia (A40-A41)	7	0.7	*
11	Pregnancy, childbirth and the puerperium (O00-O99)	5	0.5	*
11	Legal intervention (Y35,Y89.0)	5	0.5	*
13	Human immunodeficiency virus (HIV) disease (B20-B24)	3	0.3	*
13	Chronic lower respiratory diseases (J40-J47)	3	0.3	*
13	Pneumonitis due to solids and liquids (J69)	3	0.3	*
13	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	3	0.3	*
...	All other causes (residual)	145	15.4	20.8

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2014.

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex:
United States, 2014

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, both sexes, 35-44 years				
...	All causes	1,320	100.0	222.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	354	26.8	59.7
2	Chronic liver disease and cirrhosis (K70,K73-K74)	175	13.3	29.5
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	145	11.0	24.4
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	102	7.7	17.2
5	Malignant neoplasms (C00-C97)	97	7.3	16.4
6	Diabetes mellitus (E10-E14)	53	4.0	8.9
7	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	44	3.3	7.4
8	Influenza and pneumonia (J09-J18)	32	2.4	5.4
9	Septicemia (A40-A41)	25	1.9	4.2
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	13	1.0	*
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	10	0.8	*
11	Cerebrovascular diseases (I60-I69)	10	0.8	*
13	Viral hepatitis (B15-B19)	8	0.6	*
13	Chronic lower respiratory diseases (J40-J47)	8	0.6	*
15	Pregnancy, childbirth and the puerperium (O00-O99)	6	0.5	*
...	All other causes (residual)	238	18.0	40.1
American Indian, both sexes, 45-54 years				
...	All causes	2,376	100.0	431.0
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	376	15.8	68.2
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	375	15.8	68.0
3	Malignant neoplasms (C00-C97)	347	14.6	62.9
4	Chronic liver disease and cirrhosis (K70,K73-K74)	280	11.8	50.8
5	Diabetes mellitus (E10-E14)	125	5.3	22.7
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	61	2.6	11.1
7	Cerebrovascular diseases (I60-I69)	54	2.3	9.8
7	Influenza and pneumonia (J09-J18)	54	2.3	9.8
9	Chronic lower respiratory diseases (J40-J47)	47	2.0	8.5
10	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	42	1.8	7.6
11	Septicemia (A40-A41)	40	1.7	7.3
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	40	1.7	7.3
13	Viral hepatitis (B15-B19)	34	1.4	6.2
14	Human immunodeficiency virus (HIV) disease (B20-B24)	24	1.0	4.4
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	10	0.4	*
...	All other causes (residual)	467	19.7	84.7

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2014.

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex:
United States, 2014

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, both sexes, 55-64 years				
...	All causes	3,302	100.0	787.9
1	Malignant neoplasms (C00-C97)	749	22.7	178.7
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	634	19.2	151.3
3	Chronic liver disease and cirrhosis (K70,K73-K74)	260	7.9	62.0
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	250	7.6	59.7
5	Diabetes mellitus (E10-E14)	236	7.1	56.3
6	Chronic lower respiratory diseases (J40-J47)	134	4.1	32.0
7	Cerebrovascular diseases (I60-I69)	107	3.2	25.5
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	64	1.9	15.3
9	Viral hepatitis (B15-B19)	58	1.8	13.8
10	Influenza and pneumonia (J09-J18)	57	1.7	13.6
11	Septicemia (A40-A41)	56	1.7	13.4
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	36	1.1	8.6
13	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	30	0.9	7.2
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	17	0.5	*
15	Pneumonitis due to solids and liquids (J69)	16	0.5	*
...	All other causes (residual)	598	18.1	142.7
American Indian, both sexes, 65-74 years				
...	All causes	3,430	100.0	1,585.8
1	Malignant neoplasms (C00-C97)	942	27.5	435.5
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	760	22.2	351.4
3	Diabetes mellitus (E10-E14)	241	7.0	111.4
4	Chronic lower respiratory diseases (J40-J47)	232	6.8	107.3
5	Cerebrovascular diseases (I60-I69)	136	4.0	62.9
6	Chronic liver disease and cirrhosis (K70,K73-K74)	121	3.5	55.9
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	106	3.1	49.0
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	79	2.3	36.5
9	Influenza and pneumonia (J09-J18)	68	2.0	31.4
10	Septicemia (A40-A41)	61	1.8	28.2
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	36	1.0	16.6
12	Alzheimer's disease (G30)	26	0.8	12.0
13	Pneumonitis due to solids and liquids (J69)	20	0.6	9.2
14	Enterocolitis due to <i>Clostridium difficile</i> (A04.7)	17	0.5	*
15	Viral hepatitis (B15-B19)	16	0.5	*
15	Parkinson's disease (G20-G21)	16	0.5	*
...	All other causes (residual)	553	16.1	255.7

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2014.

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex:
United States, 2014

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, both sexes, 75-84 years				
...	All causes	3,149	100.0	3,598.0
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	695	22.1	794.1
2	Malignant neoplasms (C00-C97)	686	21.8	783.8
3	Chronic lower respiratory diseases (J40-J47)	221	7.0	252.5
4	Diabetes mellitus (E10-E14)	180	5.7	205.7
5	Cerebrovascular diseases (I60-I69)	164	5.2	187.4
6	Alzheimer's disease (G30)	104	3.3	118.8
7	Influenza and pneumonia (J09-J18)	89	2.8	101.7
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	87	2.8	99.4
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	86	2.7	98.3
10	Septicemia (A40-A41)	59	1.9	67.4
11	Parkinson's disease (G20-G21)	39	1.2	44.6
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	38	1.2	43.4
13	Pneumonitis due to solids and liquids (J69)	32	1.0	36.6
14	Chronic liver disease and cirrhosis (K70,K73-K74)	29	0.9	33.1
15	Aortic aneurysm and dissection (I71)	20	0.6	22.9
...	All other causes (residual)	620	19.7	708.4
American Indian, both sexes, 85 years and over				
...	All causes	2,358	100.0	8,149.9
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	609	25.8	2,104.9
2	Malignant neoplasms (C00-C97)	282	12.0	974.7
3	Alzheimer's disease (G30)	168	7.1	580.7
4	Cerebrovascular diseases (I60-I69)	163	6.9	563.4
5	Chronic lower respiratory diseases (J40-J47)	134	5.7	463.1
6	Diabetes mellitus (E10-E14)	92	3.9	318.0
7	Influenza and pneumonia (J09-J18)	87	3.7	300.7
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	65	2.8	224.7
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	52	2.2	179.7
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	46	2.0	159.0
11	Septicemia (A40-A41)	34	1.4	117.5
12	Parkinson's disease (G20-G21)	27	1.1	93.3
13	Pneumonitis due to solids and liquids (J69)	22	0.9	76.0
14	Nutritional deficiencies (E40-E64)	15	0.6	*
15	Chronic liver disease and cirrhosis (K70,K73-K74)	10	0.4	*
15	Enterocolitis due to <i>Clostridium difficile</i> (A04.7)	10	0.4	*
...	All other causes (residual)	542	23.0	1,873.3

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2014.

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex:
United States, 2014

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, male, all ages				
...	All causes	9,829	100.0	433.2
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,915	19.5	84.4
2	Malignant neoplasms (C00-C97)	1,727	17.6	76.1
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,331	13.5	58.7
4	Diabetes mellitus (E10-E14)	504	5.1	22.2
5	Chronic liver disease and cirrhosis (K70,K73-K74)	481	4.9	21.2
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	363	3.7	16.0
7	Chronic lower respiratory diseases (J40-J47)	360	3.7	15.9
8	Cerebrovascular diseases (I60-I69)	296	3.0	13.0
9	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	209	2.1	9.2
10	Influenza and pneumonia (J09-J18)	195	2.0	8.6
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	163	1.7	7.2
12	Septicemia (A40-A41)	123	1.3	5.4
13	Viral hepatitis (B15-B19)	86	0.9	3.8
14	Alzheimer's disease (G30)	80	0.8	3.5
14	Certain conditions originating in the perinatal period (P00-P96)	80	0.8	3.5
...	All other causes (residual)	1,916	19.5	84.4
American Indian, male, 1-4 years				
...	All causes	64	100.0	41.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	24	37.5	15.4
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	8	12.5	*
3	Malignant neoplasms (C00-C97)	5	7.8	*
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	3	4.7	*
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	3	4.7	*
6	Chronic lower respiratory diseases (J40-J47)	2	3.1	*
7	Meningococcal infection (A39)	1	1.6	*
7	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1	1.6	*
7	Cerebrovascular diseases (I60-I69)	1	1.6	*
7	Certain conditions originating in the perinatal period (P00-P96)	1	1.6	*
...	All other causes (residual)	15	23.4	*

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2014.

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex:
United States, 2014

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, male, 5-14 years				
...	All causes	48	100.0	12.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	23	47.9	5.9
2	Malignant neoplasms (C00-C97)	4	8.3	*
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	4	8.3	*
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	3	6.3	*
5	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	2	4.2	*
6	Cerebrovascular diseases (I60-I69)	1	2.1	*
6	Influenza and pneumonia (J09-J18)	1	2.1	*
6	Chronic lower respiratory diseases (J40-J47)	1	2.1	*
6	Diseases of appendix (K35-K38)	1	2.1	*
6	Complications of medical and surgical care (Y40-Y84,Y88)	1	2.1	*
...	All other causes (residual)	7	14.6	*
American Indian, male, 15-24 years				
...	All causes	410	100.0	103.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	190	46.3	48.1
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	93	22.7	23.5
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	48	11.7	12.1
4	Malignant neoplasms (C00-C97)	10	2.4	*
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	10	2.4	*
6	Influenza and pneumonia (J09-J18)	4	1.0	*
7	Chronic lower respiratory diseases (J40-J47)	3	0.7	*
7	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	3	0.7	*
7	Legal intervention (Y35,Y89.0)	3	0.7	*
10	Chronic liver disease and cirrhosis (K70,K73-K74)	2	0.5	*
11	Diabetes mellitus (E10-E14)	1	0.2	*
11	Meningitis (G00,G03)	1	0.2	*
11	Cerebrovascular diseases (I60-I69)	1	0.2	*
11	Aortic aneurysm and dissection (I71)	1	0.2	*
...	All other causes (residual)	40	9.8	10.1

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2014.

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex:
United States, 2014

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, male, 25-34 years				
...	All causes	648	100.0	179.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	272	42.0	75.2
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	100	15.4	27.6
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	56	8.6	15.5
4	Chronic liver disease and cirrhosis (K70,K73-K74)	41	6.3	11.3
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	35	5.4	9.7
6	Malignant neoplasms (C00-C97)	17	2.6	*
7	Diabetes mellitus (E10-E14)	10	1.5	*
8	Cerebrovascular diseases (I60-I69)	6	0.9	*
8	Influenza and pneumonia (J09-J18)	6	0.9	*
10	Legal intervention (Y35,Y89.0)	5	0.8	*
11	Septicemia (A40-A41)	3	0.5	*
11	Pneumonitis due to solids and liquids (J69)	3	0.5	*
11	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	3	0.5	*
14	Human immunodeficiency virus (HIV) disease (B20-B24)	2	0.3	*
15	Viral hepatitis (B15-B19)	1	0.2	*
15	Anemias (D50-D64)	1	0.2	*
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1	0.2	*
15	Aortic aneurysm and dissection (I71)	1	0.2	*
15	Peptic ulcer (K25-K28)	1	0.2	*
15	Enterocolitis due to <i>Clostridium difficile</i> (A04.7)	1	0.2	*
...	All other causes (residual)	83	12.8	22.9
American Indian, male, 35-44 years				
...	All causes	799	100.0	264.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	233	29.2	77.1
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	99	12.4	32.8
3	Chronic liver disease and cirrhosis (K70,K73-K74)	84	10.5	27.8
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	74	9.3	24.5
5	Malignant neoplasms (C00-C97)	46	5.8	15.2
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	34	4.3	11.3
7	Diabetes mellitus (E10-E14)	32	4.0	10.6
8	Influenza and pneumonia (J09-J18)	16	2.0	*
9	Septicemia (A40-A41)	13	1.6	*
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	8	1.0	*
11	Cerebrovascular diseases (I60-I69)	7	0.9	*
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	5	0.6	*
13	Viral hepatitis (B15-B19)	4	0.5	*
13	Human immunodeficiency virus (HIV) disease (B20-B24)	4	0.5	*
15	Chronic lower respiratory diseases (J40-J47)	3	0.4	*
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	3	0.4	*
15	Legal intervention (Y35,Y89.0)	3	0.4	*
...	All other causes (residual)	131	16.4	43.4

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2014.

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex:
United States, 2014

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, male, 45-54 years				
...	All causes	1,394	100.0	508.5
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	263	18.9	95.9
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	260	18.7	94.8
3	Malignant neoplasms (C00-C97)	161	11.5	58.7
4	Chronic liver disease and cirrhosis (K70,K73-K74)	137	9.8	50.0
5	Diabetes mellitus (E10-E14)	73	5.2	26.6
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	49	3.5	17.9
7	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	33	2.4	12.0
8	Influenza and pneumonia (J09-J18)	28	2.0	10.2
9	Viral hepatitis (B15-B19)	27	1.9	9.8
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	24	1.7	8.8
11	Cerebrovascular diseases (I60-I69)	23	1.6	8.4
12	Chronic lower respiratory diseases (J40-J47)	21	1.5	7.7
13	Septicemia (A40-A41)	19	1.4	*
14	Human immunodeficiency virus (HIV) disease (B20-B24)	16	1.1	*
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	7	0.5	*
...	All other causes (residual)	253	18.1	92.3
American Indian, male, 55-64 years				
...	All causes	1,981	100.0	984.7
1	Malignant neoplasms (C00-C97)	445	22.5	221.2
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	437	22.1	217.2
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	171	8.6	85.0
4	Chronic liver disease and cirrhosis (K70,K73-K74)	140	7.1	69.6
5	Diabetes mellitus (E10-E14)	138	7.0	68.6
6	Chronic lower respiratory diseases (J40-J47)	63	3.2	31.3
7	Cerebrovascular diseases (I60-I69)	60	3.0	29.8
8	Viral hepatitis (B15-B19)	43	2.2	21.4
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	30	1.5	14.9
10	Influenza and pneumonia (J09-J18)	29	1.5	14.4
11	Septicemia (A40-A41)	25	1.3	12.4
12	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	23	1.2	11.4
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	18	0.9	*
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	13	0.7	*
15	Human immunodeficiency virus (HIV) disease (B20-B24)	10	0.5	*
15	Pneumonitis due to solids and liquids (J69)	10	0.5	*
...	All other causes (residual)	326	16.5	162.0

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2014.

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex:
United States, 2014

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, male, 65-74 years				
...	All causes	1,863	100.0	1,830.2
1	Malignant neoplasms (C00-C97)	530	28.4	520.7
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	452	24.3	444.0
3	Diabetes mellitus (E10-E14)	128	6.9	125.7
4	Chronic lower respiratory diseases (J40-J47)	108	5.8	106.1
5	Cerebrovascular diseases (I60-I69)	76	4.1	74.7
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	62	3.3	60.9
7	Chronic liver disease and cirrhosis (K70,K73-K74)	60	3.2	58.9
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	40	2.1	39.3
9	Influenza and pneumonia (J09-J18)	31	1.7	30.5
10	Septicemia (A40-A41)	25	1.3	24.6
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	18	1.0	*
12	Pneumonitis due to solids and liquids (J69)	14	0.8	*
13	Viral hepatitis (B15-B19)	11	0.6	*
13	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	11	0.6	*
15	Parkinson's disease (G20-G21)	10	0.5	*
...	All other causes (residual)	287	15.4	281.9
American Indian, male, 75-84 years				
...	All causes	1,529	100.0	4,097.9
1	Malignant neoplasms (C00-C97)	379	24.8	1,015.8
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	365	23.9	978.2
3	Chronic lower respiratory diseases (J40-J47)	104	6.8	278.7
4	Diabetes mellitus (E10-E14)	85	5.6	227.8
5	Cerebrovascular diseases (I60-I69)	72	4.7	193.0
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	47	3.1	126.0
7	Influenza and pneumonia (J09-J18)	42	2.7	112.6
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	38	2.5	101.8
9	Alzheimer's disease (G30)	32	2.1	85.8
10	Septicemia (A40-A41)	21	1.4	56.3
11	Parkinson's disease (G20-G21)	19	1.2	*
12	Pneumonitis due to solids and liquids (J69)	17	1.1	*
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	16	1.0	*
14	Chronic liver disease and cirrhosis (K70,K73-K74)	11	0.7	*
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	7	0.5	*
15	Aortic aneurysm and dissection (I71)	7	0.5	*
15	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	7	0.5	*
15	Enterocolitis due to <i>Clostridium difficile</i> (A04.7)	7	0.5	*
...	All other causes (residual)	253	16.5	678.1

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2014.

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex:
United States, 2014

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, male, 85 years and over				
...	All causes	891	100.0	8,610.4
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	250	28.1	2,415.9
2	Malignant neoplasms (C00-C97)	130	14.6	1,256.3
3	Chronic lower respiratory diseases (J40-J47)	55	6.2	531.5
4	Cerebrovascular diseases (I60-I69)	48	5.4	463.9
5	Alzheimer's disease (G30)	38	4.3	367.2
6	Diabetes mellitus (E10-E14)	37	4.2	357.6
7	Influenza and pneumonia (J09-J18)	36	4.0	347.9
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	34	3.8	328.6
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	22	2.5	212.6
10	Parkinson's disease (G20-G21)	17	1.9	*
11	Septicemia (A40-A41)	16	1.8	*
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	11	1.2	*
13	Pneumonitis due to solids and liquids (J69)	10	1.1	*
14	Enterocolitis due to <i>Clostridium difficile</i> (A04.7)	7	0.8	*
15	Atherosclerosis (I70)	6	0.7	*
15	Chronic liver disease and cirrhosis (K70,K73-K74)	6	0.7	*
...	All other causes (residual)	168	18.9	1,623.5
American Indian, female, all ages				
...	All causes	8,179	100.0	363.5
1	Malignant neoplasms (C00-C97)	1,426	17.4	63.4
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,373	16.8	61.0
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	665	8.1	29.6
4	Chronic liver disease and cirrhosis (K70,K73-K74)	470	5.7	20.9
5	Diabetes mellitus (E10-E14)	441	5.4	19.6
6	Chronic lower respiratory diseases (J40-J47)	428	5.2	19.0
7	Cerebrovascular diseases (I60-I69)	353	4.3	15.7
8	Alzheimer's disease (G30)	224	2.7	10.0
9	Influenza and pneumonia (J09-J18)	217	2.7	9.6
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	175	2.1	7.8
11	Septicemia (A40-A41)	166	2.0	7.4
12	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	126	1.5	5.6
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	101	1.2	4.5
14	Certain conditions originating in the perinatal period (P00-P96)	62	0.8	2.8
15	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	55	0.7	2.4
...	All other causes (residual)	1,897	23.2	84.3

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2014.

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex:
United States, 2014

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, female, 1-4 years				
...	All causes	31	100.0	20.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	12	38.7	*
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	3	9.7	*
3	Septicemia (A40-A41)	2	6.5	*
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	1	3.2	*
4	Influenza and pneumonia (J09-J18)	1	3.2	*
4	Certain conditions originating in the perinatal period (P00-P96)	1	3.2	*
...	All other causes (residual)	11	35.5	*
American Indian, female, 5-14 years				
...	All causes	41	100.0	10.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	14	34.1	*
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	8	19.5	*
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	4	9.8	*
4	Malignant neoplasms (C00-C97)	3	7.3	*
4	Influenza and pneumonia (J09-J18)	3	7.3	*
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	2	4.9	*
7	Diabetes mellitus (E10-E14)	1	2.4	*
7	Diseases of heart (I00-I09,I11,I13,I20-I51)	1	2.4	*
...	All other causes (residual)	5	12.2	*
American Indian, female, 15-24 years				
...	All causes	176	100.0	46.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	92	52.3	24.5
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	36	20.5	9.6
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	8	4.5	*
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	5	2.8	*
5	Malignant neoplasms (C00-C97)	4	2.3	*
5	Pregnancy, childbirth and the puerperium (O00-O99)	4	2.3	*
7	Diseases of heart (I00-I09,I11,I13,I20-I51)	3	1.7	*
8	Septicemia (A40-A41)	2	1.1	*
8	Chronic lower respiratory diseases (J40-J47)	2	1.1	*
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1	0.6	*
10	Pneumonitis due to solids and liquids (J69)	1	0.6	*
10	Chronic liver disease and cirrhosis (K70,K73-K74)	1	0.6	*
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1	0.6	*
...	All other causes (residual)	16	9.1	*

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2014.

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex:
United States, 2014

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, female, 25-34 years				
...	All causes	295	100.0	88.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	108	36.6	32.2
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	33	11.2	9.8
3	Chronic liver disease and cirrhosis (K70,K73-K74)	32	10.8	9.5
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	16	5.4	*
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	15	5.1	*
6	Malignant neoplasms (C00-C97)	7	2.4	*
7	Diabetes mellitus (E10-E14)	6	2.0	*
8	Influenza and pneumonia (J09-J18)	5	1.7	*
8	Pregnancy, childbirth and the puerperium (O00-O99)	5	1.7	*
10	Septicemia (A40-A41)	4	1.4	*
10	Cerebrovascular diseases (I60-I69)	4	1.4	*
12	Chronic lower respiratory diseases (J40-J47)	3	1.0	*
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2	0.7	*
14	Viral hepatitis (B15-B19)	1	0.3	*
14	Human immunodeficiency virus (HIV) disease (B20-B24)	1	0.3	*
14	Aortic aneurysm and dissection (I71)	1	0.3	*
14	Infections of kidney (N10-N12,N13.6,N15.1)	1	0.3	*
...	All other causes (residual)	51	17.3	15.2
American Indian, female, 35-44 years				
...	All causes	521	100.0	179.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	121	23.2	41.6
2	Chronic liver disease and cirrhosis (K70,K73-K74)	91	17.5	31.3
3	Malignant neoplasms (C00-C97)	51	9.8	17.5
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	46	8.8	15.8
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	28	5.4	9.6
6	Diabetes mellitus (E10-E14)	21	4.0	7.2
7	Influenza and pneumonia (J09-J18)	16	3.1	*
8	Septicemia (A40-A41)	12	2.3	*
9	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	10	1.9	*
10	Pregnancy, childbirth and the puerperium (O00-O99)	6	1.2	*
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	5	1.0	*
11	Chronic lower respiratory diseases (J40-J47)	5	1.0	*
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	5	1.0	*
14	Viral hepatitis (B15-B19)	4	0.8	*
15	Cerebrovascular diseases (I60-I69)	3	0.6	*
15	Pneumonitis due to solids and liquids (J69)	3	0.6	*
...	All other causes (residual)	94	18.0	32.3

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2014.

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex:
United States, 2014

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, female, 45-54 years				
...	All causes	982	100.0	354.3
1	Malignant neoplasms (C00-C97)	186	18.9	67.1
2	Chronic liver disease and cirrhosis (K70,K73-K74)	143	14.6	51.6
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	115	11.7	41.5
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	113	11.5	40.8
5	Diabetes mellitus (E10-E14)	52	5.3	18.8
6	Cerebrovascular diseases (I60-I69)	31	3.2	11.2
7	Influenza and pneumonia (J09-J18)	26	2.6	9.4
7	Chronic lower respiratory diseases (J40-J47)	26	2.6	9.4
9	Septicemia (A40-A41)	21	2.1	7.6
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	16	1.6	*
11	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	12	1.2	*
12	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	9	0.9	*
13	Human immunodeficiency virus (HIV) disease (B20-B24)	8	0.8	*
14	Viral hepatitis (B15-B19)	7	0.7	*
15	Enterocolitis due to <i>Clostridium difficile</i> (A04.7)	5	0.5	*
...	All other causes (residual)	212	21.6	76.5
American Indian, female, 55-64 years				
...	All causes	1,321	100.0	606.2
1	Malignant neoplasms (C00-C97)	304	23.0	139.5
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	197	14.9	90.4
3	Chronic liver disease and cirrhosis (K70,K73-K74)	120	9.1	55.1
4	Diabetes mellitus (E10-E14)	98	7.4	45.0
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	79	6.0	36.3
6	Chronic lower respiratory diseases (J40-J47)	71	5.4	32.6
7	Cerebrovascular diseases (I60-I69)	47	3.6	21.6
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	34	2.6	15.6
9	Septicemia (A40-A41)	31	2.3	14.2
10	Influenza and pneumonia (J09-J18)	28	2.1	12.8
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	18	1.4	*
12	Viral hepatitis (B15-B19)	15	1.1	*
13	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	7	0.5	*
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	6	0.5	*
14	Pneumonitis due to solids and liquids (J69)	6	0.5	*
...	All other causes (residual)	260	19.7	119.3

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2014.

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex:
United States, 2014

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, female, 65-74 years				
...	All causes	1,567	100.0	1,368.6
1	Malignant neoplasms (C00-C97)	412	26.3	359.8
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	308	19.7	269.0
3	Chronic lower respiratory diseases (J40-J47)	124	7.9	108.3
4	Diabetes mellitus (E10-E14)	113	7.2	98.7
5	Chronic liver disease and cirrhosis (K70,K73-K74)	61	3.9	53.3
6	Cerebrovascular diseases (I60-I69)	60	3.8	52.4
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	44	2.8	38.4
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	39	2.5	34.1
9	Influenza and pneumonia (J09-J18)	37	2.4	32.3
10	Septicemia (A40-A41)	36	2.3	31.4
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	18	1.1	*
12	Alzheimer's disease (G30)	17	1.1	*
13	Enterocolitis due to <i>Clostridium difficile</i> (A04.7)	8	0.5	*
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	7	0.4	*
15	Parkinson's disease (G20-G21)	6	0.4	*
15	Pneumonitis due to solids and liquids (J69)	6	0.4	*
...	All other causes (residual)	271	17.3	236.7
American Indian, female, 75-84 years				
...	All causes	1,620	100.0	3,226.6
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	330	20.4	657.3
2	Malignant neoplasms (C00-C97)	307	19.0	611.5
3	Chronic lower respiratory diseases (J40-J47)	117	7.2	233.0
4	Diabetes mellitus (E10-E14)	95	5.9	189.2
5	Cerebrovascular diseases (I60-I69)	92	5.7	183.2
6	Alzheimer's disease (G30)	72	4.4	143.4
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	48	3.0	95.6
8	Influenza and pneumonia (J09-J18)	47	2.9	93.6
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	40	2.5	79.7
10	Septicemia (A40-A41)	38	2.3	75.7
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	22	1.4	43.8
12	Parkinson's disease (G20-G21)	20	1.2	39.8
13	Chronic liver disease and cirrhosis (K70,K73-K74)	18	1.1	*
14	Pneumonitis due to solids and liquids (J69)	15	0.9	*
15	Aortic aneurysm and dissection (I71)	13	0.8	*
...	All other causes (residual)	346	21.4	689.1

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2014.

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex:
United States, 2014

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, female, 85 years and over				
...	All causes	1,467	100.0	7,893.5
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	359	24.5	1,931.7
2	Malignant neoplasms (C00-C97)	152	10.4	817.9
3	Alzheimer's disease (G30)	130	8.9	699.5
4	Cerebrovascular diseases (I60-I69)	115	7.8	618.8
5	Chronic lower respiratory diseases (J40-J47)	79	5.4	425.1
6	Diabetes mellitus (E10-E14)	55	3.7	295.9
7	Influenza and pneumonia (J09-J18)	51	3.5	274.4
8	Essential hypertension and hypertensive renal disease (I10,I12,I15)	35	2.4	188.3
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	31	2.1	166.8
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	30	2.0	161.4
11	Septicemia (A40-A41)	18	1.2	*
12	Pneumonitis due to solids and liquids (J69)	12	0.8	*
13	Nutritional deficiencies (E40-E64)	11	0.7	*
14	Parkinson's disease (G20-G21)	10	0.7	*
15	Hernia (K40-K46)	7	0.5	*
...	All other causes (residual)	372	25.4	2,001.6
Asian or Pacific Islander, both sexes, all ages				
...	All causes	61,570	100.0	317.4
1	Malignant neoplasms (C00-C97)	16,524	26.8	85.2
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	13,270	21.6	68.4
3	Cerebrovascular diseases (I60-I69)	4,331	7.0	22.3
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,646	4.3	13.6
5	Diabetes mellitus (E10-E14)	2,367	3.8	12.2
6	Influenza and pneumonia (J09-J18)	1,911	3.1	9.9
7	Chronic lower respiratory diseases (J40-J47)	1,838	3.0	9.5
8	Alzheimer's disease (G30)	1,680	2.7	8.7
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,246	2.0	6.4
10	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,188	1.9	6.1
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,007	1.6	5.2
12	Septicemia (A40-A41)	753	1.2	3.9
13	Parkinson's disease (G20-G21)	611	1.0	3.1
14	Chronic liver disease and cirrhosis (K70,K73-K74)	608	1.0	3.1
15	Certain conditions originating in the perinatal period (P00-P96)	508	0.8	2.6
...	All other causes (residual)	11,082	18.0	57.1

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2014.

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex:
United States, 2014

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, both sexes, 1-4 years				
...	All causes	134	100.0	13.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	30	22.4	3.0
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	18	13.4	*
3	Malignant neoplasms (C00-C97)	12	9.0	*
4	Influenza and pneumonia (J09-J18)	7	5.2	*
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	6	4.5	*
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	5	3.7	*
7	Cerebrovascular diseases (I60-I69)	3	2.2	*
7	Chronic lower respiratory diseases (J40-J47)	3	2.2	*
9	Septicemia (A40-A41)	2	1.5	*
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2	1.5	*
11	Anemias (D50-D64)	1	0.7	*
11	Hernia (K40-K46)	1	0.7	*
...	All other causes (residual)	44	32.8	4.4
Asian or Pacific Islander, both sexes, 5-14 years				
...	All causes	204	100.0	8.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	40	19.6	1.6
2	Malignant neoplasms (C00-C97)	33	16.2	1.3
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	18	8.8	*
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	13	6.4	*
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	10	4.9	*
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	7	3.4	*
7	Septicemia (A40-A41)	6	2.9	*
8	Influenza and pneumonia (J09-J18)	4	2.0	*
8	Chronic lower respiratory diseases (J40-J47)	4	2.0	*
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	3	1.5	*
10	Anemias (D50-D64)	3	1.5	*
10	Cerebrovascular diseases (I60-I69)	3	1.5	*
13	Nutritional deficiencies (E40-E64)	1	0.5	*
13	Atherosclerosis (I70)	1	0.5	*
13	Peptic ulcer (K25-K28)	1	0.5	*
13	Diseases of appendix (K35-K38)	1	0.5	*
13	Certain conditions originating in the perinatal period (P00-P96)	1	0.5	*
13	Enterocolitis due to <i>Clostridium difficile</i> (A04.7)	1	0.5	*
...	All other causes (residual)	54	26.5	2.2

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2014.

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex:
United States, 2014

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, both sexes, 15-24 years				
...	All causes	843	100.0	31.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	272	32.3	10.1
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	229	27.2	8.5
3	Malignant neoplasms (C00-C97)	74	8.8	2.8
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	61	7.2	2.3
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	35	4.2	1.3
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	14	1.7	*
7	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	10	1.2	*
8	Cerebrovascular diseases (I60-I69)	7	0.8	*
9	Chronic lower respiratory diseases (J40-J47)	6	0.7	*
10	Septicemia (A40-A41)	3	0.4	*
10	Influenza and pneumonia (J09-J18)	3	0.4	*
10	Pneumonitis due to solids and liquids (J69)	3	0.4	*
10	Pregnancy, childbirth and the puerperium (O00-O99)	3	0.4	*
10	Legal intervention (Y35,Y89.0)	3	0.4	*
15	Diabetes mellitus (E10-E14)	2	0.2	*
15	Meningitis (G00,G03)	2	0.2	*
15	Chronic liver disease and cirrhosis (K70,K73-K74)	2	0.2	*
...	All other causes (residual)	114	13.5	4.3
Asian or Pacific Islander, both sexes, 25-34 years				
...	All causes	1,277	100.0	38.9
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	350	27.4	10.7
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	235	18.4	7.2
3	Malignant neoplasms (C00-C97)	200	15.7	6.1
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	104	8.1	3.2
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	60	4.7	1.8
6	Cerebrovascular diseases (I60-I69)	27	2.1	0.8
7	Diabetes mellitus (E10-E14)	19	1.5	*
8	Chronic liver disease and cirrhosis (K70,K73-K74)	17	1.3	*
8	Pregnancy, childbirth and the puerperium (O00-O99)	17	1.3	*
10	Influenza and pneumonia (J09-J18)	15	1.2	*
11	Septicemia (A40-A41)	13	1.0	*
11	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	13	1.0	*
13	Human immunodeficiency virus (HIV) disease (B20-B24)	10	0.8	*
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	10	0.8	*
15	Chronic lower respiratory diseases (J40-J47)	8	0.6	*
...	All other causes (residual)	179	14.0	5.5

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2014.

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex:
United States, 2014

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, both sexes, 35-44 years				
...	All causes	2,005	100.0	65.0
1	Malignant neoplasms (C00-C97)	584	29.1	18.9
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	313	15.6	10.2
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	251	12.5	8.1
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	192	9.6	6.2
5	Cerebrovascular diseases (I60-I69)	90	4.5	2.9
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	56	2.8	1.8
7	Chronic liver disease and cirrhosis (K70,K73-K74)	51	2.5	1.7
8	Diabetes mellitus (E10-E14)	44	2.2	1.4
9	Influenza and pneumonia (J09-J18)	29	1.4	0.9
10	Chronic lower respiratory diseases (J40-J47)	24	1.2	0.8
11	Pregnancy, childbirth and the puerperium (O00-O99)	22	1.1	0.7
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	19	0.9	*
13	Human immunodeficiency virus (HIV) disease (B20-B24)	18	0.9	*
14	Aortic aneurysm and dissection (I71)	16	0.8	*
15	Viral hepatitis (B15-B19)	14	0.7	*
...	All other causes (residual)	282	14.1	9.1
Asian or Pacific Islander, both sexes, 45-54 years				
...	All causes	4,229	100.0	164.5
1	Malignant neoplasms (C00-C97)	1,654	39.1	64.3
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	806	19.1	31.3
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	259	6.1	10.1
4	Cerebrovascular diseases (I60-I69)	249	5.9	9.7
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	214	5.1	8.3
6	Diabetes mellitus (E10-E14)	132	3.1	5.1
7	Chronic liver disease and cirrhosis (K70,K73-K74)	90	2.1	3.5
8	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	53	1.3	2.1
9	Influenza and pneumonia (J09-J18)	51	1.2	2.0
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	48	1.1	1.9
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	47	1.1	1.8
12	Viral hepatitis (B15-B19)	40	0.9	1.6
13	Septicemia (A40-A41)	38	0.9	1.5
14	Chronic lower respiratory diseases (J40-J47)	29	0.7	1.1
15	Aortic aneurysm and dissection (I71)	27	0.6	1.1
...	All other causes (residual)	492	11.6	19.1

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2014.

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex:
United States, 2014

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, both sexes, 55-64 years				
...	All causes	7,954	100.0	390.7
1	Malignant neoplasms (C00-C97)	3,362	42.3	165.1
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,511	19.0	74.2
3	Cerebrovascular diseases (I60-I69)	481	6.0	23.6
4	Diabetes mellitus (E10-E14)	356	4.5	17.5
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	305	3.8	15.0
6	Chronic liver disease and cirrhosis (K70,K73-K74)	149	1.9	7.3
7	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	148	1.9	7.3
8	Influenza and pneumonia (J09-J18)	144	1.8	7.1
9	Chronic lower respiratory diseases (J40-J47)	126	1.6	6.2
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	112	1.4	5.5
11	Viral hepatitis (B15-B19)	99	1.2	4.9
12	Septicemia (A40-A41)	93	1.2	4.6
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	87	1.1	4.3
14	Aortic aneurysm and dissection (I71)	58	0.7	2.8
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	41	0.5	2.0
...	All other causes (residual)	882	11.1	43.3
Asian or Pacific Islander, both sexes, 65-74 years				
...	All causes	10,728	100.0	883.0
1	Malignant neoplasms (C00-C97)	4,046	37.7	333.0
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,202	20.5	181.2
3	Cerebrovascular diseases (I60-I69)	712	6.6	58.6
4	Diabetes mellitus (E10-E14)	524	4.9	43.1
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	296	2.8	24.4
6	Chronic lower respiratory diseases (J40-J47)	269	2.5	22.1
7	Influenza and pneumonia (J09-J18)	253	2.4	20.8
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	236	2.2	19.4
9	Septicemia (A40-A41)	164	1.5	13.5
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	146	1.4	12.0
11	Chronic liver disease and cirrhosis (K70,K73-K74)	128	1.2	10.5
12	Parkinson's disease (G20-G21)	92	0.9	7.6
13	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	90	0.8	7.4
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	69	0.6	5.7
15	Viral hepatitis (B15-B19)	66	0.6	5.4
...	All other causes (residual)	1,435	13.4	118.1

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2014.

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex:
United States, 2014

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, both sexes, 75-84 years				
...	All causes	14,979	100.0	2,584.4
1	Malignant neoplasms (C00-C97)	4,063	27.1	701.0
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,412	22.8	588.7
3	Cerebrovascular diseases (I60-I69)	1,195	8.0	206.2
4	Diabetes mellitus (E10-E14)	690	4.6	119.0
5	Chronic lower respiratory diseases (J40-J47)	616	4.1	106.3
6	Influenza and pneumonia (J09-J18)	521	3.5	89.9
7	Alzheimer's disease (G30)	402	2.7	69.4
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	374	2.5	64.5
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	369	2.5	63.7
10	Parkinson's disease (G20-G21)	244	1.6	42.1
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	242	1.6	41.8
12	Septicemia (A40-A41)	203	1.4	35.0
13	Chronic liver disease and cirrhosis (K70,K73-K74)	130	0.9	22.4
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	127	0.8	21.9
15	Pneumonitis due to solids and liquids (J69)	121	0.8	20.9
...	All other causes (residual)	2,270	15.2	391.7
Asian or Pacific Islander, both sexes, 85 years and over				
...	All causes	18,319	100.0	8,427.6
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	4,847	26.5	2,229.9
2	Malignant neoplasms (C00-C97)	2,496	13.6	1,148.3
3	Cerebrovascular diseases (I60-I69)	1,560	8.5	717.7
4	Alzheimer's disease (G30)	1,212	6.6	557.6
5	Influenza and pneumonia (J09-J18)	877	4.8	403.5
6	Chronic lower respiratory diseases (J40-J47)	753	4.1	346.4
7	Diabetes mellitus (E10-E14)	600	3.3	276.0
8	Essential hypertension and hypertensive renal disease (I10,I12,I15)	471	2.6	216.7
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	457	2.5	210.2
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	445	2.4	204.7
11	Parkinson's disease (G20-G21)	260	1.4	119.6
12	Pneumonitis due to solids and liquids (J69)	253	1.4	116.4
13	Septicemia (A40-A41)	210	1.1	96.6
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	121	0.7	55.7
15	Aortic aneurysm and dissection (I71)	82	0.4	37.7
...	All other causes (residual)	3,675	20.1	1,690.7

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2014.

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex:
United States, 2014

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, male, all ages				
...	All causes	31,686	100.0	341.3
1	Malignant neoplasms (C00-C97)	8,371	26.4	90.2
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	7,279	23.0	78.4
3	Cerebrovascular diseases (I60-I69)	1,923	6.1	20.7
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,665	5.3	17.9
5	Diabetes mellitus (E10-E14)	1,235	3.9	13.3
6	Chronic lower respiratory diseases (J40-J47)	1,072	3.4	11.5
7	Influenza and pneumonia (J09-J18)	1,012	3.2	10.9
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	833	2.6	9.0
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	652	2.1	7.0
10	Alzheimer's disease (G30)	508	1.6	5.5
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	435	1.4	4.7
12	Septicemia (A40-A41)	394	1.2	4.2
13	Chronic liver disease and cirrhosis (K70,K73-K74)	382	1.2	4.1
14	Parkinson's disease (G20-G21)	362	1.1	3.9
15	Certain conditions originating in the perinatal period (P00-P96)	281	0.9	3.0
...	All other causes (residual)	5,282	16.7	56.9
Asian or Pacific Islander, male, 1-4 years				
...	All causes	73	100.0	14.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	20	27.4	3.9
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	11	15.1	*
3	Malignant neoplasms (C00-C97)	7	9.6	*
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	5	6.8	*
5	Chronic lower respiratory diseases (J40-J47)	3	4.1	*
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	2	2.7	*
7	Septicemia (A40-A41)	1	1.4	*
7	Anemias (D50-D64)	1	1.4	*
7	Cerebrovascular diseases (I60-I69)	1	1.4	*
7	Influenza and pneumonia (J09-J18)	1	1.4	*
...	All other causes (residual)	21	28.8	4.1

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2014.

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex:
United States, 2014

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, male, 5-14 years				
...	All causes	124	100.0	9.9
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	25	20.2	2.0
2	Malignant neoplasms (C00-C97)	20	16.1	1.6
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	9	7.3	*
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	8	6.5	*
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	6	4.8	*
6	Septicemia (A40-A41)	5	4.0	*
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	5	4.0	*
8	Cerebrovascular diseases (I60-I69)	3	2.4	*
8	Influenza and pneumonia (J09-J18)	3	2.4	*
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2	1.6	*
10	Chronic lower respiratory diseases (J40-J47)	2	1.6	*
12	Anemias (D50-D64)	1	0.8	*
12	Atherosclerosis (I70)	1	0.8	*
12	Peptic ulcer (K25-K28)	1	0.8	*
...	All other causes (residual)	33	26.6	2.6
Asian or Pacific Islander, male, 15-24 years				
...	All causes	608	100.0	44.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	208	34.2	15.3
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	175	28.8	12.9
3	Malignant neoplasms (C00-C97)	47	7.7	3.5
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	46	7.6	3.4
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	26	4.3	1.9
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	8	1.3	*
7	Chronic lower respiratory diseases (J40-J47)	5	0.8	*
8	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	4	0.7	*
8	Cerebrovascular diseases (I60-I69)	4	0.7	*
10	Pneumonitis due to solids and liquids (J69)	3	0.5	*
10	Legal intervention (Y35,Y89.0)	3	0.5	*
12	Septicemia (A40-A41)	2	0.3	*
12	Diabetes mellitus (E10-E14)	2	0.3	*
12	Chronic liver disease and cirrhosis (K70,K73-K74)	2	0.3	*
15	Parkinson's disease (G20-G21)	1	0.2	*
15	Influenza and pneumonia (J09-J18)	1	0.2	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1	0.2	*
15	Certain conditions originating in the perinatal period (P00-P96)	1	0.2	*
...	All other causes (residual)	69	11.3	5.1

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2014.

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex:
United States, 2014

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, male, 25-34 years				
...	All causes	855	100.0	54.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	266	31.1	16.8
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	171	20.0	10.8
3	Malignant neoplasms (C00-C97)	94	11.0	6.0
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	84	9.8	5.3
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	52	6.1	3.3
6	Cerebrovascular diseases (I60-I69)	16	1.9	*
7	Diabetes mellitus (E10-E14)	15	1.8	*
8	Chronic liver disease and cirrhosis (K70,K73-K74)	11	1.3	*
9	Human immunodeficiency virus (HIV) disease (B20-B24)	10	1.2	*
10	Septicemia (A40-A41)	7	0.8	*
10	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	7	0.8	*
12	Viral hepatitis (B15-B19)	6	0.7	*
12	Influenza and pneumonia (J09-J18)	6	0.7	*
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	6	0.7	*
15	Aortic aneurysm and dissection (I71)	5	0.6	*
...	All other causes (residual)	99	11.6	6.3
Asian or Pacific Islander, male, 35-44 years				
...	All causes	1,202	100.0	83.0
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	249	20.7	17.2
2	Malignant neoplasms (C00-C97)	230	19.1	15.9
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	183	15.2	12.6
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	123	10.2	8.5
5	Cerebrovascular diseases (I60-I69)	57	4.7	3.9
6	Chronic liver disease and cirrhosis (K70,K73-K74)	40	3.3	2.8
7	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	38	3.2	2.6
8	Diabetes mellitus (E10-E14)	26	2.2	1.8
9	Influenza and pneumonia (J09-J18)	18	1.5	*
10	Aortic aneurysm and dissection (I71)	15	1.2	*
10	Chronic lower respiratory diseases (J40-J47)	15	1.2	*
12	Human immunodeficiency virus (HIV) disease (B20-B24)	14	1.2	*
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	10	0.8	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	9	0.7	*
15	Viral hepatitis (B15-B19)	8	0.7	*
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	8	0.7	*
...	All other causes (residual)	159	13.2	11.0

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2014.

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex:
United States, 2014

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, male, 45-54 years				
...	All causes	2,558	100.0	212.3
1	Malignant neoplasms (C00-C97)	778	30.4	64.6
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	644	25.2	53.4
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	179	7.0	14.9
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	146	5.7	12.1
5	Cerebrovascular diseases (I60-I69)	132	5.2	11.0
6	Diabetes mellitus (E10-E14)	87	3.4	7.2
7	Chronic liver disease and cirrhosis (K70,K73-K74)	71	2.8	5.9
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	34	1.3	2.8
9	Essential hypertension and hypertensive renal disease (I10,I12,I15)	33	1.3	2.7
10	Viral hepatitis (B15-B19)	29	1.1	2.4
10	Influenza and pneumonia (J09-J18)	29	1.1	2.4
10	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	29	1.1	2.4
13	Septicemia (A40-A41)	22	0.9	1.8
14	Aortic aneurysm and dissection (I71)	21	0.8	1.7
14	Chronic lower respiratory diseases (J40-J47)	21	0.8	1.7
...	All other causes (residual)	303	11.8	25.1
Asian or Pacific Islander, male, 55-64 years				
...	All causes	4,808	100.0	519.9
1	Malignant neoplasms (C00-C97)	1,764	36.7	190.7
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,123	23.4	121.4
3	Cerebrovascular diseases (I60-I69)	277	5.8	30.0
4	Diabetes mellitus (E10-E14)	228	4.7	24.7
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	203	4.2	22.0
6	Chronic liver disease and cirrhosis (K70,K73-K74)	114	2.4	12.3
7	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	111	2.3	12.0
8	Influenza and pneumonia (J09-J18)	92	1.9	9.9
9	Chronic lower respiratory diseases (J40-J47)	83	1.7	9.0
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	69	1.4	7.5
11	Viral hepatitis (B15-B19)	68	1.4	7.4
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	54	1.1	5.8
13	Septicemia (A40-A41)	51	1.1	5.5
14	Aortic aneurysm and dissection (I71)	41	0.9	4.4
15	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	25	0.5	2.7
...	All other causes (residual)	505	10.5	54.6

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2014.

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex:
United States, 2014

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, male, 65-74 years				
...	All causes	6,037	100.0	1,107.1
1	Malignant neoplasms (C00-C97)	2,179	36.1	399.6
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,366	22.6	250.5
3	Cerebrovascular diseases (I60-I69)	362	6.0	66.4
4	Diabetes mellitus (E10-E14)	297	4.9	54.5
5	Chronic lower respiratory diseases (J40-J47)	173	2.9	31.7
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	173	2.9	31.7
7	Influenza and pneumonia (J09-J18)	152	2.5	27.9
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	127	2.1	23.3
9	Septicemia (A40-A41)	101	1.7	18.5
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	73	1.2	13.4
10	Chronic liver disease and cirrhosis (K70,K73-K74)	73	1.2	13.4
12	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	60	1.0	11.0
13	Parkinson's disease (G20-G21)	56	0.9	10.3
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	42	0.7	7.7
15	Pneumonitis due to solids and liquids (J69)	34	0.6	6.2
...	All other causes (residual)	769	12.7	141.0
Asian or Pacific Islander, male, 75-84 years				
...	All causes	7,563	100.0	3,047.8
1	Malignant neoplasms (C00-C97)	2,084	27.6	839.8
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,803	23.8	726.6
3	Cerebrovascular diseases (I60-I69)	539	7.1	217.2
4	Chronic lower respiratory diseases (J40-J47)	352	4.7	141.9
5	Diabetes mellitus (E10-E14)	334	4.4	134.6
6	Influenza and pneumonia (J09-J18)	279	3.7	112.4
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	207	2.7	83.4
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	203	2.7	81.8
9	Parkinson's disease (G20-G21)	149	2.0	60.0
10	Alzheimer's disease (G30)	141	1.9	56.8
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	108	1.4	43.5
12	Septicemia (A40-A41)	107	1.4	43.1
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	72	1.0	29.0
14	Pneumonitis due to solids and liquids (J69)	70	0.9	28.2
15	Aortic aneurysm and dissection (I71)	55	0.7	22.2
15	Chronic liver disease and cirrhosis (K70,K73-K74)	55	0.7	22.2
...	All other causes (residual)	1,005	13.3	405.0

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2014.

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex:
United States, 2014

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, male, 85 years and over				
...	All causes	7,370	100.0	9,263.1
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,957	26.6	2,459.7
2	Malignant neoplasms (C00-C97)	1,168	15.8	1,468.0
3	Cerebrovascular diseases (I60-I69)	530	7.2	666.1
4	Influenza and pneumonia (J09-J18)	427	5.8	536.7
5	Chronic lower respiratory diseases (J40-J47)	415	5.6	521.6
6	Alzheimer's disease (G30)	340	4.6	427.3
7	Diabetes mellitus (E10-E14)	246	3.3	309.2
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	199	2.7	250.1
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	198	2.7	248.9
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	156	2.1	196.1
11	Parkinson's disease (G20-G21)	148	2.0	186.0
12	Pneumonitis due to solids and liquids (J69)	137	1.9	172.2
13	Septicemia (A40-A41)	89	1.2	111.9
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	53	0.7	66.6
15	Tuberculosis (A16-A19)	26	0.4	32.7
...	All other causes (residual)	1,281	17.4	1,610.0
Asian or Pacific Islander, female, all ages				
...	All causes	29,884	100.0	295.5
1	Malignant neoplasms (C00-C97)	8,153	27.3	80.6
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	5,991	20.0	59.2
3	Cerebrovascular diseases (I60-I69)	2,408	8.1	23.8
4	Alzheimer's disease (G30)	1,172	3.9	11.6
5	Diabetes mellitus (E10-E14)	1,132	3.8	11.2
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	981	3.3	9.7
7	Influenza and pneumonia (J09-J18)	899	3.0	8.9
8	Chronic lower respiratory diseases (J40-J47)	766	2.6	7.6
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	594	2.0	5.9
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	572	1.9	5.7
11	Septicemia (A40-A41)	359	1.2	3.5
12	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	355	1.2	3.5
13	Parkinson's disease (G20-G21)	249	0.8	2.5
14	Certain conditions originating in the perinatal period (P00-P96)	227	0.8	2.2
15	Chronic liver disease and cirrhosis (K70,K73-K74)	226	0.8	2.2
...	All other causes (residual)	5,800	19.4	57.3

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2014.

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex:
United States, 2014

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, female, 1-4 years				
...	All causes	61	100.0	12.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	10	16.4	*
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	7	11.5	*
3	Influenza and pneumonia (J09-J18)	6	9.8	*
4	Malignant neoplasms (C00-C97)	5	8.2	*
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	3	4.9	*
6	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2	3.3	*
6	Cerebrovascular diseases (I60-I69)	2	3.3	*
8	Septicemia (A40-A41)	1	1.6	*
8	Diseases of heart (I00-I09,I11,I13,I20-I51)	1	1.6	*
8	Hernia (K40-K46)	1	1.6	*
...	All other causes (residual)	23	37.7	4.7
Asian or Pacific Islander, female, 5-14 years				
...	All causes	80	100.0	6.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	15	18.8	*
2	Malignant neoplasms (C00-C97)	13	16.3	*
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	10	12.5	*
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	4	5.0	*
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	4	5.0	*
6	Anemias (D50-D64)	2	2.5	*
6	Chronic lower respiratory diseases (J40-J47)	2	2.5	*
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	2	2.5	*
9	Septicemia (A40-A41)	1	1.3	*
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1	1.3	*
9	Nutritional deficiencies (E40-E64)	1	1.3	*
9	Influenza and pneumonia (J09-J18)	1	1.3	*
9	Diseases of appendix (K35-K38)	1	1.3	*
9	Certain conditions originating in the perinatal period (P00-P96)	1	1.3	*
9	Enterocolitis due to <i>Clostridium difficile</i> (A04.7)	1	1.3	*
...	All other causes (residual)	21	26.3	1.7

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2014.

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex:
United States, 2014

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, female, 15-24 years				
...	All causes	235	100.0	17.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	64	27.2	4.8
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	54	23.0	4.1
3	Malignant neoplasms (C00-C97)	27	11.5	2.0
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	15	6.4	*
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	9	3.8	*
6	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	6	2.6	*
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	6	2.6	*
8	Cerebrovascular diseases (I60-I69)	3	1.3	*
8	Pregnancy, childbirth and the puerperium (O00-O99)	3	1.3	*
10	Meningitis (G00,G03)	2	0.9	*
10	Influenza and pneumonia (J09-J18)	2	0.9	*
12	Septicemia (A40-A41)	1	0.4	*
12	Human immunodeficiency virus (HIV) disease (B20-B24)	1	0.4	*
12	Anemias (D50-D64)	1	0.4	*
12	Chronic lower respiratory diseases (J40-J47)	1	0.4	*
...	All other causes (residual)	40	17.0	3.0
Asian or Pacific Islander, female, 25-34 years				
...	All causes	422	100.0	24.8
1	Malignant neoplasms (C00-C97)	106	25.1	6.2
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	84	19.9	4.9
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	64	15.2	3.8
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	20	4.7	1.2
5	Pregnancy, childbirth and the puerperium (O00-O99)	17	4.0	*
6	Cerebrovascular diseases (I60-I69)	11	2.6	*
7	Influenza and pneumonia (J09-J18)	9	2.1	*
8	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	8	1.9	*
9	Septicemia (A40-A41)	6	1.4	*
9	Chronic liver disease and cirrhosis (K70,K73-K74)	6	1.4	*
9	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	6	1.4	*
12	Chronic lower respiratory diseases (J40-J47)	5	1.2	*
13	Diabetes mellitus (E10-E14)	4	0.9	*
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	4	0.9	*
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2	0.5	*
...	All other causes (residual)	70	16.6	4.1

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2014.

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex:
United States, 2014

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, female, 35-44 years				
...	All causes	803	100.0	49.1
1	Malignant neoplasms (C00-C97)	354	44.1	21.7
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	69	8.6	4.2
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	68	8.5	4.2
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	64	8.0	3.9
5	Cerebrovascular diseases (I60-I69)	33	4.1	2.0
6	Pregnancy, childbirth and the puerperium (O00-O99)	22	2.7	1.3
7	Diabetes mellitus (E10-E14)	18	2.2	*
7	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	18	2.2	*
9	Influenza and pneumonia (J09-J18)	11	1.4	*
9	Chronic liver disease and cirrhosis (K70,K73-K74)	11	1.4	*
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	10	1.2	*
12	Chronic lower respiratory diseases (J40-J47)	9	1.1	*
13	Septicemia (A40-A41)	7	0.9	*
14	Viral hepatitis (B15-B19)	6	0.7	*
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	5	0.6	*
...	All other causes (residual)	98	12.2	6.0
Asian or Pacific Islander, female, 45-54 years				
...	All causes	1,671	100.0	122.3
1	Malignant neoplasms (C00-C97)	876	52.4	64.1
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	162	9.7	11.9
3	Cerebrovascular diseases (I60-I69)	117	7.0	8.6
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	80	4.8	5.9
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	68	4.1	5.0
6	Diabetes mellitus (E10-E14)	45	2.7	3.3
7	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	24	1.4	1.8
8	Influenza and pneumonia (J09-J18)	22	1.3	1.6
9	Chronic liver disease and cirrhosis (K70,K73-K74)	19	1.1	*
10	Septicemia (A40-A41)	16	1.0	*
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	15	0.9	*
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	13	0.8	*
13	Viral hepatitis (B15-B19)	11	0.7	*
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	8	0.5	*
14	Chronic lower respiratory diseases (J40-J47)	8	0.5	*
...	All other causes (residual)	187	11.2	13.7

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2014.

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex:
United States, 2014

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, female, 55-64 years				
...	All causes	3,146	100.0	283.2
1	Malignant neoplasms (C00-C97)	1,598	50.8	143.8
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	388	12.3	34.9
3	Cerebrovascular diseases (I60-I69)	204	6.5	18.4
4	Diabetes mellitus (E10-E14)	128	4.1	11.5
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	102	3.2	9.2
6	Influenza and pneumonia (J09-J18)	52	1.7	4.7
7	Chronic lower respiratory diseases (J40-J47)	43	1.4	3.9
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	43	1.4	3.9
9	Septicemia (A40-A41)	42	1.3	3.8
10	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	37	1.2	3.3
11	Chronic liver disease and cirrhosis (K70,K73-K74)	35	1.1	3.2
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	33	1.0	3.0
13	Viral hepatitis (B15-B19)	31	1.0	2.8
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	21	0.7	1.9
15	Aortic aneurysm and dissection (I71)	17	0.5	*
...	All other causes (residual)	372	11.8	33.5
Asian or Pacific Islander, female, 65-74 years				
...	All causes	4,691	100.0	700.5
1	Malignant neoplasms (C00-C97)	1,867	39.8	278.8
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	836	17.8	124.8
3	Cerebrovascular diseases (I60-I69)	350	7.5	52.3
4	Diabetes mellitus (E10-E14)	227	4.8	33.9
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	123	2.6	18.4
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	109	2.3	16.3
7	Influenza and pneumonia (J09-J18)	101	2.2	15.1
8	Chronic lower respiratory diseases (J40-J47)	96	2.0	14.3
9	Essential hypertension and hypertensive renal disease (I10,I12,I15)	73	1.6	10.9
10	Septicemia (A40-A41)	63	1.3	9.4
11	Chronic liver disease and cirrhosis (K70,K73-K74)	55	1.2	8.2
12	Viral hepatitis (B15-B19)	36	0.8	5.4
12	Parkinson's disease (G20-G21)	36	0.8	5.4
14	Alzheimer's disease (G30)	31	0.7	4.6
15	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	30	0.6	4.5
...	All other causes (residual)	658	14.0	98.3

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2014.

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex:
United States, 2014

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, female, 75-84 years				
...	All causes	7,416	100.0	2,237.5
1	Malignant neoplasms (C00-C97)	1,979	26.7	597.1
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,609	21.7	485.4
3	Cerebrovascular diseases (I60-I69)	656	8.8	197.9
4	Diabetes mellitus (E10-E14)	356	4.8	107.4
5	Chronic lower respiratory diseases (J40-J47)	264	3.6	79.7
6	Alzheimer's disease (G30)	261	3.5	78.7
7	Influenza and pneumonia (J09-J18)	242	3.3	73.0
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	167	2.3	50.4
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	166	2.2	50.1
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	134	1.8	40.4
11	Septicemia (A40-A41)	96	1.3	29.0
12	Parkinson's disease (G20-G21)	95	1.3	28.7
13	Chronic liver disease and cirrhosis (K70,K73-K74)	75	1.0	22.6
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	55	0.7	16.6
15	Pneumonitis due to solids and liquids (J69)	51	0.7	15.4
...	All other causes (residual)	1,210	16.3	365.1
Asian or Pacific Islander, female, 85 years and over				
...	All causes	10,949	100.0	7,945.3
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,890	26.4	2,097.2
2	Malignant neoplasms (C00-C97)	1,328	12.1	963.7
3	Cerebrovascular diseases (I60-I69)	1,030	9.4	747.4
4	Alzheimer's disease (G30)	872	8.0	632.8
5	Influenza and pneumonia (J09-J18)	450	4.1	326.5
6	Diabetes mellitus (E10-E14)	354	3.2	256.9
7	Chronic lower respiratory diseases (J40-J47)	338	3.1	245.3
8	Essential hypertension and hypertensive renal disease (I10,I12,I15)	315	2.9	228.6
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	258	2.4	187.2
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	247	2.3	179.2
11	Septicemia (A40-A41)	121	1.1	87.8
12	Pneumonitis due to solids and liquids (J69)	116	1.1	84.2
13	Parkinson's disease (G20-G21)	112	1.0	81.3
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	68	0.6	49.3
15	Aortic aneurysm and dissection (I71)	61	0.6	44.3
...	All other causes (residual)	2,389	21.8	1,733.6

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2014.