

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), race, sex, and age	Number	Percent of total deaths	Rate
All races, both sexes, all ages				
...	All causes	2,468,435	100.0	799.5
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	597,689	24.2	193.6
2	Malignant neoplasms (C00-C97)	574,743	23.3	186.2
3	Chronic lower respiratory diseases (J40-J47)	138,080	5.6	44.7
4	Cerebrovascular diseases (I60-I69)	129,476	5.2	41.9
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	120,859	4.9	39.1
6	Alzheimer's disease (G30)	83,494	3.4	27.0
7	Diabetes mellitus (E10-E14)	69,071	2.8	22.4
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	50,476	2.0	16.3
9	Influenza and pneumonia (J09-J18)	50,097	2.0	16.2
10	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	38,364	1.6	12.4
11	Septicemia (A40-A41)	34,812	1.4	11.3
12	Chronic liver disease and cirrhosis (K70,K73-K74)	31,903	1.3	10.3
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	26,634	1.1	8.6
14	Parkinson's disease (G20-G21)	22,032	0.9	7.1
15	Pneumonitis due to solids and liquids (J69)	17,011	0.7	5.5
...	All other causes (Residual)	483,694	19.6	156.7
All races, both sexes, 1-4 years				
...	All causes	4,316	100.0	26.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,394	32.3	8.6
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	507	11.7	3.1
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	385	8.9	2.4
4	Malignant neoplasms (C00-C97)	346	8.0	2.1
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	159	3.7	1.0
6	Influenza and pneumonia (J09-J18)	91	2.1	0.6
7	Septicemia (A40-A41)	62	1.4	0.4
8	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	59	1.4	0.4
9	Certain conditions originating in the perinatal period (P00-P96)	52	1.2	0.3
10	Chronic lower respiratory diseases (J40-J47)	51	1.2	0.3
11	Cerebrovascular diseases (I60-I69)	50	1.2	0.3
12	Anemias (D50-D64)	29	0.7	0.2
13	Meningitis (G00,G03)	22	0.5	0.1
14	Complications of medical and surgical care (Y40-Y84,Y88)	19	0.4	*
15	Pneumonitis due to solids and liquids (J69)	13	0.3	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	13	0.3	*
...	All other causes (Residual)	1,064	24.7	6.5

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), race, sex, and age	Number	Percent of total deaths	Rate
All races, both sexes, 5-14 years				
...	All causes	5,279	100.0	12.9
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,643	31.1	4.0
2	Malignant neoplasms (C00-C97)	916	17.4	2.2
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	298	5.6	0.7
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	274	5.2	0.7
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	261	4.9	0.6
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	185	3.5	0.5
7	Chronic lower respiratory diseases (J40-J47)	133	2.5	0.3
8	Cerebrovascular diseases (I60-I69)	90	1.7	0.2
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	82	1.6	0.2
10	Influenza and pneumonia (J09-J18)	71	1.3	0.2
11	Septicemia (A40-A41)	67	1.3	0.2
12	Anemias (D50-D64)	26	0.5	0.1
12	Diabetes mellitus (E10-E14)	26	0.5	0.1
14	Certain conditions originating in the perinatal period (P00-P96)	25	0.5	0.1
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	22	0.4	0.1
15	Complications of medical and surgical care (Y40-Y84,Y88)	22	0.4	0.1
...	All other causes (Residual)	1,138	21.6	2.8
All races, both sexes, 15-24 years				
...	All causes	29,551	100.0	67.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	12,341	41.8	28.3
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	4,678	15.8	10.7
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	4,600	15.6	10.5
4	Malignant neoplasms (C00-C97)	1,604	5.4	3.7
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,028	3.5	2.4
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	412	1.4	0.9
7	Cerebrovascular diseases (I60-I69)	190	0.6	0.4
8	Influenza and pneumonia (J09-J18)	181	0.6	0.4
9	Diabetes mellitus (E10-E14)	165	0.6	0.4
10	Pregnancy, childbirth and the puerperium (O00-O99)	163	0.6	0.4
11	Chronic lower respiratory diseases (J40-J47)	149	0.5	0.3
12	Human immunodeficiency virus (HIV) disease (B20-B24)	147	0.5	0.3
13	Septicemia (A40-A41)	141	0.5	0.3
14	Anemias (D50-D64)	109	0.4	0.2
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	93	0.3	0.2
...	All other causes (Residual)	3,550	12.0	8.1

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), race, sex, and age	Number	Percent of total deaths	Rate
All races, both sexes, 25-44 years				
...	All causes	112,292	100.0	136.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	29,365	26.2	35.8
2	Malignant neoplasms (C00-C97)	15,428	13.7	18.8
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	13,816	12.3	16.8
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	12,306	11.0	15.0
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	6,731	6.0	8.2
6	Chronic liver disease and cirrhosis (K70,K73-K74)	2,910	2.6	3.5
7	Human immunodeficiency virus (HIV) disease (B20-B24)	2,639	2.4	3.2
8	Cerebrovascular diseases (I60-I69)	2,421	2.2	2.9
9	Diabetes mellitus (E10-E14)	2,395	2.1	2.9
10	Influenza and pneumonia (J09-J18)	1,158	1.0	1.4
11	Septicemia (A40-A41)	1,079	1.0	1.3
12	Chronic lower respiratory diseases (J40-J47)	981	0.9	1.2
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	969	0.9	1.2
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	846	0.8	1.0
15	Pregnancy, childbirth and the puerperium (O00-O99)	590	0.5	0.7
...	All other causes (Residual)	18,658	16.6	22.7
All races, both sexes, 45-64 years				
...	All causes	494,009	100.0	606.2
1	Malignant neoplasms (C00-C97)	159,712	32.3	196.0
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	104,806	21.2	128.6
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	33,690	6.8	41.3
4	Chronic lower respiratory diseases (J40-J47)	18,694	3.8	22.9
5	Chronic liver disease and cirrhosis (K70,K73-K74)	18,415	3.7	22.6
6	Diabetes mellitus (E10-E14)	17,287	3.5	21.2
7	Cerebrovascular diseases (I60-I69)	16,603	3.4	20.4
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	15,183	3.1	18.6
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	7,304	1.5	9.0
10	Septicemia (A40-A41)	6,937	1.4	8.5
11	Viral hepatitis (B15-B19)	5,594	1.1	6.9
12	Influenza and pneumonia (J09-J18)	5,553	1.1	6.8
13	Human immunodeficiency virus (HIV) disease (B20-B24)	4,945	1.0	6.1
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	4,056	0.8	5.0
15	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	3,062	0.6	3.8
...	All other causes (Residual)	72,168	14.6	88.6

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), race, sex, and age	Number	Percent of total deaths	Rate
All races, both sexes, 65 years and over				
...	All causes	1,798,276	100.0	4,465.8
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	477,338	26.5	1,185.4
2	Malignant neoplasms (C00-C97)	396,670	22.1	985.1
3	Chronic lower respiratory diseases (J40-J47)	118,031	6.6	293.1
4	Cerebrovascular diseases (I60-I69)	109,990	6.1	273.1
5	Alzheimer's disease (G30)	82,616	4.6	205.2
6	Diabetes mellitus (E10-E14)	49,191	2.7	122.2
7	Influenza and pneumonia (J09-J18)	42,846	2.4	106.4
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	41,994	2.3	104.3
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	41,300	2.3	102.6
10	Septicemia (A40-A41)	26,310	1.5	65.3
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	22,047	1.2	54.8
12	Parkinson's disease (G20-G21)	21,448	1.2	53.3
13	Pneumonitis due to solids and liquids (J69)	15,226	0.8	37.8
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	12,004	0.7	29.8
15	Chronic liver disease and cirrhosis (K70,K73-K74)	10,535	0.6	26.2
...	All other causes (Residual)	330,730	18.4	821.3
All races, male, all ages				
...	All causes	1,232,432	100.0	812.0
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	307,384	24.9	202.5
2	Malignant neoplasms (C00-C97)	301,037	24.4	198.3
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	75,921	6.2	50.0
4	Chronic lower respiratory diseases (J40-J47)	65,423	5.3	43.1
5	Cerebrovascular diseases (I60-I69)	52,367	4.2	34.5
6	Diabetes mellitus (E10-E14)	35,490	2.9	23.4
7	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	30,277	2.5	19.9
8	Alzheimer's disease (G30)	25,364	2.1	16.7
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	24,865	2.0	16.4
10	Influenza and pneumonia (J09-J18)	23,615	1.9	15.6
11	Chronic liver disease and cirrhosis (K70,K73-K74)	20,798	1.7	13.7
12	Septicemia (A40-A41)	16,069	1.3	10.6
13	Parkinson's disease (G20-G21)	12,871	1.0	8.5
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	12,774	1.0	8.4
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	10,846	0.9	7.1
...	All other causes (Residual)	217,331	17.6	143.2

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), race, sex, and age	Number	Percent of total deaths	Rate
All races, male, 1-4 years				
...	All causes	2,460	100.0	29.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	868	35.3	10.5
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	231	9.4	2.8
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	231	9.4	2.8
4	Malignant neoplasms (C00-C97)	197	8.0	2.4
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	90	3.7	1.1
6	Influenza and pneumonia (J09-J18)	50	2.0	0.6
7	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	32	1.3	0.4
8	Septicemia (A40-A41)	30	1.2	0.4
8	Certain conditions originating in the perinatal period (P00-P96)	30	1.2	0.4
10	Chronic lower respiratory diseases (J40-J47)	28	1.1	0.3
11	Cerebrovascular diseases (I60-I69)	26	1.1	0.3
12	Anemias (D50-D64)	17	0.7	*
13	Meningitis (G00,G03)	13	0.5	*
13	Complications of medical and surgical care (Y40-Y84,Y88)	13	0.5	*
15	Pneumonitis due to solids and liquids (J69)	9	0.4	*
...	All other causes (Residual)	595	24.2	7.2
All races, male, 5-14 years				
...	All causes	3,054	100.0	14.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,052	34.4	5.0
2	Malignant neoplasms (C00-C97)	482	15.8	2.3
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	187	6.1	0.9
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	166	5.4	0.8
5	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	148	4.8	0.7
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	96	3.1	0.5
7	Chronic lower respiratory diseases (J40-J47)	90	2.9	0.4
8	Cerebrovascular diseases (I60-I69)	54	1.8	0.3
9	Septicemia (A40-A41)	38	1.2	0.2
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	35	1.1	0.2
11	Influenza and pneumonia (J09-J18)	33	1.1	0.2
12	Certain conditions originating in the perinatal period (P00-P96)	16	0.5	*
13	Diabetes mellitus (E10-E14)	14	0.5	*
14	Anemias (D50-D64)	11	0.4	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	11	0.4	*
...	All other causes (Residual)	621	20.3	3.0

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), race, sex, and age	Number	Percent of total deaths	Rate
All races, male, 15-24 years				
...	All causes	21,790	100.0	97.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	9,088	41.7	40.7
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	4,067	18.7	18.2
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	3,771	17.3	16.9
4	Malignant neoplasms (C00-C97)	997	4.6	4.5
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	719	3.3	3.2
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	251	1.2	1.1
7	Cerebrovascular diseases (I60-I69)	107	0.5	0.5
8	Influenza and pneumonia (J09-J18)	105	0.5	0.5
9	Human immunodeficiency virus (HIV) disease (B20-B24)	100	0.5	0.4
10	Chronic lower respiratory diseases (J40-J47)	96	0.4	0.4
11	Diabetes mellitus (E10-E14)	90	0.4	0.4
12	Septicemia (A40-A41)	76	0.3	0.3
13	Legal intervention (Y35,Y89.0)	72	0.3	0.3
14	Anemias (D50-D64)	70	0.3	0.3
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	58	0.3	0.3
...	All other causes (Residual)	2,123	9.7	9.5
All races, male, 25-44 years				
...	All causes	72,626	100.0	176.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	20,999	28.9	51.1
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	9,676	13.3	23.6
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	9,578	13.2	23.3
4	Malignant neoplasms (C00-C97)	6,930	9.5	16.9
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	5,452	7.5	13.3
6	Chronic liver disease and cirrhosis (K70,K73-K74)	1,919	2.6	4.7
7	Human immunodeficiency virus (HIV) disease (B20-B24)	1,776	2.4	4.3
8	Diabetes mellitus (E10-E14)	1,405	1.9	3.4
9	Cerebrovascular diseases (I60-I69)	1,293	1.8	3.1
10	Influenza and pneumonia (J09-J18)	658	0.9	1.6
11	Septicemia (A40-A41)	546	0.8	1.3
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	512	0.7	1.2
13	Chronic lower respiratory diseases (J40-J47)	504	0.7	1.2
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	484	0.7	1.2
15	Aortic aneurysm and dissection (I71)	343	0.5	0.8
...	All other causes (Residual)	10,551	14.5	25.7

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), race, sex, and age	Number	Percent of total deaths	Rate
All races, male, 45-64 years				
...	All causes	301,313	100.0	758.1
1	Malignant neoplasms (C00-C97)	85,888	28.5	216.1
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	73,520	24.4	185.0
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	22,787	7.6	57.3
4	Chronic liver disease and cirrhosis (K70,K73-K74)	12,843	4.3	32.3
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	11,602	3.9	29.2
6	Diabetes mellitus (E10-E14)	10,650	3.5	26.8
7	Chronic lower respiratory diseases (J40-J47)	9,472	3.1	23.8
8	Cerebrovascular diseases (I60-I69)	9,408	3.1	23.7
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	4,130	1.4	10.4
10	Viral hepatitis (B15-B19)	3,965	1.3	10.0
11	Human immunodeficiency virus (HIV) disease (B20-B24)	3,730	1.2	9.4
12	Septicemia (A40-A41)	3,699	1.2	9.3
13	Influenza and pneumonia (J09-J18)	3,206	1.1	8.1
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	2,507	0.8	6.3
15	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	2,227	0.7	5.6
...	All other causes (Residual)	41,679	13.8	104.9
All races, male, 65 years and over				
...	All causes	817,400	100.0	4,707.7
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	223,165	27.3	1,285.3
2	Malignant neoplasms (C00-C97)	206,507	25.3	1,189.4
3	Chronic lower respiratory diseases (J40-J47)	55,205	6.8	317.9
4	Cerebrovascular diseases (I60-I69)	41,414	5.1	238.5
5	Alzheimer's disease (G30)	24,995	3.1	144.0
6	Diabetes mellitus (E10-E14)	23,327	2.9	134.3
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	20,479	2.5	117.9
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	20,120	2.5	115.9
9	Influenza and pneumonia (J09-J18)	19,450	2.4	112.0
10	Parkinson's disease (G20-G21)	12,480	1.5	71.9
11	Septicemia (A40-A41)	11,560	1.4	66.6
12	Pneumonitis due to solids and liquids (J69)	8,089	1.0	46.6
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	8,017	1.0	46.2
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	6,082	0.7	35.0
15	Chronic liver disease and cirrhosis (K70,K73-K74)	6,014	0.7	34.6
...	All other causes (Residual)	130,496	16.0	751.6

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), race, sex, and age	Number	Percent of total deaths	Rate
All races, female, all ages				
...	All causes	1,236,003	100.0	787.4
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	290,305	23.5	184.9
2	Malignant neoplasms (C00-C97)	273,706	22.1	174.4
3	Cerebrovascular diseases (I60-I69)	77,109	6.2	49.1
4	Chronic lower respiratory diseases (J40-J47)	72,657	5.9	46.3
5	Alzheimer's disease (G30)	58,130	4.7	37.0
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	44,938	3.6	28.6
7	Diabetes mellitus (E10-E14)	33,581	2.7	21.4
8	Influenza and pneumonia (J09-J18)	26,482	2.1	16.9
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	25,611	2.1	16.3
10	Septicemia (A40-A41)	18,743	1.5	11.9
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	15,788	1.3	10.1
12	Chronic liver disease and cirrhosis (K70,K73-K74)	11,105	0.9	7.1
13	Parkinson's disease (G20-G21)	9,161	0.7	5.8
14	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	8,087	0.7	5.2
15	Pneumonitis due to solids and liquids (J69)	7,803	0.6	5.0
...	All other causes (Residual)	262,797	21.3	167.4
All races, female, 1-4 years				
...	All causes	1,856	100.0	23.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	526	28.3	6.6
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	276	14.9	3.5
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	154	8.3	1.9
4	Malignant neoplasms (C00-C97)	149	8.0	1.9
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	69	3.7	0.9
6	Influenza and pneumonia (J09-J18)	41	2.2	0.5
7	Septicemia (A40-A41)	32	1.7	0.4
8	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	27	1.5	0.3
9	Cerebrovascular diseases (I60-I69)	24	1.3	0.3
10	Chronic lower respiratory diseases (J40-J47)	23	1.2	0.3
11	Certain conditions originating in the perinatal period (P00-P96)	22	1.2	0.3
12	Anemias (D50-D64)	12	0.6	*
13	Meningitis (G00,G03)	9	0.5	*
14	Meningococcal infection (A39)	7	0.4	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	7	0.4	*
...	All other causes (Residual)	478	25.8	6.0

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), race, sex, and age	Number	Percent of total deaths	Rate
All races, female, 5-14 years				
...	All causes	2,225	100.0	11.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	591	26.6	2.9
2	Malignant neoplasms (C00-C97)	434	19.5	2.2
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	150	6.7	0.7
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	95	4.3	0.5
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	89	4.0	0.4
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	87	3.9	0.4
7	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	47	2.1	0.2
8	Chronic lower respiratory diseases (J40-J47)	43	1.9	0.2
9	Influenza and pneumonia (J09-J18)	38	1.7	0.2
10	Cerebrovascular diseases (I60-I69)	36	1.6	0.2
11	Septicemia (A40-A41)	29	1.3	0.1
12	Anemias (D50-D64)	15	0.7	*
13	Complications of medical and surgical care (Y40-Y84,Y88)	14	0.6	*
14	Diabetes mellitus (E10-E14)	12	0.5	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	11	0.5	*
...	All other causes (Residual)	534	24.0	2.7
All races, female, 15-24 years				
...	All causes	7,761	100.0	36.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,253	41.9	15.3
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	829	10.7	3.9
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	611	7.9	2.9
4	Malignant neoplasms (C00-C97)	607	7.8	2.8
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	309	4.0	1.5
6	Pregnancy, childbirth and the puerperium (O00-O99)	163	2.1	0.8
7	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	161	2.1	0.8
8	Cerebrovascular diseases (I60-I69)	83	1.1	0.4
9	Influenza and pneumonia (J09-J18)	76	1.0	0.4
10	Diabetes mellitus (E10-E14)	75	1.0	0.4
11	Septicemia (A40-A41)	65	0.8	0.3
12	Chronic lower respiratory diseases (J40-J47)	53	0.7	0.2
13	Human immunodeficiency virus (HIV) disease (B20-B24)	47	0.6	0.2
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	42	0.5	0.2
15	Anemias (D50-D64)	39	0.5	0.2
...	All other causes (Residual)	1,348	17.4	6.3

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), race, sex, and age	Number	Percent of total deaths	Rate
All races, female, 25-44 years				
...	All causes	39,666	100.0	96.6
1	Malignant neoplasms (C00-C97)	8,498	21.4	20.7
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	8,366	21.1	20.4
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	4,238	10.7	10.3
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,630	6.6	6.4
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,279	3.2	3.1
6	Cerebrovascular diseases (I60-I69)	1,128	2.8	2.7
7	Chronic liver disease and cirrhosis (K70,K73-K74)	991	2.5	2.4
8	Diabetes mellitus (E10-E14)	990	2.5	2.4
9	Human immunodeficiency virus (HIV) disease (B20-B24)	863	2.2	2.1
10	Pregnancy, childbirth and the puerperium (O00-O99)	590	1.5	1.4
11	Septicemia (A40-A41)	533	1.3	1.3
12	Influenza and pneumonia (J09-J18)	500	1.3	1.2
13	Chronic lower respiratory diseases (J40-J47)	477	1.2	1.2
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	457	1.2	1.1
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	362	0.9	0.9
...	All other causes (Residual)	7,764	19.6	18.9
All races, female, 45-64 years				
...	All causes	192,696	100.0	461.6
1	Malignant neoplasms (C00-C97)	73,824	38.3	176.8
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	31,286	16.2	74.9
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	10,903	5.7	26.1
4	Chronic lower respiratory diseases (J40-J47)	9,222	4.8	22.1
5	Cerebrovascular diseases (I60-I69)	7,195	3.7	17.2
6	Diabetes mellitus (E10-E14)	6,637	3.4	15.9
7	Chronic liver disease and cirrhosis (K70,K73-K74)	5,572	2.9	13.3
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	3,581	1.9	8.6
9	Septicemia (A40-A41)	3,238	1.7	7.8
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3,174	1.6	7.6
11	Influenza and pneumonia (J09-J18)	2,347	1.2	5.6
12	Viral hepatitis (B15-B19)	1,629	0.8	3.9
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,549	0.8	3.7
14	Human immunodeficiency virus (HIV) disease (B20-B24)	1,215	0.6	2.9
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	947	0.5	2.3
...	All other causes (Residual)	30,377	15.8	72.8

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), race, sex, and age	Number	Percent of total deaths	Rate
All races, female, 65 years and over				
...	All causes	980,876	100.0	4,282.4
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	254,173	25.9	1,109.7
2	Malignant neoplasms (C00-C97)	190,163	19.4	830.2
3	Cerebrovascular diseases (I60-I69)	68,576	7.0	299.4
4	Chronic lower respiratory diseases (J40-J47)	62,826	6.4	274.3
5	Alzheimer's disease (G30)	57,621	5.9	251.6
6	Diabetes mellitus (E10-E14)	25,864	2.6	112.9
7	Influenza and pneumonia (J09-J18)	23,396	2.4	102.1
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	21,874	2.2	95.5
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	20,821	2.1	90.9
10	Septicemia (A40-A41)	14,750	1.5	64.4
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	14,030	1.4	61.3
12	Parkinson's disease (G20-G21)	8,968	0.9	39.2
13	Pneumonitis due to solids and liquids (J69)	7,137	0.7	31.2
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	5,922	0.6	25.9
15	Chronic liver disease and cirrhosis (K70,K73-K74)	4,521	0.5	19.7
...	All other causes (Residual)	200,234	20.4	874.2
White, both sexes, all ages				
...	All causes	2,114,749	100.0	861.7
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	514,323	24.3	209.6
2	Malignant neoplasms (C00-C97)	491,686	23.3	200.3
3	Chronic lower respiratory diseases (J40-J47)	127,176	6.0	51.8
4	Cerebrovascular diseases (I60-I69)	109,119	5.2	44.5
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	104,945	5.0	42.8
6	Alzheimer's disease (G30)	76,928	3.6	31.3
7	Diabetes mellitus (E10-E14)	54,250	2.6	22.1
8	Influenza and pneumonia (J09-J18)	43,296	2.0	17.6
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	40,205	1.9	16.4
10	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	34,690	1.6	14.1
11	Chronic liver disease and cirrhosis (K70,K73-K74)	28,014	1.3	11.4
12	Septicemia (A40-A41)	27,985	1.3	11.4
13	Parkinson's disease (G20-G21)	20,792	1.0	8.5
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	20,560	1.0	8.4
15	Pneumonitis due to solids and liquids (J69)	15,235	0.7	6.2
...	All other causes (Residual)	405,545	19.2	165.2

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), race, sex, and age	Number	Percent of total deaths	Rate
White, both sexes, 1-4 years				
...	All causes	3,015	100.0	24.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,054	35.0	8.6
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	348	11.5	2.8
3	Malignant neoplasms (C00-C97)	269	8.9	2.2
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	214	7.1	1.7
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	95	3.2	0.8
6	Influenza and pneumonia (J09-J18)	61	2.0	0.5
7	Septicemia (A40-A41)	47	1.6	0.4
8	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	43	1.4	0.4
9	Certain conditions originating in the perinatal period (P00-P96)	34	1.1	0.3
10	Cerebrovascular diseases (I60-I69)	30	1.0	0.2
11	Chronic lower respiratory diseases (J40-J47)	23	0.8	0.2
12	Meningitis (G00,G03)	16	0.5	*
13	Anemias (D50-D64)	14	0.5	*
14	Complications of medical and surgical care (Y40-Y84,Y88)	13	0.4	*
15	Acute bronchitis and bronchiolitis (J20-J21)	10	0.3	*
...	All other causes (Residual)	744	24.7	6.1
White, both sexes, 5-14 years				
...	All causes	3,841	100.0	12.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,248	32.5	4.0
2	Malignant neoplasms (C00-C97)	707	18.4	2.3
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	214	5.6	0.7
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	209	5.4	0.7
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	148	3.9	0.5
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	116	3.0	0.4
7	Cerebrovascular diseases (I60-I69)	61	1.6	0.2
8	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	60	1.6	0.2
9	Influenza and pneumonia (J09-J18)	54	1.4	0.2
10	Chronic lower respiratory diseases (J40-J47)	53	1.4	0.2
11	Septicemia (A40-A41)	44	1.1	0.1
12	Diabetes mellitus (E10-E14)	19	0.5	*
13	Certain conditions originating in the perinatal period (P00-P96)	17	0.4	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	15	0.4	*
15	Complications of medical and surgical care (Y40-Y84,Y88)	14	0.4	*
...	All other causes (Residual)	862	22.4	2.8

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), race, sex, and age	Number	Percent of total deaths	Rate
White, both sexes, 15-24 years				
...	All causes	21,509	100.0	64.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	10,358	48.2	31.2
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	3,792	17.6	11.4
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,724	8.0	5.2
4	Malignant neoplasms (C00-C97)	1,243	5.8	3.7
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	639	3.0	1.9
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	304	1.4	0.9
7	Influenza and pneumonia (J09-J18)	134	0.6	0.4
8	Cerebrovascular diseases (I60-I69)	129	0.6	0.4
9	Diabetes mellitus (E10-E14)	113	0.5	0.3
10	Pregnancy, childbirth and the puerperium (O00-O99)	106	0.5	0.3
11	Septicemia (A40-A41)	95	0.4	0.3
12	Chronic lower respiratory diseases (J40-J47)	82	0.4	0.2
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	68	0.3	0.2
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	44	0.2	0.1
15	Legal intervention (Y35,Y89.0)	41	0.2	0.1
...	All other causes (Residual)	2,637	12.3	7.9
White, both sexes, 25-44 years				
...	All causes	84,485	100.0	132.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	25,083	29.7	39.3
2	Malignant neoplasms (C00-C97)	11,858	14.0	18.6
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	10,772	12.8	16.9
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	9,644	11.4	15.1
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	2,996	3.5	4.7
6	Chronic liver disease and cirrhosis (K70,K73-K74)	2,419	2.9	3.8
7	Diabetes mellitus (E10-E14)	1,581	1.9	2.5
8	Cerebrovascular diseases (I60-I69)	1,566	1.9	2.5
9	Human immunodeficiency virus (HIV) disease (B20-B24)	1,051	1.2	1.6
10	Influenza and pneumonia (J09-J18)	871	1.0	1.4
11	Septicemia (A40-A41)	736	0.9	1.2
12	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	671	0.8	1.1
13	Chronic lower respiratory diseases (J40-J47)	669	0.8	1.0
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	557	0.7	0.9
15	Pregnancy, childbirth and the puerperium (O00-O99)	344	0.4	0.5
...	All other causes (Residual)	13,667	16.2	21.4

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), race, sex, and age	Number	Percent of total deaths	Rate
White, both sexes, 45-64 years				
...	All causes	392,632	100.0	587.5
1	Malignant neoplasms (C00-C97)	129,190	32.9	193.3
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	81,037	20.6	121.3
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	28,647	7.3	42.9
4	Chronic lower respiratory diseases (J40-J47)	16,304	4.2	24.4
5	Chronic liver disease and cirrhosis (K70,K73-K74)	15,913	4.1	23.8
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	14,199	3.6	21.2
7	Diabetes mellitus (E10-E14)	12,681	3.2	19.0
8	Cerebrovascular diseases (I60-I69)	11,389	2.9	17.0
9	Septicemia (A40-A41)	5,042	1.3	7.5
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	4,709	1.2	7.0
11	Viral hepatitis (B15-B19)	4,559	1.2	6.8
12	Influenza and pneumonia (J09-J18)	4,367	1.1	6.5
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	2,506	0.6	3.7
14	Human immunodeficiency virus (HIV) disease (B20-B24)	2,197	0.6	3.3
15	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,921	0.5	2.9
...	All other causes (Residual)	57,971	14.8	86.7
White, both sexes, 65 years and over				
...	All causes	1,593,218	100.0	4,555.8
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	422,559	26.5	1,208.3
2	Malignant neoplasms (C00-C97)	348,368	21.9	996.2
3	Chronic lower respiratory diseases (J40-J47)	110,020	6.9	314.6
4	Cerebrovascular diseases (I60-I69)	95,868	6.0	274.1
5	Alzheimer's disease (G30)	76,124	4.8	217.7
6	Diabetes mellitus (E10-E14)	39,853	2.5	114.0
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	37,839	2.4	108.2
8	Influenza and pneumonia (J09-J18)	37,706	2.4	107.8
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	34,812	2.2	99.5
10	Septicemia (A40-A41)	21,906	1.4	62.6
11	Parkinson's disease (G20-G21)	20,274	1.3	58.0
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	17,786	1.1	50.9
13	Pneumonitis due to solids and liquids (J69)	13,785	0.9	39.4
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	10,929	0.7	31.3
15	Chronic liver disease and cirrhosis (K70,K73-K74)	9,649	0.6	27.6
...	All other causes (Residual)	295,740	18.6	845.7

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), race, sex, and age	Number	Percent of total deaths	Rate
White, male, all ages				
...	All causes	1,051,514	100.0	866.1
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	264,425	25.1	217.8
2	Malignant neoplasms (C00-C97)	258,272	24.6	212.7
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	65,360	6.2	53.8
4	Chronic lower respiratory diseases (J40-J47)	59,632	5.7	49.1
5	Cerebrovascular diseases (I60-I69)	43,424	4.1	35.8
6	Diabetes mellitus (E10-E14)	28,486	2.7	23.5
7	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	27,422	2.6	22.6
8	Alzheimer's disease (G30)	23,442	2.2	19.3
9	Influenza and pneumonia (J09-J18)	20,238	1.9	16.7
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	20,172	1.9	16.6
11	Chronic liver disease and cirrhosis (K70,K73-K74)	18,352	1.7	15.1
12	Septicemia (A40-A41)	12,976	1.2	10.7
13	Parkinson's disease (G20-G21)	12,173	1.2	10.0
14	Pneumonitis due to solids and liquids (J69)	8,264	0.8	6.8
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	8,229	0.8	6.8
...	All other causes (Residual)	180,647	17.2	148.8
White, male, 1-4 years				
...	All causes	1,718	100.0	27.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	653	38.0	10.4
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	160	9.3	2.5
3	Malignant neoplasms (C00-C97)	149	8.7	2.4
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	130	7.6	2.1
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	53	3.1	0.8
6	Influenza and pneumonia (J09-J18)	33	1.9	0.5
7	Septicemia (A40-A41)	23	1.3	0.4
7	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	23	1.3	0.4
9	Certain conditions originating in the perinatal period (P00-P96)	19	1.1	*
10	Cerebrovascular diseases (I60-I69)	18	1.0	*
11	Chronic lower respiratory diseases (J40-J47)	14	0.8	*
12	Meningitis (G00,G03)	11	0.6	*
13	Complications of medical and surgical care (Y40-Y84,Y88)	9	0.5	*
14	Anemias (D50-D64)	8	0.5	*
15	Acute bronchitis and bronchiolitis (J20-J21)	7	0.4	*
...	All other causes (Residual)	408	23.7	6.5

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), race, sex, and age	Number	Percent of total deaths	Rate
White, male, 5-14 years				
...	All causes	2,222	100.0	13.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	785	35.3	4.9
2	Malignant neoplasms (C00-C97)	386	17.4	2.4
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	151	6.8	0.9
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	104	4.7	0.6
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	90	4.1	0.6
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	54	2.4	0.3
7	Cerebrovascular diseases (I60-I69)	36	1.6	0.2
8	Chronic lower respiratory diseases (J40-J47)	31	1.4	0.2
9	Septicemia (A40-A41)	26	1.2	0.2
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	25	1.1	0.2
11	Influenza and pneumonia (J09-J18)	23	1.0	0.1
12	Certain conditions originating in the perinatal period (P00-P96)	13	0.6	*
13	Diabetes mellitus (E10-E14)	11	0.5	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	7	0.3	*
14	Complications of medical and surgical care (Y40-Y84,Y88)	7	0.3	*
...	All other causes (Residual)	473	21.3	2.9
White, male, 15-24 years				
...	All causes	15,661	100.0	91.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	7,620	48.7	44.6
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	3,116	19.9	18.3
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,399	8.9	8.2
4	Malignant neoplasms (C00-C97)	788	5.0	4.6
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	471	3.0	2.8
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	184	1.2	1.1
7	Cerebrovascular diseases (I60-I69)	80	0.5	0.5
8	Influenza and pneumonia (J09-J18)	76	0.5	0.4
9	Diabetes mellitus (E10-E14)	63	0.4	0.4
10	Chronic lower respiratory diseases (J40-J47)	49	0.3	0.3
11	Septicemia (A40-A41)	46	0.3	0.3
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	41	0.3	0.2
13	Legal intervention (Y35,Y89.0)	40	0.3	0.2
14	Human immunodeficiency virus (HIV) disease (B20-B24)	27	0.2	0.2
14	Aortic aneurysm and dissection (I71)	27	0.2	0.2
...	All other causes (Residual)	1,634	10.4	9.6

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), race, sex, and age	Number	Percent of total deaths	Rate
White, male, 25-44 years				
...	All causes	55,369	100.0	171.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	17,960	32.4	55.5
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	8,476	15.3	26.2
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	6,915	12.5	21.4
4	Malignant neoplasms (C00-C97)	5,491	9.9	17.0
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	2,228	4.0	6.9
6	Chronic liver disease and cirrhosis (K70,K73-K74)	1,638	3.0	5.1
7	Diabetes mellitus (E10-E14)	939	1.7	2.9
8	Cerebrovascular diseases (I60-I69)	838	1.5	2.6
9	Human immunodeficiency virus (HIV) disease (B20-B24)	815	1.5	2.5
10	Influenza and pneumonia (J09-J18)	484	0.9	1.5
11	Septicemia (A40-A41)	384	0.7	1.2
12	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	383	0.7	1.2
13	Chronic lower respiratory diseases (J40-J47)	325	0.6	1.0
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	295	0.5	0.9
15	Aortic aneurysm and dissection (I71)	222	0.4	0.7
...	All other causes (Residual)	7,976	14.4	24.7
White, male, 45-64 years				
...	All causes	242,313	100.0	735.7
1	Malignant neoplasms (C00-C97)	69,890	28.8	212.2
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	58,270	24.0	176.9
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	19,319	8.0	58.7
4	Chronic liver disease and cirrhosis (K70,K73-K74)	11,188	4.6	34.0
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	10,858	4.5	33.0
6	Chronic lower respiratory diseases (J40-J47)	8,232	3.4	25.0
7	Diabetes mellitus (E10-E14)	8,040	3.3	24.4
8	Cerebrovascular diseases (I60-I69)	6,506	2.7	19.8
9	Viral hepatitis (B15-B19)	3,250	1.3	9.9
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2,745	1.1	8.3
11	Septicemia (A40-A41)	2,717	1.1	8.2
12	Influenza and pneumonia (J09-J18)	2,503	1.0	7.6
13	Human immunodeficiency virus (HIV) disease (B20-B24)	1,859	0.8	5.6
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,641	0.7	5.0
15	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,345	0.6	4.1
...	All other causes (Residual)	33,950	14.0	103.1

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), race, sex, and age	Number	Percent of total deaths	Rate
White, male, 65 years and over				
...	All causes	725,294	100.0	4,768.8
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	198,524	27.4	1,305.3
2	Malignant neoplasms (C00-C97)	181,539	25.0	1,193.6
3	Chronic lower respiratory diseases (J40-J47)	50,967	7.0	335.1
4	Cerebrovascular diseases (I60-I69)	35,906	5.0	236.1
5	Alzheimer's disease (G30)	23,105	3.2	151.9
6	Diabetes mellitus (E10-E14)	19,432	2.7	127.8
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	18,612	2.6	122.4
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	17,069	2.4	112.2
9	Influenza and pneumonia (J09-J18)	17,061	2.4	112.2
10	Parkinson's disease (G20-G21)	11,829	1.6	77.8
11	Septicemia (A40-A41)	9,710	1.3	63.8
12	Pneumonitis due to solids and liquids (J69)	7,352	1.0	48.3
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	6,414	0.9	42.2
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	5,611	0.8	36.9
15	Chronic liver disease and cirrhosis (K70,K73-K74)	5,509	0.8	36.2
...	All other causes (Residual)	116,654	16.1	767.0
White, female, all ages				
...	All causes	1,063,235	100.0	857.3
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	249,898	23.5	201.5
2	Malignant neoplasms (C00-C97)	233,414	22.0	188.2
3	Chronic lower respiratory diseases (J40-J47)	67,544	6.4	54.5
4	Cerebrovascular diseases (I60-I69)	65,695	6.2	53.0
5	Alzheimer's disease (G30)	53,486	5.0	43.1
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	39,585	3.7	31.9
7	Diabetes mellitus (E10-E14)	25,764	2.4	20.8
8	Influenza and pneumonia (J09-J18)	23,058	2.2	18.6
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	20,033	1.9	16.2
10	Septicemia (A40-A41)	15,009	1.4	12.1
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	12,331	1.2	9.9
12	Chronic liver disease and cirrhosis (K70,K73-K74)	9,662	0.9	7.8
13	Parkinson's disease (G20-G21)	8,619	0.8	6.9
14	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	7,268	0.7	5.9
15	Pneumonitis due to solids and liquids (J69)	6,971	0.7	5.6
...	All other causes (Residual)	224,898	21.2	181.3

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), race, sex, and age	Number	Percent of total deaths	Rate
White, female, 1-4 years				
...	All causes	1,297	100.0	21.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	401	30.9	6.7
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	188	14.5	3.1
3	Malignant neoplasms (C00-C97)	120	9.3	2.0
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	84	6.5	1.4
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	42	3.2	0.7
6	Influenza and pneumonia (J09-J18)	28	2.2	0.5
7	Septicemia (A40-A41)	24	1.9	0.4
8	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	20	1.5	0.3
9	Certain conditions originating in the perinatal period (P00-P96)	15	1.2	*
10	Cerebrovascular diseases (I60-I69)	12	0.9	*
11	Chronic lower respiratory diseases (J40-J47)	9	0.7	*
12	Meningococcal infection (A39)	7	0.5	*
13	Anemias (D50-D64)	6	0.5	*
14	Meningitis (G00,G03)	5	0.4	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	4	0.3	*
15	Complications of medical and surgical care (Y40-Y84,Y88)	4	0.3	*
...	All other causes (Residual)	328	25.3	5.5
White, female, 5-14 years				
...	All causes	1,619	100.0	10.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	463	28.6	3.0
2	Malignant neoplasms (C00-C97)	321	19.8	2.1
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	110	6.8	0.7
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	62	3.8	0.4
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	58	3.6	0.4
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	58	3.6	0.4
7	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	35	2.2	0.2
8	Influenza and pneumonia (J09-J18)	31	1.9	0.2
9	Cerebrovascular diseases (I60-I69)	25	1.5	0.2
10	Chronic lower respiratory diseases (J40-J47)	22	1.4	0.1
11	Septicemia (A40-A41)	18	1.1	*
12	Diabetes mellitus (E10-E14)	8	0.5	*
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	8	0.5	*
14	Complications of medical and surgical care (Y40-Y84,Y88)	7	0.4	*
15	Anemias (D50-D64)	6	0.4	*
15	Meningitis (G00,G03)	6	0.4	*
15	Pneumonitis due to solids and liquids (J69)	6	0.4	*
...	All other causes (Residual)	375	23.2	2.5

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), race, sex, and age	Number	Percent of total deaths	Rate
White, female, 15-24 years				
...	All causes	5,848	100.0	36.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,738	46.8	17.0
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	676	11.6	4.2
3	Malignant neoplasms (C00-C97)	455	7.8	2.8
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	325	5.6	2.0
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	168	2.9	1.0
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	120	2.1	0.7
7	Pregnancy, childbirth and the puerperium (O00-O99)	106	1.8	0.7
8	Influenza and pneumonia (J09-J18)	58	1.0	0.4
9	Diabetes mellitus (E10-E14)	50	0.9	0.3
10	Septicemia (A40-A41)	49	0.8	0.3
10	Cerebrovascular diseases (I60-I69)	49	0.8	0.3
12	Chronic lower respiratory diseases (J40-J47)	33	0.6	0.2
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	27	0.5	0.2
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	25	0.4	0.2
15	Complications of medical and surgical care (Y40-Y84,Y88)	14	0.2	*
...	All other causes (Residual)	955	16.3	5.9
White, female, 25-44 years				
...	All causes	29,116	100.0	92.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	7,123	24.5	22.6
2	Malignant neoplasms (C00-C97)	6,367	21.9	20.2
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,729	9.4	8.7
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,296	7.9	7.3
5	Chronic liver disease and cirrhosis (K70,K73-K74)	781	2.7	2.5
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	768	2.6	2.4
7	Cerebrovascular diseases (I60-I69)	728	2.5	2.3
8	Diabetes mellitus (E10-E14)	642	2.2	2.0
9	Influenza and pneumonia (J09-J18)	387	1.3	1.2
10	Septicemia (A40-A41)	352	1.2	1.1
11	Chronic lower respiratory diseases (J40-J47)	344	1.2	1.1
11	Pregnancy, childbirth and the puerperium (O00-O99)	344	1.2	1.1
13	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	288	1.0	0.9
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	262	0.9	0.8
15	Human immunodeficiency virus (HIV) disease (B20-B24)	236	0.8	0.7
...	All other causes (Residual)	5,469	18.8	17.4

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), race, sex, and age	Number	Percent of total deaths	Rate
White, female, 45-64 years				
...	All causes	150,319	100.0	443.5
1	Malignant neoplasms (C00-C97)	59,300	39.4	174.9
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	22,767	15.1	67.2
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	9,328	6.2	27.5
4	Chronic lower respiratory diseases (J40-J47)	8,072	5.4	23.8
5	Cerebrovascular diseases (I60-I69)	4,883	3.2	14.4
6	Chronic liver disease and cirrhosis (K70,K73-K74)	4,725	3.1	13.9
7	Diabetes mellitus (E10-E14)	4,641	3.1	13.7
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	3,341	2.2	9.9
9	Septicemia (A40-A41)	2,325	1.5	6.9
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,964	1.3	5.8
11	Influenza and pneumonia (J09-J18)	1,864	1.2	5.5
12	Viral hepatitis (B15-B19)	1,309	0.9	3.9
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	865	0.6	2.6
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	762	0.5	2.2
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	592	0.4	1.7
...	All other causes (Residual)	23,581	15.7	69.6
White, female, 65 years and over				
...	All causes	867,924	100.0	4,391.9
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	224,035	25.8	1,133.7
2	Malignant neoplasms (C00-C97)	166,829	19.2	844.2
3	Cerebrovascular diseases (I60-I69)	59,962	6.9	303.4
4	Chronic lower respiratory diseases (J40-J47)	59,053	6.8	298.8
5	Alzheimer's disease (G30)	53,019	6.1	268.3
6	Influenza and pneumonia (J09-J18)	20,645	2.4	104.5
7	Diabetes mellitus (E10-E14)	20,421	2.4	103.3
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	19,227	2.2	97.3
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	17,743	2.0	89.8
10	Septicemia (A40-A41)	12,196	1.4	61.7
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	11,372	1.3	57.5
12	Parkinson's disease (G20-G21)	8,445	1.0	42.7
13	Pneumonitis due to solids and liquids (J69)	6,433	0.7	32.6
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	5,318	0.6	26.9
15	Chronic liver disease and cirrhosis (K70,K73-K74)	4,140	0.5	20.9
...	All other causes (Residual)	179,086	20.6	906.2

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), race, sex, and age	Number	Percent of total deaths	Rate
Black, both sexes, all ages				
...	All causes	286,959	100.0	682.2
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	69,083	24.1	164.2
2	Malignant neoplasms (C00-C97)	65,930	23.0	156.7
3	Cerebrovascular diseases (I60-I69)	15,965	5.6	38.0
4	Diabetes mellitus (E10-E14)	12,126	4.2	28.8
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	12,069	4.2	28.7
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	8,841	3.1	21.0
7	Chronic lower respiratory diseases (J40-J47)	8,715	3.0	20.7
8	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	7,818	2.7	18.6
9	Septicemia (A40-A41)	6,001	2.1	14.3
10	Alzheimer's disease (G30)	5,220	1.8	12.4
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	5,116	1.8	12.2
12	Influenza and pneumonia (J09-J18)	4,936	1.7	11.7
13	Human immunodeficiency virus (HIV) disease (B20-B24)	4,662	1.6	11.1
14	Certain conditions originating in the perinatal period (P00-P96)	4,031	1.4	9.6
15	Chronic liver disease and cirrhosis (K70,K73-K74)	2,635	0.9	6.3
...	All other causes (Residual)	53,811	18.8	127.9
Black, both sexes, 1-4 years				
...	All causes	1,041	100.0	38.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	267	25.6	9.8
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	156	15.0	5.7
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	127	12.2	4.6
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	51	4.9	1.9
5	Malignant neoplasms (C00-C97)	48	4.6	1.8
6	Influenza and pneumonia (J09-J18)	25	2.4	0.9
7	Chronic lower respiratory diseases (J40-J47)	21	2.0	0.8
8	Certain conditions originating in the perinatal period (P00-P96)	16	1.5	*
9	Cerebrovascular diseases (I60-I69)	15	1.4	*
10	Anemias (D50-D64)	14	1.3	*
11	Septicemia (A40-A41)	12	1.2	*
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	10	1.0	*
13	Complications of medical and surgical care (Y40-Y84,Y88)	5	0.5	*
14	Meningitis (G00,G03)	4	0.4	*
14	Pneumonitis due to solids and liquids (J69)	4	0.4	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	4	0.4	*
...	All other causes (Residual)	262	25.2	9.6

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), race, sex, and age	Number	Percent of total deaths	Rate
Black, both sexes, 5-14 years				
...	All causes	1,145	100.0	17.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	319	27.9	4.8
2	Malignant neoplasms (C00-C97)	146	12.8	2.2
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	101	8.8	1.5
4	Chronic lower respiratory diseases (J40-J47)	72	6.3	1.1
5	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	63	5.5	0.9
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	61	5.3	0.9
7	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	44	3.8	0.7
8	Cerebrovascular diseases (I60-I69)	22	1.9	0.3
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	17	1.5	*
10	Septicemia (A40-A41)	15	1.3	*
11	Anemias (D50-D64)	14	1.2	*
12	Influenza and pneumonia (J09-J18)	12	1.0	*
13	Diabetes mellitus (E10-E14)	7	0.6	*
14	Certain conditions originating in the perinatal period (P00-P96)	6	0.5	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	5	0.4	*
...	All other causes (Residual)	241	21.0	3.6
Black, both sexes, 15-24 years				
...	All causes	6,675	100.0	93.2
1	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	2,816	42.2	39.3
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,457	21.8	20.4
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	469	7.0	6.6
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	336	5.0	4.7
5	Malignant neoplasms (C00-C97)	267	4.0	3.7
6	Human immunodeficiency virus (HIV) disease (B20-B24)	111	1.7	1.6
7	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	91	1.4	1.3
8	Anemias (D50-D64)	79	1.2	1.1
9	Chronic lower respiratory diseases (J40-J47)	62	0.9	0.9
10	Diabetes mellitus (E10-E14)	51	0.8	0.7
10	Pregnancy, childbirth and the puerperium (O00-O99)	51	0.8	0.7
12	Cerebrovascular diseases (I60-I69)	49	0.7	0.7
13	Septicemia (A40-A41)	41	0.6	0.6
14	Influenza and pneumonia (J09-J18)	39	0.6	0.5
15	Legal intervention (Y35,Y89.0)	28	0.4	0.4
...	All other causes (Residual)	728	10.9	10.2

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), race, sex, and age	Number	Percent of total deaths	Rate
Black, both sexes, 25-44 years				
...	All causes	22,936	100.0	200.0
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,640	15.9	31.7
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	3,489	15.2	30.4
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,214	14.0	28.0
4	Malignant neoplasms (C00-C97)	2,700	11.8	23.5
5	Human immunodeficiency virus (HIV) disease (B20-B24)	1,549	6.8	13.5
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	956	4.2	8.3
7	Diabetes mellitus (E10-E14)	723	3.2	6.3
8	Cerebrovascular diseases (I60-I69)	696	3.0	6.1
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	376	1.6	3.3
10	Septicemia (A40-A41)	298	1.3	2.6
11	Chronic lower respiratory diseases (J40-J47)	280	1.2	2.4
12	Anemias (D50-D64)	241	1.1	2.1
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	230	1.0	2.0
14	Influenza and pneumonia (J09-J18)	229	1.0	2.0
15	Chronic liver disease and cirrhosis (K70,K73-K74)	217	0.9	1.9
...	All other causes (Residual)	4,098	17.9	35.7
Black, both sexes, 45-64 years				
...	All causes	85,909	100.0	877.2
1	Malignant neoplasms (C00-C97)	24,936	29.0	254.6
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	20,945	24.4	213.9
3	Cerebrovascular diseases (I60-I69)	4,378	5.1	44.7
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	4,074	4.7	41.6
5	Diabetes mellitus (E10-E14)	3,883	4.5	39.6
6	Human immunodeficiency virus (HIV) disease (B20-B24)	2,664	3.1	27.2
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2,338	2.7	23.9
8	Chronic lower respiratory diseases (J40-J47)	2,112	2.5	21.6
9	Chronic liver disease and cirrhosis (K70,K73-K74)	1,843	2.1	18.8
10	Septicemia (A40-A41)	1,712	2.0	17.5
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,416	1.6	14.5
12	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,016	1.2	10.4
13	Influenza and pneumonia (J09-J18)	991	1.2	10.1
14	Viral hepatitis (B15-B19)	816	0.9	8.3
15	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	543	0.6	5.5
...	All other causes (Residual)	12,242	14.2	125.0

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), race, sex, and age	Number	Percent of total deaths	Rate
Black, both sexes, 65 years and over				
...	All causes	161,829	100.0	4,569.0
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	43,949	27.2	1,240.8
2	Malignant neoplasms (C00-C97)	37,824	23.4	1,067.9
3	Cerebrovascular diseases (I60-I69)	10,757	6.6	303.7
4	Diabetes mellitus (E10-E14)	7,458	4.6	210.6
5	Chronic lower respiratory diseases (J40-J47)	6,159	3.8	173.9
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	6,061	3.7	171.1
7	Alzheimer's disease (G30)	5,159	3.2	145.7
8	Septicemia (A40-A41)	3,833	2.4	108.2
9	Influenza and pneumonia (J09-J18)	3,567	2.2	100.7
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	3,457	2.1	97.6
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,382	1.5	67.3
12	Pneumonitis due to solids and liquids (J69)	1,093	0.7	30.9
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	769	0.5	21.7
14	Parkinson's disease (G20-G21)	756	0.5	21.3
15	Chronic liver disease and cirrhosis (K70,K73-K74)	570	0.4	16.1
...	All other causes (Residual)	28,035	17.3	791.5
Black, male, all ages				
...	All causes	145,802	100.0	725.4
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	35,089	24.1	174.6
2	Malignant neoplasms (C00-C97)	33,967	23.3	169.0
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	8,074	5.5	40.2
4	Cerebrovascular diseases (I60-I69)	6,938	4.8	34.5
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	6,704	4.6	33.4
6	Diabetes mellitus (E10-E14)	5,640	3.9	28.1
7	Chronic lower respiratory diseases (J40-J47)	4,532	3.1	22.5
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	4,016	2.8	20.0
9	Human immunodeficiency virus (HIV) disease (B20-B24)	3,047	2.1	15.2
10	Septicemia (A40-A41)	2,691	1.8	13.4
11	Influenza and pneumonia (J09-J18)	2,380	1.6	11.8
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	2,218	1.5	11.0
13	Certain conditions originating in the perinatal period (P00-P96)	2,216	1.5	11.0
14	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,755	1.2	8.7
15	Chronic liver disease and cirrhosis (K70,K73-K74)	1,715	1.2	8.5
...	All other causes (Residual)	24,820	17.0	123.5

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), race, sex, and age	Number	Percent of total deaths	Rate
Black, male, 1-4 years				
...	All causes	595	100.0	42.9
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	170	28.6	12.2
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	93	15.6	6.7
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	57	9.6	4.1
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	29	4.9	2.1
5	Malignant neoplasms (C00-C97)	27	4.5	1.9
6	Influenza and pneumonia (J09-J18)	15	2.5	*
7	Chronic lower respiratory diseases (J40-J47)	11	1.8	*
8	Anemias (D50-D64)	9	1.5	*
8	Certain conditions originating in the perinatal period (P00-P96)	9	1.5	*
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	6	1.0	*
11	Septicemia (A40-A41)	5	0.8	*
11	Cerebrovascular diseases (I60-I69)	5	0.8	*
13	Meningococcal infection (A39)	3	0.5	*
13	Pneumonitis due to solids and liquids (J69)	3	0.5	*
13	Complications of medical and surgical care (Y40-Y84,Y88)	3	0.5	*
...	All other causes (Residual)	150	25.2	10.8
Black, male, 5-14 years				
...	All causes	667	100.0	19.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	213	31.9	6.2
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	69	10.3	2.0
3	Malignant neoplasms (C00-C97)	65	9.7	1.9
4	Chronic lower respiratory diseases (J40-J47)	52	7.8	1.5
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	39	5.8	1.1
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	35	5.2	1.0
7	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	23	3.4	0.7
8	Cerebrovascular diseases (I60-I69)	14	2.1	*
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	9	1.3	*
10	Septicemia (A40-A41)	8	1.2	*
10	Influenza and pneumonia (J09-J18)	8	1.2	*
12	Anemias (D50-D64)	6	0.9	*
13	Diabetes mellitus (E10-E14)	3	0.4	*
13	Certain conditions originating in the perinatal period (P00-P96)	3	0.4	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2	0.3	*
...	All other causes (Residual)	118	17.7	3.5

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), race, sex, and age	Number	Percent of total deaths	Rate
Black, male, 15-24 years				
...	All causes	5,129	100.0	142.8
1	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	2,548	49.7	71.0
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,071	20.9	29.8
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	397	7.7	11.1
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	211	4.1	5.9
5	Malignant neoplasms (C00-C97)	149	2.9	4.1
6	Human immunodeficiency virus (HIV) disease (B20-B24)	71	1.4	2.0
7	Anemias (D50-D64)	55	1.1	1.5
7	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	55	1.1	1.5
9	Chronic lower respiratory diseases (J40-J47)	43	0.8	1.2
10	Legal intervention (Y35,Y89.0)	27	0.5	0.8
11	Septicemia (A40-A41)	26	0.5	0.7
11	Diabetes mellitus (E10-E14)	26	0.5	0.7
13	Influenza and pneumonia (J09-J18)	24	0.5	0.7
14	Cerebrovascular diseases (I60-I69)	21	0.4	0.6
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	12	0.2	*
...	All other causes (Residual)	393	7.7	10.9
Black, male, 25-44 years				
...	All causes	14,186	100.0	260.8
1	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	3,043	21.5	55.9
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,291	16.1	42.1
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,251	15.9	41.4
4	Malignant neoplasms (C00-C97)	1,051	7.4	19.3
5	Human immunodeficiency virus (HIV) disease (B20-B24)	929	6.5	17.1
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	788	5.6	14.5
7	Diabetes mellitus (E10-E14)	417	2.9	7.7
8	Cerebrovascular diseases (I60-I69)	364	2.6	6.7
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	199	1.4	3.7
10	Chronic lower respiratory diseases (J40-J47)	160	1.1	2.9
11	Septicemia (A40-A41)	140	1.0	2.6
12	Influenza and pneumonia (J09-J18)	135	1.0	2.5
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	127	0.9	2.3
14	Anemias (D50-D64)	118	0.8	2.2
14	Chronic liver disease and cirrhosis (K70,K73-K74)	118	0.8	2.2
...	All other causes (Residual)	2,055	14.5	37.8

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), race, sex, and age	Number	Percent of total deaths	Rate
Black, male, 45-64 years				
...	All causes	49,851	100.0	1,098.0
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	13,191	26.5	290.6
2	Malignant neoplasms (C00-C97)	13,186	26.5	290.4
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,825	5.7	62.2
4	Cerebrovascular diseases (I60-I69)	2,435	4.9	53.6
5	Diabetes mellitus (E10-E14)	2,146	4.3	47.3
6	Human immunodeficiency virus (HIV) disease (B20-B24)	1,807	3.6	39.8
7	Chronic liver disease and cirrhosis (K70,K73-K74)	1,249	2.5	27.5
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,246	2.5	27.4
9	Chronic lower respiratory diseases (J40-J47)	1,082	2.2	23.8
10	Septicemia (A40-A41)	893	1.8	19.7
11	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	797	1.6	17.6
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	789	1.6	17.4
13	Influenza and pneumonia (J09-J18)	574	1.2	12.6
14	Viral hepatitis (B15-B19)	563	1.1	12.4
15	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	432	0.9	9.5
...	All other causes (Residual)	6,636	13.3	146.2
Black, male, 65 years and over				
...	All causes	71,240	100.0	5,118.1
1	Malignant neoplasms (C00-C97)	19,485	27.4	1,399.8
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	19,301	27.1	1,386.6
3	Cerebrovascular diseases (I60-I69)	4,080	5.7	293.1
4	Chronic lower respiratory diseases (J40-J47)	3,177	4.5	228.2
5	Diabetes mellitus (E10-E14)	3,045	4.3	218.8
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2,543	3.6	182.7
7	Influenza and pneumonia (J09-J18)	1,581	2.2	113.6
8	Septicemia (A40-A41)	1,572	2.2	112.9
9	Alzheimer's disease (G30)	1,460	2.0	104.9
10	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,300	1.8	93.4
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,293	1.8	92.9
12	Pneumonitis due to solids and liquids (J69)	544	0.8	39.1
13	Parkinson's disease (G20-G21)	420	0.6	30.2
14	Chronic liver disease and cirrhosis (K70,K73-K74)	346	0.5	24.9
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	318	0.4	22.8
...	All other causes (Residual)	10,775	15.1	774.1

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), race, sex, and age	Number	Percent of total deaths	Rate
Black, female, all ages				
...	All causes	141,157	100.0	642.7
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	33,994	24.1	154.8
2	Malignant neoplasms (C00-C97)	31,963	22.6	145.5
3	Cerebrovascular diseases (I60-I69)	9,027	6.4	41.1
4	Diabetes mellitus (E10-E14)	6,486	4.6	29.5
5	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	4,825	3.4	22.0
6	Chronic lower respiratory diseases (J40-J47)	4,183	3.0	19.0
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,995	2.8	18.2
8	Alzheimer's disease (G30)	3,732	2.6	17.0
9	Septicemia (A40-A41)	3,310	2.3	15.1
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	2,898	2.1	13.2
11	Influenza and pneumonia (J09-J18)	2,556	1.8	11.6
12	Certain conditions originating in the perinatal period (P00-P96)	1,815	1.3	8.3
13	Human immunodeficiency virus (HIV) disease (B20-B24)	1,615	1.1	7.4
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,114	0.8	5.1
15	Chronic liver disease and cirrhosis (K70,K73-K74)	920	0.7	4.2
...	All other causes (Residual)	28,724	20.3	130.8
Black, female, 1-4 years				
...	All causes	446	100.0	33.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	97	21.7	7.2
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	70	15.7	5.2
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	63	14.1	4.7
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	22	4.9	1.6
5	Malignant neoplasms (C00-C97)	21	4.7	1.6
6	Cerebrovascular diseases (I60-I69)	10	2.2	*
6	Influenza and pneumonia (J09-J18)	10	2.2	*
6	Chronic lower respiratory diseases (J40-J47)	10	2.2	*
9	Septicemia (A40-A41)	7	1.6	*
9	Certain conditions originating in the perinatal period (P00-P96)	7	1.6	*
11	Anemias (D50-D64)	5	1.1	*
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	4	0.9	*
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3	0.7	*
14	Meningitis (G00,G03)	2	0.4	*
14	Complications of medical and surgical care (Y40-Y84,Y88)	2	0.4	*
...	All other causes (Residual)	113	25.3	8.4

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), race, sex, and age	Number	Percent of total deaths	Rate
Black, female, 5-14 years				
...	All causes	478	100.0	14.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	106	22.2	3.2
2	Malignant neoplasms (C00-C97)	81	16.9	2.5
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	32	6.7	1.0
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	28	5.9	0.9
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	22	4.6	0.7
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	21	4.4	0.6
7	Chronic lower respiratory diseases (J40-J47)	20	4.2	0.6
8	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	8	1.7	*
8	Anemias (D50-D64)	8	1.7	*
8	Cerebrovascular diseases (I60-I69)	8	1.7	*
11	Septicemia (A40-A41)	7	1.5	*
12	Diabetes mellitus (E10-E14)	4	0.8	*
12	Influenza and pneumonia (J09-J18)	4	0.8	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3	0.6	*
14	Certain conditions originating in the perinatal period (P00-P96)	3	0.6	*
14	Complications of medical and surgical care (Y40-Y84,Y88)	3	0.6	*
...	All other causes (Residual)	120	25.1	3.6
Black, female, 15-24 years				
...	All causes	1,546	100.0	43.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	386	25.0	10.8
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	268	17.3	7.5
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	125	8.1	3.5
4	Malignant neoplasms (C00-C97)	118	7.6	3.3
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	72	4.7	2.0
6	Pregnancy, childbirth and the puerperium (O00-O99)	51	3.3	1.4
7	Human immunodeficiency virus (HIV) disease (B20-B24)	40	2.6	1.1
8	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	36	2.3	1.0
9	Cerebrovascular diseases (I60-I69)	28	1.8	0.8
10	Diabetes mellitus (E10-E14)	25	1.6	0.7
11	Anemias (D50-D64)	24	1.6	0.7
12	Chronic lower respiratory diseases (J40-J47)	19	1.2	*
13	Septicemia (A40-A41)	15	1.0	*
13	Influenza and pneumonia (J09-J18)	15	1.0	*
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	15	1.0	*
...	All other causes (Residual)	309	20.0	8.7

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), race, sex, and age	Number	Percent of total deaths	Rate
Black, female, 25-44 years				
...	All causes	8,750	100.0	145.2
1	Malignant neoplasms (C00-C97)	1,649	18.8	27.4
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,389	15.9	23.0
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	923	10.5	15.3
4	Human immunodeficiency virus (HIV) disease (B20-B24)	620	7.1	10.3
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	446	5.1	7.4
6	Cerebrovascular diseases (I60-I69)	332	3.8	5.5
7	Diabetes mellitus (E10-E14)	306	3.5	5.1
8	Pregnancy, childbirth and the puerperium (O00-O99)	198	2.3	3.3
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	177	2.0	2.9
10	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	168	1.9	2.8
11	Septicemia (A40-A41)	158	1.8	2.6
12	Anemias (D50-D64)	123	1.4	2.0
13	Chronic lower respiratory diseases (J40-J47)	120	1.4	2.0
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	103	1.2	1.7
15	Chronic liver disease and cirrhosis (K70,K73-K74)	99	1.1	1.6
...	All other causes (Residual)	1,939	22.2	32.2
Black, female, 45-64 years				
...	All causes	36,058	100.0	686.4
1	Malignant neoplasms (C00-C97)	11,750	32.6	223.7
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	7,754	21.5	147.6
3	Cerebrovascular diseases (I60-I69)	1,943	5.4	37.0
4	Diabetes mellitus (E10-E14)	1,737	4.8	33.1
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,249	3.5	23.8
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,092	3.0	20.8
7	Chronic lower respiratory diseases (J40-J47)	1,030	2.9	19.6
8	Human immunodeficiency virus (HIV) disease (B20-B24)	857	2.4	16.3
9	Septicemia (A40-A41)	819	2.3	15.6
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	627	1.7	11.9
11	Chronic liver disease and cirrhosis (K70,K73-K74)	594	1.6	11.3
12	Influenza and pneumonia (J09-J18)	417	1.2	7.9
13	Viral hepatitis (B15-B19)	253	0.7	4.8
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	219	0.6	4.2
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	152	0.4	2.9
...	All other causes (Residual)	5,565	15.4	105.9

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), race, sex, and age	Number	Percent of total deaths	Rate
Black, female, 65 years and over				
...	All causes	90,589	100.0	4,213.5
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	24,648	27.2	1,146.4
2	Malignant neoplasms (C00-C97)	18,339	20.2	853.0
3	Cerebrovascular diseases (I60-I69)	6,677	7.4	310.6
4	Diabetes mellitus (E10-E14)	4,413	4.9	205.3
5	Alzheimer's disease (G30)	3,699	4.1	172.0
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3,518	3.9	163.6
7	Chronic lower respiratory diseases (J40-J47)	2,982	3.3	138.7
8	Septicemia (A40-A41)	2,261	2.5	105.2
9	Essential hypertension and hypertensive renal disease (I10,I12,I15)	2,164	2.4	100.7
10	Influenza and pneumonia (J09-J18)	1,986	2.2	92.4
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,082	1.2	50.3
12	Pneumonitis due to solids and liquids (J69)	549	0.6	25.5
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	451	0.5	21.0
14	Parkinson's disease (G20-G21)	336	0.4	15.6
15	Atherosclerosis (I70)	305	0.3	14.2
...	All other causes (Residual)	17,179	19.0	799.0
American Indian, both sexes, all ages				
...	All causes	15,565	100.0	365.1
1	Malignant neoplasms (C00-C97)	2,962	19.0	69.5
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,793	17.9	65.5
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,701	10.9	39.9
4	Diabetes mellitus (E10-E14)	857	5.5	20.1
5	Chronic liver disease and cirrhosis (K70,K73-K74)	787	5.1	18.5
6	Chronic lower respiratory diseases (J40-J47)	702	4.5	16.5
7	Cerebrovascular diseases (I60-I69)	559	3.6	13.1
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	469	3.0	11.0
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	339	2.2	8.0
10	Influenza and pneumonia (J09-J18)	326	2.1	7.6
11	Alzheimer's disease (G30)	264	1.7	6.2
12	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	257	1.7	6.0
13	Septicemia (A40-A41)	244	1.6	5.7
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	144	0.9	3.4
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	130	0.8	3.0
...	All other causes (Residual)	3,031	19.5	71.1

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, both sexes, 1-4 years				
...	All causes	93	100.0	29.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	31	33.3	9.8
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	10	10.8	*
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	9	9.7	*
4	Malignant neoplasms (C00-C97)	6	6.5	*
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	5	5.4	*
6	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2	2.2	*
6	Chronic lower respiratory diseases (J40-J47)	2	2.2	*
8	Septicemia (A40-A41)	1	1.1	*
8	Meningitis (G00,G03)	1	1.1	*
8	Cerebrovascular diseases (I60-I69)	1	1.1	*
8	Influenza and pneumonia (J09-J18)	1	1.1	*
8	Acute bronchitis and bronchiolitis (J20-J21)	1	1.1	*
8	Pneumonitis due to solids and liquids (J69)	1	1.1	*
8	Certain conditions originating in the perinatal period (P00-P96)	1	1.1	*
8	Legal intervention (Y35,Y89.0)	1	1.1	*
8	Complications of medical and surgical care (Y40-Y84,Y88)	1	1.1	*
...	All other causes (Residual)	19	20.4	*
American Indian, both sexes, 5-14 years				
...	All causes	108	100.0	14.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	47	43.5	6.3
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	14	13.0	*
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	10	9.3	*
4	Malignant neoplasms (C00-C97)	8	7.4	*
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	4	3.7	*
6	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	3	2.8	*
7	Diseases of heart (I00-I09,I11,I13,I20-I51)	2	1.9	*
7	Cerebrovascular diseases (I60-I69)	2	1.9	*
7	Influenza and pneumonia (J09-J18)	2	1.9	*
10	Septicemia (A40-A41)	1	0.9	*
10	Chronic lower respiratory diseases (J40-J47)	1	0.9	*
10	Complications of medical and surgical care (Y40-Y84,Y88)	1	0.9	*
...	All other causes (Residual)	13	12.0	*

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, both sexes, 15-24 years				
...	All causes	615	100.0	81.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	270	43.9	35.7
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	158	25.7	20.9
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	75	12.2	9.9
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	16	2.6	*
5	Malignant neoplasms (C00-C97)	14	2.3	*
6	Pregnancy, childbirth and the puerperium (O00-O99)	5	0.8	*
7	Chronic liver disease and cirrhosis (K70,K73-K74)	4	0.7	*
7	Legal intervention (Y35,Y89.0)	4	0.7	*
9	Cerebrovascular diseases (I60-I69)	3	0.5	*
9	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	3	0.5	*
11	Septicemia (A40-A41)	2	0.3	*
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	2	0.3	*
11	Influenza and pneumonia (J09-J18)	2	0.3	*
11	Chronic lower respiratory diseases (J40-J47)	2	0.3	*
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2	0.3	*
...	All other causes (Residual)	53	8.6	7.0
American Indian, both sexes, 25-44 years				
...	All causes	1,958	100.0	160.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	612	31.3	50.0
2	Chronic liver disease and cirrhosis (K70,K73-K74)	219	11.2	17.9
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	183	9.3	14.9
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	175	8.9	14.3
5	Malignant neoplasms (C00-C97)	124	6.3	10.1
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	121	6.2	9.9
7	Diabetes mellitus (E10-E14)	49	2.5	4.0
8	Influenza and pneumonia (J09-J18)	26	1.3	2.1
9	Human immunodeficiency virus (HIV) disease (B20-B24)	21	1.1	1.7
9	Cerebrovascular diseases (I60-I69)	21	1.1	1.7
11	Septicemia (A40-A41)	19	1.0	*
12	Chronic lower respiratory diseases (J40-J47)	14	0.7	*
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	12	0.6	*
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	11	0.6	*
15	Viral hepatitis (B15-B19)	10	0.5	*
...	All other causes (Residual)	341	17.4	27.9

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, both sexes, 45-64 years				
...	All causes	4,907	100.0	555.4
1	Malignant neoplasms (C00-C97)	1,122	22.9	127.0
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	884	18.0	100.1
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	486	9.9	55.0
4	Chronic liver disease and cirrhosis (K70,K73-K74)	445	9.1	50.4
5	Diabetes mellitus (E10-E14)	324	6.6	36.7
6	Chronic lower respiratory diseases (J40-J47)	164	3.3	18.6
7	Cerebrovascular diseases (I60-I69)	137	2.8	15.5
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	105	2.1	11.9
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	93	1.9	10.5
10	Septicemia (A40-A41)	81	1.7	9.2
11	Viral hepatitis (B15-B19)	76	1.5	8.6
11	Influenza and pneumonia (J09-J18)	76	1.5	8.6
13	Human immunodeficiency virus (HIV) disease (B20-B24)	39	0.8	4.4
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	38	0.8	4.3
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	36	0.7	4.1
...	All other causes (Residual)	801	16.3	90.7
American Indian, both sexes, 65 years and over				
...	All causes	7,530	100.0	2,950.5
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,697	22.5	664.9
2	Malignant neoplasms (C00-C97)	1,686	22.4	660.6
3	Chronic lower respiratory diseases (J40-J47)	515	6.8	201.8
4	Diabetes mellitus (E10-E14)	483	6.4	189.3
5	Cerebrovascular diseases (I60-I69)	393	5.2	154.0
6	Alzheimer's disease (G30)	261	3.5	102.3
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	230	3.1	90.1
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	229	3.0	89.7
9	Influenza and pneumonia (J09-J18)	210	2.8	82.3
10	Septicemia (A40-A41)	137	1.8	53.7
11	Chronic liver disease and cirrhosis (K70,K73-K74)	119	1.6	46.6
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	98	1.3	38.4
13	Parkinson's disease (G20-G21)	60	0.8	23.5
14	Pneumonitis due to solids and liquids (J69)	56	0.7	21.9
15	Aortic aneurysm and dissection (I71)	39	0.5	15.3
...	All other causes (Residual)	1,317	17.5	516.0

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, male, all ages				
...	All causes	8,516	100.0	397.5
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,608	18.9	75.0
2	Malignant neoplasms (C00-C97)	1,588	18.6	74.1
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,150	13.5	53.7
4	Diabetes mellitus (E10-E14)	432	5.1	20.2
5	Chronic liver disease and cirrhosis (K70,K73-K74)	429	5.0	20.0
6	Chronic lower respiratory diseases (J40-J47)	349	4.1	16.3
7	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	344	4.0	16.1
8	Cerebrovascular diseases (I60-I69)	258	3.0	12.0
9	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	204	2.4	9.5
10	Influenza and pneumonia (J09-J18)	172	2.0	8.0
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	153	1.8	7.1
12	Septicemia (A40-A41)	106	1.2	4.9
13	Alzheimer's disease (G30)	89	1.0	4.2
14	Certain conditions originating in the perinatal period (P00-P96)	76	0.9	3.5
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	71	0.8	3.3
...	All other causes (Residual)	1,487	17.5	69.4
American Indian, male, 1-4 years				
...	All causes	55	100.0	34.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	19	34.5	*
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	7	12.7	*
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	6	10.9	*
4	Malignant neoplasms (C00-C97)	4	7.3	*
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	3	5.5	*
6	Septicemia (A40-A41)	1	1.8	*
6	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1	1.8	*
6	Chronic lower respiratory diseases (J40-J47)	1	1.8	*
6	Certain conditions originating in the perinatal period (P00-P96)	1	1.8	*
6	Complications of medical and surgical care (Y40-Y84,Y88)	1	1.8	*
...	All other causes (Residual)	11	20.0	*

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, male, 5-14 years				
...	All causes	69	100.0	18.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	35	50.7	9.2
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	8	11.6	*
3	Malignant neoplasms (C00-C97)	4	5.8	*
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	4	5.8	*
5	Cerebrovascular diseases (I60-I69)	2	2.9	*
6	Septicemia (A40-A41)	1	1.4	*
6	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1	1.4	*
6	Influenza and pneumonia (J09-J18)	1	1.4	*
6	Chronic lower respiratory diseases (J40-J47)	1	1.4	*
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1	1.4	*
...	All other causes (Residual)	11	15.9	*
American Indian, male, 15-24 years				
...	All causes	456	100.0	116.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	201	44.1	51.3
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	120	26.3	30.6
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	69	15.1	17.6
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	11	2.4	*
5	Malignant neoplasms (C00-C97)	8	1.8	*
6	Legal intervention (Y35,Y89.0)	4	0.9	*
7	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	3	0.7	*
8	Cerebrovascular diseases (I60-I69)	2	0.4	*
8	Chronic liver disease and cirrhosis (K70,K73-K74)	2	0.4	*
10	Septicemia (A40-A41)	1	0.2	*
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1	0.2	*
10	Diabetes mellitus (E10-E14)	1	0.2	*
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1	0.2	*
10	Influenza and pneumonia (J09-J18)	1	0.2	*
10	Chronic lower respiratory diseases (J40-J47)	1	0.2	*
10	Pneumonitis due to solids and liquids (J69)	1	0.2	*
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1	0.2	*
10	Complications of medical and surgical care (Y40-Y84,Y88)	1	0.2	*
...	All other causes (Residual)	27	5.9	6.9

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, male, 25-44 years				
...	All causes	1,274	100.0	203.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	404	31.7	64.5
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	134	10.5	21.4
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	131	10.3	20.9
4	Chronic liver disease and cirrhosis (K70,K73-K74)	123	9.7	19.6
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	93	7.3	14.8
6	Malignant neoplasms (C00-C97)	64	5.0	10.2
7	Diabetes mellitus (E10-E14)	27	2.1	4.3
8	Influenza and pneumonia (J09-J18)	18	1.4	*
9	Human immunodeficiency virus (HIV) disease (B20-B24)	17	1.3	*
10	Cerebrovascular diseases (I60-I69)	12	0.9	*
11	Septicemia (A40-A41)	11	0.9	*
12	Chronic lower respiratory diseases (J40-J47)	10	0.8	*
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	9	0.7	*
13	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	9	0.7	*
15	Viral hepatitis (B15-B19)	8	0.6	*
...	All other causes (Residual)	204	16.0	32.6
American Indian, male, 45-64 years				
...	All causes	2,901	100.0	672.4
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	622	21.4	144.2
2	Malignant neoplasms (C00-C97)	602	20.8	139.5
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	341	11.8	79.0
4	Chronic liver disease and cirrhosis (K70,K73-K74)	244	8.4	56.6
5	Diabetes mellitus (E10-E14)	192	6.6	44.5
6	Chronic lower respiratory diseases (J40-J47)	84	2.9	19.5
7	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	77	2.7	17.8
8	Cerebrovascular diseases (I60-I69)	68	2.3	15.8
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	53	1.8	12.3
10	Influenza and pneumonia (J09-J18)	51	1.8	11.8
11	Viral hepatitis (B15-B19)	47	1.6	10.9
12	Septicemia (A40-A41)	34	1.2	7.9
13	Human immunodeficiency virus (HIV) disease (B20-B24)	30	1.0	7.0
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	28	1.0	6.5
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	22	0.8	5.1
...	All other causes (Residual)	406	14.0	94.1

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, male, 65 years and over				
...	All causes	3,548	100.0	3,143.9
1	Malignant neoplasms (C00-C97)	905	25.5	801.9
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	834	23.5	739.0
3	Chronic lower respiratory diseases (J40-J47)	248	7.0	219.8
4	Diabetes mellitus (E10-E14)	212	6.0	187.9
5	Cerebrovascular diseases (I60-I69)	172	4.8	152.4
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	134	3.8	118.7
7	Influenza and pneumonia (J09-J18)	96	2.7	85.1
8	Alzheimer's disease (G30)	89	2.5	78.9
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	88	2.5	78.0
10	Chronic liver disease and cirrhosis (K70,K73-K74)	60	1.7	53.2
11	Septicemia (A40-A41)	56	1.6	49.6
12	Parkinson's disease (G20-G21)	38	1.1	33.7
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	36	1.0	31.9
14	Pneumonitis due to solids and liquids (J69)	25	0.7	22.2
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	20	0.6	17.7
...	All other causes (Residual)	535	15.1	474.1
American Indian, female, all ages				
...	All causes	7,049	100.0	332.4
1	Malignant neoplasms (C00-C97)	1,374	19.5	64.8
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,185	16.8	55.9
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	551	7.8	26.0
4	Diabetes mellitus (E10-E14)	425	6.0	20.0
5	Chronic liver disease and cirrhosis (K70,K73-K74)	358	5.1	16.9
6	Chronic lower respiratory diseases (J40-J47)	353	5.0	16.6
7	Cerebrovascular diseases (I60-I69)	301	4.3	14.2
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	186	2.6	8.8
9	Alzheimer's disease (G30)	175	2.5	8.3
10	Influenza and pneumonia (J09-J18)	154	2.2	7.3
11	Septicemia (A40-A41)	138	2.0	6.5
12	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	125	1.8	5.9
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	82	1.2	3.9
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	59	0.8	2.8
15	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	53	0.8	2.5
...	All other causes (Residual)	1,530	21.7	72.1

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, female, 1-4 years				
...	All causes	38	100.0	24.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	12	31.6	*
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	3	7.9	*
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	3	7.9	*
4	Malignant neoplasms (C00-C97)	2	5.3	*
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	2	5.3	*
6	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1	2.6	*
6	Meningitis (G00,G03)	1	2.6	*
6	Cerebrovascular diseases (I60-I69)	1	2.6	*
6	Influenza and pneumonia (J09-J18)	1	2.6	*
6	Acute bronchitis and bronchiolitis (J20-J21)	1	2.6	*
6	Chronic lower respiratory diseases (J40-J47)	1	2.6	*
6	Pneumonitis due to solids and liquids (J69)	1	2.6	*
6	Legal intervention (Y35,Y89.0)	1	2.6	*
...	All other causes (Residual)	8	21.1	*
American Indian, female, 5-14 years				
...	All causes	39	100.0	10.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	12	30.8	*
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	6	15.4	*
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	6	15.4	*
4	Malignant neoplasms (C00-C97)	4	10.3	*
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	3	7.7	*
6	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2	5.1	*
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	2	5.1	*
8	Influenza and pneumonia (J09-J18)	1	2.6	*
8	Complications of medical and surgical care (Y40-Y84,Y88)	1	2.6	*
...	All other causes (Residual)	2	5.1	*

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, female, 15-24 years				
...	All causes	159	100.0	43.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	69	43.4	18.9
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	38	23.9	10.4
3	Malignant neoplasms (C00-C97)	6	3.8	*
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	6	3.8	*
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	5	3.1	*
5	Pregnancy, childbirth and the puerperium (O00-O99)	5	3.1	*
7	Chronic liver disease and cirrhosis (K70,K73-K74)	2	1.3	*
8	Septicemia (A40-A41)	1	0.6	*
8	Anemias (D50-D64)	1	0.6	*
8	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1	0.6	*
8	Cerebrovascular diseases (I60-I69)	1	0.6	*
8	Influenza and pneumonia (J09-J18)	1	0.6	*
8	Chronic lower respiratory diseases (J40-J47)	1	0.6	*
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1	0.6	*
...	All other causes (Residual)	21	13.2	5.8
American Indian, female, 25-44 years				
...	All causes	684	100.0	114.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	208	30.4	34.8
2	Chronic liver disease and cirrhosis (K70,K73-K74)	96	14.0	16.1
3	Malignant neoplasms (C00-C97)	60	8.8	10.0
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	49	7.2	8.2
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	44	6.4	7.4
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	28	4.1	4.7
7	Diabetes mellitus (E10-E14)	22	3.2	3.7
8	Cerebrovascular diseases (I60-I69)	9	1.3	*
9	Septicemia (A40-A41)	8	1.2	*
9	Influenza and pneumonia (J09-J18)	8	1.2	*
9	Pregnancy, childbirth and the puerperium (O00-O99)	8	1.2	*
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	5	0.7	*
13	Human immunodeficiency virus (HIV) disease (B20-B24)	4	0.6	*
13	Chronic lower respiratory diseases (J40-J47)	4	0.6	*
15	Anemias (D50-D64)	3	0.4	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3	0.4	*
...	All other causes (Residual)	125	18.3	20.9

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, female, 45-64 years				
...	All causes	2,006	100.0	443.8
1	Malignant neoplasms (C00-C97)	520	25.9	115.0
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	262	13.1	58.0
3	Chronic liver disease and cirrhosis (K70,K73-K74)	201	10.0	44.5
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	145	7.2	32.1
5	Diabetes mellitus (E10-E14)	132	6.6	29.2
6	Chronic lower respiratory diseases (J40-J47)	80	4.0	17.7
7	Cerebrovascular diseases (I60-I69)	69	3.4	15.3
8	Septicemia (A40-A41)	47	2.3	10.4
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	40	2.0	8.8
10	Viral hepatitis (B15-B19)	29	1.4	6.4
11	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	28	1.4	6.2
12	Influenza and pneumonia (J09-J18)	25	1.2	5.5
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	14	0.7	*
14	Pneumonitis due to solids and liquids (J69)	10	0.5	*
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	10	0.5	*
...	All other causes (Residual)	394	19.6	87.2
American Indian, female, 65 years and over				
...	All causes	3,982	100.0	2,797.1
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	863	21.7	606.2
2	Malignant neoplasms (C00-C97)	781	19.6	548.6
3	Diabetes mellitus (E10-E14)	271	6.8	190.4
4	Chronic lower respiratory diseases (J40-J47)	267	6.7	187.6
5	Cerebrovascular diseases (I60-I69)	221	5.5	155.2
6	Alzheimer's disease (G30)	172	4.3	120.8
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	142	3.6	99.7
8	Influenza and pneumonia (J09-J18)	114	2.9	80.1
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	95	2.4	66.7
10	Septicemia (A40-A41)	81	2.0	56.9
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	62	1.6	43.6
12	Chronic liver disease and cirrhosis (K70,K73-K74)	59	1.5	41.4
13	Pneumonitis due to solids and liquids (J69)	31	0.8	21.8
14	Parkinson's disease (G20-G21)	22	0.6	15.5
15	Aortic aneurysm and dissection (I71)	20	0.5	14.0
...	All other causes (Residual)	781	19.6	548.6

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, both sexes, all ages				
...	All causes	51,162	100.0	301.1
1	Malignant neoplasms (C00-C97)	14,165	27.7	83.4
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	11,490	22.5	67.6
3	Cerebrovascular diseases (I60-I69)	3,833	7.5	22.6
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,144	4.2	12.6
5	Diabetes mellitus (E10-E14)	1,838	3.6	10.8
6	Influenza and pneumonia (J09-J18)	1,539	3.0	9.1
7	Chronic lower respiratory diseases (J40-J47)	1,487	2.9	8.8
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,091	2.1	6.4
9	Alzheimer's disease (G30)	1,082	2.1	6.4
10	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,061	2.1	6.2
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	814	1.6	4.8
12	Septicemia (A40-A41)	582	1.1	3.4
13	Chronic liver disease and cirrhosis (K70,K73-K74)	467	0.9	2.7
13	Certain conditions originating in the perinatal period (P00-P96)	467	0.9	2.7
15	Parkinson's disease (G20-G21)	374	0.7	2.2
...	All other causes (Residual)	8,728	17.1	51.4
Asian or Pacific Islander, both sexes, 1-4 years				
...	All causes	167	100.0	17.9
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	42	25.1	4.5
2	Malignant neoplasms (C00-C97)	23	13.8	2.5
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	23	13.8	2.5
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	8	4.8	*
5	Chronic lower respiratory diseases (J40-J47)	5	3.0	*
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	5	3.0	*
7	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	4	2.4	*
7	Cerebrovascular diseases (I60-I69)	4	2.4	*
7	Influenza and pneumonia (J09-J18)	4	2.4	*
10	Septicemia (A40-A41)	2	1.2	*
11	Anemias (D50-D64)	1	0.6	*
11	Meningitis (G00,G03)	1	0.6	*
11	Acute bronchitis and bronchiolitis (J20-J21)	1	0.6	*
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1	0.6	*
11	Certain conditions originating in the perinatal period (P00-P96)	1	0.6	*
...	All other causes (Residual)	42	25.1	4.5

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, both sexes, 5-14 years				
...	All causes	185	100.0	8.2
1	Malignant neoplasms (C00-C97)	55	29.7	2.4
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	29	15.7	1.3
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	11	5.9	*
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	8	4.3	*
5	Septicemia (A40-A41)	7	3.8	*
5	Chronic lower respiratory diseases (J40-J47)	7	3.8	*
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	7	3.8	*
8	Diseases of heart (I00-I09,I11,I13,I20-I51)	6	3.2	*
9	Cerebrovascular diseases (I60-I69)	5	2.7	*
10	Influenza and pneumonia (J09-J18)	3	1.6	*
10	Complications of medical and surgical care (Y40-Y84,Y88)	3	1.6	*
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2	1.1	*
12	Anemias (D50-D64)	2	1.1	*
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2	1.1	*
12	Certain conditions originating in the perinatal period (P00-P96)	2	1.1	*
...	All other causes (Residual)	36	19.5	1.6
Asian or Pacific Islander, both sexes, 15-24 years				
...	All causes	752	100.0	30.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	256	34.0	10.3
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	181	24.1	7.3
3	Malignant neoplasms (C00-C97)	80	10.6	3.2
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	63	8.4	2.5
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	37	4.9	1.5
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	14	1.9	*
7	Cerebrovascular diseases (I60-I69)	9	1.2	*
8	Influenza and pneumonia (J09-J18)	6	0.8	*
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	5	0.7	*
10	Septicemia (A40-A41)	3	0.4	*
10	Anemias (D50-D64)	3	0.4	*
10	Chronic lower respiratory diseases (J40-J47)	3	0.4	*
13	Tuberculosis (A16-A19)	2	0.3	*
13	Human immunodeficiency virus (HIV) disease (B20-B24)	2	0.3	*
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2	0.3	*
13	Complications of medical and surgical care (Y40-Y84,Y88)	2	0.3	*
...	All other causes (Residual)	84	11.2	3.4

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, both sexes, 25-44 years				
...	All causes	2,913	100.0	52.0
1	Malignant neoplasms (C00-C97)	746	25.6	13.3
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	456	15.7	8.1
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	403	13.8	7.2
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	349	12.0	6.2
5	Cerebrovascular diseases (I60-I69)	138	4.7	2.5
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	125	4.3	2.2
7	Chronic liver disease and cirrhosis (K70,K73-K74)	55	1.9	1.0
8	Diabetes mellitus (E10-E14)	42	1.4	0.7
9	Pregnancy, childbirth and the puerperium (O00-O99)	40	1.4	0.7
10	Influenza and pneumonia (J09-J18)	32	1.1	0.6
11	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	30	1.0	0.5
12	Septicemia (A40-A41)	26	0.9	0.5
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	24	0.8	0.4
14	Aortic aneurysm and dissection (I71)	23	0.8	0.4
15	Human immunodeficiency virus (HIV) disease (B20-B24)	18	0.6	*
15	Chronic lower respiratory diseases (J40-J47)	18	0.6	*
...	All other causes (Residual)	388	13.3	6.9
Asian or Pacific Islander, both sexes, 45-64 years				
...	All causes	10,561	100.0	265.4
1	Malignant neoplasms (C00-C97)	4,464	42.3	112.2
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,940	18.4	48.8
3	Cerebrovascular diseases (I60-I69)	699	6.6	17.6
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	483	4.6	12.1
5	Diabetes mellitus (E10-E14)	399	3.8	10.0
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	336	3.2	8.4
7	Chronic liver disease and cirrhosis (K70,K73-K74)	214	2.0	5.4
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	164	1.6	4.1
9	Viral hepatitis (B15-B19)	143	1.4	3.6
10	Influenza and pneumonia (J09-J18)	119	1.1	3.0
11	Chronic lower respiratory diseases (J40-J47)	114	1.1	2.9
12	Septicemia (A40-A41)	102	1.0	2.6
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	98	0.9	2.5
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	87	0.8	2.2
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	52	0.5	1.3
...	All other causes (Residual)	1,147	10.9	28.8

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, both sexes, 65 years and over				
...	All causes	35,699	100.0	2,380.5
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	9,133	25.6	609.0
2	Malignant neoplasms (C00-C97)	8,792	24.6	586.3
3	Cerebrovascular diseases (I60-I69)	2,972	8.3	198.2
4	Diabetes mellitus (E10-E14)	1,397	3.9	93.2
5	Influenza and pneumonia (J09-J18)	1,363	3.8	90.9
6	Chronic lower respiratory diseases (J40-J47)	1,337	3.7	89.2
7	Alzheimer's disease (G30)	1,072	3.0	71.5
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	891	2.5	59.4
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	850	2.4	56.7
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	706	2.0	47.1
11	Septicemia (A40-A41)	434	1.2	28.9
12	Parkinson's disease (G20-G21)	358	1.0	23.9
13	Pneumonitis due to solids and liquids (J69)	292	0.8	19.5
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	271	0.8	18.1
15	Aortic aneurysm and dissection (I71)	203	0.6	13.5
...	All other causes (Residual)	5,628	15.8	375.3
Asian or Pacific Islander, male, all ages				
...	All causes	26,600	100.0	327.0
1	Malignant neoplasms (C00-C97)	7,210	27.1	88.6
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	6,262	23.5	77.0
3	Cerebrovascular diseases (I60-I69)	1,747	6.6	21.5
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,337	5.0	16.4
5	Diabetes mellitus (E10-E14)	932	3.5	11.5
6	Chronic lower respiratory diseases (J40-J47)	910	3.4	11.2
7	Influenza and pneumonia (J09-J18)	825	3.1	10.1
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	756	2.8	9.3
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	524	2.0	6.4
10	Alzheimer's disease (G30)	345	1.3	4.2
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	337	1.3	4.1
12	Chronic liver disease and cirrhosis (K70,K73-K74)	302	1.1	3.7
13	Septicemia (A40-A41)	296	1.1	3.6
14	Certain conditions originating in the perinatal period (P00-P96)	278	1.0	3.4
15	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	218	0.8	2.7
...	All other causes (Residual)	4,321	16.2	53.1

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, male, 1-4 years				
...	All causes	92	100.0	19.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	26	28.3	5.5
2	Malignant neoplasms (C00-C97)	17	18.5	*
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	8	8.7	*
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	5	5.4	*
5	Cerebrovascular diseases (I60-I69)	3	3.3	*
6	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2	2.2	*
6	Influenza and pneumonia (J09-J18)	2	2.2	*
6	Chronic lower respiratory diseases (J40-J47)	2	2.2	*
9	Septicemia (A40-A41)	1	1.1	*
9	Acute bronchitis and bronchiolitis (J20-J21)	1	1.1	*
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1	1.1	*
9	Certain conditions originating in the perinatal period (P00-P96)	1	1.1	*
9	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1	1.1	*
...	All other causes (Residual)	22	23.9	4.6
Asian or Pacific Islander, male, 5-14 years				
...	All causes	96	100.0	8.4
1	Malignant neoplasms (C00-C97)	27	28.1	2.4
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	19	19.8	*
3	Chronic lower respiratory diseases (J40-J47)	6	6.3	*
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	6	6.3	*
5	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	5	5.2	*
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	5	5.2	*
7	Septicemia (A40-A41)	3	3.1	*
7	Diseases of heart (I00-I09,I11,I13,I20-I51)	3	3.1	*
9	Cerebrovascular diseases (I60-I69)	2	2.1	*
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2	2.1	*
11	Anemias (D50-D64)	1	1.0	*
11	Influenza and pneumonia (J09-J18)	1	1.0	*
...	All other causes (Residual)	16	16.7	*

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, male, 15-24 years				
...	All causes	544	100.0	43.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	196	36.0	15.5
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	138	25.4	10.9
3	Malignant neoplasms (C00-C97)	52	9.6	4.1
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	51	9.4	4.0
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	26	4.8	2.1
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	9	1.7	*
7	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	4	0.7	*
7	Cerebrovascular diseases (I60-I69)	4	0.7	*
7	Influenza and pneumonia (J09-J18)	4	0.7	*
10	Septicemia (A40-A41)	3	0.6	*
10	Chronic lower respiratory diseases (J40-J47)	3	0.6	*
12	Human immunodeficiency virus (HIV) disease (B20-B24)	2	0.4	*
13	Tuberculosis (A16-A19)	1	0.2	*
13	Anemias (D50-D64)	1	0.2	*
13	Chronic liver disease and cirrhosis (K70,K73-K74)	1	0.2	*
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1	0.2	*
13	Legal intervention (Y35,Y89.0)	1	0.2	*
13	Complications of medical and surgical care (Y40-Y84,Y88)	1	0.2	*
...	All other causes (Residual)	46	8.5	3.6
Asian or Pacific Islander, male, 25-44 years				
...	All causes	1,797	100.0	67.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	344	19.1	13.0
2	Malignant neoplasms (C00-C97)	324	18.0	12.2
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	281	15.6	10.6
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	278	15.5	10.5
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	88	4.9	3.3
6	Cerebrovascular diseases (I60-I69)	79	4.4	3.0
7	Chronic liver disease and cirrhosis (K70,K73-K74)	40	2.2	1.5
8	Diabetes mellitus (E10-E14)	22	1.2	0.8
9	Influenza and pneumonia (J09-J18)	21	1.2	0.8
10	Aortic aneurysm and dissection (I71)	16	0.9	*
10	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	16	0.9	*
12	Human immunodeficiency virus (HIV) disease (B20-B24)	15	0.8	*
13	Viral hepatitis (B15-B19)	12	0.7	*
14	Septicemia (A40-A41)	11	0.6	*
15	Chronic lower respiratory diseases (J40-J47)	9	0.5	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	9	0.5	*
...	All other causes (Residual)	232	12.9	8.7

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, male, 45-64 years				
...	All causes	6,248	100.0	340.3
1	Malignant neoplasms (C00-C97)	2,210	35.4	120.4
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,437	23.0	78.3
3	Cerebrovascular diseases (I60-I69)	399	6.4	21.7
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	302	4.8	16.4
5	Diabetes mellitus (E10-E14)	272	4.4	14.8
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	235	3.8	12.8
7	Chronic liver disease and cirrhosis (K70,K73-K74)	162	2.6	8.8
8	Viral hepatitis (B15-B19)	105	1.7	5.7
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	86	1.4	4.7
10	Influenza and pneumonia (J09-J18)	78	1.2	4.2
11	Chronic lower respiratory diseases (J40-J47)	74	1.2	4.0
12	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	57	0.9	3.1
13	Septicemia (A40-A41)	55	0.9	3.0
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	55	0.9	3.0
15	Human immunodeficiency virus (HIV) disease (B20-B24)	34	0.5	1.9
...	All other causes (Residual)	687	11.0	37.4
Asian or Pacific Islander, male, 65 years and over				
...	All causes	17,318	100.0	2,668.4
1	Malignant neoplasms (C00-C97)	4,578	26.4	705.4
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	4,506	26.0	694.3
3	Cerebrovascular diseases (I60-I69)	1,256	7.3	193.5
4	Chronic lower respiratory diseases (J40-J47)	813	4.7	125.3
5	Influenza and pneumonia (J09-J18)	712	4.1	109.7
6	Diabetes mellitus (E10-E14)	638	3.7	98.3
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	433	2.5	66.7
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	420	2.4	64.7
9	Alzheimer's disease (G30)	341	2.0	52.5
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	274	1.6	42.2
11	Septicemia (A40-A41)	222	1.3	34.2
12	Parkinson's disease (G20-G21)	193	1.1	29.7
13	Pneumonitis due to solids and liquids (J69)	168	1.0	25.9
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	133	0.8	20.5
15	Aortic aneurysm and dissection (I71)	110	0.6	16.9
...	All other causes (Residual)	2,521	14.6	388.4

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, female, all ages				
...	All causes	24,562	100.0	277.3
1	Malignant neoplasms (C00-C97)	6,955	28.3	78.5
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	5,228	21.3	59.0
3	Cerebrovascular diseases (I60-I69)	2,086	8.5	23.5
4	Diabetes mellitus (E10-E14)	906	3.7	10.2
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	807	3.3	9.1
6	Alzheimer's disease (G30)	737	3.0	8.3
7	Influenza and pneumonia (J09-J18)	714	2.9	8.1
8	Chronic lower respiratory diseases (J40-J47)	577	2.3	6.5
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	567	2.3	6.4
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	477	1.9	5.4
11	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	305	1.2	3.4
12	Septicemia (A40-A41)	286	1.2	3.2
13	Certain conditions originating in the perinatal period (P00-P96)	189	0.8	2.1
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	178	0.7	2.0
15	Parkinson's disease (G20-G21)	168	0.7	1.9
...	All other causes (Residual)	4,382	17.8	49.5
Asian or Pacific Islander, female, 1-4 years				
...	All causes	75	100.0	16.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	16	21.3	*
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	15	20.0	*
3	Malignant neoplasms (C00-C97)	6	8.0	*
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	4	5.3	*
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	3	4.0	*
5	Chronic lower respiratory diseases (J40-J47)	3	4.0	*
7	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2	2.7	*
7	Influenza and pneumonia (J09-J18)	2	2.7	*
9	Septicemia (A40-A41)	1	1.3	*
9	Anemias (D50-D64)	1	1.3	*
9	Meningitis (G00,G03)	1	1.3	*
9	Cerebrovascular diseases (I60-I69)	1	1.3	*
...	All other causes (Residual)	20	26.7	4.3

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, female, 5-14 years				
...	All causes	89	100.0	7.9
1	Malignant neoplasms (C00-C97)	28	31.5	2.5
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	10	11.2	*
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	6	6.7	*
4	Septicemia (A40-A41)	4	4.5	*
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	3	3.4	*
5	Cerebrovascular diseases (I60-I69)	3	3.4	*
5	Complications of medical and surgical care (Y40-Y84,Y88)	3	3.4	*
8	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2	2.2	*
8	Influenza and pneumonia (J09-J18)	2	2.2	*
8	Certain conditions originating in the perinatal period (P00-P96)	2	2.2	*
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2	2.2	*
8	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	2	2.2	*
13	Meningococcal infection (A39)	1	1.1	*
13	Anemias (D50-D64)	1	1.1	*
13	Chronic lower respiratory diseases (J40-J47)	1	1.1	*
...	All other causes (Residual)	19	21.3	*
Asian or Pacific Islander, female, 15-24 years				
...	All causes	208	100.0	17.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	60	28.8	4.9
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	43	20.7	3.5
3	Malignant neoplasms (C00-C97)	28	13.5	2.3
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	12	5.8	*
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	11	5.3	*
6	Cerebrovascular diseases (I60-I69)	5	2.4	*
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	5	2.4	*
8	Anemias (D50-D64)	2	1.0	*
8	Influenza and pneumonia (J09-J18)	2	1.0	*
10	Tuberculosis (A16-A19)	1	0.5	*
10	Scarlet fever and erysipelas (A38,A46)	1	0.5	*
10	Meningococcal infection (A39)	1	0.5	*
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1	0.5	*
10	Aortic aneurysm and dissection (I71)	1	0.5	*
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1	0.5	*
10	Pregnancy, childbirth and the puerperium (O00-O99)	1	0.5	*
10	Complications of medical and surgical care (Y40-Y84,Y88)	1	0.5	*
...	All other causes (Residual)	32	15.4	2.6

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, female, 25-44 years				
...	All causes	1,116	100.0	37.9
1	Malignant neoplasms (C00-C97)	422	37.8	14.3
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	122	10.9	4.1
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	112	10.0	3.8
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	71	6.4	2.4
5	Cerebrovascular diseases (I60-I69)	59	5.3	2.0
6	Pregnancy, childbirth and the puerperium (O00-O99)	40	3.6	1.4
7	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	37	3.3	1.3
8	Diabetes mellitus (E10-E14)	20	1.8	0.7
9	Septicemia (A40-A41)	15	1.3	*
9	Chronic liver disease and cirrhosis (K70,K73-K74)	15	1.3	*
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	15	1.3	*
12	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	14	1.3	*
13	Influenza and pneumonia (J09-J18)	11	1.0	*
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	9	0.8	*
14	Chronic lower respiratory diseases (J40-J47)	9	0.8	*
...	All other causes (Residual)	145	13.0	4.9
Asian or Pacific Islander, female, 45-64 years				
...	All causes	4,313	100.0	201.2
1	Malignant neoplasms (C00-C97)	2,254	52.3	105.2
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	503	11.7	23.5
3	Cerebrovascular diseases (I60-I69)	300	7.0	14.0
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	181	4.2	8.4
5	Diabetes mellitus (E10-E14)	127	2.9	5.9
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	101	2.3	4.7
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	78	1.8	3.6
8	Chronic liver disease and cirrhosis (K70,K73-K74)	52	1.2	2.4
9	Septicemia (A40-A41)	47	1.1	2.2
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	43	1.0	2.0
11	Influenza and pneumonia (J09-J18)	41	1.0	1.9
12	Chronic lower respiratory diseases (J40-J47)	40	0.9	1.9
13	Viral hepatitis (B15-B19)	38	0.9	1.8
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	30	0.7	1.4
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	26	0.6	1.2
...	All other causes (Residual)	452	10.5	21.1

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), race, sex, and age	Number	Percent of total deaths	Rate
	Asian or Pacific Islander, female, 65 years and over			
...	All causes	18,381	100.0	2,160.8
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	4,627	25.2	543.9
2	Malignant neoplasms (C00-C97)	4,214	22.9	495.4
3	Cerebrovascular diseases (I60-I69)	1,716	9.3	201.7
4	Diabetes mellitus (E10-E14)	759	4.1	89.2
5	Alzheimer's disease (G30)	731	4.0	85.9
6	Influenza and pneumonia (J09-J18)	651	3.5	76.5
7	Chronic lower respiratory diseases (J40-J47)	524	2.9	61.6
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	471	2.6	55.4
9	Essential hypertension and hypertensive renal disease (I10,I12,I15)	432	2.4	50.8
10	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	417	2.3	49.0
11	Septicemia (A40-A41)	212	1.2	24.9
12	Parkinson's disease (G20-G21)	165	0.9	19.4
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	138	0.8	16.2
14	Pneumonitis due to solids and liquids (J69)	124	0.7	14.6
15	Chronic liver disease and cirrhosis (K70,K73-K74)	98	0.5	11.5
...	All other causes (Residual)	3,102	16.9	364.7

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable