

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2005

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
White, non-Hispanic, female, 5-9 years				
...	All causes	701	100.0	12.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	242	34.5	4.3
2	Malignant neoplasms (C00-C97)	133	19.0	2.4
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	60	8.6	1.1
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	25	3.6	0.4
5	Influenza and pneumonia (J10-J18)	18	2.6	*
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	16	2.3	*
7	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	13	1.9	*
7	Cerebrovascular diseases (I60-I69)	13	1.9	*
9	Chronic lower respiratory diseases (J40-J47)	8	1.1	*
10	Septicemia (A40-A41)	7	1.0	*
11	Meningitis (G00,G03)	6	0.9	*
12	Certain conditions originating in the perinatal period (P00-P96)	4	0.6	*
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3	0.4	*
14	Anemias (D50-D64)	2	0.3	*
14	Diabetes mellitus (E10-E14)	2	0.3	*
14	Nutritional deficiencies (E40-E64)	2	0.3	*
14	Complications of medical and surgical care (Y40-Y84,Y88)	2	0.3	*
...	All other causes (Residual)	145	20.7	2.6
White, non-Hispanic, female, 10-14 years				
...	All causes	817	100.0	13.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	295	36.1	4.8
2	Malignant neoplasms (C00-C97)	131	16.0	2.1
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	46	5.6	0.8
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	37	4.5	0.6
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	34	4.2	0.6
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	32	3.9	0.5
7	Influenza and pneumonia (J10-J18)	20	2.4	0.3
8	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	12	1.5	*
9	Septicemia (A40-A41)	11	1.3	*
9	Diabetes mellitus (E10-E14)	11	1.3	*
11	Cerebrovascular diseases (I60-I69)	6	0.7	*
11	Chronic lower respiratory diseases (J40-J47)	6	0.7	*
11	Certain conditions originating in the perinatal period (P00-P96)	6	0.7	*
14	Anemias (D50-D64)	4	0.5	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	4	0.5	*
...	All other causes (Residual)	162	19.8	2.6

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2005

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
White, non-Hispanic, female, 15-19 years				
...	All causes	2,421	100.0	37.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,402	57.9	21.8
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	211	8.7	3.3
3	Malignant neoplasms (C00-C97)	182	7.5	2.8
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	96	4.0	1.5
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	75	3.1	1.2
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	44	1.8	0.7
7	Cerebrovascular diseases (I60-I69)	21	0.9	0.3
8	Diabetes mellitus (E10-E14)	18	0.7	*
9	Septicemia (A40-A41)	14	0.6	*
9	Influenza and pneumonia (J10-J18)	14	0.6	*
11	Chronic lower respiratory diseases (J40-J47)	11	0.5	*
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	10	0.4	*
12	Pregnancy, childbirth and the puerperium (O00-099)	10	0.4	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	9	0.4	*
15	Meningococcal infection (A39)	7	0.3	*
15	Pneumonitis due to solids and liquids (J69)	7	0.3	*
...	All other causes (Residual)	290	12.0	4.5
White, non-Hispanic, female, 20-24 years				
...	All causes	2,995	100.0	46.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,367	45.6	21.4
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	294	9.8	4.6
3	Malignant neoplasms (C00-C97)	238	7.9	3.7
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	160	5.3	2.5
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	107	3.6	1.7
6	Pregnancy, childbirth and the puerperium (O00-099)	66	2.2	1.0
7	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	59	2.0	0.9
8	Cerebrovascular diseases (I60-I69)	31	1.0	0.5
9	Diabetes mellitus (E10-E14)	28	0.9	0.4
10	Influenza and pneumonia (J10-J18)	27	0.9	0.4
11	Septicemia (A40-A41)	17	0.6	*
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	14	0.5	*
13	Chronic lower respiratory diseases (J40-J47)	13	0.4	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	12	0.4	*
15	Human immunodeficiency virus (HIV) disease (B20-B24)	8	0.3	*
15	Pneumonitis due to solids and liquids (J69)	8	0.3	*
...	All other causes (Residual)	546	18.2	8.5

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2005

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
White, non-Hispanic, female, 25-29 years				
...	All causes	3,118	100.0	52.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,138	36.5	19.1
2	Malignant neoplasms (C00-C97)	326	10.5	5.5
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	295	9.5	5.0
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	169	5.4	2.8
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	161	5.2	2.7
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	64	2.1	1.1
7	Pregnancy, childbirth and the puerperium (O00-099)	56	1.8	0.9
8	Influenza and pneumonia (J10-J18)	46	1.5	0.8
9	Diabetes mellitus (E10-E14)	40	1.3	0.7
9	Cerebrovascular diseases (I60-I69)	40	1.3	0.7
11	Septicemia (A40-A41)	31	1.0	0.5
12	Chronic lower respiratory diseases (J40-J47)	28	0.9	0.5
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	28	0.9	0.5
14	Human immunodeficiency virus (HIV) disease (B20-B24)	21	0.7	0.4
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	18	0.6	*
...	All other causes (Residual)	657	21.1	11.0
White, non-Hispanic, female, 30-34 years				
...	All causes	4,342	100.0	71.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,243	28.6	20.5
2	Malignant neoplasms (C00-C97)	746	17.2	12.3
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	418	9.6	6.9
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	367	8.5	6.1
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	148	3.4	2.4
6	Cerebrovascular diseases (I60-I69)	87	2.0	1.4
7	Diabetes mellitus (E10-E14)	77	1.8	1.3
8	Pregnancy, childbirth and the puerperium (O00-099)	65	1.5	1.1
9	Human immunodeficiency virus (HIV) disease (B20-B24)	61	1.4	1.0
9	Influenza and pneumonia (J10-J18)	61	1.4	1.0
11	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	60	1.4	1.0
12	Septicemia (A40-A41)	52	1.2	0.9
13	Chronic liver disease and cirrhosis (K70,K73-K74)	51	1.2	0.8
14	Chronic lower respiratory diseases (J40-J47)	44	1.0	0.7
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	32	0.7	0.5
...	All other causes (Residual)	830	19.1	13.7

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2005

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
White, non-Hispanic, female, 35-39 years				
...	All causes	7,061	100.0	104.4
1	Malignant neoplasms (C00-C97)	1,634	23.1	24.2
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,524	21.6	22.5
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	720	10.2	10.6
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	522	7.4	7.7
5	Diabetes mellitus (E10-E14)	194	2.7	2.9
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	182	2.6	2.7
7	Cerebrovascular diseases (I60-I69)	179	2.5	2.6
8	Chronic liver disease and cirrhosis (K70,K73-K74)	169	2.4	2.5
9	Septicemia (A40-A41)	93	1.3	1.4
10	Chronic lower respiratory diseases (J40-J47)	90	1.3	1.3
11	Human immunodeficiency virus (HIV) disease (B20-B24)	84	1.2	1.2
12	Influenza and pneumonia (J10-J18)	83	1.2	1.2
13	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	79	1.1	1.2
14	Pregnancy, childbirth and the puerperium (O00-099)	58	0.8	0.9
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	53	0.8	0.8
...	All other causes (Residual)	1,397	19.8	20.7
White, non-Hispanic, female, 40-44 years				
...	All causes	13,006	100.0	165.0
1	Malignant neoplasms (C00-C97)	3,862	29.7	49.0
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,143	16.5	27.2
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,513	11.6	19.2
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	748	5.8	9.5
5	Chronic liver disease and cirrhosis (K70,K73-K74)	478	3.7	6.1
6	Cerebrovascular diseases (I60-I69)	398	3.1	5.0
7	Diabetes mellitus (E10-E14)	261	2.0	3.3
8	Chronic lower respiratory diseases (J40-J47)	229	1.8	2.9
9	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	207	1.6	2.6
10	Influenza and pneumonia (J10-J18)	176	1.4	2.2
11	Septicemia (A40-A41)	146	1.1	1.9
11	Human immunodeficiency virus (HIV) disease (B20-B24)	146	1.1	1.9
13	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	107	0.8	1.4
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	79	0.6	1.0
15	Viral hepatitis (B15-B19)	76	0.6	1.0
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	76	0.6	1.0
...	All other causes (Residual)	2,361	18.2	29.9

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2005

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
White, non-Hispanic, female, 45-49 years				
...	All causes	20,183	100.0	248.0
1	Malignant neoplasms (C00-C97)	7,134	35.3	87.6
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,711	13.4	33.3
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,307	11.4	28.3
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	842	4.2	10.3
5	Chronic liver disease and cirrhosis (K70,K73-K74)	761	3.8	9.3
6	Cerebrovascular diseases (I60-I69)	672	3.3	8.3
7	Chronic lower respiratory diseases (J40-J47)	552	2.7	6.8
8	Diabetes mellitus (E10-E14)	452	2.2	5.6
9	Influenza and pneumonia (J10-J18)	263	1.3	3.2
10	Septicemia (A40-A41)	242	1.2	3.0
11	Viral hepatitis (B15-B19)	175	0.9	2.1
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	163	0.8	2.0
13	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	157	0.8	1.9
14	Human immunodeficiency virus (HIV) disease (B20-B24)	124	0.6	1.5
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	116	0.6	1.4
...	All other causes (Residual)	3,512	17.4	43.1
White, non-Hispanic, female, 50-54 years				
...	All causes	26,626	100.0	353.5
1	Malignant neoplasms (C00-C97)	10,725	40.3	142.4
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,939	14.8	52.3
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,788	6.7	23.7
4	Chronic lower respiratory diseases (J40-J47)	940	3.5	12.5
5	Cerebrovascular diseases (I60-I69)	921	3.5	12.2
6	Diabetes mellitus (E10-E14)	823	3.1	10.9
7	Chronic liver disease and cirrhosis (K70,K73-K74)	763	2.9	10.1
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	710	2.7	9.4
9	Septicemia (A40-A41)	392	1.5	5.2
10	Influenza and pneumonia (J10-J18)	339	1.3	4.5
11	Viral hepatitis (B15-B19)	240	0.9	3.2
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	240	0.9	3.2
13	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	150	0.6	2.0
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	134	0.5	1.8
15	Human immunodeficiency virus (HIV) disease (B20-B24)	107	0.4	1.4
...	All other causes (Residual)	4,415	16.6	58.6

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2005

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
White, non-Hispanic, female, 55-59 years				
...	All causes	36,138	100.0	531.6
1	Malignant neoplasms (C00-C97)	15,700	43.4	231.0
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	5,821	16.1	85.6
3	Chronic lower respiratory diseases (J40-J47)	1,950	5.4	28.7
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,446	4.0	21.3
5	Diabetes mellitus (E10-E14)	1,267	3.5	18.6
6	Cerebrovascular diseases (I60-I69)	1,199	3.3	17.6
7	Chronic liver disease and cirrhosis (K70,K73-K74)	805	2.2	11.8
8	Septicemia (A40-A41)	574	1.6	8.4
9	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	535	1.5	7.9
10	Influenza and pneumonia (J10-J18)	460	1.3	6.8
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	424	1.2	6.2
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	212	0.6	3.1
13	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	158	0.4	2.3
14	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	156	0.4	2.3
15	Viral hepatitis (B15-B19)	139	0.4	2.0
...	All other causes (Residual)	5,292	14.6	77.9
White, non-Hispanic, female, 60-64 years				
...	All causes	45,711	100.0	864.3
1	Malignant neoplasms (C00-C97)	19,789	43.3	374.2
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	8,212	18.0	155.3
3	Chronic lower respiratory diseases (J40-J47)	3,479	7.6	65.8
4	Diabetes mellitus (E10-E14)	1,642	3.6	31.0
5	Cerebrovascular diseases (I60-I69)	1,618	3.5	30.6
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,208	2.6	22.8
7	Chronic liver disease and cirrhosis (K70,K73-K74)	715	1.6	13.5
8	Septicemia (A40-A41)	651	1.4	12.3
9	Influenza and pneumonia (J10-J18)	598	1.3	11.3
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	548	1.2	10.4
11	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	334	0.7	6.3
12	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	245	0.5	4.6
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	223	0.5	4.2
14	Alzheimer's disease (G30)	208	0.5	3.9
15	Aortic aneurysm and dissection (I71)	198	0.4	3.7
...	All other causes (Residual)	6,043	13.2	114.3

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2005

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
White, non-Hispanic, female, 65-69 years				
...	All causes	56,712	100.0	1,346.2
1	Malignant neoplasms (C00-C97)	22,489	39.7	533.8
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	11,009	19.4	261.3
3	Chronic lower respiratory diseases (J40-J47)	5,182	9.1	123.0
4	Cerebrovascular diseases (I60-I69)	2,435	4.3	57.8
5	Diabetes mellitus (E10-E14)	2,070	3.7	49.1
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,179	2.1	28.0
7	Septicemia (A40-A41)	877	1.5	20.8
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	849	1.5	20.2
9	Influenza and pneumonia (J10-J18)	807	1.4	19.2
10	Chronic liver disease and cirrhosis (K70,K73-K74)	744	1.3	17.7
11	Alzheimer's disease (G30)	439	0.8	10.4
12	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	377	0.7	8.9
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	321	0.6	7.6
14	Aortic aneurysm and dissection (I71)	304	0.5	7.2
15	Pneumonitis due to solids and liquids (J69)	198	0.3	4.7
...	All other causes (Residual)	7,432	13.1	176.4
White, non-Hispanic, female, 70-74 years				
...	All causes	80,729	100.0	2,162.2
1	Malignant neoplasms (C00-C97)	27,923	34.6	747.9
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	17,169	21.3	459.9
3	Chronic lower respiratory diseases (J40-J47)	7,992	9.9	214.1
4	Cerebrovascular diseases (I60-I69)	4,207	5.2	112.7
5	Diabetes mellitus (E10-E14)	2,815	3.5	75.4
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,479	1.8	39.6
7	Influenza and pneumonia (J10-J18)	1,462	1.8	39.2
8	Alzheimer's disease (G30)	1,389	1.7	37.2
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,372	1.7	36.7
10	Septicemia (A40-A41)	1,287	1.6	34.5
11	Chronic liver disease and cirrhosis (K70,K73-K74)	837	1.0	22.4
12	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	621	0.8	16.6
13	Parkinson's disease (G20-G21)	517	0.6	13.8
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	494	0.6	13.2
15	Aortic aneurysm and dissection (I71)	480	0.6	12.9
...	All other causes (Residual)	10,685	13.2	286.2

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2005

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
White, non-Hispanic, female, 75-79 years				
...	All causes	122,525	100.0	3,481.1
1	Malignant neoplasms (C00-C97)	33,660	27.5	956.3
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	29,411	24.0	835.6
3	Chronic lower respiratory diseases (J40-J47)	11,285	9.2	320.6
4	Cerebrovascular diseases (I60-I69)	8,090	6.6	229.8
5	Alzheimer's disease (G30)	4,031	3.3	114.5
6	Diabetes mellitus (E10-E14)	4,022	3.3	114.3
7	Influenza and pneumonia (J10-J18)	2,757	2.3	78.3
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2,301	1.9	65.4
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,260	1.8	64.2
10	Septicemia (A40-A41)	1,894	1.5	53.8
11	Parkinson's disease (G20-G21)	1,239	1.0	35.2
12	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	1,204	1.0	34.2
13	Aortic aneurysm and dissection (I71)	851	0.7	24.2
14	Chronic liver disease and cirrhosis (K70,K73-K74)	831	0.7	23.6
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	794	0.6	22.6
...	All other causes (Residual)	17,895	14.6	508.4
White, non-Hispanic, female, 80-84 years				
...	All causes	173,579	100.0	5,892.3
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	48,409	27.9	1,643.3
2	Malignant neoplasms (C00-C97)	34,198	19.7	1,160.9
3	Cerebrovascular diseases (I60-I69)	14,361	8.3	487.5
4	Chronic lower respiratory diseases (J40-J47)	12,683	7.3	430.5
5	Alzheimer's disease (G30)	9,112	5.2	309.3
6	Influenza and pneumonia (J10-J18)	5,114	2.9	173.6
7	Diabetes mellitus (E10-E14)	4,969	2.9	168.7
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,371	1.9	114.4
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3,195	1.8	108.5
10	Septicemia (A40-A41)	2,652	1.5	90.0
11	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	2,048	1.2	69.5
12	Parkinson's disease (G20-G21)	2,042	1.2	69.3
13	Pneumonitis due to solids and liquids (J69)	1,376	0.8	46.7
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1,103	0.6	37.4
15	Atherosclerosis (I70)	1,083	0.6	36.8
...	All other causes (Residual)	27,863	16.1	945.8

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2005

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
White, non-Hispanic, female, 85-89 years				
...	All causes	186,328	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	59,514	31.9	---
2	Malignant neoplasms (C00-C97)	24,466	13.1	---
3	Cerebrovascular diseases (I60-I69)	16,842	9.0	---
4	Alzheimer's disease (G30)	12,785	6.9	---
5	Chronic lower respiratory diseases (J40-J47)	10,225	5.5	---
6	Influenza and pneumonia (J10-J18)	6,763	3.6	---
7	Diabetes mellitus (E10-E14)	4,455	2.4	---
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,644	2.0	---
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3,427	1.8	---
10	Septicemia (A40-A41)	2,582	1.4	---
11	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	2,547	1.4	---
12	Parkinson's disease (G20-G21)	2,000	1.1	---
13	Pneumonitis due to solids and liquids (J69)	1,760	0.9	---
14	Atherosclerosis (I70)	1,601	0.9	---
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1,145	0.6	---
...	All other causes (Residual)	32,572	17.5	---
White, non-Hispanic, female, 90-94 years				
...	All causes	145,875	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	52,511	36.0	---
2	Cerebrovascular diseases (I60-I69)	13,448	9.2	---
3	Malignant neoplasms (C00-C97)	12,114	8.3	---
4	Alzheimer's disease (G30)	11,472	7.9	---
5	Influenza and pneumonia (J10-J18)	6,337	4.3	---
6	Chronic lower respiratory diseases (J40-J47)	5,539	3.8	---
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,857	2.0	---
8	Diabetes mellitus (E10-E14)	2,612	1.8	---
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2,554	1.8	---
10	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	2,493	1.7	---
11	Septicemia (A40-A41)	1,813	1.2	---
12	Atherosclerosis (I70)	1,624	1.1	---
13	Pneumonitis due to solids and liquids (J69)	1,533	1.1	---
14	Parkinson's disease (G20-G21)	1,085	0.7	---
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	765	0.5	---
...	All other causes (Residual)	27,118	18.6	---

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2005

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
White, non-Hispanic, female, 95-99 years				
...	All causes	63,498	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	24,315	38.3	---
2	Cerebrovascular diseases (I60-I69)	5,379	8.5	---
3	Alzheimer's disease (G30)	5,219	8.2	---
4	Malignant neoplasms (C00-C97)	3,432	5.4	---
5	Influenza and pneumonia (J10-J18)	3,258	5.1	---
6	Chronic lower respiratory diseases (J40-J47)	1,789	2.8	---
7	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	1,287	2.0	---
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,168	1.8	---
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,019	1.6	---
10	Atherosclerosis (I70)	907	1.4	---
11	Diabetes mellitus (E10-E14)	896	1.4	---
12	Pneumonitis due to solids and liquids (J69)	691	1.1	---
13	Septicemia (A40-A41)	664	1.0	---
14	Parkinson's disease (G20-G21)	274	0.4	---
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	249	0.4	---
...	All other causes (Residual)	12,951	20.4	---
White, non-Hispanic, female, 100 years and over				
...	All causes	13,750	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	5,650	41.1	---
2	Cerebrovascular diseases (I60-I69)	1,056	7.7	---
3	Alzheimer's disease (G30)	951	6.9	---
4	Influenza and pneumonia (J10-J18)	868	6.3	---
5	Malignant neoplasms (C00-C97)	480	3.5	---
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	279	2.0	---
7	Chronic lower respiratory diseases (J40-J47)	261	1.9	---
8	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	260	1.9	---
9	Atherosclerosis (I70)	243	1.8	---
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	198	1.4	---
11	Diabetes mellitus (E10-E14)	138	1.0	---
12	Pneumonitis due to solids and liquids (J69)	131	1.0	---
13	Septicemia (A40-A41)	105	0.8	---
14	Anemias (D50-D64)	70	0.5	---
15	Nutritional deficiencies (E40-E64)	43	0.3	---
...	All other causes (Residual)	3,017	21.9	---

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2005

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Black, non-Hispanic, both sexes, all ages				
...	All causes	289,163	100.0	774.4
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	73,302	25.3	196.3
2	Malignant neoplasms (C00-C97)	62,510	21.6	167.4
3	Cerebrovascular diseases (I60-I69)	17,378	6.0	46.5
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	13,393	4.6	35.9
5	Diabetes mellitus (E10-E14)	12,835	4.4	34.4
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	8,522	2.9	22.8
7	Chronic lower respiratory diseases (J40-J47)	8,136	2.8	21.8
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	8,017	2.8	21.5
9	Human immunodeficiency virus (HIV) disease (B20-B24)	6,908	2.4	18.5
10	Septicemia (A40-A41)	6,159	2.1	16.5
11	Influenza and pneumonia (J10-J18)	5,716	2.0	15.3
12	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	4,906	1.7	13.1
13	Certain conditions originating in the perinatal period (P00-P96)	4,837	1.7	13.0
14	Alzheimer's disease (G30)	4,579	1.6	12.3
15	Chronic liver disease and cirrhosis (K70,K73-K74)	2,495	0.9	6.7
...	All other causes (Residual)	49,470	17.1	132.5
Black, non-Hispanic, both sexes, 1-4 years				
...	All causes	1,065	100.0	42.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	316	29.7	12.6
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	138	13.0	5.5
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	104	9.8	4.1
4	Malignant neoplasms (C00-C97)	62	5.8	2.5
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	45	4.2	1.8
6	Chronic lower respiratory diseases (J40-J47)	30	2.8	1.2
7	Influenza and pneumonia (J10-J18)	28	2.6	1.1
8	Certain conditions originating in the perinatal period (P00-P96)	23	2.2	0.9
9	Septicemia (A40-A41)	20	1.9	0.8
10	Cerebrovascular diseases (I60-I69)	14	1.3	*
11	Anemias (D50-D64)	11	1.0	*
12	Meningitis (G00,G03)	6	0.6	*
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	5	0.5	*
13	Pneumonitis due to solids and liquids (J69)	5	0.5	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3	0.3	*
...	All other causes (Residual)	255	23.9	10.2

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2005

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Black, non-Hispanic, both sexes, 5-9 years				
...	All causes	645	100.0	21.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	235	36.4	8.0
2	Malignant neoplasms (C00-C97)	79	12.2	2.7
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	44	6.8	1.5
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	36	5.6	1.2
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	34	5.3	1.2
6	Chronic lower respiratory diseases (J40-J47)	22	3.4	0.7
7	Cerebrovascular diseases (I60-I69)	13	2.0	*
8	Septicemia (A40-A41)	11	1.7	*
8	Influenza and pneumonia (J10-J18)	11	1.7	*
10	Anemias (D50-D64)	8	1.2	*
11	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	7	1.1	*
12	Meningitis (G00,G03)	3	0.5	*
13	Human immunodeficiency virus (HIV) disease (B20-B24)	2	0.3	*
13	Pneumonitis due to solids and liquids (J69)	2	0.3	*
13	Certain conditions originating in the perinatal period (P00-P96)	2	0.3	*
...	All other causes (Residual)	136	21.1	4.6
Black, non-Hispanic, both sexes, 10-14 years				
...	All causes	855	100.0	25.9
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	262	30.6	7.9
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	101	11.8	3.1
3	Malignant neoplasms (C00-C97)	86	10.1	2.6
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	44	5.1	1.3
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	41	4.8	1.2
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	40	4.7	1.2
7	Chronic lower respiratory diseases (J40-J47)	29	3.4	0.9
8	Diabetes mellitus (E10-E14)	12	1.4	*
9	Influenza and pneumonia (J10-J18)	11	1.3	*
10	Cerebrovascular diseases (I60-I69)	10	1.2	*
11	Septicemia (A40-A41)	9	1.1	*
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	7	0.8	*
12	Anemias (D50-D64)	7	0.8	*
14	Human immunodeficiency virus (HIV) disease (B20-B24)	6	0.7	*
15	Meningitis (G00,G03)	4	0.5	*
15	Diseases of appendix (K35-K38)	4	0.5	*
...	All other causes (Residual)	182	21.3	5.5

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2005

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Black, non-Hispanic, both sexes, 15-19 years				
...	All causes	2,793	100.0	86.2
1	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,120	40.1	34.6
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	754	27.0	23.3
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	145	5.2	4.5
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	116	4.2	3.6
5	Malignant neoplasms (C00-C97)	101	3.6	3.1
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	55	2.0	1.7
7	Anemias (D50-D64)	30	1.1	0.9
8	Chronic lower respiratory diseases (J40-J47)	25	0.9	0.8
9	Diabetes mellitus (E10-E14)	24	0.9	0.7
10	Human immunodeficiency virus (HIV) disease (B20-B24)	22	0.8	0.7
11	Septicemia (A40-A41)	20	0.7	0.6
11	Pregnancy, childbirth and the puerperium (O00-099)	20	0.7	0.6
13	Cerebrovascular diseases (I60-I69)	19	0.7	*
14	Influenza and pneumonia (J10-J18)	13	0.5	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	12	0.4	*
...	All other causes (Residual)	317	11.3	9.8
Black, non-Hispanic, both sexes, 20-24 years				
...	All causes	4,451	100.0	147.6
1	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,910	42.9	63.4
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	980	22.0	32.5
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	283	6.4	9.4
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	217	4.9	7.2
5	Malignant neoplasms (C00-C97)	158	3.5	5.2
6	Human immunodeficiency virus (HIV) disease (B20-B24)	86	1.9	2.9
7	Diabetes mellitus (E10-E14)	49	1.1	1.6
7	Chronic lower respiratory diseases (J40-J47)	49	1.1	1.6
7	Pregnancy, childbirth and the puerperium (O00-099)	49	1.1	1.6
10	Anemias (D50-D64)	47	1.1	1.6
11	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	44	1.0	1.5
12	Cerebrovascular diseases (I60-I69)	32	0.7	1.1
13	Septicemia (A40-A41)	24	0.5	0.8
14	Influenza and pneumonia (J10-J18)	18	0.4	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	17	0.4	*
...	All other causes (Residual)	488	11.0	16.2

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2005

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Black, non-Hispanic, both sexes, 25-29 years				
...	All causes	4,559	100.0	168.3
1	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,492	32.7	55.1
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	826	18.1	30.5
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	370	8.1	13.7
4	Human immunodeficiency virus (HIV) disease (B20-B24)	274	6.0	10.1
5	Malignant neoplasms (C00-C97)	256	5.6	9.4
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	249	5.5	9.2
7	Diabetes mellitus (E10-E14)	80	1.8	3.0
8	Anemias (D50-D64)	65	1.4	2.4
8	Pregnancy, childbirth and the puerperium (O00-099)	65	1.4	2.4
10	Cerebrovascular diseases (I60-I69)	64	1.4	2.4
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	46	1.0	1.7
11	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	46	1.0	1.7
13	Septicemia (A40-A41)	44	1.0	1.6
14	Chronic lower respiratory diseases (J40-J47)	34	0.7	1.3
15	Influenza and pneumonia (J10-J18)	27	0.6	1.0
...	All other causes (Residual)	621	13.6	22.9
Black, non-Hispanic, both sexes, 30-34 years				
...	All causes	5,291	100.0	204.6
1	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,046	19.8	40.4
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	928	17.5	35.9
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	638	12.1	24.7
4	Human immunodeficiency virus (HIV) disease (B20-B24)	550	10.4	21.3
5	Malignant neoplasms (C00-C97)	418	7.9	16.2
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	225	4.3	8.7
7	Diabetes mellitus (E10-E14)	138	2.6	5.3
8	Cerebrovascular diseases (I60-I69)	99	1.9	3.8
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	74	1.4	2.9
10	Anemias (D50-D64)	66	1.2	2.6
11	Septicemia (A40-A41)	62	1.2	2.4
11	Chronic lower respiratory diseases (J40-J47)	62	1.2	2.4
13	Influenza and pneumonia (J10-J18)	50	0.9	1.9
13	Pregnancy, childbirth and the puerperium (O00-099)	50	0.9	1.9
15	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	36	0.7	1.4
...	All other causes (Residual)	849	16.0	32.8

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2005

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Black, non-Hispanic, both sexes, 35-39 years				
...	All causes	6,948	100.0	263.2
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,142	16.4	43.3
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	977	14.1	37.0
3	Human immunodeficiency virus (HIV) disease (B20-B24)	943	13.6	35.7
4	Malignant neoplasms (C00-C97)	765	11.0	29.0
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	725	10.4	27.5
6	Diabetes mellitus (E10-E14)	221	3.2	8.4
7	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	197	2.8	7.5
8	Cerebrovascular diseases (I60-I69)	187	2.7	7.1
9	Septicemia (A40-A41)	99	1.4	3.8
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	98	1.4	3.7
11	Influenza and pneumonia (J10-J18)	70	1.0	2.7
11	Chronic lower respiratory diseases (J40-J47)	70	1.0	2.7
13	Anemias (D50-D64)	61	0.9	2.3
14	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	60	0.9	2.3
15	Chronic liver disease and cirrhosis (K70,K73-K74)	56	0.8	2.1
...	All other causes (Residual)	1,277	18.4	48.4
Black, non-Hispanic, both sexes, 40-44 years				
...	All causes	10,982	100.0	389.7
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,023	18.4	71.8
2	Malignant neoplasms (C00-C97)	1,756	16.0	62.3
3	Human immunodeficiency virus (HIV) disease (B20-B24)	1,355	12.3	48.1
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,184	10.8	42.0
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	626	5.7	22.2
6	Cerebrovascular diseases (I60-I69)	496	4.5	17.6
7	Diabetes mellitus (E10-E14)	313	2.9	11.1
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	218	2.0	7.7
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	214	1.9	7.6
10	Chronic liver disease and cirrhosis (K70,K73-K74)	178	1.6	6.3
11	Septicemia (A40-A41)	159	1.4	5.6
12	Chronic lower respiratory diseases (J40-J47)	147	1.3	5.2
13	Influenza and pneumonia (J10-J18)	144	1.3	5.1
14	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	134	1.2	4.8
15	Anemias (D50-D64)	83	0.8	2.9
...	All other causes (Residual)	1,952	17.8	69.3

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2005

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Black, non-Hispanic, both sexes, 45-49 years				
...	All causes	16,504	100.0	616.8
1	Malignant neoplasms (C00-C97)	3,540	21.4	132.3
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,511	21.3	131.2
3	Human immunodeficiency virus (HIV) disease (B20-B24)	1,398	8.5	52.2
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,349	8.2	50.4
5	Cerebrovascular diseases (I60-I69)	838	5.1	31.3
6	Diabetes mellitus (E10-E14)	567	3.4	21.2
7	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	471	2.9	17.6
8	Chronic liver disease and cirrhosis (K70,K73-K74)	392	2.4	14.6
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	333	2.0	12.4
10	Chronic lower respiratory diseases (J40-J47)	289	1.8	10.8
11	Septicemia (A40-A41)	275	1.7	10.3
12	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	236	1.4	8.8
13	Influenza and pneumonia (J10-J18)	216	1.3	8.1
14	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	167	1.0	6.2
15	Viral hepatitis (B15-B19)	130	0.8	4.9
...	All other causes (Residual)	2,792	16.9	104.3
Black, non-Hispanic, both sexes, 50-54 years				
...	All causes	20,794	100.0	935.4
1	Malignant neoplasms (C00-C97)	5,265	25.3	236.9
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	5,102	24.5	229.5
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,298	6.2	58.4
4	Cerebrovascular diseases (I60-I69)	1,148	5.5	51.6
5	Human immunodeficiency virus (HIV) disease (B20-B24)	1,077	5.2	48.5
6	Diabetes mellitus (E10-E14)	829	4.0	37.3
7	Chronic liver disease and cirrhosis (K70,K73-K74)	492	2.4	22.1
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	412	2.0	18.5
9	Chronic lower respiratory diseases (J40-J47)	408	2.0	18.4
10	Septicemia (A40-A41)	379	1.8	17.0
11	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	314	1.5	14.1
12	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	291	1.4	13.1
13	Influenza and pneumonia (J10-J18)	278	1.3	12.5
14	Viral hepatitis (B15-B19)	219	1.1	9.9
15	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	142	0.7	6.4
...	All other causes (Residual)	3,140	15.1	141.3

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2005

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Black, non-Hispanic, both sexes, 55-59 years				
...	All causes	22,417	100.0	1,281.3
1	Malignant neoplasms (C00-C97)	6,700	29.9	383.0
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	5,832	26.0	333.3
3	Cerebrovascular diseases (I60-I69)	1,178	5.3	67.3
4	Diabetes mellitus (E10-E14)	1,141	5.1	65.2
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	877	3.9	50.1
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	644	2.9	36.8
7	Human immunodeficiency virus (HIV) disease (B20-B24)	605	2.7	34.6
8	Chronic lower respiratory diseases (J40-J47)	545	2.4	31.2
9	Septicemia (A40-A41)	455	2.0	26.0
10	Chronic liver disease and cirrhosis (K70,K73-K74)	436	1.9	24.9
11	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	328	1.5	18.7
12	Influenza and pneumonia (J10-J18)	301	1.3	17.2
13	Viral hepatitis (B15-B19)	163	0.7	9.3
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	147	0.7	8.4
15	Aortic aneurysm and dissection (I71)	98	0.4	5.6
...	All other causes (Residual)	2,967	13.2	169.6
Black, non-Hispanic, both sexes, 60-64 years				
...	All causes	22,520	100.0	1,846.9
1	Malignant neoplasms (C00-C97)	7,135	31.7	585.2
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	6,145	27.3	504.0
3	Diabetes mellitus (E10-E14)	1,274	5.7	104.5
4	Cerebrovascular diseases (I60-I69)	1,242	5.5	101.9
5	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	696	3.1	57.1
6	Chronic lower respiratory diseases (J40-J47)	650	2.9	53.3
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	564	2.5	46.3
8	Septicemia (A40-A41)	479	2.1	39.3
9	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	356	1.6	29.2
10	Chronic liver disease and cirrhosis (K70,K73-K74)	315	1.4	25.8
11	Influenza and pneumonia (J10-J18)	302	1.3	24.8
12	Human immunodeficiency virus (HIV) disease (B20-B24)	297	1.3	24.4
13	Aortic aneurysm and dissection (I71)	109	0.5	8.9
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	97	0.4	8.0
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	94	0.4	7.7
...	All other causes (Residual)	2,765	12.3	226.8

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2005

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Black, non-Hispanic, both sexes, 65-69 years				
...	All causes	24,098	100.0	2,511.2
1	Malignant neoplasms (C00-C97)	7,656	31.8	797.8
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	6,460	26.8	673.2
3	Diabetes mellitus (E10-E14)	1,473	6.1	153.5
4	Cerebrovascular diseases (I60-I69)	1,412	5.9	147.1
5	Chronic lower respiratory diseases (J40-J47)	898	3.7	93.6
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	782	3.2	81.5
7	Septicemia (A40-A41)	545	2.3	56.8
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	465	1.9	48.5
9	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	435	1.8	45.3
10	Influenza and pneumonia (J10-J18)	384	1.6	40.0
11	Chronic liver disease and cirrhosis (K70,K73-K74)	195	0.8	20.3
12	Human immunodeficiency virus (HIV) disease (B20-B24)	161	0.7	16.8
13	Aortic aneurysm and dissection (I71)	131	0.5	13.7
14	Alzheimer's disease (G30)	102	0.4	10.6
15	Pneumonitis due to solids and liquids (J69)	101	0.4	10.5
...	All other causes (Residual)	2,898	12.0	302.0
Black, non-Hispanic, both sexes, 70-74 years				
...	All causes	27,068	100.0	3,572.9
1	Malignant neoplasms (C00-C97)	7,904	29.2	1,043.3
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	7,380	27.3	974.1
3	Cerebrovascular diseases (I60-I69)	1,782	6.6	235.2
4	Diabetes mellitus (E10-E14)	1,628	6.0	214.9
5	Chronic lower respiratory diseases (J40-J47)	1,100	4.1	145.2
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	962	3.6	127.0
7	Septicemia (A40-A41)	626	2.3	82.6
8	Influenza and pneumonia (J10-J18)	530	2.0	70.0
9	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	513	1.9	67.7
10	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	456	1.7	60.2
11	Alzheimer's disease (G30)	288	1.1	38.0
12	Chronic liver disease and cirrhosis (K70,K73-K74)	157	0.6	20.7
13	Pneumonitis due to solids and liquids (J69)	131	0.5	17.3
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	127	0.5	16.8
15	Aortic aneurysm and dissection (I71)	123	0.5	16.2
...	All other causes (Residual)	3,361	12.4	443.6

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2005

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Black, non-Hispanic, both sexes, 75-79 years				
...	All causes	30,563	100.0	5,236.9
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	8,844	28.9	1,515.4
2	Malignant neoplasms (C00-C97)	7,688	25.2	1,317.3
3	Cerebrovascular diseases (I60-I69)	2,201	7.2	377.1
4	Diabetes mellitus (E10-E14)	1,673	5.5	286.7
5	Chronic lower respiratory diseases (J40-J47)	1,305	4.3	223.6
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,050	3.4	179.9
7	Septicemia (A40-A41)	747	2.4	128.0
8	Influenza and pneumonia (J10-J18)	684	2.2	117.2
9	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	600	2.0	102.8
10	Alzheimer's disease (G30)	593	1.9	101.6
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	511	1.7	87.6
12	Pneumonitis due to solids and liquids (J69)	244	0.8	41.8
13	Aortic aneurysm and dissection (I71)	152	0.5	26.0
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	142	0.5	24.3
15	Parkinson's disease (G20-G21)	134	0.4	23.0
...	All other causes (Residual)	3,995	13.1	684.5
Black, non-Hispanic, both sexes, 80-84 years				
...	All causes	30,756	100.0	7,641.2
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	9,274	30.2	2,304.1
2	Malignant neoplasms (C00-C97)	6,468	21.0	1,607.0
3	Cerebrovascular diseases (I60-I69)	2,490	8.1	618.6
4	Diabetes mellitus (E10-E14)	1,532	5.0	380.6
5	Chronic lower respiratory diseases (J40-J47)	1,109	3.6	275.5
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,040	3.4	258.4
7	Alzheimer's disease (G30)	986	3.2	245.0
8	Septicemia (A40-A41)	830	2.7	206.2
9	Influenza and pneumonia (J10-J18)	813	2.6	202.0
10	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	626	2.0	155.5
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	453	1.5	112.5
12	Pneumonitis due to solids and liquids (J69)	281	0.9	69.8
13	Parkinson's disease (G20-G21)	205	0.7	50.9
14	Aortic aneurysm and dissection (I71)	155	0.5	38.5
15	Atherosclerosis (I70)	139	0.5	34.5
...	All other causes (Residual)	4,355	14.2	1,082.0

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2005

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Black, non-Hispanic, both sexes, 85-89 years				
...	All causes	23,772	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	7,484	31.5	---
2	Malignant neoplasms (C00-C97)	3,949	16.6	---
3	Cerebrovascular diseases (I60-I69)	2,038	8.6	---
4	Alzheimer's disease (G30)	1,094	4.6	---
5	Diabetes mellitus (E10-E14)	1,042	4.4	---
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	826	3.5	---
7	Influenza and pneumonia (J10-J18)	774	3.3	---
8	Chronic lower respiratory diseases (J40-J47)	755	3.2	---
9	Septicemia (A40-A41)	624	2.6	---
10	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	565	2.4	---
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	312	1.3	---
12	Pneumonitis due to solids and liquids (J69)	263	1.1	---
13	Atherosclerosis (I70)	147	0.6	---
14	Parkinson's disease (G20-G21)	128	0.5	---
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	116	0.5	---
...	All other causes (Residual)	3,655	15.4	---
Black, non-Hispanic, both sexes, 90-94 years				
...	All causes	15,851	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	5,308	33.5	---
2	Malignant neoplasms (C00-C97)	1,835	11.6	---
3	Cerebrovascular diseases (I60-I69)	1,400	8.8	---
4	Alzheimer's disease (G30)	915	5.8	---
5	Diabetes mellitus (E10-E14)	606	3.8	---
6	Influenza and pneumonia (J10-J18)	586	3.7	---
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	500	3.2	---
8	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	431	2.7	---
9	Chronic lower respiratory diseases (J40-J47)	406	2.6	---
10	Septicemia (A40-A41)	405	2.6	---
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	217	1.4	---
12	Pneumonitis due to solids and liquids (J69)	191	1.2	---
13	Atherosclerosis (I70)	122	0.8	---
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	58	0.4	---
14	Aortic aneurysm and dissection (I71)	58	0.4	---
...	All other causes (Residual)	2,813	17.7	---

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2005

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Black, non-Hispanic, both sexes, 95-99 years				
...	All causes	6,736	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,417	35.9	---
2	Malignant neoplasms (C00-C97)	578	8.6	---
3	Cerebrovascular diseases (I60-I69)	508	7.5	---
4	Alzheimer's disease (G30)	415	6.2	---
5	Influenza and pneumonia (J10-J18)	270	4.0	---
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	201	3.0	---
7	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	197	2.9	---
8	Diabetes mellitus (E10-E14)	185	2.7	---
9	Septicemia (A40-A41)	172	2.6	---
10	Chronic lower respiratory diseases (J40-J47)	157	2.3	---
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	95	1.4	---
12	Pneumonitis due to solids and liquids (J69)	91	1.4	---
13	Atherosclerosis (I70)	63	0.9	---
14	Nutritional deficiencies (E40-E64)	36	0.5	---
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	26	0.4	---
...	All other causes (Residual)	1,325	19.7	---
Black, non-Hispanic, both sexes, 100 years and over				
...	All causes	2,122	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	807	38.0	---
2	Cerebrovascular diseases (I60-I69)	164	7.7	---
3	Alzheimer's disease (G30)	135	6.4	---
4	Influenza and pneumonia (J10-J18)	111	5.2	---
5	Malignant neoplasms (C00-C97)	99	4.7	---
6	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	71	3.3	---
7	Septicemia (A40-A41)	47	2.2	---
7	Diabetes mellitus (E10-E14)	47	2.2	---
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	47	2.2	---
10	Chronic lower respiratory diseases (J40-J47)	34	1.6	---
11	Atherosclerosis (I70)	33	1.6	---
12	Pneumonitis due to solids and liquids (J69)	24	1.1	---
13	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	23	1.1	---
14	Nutritional deficiencies (E40-E64)	11	0.5	---
15	Parkinson's disease (G20-G21)	5	0.2	---
...	All other causes (Residual)	464	21.9	---

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2005

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Black, non-Hispanic, male, all ages				
...	All causes	147,010	100.0	825.7
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	35,874	24.4	201.5
2	Malignant neoplasms (C00-C97)	32,363	22.0	181.8
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	9,130	6.2	51.3
4	Cerebrovascular diseases (I60-I69)	7,448	5.1	41.8
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	7,288	5.0	40.9
6	Diabetes mellitus (E10-E14)	5,666	3.9	31.8
7	Human immunodeficiency virus (HIV) disease (B20-B24)	4,600	3.1	25.8
8	Chronic lower respiratory diseases (J40-J47)	4,406	3.0	24.7
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3,612	2.5	20.3
10	Septicemia (A40-A41)	2,755	1.9	15.5
11	Certain conditions originating in the perinatal period (P00-P96)	2,712	1.8	15.2
12	Influenza and pneumonia (J10-J18)	2,697	1.8	15.1
13	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	2,108	1.4	11.8
14	Chronic liver disease and cirrhosis (K70,K73-K74)	1,634	1.1	9.2
15	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,592	1.1	8.9
...	All other causes (Residual)	23,125	15.7	129.9
Black, non-Hispanic, male, 1-4 years				
...	All causes	606	100.0	47.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	179	29.5	14.1
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	88	14.5	6.9
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	46	7.6	3.6
4	Malignant neoplasms (C00-C97)	36	5.9	2.8
5	Chronic lower respiratory diseases (J40-J47)	21	3.5	1.6
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	19	3.1	*
7	Influenza and pneumonia (J10-J18)	16	2.6	*
7	Certain conditions originating in the perinatal period (P00-P96)	16	2.6	*
9	Septicemia (A40-A41)	13	2.1	*
10	Cerebrovascular diseases (I60-I69)	10	1.7	*
11	Anemias (D50-D64)	5	0.8	*
12	Meningitis (G00,G03)	3	0.5	*
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3	0.5	*
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2	0.3	*
14	Pneumonitis due to solids and liquids (J69)	2	0.3	*
...	All other causes (Residual)	147	24.3	11.5

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2005

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Black, non-Hispanic, male, 5-9 years				
...	All causes	354	100.0	23.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	132	37.3	8.8
2	Malignant neoplasms (C00-C97)	46	13.0	3.1
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	24	6.8	1.6
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	16	4.5	*
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	14	4.0	*
6	Chronic lower respiratory diseases (J40-J47)	12	3.4	*
7	Septicemia (A40-A41)	9	2.5	*
8	Cerebrovascular diseases (I60-I69)	8	2.3	*
9	Influenza and pneumonia (J10-J18)	7	2.0	*
10	Anemias (D50-D64)	5	1.4	*
11	Meningitis (G00,G03)	3	0.8	*
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2	0.6	*
13	Nutritional deficiencies (E40-E64)	1	0.3	*
13	Acute bronchitis and bronchiolitis (J20-J21)	1	0.3	*
13	Hernia (K40-K46)	1	0.3	*
13	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1	0.3	*
13	Complications of medical and surgical care (Y40-Y84,Y88)	1	0.3	*
...	All other causes (Residual)	71	20.1	4.7
Black, non-Hispanic, male, 10-14 years				
...	All causes	531	100.0	31.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	180	33.9	10.7
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	69	13.0	4.1
3	Malignant neoplasms (C00-C97)	52	9.8	3.1
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	26	4.9	1.5
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	21	4.0	1.3
5	Chronic lower respiratory diseases (J40-J47)	21	4.0	1.3
5	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	21	4.0	1.3
8	Cerebrovascular diseases (I60-I69)	6	1.1	*
8	Influenza and pneumonia (J10-J18)	6	1.1	*
10	Human immunodeficiency virus (HIV) disease (B20-B24)	5	0.9	*
11	Septicemia (A40-A41)	4	0.8	*
11	Anemias (D50-D64)	4	0.8	*
13	Diabetes mellitus (E10-E14)	3	0.6	*
13	Meningitis (G00,G03)	3	0.6	*
13	Diseases of appendix (K35-K38)	3	0.6	*
...	All other causes (Residual)	107	20.2	6.4

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2005

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Black, non-Hispanic, male, 15-19 years				
...	All causes	2,155	100.0	131.3
1	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,018	47.2	62.0
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	546	25.3	33.3
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	122	5.7	7.4
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	85	3.9	5.2
5	Malignant neoplasms (C00-C97)	56	2.6	3.4
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	36	1.7	2.2
7	Chronic lower respiratory diseases (J40-J47)	20	0.9	1.2
8	Anemias (D50-D64)	18	0.8	*
9	Diabetes mellitus (E10-E14)	17	0.8	*
10	Septicemia (A40-A41)	13	0.6	*
11	Legal intervention (Y35,Y89.0)	11	0.5	*
12	Human immunodeficiency virus (HIV) disease (B20-B24)	10	0.5	*
13	Influenza and pneumonia (J10-J18)	9	0.4	*
14	Cerebrovascular diseases (I60-I69)	8	0.4	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	4	0.2	*
...	All other causes (Residual)	182	8.4	11.1
Black, non-Hispanic, male, 20-24 years				
...	All causes	3,460	100.0	227.8
1	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,732	50.1	114.0
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	755	21.8	49.7
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	253	7.3	16.7
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	143	4.1	9.4
5	Malignant neoplasms (C00-C97)	95	2.7	6.3
6	Human immunodeficiency virus (HIV) disease (B20-B24)	47	1.4	3.1
7	Anemias (D50-D64)	27	0.8	1.8
7	Chronic lower respiratory diseases (J40-J47)	27	0.8	1.8
7	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	27	0.8	1.8
10	Diabetes mellitus (E10-E14)	22	0.6	1.4
11	Cerebrovascular diseases (I60-I69)	16	0.5	*
12	Influenza and pneumonia (J10-J18)	14	0.4	*
13	Legal intervention (Y35,Y89.0)	13	0.4	*
14	Septicemia (A40-A41)	11	0.3	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	7	0.2	*
...	All other causes (Residual)	271	7.8	17.8

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2005

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Black, non-Hispanic, male, 25-29 years				
...	All causes	3,268	100.0	248.7
1	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,349	41.3	102.7
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	642	19.6	48.9
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	227	6.9	17.3
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	207	6.3	15.8
5	Human immunodeficiency virus (HIV) disease (B20-B24)	152	4.7	11.6
6	Malignant neoplasms (C00-C97)	124	3.8	9.4
7	Diabetes mellitus (E10-E14)	41	1.3	3.1
8	Cerebrovascular diseases (I60-I69)	36	1.1	2.7
9	Anemias (D50-D64)	34	1.0	2.6
10	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	29	0.9	2.2
11	Septicemia (A40-A41)	23	0.7	1.8
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	23	0.7	1.8
13	Chronic lower respiratory diseases (J40-J47)	21	0.6	1.6
14	Influenza and pneumonia (J10-J18)	15	0.5	*
15	Legal intervention (Y35,Y89.0)	12	0.4	*
...	All other causes (Residual)	333	10.2	25.3
Black, non-Hispanic, male, 30-34 years				
...	All causes	3,438	100.0	280.3
1	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	921	26.8	75.1
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	690	20.1	56.3
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	397	11.5	32.4
4	Human immunodeficiency virus (HIV) disease (B20-B24)	303	8.8	24.7
5	Malignant neoplasms (C00-C97)	189	5.5	15.4
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	187	5.4	15.2
7	Diabetes mellitus (E10-E14)	86	2.5	7.0
8	Cerebrovascular diseases (I60-I69)	53	1.5	4.3
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	37	1.1	3.0
10	Anemias (D50-D64)	35	1.0	2.9
11	Chronic lower respiratory diseases (J40-J47)	31	0.9	2.5
12	Influenza and pneumonia (J10-J18)	29	0.8	2.4
13	Legal intervention (Y35,Y89.0)	27	0.8	2.2
14	Septicemia (A40-A41)	26	0.8	2.1
15	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	21	0.6	1.7
...	All other causes (Residual)	406	11.8	33.1

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2005

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Black, non-Hispanic, male, 35-39 years				
...	All causes	4,172	100.0	334.8
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	726	17.4	58.3
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	709	17.0	56.9
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	577	13.8	46.3
4	Human immunodeficiency virus (HIV) disease (B20-B24)	576	13.8	46.2
5	Malignant neoplasms (C00-C97)	280	6.7	22.5
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	156	3.7	12.5
7	Diabetes mellitus (E10-E14)	140	3.4	11.2
8	Cerebrovascular diseases (I60-I69)	92	2.2	7.4
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	48	1.2	3.9
10	Septicemia (A40-A41)	46	1.1	3.7
11	Influenza and pneumonia (J10-J18)	43	1.0	3.5
12	Aortic aneurysm and dissection (I71)	38	0.9	3.0
13	Anemias (D50-D64)	35	0.8	2.8
13	Chronic lower respiratory diseases (J40-J47)	35	0.8	2.8
15	Chronic liver disease and cirrhosis (K70,K73-K74)	33	0.8	2.6
...	All other causes (Residual)	638	15.3	51.2
Black, non-Hispanic, male, 40-44 years				
...	All causes	6,279	100.0	476.2
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,243	19.8	94.3
2	Human immunodeficiency virus (HIV) disease (B20-B24)	878	14.0	66.6
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	812	12.9	61.6
4	Malignant neoplasms (C00-C97)	680	10.8	51.6
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	498	7.9	37.8
6	Cerebrovascular diseases (I60-I69)	237	3.8	18.0
7	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	177	2.8	13.4
8	Diabetes mellitus (E10-E14)	167	2.7	12.7
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	116	1.8	8.8
10	Chronic liver disease and cirrhosis (K70,K73-K74)	108	1.7	8.2
11	Influenza and pneumonia (J10-J18)	84	1.3	6.4
12	Septicemia (A40-A41)	75	1.2	5.7
13	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	72	1.1	5.5
14	Chronic lower respiratory diseases (J40-J47)	68	1.1	5.2
15	Aortic aneurysm and dissection (I71)	43	0.7	3.3
...	All other causes (Residual)	1,021	16.3	77.4

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2005

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Black, non-Hispanic, male, 45-49 years				
...	All causes	9,622	100.0	772.6
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,237	23.2	179.6
2	Malignant neoplasms (C00-C97)	1,619	16.8	130.0
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	954	9.9	76.6
4	Human immunodeficiency virus (HIV) disease (B20-B24)	951	9.9	76.4
5	Cerebrovascular diseases (I60-I69)	408	4.2	32.8
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	375	3.9	30.1
7	Diabetes mellitus (E10-E14)	329	3.4	26.4
8	Chronic liver disease and cirrhosis (K70,K73-K74)	268	2.8	21.5
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	203	2.1	16.3
10	Chronic lower respiratory diseases (J40-J47)	141	1.5	11.3
11	Septicemia (A40-A41)	139	1.4	11.2
12	Influenza and pneumonia (J10-J18)	132	1.4	10.6
13	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	130	1.4	10.4
14	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	127	1.3	10.2
15	Viral hepatitis (B15-B19)	86	0.9	6.9
...	All other causes (Residual)	1,523	15.8	122.3
Black, non-Hispanic, male, 50-54 years				
...	All causes	12,324	100.0	1,211.4
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,260	26.5	320.4
2	Malignant neoplasms (C00-C97)	2,690	21.8	264.4
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	943	7.7	92.7
4	Human immunodeficiency virus (HIV) disease (B20-B24)	777	6.3	76.4
5	Cerebrovascular diseases (I60-I69)	631	5.1	62.0
6	Diabetes mellitus (E10-E14)	443	3.6	43.5
7	Chronic liver disease and cirrhosis (K70,K73-K74)	331	2.7	32.5
8	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	249	2.0	24.5
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	235	1.9	23.1
10	Chronic lower respiratory diseases (J40-J47)	210	1.7	20.6
11	Septicemia (A40-A41)	205	1.7	20.2
12	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	180	1.5	17.7
13	Influenza and pneumonia (J10-J18)	167	1.4	16.4
14	Viral hepatitis (B15-B19)	148	1.2	14.5
15	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	108	0.9	10.6
...	All other causes (Residual)	1,747	14.2	171.7

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2005

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Black, non-Hispanic, male, 55-59 years				
...	All causes	13,447	100.0	1,701.2
1	Malignant neoplasms (C00-C97)	3,720	27.7	470.6
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,708	27.6	469.1
3	Cerebrovascular diseases (I60-I69)	681	5.1	86.2
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	642	4.8	81.2
5	Diabetes mellitus (E10-E14)	632	4.7	80.0
6	Human immunodeficiency virus (HIV) disease (B20-B24)	467	3.5	59.1
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	373	2.8	47.2
8	Chronic liver disease and cirrhosis (K70,K73-K74)	308	2.3	39.0
9	Chronic lower respiratory diseases (J40-J47)	303	2.3	38.3
10	Septicemia (A40-A41)	248	1.8	31.4
11	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	197	1.5	24.9
12	Influenza and pneumonia (J10-J18)	193	1.4	24.4
13	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	122	0.9	15.4
14	Viral hepatitis (B15-B19)	114	0.8	14.4
15	Aortic aneurysm and dissection (I71)	69	0.5	8.7
...	All other causes (Residual)	1,670	12.4	211.3
Black, non-Hispanic, male, 60-64 years				
...	All causes	12,949	100.0	2,427.3
1	Malignant neoplasms (C00-C97)	3,975	30.7	745.1
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,696	28.5	692.8
3	Cerebrovascular diseases (I60-I69)	730	5.6	136.8
4	Diabetes mellitus (E10-E14)	642	5.0	120.3
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	429	3.3	80.4
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	373	2.9	69.9
7	Chronic lower respiratory diseases (J40-J47)	357	2.8	66.9
8	Septicemia (A40-A41)	237	1.8	44.4
9	Human immunodeficiency virus (HIV) disease (B20-B24)	222	1.7	41.6
10	Chronic liver disease and cirrhosis (K70,K73-K74)	220	1.7	41.2
11	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	198	1.5	37.1
12	Influenza and pneumonia (J10-J18)	179	1.4	33.6
13	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	71	0.5	13.3
14	Aortic aneurysm and dissection (I71)	65	0.5	12.2
15	Viral hepatitis (B15-B19)	57	0.4	10.7
...	All other causes (Residual)	1,498	11.6	280.8

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2005

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Black, non-Hispanic, male, 65-69 years				
...	All causes	13,198	100.0	3,232.7
1	Malignant neoplasms (C00-C97)	4,265	32.3	1,044.7
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,646	27.6	893.1
3	Cerebrovascular diseases (I60-I69)	768	5.8	188.1
4	Diabetes mellitus (E10-E14)	673	5.1	164.8
5	Chronic lower respiratory diseases (J40-J47)	525	4.0	128.6
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	332	2.5	81.3
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	305	2.3	74.7
8	Septicemia (A40-A41)	282	2.1	69.1
9	Influenza and pneumonia (J10-J18)	224	1.7	54.9
10	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	218	1.7	53.4
11	Chronic liver disease and cirrhosis (K70,K73-K74)	127	1.0	31.1
12	Human immunodeficiency virus (HIV) disease (B20-B24)	123	0.9	30.1
13	Aortic aneurysm and dissection (I71)	79	0.6	19.4
14	Pneumonitis due to solids and liquids (J69)	63	0.5	15.4
15	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	53	0.4	13.0
...	All other causes (Residual)	1,515	11.5	371.1
Black, non-Hispanic, male, 70-74 years				
...	All causes	14,105	100.0	4,585.9
1	Malignant neoplasms (C00-C97)	4,365	30.9	1,419.2
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,940	27.9	1,281.0
3	Cerebrovascular diseases (I60-I69)	868	6.2	282.2
4	Diabetes mellitus (E10-E14)	740	5.2	240.6
5	Chronic lower respiratory diseases (J40-J47)	621	4.4	201.9
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	418	3.0	135.9
7	Septicemia (A40-A41)	304	2.2	98.8
8	Influenza and pneumonia (J10-J18)	286	2.0	93.0
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	284	2.0	92.3
10	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	247	1.8	80.3
11	Alzheimer's disease (G30)	114	0.8	37.1
12	Chronic liver disease and cirrhosis (K70,K73-K74)	98	0.7	31.9
13	Pneumonitis due to solids and liquids (J69)	80	0.6	26.0
14	Parkinson's disease (G20-G21)	59	0.4	19.2
15	Human immunodeficiency virus (HIV) disease (B20-B24)	56	0.4	18.2
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	56	0.4	18.2
...	All other causes (Residual)	1,569	11.1	510.1

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2005

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Black, non-Hispanic, male, 75-79 years				
...	All causes	14,530	100.0	6,672.3
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	4,149	28.6	1,905.3
2	Malignant neoplasms (C00-C97)	3,989	27.5	1,831.8
3	Cerebrovascular diseases (I60-I69)	899	6.2	412.8
4	Chronic lower respiratory diseases (J40-J47)	761	5.2	349.5
5	Diabetes mellitus (E10-E14)	687	4.7	315.5
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	464	3.2	213.1
7	Septicemia (A40-A41)	319	2.2	146.5
8	Influenza and pneumonia (J10-J18)	318	2.2	146.0
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	283	1.9	130.0
10	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	277	1.9	127.2
11	Alzheimer's disease (G30)	228	1.6	104.7
12	Pneumonitis due to solids and liquids (J69)	139	1.0	63.8
13	Parkinson's disease (G20-G21)	80	0.6	36.7
14	Aortic aneurysm and dissection (I71)	75	0.5	34.4
15	Chronic liver disease and cirrhosis (K70,K73-K74)	72	0.5	33.1
...	All other causes (Residual)	1,790	12.3	822.0
Black, non-Hispanic, male, 80-84 years				
...	All causes	13,193	100.0	9,492.7
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,803	28.8	2,736.4
2	Malignant neoplasms (C00-C97)	3,315	25.1	2,385.2
3	Cerebrovascular diseases (I60-I69)	922	7.0	663.4
4	Chronic lower respiratory diseases (J40-J47)	599	4.5	431.0
5	Diabetes mellitus (E10-E14)	549	4.2	395.0
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	427	3.2	307.2
7	Influenza and pneumonia (J10-J18)	365	2.8	262.6
8	Septicemia (A40-A41)	335	2.5	241.0
9	Alzheimer's disease (G30)	317	2.4	228.1
10	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	235	1.8	169.1
11	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	219	1.7	157.6
12	Pneumonitis due to solids and liquids (J69)	150	1.1	107.9
13	Parkinson's disease (G20-G21)	116	0.9	83.5
14	Aortic aneurysm and dissection (I71)	65	0.5	46.8
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	61	0.5	43.9
...	All other causes (Residual)	1,715	13.0	1,234.0

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2005

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Black, non-Hispanic, male, 85-89 years				
...	All causes	8,526	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,517	29.5	---
2	Malignant neoplasms (C00-C97)	1,850	21.7	---
3	Cerebrovascular diseases (I60-I69)	628	7.4	---
4	Chronic lower respiratory diseases (J40-J47)	382	4.5	---
5	Diabetes mellitus (E10-E14)	309	3.6	---
6	Alzheimer's disease (G30)	305	3.6	---
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	292	3.4	---
8	Influenza and pneumonia (J10-J18)	289	3.4	---
9	Septicemia (A40-A41)	249	2.9	---
10	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	168	2.0	---
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	133	1.6	---
12	Pneumonitis due to solids and liquids (J69)	117	1.4	---
13	Parkinson's disease (G20-G21)	68	0.8	---
14	Aortic aneurysm and dissection (I71)	39	0.5	---
15	Atherosclerosis (I70)	38	0.4	---
...	All other causes (Residual)	1,142	13.4	---
Black, non-Hispanic, male, 90-94 years				
...	All causes	4,367	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,388	31.8	---
2	Malignant neoplasms (C00-C97)	759	17.4	---
3	Cerebrovascular diseases (I60-I69)	318	7.3	---
4	Alzheimer's disease (G30)	187	4.3	---
5	Chronic lower respiratory diseases (J40-J47)	184	4.2	---
6	Influenza and pneumonia (J10-J18)	176	4.0	---
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	154	3.5	---
8	Diabetes mellitus (E10-E14)	142	3.3	---
9	Septicemia (A40-A41)	107	2.5	---
10	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	91	2.1	---
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	72	1.6	---
12	Pneumonitis due to solids and liquids (J69)	66	1.5	---
13	Atherosclerosis (I70)	29	0.7	---
14	Parkinson's disease (G20-G21)	26	0.6	---
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	14	0.3	---
15	Aortic aneurysm and dissection (I71)	14	0.3	---
...	All other causes (Residual)	640	14.7	---

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2005

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Black, non-Hispanic, male, 95-99 years				
...	All causes	1,466	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	467	31.9	---
2	Malignant neoplasms (C00-C97)	219	14.9	---
3	Cerebrovascular diseases (I60-I69)	86	5.9	---
4	Influenza and pneumonia (J10-J18)	74	5.0	---
5	Chronic lower respiratory diseases (J40-J47)	55	3.8	---
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	53	3.6	---
7	Alzheimer's disease (G30)	49	3.3	---
8	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	41	2.8	---
9	Diabetes mellitus (E10-E14)	39	2.7	---
10	Septicemia (A40-A41)	36	2.5	---
11	Pneumonitis due to solids and liquids (J69)	29	2.0	---
12	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	28	1.9	---
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	9	0.6	---
13	Atherosclerosis (I70)	9	0.6	---
15	Parkinson's disease (G20-G21)	8	0.5	---
...	All other causes (Residual)	264	18.0	---
Black, non-Hispanic, male, 100 years and over				
...	All causes	331	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	123	37.2	---
2	Malignant neoplasms (C00-C97)	31	9.4	---
3	Influenza and pneumonia (J10-J18)	26	7.9	---
4	Cerebrovascular diseases (I60-I69)	17	5.1	---
5	Alzheimer's disease (G30)	13	3.9	---
5	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	13	3.9	---
7	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	9	2.7	---
8	Septicemia (A40-A41)	7	2.1	---
9	Chronic lower respiratory diseases (J40-J47)	6	1.8	---
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	6	1.8	---
11	Diabetes mellitus (E10-E14)	5	1.5	---
11	Atherosclerosis (I70)	5	1.5	---
13	Pneumonitis due to solids and liquids (J69)	4	1.2	---
14	Hernia (K40-K46)	3	0.9	---
15	Nutritional deficiencies (E40-E64)	2	0.6	---
15	Parkinson's disease (G20-G21)	2	0.6	---
...	All other causes (Residual)	59	17.8	---

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2005

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Black, non-Hispanic, female, all ages				
...	All causes	142,153	100.0	727.6
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	37,428	26.3	191.6
2	Malignant neoplasms (C00-C97)	30,147	21.2	154.3
3	Cerebrovascular diseases (I60-I69)	9,930	7.0	50.8
4	Diabetes mellitus (E10-E14)	7,169	5.0	36.7
5	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	4,405	3.1	22.5
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	4,263	3.0	21.8
7	Chronic lower respiratory diseases (J40-J47)	3,730	2.6	19.1
8	Septicemia (A40-A41)	3,404	2.4	17.4
9	Alzheimer's disease (G30)	3,296	2.3	16.9
10	Influenza and pneumonia (J10-J18)	3,019	2.1	15.5
11	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	2,798	2.0	14.3
12	Human immunodeficiency virus (HIV) disease (B20-B24)	2,308	1.6	11.8
13	Certain conditions originating in the perinatal period (P00-P96)	2,125	1.5	10.9
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,234	0.9	6.3
15	Chronic liver disease and cirrhosis (K70,K73-K74)	861	0.6	4.4
...	All other causes (Residual)	26,036	18.3	133.3
Black, non-Hispanic, female, 1-4 years				
...	All causes	459	100.0	37.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	137	29.8	11.1
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	58	12.6	4.7
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	50	10.9	4.1
4	Malignant neoplasms (C00-C97)	26	5.7	2.1
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	26	5.7	2.1
6	Influenza and pneumonia (J10-J18)	12	2.6	*
7	Chronic lower respiratory diseases (J40-J47)	9	2.0	*
8	Septicemia (A40-A41)	7	1.5	*
8	Certain conditions originating in the perinatal period (P00-P96)	7	1.5	*
10	Anemias (D50-D64)	6	1.3	*
11	Cerebrovascular diseases (I60-I69)	4	0.9	*
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	3	0.7	*
12	Meningitis (G00,G03)	3	0.7	*
12	Pneumonitis due to solids and liquids (J69)	3	0.7	*
15	Complications of medical and surgical care (Y40-Y84,Y88)	2	0.4	*
...	All other causes (Residual)	106	23.1	8.6

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2005

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Black, non-Hispanic, female, 5-9 years				
...	All causes	291	100.0	20.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	103	35.4	7.1
2	Malignant neoplasms (C00-C97)	33	11.3	2.3
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	20	6.9	1.4
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	20	6.9	1.4
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	20	6.9	1.4
6	Chronic lower respiratory diseases (J40-J47)	10	3.4	*
7	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	5	1.7	*
7	Cerebrovascular diseases (I60-I69)	5	1.7	*
9	Influenza and pneumonia (J10-J18)	4	1.4	*
10	Anemias (D50-D64)	3	1.0	*
11	Septicemia (A40-A41)	2	0.7	*
11	Human immunodeficiency virus (HIV) disease (B20-B24)	2	0.7	*
11	Pneumonitis due to solids and liquids (J69)	2	0.7	*
11	Certain conditions originating in the perinatal period (P00-P96)	2	0.7	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1	0.3	*
...	All other causes (Residual)	59	20.3	4.1
Black, non-Hispanic, female, 10-14 years				
...	All causes	324	100.0	19.9
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	82	25.3	5.0
2	Malignant neoplasms (C00-C97)	34	10.5	2.1
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	32	9.9	2.0
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	23	7.1	1.4
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	20	6.2	1.2
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	14	4.3	*
7	Diabetes mellitus (E10-E14)	9	2.8	*
8	Chronic lower respiratory diseases (J40-J47)	8	2.5	*
9	Septicemia (A40-A41)	5	1.5	*
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	5	1.5	*
9	Influenza and pneumonia (J10-J18)	5	1.5	*
12	Cerebrovascular diseases (I60-I69)	4	1.2	*
13	Anemias (D50-D64)	3	0.9	*
14	Syphilis (A50-A53)	1	0.3	*
14	Human immunodeficiency virus (HIV) disease (B20-B24)	1	0.3	*
14	Meningitis (G00,G03)	1	0.3	*
14	Diseases of appendix (K35-K38)	1	0.3	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1	0.3	*
14	Certain conditions originating in the perinatal period (P00-P96)	1	0.3	*
14	Complications of medical and surgical care (Y40-Y84,Y88)	1	0.3	*
...	All other causes (Residual)	73	22.5	4.5

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2005

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Black, non-Hispanic, female, 15-19 years				
...	All causes	638	100.0	39.9
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	208	32.6	13.0
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	102	16.0	6.4
3	Malignant neoplasms (C00-C97)	45	7.1	2.8
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	31	4.9	1.9
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	23	3.6	1.4
6	Pregnancy, childbirth and the puerperium (000-099)	20	3.1	1.3
7	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	19	3.0	*
8	Human immunodeficiency virus (HIV) disease (B20-B24)	12	1.9	*
8	Anemias (D50-D64)	12	1.9	*
10	Cerebrovascular diseases (I60-I69)	11	1.7	*
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	8	1.3	*
12	Septicemia (A40-A41)	7	1.1	*
12	Diabetes mellitus (E10-E14)	7	1.1	*
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	5	0.8	*
14	Chronic lower respiratory diseases (J40-J47)	5	0.8	*
...	All other causes (Residual)	123	19.3	7.7
Black, non-Hispanic, female, 20-24 years				
...	All causes	991	100.0	66.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	225	22.7	15.0
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	178	18.0	11.9
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	74	7.5	4.9
4	Malignant neoplasms (C00-C97)	63	6.4	4.2
5	Pregnancy, childbirth and the puerperium (000-099)	49	4.9	3.3
6	Human immunodeficiency virus (HIV) disease (B20-B24)	39	3.9	2.6
7	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	30	3.0	2.0
8	Diabetes mellitus (E10-E14)	27	2.7	1.8
9	Chronic lower respiratory diseases (J40-J47)	22	2.2	1.5
10	Anemias (D50-D64)	20	2.0	1.3
11	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	17	1.7	*
12	Cerebrovascular diseases (I60-I69)	16	1.6	*
13	Septicemia (A40-A41)	13	1.3	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	10	1.0	*
15	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	4	0.4	*
15	Influenza and pneumonia (J10-J18)	4	0.4	*
...	All other causes (Residual)	200	20.2	13.4

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2005

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Black, non-Hispanic, female, 25-29 years				
...	All causes	1,291	100.0	92.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	184	14.3	13.2
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	143	11.1	10.2
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	143	11.1	10.2
4	Malignant neoplasms (C00-C97)	132	10.2	9.5
5	Human immunodeficiency virus (HIV) disease (B20-B24)	122	9.5	8.7
6	Pregnancy, childbirth and the puerperium (O00-099)	65	5.0	4.7
7	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	42	3.3	3.0
8	Diabetes mellitus (E10-E14)	39	3.0	2.8
9	Anemias (D50-D64)	31	2.4	2.2
10	Cerebrovascular diseases (I60-I69)	28	2.2	2.0
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	23	1.8	1.6
12	Septicemia (A40-A41)	21	1.6	1.5
13	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	17	1.3	*
14	Chronic lower respiratory diseases (J40-J47)	13	1.0	*
15	Influenza and pneumonia (J10-J18)	12	0.9	*
...	All other causes (Residual)	276	21.4	19.8
Black, non-Hispanic, female, 30-34 years				
...	All causes	1,853	100.0	136.3
1	Human immunodeficiency virus (HIV) disease (B20-B24)	247	13.3	18.2
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	241	13.0	17.7
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	238	12.8	17.5
4	Malignant neoplasms (C00-C97)	229	12.4	16.8
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	125	6.7	9.2
6	Diabetes mellitus (E10-E14)	52	2.8	3.8
7	Pregnancy, childbirth and the puerperium (O00-099)	50	2.7	3.7
8	Cerebrovascular diseases (I60-I69)	46	2.5	3.4
9	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	38	2.1	2.8
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	37	2.0	2.7
11	Septicemia (A40-A41)	36	1.9	2.6
12	Anemias (D50-D64)	31	1.7	2.3
12	Chronic lower respiratory diseases (J40-J47)	31	1.7	2.3
14	Influenza and pneumonia (J10-J18)	21	1.1	1.5
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	21	1.1	1.5
...	All other causes (Residual)	410	22.1	30.2

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2005

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Black, non-Hispanic, female, 35-39 years				
...	All causes	2,776	100.0	199.2
1	Malignant neoplasms (C00-C97)	485	17.5	34.8
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	416	15.0	29.9
3	Human immunodeficiency virus (HIV) disease (B20-B24)	367	13.2	26.3
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	268	9.7	19.2
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	148	5.3	10.6
6	Cerebrovascular diseases (I60-I69)	95	3.4	6.8
7	Diabetes mellitus (E10-E14)	81	2.9	5.8
8	Septicemia (A40-A41)	53	1.9	3.8
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	50	1.8	3.6
10	Pregnancy, childbirth and the puerperium (O00-099)	41	1.5	2.9
10	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	41	1.5	2.9
12	Chronic lower respiratory diseases (J40-J47)	35	1.3	2.5
13	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	29	1.0	2.1
14	Influenza and pneumonia (J10-J18)	27	1.0	1.9
15	Anemias (D50-D64)	26	0.9	1.9
...	All other causes (Residual)	614	22.1	44.1
Black, non-Hispanic, female, 40-44 years				
...	All causes	4,703	100.0	313.6
1	Malignant neoplasms (C00-C97)	1,076	22.9	71.7
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	780	16.6	52.0
3	Human immunodeficiency virus (HIV) disease (B20-B24)	477	10.1	31.8
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	372	7.9	24.8
5	Cerebrovascular diseases (I60-I69)	259	5.5	17.3
6	Diabetes mellitus (E10-E14)	146	3.1	9.7
7	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	128	2.7	8.5
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	98	2.1	6.5
9	Septicemia (A40-A41)	84	1.8	5.6
10	Chronic lower respiratory diseases (J40-J47)	79	1.7	5.3
11	Chronic liver disease and cirrhosis (K70,K73-K74)	70	1.5	4.7
12	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	62	1.3	4.1
13	Influenza and pneumonia (J10-J18)	60	1.3	4.0
14	Anemias (D50-D64)	41	0.9	2.7
14	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	41	0.9	2.7
...	All other causes (Residual)	930	19.8	62.0

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2005

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Black, non-Hispanic, female, 45-49 years				
...	All causes	6,882	100.0	481.1
1	Malignant neoplasms (C00-C97)	1,921	27.9	134.3
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,274	18.5	89.1
3	Human immunodeficiency virus (HIV) disease (B20-B24)	447	6.5	31.2
4	Cerebrovascular diseases (I60-I69)	430	6.2	30.1
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	395	5.7	27.6
6	Diabetes mellitus (E10-E14)	238	3.5	16.6
7	Chronic lower respiratory diseases (J40-J47)	148	2.2	10.3
8	Septicemia (A40-A41)	136	2.0	9.5
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	130	1.9	9.1
10	Chronic liver disease and cirrhosis (K70,K73-K74)	124	1.8	8.7
11	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	109	1.6	7.6
12	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	96	1.4	6.7
13	Influenza and pneumonia (J10-J18)	84	1.2	5.9
14	Viral hepatitis (B15-B19)	44	0.6	3.1
15	Anemias (D50-D64)	38	0.6	2.7
...	All other causes (Residual)	1,268	18.4	88.6
Black, non-Hispanic, female, 50-54 years				
...	All causes	8,470	100.0	702.6
1	Malignant neoplasms (C00-C97)	2,575	30.4	213.6
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,842	21.7	152.8
3	Cerebrovascular diseases (I60-I69)	517	6.1	42.9
4	Diabetes mellitus (E10-E14)	386	4.6	32.0
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	355	4.2	29.4
6	Human immunodeficiency virus (HIV) disease (B20-B24)	300	3.5	24.9
7	Chronic lower respiratory diseases (J40-J47)	198	2.3	16.4
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	177	2.1	14.7
9	Septicemia (A40-A41)	174	2.1	14.4
10	Chronic liver disease and cirrhosis (K70,K73-K74)	161	1.9	13.4
11	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	111	1.3	9.2
11	Influenza and pneumonia (J10-J18)	111	1.3	9.2
13	Viral hepatitis (B15-B19)	71	0.8	5.9
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	65	0.8	5.4
15	Anemias (D50-D64)	39	0.5	3.2
...	All other causes (Residual)	1,388	16.4	115.1

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2005

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Black, non-Hispanic, female, 55-59 years				
...	All causes	8,970	100.0	935.2
1	Malignant neoplasms (C00-C97)	2,980	33.2	310.7
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,124	23.7	221.5
3	Diabetes mellitus (E10-E14)	509	5.7	53.1
4	Cerebrovascular diseases (I60-I69)	497	5.5	51.8
5	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	271	3.0	28.3
6	Chronic lower respiratory diseases (J40-J47)	242	2.7	25.2
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	235	2.6	24.5
8	Septicemia (A40-A41)	207	2.3	21.6
9	Human immunodeficiency virus (HIV) disease (B20-B24)	138	1.5	14.4
10	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	131	1.5	13.7
11	Chronic liver disease and cirrhosis (K70,K73-K74)	128	1.4	13.3
12	Influenza and pneumonia (J10-J18)	108	1.2	11.3
13	Viral hepatitis (B15-B19)	49	0.5	5.1
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	36	0.4	3.8
15	Anemias (D50-D64)	29	0.3	3.0
15	Aortic aneurysm and dissection (I71)	29	0.3	3.0
...	All other causes (Residual)	1,257	14.0	131.1
Black, non-Hispanic, female, 60-64 years				
...	All causes	9,571	100.0	1,395.5
1	Malignant neoplasms (C00-C97)	3,160	33.0	460.7
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,449	25.6	357.1
3	Diabetes mellitus (E10-E14)	632	6.6	92.1
4	Cerebrovascular diseases (I60-I69)	512	5.3	74.7
5	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	323	3.4	47.1
6	Chronic lower respiratory diseases (J40-J47)	293	3.1	42.7
7	Septicemia (A40-A41)	242	2.5	35.3
8	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	158	1.7	23.0
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	135	1.4	19.7
10	Influenza and pneumonia (J10-J18)	123	1.3	17.9
11	Chronic liver disease and cirrhosis (K70,K73-K74)	95	1.0	13.9
12	Human immunodeficiency virus (HIV) disease (B20-B24)	75	0.8	10.9
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	44	0.5	6.4
13	Aortic aneurysm and dissection (I71)	44	0.5	6.4
15	Viral hepatitis (B15-B19)	33	0.3	4.8
...	All other causes (Residual)	1,253	13.1	182.7

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2005

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Black, non-Hispanic, female, 65-69 years				
...	All causes	10,900	100.0	1,977.0
1	Malignant neoplasms (C00-C97)	3,391	31.1	615.0
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,814	25.8	510.4
3	Diabetes mellitus (E10-E14)	800	7.3	145.1
4	Cerebrovascular diseases (I60-I69)	644	5.9	116.8
5	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	450	4.1	81.6
6	Chronic lower respiratory diseases (J40-J47)	373	3.4	67.7
7	Septicemia (A40-A41)	263	2.4	47.7
8	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	217	2.0	39.4
9	Influenza and pneumonia (J10-J18)	160	1.5	29.0
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	160	1.5	29.0
11	Chronic liver disease and cirrhosis (K70,K73-K74)	68	0.6	12.3
12	Alzheimer's disease (G30)	53	0.5	9.6
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	52	0.5	9.4
13	Aortic aneurysm and dissection (I71)	52	0.5	9.4
15	Human immunodeficiency virus (HIV) disease (B20-B24)	38	0.3	6.9
15	Pneumonitis due to solids and liquids (J69)	38	0.3	6.9
...	All other causes (Residual)	1,327	12.2	240.7
Black, non-Hispanic, female, 70-74 years				
...	All causes	12,963	100.0	2,880.6
1	Malignant neoplasms (C00-C97)	3,539	27.3	786.4
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,440	26.5	764.4
3	Cerebrovascular diseases (I60-I69)	914	7.1	203.1
4	Diabetes mellitus (E10-E14)	888	6.9	197.3
5	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	544	4.2	120.9
6	Chronic lower respiratory diseases (J40-J47)	479	3.7	106.4
7	Septicemia (A40-A41)	322	2.5	71.6
8	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	266	2.1	59.1
9	Influenza and pneumonia (J10-J18)	244	1.9	54.2
10	Alzheimer's disease (G30)	174	1.3	38.7
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	172	1.3	38.2
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	71	0.5	15.8
13	Aortic aneurysm and dissection (I71)	70	0.5	15.6
14	Chronic liver disease and cirrhosis (K70,K73-K74)	59	0.5	13.1
15	Pneumonitis due to solids and liquids (J69)	51	0.4	11.3
...	All other causes (Residual)	1,730	13.3	384.4

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2005

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Black, non-Hispanic, female, 75-79 years				
...	All causes	16,033	100.0	4,382.4
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	4,695	29.3	1,283.3
2	Malignant neoplasms (C00-C97)	3,699	23.1	1,011.1
3	Cerebrovascular diseases (I60-I69)	1,302	8.1	355.9
4	Diabetes mellitus (E10-E14)	986	6.1	269.5
5	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	586	3.7	160.2
6	Chronic lower respiratory diseases (J40-J47)	544	3.4	148.7
7	Septicemia (A40-A41)	428	2.7	117.0
8	Influenza and pneumonia (J10-J18)	366	2.3	100.0
9	Alzheimer's disease (G30)	365	2.3	99.8
10	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	323	2.0	88.3
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	228	1.4	62.3
12	Pneumonitis due to solids and liquids (J69)	105	0.7	28.7
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	80	0.5	21.9
14	Aortic aneurysm and dissection (I71)	77	0.5	21.0
15	Atherosclerosis (I70)	60	0.4	16.4
...	All other causes (Residual)	2,189	13.7	598.3
Black, non-Hispanic, female, 80-84 years				
...	All causes	17,563	100.0	6,664.8
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	5,471	31.2	2,076.1
2	Malignant neoplasms (C00-C97)	3,153	18.0	1,196.5
3	Cerebrovascular diseases (I60-I69)	1,568	8.9	595.0
4	Diabetes mellitus (E10-E14)	983	5.6	373.0
5	Alzheimer's disease (G30)	669	3.8	253.9
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	613	3.5	232.6
7	Chronic lower respiratory diseases (J40-J47)	510	2.9	193.5
8	Septicemia (A40-A41)	495	2.8	187.8
9	Influenza and pneumonia (J10-J18)	448	2.6	170.0
10	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	407	2.3	154.4
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	218	1.2	82.7
12	Pneumonitis due to solids and liquids (J69)	131	0.7	49.7
13	Aortic aneurysm and dissection (I71)	90	0.5	34.2
14	Parkinson's disease (G20-G21)	89	0.5	33.8
15	Atherosclerosis (I70)	82	0.5	31.1
...	All other causes (Residual)	2,636	15.0	1,000.3

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2005

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Black, non-Hispanic, female, 85-89 years				
...	All causes	15,246	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	4,967	32.6	---
2	Malignant neoplasms (C00-C97)	2,099	13.8	---
3	Cerebrovascular diseases (I60-I69)	1,410	9.2	---
4	Alzheimer's disease (G30)	789	5.2	---
5	Diabetes mellitus (E10-E14)	733	4.8	---
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	534	3.5	---
7	Influenza and pneumonia (J10-J18)	485	3.2	---
8	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	397	2.6	---
9	Septicemia (A40-A41)	375	2.5	---
10	Chronic lower respiratory diseases (J40-J47)	373	2.4	---
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	179	1.2	---
12	Pneumonitis due to solids and liquids (J69)	146	1.0	---
13	Atherosclerosis (I70)	109	0.7	---
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	80	0.5	---
15	Aortic aneurysm and dissection (I71)	71	0.5	---
...	All other causes (Residual)	2,499	16.4	---
Black, non-Hispanic, female, 90-94 years				
...	All causes	11,484	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,920	34.1	---
2	Cerebrovascular diseases (I60-I69)	1,082	9.4	---
3	Malignant neoplasms (C00-C97)	1,076	9.4	---
4	Alzheimer's disease (G30)	728	6.3	---
5	Diabetes mellitus (E10-E14)	464	4.0	---
6	Influenza and pneumonia (J10-J18)	410	3.6	---
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	346	3.0	---
8	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	340	3.0	---
9	Septicemia (A40-A41)	298	2.6	---
10	Chronic lower respiratory diseases (J40-J47)	222	1.9	---
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	145	1.3	---
12	Pneumonitis due to solids and liquids (J69)	125	1.1	---
13	Atherosclerosis (I70)	93	0.8	---
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	44	0.4	---
14	Aortic aneurysm and dissection (I71)	44	0.4	---
...	All other causes (Residual)	2,147	18.7	---

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2005

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Black, non-Hispanic, female, 95-99 years				
...	All causes	5,270	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,950	37.0	---
2	Cerebrovascular diseases (I60-I69)	422	8.0	---
3	Alzheimer's disease (G30)	366	6.9	---
4	Malignant neoplasms (C00-C97)	359	6.8	---
5	Influenza and pneumonia (J10-J18)	196	3.7	---
6	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	156	3.0	---
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	148	2.8	---
8	Diabetes mellitus (E10-E14)	146	2.8	---
9	Septicemia (A40-A41)	136	2.6	---
10	Chronic lower respiratory diseases (J40-J47)	102	1.9	---
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	67	1.3	---
12	Pneumonitis due to solids and liquids (J69)	62	1.2	---
13	Atherosclerosis (I70)	54	1.0	---
14	Nutritional deficiencies (E40-E64)	29	0.6	---
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	17	0.3	---
15	Anemias (D50-D64)	17	0.3	---
15	Aortic aneurysm and dissection (I71)	17	0.3	---
...	All other causes (Residual)	1,026	19.5	---
Black, non-Hispanic, female, 100 years and over				
...	All causes	1,791	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	684	38.2	---
2	Cerebrovascular diseases (I60-I69)	147	8.2	---
3	Alzheimer's disease (G30)	122	6.8	---
4	Influenza and pneumonia (J10-J18)	85	4.7	---
5	Malignant neoplasms (C00-C97)	68	3.8	---
6	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	62	3.5	---
7	Diabetes mellitus (E10-E14)	42	2.3	---
8	Septicemia (A40-A41)	40	2.2	---
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	34	1.9	---
10	Atherosclerosis (I70)	28	1.6	---
10	Chronic lower respiratory diseases (J40-J47)	28	1.6	---
12	Pneumonitis due to solids and liquids (J69)	20	1.1	---
13	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	17	0.9	---
14	Nutritional deficiencies (E40-E64)	9	0.5	---
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	4	0.2	---
15	Anemias (D50-D64)	4	0.2	---
...	All other causes (Residual)	397	22.2	---

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available