

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, both sexes, all ages				
...	All causes	149,635	100.0	287.5
1	Malignant neoplasms (C00-C97)	32,381	21.6	62.2
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	30,385	20.3	58.4
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	11,166	7.5	21.5
4	Cerebrovascular diseases (I60-I69)	7,552	5.0	14.5
5	Diabetes mellitus (E10-E14)	7,054	4.7	13.6
6	Chronic liver disease and cirrhosis (K70,K73-K74)	4,774	3.2	9.2
7	Chronic lower respiratory diseases (J40-J47)	4,355	2.9	8.4
8	Alzheimer's disease (G30)	3,599	2.4	6.9
9	Influenza and pneumonia (J09-J18)	3,256	2.2	6.3
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2,799	1.9	5.4
11	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	2,759	1.8	5.3
12	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,720	1.8	5.2
13	Certain conditions originating in the perinatal period (P00-P96)	2,387	1.6	4.6
14	Septicemia (A40-A41)	2,168	1.4	4.2
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	1,803	1.2	3.5
...	All other causes (Residual)	30,477	20.4	58.6
Hispanic, both sexes, 1-4 years				
...	All causes	972	100.0	23.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	317	32.6	7.7
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	126	13.0	3.0
3	Malignant neoplasms (C00-C97)	97	10.0	2.3
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	94	9.7	2.3
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	35	3.6	0.8
6	Influenza and pneumonia (J09-J18)	21	2.2	0.5
7	Septicemia (A40-A41)	16	1.6	*
8	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	10	1.0	*
8	Cerebrovascular diseases (I60-I69)	10	1.0	*
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	7	0.7	*
11	Complications of medical and surgical care (Y40-Y84,Y88)	6	0.6	*
12	Certain conditions originating in the perinatal period (P00-P96)	5	0.5	*
13	Acute bronchitis and bronchiolitis (J20-J21)	4	0.4	*
13	Chronic lower respiratory diseases (J40-J47)	4	0.4	*
15	Meningitis (G00,G03)	3	0.3	*
...	All other causes (Residual)	217	22.3	5.2

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, both sexes, 5-9 years				
...	All causes	490	100.0	10.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	145	29.6	3.0
2	Malignant neoplasms (C00-C97)	100	20.4	2.0
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	48	9.8	1.0
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	25	5.1	0.5
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	22	4.5	0.4
6	Influenza and pneumonia (J09-J18)	15	3.1	*
7	Cerebrovascular diseases (I60-I69)	11	2.2	*
8	Chronic lower respiratory diseases (J40-J47)	9	1.8	*
9	Septicemia (A40-A41)	6	1.2	*
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	4	0.8	*
11	Certain conditions originating in the perinatal period (P00-P96)	3	0.6	*
12	Meningitis (G00,G03)	2	0.4	*
12	Diseases of appendix (K35-K38)	2	0.4	*
12	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2	0.4	*
12	Complications of medical and surgical care (Y40-Y84,Y88)	2	0.4	*
...	All other causes (Residual)	94	19.2	1.9
Hispanic, both sexes, 10-14 years				
...	All causes	562	100.0	12.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	133	23.7	2.9
2	Malignant neoplasms (C00-C97)	100	17.8	2.2
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	56	10.0	1.2
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	44	7.8	1.0
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	32	5.7	0.7
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	20	3.6	0.4
7	Chronic lower respiratory diseases (J40-J47)	14	2.5	*
8	Cerebrovascular diseases (I60-I69)	11	2.0	*
9	Influenza and pneumonia (J09-J18)	7	1.2	*
10	Diabetes mellitus (E10-E14)	6	1.1	*
11	Anemias (D50-D64)	5	0.9	*
12	Septicemia (A40-A41)	4	0.7	*
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	3	0.5	*
14	Meningococcal infection (A39)	2	0.4	*
14	Meningitis (G00,G03)	2	0.4	*
14	Pneumonitis due to solids and liquids (J69)	2	0.4	*
14	Certain conditions originating in the perinatal period (P00-P96)	2	0.4	*
...	All other causes (Residual)	119	21.2	2.6

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, both sexes, 15-19 years				
...	All causes	1,836	100.0	40.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	664	36.2	14.5
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	381	20.8	8.3
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	249	13.6	5.4
4	Malignant neoplasms (C00-C97)	172	9.4	3.8
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	50	2.7	1.1
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	30	1.6	0.7
7	Cerebrovascular diseases (I60-I69)	19	1.0	*
8	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	9	0.5	*
8	Pregnancy, childbirth and the puerperium (O00-O99)	9	0.5	*
8	Legal intervention (Y35,Y89.0)	9	0.5	*
11	Anemias (D50-D64)	8	0.4	*
11	Diabetes mellitus (E10-E14)	8	0.4	*
13	Septicemia (A40-A41)	7	0.4	*
13	Chronic lower respiratory diseases (J40-J47)	7	0.4	*
15	Influenza and pneumonia (J09-J18)	6	0.3	*
...	All other causes (Residual)	208	11.3	4.5
Hispanic, both sexes, 20-24 years				
...	All causes	2,990	100.0	66.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,247	41.7	27.8
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	552	18.5	12.3
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	386	12.9	8.6
4	Malignant neoplasms (C00-C97)	211	7.1	4.7
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	106	3.5	2.4
6	Cerebrovascular diseases (I60-I69)	34	1.1	0.8
7	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	26	0.9	0.6
8	Legal intervention (Y35,Y89.0)	25	0.8	0.6
9	Influenza and pneumonia (J09-J18)	22	0.7	0.5
9	Pregnancy, childbirth and the puerperium (O00-O99)	22	0.7	0.5
11	Human immunodeficiency virus (HIV) disease (B20-B24)	19	0.6	*
12	Diabetes mellitus (E10-E14)	11	0.4	*
13	Chronic lower respiratory diseases (J40-J47)	10	0.3	*
14	Septicemia (A40-A41)	9	0.3	*
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	9	0.3	*
...	All other causes (Residual)	301	10.1	6.7

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, both sexes, 25-29 years				
...	All causes	2,862	100.0	65.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,067	37.3	24.4
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	437	15.3	10.0
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	342	11.9	7.8
4	Malignant neoplasms (C00-C97)	225	7.9	5.2
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	142	5.0	3.3
6	Cerebrovascular diseases (I60-I69)	40	1.4	0.9
7	Influenza and pneumonia (J09-J18)	33	1.2	0.8
8	Human immunodeficiency virus (HIV) disease (B20-B24)	32	1.1	0.7
9	Pregnancy, childbirth and the puerperium (O00-O99)	30	1.0	0.7
9	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	30	1.0	0.7
11	Diabetes mellitus (E10-E14)	29	1.0	0.7
12	Chronic liver disease and cirrhosis (K70,K73-K74)	26	0.9	0.6
13	Septicemia (A40-A41)	18	0.6	*
13	Legal intervention (Y35,Y89.0)	18	0.6	*
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	11	0.4	*
...	All other causes (Residual)	382	13.3	8.7
Hispanic, both sexes, 30-34 years				
...	All causes	3,199	100.0	75.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,029	32.2	24.3
2	Malignant neoplasms (C00-C97)	370	11.6	8.7
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	345	10.8	8.1
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	312	9.8	7.4
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	244	7.6	5.8
6	Human immunodeficiency virus (HIV) disease (B20-B24)	87	2.7	2.1
7	Chronic liver disease and cirrhosis (K70,K73-K74)	80	2.5	1.9
8	Cerebrovascular diseases (I60-I69)	70	2.2	1.7
9	Diabetes mellitus (E10-E14)	49	1.5	1.2
10	Influenza and pneumonia (J09-J18)	46	1.4	1.1
11	Septicemia (A40-A41)	33	1.0	0.8
12	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	26	0.8	0.6
13	Pregnancy, childbirth and the puerperium (O00-O99)	25	0.8	0.6
14	Legal intervention (Y35,Y89.0)	16	0.5	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	15	0.5	*
...	All other causes (Residual)	452	14.1	10.7

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, both sexes, 35-39 years				
...	All causes	3,556	100.0	90.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	905	25.4	23.0
2	Malignant neoplasms (C00-C97)	585	16.5	14.9
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	345	9.7	8.8
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	283	8.0	7.2
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	254	7.1	6.5
6	Chronic liver disease and cirrhosis (K70,K73-K74)	171	4.8	4.3
7	Human immunodeficiency virus (HIV) disease (B20-B24)	112	3.1	2.8
8	Cerebrovascular diseases (I60-I69)	88	2.5	2.2
9	Diabetes mellitus (E10-E14)	86	2.4	2.2
10	Influenza and pneumonia (J09-J18)	40	1.1	1.0
11	Septicemia (A40-A41)	37	1.0	0.9
12	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	27	0.8	0.7
13	Pregnancy, childbirth and the puerperium (O00-O99)	23	0.6	0.6
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	20	0.6	0.5
15	Chronic lower respiratory diseases (J40-J47)	18	0.5	*
...	All other causes (Residual)	562	15.8	14.3
Hispanic, both sexes, 40-44 years				
...	All causes	4,636	100.0	129.7
1	Malignant neoplasms (C00-C97)	924	19.9	25.8
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	902	19.5	25.2
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	600	12.9	16.8
4	Chronic liver disease and cirrhosis (K70,K73-K74)	298	6.4	8.3
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	240	5.2	6.7
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	190	4.1	5.3
7	Cerebrovascular diseases (I60-I69)	167	3.6	4.7
8	Human immunodeficiency virus (HIV) disease (B20-B24)	159	3.4	4.4
9	Diabetes mellitus (E10-E14)	141	3.0	3.9
10	Influenza and pneumonia (J09-J18)	68	1.5	1.9
11	Septicemia (A40-A41)	57	1.2	1.6
12	Viral hepatitis (B15-B19)	51	1.1	1.4
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	49	1.1	1.4
14	Chronic lower respiratory diseases (J40-J47)	33	0.7	0.9
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	31	0.7	0.9
...	All other causes (Residual)	726	15.7	20.3

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, both sexes, 45-49 years				
...	All causes	6,759	100.0	215.4
1	Malignant neoplasms (C00-C97)	1,474	21.8	47.0
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,049	15.5	33.4
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	928	13.7	29.6
4	Chronic liver disease and cirrhosis (K70,K73-K74)	603	8.9	19.2
5	Cerebrovascular diseases (I60-I69)	296	4.4	9.4
6	Diabetes mellitus (E10-E14)	276	4.1	8.8
7	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	220	3.3	7.0
8	Human immunodeficiency virus (HIV) disease (B20-B24)	207	3.1	6.6
9	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	151	2.2	4.8
10	Viral hepatitis (B15-B19)	148	2.2	4.7
11	Septicemia (A40-A41)	101	1.5	3.2
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	90	1.3	2.9
13	Influenza and pneumonia (J09-J18)	87	1.3	2.8
14	Chronic lower respiratory diseases (J40-J47)	55	0.8	1.8
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	52	0.8	1.7
...	All other causes (Residual)	1,022	15.1	32.6
Hispanic, both sexes, 50-54 years				
...	All causes	8,790	100.0	340.9
1	Malignant neoplasms (C00-C97)	2,397	27.3	93.0
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,482	16.9	57.5
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	833	9.5	32.3
4	Chronic liver disease and cirrhosis (K70,K73-K74)	763	8.7	29.6
5	Diabetes mellitus (E10-E14)	395	4.5	15.3
6	Cerebrovascular diseases (I60-I69)	373	4.2	14.5
7	Viral hepatitis (B15-B19)	220	2.5	8.5
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	203	2.3	7.9
9	Human immunodeficiency virus (HIV) disease (B20-B24)	189	2.2	7.3
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	141	1.6	5.5
11	Influenza and pneumonia (J09-J18)	117	1.3	4.5
12	Septicemia (A40-A41)	116	1.3	4.5
13	Chronic lower respiratory diseases (J40-J47)	97	1.1	3.8
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	84	1.0	3.3
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	79	0.9	3.1
...	All other causes (Residual)	1,301	14.8	50.5

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, both sexes, 55-59 years				
...	All causes	10,011	100.0	506.9
1	Malignant neoplasms (C00-C97)	3,055	30.5	154.7
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,913	19.1	96.9
3	Chronic liver disease and cirrhosis (K70,K73-K74)	764	7.6	38.7
4	Diabetes mellitus (E10-E14)	584	5.8	29.6
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	569	5.7	28.8
6	Cerebrovascular diseases (I60-I69)	432	4.3	21.9
7	Viral hepatitis (B15-B19)	276	2.8	14.0
8	Septicemia (A40-A41)	180	1.8	9.1
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	171	1.7	8.7
10	Chronic lower respiratory diseases (J40-J47)	166	1.7	8.4
11	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	136	1.4	6.9
12	Human immunodeficiency virus (HIV) disease (B20-B24)	125	1.2	6.3
13	Influenza and pneumonia (J09-J18)	116	1.2	5.9
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	82	0.8	4.2
15	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	63	0.6	3.2
...	All other causes (Residual)	1,379	13.8	69.8
Hispanic, both sexes, 60-64 years				
...	All causes	10,947	100.0	738.3
1	Malignant neoplasms (C00-C97)	3,571	32.6	240.9
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,226	20.3	150.1
3	Diabetes mellitus (E10-E14)	683	6.2	46.1
4	Chronic liver disease and cirrhosis (K70,K73-K74)	583	5.3	39.3
5	Cerebrovascular diseases (I60-I69)	529	4.8	35.7
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	448	4.1	30.2
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	241	2.2	16.3
8	Chronic lower respiratory diseases (J40-J47)	224	2.0	15.1
9	Viral hepatitis (B15-B19)	197	1.8	13.3
10	Septicemia (A40-A41)	183	1.7	12.3
11	Influenza and pneumonia (J09-J18)	156	1.4	10.5
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	94	0.9	6.3
13	Human immunodeficiency virus (HIV) disease (B20-B24)	90	0.8	6.1
13	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	90	0.8	6.1
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	66	0.6	4.5
...	All other causes (Residual)	1,566	14.3	105.6

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, both sexes, 65-69 years				
...	All causes	11,353	100.0	1,114.1
1	Malignant neoplasms (C00-C97)	3,761	33.1	369.1
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,428	21.4	238.3
3	Diabetes mellitus (E10-E14)	803	7.1	78.8
4	Cerebrovascular diseases (I60-I69)	516	4.5	50.6
5	Chronic liver disease and cirrhosis (K70,K73-K74)	462	4.1	45.3
6	Chronic lower respiratory diseases (J40-J47)	320	2.8	31.4
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	319	2.8	31.3
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	273	2.4	26.8
9	Influenza and pneumonia (J09-J18)	195	1.7	19.1
10	Septicemia (A40-A41)	191	1.7	18.7
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	111	1.0	10.9
12	Viral hepatitis (B15-B19)	100	0.9	9.8
13	Alzheimer's disease (G30)	84	0.7	8.2
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	58	0.5	5.7
15	Pneumonitis due to solids and liquids (J69)	51	0.4	5.0
15	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	51	0.4	5.0
...	All other causes (Residual)	1,630	14.4	160.0
Hispanic, both sexes, 70-74 years				
...	All causes	12,631	100.0	1,715.1
1	Malignant neoplasms (C00-C97)	3,766	29.8	511.4
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,778	22.0	377.2
3	Diabetes mellitus (E10-E14)	898	7.1	121.9
4	Cerebrovascular diseases (I60-I69)	690	5.5	93.7
5	Chronic lower respiratory diseases (J40-J47)	479	3.8	65.0
6	Chronic liver disease and cirrhosis (K70,K73-K74)	367	2.9	49.8
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	334	2.6	45.4
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	272	2.2	36.9
9	Influenza and pneumonia (J09-J18)	254	2.0	34.5
10	Septicemia (A40-A41)	226	1.8	30.7
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	161	1.3	21.9
12	Alzheimer's disease (G30)	135	1.1	18.3
13	Parkinson's disease (G20-G21)	116	0.9	15.8
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	80	0.6	10.9
15	Viral hepatitis (B15-B19)	69	0.5	9.4
...	All other causes (Residual)	2,006	15.9	272.4

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, both sexes, 75-79 years				
...	All causes	14,921	100.0	2,788.7
1	Malignant neoplasms (C00-C97)	3,935	26.4	735.4
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,500	23.5	654.1
3	Cerebrovascular diseases (I60-I69)	979	6.6	183.0
4	Diabetes mellitus (E10-E14)	945	6.3	176.6
5	Chronic lower respiratory diseases (J40-J47)	654	4.4	122.2
6	Alzheimer's disease (G30)	397	2.7	74.2
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	363	2.4	67.8
8	Influenza and pneumonia (J09-J18)	362	2.4	67.7
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	295	2.0	55.1
10	Chronic liver disease and cirrhosis (K70,K73-K74)	281	1.9	52.5
11	Septicemia (A40-A41)	242	1.6	45.2
12	Parkinson's disease (G20-G21)	197	1.3	36.8
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	191	1.3	35.7
14	Pneumonitis due to solids and liquids (J69)	99	0.7	18.5
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	97	0.7	18.1
...	All other causes (Residual)	2,384	16.0	445.6
Hispanic, both sexes, 80-84 years				
...	All causes	17,700	100.0	4,723.1
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	4,445	25.1	1,186.1
2	Malignant neoplasms (C00-C97)	3,795	21.4	1,012.7
3	Cerebrovascular diseases (I60-I69)	1,192	6.7	318.1
4	Diabetes mellitus (E10-E14)	972	5.5	259.4
5	Chronic lower respiratory diseases (J40-J47)	806	4.6	215.1
6	Alzheimer's disease (G30)	763	4.3	203.6
7	Influenza and pneumonia (J09-J18)	521	2.9	139.0
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	407	2.3	108.6
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	355	2.0	94.7
10	Septicemia (A40-A41)	293	1.7	78.2
11	Parkinson's disease (G20-G21)	279	1.6	74.4
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	274	1.5	73.1
13	Chronic liver disease and cirrhosis (K70,K73-K74)	229	1.3	61.1
14	Pneumonitis due to solids and liquids (J69)	127	0.7	33.9
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	119	0.7	31.8
...	All other causes (Residual)	3,123	17.6	833.3

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, both sexes, 85-89 years				
...	All causes	16,172	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	4,508	27.9	---
2	Malignant neoplasms (C00-C97)	2,520	15.6	---
3	Cerebrovascular diseases (I60-I69)	1,151	7.1	---
4	Alzheimer's disease (G30)	998	6.2	---
5	Chronic lower respiratory diseases (J40-J47)	821	5.1	---
6	Diabetes mellitus (E10-E14)	710	4.4	---
7	Influenza and pneumonia (J09-J18)	549	3.4	---
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	385	2.4	---
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	290	1.8	---
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	273	1.7	---
11	Septicemia (A40-A41)	251	1.6	---
12	Parkinson's disease (G20-G21)	249	1.5	---
13	Pneumonitis due to solids and liquids (J69)	169	1.0	---
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	101	0.6	---
15	Chronic liver disease and cirrhosis (K70,K73-K74)	90	0.6	---
...	All other causes (Residual)	3,107	19.2	---
Hispanic, both sexes, 90-94 years				
...	All causes	9,716	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,897	29.8	---
2	Malignant neoplasms (C00-C97)	1,005	10.3	---
3	Alzheimer's disease (G30)	745	7.7	---
4	Cerebrovascular diseases (I60-I69)	646	6.6	---
5	Chronic lower respiratory diseases (J40-J47)	431	4.4	---
6	Influenza and pneumonia (J09-J18)	378	3.9	---
7	Diabetes mellitus (E10-E14)	362	3.7	---
8	Essential hypertension and hypertensive renal disease (I10,I12,I15)	206	2.1	---
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	194	2.0	---
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	193	2.0	---
11	Parkinson's disease (G20-G21)	121	1.2	---
12	Septicemia (A40-A41)	112	1.2	---
13	Pneumonitis due to solids and liquids (J69)	85	0.9	---
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	53	0.5	---
15	Enterocolitis due to <i>Clostridium difficile</i> (A04.7)	44	0.5	---
...	All other causes (Residual)	2,244	23.1	---

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, both sexes, 95-99 years				
...	All causes	3,751	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,194	31.8	---
2	Alzheimer's disease (G30)	319	8.5	---
3	Malignant neoplasms (C00-C97)	262	7.0	---
4	Cerebrovascular diseases (I60-I69)	227	6.1	---
5	Influenza and pneumonia (J09-J18)	186	5.0	---
6	Chronic lower respiratory diseases (J40-J47)	153	4.1	---
7	Essential hypertension and hypertensive renal disease (I10,I12,I15)	91	2.4	---
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	79	2.1	---
9	Diabetes mellitus (E10-E14)	78	2.1	---
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	62	1.7	---
11	Septicemia (A40-A41)	46	1.2	---
12	Pneumonitis due to solids and liquids (J69)	35	0.9	---
13	Atherosclerosis (I70)	22	0.6	---
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	16	0.4	---
15	Parkinson's disease (G20-G21)	14	0.4	---
15	Cholelithiasis and other disorders of gallbladder (K80-K82)	14	0.4	---
...	All other causes (Residual)	953	25.4	---
Hispanic, both sexes, 100 years and over				
...	All causes	949	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	333	35.1	---
2	Alzheimer's disease (G30)	83	8.7	---
3	Cerebrovascular diseases (I60-I69)	50	5.3	---
4	Influenza and pneumonia (J09-J18)	47	5.0	---
5	Malignant neoplasms (C00-C97)	41	4.3	---
6	Essential hypertension and hypertensive renal disease (I10,I12,I15)	27	2.8	---
6	Chronic lower respiratory diseases (J40-J47)	27	2.8	---
8	Diabetes mellitus (E10-E14)	16	1.7	---
9	Pneumonitis due to solids and liquids (J69)	15	1.6	---
10	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	13	1.4	---
11	Atherosclerosis (I70)	11	1.2	---
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	11	1.2	---
13	Septicemia (A40-A41)	7	0.7	---
13	Nutritional deficiencies (E40-E64)	7	0.7	---
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	5	0.5	---
...	All other causes (Residual)	256	27.0	---

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, male, all ages				
...	All causes	81,887	100.0	309.7
1	Malignant neoplasms (C00-C97)	17,185	21.0	65.0
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	16,526	20.2	62.5
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	8,098	9.9	30.6
4	Diabetes mellitus (E10-E14)	3,726	4.6	14.1
5	Cerebrovascular diseases (I60-I69)	3,528	4.3	13.3
6	Chronic liver disease and cirrhosis (K70,K73-K74)	3,311	4.0	12.5
7	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	2,279	2.8	8.6
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,224	2.7	8.4
9	Chronic lower respiratory diseases (J40-J47)	2,222	2.7	8.4
10	Influenza and pneumonia (J09-J18)	1,606	2.0	6.1
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,388	1.7	5.2
12	Certain conditions originating in the perinatal period (P00-P96)	1,297	1.6	4.9
13	Alzheimer's disease (G30)	1,151	1.4	4.4
14	Septicemia (A40-A41)	1,048	1.3	4.0
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	958	1.2	3.6
...	All other causes (Residual)	15,340	18.7	58.0
Hispanic, male, 1-4 years				
...	All causes	539	100.0	25.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	187	34.7	8.8
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	70	13.0	3.3
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	55	10.2	2.6
4	Malignant neoplasms (C00-C97)	51	9.5	2.4
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	20	3.7	0.9
6	Influenza and pneumonia (J09-J18)	10	1.9	*
7	Septicemia (A40-A41)	8	1.5	*
8	Cerebrovascular diseases (I60-I69)	7	1.3	*
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	4	0.7	*
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	3	0.6	*
10	Meningitis (G00,G03)	3	0.6	*
10	Acute bronchitis and bronchiolitis (J20-J21)	3	0.6	*
10	Certain conditions originating in the perinatal period (P00-P96)	3	0.6	*
10	Complications of medical and surgical care (Y40-Y84,Y88)	3	0.6	*
15	Chronic lower respiratory diseases (J40-J47)	2	0.4	*
...	All other causes (Residual)	110	20.4	5.2

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, male, 5-9 years				
...	All causes	271	100.0	10.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	94	34.7	3.8
2	Malignant neoplasms (C00-C97)	50	18.5	2.0
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	25	9.2	1.0
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	11	4.1	*
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	8	3.0	*
6	Influenza and pneumonia (J09-J18)	6	2.2	*
7	Cerebrovascular diseases (I60-I69)	5	1.8	*
7	Chronic lower respiratory diseases (J40-J47)	5	1.8	*
9	Septicemia (A40-A41)	3	1.1	*
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2	0.7	*
10	Meningitis (G00,G03)	2	0.7	*
10	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2	0.7	*
13	Anemias (D50-D64)	1	0.4	*
13	Diseases of appendix (K35-K38)	1	0.4	*
13	Certain conditions originating in the perinatal period (P00-P96)	1	0.4	*
...	All other causes (Residual)	55	20.3	2.2
Hispanic, male, 10-14 years				
...	All causes	337	100.0	14.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	88	26.1	3.7
2	Malignant neoplasms (C00-C97)	58	17.2	2.5
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	36	10.7	1.5
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	25	7.4	1.1
5	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	23	6.8	1.0
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	9	2.7	*
7	Cerebrovascular diseases (I60-I69)	7	2.1	*
7	Chronic lower respiratory diseases (J40-J47)	7	2.1	*
9	Anemias (D50-D64)	4	1.2	*
10	Influenza and pneumonia (J09-J18)	3	0.9	*
11	Septicemia (A40-A41)	2	0.6	*
11	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2	0.6	*
11	Diabetes mellitus (E10-E14)	2	0.6	*
14	Tuberculosis (A16-A19)	1	0.3	*
14	Meningococcal infection (A39)	1	0.3	*
14	Nutritional deficiencies (E40-E64)	1	0.3	*
14	Meningitis (G00,G03)	1	0.3	*
14	Aortic aneurysm and dissection (I71)	1	0.3	*
14	Diseases of appendix (K35-K38)	1	0.3	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1	0.3	*
...	All other causes (Residual)	64	19.0	2.7

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, male, 15-19 years				
...	All causes	1,385	100.0	58.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	518	37.4	21.9
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	346	25.0	14.6
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	187	13.5	7.9
4	Malignant neoplasms (C00-C97)	105	7.6	4.4
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	29	2.1	1.2
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	20	1.4	0.8
7	Cerebrovascular diseases (I60-I69)	13	0.9	*
8	Legal intervention (Y35,Y89.0)	9	0.6	*
9	Anemias (D50-D64)	6	0.4	*
9	Diabetes mellitus (E10-E14)	6	0.4	*
9	Influenza and pneumonia (J09-J18)	6	0.4	*
12	Chronic lower respiratory diseases (J40-J47)	4	0.3	*
13	Septicemia (A40-A41)	3	0.2	*
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	3	0.2	*
13	Chronic liver disease and cirrhosis (K70,K73-K74)	3	0.2	*
...	All other causes (Residual)	127	9.2	5.4
Hispanic, male, 20-24 years				
...	All causes	2,338	100.0	98.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	999	42.7	42.0
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	489	20.9	20.6
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	323	13.8	13.6
4	Malignant neoplasms (C00-C97)	145	6.2	6.1
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	69	3.0	2.9
6	Legal intervention (Y35,Y89.0)	24	1.0	1.0
7	Cerebrovascular diseases (I60-I69)	19	0.8	*
8	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	17	0.7	*
9	Influenza and pneumonia (J09-J18)	11	0.5	*
10	Human immunodeficiency virus (HIV) disease (B20-B24)	10	0.4	*
11	Chronic lower respiratory diseases (J40-J47)	9	0.4	*
12	Septicemia (A40-A41)	5	0.2	*
12	Chronic liver disease and cirrhosis (K70,K73-K74)	5	0.2	*
14	Meningitis (G00,G03)	4	0.2	*
15	Anemias (D50-D64)	3	0.1	*
15	Diabetes mellitus (E10-E14)	3	0.1	*
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	3	0.1	*
15	Aortic aneurysm and dissection (I71)	3	0.1	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3	0.1	*
...	All other causes (Residual)	194	8.3	8.2

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, male, 25-29 years				
...	All causes	2,147	100.0	92.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	870	40.5	37.5
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	372	17.3	16.0
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	279	13.0	12.0
4	Malignant neoplasms (C00-C97)	122	5.7	5.3
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	103	4.8	4.4
6	Human immunodeficiency virus (HIV) disease (B20-B24)	25	1.2	1.1
6	Cerebrovascular diseases (I60-I69)	25	1.2	1.1
8	Influenza and pneumonia (J09-J18)	20	0.9	0.9
9	Chronic liver disease and cirrhosis (K70,K73-K74)	18	0.8	*
9	Legal intervention (Y35,Y89.0)	18	0.8	*
11	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	16	0.7	*
12	Diabetes mellitus (E10-E14)	15	0.7	*
13	Septicemia (A40-A41)	9	0.4	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	8	0.4	*
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	5	0.2	*
...	All other causes (Residual)	242	11.3	10.4
Hispanic, male, 30-34 years				
...	All causes	2,320	100.0	104.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	821	35.4	37.1
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	292	12.6	13.2
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	267	11.5	12.1
4	Malignant neoplasms (C00-C97)	182	7.8	8.2
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	180	7.8	8.1
6	Human immunodeficiency virus (HIV) disease (B20-B24)	71	3.1	3.2
7	Chronic liver disease and cirrhosis (K70,K73-K74)	58	2.5	2.6
8	Cerebrovascular diseases (I60-I69)	46	2.0	2.1
9	Influenza and pneumonia (J09-J18)	32	1.4	1.4
10	Diabetes mellitus (E10-E14)	29	1.3	1.3
11	Septicemia (A40-A41)	21	0.9	0.9
12	Legal intervention (Y35,Y89.0)	16	0.7	*
13	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	15	0.6	*
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	8	0.3	*
14	Chronic lower respiratory diseases (J40-J47)	8	0.3	*
...	All other causes (Residual)	274	11.8	12.4

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, male, 35-39 years				
...	All causes	2,427	100.0	120.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	705	29.0	35.0
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	250	10.3	12.4
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	246	10.1	12.2
4	Malignant neoplasms (C00-C97)	245	10.1	12.2
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	201	8.3	10.0
6	Chronic liver disease and cirrhosis (K70,K73-K74)	141	5.8	7.0
7	Human immunodeficiency virus (HIV) disease (B20-B24)	84	3.5	4.2
8	Diabetes mellitus (E10-E14)	55	2.3	2.7
9	Cerebrovascular diseases (I60-I69)	53	2.2	2.6
10	Influenza and pneumonia (J09-J18)	24	1.0	1.2
11	Septicemia (A40-A41)	19	0.8	*
11	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	19	0.8	*
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	15	0.6	*
14	Aortic aneurysm and dissection (I71)	14	0.6	*
15	Viral hepatitis (B15-B19)	8	0.3	*
15	Pneumonitis due to solids and liquids (J69)	8	0.3	*
15	Legal intervention (Y35,Y89.0)	8	0.3	*
...	All other causes (Residual)	332	13.7	16.5
Hispanic, male, 40-44 years				
...	All causes	3,051	100.0	166.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	681	22.3	37.2
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	451	14.8	24.7
3	Malignant neoplasms (C00-C97)	386	12.7	21.1
4	Chronic liver disease and cirrhosis (K70,K73-K74)	235	7.7	12.8
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	187	6.1	10.2
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	144	4.7	7.9
7	Human immunodeficiency virus (HIV) disease (B20-B24)	127	4.2	6.9
8	Cerebrovascular diseases (I60-I69)	103	3.4	5.6
9	Diabetes mellitus (E10-E14)	92	3.0	5.0
10	Viral hepatitis (B15-B19)	42	1.4	2.3
11	Influenza and pneumonia (J09-J18)	40	1.3	2.2
12	Septicemia (A40-A41)	32	1.0	1.7
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	30	1.0	1.6
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	19	0.6	*
15	Aortic aneurysm and dissection (I71)	15	0.5	*
15	Chronic lower respiratory diseases (J40-J47)	15	0.5	*
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	15	0.5	*
...	All other causes (Residual)	437	14.3	23.9

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, male, 45-49 years				
...	All causes	4,476	100.0	281.3
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	773	17.3	48.6
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	720	16.1	45.2
3	Malignant neoplasms (C00-C97)	715	16.0	44.9
4	Chronic liver disease and cirrhosis (K70,K73-K74)	449	10.0	28.2
5	Diabetes mellitus (E10-E14)	179	4.0	11.2
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	176	3.9	11.1
7	Cerebrovascular diseases (I60-I69)	172	3.8	10.8
8	Human immunodeficiency virus (HIV) disease (B20-B24)	154	3.4	9.7
9	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	120	2.7	7.5
10	Viral hepatitis (B15-B19)	118	2.6	7.4
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	61	1.4	3.8
12	Influenza and pneumonia (J09-J18)	60	1.3	3.8
13	Septicemia (A40-A41)	51	1.1	3.2
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	32	0.7	2.0
15	Chronic lower respiratory diseases (J40-J47)	29	0.6	1.8
...	All other causes (Residual)	667	14.9	41.9
Hispanic, male, 50-54 years				
...	All causes	5,795	100.0	451.6
1	Malignant neoplasms (C00-C97)	1,274	22.0	99.3
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,124	19.4	87.6
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	627	10.8	48.9
4	Chronic liver disease and cirrhosis (K70,K73-K74)	599	10.3	46.7
5	Diabetes mellitus (E10-E14)	260	4.5	20.3
6	Cerebrovascular diseases (I60-I69)	227	3.9	17.7
7	Viral hepatitis (B15-B19)	169	2.9	13.2
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	159	2.7	12.4
9	Human immunodeficiency virus (HIV) disease (B20-B24)	141	2.4	11.0
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	84	1.4	6.5
11	Influenza and pneumonia (J09-J18)	74	1.3	5.8
12	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	71	1.2	5.5
13	Septicemia (A40-A41)	62	1.1	4.8
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	53	0.9	4.1
15	Chronic lower respiratory diseases (J40-J47)	47	0.8	3.7
...	All other causes (Residual)	824	14.2	64.2

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, male, 55-59 years				
...	All causes	6,406	100.0	669.2
1	Malignant neoplasms (C00-C97)	1,697	26.5	177.3
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,346	21.0	140.6
3	Chronic liver disease and cirrhosis (K70,K73-K74)	582	9.1	60.8
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	428	6.7	44.7
5	Diabetes mellitus (E10-E14)	373	5.8	39.0
6	Cerebrovascular diseases (I60-I69)	262	4.1	27.4
7	Viral hepatitis (B15-B19)	205	3.2	21.4
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	112	1.7	11.7
9	Septicemia (A40-A41)	106	1.7	11.1
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	103	1.6	10.8
11	Chronic lower respiratory diseases (J40-J47)	96	1.5	10.0
12	Human immunodeficiency virus (HIV) disease (B20-B24)	91	1.4	9.5
13	Influenza and pneumonia (J09-J18)	70	1.1	7.3
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	56	0.9	5.8
15	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	49	0.8	5.1
...	All other causes (Residual)	830	13.0	86.7
Hispanic, male, 60-64 years				
...	All causes	6,706	100.0	957.4
1	Malignant neoplasms (C00-C97)	2,021	30.1	288.5
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,510	22.5	215.6
3	Chronic liver disease and cirrhosis (K70,K73-K74)	418	6.2	59.7
4	Diabetes mellitus (E10-E14)	389	5.8	55.5
5	Cerebrovascular diseases (I60-I69)	321	4.8	45.8
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	311	4.6	44.4
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	140	2.1	20.0
8	Viral hepatitis (B15-B19)	137	2.0	19.6
9	Chronic lower respiratory diseases (J40-J47)	122	1.8	17.4
10	Septicemia (A40-A41)	113	1.7	16.1
11	Influenza and pneumonia (J09-J18)	91	1.4	13.0
12	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	77	1.1	11.0
13	Human immunodeficiency virus (HIV) disease (B20-B24)	70	1.0	10.0
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	52	0.8	7.4
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	39	0.6	5.6
...	All other causes (Residual)	895	13.3	127.8

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, male, 65-69 years				
...	All causes	6,612	100.0	1,426.9
1	Malignant neoplasms (C00-C97)	2,111	31.9	455.6
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,572	23.8	339.2
3	Diabetes mellitus (E10-E14)	470	7.1	101.4
4	Chronic liver disease and cirrhosis (K70,K73-K74)	300	4.5	64.7
5	Cerebrovascular diseases (I60-I69)	279	4.2	60.2
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	204	3.1	44.0
7	Chronic lower respiratory diseases (J40-J47)	190	2.9	41.0
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	134	2.0	28.9
9	Influenza and pneumonia (J09-J18)	103	1.6	22.2
10	Septicemia (A40-A41)	98	1.5	21.1
11	Viral hepatitis (B15-B19)	59	0.9	12.7
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	57	0.9	12.3
13	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	43	0.7	9.3
14	Human immunodeficiency virus (HIV) disease (B20-B24)	41	0.6	8.8
15	Pneumonitis due to solids and liquids (J69)	33	0.5	7.1
...	All other causes (Residual)	918	13.9	198.1
Hispanic, male, 70-74 years				
...	All causes	6,928	100.0	2,146.7
1	Malignant neoplasms (C00-C97)	2,041	29.5	632.4
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,674	24.2	518.7
3	Diabetes mellitus (E10-E14)	492	7.1	152.5
4	Cerebrovascular diseases (I60-I69)	353	5.1	109.4
5	Chronic lower respiratory diseases (J40-J47)	263	3.8	81.5
6	Chronic liver disease and cirrhosis (K70,K73-K74)	200	2.9	62.0
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	160	2.3	49.6
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	145	2.1	44.9
9	Influenza and pneumonia (J09-J18)	143	2.1	44.3
10	Septicemia (A40-A41)	122	1.8	37.8
11	Parkinson's disease (G20-G21)	90	1.3	27.9
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	90	1.3	27.9
13	Alzheimer's disease (G30)	66	1.0	20.5
14	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	43	0.6	13.3
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	36	0.5	11.2
...	All other causes (Residual)	1,010	14.6	313.0

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, male, 75-79 years				
...	All causes	7,760	100.0	3,459.5
1	Malignant neoplasms (C00-C97)	2,124	27.4	946.9
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,931	24.9	860.9
3	Diabetes mellitus (E10-E14)	478	6.2	213.1
4	Cerebrovascular diseases (I60-I69)	464	6.0	206.9
5	Chronic lower respiratory diseases (J40-J47)	377	4.9	168.1
6	Influenza and pneumonia (J09-J18)	174	2.2	77.6
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	174	2.2	77.6
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	174	2.2	77.6
9	Alzheimer's disease (G30)	148	1.9	66.0
10	Chronic liver disease and cirrhosis (K70,K73-K74)	146	1.9	65.1
11	Parkinson's disease (G20-G21)	130	1.7	58.0
12	Septicemia (A40-A41)	104	1.3	46.4
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	97	1.3	43.2
14	Pneumonitis due to solids and liquids (J69)	65	0.8	29.0
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	59	0.8	26.3
...	All other causes (Residual)	1,115	14.4	497.1
Hispanic, male, 80-84 years				
...	All causes	8,321	100.0	5,571.1
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,152	25.9	1,440.8
2	Malignant neoplasms (C00-C97)	2,003	24.1	1,341.1
3	Cerebrovascular diseases (I60-I69)	453	5.4	303.3
4	Diabetes mellitus (E10-E14)	447	5.4	299.3
5	Chronic lower respiratory diseases (J40-J47)	411	4.9	275.2
6	Alzheimer's disease (G30)	283	3.4	189.5
7	Influenza and pneumonia (J09-J18)	255	3.1	170.7
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	185	2.2	123.9
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	184	2.2	123.2
10	Parkinson's disease (G20-G21)	149	1.8	99.8
11	Septicemia (A40-A41)	121	1.5	81.0
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	109	1.3	73.0
13	Chronic liver disease and cirrhosis (K70,K73-K74)	102	1.2	68.3
14	Pneumonitis due to solids and liquids (J69)	72	0.9	48.2
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	65	0.8	43.5
...	All other causes (Residual)	1,330	16.0	890.5

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, male, 85-89 years				
...	All causes	6,828	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,923	28.2	---
2	Malignant neoplasms (C00-C97)	1,261	18.5	---
3	Cerebrovascular diseases (I60-I69)	455	6.7	---
4	Chronic lower respiratory diseases (J40-J47)	382	5.6	---
5	Alzheimer's disease (G30)	317	4.6	---
6	Diabetes mellitus (E10-E14)	285	4.2	---
7	Influenza and pneumonia (J09-J18)	241	3.5	---
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	185	2.7	---
9	Parkinson's disease (G20-G21)	130	1.9	---
10	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	129	1.9	---
11	Septicemia (A40-A41)	100	1.5	---
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	97	1.4	---
13	Pneumonitis due to solids and liquids (J69)	80	1.2	---
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown	56	0.8	---
15	Chronic liver disease and cirrhosis (K70,K73-K74)	31	0.5	---
...	All other causes (Residual)	1,156	16.9	---
Hispanic, male, 90-94 years				
...	All causes	3,398	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	973	28.6	---
2	Malignant neoplasms (C00-C97)	467	13.7	---
3	Alzheimer's disease (G30)	212	6.2	---
4	Cerebrovascular diseases (I60-I69)	197	5.8	---
5	Chronic lower respiratory diseases (J40-J47)	172	5.1	---
6	Influenza and pneumonia (J09-J18)	148	4.4	---
7	Diabetes mellitus (E10-E14)	117	3.4	---
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	79	2.3	---
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	76	2.2	---
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	70	2.1	---
11	Parkinson's disease (G20-G21)	61	1.8	---
12	Septicemia (A40-A41)	40	1.2	---
13	Pneumonitis due to solids and liquids (J69)	31	0.9	---
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	17	0.5	---
14	Chronic liver disease and cirrhosis (K70,K73-K74)	17	0.5	---
...	All other causes (Residual)	721	21.2	---

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, male, 95-99 years				
...	All causes	1,002	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	317	31.6	---
2	Malignant neoplasms (C00-C97)	104	10.4	---
3	Influenza and pneumonia (J09-J18)	62	6.2	---
4	Chronic lower respiratory diseases (J40-J47)	60	6.0	---
5	Alzheimer's disease (G30)	54	5.4	---
6	Cerebrovascular diseases (I60-I69)	47	4.7	---
7	Diabetes mellitus (E10-E14)	27	2.7	---
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	22	2.2	---
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	21	2.1	---
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	18	1.8	---
11	Pneumonitis due to solids and liquids (J69)	14	1.4	---
12	Septicemia (A40-A41)	12	1.2	---
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	8	0.8	---
14	Parkinson's disease (G20-G21)	6	0.6	---
14	Hernia (K40-K46)	6	0.6	---
...	All other causes (Residual)	224	22.4	---
Hispanic, male, 100 years and over				
...	All causes	214	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	74	34.6	---
2	Malignant neoplasms (C00-C97)	15	7.0	---
3	Cerebrovascular diseases (I60-I69)	12	5.6	---
3	Influenza and pneumonia (J09-J18)	12	5.6	---
5	Chronic lower respiratory diseases (J40-J47)	11	5.1	---
6	Alzheimer's disease (G30)	10	4.7	---
7	Pneumonitis due to solids and liquids (J69)	7	3.3	---
8	Diabetes mellitus (E10-E14)	6	2.8	---
8	Essential hypertension and hypertensive renal disease (I10,I12,I15)	6	2.8	---
10	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	5	2.3	---
11	Nutritional deficiencies (E40-E64)	3	1.4	---
12	Septicemia (A40-A41)	2	0.9	---
12	Parkinson's disease (G20-G21)	2	0.9	---
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2	0.9	---
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1	0.5	---
15	Atherosclerosis (I70)	1	0.5	---
15	Peptic ulcer (K25-K28)	1	0.5	---
15	Chronic liver disease and cirrhosis (K70,K73-K74)	1	0.5	---
...	All other causes (Residual)	43	20.1	---

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, female, all ages				
...	All causes	67,748	100.0	264.6
1	Malignant neoplasms (C00-C97)	15,196	22.4	59.4
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	13,859	20.5	54.1
3	Cerebrovascular diseases (I60-I69)	4,024	5.9	15.7
4	Diabetes mellitus (E10-E14)	3,328	4.9	13.0
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,068	4.5	12.0
6	Alzheimer's disease (G30)	2,448	3.6	9.6
7	Chronic lower respiratory diseases (J40-J47)	2,133	3.1	8.3
8	Influenza and pneumonia (J09-J18)	1,650	2.4	6.4
9	Chronic liver disease and cirrhosis (K70,K73-K74)	1,463	2.2	5.7
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,411	2.1	5.5
11	Septicemia (A40-A41)	1,120	1.7	4.4
12	Certain conditions originating in the perinatal period (P00-P96)	1,090	1.6	4.3
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	927	1.4	3.6
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	845	1.2	3.3
15	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	496	0.7	1.9
...	All other causes (Residual)	14,690	21.7	57.4
Hispanic, female, 1-4 years				
...	All causes	433	100.0	21.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	130	30.0	6.4
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	56	12.9	2.8
3	Malignant neoplasms (C00-C97)	46	10.6	2.3
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	39	9.0	1.9
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	15	3.5	*
6	Influenza and pneumonia (J09-J18)	11	2.5	*
7	Septicemia (A40-A41)	8	1.8	*
8	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	7	1.6	*
9	Cerebrovascular diseases (I60-I69)	3	0.7	*
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3	0.7	*
9	Complications of medical and surgical care (Y40-Y84,Y88)	3	0.7	*
12	Chronic lower respiratory diseases (J40-J47)	2	0.5	*
12	Diseases of appendix (K35-K38)	2	0.5	*
12	Hernia (K40-K46)	2	0.5	*
12	Certain conditions originating in the perinatal period (P00-P96)	2	0.5	*
...	All other causes (Residual)	104	24.0	5.1

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, female, 5-9 years				
...	All causes	219	100.0	9.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	51	23.3	2.1
2	Malignant neoplasms (C00-C97)	50	22.8	2.1
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	23	10.5	1.0
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	17	7.8	*
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	11	5.0	*
6	Influenza and pneumonia (J09-J18)	9	4.1	*
7	Cerebrovascular diseases (I60-I69)	6	2.7	*
8	Chronic lower respiratory diseases (J40-J47)	4	1.8	*
9	Septicemia (A40-A41)	3	1.4	*
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2	0.9	*
10	Certain conditions originating in the perinatal period (P00-P96)	2	0.9	*
10	Complications of medical and surgical care (Y40-Y84,Y88)	2	0.9	*
13	Meningococcal infection (A39)	1	0.5	*
13	Aortic aneurysm and dissection (I71)	1	0.5	*
13	Acute bronchitis and bronchiolitis (J20-J21)	1	0.5	*
13	Diseases of appendix (K35-K38)	1	0.5	*
...	All other causes (Residual)	35	16.0	1.5
Hispanic, female, 10-14 years				
...	All causes	225	100.0	9.9
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	45	20.0	2.0
2	Malignant neoplasms (C00-C97)	42	18.7	1.9
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	21	9.3	0.9
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	20	8.9	0.9
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	11	4.9	*
6	Chronic lower respiratory diseases (J40-J47)	7	3.1	*
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	7	3.1	*
8	Diabetes mellitus (E10-E14)	4	1.8	*
8	Cerebrovascular diseases (I60-I69)	4	1.8	*
8	Influenza and pneumonia (J09-J18)	4	1.8	*
11	Septicemia (A40-A41)	2	0.9	*
11	Pneumonitis due to solids and liquids (J69)	2	0.9	*
11	Certain conditions originating in the perinatal period (P00-P96)	2	0.9	*
14	Meningococcal infection (A39)	1	0.4	*
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1	0.4	*
14	Anemias (D50-D64)	1	0.4	*
14	Meningitis (G00,G03)	1	0.4	*
14	Legal intervention (Y35,Y89.0)	1	0.4	*
...	All other causes (Residual)	49	21.8	2.2

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, female, 15-19 years				
...	All causes	451	100.0	20.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	146	32.4	6.6
2	Malignant neoplasms (C00-C97)	67	14.9	3.0
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	62	13.7	2.8
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	35	7.8	1.6
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	21	4.7	0.9
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	10	2.2	*
7	Pregnancy, childbirth and the puerperium (O00-O99)	9	2.0	*
8	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	6	1.3	*
8	Cerebrovascular diseases (I60-I69)	6	1.3	*
10	Septicemia (A40-A41)	4	0.9	*
11	Chronic lower respiratory diseases (J40-J47)	3	0.7	*
12	Anemias (D50-D64)	2	0.4	*
12	Diabetes mellitus (E10-E14)	2	0.4	*
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2	0.4	*
15	Viral hepatitis (B15-B19)	1	0.2	*
15	Human immunodeficiency virus (HIV) disease (B20-B24)	1	0.2	*
15	Chronic liver disease and cirrhosis (K70,K73-K74)	1	0.2	*
15	Certain conditions originating in the perinatal period (P00-P96)	1	0.2	*
...	All other causes (Residual)	72	16.0	3.3
Hispanic, female, 20-24 years				
...	All causes	652	100.0	31.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	248	38.0	11.8
2	Malignant neoplasms (C00-C97)	66	10.1	3.1
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	63	9.7	3.0
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	63	9.7	3.0
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	37	5.7	1.8
6	Pregnancy, childbirth and the puerperium (O00-O99)	22	3.4	1.0
7	Cerebrovascular diseases (I60-I69)	15	2.3	*
8	Influenza and pneumonia (J09-J18)	11	1.7	*
9	Human immunodeficiency virus (HIV) disease (B20-B24)	9	1.4	*
9	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	9	1.4	*
11	Diabetes mellitus (E10-E14)	8	1.2	*
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	7	1.1	*
13	Septicemia (A40-A41)	4	0.6	*
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	4	0.6	*
15	Anemias (D50-D64)	2	0.3	*
...	All other causes (Residual)	84	12.9	4.0

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, female, 25-29 years				
...	All causes	715	100.0	34.9
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	197	27.6	9.6
2	Malignant neoplasms (C00-C97)	103	14.4	5.0
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	65	9.1	3.2
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	63	8.8	3.1
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	39	5.5	1.9
6	Pregnancy, childbirth and the puerperium (O00-O99)	30	4.2	1.5
7	Cerebrovascular diseases (I60-I69)	15	2.1	*
8	Diabetes mellitus (E10-E14)	14	2.0	*
8	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	14	2.0	*
10	Influenza and pneumonia (J09-J18)	13	1.8	*
11	Septicemia (A40-A41)	9	1.3	*
12	Chronic liver disease and cirrhosis (K70,K73-K74)	8	1.1	*
13	Human immunodeficiency virus (HIV) disease (B20-B24)	7	1.0	*
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	6	0.8	*
15	Chronic lower respiratory diseases (J40-J47)	5	0.7	*
...	All other causes (Residual)	127	17.8	6.2
Hispanic, female, 30-34 years				
...	All causes	879	100.0	43.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	208	23.7	10.3
2	Malignant neoplasms (C00-C97)	188	21.4	9.3
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	64	7.3	3.2
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	53	6.0	2.6
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	45	5.1	2.2
6	Pregnancy, childbirth and the puerperium (O00-O99)	25	2.8	1.2
7	Cerebrovascular diseases (I60-I69)	24	2.7	1.2
8	Chronic liver disease and cirrhosis (K70,K73-K74)	22	2.5	1.1
9	Diabetes mellitus (E10-E14)	20	2.3	1.0
10	Human immunodeficiency virus (HIV) disease (B20-B24)	16	1.8	*
11	Influenza and pneumonia (J09-J18)	14	1.6	*
12	Septicemia (A40-A41)	12	1.4	*
13	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	11	1.3	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	8	0.9	*
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	6	0.7	*
...	All other causes (Residual)	163	18.5	8.1

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, female, 35-39 years				
...	All causes	1,129	100.0	58.9
1	Malignant neoplasms (C00-C97)	340	30.1	17.7
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	200	17.7	10.4
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	95	8.4	5.0
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	53	4.7	2.8
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	37	3.3	1.9
6	Cerebrovascular diseases (I60-I69)	35	3.1	1.8
7	Diabetes mellitus (E10-E14)	31	2.7	1.6
8	Chronic liver disease and cirrhosis (K70,K73-K74)	30	2.7	1.6
9	Human immunodeficiency virus (HIV) disease (B20-B24)	28	2.5	1.5
10	Pregnancy, childbirth and the puerperium (O00-O99)	23	2.0	1.2
11	Septicemia (A40-A41)	18	1.6	*
12	Influenza and pneumonia (J09-J18)	16	1.4	*
13	Chronic lower respiratory diseases (J40-J47)	11	1.0	*
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	8	0.7	*
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	7	0.6	*
...	All other causes (Residual)	197	17.4	10.3
Hispanic, female, 40-44 years				
...	All causes	1,585	100.0	90.8
1	Malignant neoplasms (C00-C97)	538	33.9	30.8
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	221	13.9	12.7
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	149	9.4	8.5
4	Cerebrovascular diseases (I60-I69)	64	4.0	3.7
5	Chronic liver disease and cirrhosis (K70,K73-K74)	63	4.0	3.6
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	53	3.3	3.0
7	Diabetes mellitus (E10-E14)	49	3.1	2.8
8	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	46	2.9	2.6
9	Human immunodeficiency virus (HIV) disease (B20-B24)	32	2.0	1.8
10	Influenza and pneumonia (J09-J18)	28	1.8	1.6
11	Septicemia (A40-A41)	25	1.6	1.4
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	19	1.2	*
13	Chronic lower respiratory diseases (J40-J47)	18	1.1	*
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	12	0.8	*
15	Viral hepatitis (B15-B19)	9	0.6	*
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	9	0.6	*
15	Pregnancy, childbirth and the puerperium (O00-O99)	9	0.6	*
...	All other causes (Residual)	241	15.2	13.8

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, female, 45-49 years				
...	All causes	2,283	100.0	147.7
1	Malignant neoplasms (C00-C97)	759	33.2	49.1
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	276	12.1	17.9
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	208	9.1	13.5
4	Chronic liver disease and cirrhosis (K70,K73-K74)	154	6.7	10.0
5	Cerebrovascular diseases (I60-I69)	124	5.4	8.0
6	Diabetes mellitus (E10-E14)	97	4.2	6.3
7	Human immunodeficiency virus (HIV) disease (B20-B24)	53	2.3	3.4
8	Septicemia (A40-A41)	50	2.2	3.2
9	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	44	1.9	2.8
10	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	31	1.4	2.0
11	Viral hepatitis (B15-B19)	30	1.3	1.9
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	29	1.3	1.9
13	Influenza and pneumonia (J09-J18)	27	1.2	1.7
14	Chronic lower respiratory diseases (J40-J47)	26	1.1	1.7
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	20	0.9	1.3
...	All other causes (Residual)	355	15.5	23.0
Hispanic, female, 50-54 years				
...	All causes	2,995	100.0	231.2
1	Malignant neoplasms (C00-C97)	1,123	37.5	86.7
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	358	12.0	27.6
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	206	6.9	15.9
4	Chronic liver disease and cirrhosis (K70,K73-K74)	164	5.5	12.7
5	Cerebrovascular diseases (I60-I69)	146	4.9	11.3
6	Diabetes mellitus (E10-E14)	135	4.5	10.4
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	57	1.9	4.4
8	Septicemia (A40-A41)	54	1.8	4.2
9	Viral hepatitis (B15-B19)	51	1.7	3.9
10	Chronic lower respiratory diseases (J40-J47)	50	1.7	3.9
11	Human immunodeficiency virus (HIV) disease (B20-B24)	48	1.6	3.7
12	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	44	1.5	3.4
13	Influenza and pneumonia (J09-J18)	43	1.4	3.3
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	26	0.9	2.0
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	23	0.8	1.8
...	All other causes (Residual)	467	15.6	36.0

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, female, 55-59 years				
...	All causes	3,605	100.0	354.2
1	Malignant neoplasms (C00-C97)	1,358	37.7	133.4
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	567	15.7	55.7
3	Diabetes mellitus (E10-E14)	211	5.9	20.7
4	Chronic liver disease and cirrhosis (K70,K73-K74)	182	5.0	17.9
5	Cerebrovascular diseases (I60-I69)	170	4.7	16.7
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	141	3.9	13.9
7	Septicemia (A40-A41)	74	2.1	7.3
8	Viral hepatitis (B15-B19)	71	2.0	7.0
9	Chronic lower respiratory diseases (J40-J47)	70	1.9	6.9
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	68	1.9	6.7
11	Influenza and pneumonia (J09-J18)	46	1.3	4.5
12	Human immunodeficiency virus (HIV) disease (B20-B24)	34	0.9	3.3
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	26	0.7	2.6
14	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	24	0.7	2.4
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	18	0.5	*
...	All other causes (Residual)	545	15.1	53.5
Hispanic, female, 60-64 years				
...	All causes	4,241	100.0	542.2
1	Malignant neoplasms (C00-C97)	1,550	36.5	198.2
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	716	16.9	91.5
3	Diabetes mellitus (E10-E14)	294	6.9	37.6
4	Cerebrovascular diseases (I60-I69)	208	4.9	26.6
5	Chronic liver disease and cirrhosis (K70,K73-K74)	165	3.9	21.1
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	137	3.2	17.5
7	Chronic lower respiratory diseases (J40-J47)	102	2.4	13.0
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	101	2.4	12.9
9	Septicemia (A40-A41)	70	1.7	8.9
10	Influenza and pneumonia (J09-J18)	65	1.5	8.3
11	Viral hepatitis (B15-B19)	60	1.4	7.7
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	42	1.0	5.4
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	27	0.6	3.5
13	Alzheimer's disease (G30)	27	0.6	3.5
15	Human immunodeficiency virus (HIV) disease (B20-B24)	20	0.5	2.6
...	All other causes (Residual)	657	15.5	84.0

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, female, 65-69 years				
...	All causes	4,741	100.0	853.3
1	Malignant neoplasms (C00-C97)	1,650	34.8	297.0
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	856	18.1	154.1
3	Diabetes mellitus (E10-E14)	333	7.0	59.9
4	Cerebrovascular diseases (I60-I69)	237	5.0	42.7
5	Chronic liver disease and cirrhosis (K70,K73-K74)	162	3.4	29.2
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	139	2.9	25.0
7	Chronic lower respiratory diseases (J40-J47)	130	2.7	23.4
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	115	2.4	20.7
9	Septicemia (A40-A41)	93	2.0	16.7
10	Influenza and pneumonia (J09-J18)	92	1.9	16.6
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	54	1.1	9.7
12	Alzheimer's disease (G30)	53	1.1	9.5
13	Viral hepatitis (B15-B19)	41	0.9	7.4
14	Enterocolitis due to <i>Clostridium difficile</i> (A04.7)	28	0.6	5.0
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	26	0.5	4.7
...	All other causes (Residual)	732	15.4	131.7
Hispanic, female, 70-74 years				
...	All causes	5,703	100.0	1,378.4
1	Malignant neoplasms (C00-C97)	1,725	30.2	416.9
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,104	19.4	266.8
3	Diabetes mellitus (E10-E14)	406	7.1	98.1
4	Cerebrovascular diseases (I60-I69)	337	5.9	81.5
5	Chronic lower respiratory diseases (J40-J47)	216	3.8	52.2
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	189	3.3	45.7
7	Chronic liver disease and cirrhosis (K70,K73-K74)	167	2.9	40.4
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	112	2.0	27.1
9	Influenza and pneumonia (J09-J18)	111	1.9	26.8
10	Septicemia (A40-A41)	104	1.8	25.1
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	71	1.2	17.2
12	Alzheimer's disease (G30)	69	1.2	16.7
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	44	0.8	10.6
14	Viral hepatitis (B15-B19)	36	0.6	8.7
15	Pneumonitis due to solids and liquids (J69)	29	0.5	7.0
...	All other causes (Residual)	983	17.2	237.6

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, female, 75-79 years				
...	All causes	7,161	100.0	2,304.5
1	Malignant neoplasms (C00-C97)	1,811	25.3	582.8
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,569	21.9	504.9
3	Cerebrovascular diseases (I60-I69)	515	7.2	165.7
4	Diabetes mellitus (E10-E14)	467	6.5	150.3
5	Chronic lower respiratory diseases (J40-J47)	277	3.9	89.1
6	Alzheimer's disease (G30)	249	3.5	80.1
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	189	2.6	60.8
8	Influenza and pneumonia (J09-J18)	188	2.6	60.5
9	Septicemia (A40-A41)	138	1.9	44.4
10	Chronic liver disease and cirrhosis (K70,K73-K74)	135	1.9	43.4
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	121	1.7	38.9
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	94	1.3	30.3
13	Parkinson's disease (G20-G21)	67	0.9	21.6
14	Enterocolitis due to <i>Clostridium difficile</i> (A04.7)	41	0.6	13.2
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	38	0.5	12.2
...	All other causes (Residual)	1,262	17.6	406.1
Hispanic, female, 80-84 years				
...	All causes	9,379	100.0	4,161.2
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,293	24.4	1,017.3
2	Malignant neoplasms (C00-C97)	1,792	19.1	795.1
3	Cerebrovascular diseases (I60-I69)	739	7.9	327.9
4	Diabetes mellitus (E10-E14)	525	5.6	232.9
5	Alzheimer's disease (G30)	480	5.1	213.0
6	Chronic lower respiratory diseases (J40-J47)	395	4.2	175.2
7	Influenza and pneumonia (J09-J18)	266	2.8	118.0
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	222	2.4	98.5
9	Septicemia (A40-A41)	172	1.8	76.3
10	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	171	1.8	75.9
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	165	1.8	73.2
12	Parkinson's disease (G20-G21)	130	1.4	57.7
13	Chronic liver disease and cirrhosis (K70,K73-K74)	127	1.4	56.3
14	Pneumonitis due to solids and liquids (J69)	55	0.6	24.4
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	54	0.6	24.0
...	All other causes (Residual)	1,793	19.1	795.5

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, female, 85-89 years				
...	All causes	9,344	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,585	27.7	---
2	Malignant neoplasms (C00-C97)	1,259	13.5	---
3	Cerebrovascular diseases (I60-I69)	696	7.4	---
4	Alzheimer's disease (G30)	681	7.3	---
5	Chronic lower respiratory diseases (J40-J47)	439	4.7	---
6	Diabetes mellitus (E10-E14)	425	4.5	---
7	Influenza and pneumonia (J09-J18)	308	3.3	---
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	200	2.1	---
9	Essential hypertension and hypertensive renal disease (I10,I12,I15)	176	1.9	---
10	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	161	1.7	---
11	Septicemia (A40-A41)	151	1.6	---
12	Parkinson's disease (G20-G21)	119	1.3	---
13	Pneumonitis due to solids and liquids (J69)	89	1.0	---
14	Chronic liver disease and cirrhosis (K70,K73-K74)	59	0.6	---
15	Enterocolitis due to <i>Clostridium difficile</i> (A04.7)	53	0.6	---
...	All other causes (Residual)	1,943	20.8	---
Hispanic, female, 90-94 years				
...	All causes	6,318	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,924	30.5	---
2	Malignant neoplasms (C00-C97)	538	8.5	---
3	Alzheimer's disease (G30)	533	8.4	---
4	Cerebrovascular diseases (I60-I69)	449	7.1	---
5	Chronic lower respiratory diseases (J40-J47)	259	4.1	---
6	Diabetes mellitus (E10-E14)	245	3.9	---
7	Influenza and pneumonia (J09-J18)	230	3.6	---
8	Essential hypertension and hypertensive renal disease (I10,I12,I15)	136	2.2	---
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	117	1.9	---
10	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	115	1.8	---
11	Septicemia (A40-A41)	72	1.1	---
12	Parkinson's disease (G20-G21)	60	0.9	---
13	Pneumonitis due to solids and liquids (J69)	54	0.9	---
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	36	0.6	---
15	Enterocolitis due to <i>Clostridium difficile</i> (A04.7)	32	0.5	---
...	All other causes (Residual)	1,518	24.0	---

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, female, 95-99 years				
...	All causes	2,749	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	877	31.9	---
2	Alzheimer's disease (G30)	265	9.6	---
3	Cerebrovascular diseases (I60-I69)	180	6.5	---
4	Malignant neoplasms (C00-C97)	158	5.7	---
5	Influenza and pneumonia (J09-J18)	124	4.5	---
6	Chronic lower respiratory diseases (J40-J47)	93	3.4	---
7	Essential hypertension and hypertensive renal disease (I10,I12,I15)	73	2.7	---
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	57	2.1	---
9	Diabetes mellitus (E10-E14)	51	1.9	---
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	41	1.5	---
11	Septicemia (A40-A41)	34	1.2	---
12	Pneumonitis due to solids and liquids (J69)	21	0.8	---
13	Atherosclerosis (I70)	19	0.7	---
14	Enterocolitis due to <i>Clostridium difficile</i> (A04.7)	12	0.4	---
15	Anemias (D50-D64)	10	0.4	---
15	Cholelithiasis and other disorders of gallbladder (K80-K82)	10	0.4	---
...	All other causes (Residual)	724	26.3	---
Hispanic, female, 100 years and over				
...	All causes	735	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	259	35.2	---
2	Alzheimer's disease (G30)	73	9.9	---
3	Cerebrovascular diseases (I60-I69)	38	5.2	---
4	Influenza and pneumonia (J09-J18)	35	4.8	---
5	Malignant neoplasms (C00-C97)	26	3.5	---
6	Essential hypertension and hypertensive renal disease (I10,I12,I15)	21	2.9	---
7	Chronic lower respiratory diseases (J40-J47)	16	2.2	---
8	Diabetes mellitus (E10-E14)	10	1.4	---
8	Atherosclerosis (I70)	10	1.4	---
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	9	1.2	---
11	Pneumonitis due to solids and liquids (J69)	8	1.1	---
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	8	1.1	---
13	Septicemia (A40-A41)	5	0.7	---
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	4	0.5	---
14	Nutritional deficiencies (E40-E64)	4	0.5	---
14	Cholelithiasis and other disorders of gallbladder (K80-K82)	4	0.5	---
...	All other causes (Residual)	205	27.9	---

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, both sexes, all ages				
...	All causes	2,006,319	100.0	1,001.0
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	482,979	24.1	241.0
2	Malignant neoplasms (C00-C97)	460,026	22.9	229.5
3	Chronic lower respiratory diseases (J40-J47)	127,035	6.3	63.4
4	Cerebrovascular diseases (I60-I69)	101,043	5.0	50.4
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	98,605	4.9	49.2
6	Alzheimer's disease (G30)	74,448	3.7	37.1
7	Diabetes mellitus (E10-E14)	50,964	2.5	25.4
8	Influenza and pneumonia (J09-J18)	43,350	2.2	21.6
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	33,492	1.7	16.7
10	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	33,031	1.6	16.5
11	Septicemia (A40-A41)	26,669	1.3	13.3
12	Chronic liver disease and cirrhosis (K70,K73-K74)	24,752	1.2	12.3
13	Parkinson's disease (G20-G21)	20,625	1.0	10.3
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	20,153	1.0	10.1
15	Pneumonitis due to solids and liquids (J69)	15,513	0.8	7.7
...	All other causes (Residual)	393,634	19.6	196.4
Non-Hispanic, white, both sexes, 1-4 years				
...	All causes	2,055	100.0	24.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	724	35.2	8.5
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	236	11.5	2.8
3	Malignant neoplasms (C00-C97)	183	8.9	2.1
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	150	7.3	1.8
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	74	3.6	0.9
6	Influenza and pneumonia (J09-J18)	45	2.2	0.5
7	Septicemia (A40-A41)	30	1.5	0.4
8	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	24	1.2	0.3
9	Chronic lower respiratory diseases (J40-J47)	19	0.9	*
10	Certain conditions originating in the perinatal period (P00-P96)	17	0.8	*
11	Cerebrovascular diseases (I60-I69)	14	0.7	*
12	Acute bronchitis and bronchiolitis (J20-J21)	8	0.4	*
13	Complications of medical and surgical care (Y40-Y84,Y88)	7	0.3	*
14	Anemias (D50-D64)	6	0.3	*
15	Meningitis (G00,G03)	5	0.2	*
...	All other causes (Residual)	513	25.0	6.0

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, both sexes, 5-9 years				
...	All causes	1,300	100.0	11.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	428	32.9	3.8
2	Malignant neoplasms (C00-C97)	254	19.5	2.3
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	87	6.7	0.8
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	61	4.7	0.5
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	41	3.2	0.4
6	Influenza and pneumonia (J09-J18)	32	2.5	0.3
7	Septicemia (A40-A41)	25	1.9	0.2
8	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	24	1.8	0.2
9	Chronic lower respiratory diseases (J40-J47)	18	1.4	*
10	Cerebrovascular diseases (I60-I69)	13	1.0	*
11	Certain conditions originating in the perinatal period (P00-P96)	10	0.8	*
12	Anemias (D50-D64)	7	0.5	*
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	4	0.3	*
14	Meningococcal infection (A39)	3	0.2	*
14	Meningitis (G00,G03)	3	0.2	*
14	Pneumonitis due to solids and liquids (J69)	3	0.2	*
...	All other causes (Residual)	287	22.1	2.6
Non-Hispanic, white, both sexes, 10-14 years				
...	All causes	1,608	100.0	13.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	545	33.9	4.7
2	Malignant neoplasms (C00-C97)	227	14.1	1.9
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	168	10.4	1.4
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	81	5.0	0.7
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	64	4.0	0.5
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	56	3.5	0.5
7	Influenza and pneumonia (J09-J18)	29	1.8	0.2
8	Cerebrovascular diseases (I60-I69)	25	1.6	0.2
9	Septicemia (A40-A41)	21	1.3	0.2
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	19	1.2	*
11	Diabetes mellitus (E10-E14)	14	0.9	*
12	Chronic lower respiratory diseases (J40-J47)	11	0.7	*
13	Pneumonitis due to solids and liquids (J69)	7	0.4	*
14	Certain conditions originating in the perinatal period (P00-P96)	6	0.4	*
15	Complications of medical and surgical care (Y40-Y84,Y88)	5	0.3	*
...	All other causes (Residual)	330	20.5	2.8

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, both sexes, 15-19 years				
...	All causes	5,955	100.0	48.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,911	48.9	23.6
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,276	21.4	10.3
3	Malignant neoplasms (C00-C97)	336	5.6	2.7
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	245	4.1	2.0
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	160	2.7	1.3
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	117	2.0	0.9
7	Influenza and pneumonia (J09-J18)	48	0.8	0.4
8	Cerebrovascular diseases (I60-I69)	36	0.6	0.3
9	Diabetes mellitus (E10-E14)	30	0.5	0.2
10	Septicemia (A40-A41)	26	0.4	0.2
11	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	23	0.4	0.2
12	Chronic lower respiratory diseases (J40-J47)	22	0.4	0.2
13	Pregnancy, childbirth and the puerperium (O00-O99)	14	0.2	*
14	Complications of medical and surgical care (Y40-Y84,Y88)	12	0.2	*
15	Pneumonitis due to solids and liquids (J69)	8	0.1	*
...	All other causes (Residual)	691	11.6	5.6
Non-Hispanic, white, both sexes, 20-24 years				
...	All causes	11,006	100.0	85.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	5,617	51.0	43.6
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,137	19.4	16.6
3	Malignant neoplasms (C00-C97)	555	5.0	4.3
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	456	4.1	3.5
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	333	3.0	2.6
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	135	1.2	1.0
7	Influenza and pneumonia (J09-J18)	90	0.8	0.7
8	Diabetes mellitus (E10-E14)	67	0.6	0.5
9	Pregnancy, childbirth and the puerperium (O00-O99)	59	0.5	0.5
10	Cerebrovascular diseases (I60-I69)	54	0.5	0.4
11	Septicemia (A40-A41)	51	0.5	0.4
12	Chronic lower respiratory diseases (J40-J47)	42	0.4	0.3
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	31	0.3	0.2
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	24	0.2	0.2
15	Pneumonitis due to solids and liquids (J69)	20	0.2	0.2
...	All other causes (Residual)	1,335	12.1	10.4

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, both sexes, 25-29 years				
...	All causes	12,551	100.0	100.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	5,725	45.6	45.8
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,280	18.2	18.2
3	Malignant neoplasms (C00-C97)	788	6.3	6.3
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	658	5.2	5.3
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	504	4.0	4.0
6	Influenza and pneumonia (J09-J18)	148	1.2	1.2
7	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	124	1.0	1.0
8	Diabetes mellitus (E10-E14)	123	1.0	1.0
9	Pregnancy, childbirth and the puerperium (O00-O99)	100	0.8	0.8
10	Chronic liver disease and cirrhosis (K70,K73-K74)	85	0.7	0.7
11	Cerebrovascular diseases (I60-I69)	84	0.7	0.7
12	Septicemia (A40-A41)	77	0.6	0.6
13	Chronic lower respiratory diseases (J40-J47)	49	0.4	0.4
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	41	0.3	0.3
15	Pneumonitis due to solids and liquids (J69)	30	0.2	0.2
15	Legal intervention (Y35,Y89.0)	30	0.2	0.2
...	All other causes (Residual)	1,705	13.6	13.6
Non-Hispanic, white, both sexes, 30-34 years				
...	All causes	14,203	100.0	118.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	5,457	38.4	45.4
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,301	16.2	19.2
3	Malignant neoplasms (C00-C97)	1,269	8.9	10.6
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,158	8.2	9.6
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	443	3.1	3.7
6	Diabetes mellitus (E10-E14)	235	1.7	2.0
7	Chronic liver disease and cirrhosis (K70,K73-K74)	204	1.4	1.7
8	Cerebrovascular diseases (I60-I69)	169	1.2	1.4
9	Influenza and pneumonia (J09-J18)	162	1.1	1.3
10	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	157	1.1	1.3
11	Septicemia (A40-A41)	109	0.8	0.9
12	Chronic lower respiratory diseases (J40-J47)	93	0.7	0.8
13	Human immunodeficiency virus (HIV) disease (B20-B24)	87	0.6	0.7
14	Pregnancy, childbirth and the puerperium (O00-O99)	76	0.5	0.6
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	62	0.4	0.5
...	All other causes (Residual)	2,221	15.6	18.5

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, both sexes, 35-39 years				
...	All causes	17,043	100.0	147.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	5,105	30.0	44.2
2	Malignant neoplasms (C00-C97)	2,366	13.9	20.5
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,331	13.7	20.2
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,141	12.6	18.5
5	Chronic liver disease and cirrhosis (K70,K73-K74)	442	2.6	3.8
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	392	2.3	3.4
7	Diabetes mellitus (E10-E14)	345	2.0	3.0
8	Cerebrovascular diseases (I60-I69)	261	1.5	2.3
9	Influenza and pneumonia (J09-J18)	199	1.2	1.7
10	Septicemia (A40-A41)	152	0.9	1.3
11	Human immunodeficiency virus (HIV) disease (B20-B24)	147	0.9	1.3
12	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	133	0.8	1.2
13	Chronic lower respiratory diseases (J40-J47)	112	0.7	1.0
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	108	0.6	0.9
15	Pregnancy, childbirth and the puerperium (O00-O99)	72	0.4	0.6
...	All other causes (Residual)	2,737	16.1	23.7
Non-Hispanic, white, both sexes, 40-44 years				
...	All causes	27,848	100.0	209.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	6,097	21.9	45.8
2	Malignant neoplasms (C00-C97)	5,097	18.3	38.3
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	4,438	15.9	33.3
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	3,033	10.9	22.8
5	Chronic liver disease and cirrhosis (K70,K73-K74)	1,149	4.1	8.6
6	Diabetes mellitus (E10-E14)	690	2.5	5.2
7	Cerebrovascular diseases (I60-I69)	563	2.0	4.2
8	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	444	1.6	3.3
9	Chronic lower respiratory diseases (J40-J47)	364	1.3	2.7
10	Influenza and pneumonia (J09-J18)	322	1.2	2.4
11	Septicemia (A40-A41)	299	1.1	2.2
12	Human immunodeficiency virus (HIV) disease (B20-B24)	278	1.0	2.1
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	192	0.7	1.4
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	163	0.6	1.2
15	Viral hepatitis (B15-B19)	141	0.5	1.1
...	All other causes (Residual)	4,578	16.4	34.4

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, both sexes, 45-49 years				
...	All causes	49,021	100.0	329.8
1	Malignant neoplasms (C00-C97)	11,502	23.5	77.4
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	9,093	18.5	61.2
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	7,740	15.8	52.1
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	3,730	7.6	25.1
5	Chronic liver disease and cirrhosis (K70,K73-K74)	2,473	5.0	16.6
6	Diabetes mellitus (E10-E14)	1,370	2.8	9.2
7	Cerebrovascular diseases (I60-I69)	1,135	2.3	7.6
8	Chronic lower respiratory diseases (J40-J47)	1,059	2.2	7.1
9	Influenza and pneumonia (J09-J18)	571	1.2	3.8
10	Septicemia (A40-A41)	525	1.1	3.5
11	Viral hepatitis (B15-B19)	507	1.0	3.4
12	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	503	1.0	3.4
13	Human immunodeficiency virus (HIV) disease (B20-B24)	464	0.9	3.1
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	343	0.7	2.3
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	269	0.5	1.8
...	All other causes (Residual)	7,737	15.8	52.1
Non-Hispanic, white, both sexes, 50-54 years				
...	All causes	78,815	100.0	493.0
1	Malignant neoplasms (C00-C97)	23,173	29.4	144.9
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	15,541	19.7	97.2
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	8,221	10.4	51.4
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	4,091	5.2	25.6
5	Chronic liver disease and cirrhosis (K70,K73-K74)	3,853	4.9	24.1
6	Chronic lower respiratory diseases (J40-J47)	2,639	3.3	16.5
7	Diabetes mellitus (E10-E14)	2,271	2.9	14.2
8	Cerebrovascular diseases (I60-I69)	1,951	2.5	12.2
9	Viral hepatitis (B15-B19)	1,068	1.4	6.7
10	Septicemia (A40-A41)	1,016	1.3	6.4
11	Influenza and pneumonia (J09-J18)	955	1.2	6.0
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	630	0.8	3.9
13	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	486	0.6	3.0
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	432	0.5	2.7
15	Human immunodeficiency virus (HIV) disease (B20-B24)	430	0.5	2.7
...	All other causes (Residual)	12,058	15.3	75.4

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, both sexes, 55-59 years				
...	All causes	104,196	100.0	700.2
1	Malignant neoplasms (C00-C97)	35,578	34.1	239.1
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	21,665	20.8	145.6
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	6,499	6.2	43.7
4	Chronic lower respiratory diseases (J40-J47)	4,743	4.6	31.9
5	Chronic liver disease and cirrhosis (K70,K73-K74)	4,297	4.1	28.9
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	3,437	3.3	23.1
7	Diabetes mellitus (E10-E14)	3,386	3.2	22.8
8	Cerebrovascular diseases (I60-I69)	2,745	2.6	18.4
9	Viral hepatitis (B15-B19)	1,414	1.4	9.5
10	Septicemia (A40-A41)	1,336	1.3	9.0
11	Influenza and pneumonia (J09-J18)	1,305	1.3	8.8
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,073	1.0	7.2
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	669	0.6	4.5
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	444	0.4	3.0
15	Aortic aneurysm and dissection (I71)	395	0.4	2.7
...	All other causes (Residual)	15,210	14.6	102.2
Non-Hispanic, white, both sexes, 60-64 years				
...	All causes	134,288	100.0	988.6
1	Malignant neoplasms (C00-C97)	49,663	37.0	365.6
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	28,799	21.4	212.0
3	Chronic lower respiratory diseases (J40-J47)	8,161	6.1	60.1
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	5,108	3.8	37.6
5	Diabetes mellitus (E10-E14)	4,770	3.6	35.1
6	Cerebrovascular diseases (I60-I69)	4,230	3.1	31.1
7	Chronic liver disease and cirrhosis (K70,K73-K74)	3,633	2.7	26.7
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,455	1.8	18.1
9	Septicemia (A40-A41)	1,966	1.5	14.5
10	Influenza and pneumonia (J09-J18)	1,752	1.3	12.9
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,698	1.3	12.5
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	901	0.7	6.6
13	Viral hepatitis (B15-B19)	897	0.7	6.6
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	678	0.5	5.0
15	Aortic aneurysm and dissection (I71)	589	0.4	4.3
...	All other causes (Residual)	18,988	14.1	139.8

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, both sexes, 65-69 years				
...	All causes	150,416	100.0	1,499.9
1	Malignant neoplasms (C00-C97)	56,142	37.3	559.8
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	31,627	21.0	315.4
3	Chronic lower respiratory diseases (J40-J47)	12,494	8.3	124.6
4	Cerebrovascular diseases (I60-I69)	5,252	3.5	52.4
5	Diabetes mellitus (E10-E14)	5,155	3.4	51.4
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,969	2.6	39.6
7	Chronic liver disease and cirrhosis (K70,K73-K74)	2,450	1.6	24.4
8	Septicemia (A40-A41)	2,195	1.5	21.9
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2,115	1.4	21.1
10	Influenza and pneumonia (J09-J18)	2,106	1.4	21.0
11	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,632	1.1	16.3
12	Alzheimer's disease (G30)	1,094	0.7	10.9
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,075	0.7	10.7
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	926	0.6	9.2
15	Aortic aneurysm and dissection (I71)	803	0.5	8.0
...	All other causes (Residual)	21,381	14.2	213.2
Non-Hispanic, white, both sexes, 70-74 years				
...	All causes	175,514	100.0	2,330.4
1	Malignant neoplasms (C00-C97)	60,672	34.6	805.6
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	37,030	21.1	491.7
3	Chronic lower respiratory diseases (J40-J47)	16,439	9.4	218.3
4	Cerebrovascular diseases (I60-I69)	7,367	4.2	97.8
5	Diabetes mellitus (E10-E14)	5,655	3.2	75.1
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	4,136	2.4	54.9
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2,947	1.7	39.1
8	Influenza and pneumonia (J09-J18)	2,854	1.6	37.9
9	Septicemia (A40-A41)	2,623	1.5	34.8
10	Alzheimer's disease (G30)	2,535	1.4	33.7
11	Chronic liver disease and cirrhosis (K70,K73-K74)	2,053	1.2	27.3
12	Parkinson's disease (G20-G21)	1,671	1.0	22.2
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,333	0.8	17.7
14	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,249	0.7	16.6
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1,204	0.7	16.0
...	All other causes (Residual)	25,746	14.7	341.8

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, both sexes, 75-79 years				
...	All causes	221,216	100.0	3,755.0
1	Malignant neoplasms (C00-C97)	63,735	28.8	1,081.8
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	49,285	22.3	836.6
3	Chronic lower respiratory diseases (J40-J47)	20,381	9.2	345.9
4	Cerebrovascular diseases (I60-I69)	11,701	5.3	198.6
5	Alzheimer's disease (G30)	6,745	3.0	114.5
6	Diabetes mellitus (E10-E14)	6,318	2.9	107.2
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	4,948	2.2	84.0
8	Influenza and pneumonia (J09-J18)	4,125	1.9	70.0
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	4,086	1.8	69.4
10	Parkinson's disease (G20-G21)	3,421	1.5	58.1
11	Septicemia (A40-A41)	3,366	1.5	57.1
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,878	0.8	31.9
13	Chronic liver disease and cirrhosis (K70,K73-K74)	1,708	0.8	29.0
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1,673	0.8	28.4
15	Pneumonitis due to solids and liquids (J69)	1,616	0.7	27.4
...	All other causes (Residual)	36,230	16.4	615.0
Non-Hispanic, white, both sexes, 80-84 years				
...	All causes	297,944	100.0	6,251.3
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	73,388	24.6	1,539.8
2	Malignant neoplasms (C00-C97)	65,506	22.0	1,374.4
3	Chronic lower respiratory diseases (J40-J47)	24,215	8.1	508.1
4	Cerebrovascular diseases (I60-I69)	18,403	6.2	386.1
5	Alzheimer's disease (G30)	14,197	4.8	297.9
6	Diabetes mellitus (E10-E14)	7,665	2.6	160.8
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	7,266	2.4	152.5
8	Influenza and pneumonia (J09-J18)	6,898	2.3	144.7
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	6,077	2.0	127.5
10	Parkinson's disease (G20-G21)	5,498	1.8	115.4
11	Septicemia (A40-A41)	4,247	1.4	89.1
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	3,069	1.0	64.4
13	Pneumonitis due to solids and liquids (J69)	2,766	0.9	58.0
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2,281	0.8	47.9
15	Aortic aneurysm and dissection (I71)	1,473	0.5	30.9
...	All other causes (Residual)	54,995	18.5	1,153.9

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, both sexes, 85-89 years				
...	All causes	333,072	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	92,997	27.9	---
2	Malignant neoplasms (C00-C97)	51,429	15.4	---
3	Cerebrovascular diseases (I60-I69)	22,525	6.8	---
4	Alzheimer's disease (G30)	21,413	6.4	---
5	Chronic lower respiratory diseases (J40-J47)	21,110	6.3	---
6	Influenza and pneumonia (J09-J18)	9,130	2.7	---
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	8,531	2.6	---
8	Diabetes mellitus (E10-E14)	7,293	2.2	---
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	7,052	2.1	---
10	Parkinson's disease (G20-G21)	5,405	1.6	---
11	Septicemia (A40-A41)	4,586	1.4	---
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	4,160	1.2	---
13	Pneumonitis due to solids and liquids (J69)	3,640	1.1	---
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2,468	0.7	---
15	Enterocolitis due to <i>Clostridium difficile</i> (A04.7)	1,675	0.5	---
...	All other causes (Residual)	69,658	20.9	---
Non-Hispanic, white, both sexes, 90-94 years				
...	All causes	241,951	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	75,639	31.3	---
2	Malignant neoplasms (C00-C97)	24,595	10.2	---
3	Alzheimer's disease (G30)	18,343	7.6	---
4	Cerebrovascular diseases (I60-I69)	16,853	7.0	---
5	Chronic lower respiratory diseases (J40-J47)	11,115	4.6	---
6	Influenza and pneumonia (J09-J18)	7,903	3.3	---
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	6,106	2.5	---
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	4,943	2.0	---
9	Diabetes mellitus (E10-E14)	4,150	1.7	---
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	3,804	1.6	---
11	Pneumonitis due to solids and liquids (J69)	3,051	1.3	---
12	Septicemia (A40-A41)	2,856	1.2	---
13	Parkinson's disease (G20-G21)	2,735	1.1	---
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1,399	0.6	---
15	Atherosclerosis (I70)	1,319	0.5	---
...	All other causes (Residual)	57,140	23.6	---

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, both sexes, 95-99 years				
...	All causes	96,269	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	32,002	33.2	---
2	Alzheimer's disease (G30)	7,921	8.2	---
3	Cerebrovascular diseases (I60-I69)	6,449	6.7	---
4	Malignant neoplasms (C00-C97)	6,192	6.4	---
5	Influenza and pneumonia (J09-J18)	3,686	3.8	---
6	Chronic lower respiratory diseases (J40-J47)	3,460	3.6	---
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,414	2.5	---
8	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,845	1.9	---
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,778	1.8	---
10	Diabetes mellitus (E10-E14)	1,230	1.3	---
11	Pneumonitis due to solids and liquids (J69)	1,163	1.2	---
12	Septicemia (A40-A41)	936	1.0	---
13	Atherosclerosis (I70)	683	0.7	---
14	Parkinson's disease (G20-G21)	596	0.6	---
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	472	0.5	---
...	All other causes (Residual)	25,442	26.4	---
Non-Hispanic, white, both sexes, 100 years and over				
...	All causes	19,148	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	6,714	35.1	---
2	Alzheimer's disease (G30)	1,426	7.4	---
3	Cerebrovascular diseases (I60-I69)	1,146	6.0	---
4	Influenza and pneumonia (J09-J18)	899	4.7	---
5	Malignant neoplasms (C00-C97)	724	3.8	---
6	Essential hypertension and hypertensive renal disease (I10,I12,I15)	518	2.7	---
7	Chronic lower respiratory diseases (J40-J47)	473	2.5	---
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	470	2.5	---
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	293	1.5	---
10	Pneumonitis due to solids and liquids (J69)	215	1.1	---
11	Atherosclerosis (I70)	193	1.0	---
12	Diabetes mellitus (E10-E14)	191	1.0	---
13	Septicemia (A40-A41)	160	0.8	---
14	Anemias (D50-D64)	84	0.4	---
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	65	0.3	---
...	All other causes (Residual)	5,577	29.1	---

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, male, all ages				
...	All causes	989,835	100.0	1,004.1
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	248,787	25.1	252.4
2	Malignant neoplasms (C00-C97)	242,140	24.5	245.6
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	60,043	6.1	60.9
4	Chronic lower respiratory diseases (J40-J47)	59,223	6.0	60.1
5	Cerebrovascular diseases (I60-I69)	39,753	4.0	40.3
6	Diabetes mellitus (E10-E14)	27,060	2.7	27.4
7	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	25,858	2.6	26.2
8	Alzheimer's disease (G30)	22,482	2.3	22.8
9	Influenza and pneumonia (J09-J18)	20,316	2.1	20.6
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	16,898	1.7	17.1
11	Chronic liver disease and cirrhosis (K70,K73-K74)	15,764	1.6	16.0
12	Septicemia (A40-A41)	12,319	1.2	12.5
13	Parkinson's disease (G20-G21)	12,134	1.2	12.3
14	Pneumonitis due to solids and liquids (J69)	8,242	0.8	8.4
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	7,926	0.8	8.0
...	All other causes (Residual)	170,890	17.3	173.4
Non-Hispanic, white, male, 1-4 years				
...	All causes	1,177	100.0	27.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	454	38.6	10.4
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	124	10.5	2.8
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	93	7.9	2.1
4	Malignant neoplasms (C00-C97)	89	7.6	2.0
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	35	3.0	0.8
6	Influenza and pneumonia (J09-J18)	22	1.9	0.5
7	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	16	1.4	*
8	Chronic lower respiratory diseases (J40-J47)	15	1.3	*
9	Septicemia (A40-A41)	12	1.0	*
10	Certain conditions originating in the perinatal period (P00-P96)	9	0.8	*
11	Cerebrovascular diseases (I60-I69)	8	0.7	*
12	Anemias (D50-D64)	4	0.3	*
12	Acute bronchitis and bronchiolitis (J20-J21)	4	0.3	*
12	Complications of medical and surgical care (Y40-Y84,Y88)	4	0.3	*
15	Meningitis (G00,G03)	3	0.3	*
...	All other causes (Residual)	285	24.2	6.5

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, male, 5-9 years				
...	All causes	746	100.0	13.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	260	34.9	4.6
2	Malignant neoplasms (C00-C97)	146	19.6	2.6
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	57	7.6	1.0
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	37	5.0	0.6
5	Influenza and pneumonia (J09-J18)	17	2.3	*
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	16	2.1	*
7	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	11	1.5	*
7	Chronic lower respiratory diseases (J40-J47)	11	1.5	*
9	Septicemia (A40-A41)	9	1.2	*
10	Cerebrovascular diseases (I60-I69)	8	1.1	*
11	Anemias (D50-D64)	4	0.5	*
11	Certain conditions originating in the perinatal period (P00-P96)	4	0.5	*
13	Meningococcal infection (A39)	3	0.4	*
13	Pneumonitis due to solids and liquids (J69)	3	0.4	*
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3	0.4	*
...	All other causes (Residual)	157	21.0	2.8
Non-Hispanic, white, male, 10-14 years				
...	All causes	981	100.0	16.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	353	36.0	5.9
2	Malignant neoplasms (C00-C97)	128	13.0	2.1
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	121	12.3	2.0
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	37	3.8	0.6
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	35	3.6	0.6
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	32	3.3	0.5
7	Influenza and pneumonia (J09-J18)	18	1.8	*
8	Septicemia (A40-A41)	15	1.5	*
9	Cerebrovascular diseases (I60-I69)	13	1.3	*
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	12	1.2	*
11	Chronic lower respiratory diseases (J40-J47)	6	0.6	*
12	Pneumonitis due to solids and liquids (J69)	5	0.5	*
13	Diabetes mellitus (E10-E14)	4	0.4	*
13	Certain conditions originating in the perinatal period (P00-P96)	4	0.4	*
15	Anemias (D50-D64)	2	0.2	*
15	Complications of medical and surgical care (Y40-Y84,Y88)	2	0.2	*
...	All other causes (Residual)	194	19.8	3.2

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, male, 15-19 years				
...	All causes	4,141	100.0	65.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,006	48.4	31.6
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,032	24.9	16.2
3	Malignant neoplasms (C00-C97)	205	5.0	3.2
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	170	4.1	2.7
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	117	2.8	1.8
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	76	1.8	1.2
7	Influenza and pneumonia (J09-J18)	28	0.7	0.4
8	Cerebrovascular diseases (I60-I69)	20	0.5	0.3
9	Septicemia (A40-A41)	18	0.4	*
10	Diabetes mellitus (E10-E14)	17	0.4	*
10	Chronic lower respiratory diseases (J40-J47)	17	0.4	*
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	11	0.3	*
13	Pneumonitis due to solids and liquids (J69)	4	0.1	*
13	Legal intervention (Y35,Y89.0)	4	0.1	*
15	Meningitis (G00,G03)	3	0.1	*
15	Peptic ulcer (K25-K28)	3	0.1	*
15	Certain conditions originating in the perinatal period (P00-P96)	3	0.1	*
15	Complications of medical and surgical care (Y40-Y84,Y88)	3	0.1	*
...	All other causes (Residual)	404	9.8	6.4
Non-Hispanic, white, male, 20-24 years				
...	All causes	8,081	100.0	123.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	4,243	52.5	64.9
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,785	22.1	27.3
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	327	4.0	5.0
4	Malignant neoplasms (C00-C97)	312	3.9	4.8
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	239	3.0	3.7
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	88	1.1	1.3
7	Influenza and pneumonia (J09-J18)	53	0.7	0.8
8	Diabetes mellitus (E10-E14)	50	0.6	0.8
9	Cerebrovascular diseases (I60-I69)	31	0.4	0.5
10	Septicemia (A40-A41)	30	0.4	0.5
10	Chronic lower respiratory diseases (J40-J47)	30	0.4	0.5
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	16	0.2	*
12	Pneumonitis due to solids and liquids (J69)	16	0.2	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	15	0.2	*
15	Legal intervention (Y35,Y89.0)	14	0.2	*
...	All other causes (Residual)	832	10.3	12.7

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, male, 25-29 years				
...	All causes	8,869	100.0	140.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	4,264	48.1	67.5
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,855	20.9	29.4
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	458	5.2	7.3
4	Malignant neoplasms (C00-C97)	433	4.9	6.9
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	353	4.0	5.6
6	Influenza and pneumonia (J09-J18)	78	0.9	1.2
7	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	73	0.8	1.2
8	Diabetes mellitus (E10-E14)	69	0.8	1.1
9	Septicemia (A40-A41)	47	0.5	0.7
10	Chronic liver disease and cirrhosis (K70,K73-K74)	44	0.5	0.7
11	Cerebrovascular diseases (I60-I69)	38	0.4	0.6
12	Chronic lower respiratory diseases (J40-J47)	30	0.3	0.5
12	Legal intervention (Y35,Y89.0)	30	0.3	0.5
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	28	0.3	0.4
15	Pneumonitis due to solids and liquids (J69)	19	0.2	*
...	All other causes (Residual)	1,050	11.8	16.6
Non-Hispanic, white, male, 30-34 years				
...	All causes	9,422	100.0	155.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,853	40.9	63.7
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,787	19.0	29.5
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	783	8.3	12.9
4	Malignant neoplasms (C00-C97)	605	6.4	10.0
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	309	3.3	5.1
6	Diabetes mellitus (E10-E14)	146	1.5	2.4
7	Chronic liver disease and cirrhosis (K70,K73-K74)	128	1.4	2.1
8	Influenza and pneumonia (J09-J18)	96	1.0	1.6
9	Cerebrovascular diseases (I60-I69)	90	1.0	1.5
10	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	87	0.9	1.4
11	Human immunodeficiency virus (HIV) disease (B20-B24)	64	0.7	1.1
12	Septicemia (A40-A41)	62	0.7	1.0
13	Chronic lower respiratory diseases (J40-J47)	46	0.5	0.8
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	35	0.4	0.6
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	35	0.4	0.6
...	All other causes (Residual)	1,296	13.8	21.4

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, male, 35-39 years				
...	All causes	10,839	100.0	186.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,471	32.0	59.7
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,776	16.4	30.6
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,518	14.0	26.1
4	Malignant neoplasms (C00-C97)	1,084	10.0	18.7
5	Chronic liver disease and cirrhosis (K70,K73-K74)	272	2.5	4.7
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	262	2.4	4.5
7	Diabetes mellitus (E10-E14)	216	2.0	3.7
8	Cerebrovascular diseases (I60-I69)	129	1.2	2.2
9	Influenza and pneumonia (J09-J18)	110	1.0	1.9
10	Human immunodeficiency virus (HIV) disease (B20-B24)	105	1.0	1.8
11	Septicemia (A40-A41)	75	0.7	1.3
12	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	73	0.7	1.3
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	71	0.7	1.2
14	Aortic aneurysm and dissection (I71)	58	0.5	1.0
15	Chronic lower respiratory diseases (J40-J47)	48	0.4	0.8
...	All other causes (Residual)	1,571	14.5	27.0
Non-Hispanic, white, male, 40-44 years				
...	All causes	17,124	100.0	256.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	4,026	23.5	60.3
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,146	18.4	47.1
3	Malignant neoplasms (C00-C97)	2,317	13.5	34.7
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,291	13.4	34.3
5	Chronic liver disease and cirrhosis (K70,K73-K74)	708	4.1	10.6
6	Diabetes mellitus (E10-E14)	424	2.5	6.3
7	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	299	1.7	4.5
8	Cerebrovascular diseases (I60-I69)	293	1.7	4.4
9	Human immunodeficiency virus (HIV) disease (B20-B24)	221	1.3	3.3
10	Chronic lower respiratory diseases (J40-J47)	176	1.0	2.6
11	Influenza and pneumonia (J09-J18)	161	0.9	2.4
12	Septicemia (A40-A41)	142	0.8	2.1
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	112	0.7	1.7
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	93	0.5	1.4
15	Viral hepatitis (B15-B19)	85	0.5	1.3
...	All other causes (Residual)	2,630	15.4	39.4

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, male, 45-49 years				
...	All causes	30,022	100.0	405.7
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	6,579	21.9	88.9
2	Malignant neoplasms (C00-C97)	5,532	18.4	74.7
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	5,094	17.0	68.8
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,777	9.2	37.5
5	Chronic liver disease and cirrhosis (K70,K73-K74)	1,561	5.2	21.1
6	Diabetes mellitus (E10-E14)	870	2.9	11.8
7	Cerebrovascular diseases (I60-I69)	615	2.0	8.3
8	Chronic lower respiratory diseases (J40-J47)	471	1.6	6.4
9	Human immunodeficiency virus (HIV) disease (B20-B24)	393	1.3	5.3
10	Viral hepatitis (B15-B19)	360	1.2	4.9
11	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	340	1.1	4.6
12	Influenza and pneumonia (J09-J18)	323	1.1	4.4
13	Septicemia (A40-A41)	262	0.9	3.5
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	210	0.7	2.8
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	188	0.6	2.5
...	All other causes (Residual)	4,447	14.8	60.1
Non-Hispanic, white, male, 50-54 years				
...	All causes	48,443	100.0	612.0
1	Malignant neoplasms (C00-C97)	11,989	24.7	151.5
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	11,352	23.4	143.4
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	5,393	11.1	68.1
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	3,062	6.3	38.7
5	Chronic liver disease and cirrhosis (K70,K73-K74)	2,547	5.3	32.2
6	Diabetes mellitus (E10-E14)	1,479	3.1	18.7
7	Chronic lower respiratory diseases (J40-J47)	1,262	2.6	15.9
8	Cerebrovascular diseases (I60-I69)	1,102	2.3	13.9
9	Viral hepatitis (B15-B19)	745	1.5	9.4
10	Septicemia (A40-A41)	568	1.2	7.2
11	Influenza and pneumonia (J09-J18)	553	1.1	7.0
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	370	0.8	4.7
13	Human immunodeficiency virus (HIV) disease (B20-B24)	355	0.7	4.5
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	321	0.7	4.1
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	273	0.6	3.4
...	All other causes (Residual)	7,072	14.6	89.3

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, male, 55-59 years				
...	All causes	64,979	100.0	890.2
1	Malignant neoplasms (C00-C97)	19,801	30.5	271.3
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	15,791	24.3	216.3
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	4,429	6.8	60.7
4	Chronic liver disease and cirrhosis (K70,K73-K74)	3,048	4.7	41.8
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,638	4.1	36.1
6	Chronic lower respiratory diseases (J40-J47)	2,481	3.8	34.0
7	Diabetes mellitus (E10-E14)	2,085	3.2	28.6
8	Cerebrovascular diseases (I60-I69)	1,594	2.5	21.8
9	Viral hepatitis (B15-B19)	1,071	1.6	14.7
10	Influenza and pneumonia (J09-J18)	767	1.2	10.5
11	Septicemia (A40-A41)	726	1.1	9.9
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	631	1.0	8.6
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	449	0.7	6.2
14	Aortic aneurysm and dissection (I71)	320	0.5	4.4
15	Human immunodeficiency virus (HIV) disease (B20-B24)	296	0.5	4.1
...	All other causes (Residual)	8,852	13.6	121.3
Non-Hispanic, white, male, 60-64 years				
...	All causes	81,044	100.0	1,226.4
1	Malignant neoplasms (C00-C97)	27,734	34.2	419.7
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	20,090	24.8	304.0
3	Chronic lower respiratory diseases (J40-J47)	4,236	5.2	64.1
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,484	4.3	52.7
5	Diabetes mellitus (E10-E14)	2,926	3.6	44.3
6	Chronic liver disease and cirrhosis (K70,K73-K74)	2,534	3.1	38.3
7	Cerebrovascular diseases (I60-I69)	2,340	2.9	35.4
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,885	2.3	28.5
9	Influenza and pneumonia (J09-J18)	1,015	1.3	15.4
10	Septicemia (A40-A41)	1,007	1.2	15.2
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	957	1.2	14.5
12	Viral hepatitis (B15-B19)	654	0.8	9.9
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	567	0.7	8.6
14	Aortic aneurysm and dissection (I71)	432	0.5	6.5
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	397	0.5	6.0
...	All other causes (Residual)	10,786	13.3	163.2

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, male, 65-69 years				
...	All causes	87,010	100.0	1,812.3
1	Malignant neoplasms (C00-C97)	30,974	35.6	645.1
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	21,016	24.2	437.7
3	Chronic lower respiratory diseases (J40-J47)	6,288	7.2	131.0
4	Diabetes mellitus (E10-E14)	3,124	3.6	65.1
5	Cerebrovascular diseases (I60-I69)	2,828	3.3	58.9
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,550	2.9	53.1
7	Chronic liver disease and cirrhosis (K70,K73-K74)	1,603	1.8	33.4
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,283	1.5	26.7
9	Influenza and pneumonia (J09-J18)	1,187	1.4	24.7
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,170	1.3	24.4
11	Septicemia (A40-A41)	1,143	1.3	23.8
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	616	0.7	12.8
13	Aortic aneurysm and dissection (I71)	567	0.7	11.8
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	558	0.6	11.6
15	Parkinson's disease (G20-G21)	513	0.6	10.7
...	All other causes (Residual)	11,590	13.3	241.4
Non-Hispanic, white, male, 70-74 years				
...	All causes	98,011	100.0	2,795.8
1	Malignant neoplasms (C00-C97)	33,463	34.1	954.5
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	23,088	23.6	658.6
3	Chronic lower respiratory diseases (J40-J47)	8,220	8.4	234.5
4	Cerebrovascular diseases (I60-I69)	3,797	3.9	108.3
5	Diabetes mellitus (E10-E14)	3,237	3.3	92.3
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,531	2.6	72.2
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,606	1.6	45.8
8	Influenza and pneumonia (J09-J18)	1,572	1.6	44.8
9	Septicemia (A40-A41)	1,346	1.4	38.4
10	Chronic liver disease and cirrhosis (K70,K73-K74)	1,199	1.2	34.2
11	Parkinson's disease (G20-G21)	1,125	1.1	32.1
12	Alzheimer's disease (G30)	1,064	1.1	30.4
13	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,034	1.1	29.5
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	718	0.7	20.5
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	644	0.7	18.4
...	All other causes (Residual)	13,367	13.6	381.3

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, male, 75-79 years				
...	All causes	116,832	100.0	4,457.5
1	Malignant neoplasms (C00-C97)	34,632	29.6	1,321.3
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	28,374	24.3	1,082.6
3	Chronic lower respiratory diseases (J40-J47)	9,800	8.4	373.9
4	Cerebrovascular diseases (I60-I69)	5,451	4.7	208.0
5	Diabetes mellitus (E10-E14)	3,411	2.9	130.1
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,781	2.4	106.1
7	Alzheimer's disease (G30)	2,735	2.3	104.3
8	Parkinson's disease (G20-G21)	2,226	1.9	84.9
9	Influenza and pneumonia (J09-J18)	2,218	1.9	84.6
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2,204	1.9	84.1
11	Septicemia (A40-A41)	1,662	1.4	63.4
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	949	0.8	36.2
13	Chronic liver disease and cirrhosis (K70,K73-K74)	941	0.8	35.9
14	Pneumonitis due to solids and liquids (J69)	918	0.8	35.0
15	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	881	0.8	33.6
...	All other causes (Residual)	17,649	15.1	673.4
Non-Hispanic, white, male, 80-84 years				
...	All causes	145,300	100.0	7,429.0
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	38,125	26.2	1,949.3
2	Malignant neoplasms (C00-C97)	34,395	23.7	1,758.6
3	Chronic lower respiratory diseases (J40-J47)	11,367	7.8	581.2
4	Cerebrovascular diseases (I60-I69)	7,590	5.2	388.1
5	Alzheimer's disease (G30)	5,187	3.6	265.2
6	Diabetes mellitus (E10-E14)	3,813	2.6	195.0
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,780	2.6	193.3
8	Influenza and pneumonia (J09-J18)	3,404	2.3	174.0
9	Parkinson's disease (G20-G21)	3,292	2.3	168.3
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3,242	2.2	165.8
11	Septicemia (A40-A41)	2,026	1.4	103.6
12	Pneumonitis due to solids and liquids (J69)	1,556	1.1	79.6
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,287	0.9	65.8
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1,247	0.9	63.8
15	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	858	0.6	43.9
...	All other causes (Residual)	24,131	16.6	1,233.8

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, male, 85-89 years				
...	All causes	139,628	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	40,897	29.3	---
2	Malignant neoplasms (C00-C97)	25,033	17.9	---
3	Chronic lower respiratory diseases (J40-J47)	9,249	6.6	---
4	Cerebrovascular diseases (I60-I69)	7,707	5.5	---
5	Alzheimer's disease (G30)	6,561	4.7	---
6	Influenza and pneumonia (J09-J18)	4,180	3.0	---
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,764	2.7	---
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3,362	2.4	---
9	Diabetes mellitus (E10-E14)	3,201	2.3	---
10	Parkinson's disease (G20-G21)	3,063	2.2	---
11	Pneumonitis due to solids and liquids (J69)	1,956	1.4	---
12	Septicemia (A40-A41)	1,816	1.3	---
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,407	1.0	---
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1,249	0.9	---
15	Aortic aneurysm and dissection (I71)	657	0.5	---
...	All other causes (Residual)	25,526	18.3	---
Non-Hispanic, white, male, 90-94 years				
...	All causes	83,837	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	27,526	32.8	---
2	Malignant neoplasms (C00-C97)	10,847	12.9	---
3	Alzheimer's disease (G30)	4,631	5.5	---
4	Cerebrovascular diseases (I60-I69)	4,586	5.5	---
5	Chronic lower respiratory diseases (J40-J47)	4,282	5.1	---
6	Influenza and pneumonia (J09-J18)	3,080	3.7	---
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,215	2.6	---
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2,192	2.6	---
9	Diabetes mellitus (E10-E14)	1,601	1.9	---
10	Pneumonitis due to solids and liquids (J69)	1,446	1.7	---
11	Parkinson's disease (G20-G21)	1,368	1.6	---
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,046	1.2	---
13	Septicemia (A40-A41)	1,031	1.2	---
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	584	0.7	---
15	Atherosclerosis (I70)	412	0.5	---
...	All other causes (Residual)	16,990	20.3	---

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, male, 95-99 years				
...	All causes	24,149	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	8,409	34.8	---
2	Malignant neoplasms (C00-C97)	2,208	9.1	---
3	Cerebrovascular diseases (I60-I69)	1,329	5.5	---
4	Alzheimer's disease (G30)	1,316	5.4	---
5	Influenza and pneumonia (J09-J18)	1,181	4.9	---
6	Chronic lower respiratory diseases (J40-J47)	1,071	4.4	---
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	677	2.8	---
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	617	2.6	---
9	Pneumonitis due to solids and liquids (J69)	466	1.9	---
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	386	1.6	---
11	Diabetes mellitus (E10-E14)	346	1.4	---
12	Septicemia (A40-A41)	252	1.0	---
13	Parkinson's disease (G20-G21)	233	1.0	---
14	Atherosclerosis (I70)	158	0.7	---
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	132	0.5	---
...	All other causes (Residual)	5,368	22.2	---
Non-Hispanic, white, male, 100 years and over				
...	All causes	3,074	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,121	36.5	---
2	Influenza and pneumonia (J09-J18)	198	6.4	---
3	Malignant neoplasms (C00-C97)	190	6.2	---
4	Alzheimer's disease (G30)	143	4.7	---
5	Cerebrovascular diseases (I60-I69)	142	4.6	---
6	Chronic lower respiratory diseases (J40-J47)	106	3.4	---
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	83	2.7	---
8	Essential hypertension and hypertensive renal disease (I10,I12,I15)	70	2.3	---
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	67	2.2	---
10	Pneumonitis due to solids and liquids (J69)	65	2.1	---
11	Diabetes mellitus (E10-E14)	38	1.2	---
12	Septicemia (A40-A41)	37	1.2	---
13	Atherosclerosis (I70)	29	0.9	---
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	20	0.7	---
15	Aortic aneurysm and dissection (I71)	15	0.5	---
...	All other causes (Residual)	750	24.4	---

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, female, all ages				
...	All causes	1,016,484	100.0	998.1
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	234,192	23.0	230.0
2	Malignant neoplasms (C00-C97)	217,886	21.4	213.9
3	Chronic lower respiratory diseases (J40-J47)	67,812	6.7	66.6
4	Cerebrovascular diseases (I60-I69)	61,290	6.0	60.2
5	Alzheimer's disease (G30)	51,966	5.1	51.0
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	38,562	3.8	37.9
7	Diabetes mellitus (E10-E14)	23,904	2.4	23.5
8	Influenza and pneumonia (J09-J18)	23,034	2.3	22.6
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	16,594	1.6	16.3
10	Septicemia (A40-A41)	14,350	1.4	14.1
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	12,227	1.2	12.0
12	Chronic liver disease and cirrhosis (K70,K73-K74)	8,988	0.9	8.8
13	Parkinson's disease (G20-G21)	8,491	0.8	8.3
14	Pneumonitis due to solids and liquids (J69)	7,271	0.7	7.1
15	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	7,173	0.7	7.0
...	All other causes (Residual)	222,744	21.9	218.7
Non-Hispanic, white, female, 1-4 years				
...	All causes	878	100.0	21.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	270	30.8	6.5
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	112	12.8	2.7
3	Malignant neoplasms (C00-C97)	94	10.7	2.3
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	57	6.5	1.4
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	39	4.4	0.9
6	Influenza and pneumonia (J09-J18)	23	2.6	0.6
7	Septicemia (A40-A41)	18	2.1	*
8	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	8	0.9	*
8	Certain conditions originating in the perinatal period (P00-P96)	8	0.9	*
10	Cerebrovascular diseases (I60-I69)	6	0.7	*
11	Acute bronchitis and bronchiolitis (J20-J21)	4	0.5	*
11	Chronic lower respiratory diseases (J40-J47)	4	0.5	*
13	Diabetes mellitus (E10-E14)	3	0.3	*
13	Complications of medical and surgical care (Y40-Y84,Y88)	3	0.3	*
15	Anemias (D50-D64)	2	0.2	*
15	Meningitis (G00,G03)	2	0.2	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2	0.2	*
...	All other causes (Residual)	223	25.4	5.4

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, female, 5-9 years				
...	All causes	554	100.0	10.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	168	30.3	3.1
2	Malignant neoplasms (C00-C97)	108	19.5	2.0
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	30	5.4	0.6
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	25	4.5	0.5
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	24	4.3	0.4
6	Septicemia (A40-A41)	16	2.9	*
7	Influenza and pneumonia (J09-J18)	15	2.7	*
8	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	13	2.3	*
9	Chronic lower respiratory diseases (J40-J47)	7	1.3	*
10	Certain conditions originating in the perinatal period (P00-P96)	6	1.1	*
11	Cerebrovascular diseases (I60-I69)	5	0.9	*
12	Anemias (D50-D64)	3	0.5	*
13	Meningitis (G00,G03)	2	0.4	*
14	Tuberculosis (A16-A19)	1	0.2	*
14	Atherosclerosis (I70)	1	0.2	*
14	Acute bronchitis and bronchiolitis (J20-J21)	1	0.2	*
14	Hernia (K40-K46)	1	0.2	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1	0.2	*
14	Enterocolitis due to <i>Clostridium difficile</i> (A04.7)	1	0.2	*
...	All other causes (Residual)	126	22.7	2.3
Non-Hispanic, white, female, 10-14 years				
...	All causes	627	100.0	11.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	192	30.6	3.4
2	Malignant neoplasms (C00-C97)	99	15.8	1.7
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	47	7.5	0.8
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	44	7.0	0.8
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	29	4.6	0.5
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	24	3.8	0.4
7	Cerebrovascular diseases (I60-I69)	12	1.9	*
8	Influenza and pneumonia (J09-J18)	11	1.8	*
9	Diabetes mellitus (E10-E14)	10	1.6	*
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	7	1.1	*
11	Septicemia (A40-A41)	6	1.0	*
12	Chronic lower respiratory diseases (J40-J47)	5	0.8	*
13	Complications of medical and surgical care (Y40-Y84,Y88)	3	0.5	*
14	Meningitis (G00,G03)	2	0.3	*
14	Pneumonitis due to solids and liquids (J69)	2	0.3	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2	0.3	*
14	Certain conditions originating in the perinatal period (P00-P96)	2	0.3	*
...	All other causes (Residual)	130	20.7	2.3

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, female, 15-19 years				
...	All causes	1,814	100.0	30.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	905	49.9	15.1
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	244	13.5	4.1
3	Malignant neoplasms (C00-C97)	131	7.2	2.2
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	75	4.1	1.2
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	43	2.4	0.7
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	41	2.3	0.7
7	Influenza and pneumonia (J09-J18)	20	1.1	0.3
8	Cerebrovascular diseases (I60-I69)	16	0.9	*
9	Pregnancy, childbirth and the puerperium (O00-O99)	14	0.8	*
10	Diabetes mellitus (E10-E14)	13	0.7	*
11	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	12	0.7	*
12	Complications of medical and surgical care (Y40-Y84,Y88)	9	0.5	*
13	Septicemia (A40-A41)	8	0.4	*
14	Chronic lower respiratory diseases (J40-J47)	5	0.3	*
15	Pneumonitis due to solids and liquids (J69)	4	0.2	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	4	0.2	*
...	All other causes (Residual)	270	14.9	4.5
Non-Hispanic, white, female, 20-24 years				
...	All causes	2,925	100.0	46.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,374	47.0	21.7
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	352	12.0	5.6
3	Malignant neoplasms (C00-C97)	243	8.3	3.8
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	129	4.4	2.0
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	94	3.2	1.5
6	Pregnancy, childbirth and the puerperium (O00-O99)	59	2.0	0.9
7	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	47	1.6	0.7
8	Influenza and pneumonia (J09-J18)	37	1.3	0.6
9	Cerebrovascular diseases (I60-I69)	23	0.8	0.4
10	Septicemia (A40-A41)	21	0.7	0.3
11	Diabetes mellitus (E10-E14)	17	0.6	*
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	15	0.5	*
13	Chronic lower respiratory diseases (J40-J47)	12	0.4	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	9	0.3	*
15	Complications of medical and surgical care (Y40-Y84,Y88)	8	0.3	*
...	All other causes (Residual)	485	16.6	7.7

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, female, 25-29 years				
...	All causes	3,682	100.0	59.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,461	39.7	23.6
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	425	11.5	6.9
3	Malignant neoplasms (C00-C97)	355	9.6	5.7
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	200	5.4	3.2
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	151	4.1	2.4
6	Pregnancy, childbirth and the puerperium (O00-O99)	100	2.7	1.6
7	Influenza and pneumonia (J09-J18)	70	1.9	1.1
8	Diabetes mellitus (E10-E14)	54	1.5	0.9
9	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	51	1.4	0.8
10	Cerebrovascular diseases (I60-I69)	46	1.2	0.7
11	Chronic liver disease and cirrhosis (K70,K73-K74)	41	1.1	0.7
12	Septicemia (A40-A41)	30	0.8	0.5
13	Chronic lower respiratory diseases (J40-J47)	19	0.5	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	16	0.4	*
15	Human immunodeficiency virus (HIV) disease (B20-B24)	15	0.4	*
...	All other causes (Residual)	648	17.6	10.5
Non-Hispanic, white, female, 30-34 years				
...	All causes	4,781	100.0	80.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,604	33.5	26.9
2	Malignant neoplasms (C00-C97)	664	13.9	11.1
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	514	10.8	8.6
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	375	7.8	6.3
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	134	2.8	2.2
6	Diabetes mellitus (E10-E14)	89	1.9	1.5
7	Cerebrovascular diseases (I60-I69)	79	1.7	1.3
8	Chronic liver disease and cirrhosis (K70,K73-K74)	76	1.6	1.3
8	Pregnancy, childbirth and the puerperium (O00-O99)	76	1.6	1.3
10	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	70	1.5	1.2
11	Influenza and pneumonia (J09-J18)	66	1.4	1.1
12	Septicemia (A40-A41)	47	1.0	0.8
12	Chronic lower respiratory diseases (J40-J47)	47	1.0	0.8
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	27	0.6	0.5
15	Human immunodeficiency virus (HIV) disease (B20-B24)	23	0.5	0.4
...	All other causes (Residual)	890	18.6	14.9

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, female, 35-39 years				
...	All causes	6,204	100.0	107.9
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,634	26.3	28.4
2	Malignant neoplasms (C00-C97)	1,282	20.7	22.3
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	623	10.0	10.8
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	555	8.9	9.7
5	Chronic liver disease and cirrhosis (K70,K73-K74)	170	2.7	3.0
6	Cerebrovascular diseases (I60-I69)	132	2.1	2.3
7	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	130	2.1	2.3
8	Diabetes mellitus (E10-E14)	129	2.1	2.2
9	Influenza and pneumonia (J09-J18)	89	1.4	1.5
10	Septicemia (A40-A41)	77	1.2	1.3
11	Pregnancy, childbirth and the puerperium (O00-O99)	72	1.2	1.3
12	Chronic lower respiratory diseases (J40-J47)	64	1.0	1.1
13	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	60	1.0	1.0
14	Human immunodeficiency virus (HIV) disease (B20-B24)	42	0.7	0.7
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	37	0.6	0.6
...	All other causes (Residual)	1,108	17.9	19.3
Non-Hispanic, white, female, 40-44 years				
...	All causes	10,724	100.0	161.5
1	Malignant neoplasms (C00-C97)	2,780	25.9	41.9
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,071	19.3	31.2
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,292	12.0	19.5
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	742	6.9	11.2
5	Chronic liver disease and cirrhosis (K70,K73-K74)	441	4.1	6.6
6	Cerebrovascular diseases (I60-I69)	270	2.5	4.1
7	Diabetes mellitus (E10-E14)	266	2.5	4.0
8	Chronic lower respiratory diseases (J40-J47)	188	1.8	2.8
9	Influenza and pneumonia (J09-J18)	161	1.5	2.4
10	Septicemia (A40-A41)	157	1.5	2.4
11	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	145	1.4	2.2
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	80	0.7	1.2
13	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	70	0.7	1.1
14	Human immunodeficiency virus (HIV) disease (B20-B24)	57	0.5	0.9
15	Viral hepatitis (B15-B19)	56	0.5	0.8
...	All other causes (Residual)	1,948	18.2	29.3

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, female, 45-49 years				
...	All causes	18,999	100.0	254.6
1	Malignant neoplasms (C00-C97)	5,970	31.4	80.0
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,646	13.9	35.5
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,514	13.2	33.7
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	953	5.0	12.8
5	Chronic liver disease and cirrhosis (K70,K73-K74)	912	4.8	12.2
6	Chronic lower respiratory diseases (J40-J47)	588	3.1	7.9
7	Cerebrovascular diseases (I60-I69)	520	2.7	7.0
8	Diabetes mellitus (E10-E14)	500	2.6	6.7
9	Septicemia (A40-A41)	263	1.4	3.5
10	Influenza and pneumonia (J09-J18)	248	1.3	3.3
11	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	163	0.9	2.2
12	Viral hepatitis (B15-B19)	147	0.8	2.0
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	133	0.7	1.8
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	113	0.6	1.5
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	91	0.5	1.2
...	All other causes (Residual)	3,238	17.0	43.4
Non-Hispanic, white, female, 50-54 years				
...	All causes	30,372	100.0	376.3
1	Malignant neoplasms (C00-C97)	11,184	36.8	138.6
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	4,189	13.8	51.9
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,828	9.3	35.0
4	Chronic lower respiratory diseases (J40-J47)	1,377	4.5	17.1
5	Chronic liver disease and cirrhosis (K70,K73-K74)	1,306	4.3	16.2
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,029	3.4	12.7
7	Cerebrovascular diseases (I60-I69)	849	2.8	10.5
8	Diabetes mellitus (E10-E14)	792	2.6	9.8
9	Septicemia (A40-A41)	448	1.5	5.6
10	Influenza and pneumonia (J09-J18)	402	1.3	5.0
11	Viral hepatitis (B15-B19)	323	1.1	4.0
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	260	0.9	3.2
13	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	165	0.5	2.0
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	159	0.5	2.0
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	159	0.5	2.0
...	All other causes (Residual)	4,902	16.1	60.7

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, female, 55-59 years				
...	All causes	39,217	100.0	517.3
1	Malignant neoplasms (C00-C97)	15,777	40.2	208.1
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	5,874	15.0	77.5
3	Chronic lower respiratory diseases (J40-J47)	2,262	5.8	29.8
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,070	5.3	27.3
5	Diabetes mellitus (E10-E14)	1,301	3.3	17.2
6	Chronic liver disease and cirrhosis (K70,K73-K74)	1,249	3.2	16.5
7	Cerebrovascular diseases (I60-I69)	1,151	2.9	15.2
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	799	2.0	10.5
9	Septicemia (A40-A41)	610	1.6	8.0
10	Influenza and pneumonia (J09-J18)	538	1.4	7.1
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	442	1.1	5.8
12	Viral hepatitis (B15-B19)	343	0.9	4.5
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	220	0.6	2.9
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	194	0.5	2.6
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	192	0.5	2.5
...	All other causes (Residual)	6,195	15.8	81.7
Non-Hispanic, white, female, 60-64 years				
...	All causes	53,244	100.0	763.3
1	Malignant neoplasms (C00-C97)	21,929	41.2	314.4
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	8,709	16.4	124.8
3	Chronic lower respiratory diseases (J40-J47)	3,925	7.4	56.3
4	Cerebrovascular diseases (I60-I69)	1,890	3.5	27.1
5	Diabetes mellitus (E10-E14)	1,844	3.5	26.4
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,624	3.1	23.3
7	Chronic liver disease and cirrhosis (K70,K73-K74)	1,099	2.1	15.8
8	Septicemia (A40-A41)	959	1.8	13.7
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	741	1.4	10.6
10	Influenza and pneumonia (J09-J18)	737	1.4	10.6
11	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	570	1.1	8.2
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	334	0.6	4.8
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	281	0.5	4.0
14	Alzheimer's disease (G30)	268	0.5	3.8
15	Viral hepatitis (B15-B19)	243	0.5	3.5
...	All other causes (Residual)	8,091	15.2	116.0

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, female, 65-69 years				
...	All causes	63,406	100.0	1,212.9
1	Malignant neoplasms (C00-C97)	25,168	39.7	481.4
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	10,611	16.7	203.0
3	Chronic lower respiratory diseases (J40-J47)	6,206	9.8	118.7
4	Cerebrovascular diseases (I60-I69)	2,424	3.8	46.4
5	Diabetes mellitus (E10-E14)	2,031	3.2	38.9
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,419	2.2	27.1
7	Septicemia (A40-A41)	1,052	1.7	20.1
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	945	1.5	18.1
9	Influenza and pneumonia (J09-J18)	919	1.4	17.6
10	Chronic liver disease and cirrhosis (K70,K73-K74)	847	1.3	16.2
11	Alzheimer's disease (G30)	592	0.9	11.3
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	459	0.7	8.8
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	368	0.6	7.0
14	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	349	0.6	6.7
15	Parkinson's disease (G20-G21)	273	0.4	5.2
...	All other causes (Residual)	9,743	15.4	186.4
Non-Hispanic, white, female, 70-74 years				
...	All causes	77,503	100.0	1,925.1
1	Malignant neoplasms (C00-C97)	27,209	35.1	675.9
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	13,942	18.0	346.3
3	Chronic lower respiratory diseases (J40-J47)	8,219	10.6	204.2
4	Cerebrovascular diseases (I60-I69)	3,570	4.6	88.7
5	Diabetes mellitus (E10-E14)	2,418	3.1	60.1
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,605	2.1	39.9
7	Alzheimer's disease (G30)	1,471	1.9	36.5
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,341	1.7	33.3
9	Influenza and pneumonia (J09-J18)	1,282	1.7	31.8
10	Septicemia (A40-A41)	1,277	1.6	31.7
11	Chronic liver disease and cirrhosis (K70,K73-K74)	854	1.1	21.2
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	689	0.9	17.1
13	Parkinson's disease (G20-G21)	546	0.7	13.6
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	486	0.6	12.1
15	Pneumonitis due to solids and liquids (J69)	371	0.5	9.2
...	All other causes (Residual)	12,223	15.8	303.6

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, female, 75-79 years				
...	All causes	104,384	100.0	3,191.9
1	Malignant neoplasms (C00-C97)	29,103	27.9	889.9
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	20,911	20.0	639.4
3	Chronic lower respiratory diseases (J40-J47)	10,581	10.1	323.5
4	Cerebrovascular diseases (I60-I69)	6,250	6.0	191.1
5	Alzheimer's disease (G30)	4,010	3.8	122.6
6	Diabetes mellitus (E10-E14)	2,907	2.8	88.9
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,167	2.1	66.3
8	Influenza and pneumonia (J09-J18)	1,907	1.8	58.3
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,882	1.8	57.5
10	Septicemia (A40-A41)	1,704	1.6	52.1
11	Parkinson's disease (G20-G21)	1,195	1.1	36.5
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,018	1.0	31.1
13	Chronic liver disease and cirrhosis (K70,K73-K74)	767	0.7	23.5
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	724	0.7	22.1
15	Pneumonitis due to solids and liquids (J69)	698	0.7	21.3
...	All other causes (Residual)	18,560	17.8	567.5
Non-Hispanic, white, female, 80-84 years				
...	All causes	152,644	100.0	5,431.6
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	35,263	23.1	1,254.8
2	Malignant neoplasms (C00-C97)	31,111	20.4	1,107.0
3	Chronic lower respiratory diseases (J40-J47)	12,848	8.4	457.2
4	Cerebrovascular diseases (I60-I69)	10,813	7.1	384.8
5	Alzheimer's disease (G30)	9,010	5.9	320.6
6	Diabetes mellitus (E10-E14)	3,852	2.5	137.1
7	Influenza and pneumonia (J09-J18)	3,494	2.3	124.3
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,486	2.3	124.0
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2,835	1.9	100.9
10	Septicemia (A40-A41)	2,221	1.5	79.0
11	Parkinson's disease (G20-G21)	2,206	1.4	78.5
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,782	1.2	63.4
13	Pneumonitis due to solids and liquids (J69)	1,210	0.8	43.1
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1,034	0.7	36.8
15	Enterocolitis due to <i>Clostridium difficile</i> (A04.7)	791	0.5	28.1
...	All other causes (Residual)	30,688	20.1	1,092.0

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, female, 85-89 years				
...	All causes	193,444	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	52,100	26.9	---
2	Malignant neoplasms (C00-C97)	26,396	13.6	---
3	Alzheimer's disease (G30)	14,852	7.7	---
4	Cerebrovascular diseases (I60-I69)	14,818	7.7	---
5	Chronic lower respiratory diseases (J40-J47)	11,861	6.1	---
6	Influenza and pneumonia (J09-J18)	4,950	2.6	---
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	4,767	2.5	---
8	Diabetes mellitus (E10-E14)	4,092	2.1	---
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3,690	1.9	---
10	Septicemia (A40-A41)	2,770	1.4	---
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	2,753	1.4	---
12	Parkinson's disease (G20-G21)	2,342	1.2	---
13	Pneumonitis due to solids and liquids (J69)	1,684	0.9	---
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown	1,219	0.6	---
15	Enterocolitis due to <i>Clostridium difficile</i> (A04.7)	1,103	0.6	---
...	All other causes (Residual)	44,047	22.8	---
Non-Hispanic, white, female, 90-94 years				
...	All causes	158,114	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	48,113	30.4	---
2	Malignant neoplasms (C00-C97)	13,748	8.7	---
3	Alzheimer's disease (G30)	13,712	8.7	---
4	Cerebrovascular diseases (I60-I69)	12,267	7.8	---
5	Chronic lower respiratory diseases (J40-J47)	6,833	4.3	---
6	Influenza and pneumonia (J09-J18)	4,823	3.1	---
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,891	2.5	---
8	Essential hypertension and hypertensive renal disease (I10,I12,I15)	2,758	1.7	---
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2,751	1.7	---
10	Diabetes mellitus (E10-E14)	2,549	1.6	---
11	Septicemia (A40-A41)	1,825	1.2	---
12	Pneumonitis due to solids and liquids (J69)	1,605	1.0	---
13	Parkinson's disease (G20-G21)	1,367	0.9	---
14	Atherosclerosis (I70)	907	0.6	---
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	815	0.5	---
...	All other causes (Residual)	40,150	25.4	---

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, female, 95-99 years				
...	All causes	72,120	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	23,593	32.7	---
2	Alzheimer's disease (G30)	6,605	9.2	---
3	Cerebrovascular diseases (I60-I69)	5,120	7.1	---
4	Malignant neoplasms (C00-C97)	3,984	5.5	---
5	Influenza and pneumonia (J09-J18)	2,505	3.5	---
6	Chronic lower respiratory diseases (J40-J47)	2,389	3.3	---
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,737	2.4	---
8	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,459	2.0	---
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,161	1.6	---
10	Diabetes mellitus (E10-E14)	884	1.2	---
11	Pneumonitis due to solids and liquids (J69)	697	1.0	---
12	Septicemia (A40-A41)	684	0.9	---
13	Atherosclerosis (I70)	525	0.7	---
14	Parkinson's disease (G20-G21)	363	0.5	---
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	340	0.5	---
...	All other causes (Residual)	20,074	27.8	---
Non-Hispanic, white, female, 100 years and over				
...	All causes	16,074	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	5,593	34.8	---
2	Alzheimer's disease (G30)	1,283	8.0	---
3	Cerebrovascular diseases (I60-I69)	1,004	6.2	---
4	Influenza and pneumonia (J09-J18)	701	4.4	---
5	Malignant neoplasms (C00-C97)	534	3.3	---
6	Essential hypertension and hypertensive renal disease (I10,I12,I15)	448	2.8	---
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	387	2.4	---
8	Chronic lower respiratory diseases (J40-J47)	367	2.3	---
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	226	1.4	---
10	Atherosclerosis (I70)	164	1.0	---
11	Diabetes mellitus (E10-E14)	153	1.0	---
12	Pneumonitis due to solids and liquids (J69)	150	0.9	---
13	Septicemia (A40-A41)	123	0.8	---
14	Anemias (D50-D64)	70	0.4	---
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	45	0.3	---
...	All other causes (Residual)	4,826	30.0	---

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, both sexes, all ages				
...	All causes	286,797	100.0	718.0
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	67,595	23.6	169.2
2	Malignant neoplasms (C00-C97)	66,158	23.1	165.6
3	Cerebrovascular diseases (I60-I69)	15,722	5.5	39.4
4	Diabetes mellitus (E10-E14)	12,771	4.5	32.0
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	12,299	4.3	30.8
6	Chronic lower respiratory diseases (J40-J47)	8,994	3.1	22.5
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	8,044	2.8	20.1
8	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	7,764	2.7	19.4
9	Septicemia (A40-A41)	5,963	2.1	14.9
10	Alzheimer's disease (G30)	5,446	1.9	13.6
11	Influenza and pneumonia (J09-J18)	5,098	1.8	12.8
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	4,965	1.7	12.4
13	Human immunodeficiency virus (HIV) disease (B20-B24)	4,138	1.4	10.4
14	Certain conditions originating in the perinatal period (P00-P96)	3,890	1.4	9.7
15	Chronic liver disease and cirrhosis (K70,K73-K74)	2,783	1.0	7.0
...	All other causes (Residual)	55,167	19.2	138.1
Non-Hispanic, black, both sexes, 1-4 years				
...	All causes	1,011	100.0	40.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	286	28.3	11.5
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	154	15.2	6.2
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	105	10.4	4.2
4	Malignant neoplasms (C00-C97)	53	5.2	2.1
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	52	5.1	2.1
6	Influenza and pneumonia (J09-J18)	32	3.2	1.3
7	Chronic lower respiratory diseases (J40-J47)	24	2.4	1.0
8	Cerebrovascular diseases (I60-I69)	17	1.7	*
9	Certain conditions originating in the perinatal period (P00-P96)	12	1.2	*
10	Septicemia (A40-A41)	11	1.1	*
11	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	10	1.0	*
12	Complications of medical and surgical care (Y40-Y84,Y88)	8	0.8	*
13	Anemias (D50-D64)	6	0.6	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3	0.3	*
15	Meningitis (G00,G03)	2	0.2	*
15	Acute bronchitis and bronchiolitis (J20-J21)	2	0.2	*
15	Pneumonitis due to solids and liquids (J69)	2	0.2	*
...	All other causes (Residual)	232	22.9	9.4

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, both sexes, 5-9 years				
...	All causes	527	100.0	17.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	157	29.8	5.2
2	Malignant neoplasms (C00-C97)	63	12.0	2.1
3	Chronic lower respiratory diseases (J40-J47)	36	6.8	1.2
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	36	6.8	1.2
5	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	35	6.6	1.2
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	21	4.0	0.7
7	Anemias (D50-D64)	15	2.8	*
8	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	12	2.3	*
8	Cerebrovascular diseases (I60-I69)	12	2.3	*
10	Influenza and pneumonia (J09-J18)	11	2.1	*
11	Septicemia (A40-A41)	6	1.1	*
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	5	0.9	*
13	Certain conditions originating in the perinatal period (P00-P96)	3	0.6	*
13	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	3	0.6	*
15	Meningitis (G00,G03)	2	0.4	*
15	Pneumonitis due to solids and liquids (J69)	2	0.4	*
15	Diseases of appendix (K35-K38)	2	0.4	*
...	All other causes (Residual)	106	20.1	3.5
Non-Hispanic, black, both sexes, 10-14 years				
...	All causes	623	100.0	19.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	156	25.0	5.0
2	Malignant neoplasms (C00-C97)	64	10.3	2.0
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	53	8.5	1.7
4	Chronic lower respiratory diseases (J40-J47)	45	7.2	1.4
5	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	43	6.9	1.4
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	40	6.4	1.3
7	Diseases of heart (I00-I09,I11,I13,I20-I51)	28	4.5	0.9
8	Influenza and pneumonia (J09-J18)	14	2.2	*
9	Cerebrovascular diseases (I60-I69)	8	1.3	*
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	7	1.1	*
10	Anemias (D50-D64)	7	1.1	*
12	Septicemia (A40-A41)	5	0.8	*
13	Diabetes mellitus (E10-E14)	4	0.6	*
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	4	0.6	*
15	Pneumonitis due to solids and liquids (J69)	3	0.5	*
15	Complications of medical and surgical care (Y40-Y84,Y88)	3	0.5	*
...	All other causes (Residual)	139	22.3	4.4

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, both sexes, 15-19 years				
...	All causes	2,335	100.0	68.4
1	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,012	43.3	29.7
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	540	23.1	15.8
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	159	6.8	4.7
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	98	4.2	2.9
5	Malignant neoplasms (C00-C97)	96	4.1	2.8
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	53	2.3	1.6
7	Chronic lower respiratory diseases (J40-J47)	36	1.5	1.1
8	Anemias (D50-D64)	18	0.8	*
9	Legal intervention (Y35,Y89.0)	16	0.7	*
10	Human immunodeficiency virus (HIV) disease (B20-B24)	13	0.6	*
11	Pregnancy, childbirth and the puerperium (O00-O99)	11	0.5	*
12	Diabetes mellitus (E10-E14)	10	0.4	*
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	9	0.4	*
13	Cerebrovascular diseases (I60-I69)	9	0.4	*
13	Influenza and pneumonia (J09-J18)	9	0.4	*
...	All other causes (Residual)	246	10.5	7.2
Non-Hispanic, black, both sexes, 20-24 years				
...	All causes	4,241	100.0	127.5
1	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,785	42.1	53.7
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	817	19.3	24.6
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	328	7.7	9.9
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	205	4.8	6.2
5	Malignant neoplasms (C00-C97)	161	3.8	4.8
6	Human immunodeficiency virus (HIV) disease (B20-B24)	87	2.1	2.6
7	Anemias (D50-D64)	53	1.2	1.6
8	Pregnancy, childbirth and the puerperium (O00-O99)	51	1.2	1.5
9	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	49	1.2	1.5
10	Chronic lower respiratory diseases (J40-J47)	45	1.1	1.4
11	Diabetes mellitus (E10-E14)	37	0.9	1.1
12	Influenza and pneumonia (J09-J18)	35	0.8	1.1
13	Legal intervention (Y35,Y89.0)	30	0.7	0.9
14	Septicemia (A40-A41)	28	0.7	0.8
15	Cerebrovascular diseases (I60-I69)	22	0.5	0.7
...	All other causes (Residual)	508	12.0	15.3

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, both sexes, 25-29 years				
...	All causes	4,246	100.0	149.8
1	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,346	31.7	47.5
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	833	19.6	29.4
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	329	7.7	11.6
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	300	7.1	10.6
5	Malignant neoplasms (C00-C97)	232	5.5	8.2
6	Human immunodeficiency virus (HIV) disease (B20-B24)	165	3.9	5.8
7	Diabetes mellitus (E10-E14)	79	1.9	2.8
8	Anemias (D50-D64)	63	1.5	2.2
9	Pregnancy, childbirth and the puerperium (O00-O99)	61	1.4	2.2
10	Influenza and pneumonia (J09-J18)	48	1.1	1.7
11	Septicemia (A40-A41)	44	1.0	1.6
12	Chronic lower respiratory diseases (J40-J47)	42	1.0	1.5
13	Cerebrovascular diseases (I60-I69)	40	0.9	1.4
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	39	0.9	1.4
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	30	0.7	1.1
...	All other causes (Residual)	595	14.0	21.0
Non-Hispanic, black, both sexes, 30-34 years				
...	All causes	4,711	100.0	173.8
1	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	956	20.3	35.3
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	799	17.0	29.5
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	612	13.0	22.6
4	Malignant neoplasms (C00-C97)	408	8.7	15.1
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	248	5.3	9.1
6	Human immunodeficiency virus (HIV) disease (B20-B24)	247	5.2	9.1
7	Diabetes mellitus (E10-E14)	147	3.1	5.4
8	Cerebrovascular diseases (I60-I69)	95	2.0	3.5
9	Anemias (D50-D64)	70	1.5	2.6
10	Pregnancy, childbirth and the puerperium (O00-O99)	63	1.3	2.3
11	Influenza and pneumonia (J09-J18)	58	1.2	2.1
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	55	1.2	2.0
13	Septicemia (A40-A41)	53	1.1	2.0
14	Chronic lower respiratory diseases (J40-J47)	48	1.0	1.8
15	Chronic liver disease and cirrhosis (K70,K73-K74)	37	0.8	1.4
...	All other causes (Residual)	815	17.3	30.1

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, both sexes, 35-39 years				
...	All causes	5,546	100.0	218.5
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	967	17.4	38.1
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	765	13.8	30.1
3	Malignant neoplasms (C00-C97)	728	13.1	28.7
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	649	11.7	25.6
5	Human immunodeficiency virus (HIV) disease (B20-B24)	346	6.2	13.6
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	224	4.0	8.8
7	Cerebrovascular diseases (I60-I69)	181	3.3	7.1
8	Diabetes mellitus (E10-E14)	160	2.9	6.3
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	106	1.9	4.2
10	Septicemia (A40-A41)	83	1.5	3.3
11	Chronic lower respiratory diseases (J40-J47)	74	1.3	2.9
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	71	1.3	2.8
13	Influenza and pneumonia (J09-J18)	61	1.1	2.4
14	Chronic liver disease and cirrhosis (K70,K73-K74)	56	1.0	2.2
15	Pregnancy, childbirth and the puerperium (O00-O99)	54	1.0	2.1
...	All other causes (Residual)	1,021	18.4	40.2
Non-Hispanic, black, both sexes, 40-44 years				
...	All causes	8,065	100.0	302.4
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,714	21.3	64.3
2	Malignant neoplasms (C00-C97)	1,361	16.9	51.0
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	894	11.1	33.5
4	Human immunodeficiency virus (HIV) disease (B20-B24)	543	6.7	20.4
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	474	5.9	17.8
6	Cerebrovascular diseases (I60-I69)	328	4.1	12.3
7	Diabetes mellitus (E10-E14)	327	4.1	12.3
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	193	2.4	7.2
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	164	2.0	6.1
10	Septicemia (A40-A41)	132	1.6	4.9
11	Chronic liver disease and cirrhosis (K70,K73-K74)	123	1.5	4.6
12	Influenza and pneumonia (J09-J18)	118	1.5	4.4
13	Chronic lower respiratory diseases (J40-J47)	117	1.5	4.4
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	115	1.4	4.3
15	Anemias (D50-D64)	58	0.7	2.2
...	All other causes (Residual)	1,404	17.4	52.6

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, both sexes, 45-49 years				
...	All causes	13,033	100.0	467.4
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,945	22.6	105.6
2	Malignant neoplasms (C00-C97)	2,859	21.9	102.5
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,110	8.5	39.8
4	Human immunodeficiency virus (HIV) disease (B20-B24)	681	5.2	24.4
5	Cerebrovascular diseases (I60-I69)	651	5.0	23.3
6	Diabetes mellitus (E10-E14)	540	4.1	19.4
7	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	423	3.2	15.2
8	Chronic liver disease and cirrhosis (K70,K73-K74)	276	2.1	9.9
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	268	2.1	9.6
10	Septicemia (A40-A41)	232	1.8	8.3
11	Chronic lower respiratory diseases (J40-J47)	218	1.7	7.8
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	203	1.6	7.3
13	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	197	1.5	7.1
14	Influenza and pneumonia (J09-J18)	149	1.1	5.3
15	Anemias (D50-D64)	97	0.7	3.5
...	All other causes (Residual)	2,184	16.8	78.3
Non-Hispanic, black, both sexes, 50-54 years				
...	All causes	20,118	100.0	735.6
1	Malignant neoplasms (C00-C97)	5,451	27.1	199.3
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	4,804	23.9	175.7
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,248	6.2	45.6
4	Cerebrovascular diseases (I60-I69)	980	4.9	35.8
5	Diabetes mellitus (E10-E14)	896	4.5	32.8
6	Human immunodeficiency virus (HIV) disease (B20-B24)	748	3.7	27.4
7	Chronic liver disease and cirrhosis (K70,K73-K74)	509	2.5	18.6
8	Chronic lower respiratory diseases (J40-J47)	475	2.4	17.4
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	457	2.3	16.7
10	Septicemia (A40-A41)	391	1.9	14.3
11	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	304	1.5	11.1
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	292	1.5	10.7
13	Influenza and pneumonia (J09-J18)	247	1.2	9.0
14	Viral hepatitis (B15-B19)	198	1.0	7.2
15	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	152	0.8	5.6
...	All other causes (Residual)	2,966	14.7	108.5

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, both sexes, 55-59 years				
...	All causes	25,203	100.0	1,096.9
1	Malignant neoplasms (C00-C97)	7,591	30.1	330.4
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	6,079	24.1	264.6
3	Cerebrovascular diseases (I60-I69)	1,265	5.0	55.1
4	Diabetes mellitus (E10-E14)	1,225	4.9	53.3
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,192	4.7	51.9
6	Chronic lower respiratory diseases (J40-J47)	638	2.5	27.8
7	Human immunodeficiency virus (HIV) disease (B20-B24)	609	2.4	26.5
7	Chronic liver disease and cirrhosis (K70,K73-K74)	609	2.4	26.5
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	587	2.3	25.5
10	Septicemia (A40-A41)	544	2.2	23.7
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	394	1.6	17.1
12	Viral hepatitis (B15-B19)	327	1.3	14.2
13	Influenza and pneumonia (J09-J18)	323	1.3	14.1
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	217	0.9	9.4
15	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	120	0.5	5.2
...	All other causes (Residual)	3,483	13.8	151.6
Non-Hispanic, black, both sexes, 60-64 years				
...	All causes	28,093	100.0	1,537.9
1	Malignant neoplasms (C00-C97)	9,216	32.8	504.5
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	6,943	24.7	380.1
3	Diabetes mellitus (E10-E14)	1,520	5.4	83.2
4	Cerebrovascular diseases (I60-I69)	1,425	5.1	78.0
5	Chronic lower respiratory diseases (J40-J47)	831	3.0	45.5
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	830	3.0	45.4
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	809	2.9	44.3
8	Septicemia (A40-A41)	575	2.0	31.5
9	Chronic liver disease and cirrhosis (K70,K73-K74)	540	1.9	29.6
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	450	1.6	24.6
11	Influenza and pneumonia (J09-J18)	377	1.3	20.6
12	Human immunodeficiency virus (HIV) disease (B20-B24)	370	1.3	20.3
13	Viral hepatitis (B15-B19)	270	1.0	14.8
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	114	0.4	6.2
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	112	0.4	6.1
...	All other causes (Residual)	3,711	13.2	203.1

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, both sexes, 65-69 years				
...	All causes	25,884	100.0	2,148.5
1	Malignant neoplasms (C00-C97)	8,496	32.8	705.2
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	6,449	24.9	535.3
3	Diabetes mellitus (E10-E14)	1,510	5.8	125.3
4	Cerebrovascular diseases (I60-I69)	1,362	5.3	113.1
5	Chronic lower respiratory diseases (J40-J47)	1,005	3.9	83.4
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	820	3.2	68.1
7	Septicemia (A40-A41)	580	2.2	48.1
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	540	2.1	44.8
9	Essential hypertension and hypertensive renal disease (I10,I12,I15)	437	1.7	36.3
10	Influenza and pneumonia (J09-J18)	379	1.5	31.5
11	Chronic liver disease and cirrhosis (K70,K73-K74)	256	1.0	21.2
12	Human immunodeficiency virus (HIV) disease (B20-B24)	176	0.7	14.6
13	Pneumonitis due to solids and liquids (J69)	124	0.5	10.3
14	Alzheimer's disease (G30)	118	0.5	9.8
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	99	0.4	8.2
...	All other causes (Residual)	3,533	13.6	293.3
Non-Hispanic, black, both sexes, 70-74 years				
...	All causes	26,860	100.0	3,036.7
1	Malignant neoplasms (C00-C97)	8,312	30.9	939.7
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	6,746	25.1	762.7
3	Cerebrovascular diseases (I60-I69)	1,642	6.1	185.6
4	Diabetes mellitus (E10-E14)	1,473	5.5	166.5
5	Chronic lower respiratory diseases (J40-J47)	1,201	4.5	135.8
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	925	3.4	104.6
7	Septicemia (A40-A41)	659	2.5	74.5
8	Essential hypertension and hypertensive renal disease (I10,I12,I15)	468	1.7	52.9
9	Influenza and pneumonia (J09-J18)	464	1.7	52.5
10	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	403	1.5	45.6
11	Alzheimer's disease (G30)	287	1.1	32.4
12	Chronic liver disease and cirrhosis (K70,K73-K74)	149	0.6	16.8
13	Pneumonitis due to solids and liquids (J69)	129	0.5	14.6
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	124	0.5	14.0
15	Parkinson's disease (G20-G21)	123	0.5	13.9
...	All other causes (Residual)	3,755	14.0	424.5

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, both sexes, 75-79 years				
...	All causes	28,320	100.0	4,399.8
1	Malignant neoplasms (C00-C97)	7,587	26.8	1,178.7
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	7,182	25.4	1,115.8
3	Cerebrovascular diseases (I60-I69)	1,888	6.7	293.3
4	Diabetes mellitus (E10-E14)	1,499	5.3	232.9
5	Chronic lower respiratory diseases (J40-J47)	1,271	4.5	197.5
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,040	3.7	161.6
7	Septicemia (A40-A41)	710	2.5	110.3
8	Alzheimer's disease (G30)	684	2.4	106.3
9	Influenza and pneumonia (J09-J18)	571	2.0	88.7
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	518	1.8	80.5
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	397	1.4	61.7
12	Pneumonitis due to solids and liquids (J69)	189	0.7	29.4
13	Parkinson's disease (G20-G21)	173	0.6	26.9
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	151	0.5	23.5
15	Chronic liver disease and cirrhosis (K70,K73-K74)	104	0.4	16.2
...	All other causes (Residual)	4,356	15.4	676.7
Non-Hispanic, black, both sexes, 80-84 years				
...	All causes	29,047	100.0	6,717.2
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	7,660	26.4	1,771.4
2	Malignant neoplasms (C00-C97)	6,394	22.0	1,478.6
3	Cerebrovascular diseases (I60-I69)	2,051	7.1	474.3
4	Diabetes mellitus (E10-E14)	1,411	4.9	326.3
5	Chronic lower respiratory diseases (J40-J47)	1,220	4.2	282.1
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,032	3.6	238.7
7	Alzheimer's disease (G30)	1,026	3.5	237.3
8	Septicemia (A40-A41)	680	2.3	157.3
9	Influenza and pneumonia (J09-J18)	648	2.2	149.9
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	585	2.0	135.3
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	363	1.2	83.9
12	Pneumonitis due to solids and liquids (J69)	217	0.7	50.2
13	Parkinson's disease (G20-G21)	209	0.7	48.3
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	179	0.6	41.4
15	Enterocolitis due to <i>Clostridium difficile</i> (A04.7)	95	0.3	22.0
...	All other causes (Residual)	5,277	18.2	1,220.3

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, both sexes, 85-89 years				
...	All causes	25,996	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	7,118	27.4	---
2	Malignant neoplasms (C00-C97)	4,399	16.9	---
3	Cerebrovascular diseases (I60-I69)	1,844	7.1	---
4	Alzheimer's disease (G30)	1,383	5.3	---
5	Diabetes mellitus (E10-E14)	1,096	4.2	---
6	Chronic lower respiratory diseases (J40-J47)	979	3.8	---
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	878	3.4	---
8	Influenza and pneumonia (J09-J18)	705	2.7	---
9	Septicemia (A40-A41)	636	2.4	---
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	596	2.3	---
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	299	1.2	---
12	Pneumonitis due to solids and liquids (J69)	245	0.9	---
13	Parkinson's disease (G20-G21)	168	0.6	---
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	132	0.5	---
15	Atherosclerosis (I70)	108	0.4	---
...	All other causes (Residual)	5,410	20.8	---
Non-Hispanic, black, both sexes, 90-94 years				
...	All causes	16,549	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	4,725	28.6	---
2	Malignant neoplasms (C00-C97)	1,977	11.9	---
3	Cerebrovascular diseases (I60-I69)	1,234	7.5	---
4	Alzheimer's disease (G30)	1,159	7.0	---
5	Diabetes mellitus (E10-E14)	579	3.5	---
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	535	3.2	---
7	Chronic lower respiratory diseases (J40-J47)	473	2.9	---
8	Essential hypertension and hypertensive renal disease (I10,I12,I15)	456	2.8	---
9	Influenza and pneumonia (J09-J18)	453	2.7	---
10	Septicemia (A40-A41)	348	2.1	---
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	213	1.3	---
12	Pneumonitis due to solids and liquids (J69)	175	1.1	---
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	83	0.5	---
14	Parkinson's disease (G20-G21)	70	0.4	---
14	Atherosclerosis (I70)	70	0.4	---
...	All other causes (Residual)	3,999	24.2	---

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, both sexes, 95-99 years				
...	All causes	7,207	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,136	29.6	---
2	Malignant neoplasms (C00-C97)	588	8.2	---
3	Alzheimer's disease (G30)	554	7.7	---
4	Cerebrovascular diseases (I60-I69)	487	6.8	---
5	Essential hypertension and hypertensive renal disease (I10,I12,I15)	237	3.3	---
6	Influenza and pneumonia (J09-J18)	230	3.2	---
7	Diabetes mellitus (E10-E14)	217	3.0	---
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	195	2.7	---
9	Chronic lower respiratory diseases (J40-J47)	161	2.2	---
10	Septicemia (A40-A41)	142	2.0	---
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	83	1.2	---
12	Pneumonitis due to solids and liquids (J69)	63	0.9	---
13	Atherosclerosis (I70)	38	0.5	---
14	Anemias (D50-D64)	25	0.3	---
14	Nutritional deficiencies (E40-E64)	25	0.3	---
...	All other causes (Residual)	2,026	28.1	---
Non-Hispanic, black, both sexes, 100 years and over				
...	All causes	2,181	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	682	31.3	---
2	Alzheimer's disease (G30)	158	7.2	---
3	Cerebrovascular diseases (I60-I69)	136	6.2	---
4	Malignant neoplasms (C00-C97)	106	4.9	---
5	Influenza and pneumonia (J09-J18)	86	3.9	---
6	Essential hypertension and hypertensive renal disease (I10,I12,I15)	84	3.9	---
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	50	2.3	---
8	Chronic lower respiratory diseases (J40-J47)	43	2.0	---
9	Diabetes mellitus (E10-E14)	41	1.9	---
10	Septicemia (A40-A41)	36	1.7	---
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	28	1.3	---
12	Atherosclerosis (I70)	22	1.0	---
12	Pneumonitis due to solids and liquids (J69)	22	1.0	---
14	Anemias (D50-D64)	12	0.6	---
15	Nutritional deficiencies (E40-E64)	9	0.4	---
...	All other causes (Residual)	666	30.5	---

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, male, all ages				
...	All causes	145,052	100.0	760.4
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	34,476	23.8	180.7
2	Malignant neoplasms (C00-C97)	33,628	23.2	176.3
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	8,219	5.7	43.1
4	Cerebrovascular diseases (I60-I69)	6,979	4.8	36.6
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	6,658	4.6	34.9
6	Diabetes mellitus (E10-E14)	5,986	4.1	31.4
7	Chronic lower respiratory diseases (J40-J47)	4,648	3.2	24.4
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3,734	2.6	19.6
9	Human immunodeficiency virus (HIV) disease (B20-B24)	2,693	1.9	14.1
10	Septicemia (A40-A41)	2,666	1.8	14.0
11	Influenza and pneumonia (J09-J18)	2,386	1.6	12.5
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	2,180	1.5	11.4
13	Certain conditions originating in the perinatal period (P00-P96)	2,171	1.5	11.4
14	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,790	1.2	9.4
15	Chronic liver disease and cirrhosis (K70,K73-K74)	1,750	1.2	9.2
...	All other causes (Residual)	25,088	17.3	131.5
Non-Hispanic, black, male, 1-4 years				
...	All causes	575	100.0	45.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	167	29.0	13.3
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	84	14.6	6.7
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	57	9.9	4.5
4	Malignant neoplasms (C00-C97)	30	5.2	2.4
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	28	4.9	2.2
6	Influenza and pneumonia (J09-J18)	18	3.1	*
7	Chronic lower respiratory diseases (J40-J47)	16	2.8	*
8	Cerebrovascular diseases (I60-I69)	14	2.4	*
9	Certain conditions originating in the perinatal period (P00-P96)	8	1.4	*
10	Septicemia (A40-A41)	6	1.0	*
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	6	1.0	*
10	Complications of medical and surgical care (Y40-Y84,Y88)	6	1.0	*
13	Anemias (D50-D64)	3	0.5	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2	0.3	*
15	Meningitis (G00,G03)	1	0.2	*
15	Acute bronchitis and bronchiolitis (J20-J21)	1	0.2	*
15	Pneumonitis due to solids and liquids (J69)	1	0.2	*
...	All other causes (Residual)	127	22.1	10.1

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, male, 5-9 years				
...	All causes	294	100.0	19.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	92	31.3	6.0
2	Malignant neoplasms (C00-C97)	36	12.2	2.4
3	Chronic lower respiratory diseases (J40-J47)	28	9.5	1.8
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	21	7.1	1.4
5	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	19	6.5	*
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	12	4.1	*
7	Anemias (D50-D64)	9	3.1	*
8	Influenza and pneumonia (J09-J18)	7	2.4	*
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	6	2.0	*
10	Cerebrovascular diseases (I60-I69)	5	1.7	*
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3	1.0	*
11	Certain conditions originating in the perinatal period (P00-P96)	3	1.0	*
13	Septicemia (A40-A41)	2	0.7	*
13	Pneumonitis due to solids and liquids (J69)	2	0.7	*
13	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2	0.7	*
...	All other causes (Residual)	47	16.0	3.1
Non-Hispanic, black, male, 10-14 years				
...	All causes	390	100.0	24.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	104	26.7	6.5
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	40	10.3	2.5
3	Malignant neoplasms (C00-C97)	37	9.5	2.3
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	30	7.7	1.9
5	Chronic lower respiratory diseases (J40-J47)	29	7.4	1.8
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	24	6.2	1.5
7	Diseases of heart (I00-I09,I11,I13,I20-I51)	13	3.3	*
8	Influenza and pneumonia (J09-J18)	8	2.1	*
9	Cerebrovascular diseases (I60-I69)	6	1.5	*
10	Anemias (D50-D64)	5	1.3	*
11	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	4	1.0	*
12	Diabetes mellitus (E10-E14)	3	0.8	*
12	Pneumonitis due to solids and liquids (J69)	3	0.8	*
12	Complications of medical and surgical care (Y40-Y84,Y88)	3	0.8	*
15	Septicemia (A40-A41)	2	0.5	*
...	All other causes (Residual)	79	20.3	4.9

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, male, 15-19 years				
...	All causes	1,781	100.0	102.5
1	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	893	50.1	51.4
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	402	22.6	23.1
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	125	7.0	7.2
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	72	4.0	4.1
5	Malignant neoplasms (C00-C97)	57	3.2	3.3
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	30	1.7	1.7
7	Chronic lower respiratory diseases (J40-J47)	26	1.5	1.5
8	Legal intervention (Y35,Y89.0)	15	0.8	*
9	Anemias (D50-D64)	8	0.4	*
10	Diabetes mellitus (E10-E14)	5	0.3	*
10	Influenza and pneumonia (J09-J18)	5	0.3	*
12	Septicemia (A40-A41)	4	0.2	*
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	4	0.2	*
12	Cerebrovascular diseases (I60-I69)	4	0.2	*
12	Aortic aneurysm and dissection (I71)	4	0.2	*
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	4	0.2	*
...	All other causes (Residual)	123	6.9	7.1
Non-Hispanic, black, male, 20-24 years				
...	All causes	3,259	100.0	197.0
1	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,615	49.6	97.6
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	610	18.7	36.9
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	283	8.7	17.1
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	125	3.8	7.6
5	Malignant neoplasms (C00-C97)	94	2.9	5.7
6	Human immunodeficiency virus (HIV) disease (B20-B24)	59	1.8	3.6
7	Anemias (D50-D64)	35	1.1	2.1
8	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	32	1.0	1.9
9	Legal intervention (Y35,Y89.0)	30	0.9	1.8
10	Chronic lower respiratory diseases (J40-J47)	28	0.9	1.7
11	Diabetes mellitus (E10-E14)	18	0.6	*
12	Influenza and pneumonia (J09-J18)	15	0.5	*
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	13	0.4	*
14	Cerebrovascular diseases (I60-I69)	12	0.4	*
15	Septicemia (A40-A41)	8	0.2	*
...	All other causes (Residual)	282	8.7	17.0

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, male, 25-29 years				
...	All causes	3,038	100.0	222.0
1	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,202	39.6	87.8
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	618	20.3	45.2
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	250	8.2	18.3
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	205	6.7	15.0
5	Human immunodeficiency virus (HIV) disease (B20-B24)	111	3.7	8.1
6	Malignant neoplasms (C00-C97)	109	3.6	8.0
7	Anemias (D50-D64)	38	1.3	2.8
8	Diabetes mellitus (E10-E14)	34	1.1	2.5
9	Chronic lower respiratory diseases (J40-J47)	31	1.0	2.3
10	Influenza and pneumonia (J09-J18)	26	0.9	1.9
11	Cerebrovascular diseases (I60-I69)	23	0.8	1.7
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	23	0.8	1.7
13	Septicemia (A40-A41)	18	0.6	*
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	17	0.6	*
15	Legal intervention (Y35,Y89.0)	15	0.5	*
...	All other causes (Residual)	318	10.5	23.2
Non-Hispanic, black, male, 30-34 years				
...	All causes	2,980	100.0	231.7
1	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	859	28.8	66.8
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	555	18.6	43.2
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	362	12.1	28.1
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	196	6.6	15.2
5	Malignant neoplasms (C00-C97)	151	5.1	11.7
6	Human immunodeficiency virus (HIV) disease (B20-B24)	129	4.3	10.0
7	Diabetes mellitus (E10-E14)	78	2.6	6.1
8	Cerebrovascular diseases (I60-I69)	44	1.5	3.4
9	Anemias (D50-D64)	37	1.2	2.9
10	Influenza and pneumonia (J09-J18)	29	1.0	2.3
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	29	1.0	2.3
12	Chronic lower respiratory diseases (J40-J47)	26	0.9	2.0
13	Chronic liver disease and cirrhosis (K70,K73-K74)	24	0.8	1.9
14	Septicemia (A40-A41)	22	0.7	1.7
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	18	0.6	*
...	All other causes (Residual)	421	14.1	32.7

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, male, 35-39 years				
...	All causes	3,288	100.0	275.4
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	591	18.0	49.5
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	552	16.8	46.2
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	550	16.7	46.1
4	Malignant neoplasms (C00-C97)	270	8.2	22.6
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	192	5.8	16.1
6	Human immunodeficiency virus (HIV) disease (B20-B24)	181	5.5	15.2
7	Diabetes mellitus (E10-E14)	86	2.6	7.2
8	Cerebrovascular diseases (I60-I69)	85	2.6	7.1
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	55	1.7	4.6
10	Septicemia (A40-A41)	49	1.5	4.1
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	42	1.3	3.5
12	Influenza and pneumonia (J09-J18)	37	1.1	3.1
13	Chronic lower respiratory diseases (J40-J47)	35	1.1	2.9
14	Anemias (D50-D64)	28	0.9	2.3
15	Aortic aneurysm and dissection (I71)	26	0.8	2.2
15	Chronic liver disease and cirrhosis (K70,K73-K74)	26	0.8	2.2
...	All other causes (Residual)	483	14.7	40.5
Non-Hispanic, black, male, 40-44 years				
...	All causes	4,581	100.0	364.5
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,122	24.5	89.3
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	625	13.6	49.7
3	Malignant neoplasms (C00-C97)	477	10.4	38.0
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	395	8.6	31.4
5	Human immunodeficiency virus (HIV) disease (B20-B24)	321	7.0	25.5
6	Diabetes mellitus (E10-E14)	188	4.1	15.0
7	Cerebrovascular diseases (I60-I69)	177	3.9	14.1
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	144	3.1	11.5
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	94	2.1	7.5
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	70	1.5	5.6
11	Septicemia (A40-A41)	67	1.5	5.3
11	Chronic liver disease and cirrhosis (K70,K73-K74)	67	1.5	5.3
13	Influenza and pneumonia (J09-J18)	56	1.2	4.5
14	Chronic lower respiratory diseases (J40-J47)	51	1.1	4.1
15	Aortic aneurysm and dissection (I71)	41	0.9	3.3
...	All other causes (Residual)	686	15.0	54.6

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, male, 45-49 years				
...	All causes	7,306	100.0	556.3
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,885	25.8	143.5
2	Malignant neoplasms (C00-C97)	1,204	16.5	91.7
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	759	10.4	57.8
4	Human immunodeficiency virus (HIV) disease (B20-B24)	445	6.1	33.9
5	Cerebrovascular diseases (I60-I69)	350	4.8	26.6
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	329	4.5	25.0
7	Diabetes mellitus (E10-E14)	302	4.1	23.0
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	161	2.2	12.3
9	Chronic liver disease and cirrhosis (K70,K73-K74)	153	2.1	11.6
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	139	1.9	10.6
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	121	1.7	9.2
12	Septicemia (A40-A41)	104	1.4	7.9
13	Chronic lower respiratory diseases (J40-J47)	93	1.3	7.1
14	Influenza and pneumonia (J09-J18)	75	1.0	5.7
15	Viral hepatitis (B15-B19)	49	0.7	3.7
...	All other causes (Residual)	1,137	15.6	86.6
Non-Hispanic, black, male, 50-54 years				
...	All causes	11,632	100.0	908.4
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,074	26.4	240.1
2	Malignant neoplasms (C00-C97)	2,701	23.2	210.9
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	854	7.3	66.7
4	Cerebrovascular diseases (I60-I69)	561	4.8	43.8
5	Human immunodeficiency virus (HIV) disease (B20-B24)	515	4.4	40.2
6	Diabetes mellitus (E10-E14)	490	4.2	38.3
7	Chronic liver disease and cirrhosis (K70,K73-K74)	322	2.8	25.1
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	260	2.2	20.3
9	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	241	2.1	18.8
10	Chronic lower respiratory diseases (J40-J47)	222	1.9	17.3
11	Septicemia (A40-A41)	210	1.8	16.4
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	176	1.5	13.7
13	Influenza and pneumonia (J09-J18)	140	1.2	10.9
14	Viral hepatitis (B15-B19)	130	1.1	10.2
15	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	123	1.1	9.6
...	All other causes (Residual)	1,613	13.9	126.0

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, male, 55-59 years				
...	All causes	14,813	100.0	1,404.2
1	Malignant neoplasms (C00-C97)	4,078	27.5	386.6
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,810	25.7	361.2
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	840	5.7	79.6
4	Cerebrovascular diseases (I60-I69)	748	5.0	70.9
5	Diabetes mellitus (E10-E14)	709	4.8	67.2
6	Human immunodeficiency virus (HIV) disease (B20-B24)	427	2.9	40.5
7	Chronic liver disease and cirrhosis (K70,K73-K74)	407	2.7	38.6
8	Chronic lower respiratory diseases (J40-J47)	333	2.2	31.6
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	333	2.2	31.6
10	Septicemia (A40-A41)	282	1.9	26.7
11	Viral hepatitis (B15-B19)	226	1.5	21.4
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	217	1.5	20.6
13	Influenza and pneumonia (J09-J18)	187	1.3	17.7
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	178	1.2	16.9
15	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	92	0.6	8.7
...	All other causes (Residual)	1,946	13.1	184.5
Non-Hispanic, black, male, 60-64 years				
...	All causes	16,458	100.0	2,006.5
1	Malignant neoplasms (C00-C97)	5,154	31.3	628.3
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	4,367	26.5	532.4
3	Diabetes mellitus (E10-E14)	825	5.0	100.6
4	Cerebrovascular diseases (I60-I69)	823	5.0	100.3
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	579	3.5	70.6
6	Chronic lower respiratory diseases (J40-J47)	468	2.8	57.1
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	451	2.7	55.0
8	Chronic liver disease and cirrhosis (K70,K73-K74)	378	2.3	46.1
9	Septicemia (A40-A41)	326	2.0	39.7
10	Human immunodeficiency virus (HIV) disease (B20-B24)	270	1.6	32.9
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	259	1.6	31.6
12	Influenza and pneumonia (J09-J18)	209	1.3	25.5
13	Viral hepatitis (B15-B19)	183	1.1	22.3
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	99	0.6	12.1
15	Pneumonitis due to solids and liquids (J69)	67	0.4	8.2
...	All other causes (Residual)	2,000	12.2	243.8

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, male, 65-69 years				
...	All causes	14,363	100.0	2,740.7
1	Malignant neoplasms (C00-C97)	4,650	32.4	887.3
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,808	26.5	726.6
3	Cerebrovascular diseases (I60-I69)	756	5.3	144.3
4	Diabetes mellitus (E10-E14)	742	5.2	141.6
5	Chronic lower respiratory diseases (J40-J47)	578	4.0	110.3
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	431	3.0	82.2
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	358	2.5	68.3
8	Septicemia (A40-A41)	281	2.0	53.6
9	Essential hypertension and hypertensive renal disease (I10,I12,I15)	249	1.7	47.5
10	Influenza and pneumonia (J09-J18)	200	1.4	38.2
11	Chronic liver disease and cirrhosis (K70,K73-K74)	166	1.2	31.7
12	Human immunodeficiency virus (HIV) disease (B20-B24)	124	0.9	23.7
13	Pneumonitis due to solids and liquids (J69)	70	0.5	13.4
14	Alzheimer's disease (G30)	58	0.4	11.1
15	Viral hepatitis (B15-B19)	57	0.4	10.9
...	All other causes (Residual)	1,835	12.8	350.1
Non-Hispanic, black, male, 70-74 years				
...	All causes	14,234	100.0	3,861.0
1	Malignant neoplasms (C00-C97)	4,531	31.8	1,229.0
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,719	26.1	1,008.8
3	Cerebrovascular diseases (I60-I69)	820	5.8	222.4
4	Diabetes mellitus (E10-E14)	738	5.2	200.2
5	Chronic lower respiratory diseases (J40-J47)	671	4.7	182.0
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	454	3.2	123.1
7	Septicemia (A40-A41)	302	2.1	81.9
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	248	1.7	67.3
9	Influenza and pneumonia (J09-J18)	238	1.7	64.6
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	224	1.6	60.8
11	Alzheimer's disease (G30)	125	0.9	33.9
12	Chronic liver disease and cirrhosis (K70,K73-K74)	96	0.7	26.0
13	Parkinson's disease (G20-G21)	81	0.6	22.0
14	Pneumonitis due to solids and liquids (J69)	78	0.5	21.2
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	56	0.4	15.2
...	All other causes (Residual)	1,853	13.0	502.6

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, male, 75-79 years				
...	All causes	13,694	100.0	5,466.7
1	Malignant neoplasms (C00-C97)	3,913	28.6	1,562.1
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,566	26.0	1,423.6
3	Cerebrovascular diseases (I60-I69)	844	6.2	336.9
4	Chronic lower respiratory diseases (J40-J47)	688	5.0	274.7
5	Diabetes mellitus (E10-E14)	635	4.6	253.5
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	468	3.4	186.8
7	Septicemia (A40-A41)	303	2.2	121.0
8	Alzheimer's disease (G30)	279	2.0	111.4
9	Influenza and pneumonia (J09-J18)	272	2.0	108.6
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	242	1.8	96.6
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	230	1.7	91.8
12	Pneumonitis due to solids and liquids (J69)	103	0.8	41.1
13	Parkinson's disease (G20-G21)	99	0.7	39.5
14	Chronic liver disease and cirrhosis (K70,K73-K74)	56	0.4	22.4
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	54	0.4	21.6
...	All other causes (Residual)	1,942	14.2	775.3
Non-Hispanic, black, male, 80-84 years				
...	All causes	12,367	100.0	8,215.3
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,296	26.7	2,189.5
2	Malignant neoplasms (C00-C97)	3,091	25.0	2,053.3
3	Cerebrovascular diseases (I60-I69)	740	6.0	491.6
4	Chronic lower respiratory diseases (J40-J47)	608	4.9	403.9
5	Diabetes mellitus (E10-E14)	533	4.3	354.1
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	404	3.3	268.4
7	Alzheimer's disease (G30)	337	2.7	223.9
8	Influenza and pneumonia (J09-J18)	304	2.5	201.9
9	Septicemia (A40-A41)	280	2.3	186.0
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	220	1.8	146.1
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	171	1.4	113.6
12	Parkinson's disease (G20-G21)	120	1.0	79.7
13	Pneumonitis due to solids and liquids (J69)	118	1.0	78.4
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	76	0.6	50.5
15	Chronic liver disease and cirrhosis (K70,K73-K74)	34	0.3	22.6
15	Enterocolitis due to <i>Clostridium difficile</i> (A04.7)	34	0.3	22.6
...	All other causes (Residual)	2,001	16.2	1,329.2

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, male, 85-89 years				
...	All causes	9,541	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,532	26.5	---
2	Malignant neoplasms (C00-C97)	2,027	21.2	---
3	Cerebrovascular diseases (I60-I69)	574	6.0	---
4	Chronic lower respiratory diseases (J40-J47)	451	4.7	---
5	Diabetes mellitus (E10-E14)	398	4.2	---
6	Alzheimer's disease (G30)	377	4.0	---
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	324	3.4	---
8	Influenza and pneumonia (J09-J18)	291	3.0	---
9	Septicemia (A40-A41)	219	2.3	---
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	177	1.9	---
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	142	1.5	---
12	Pneumonitis due to solids and liquids (J69)	116	1.2	---
13	Parkinson's disease (G20-G21)	79	0.8	---
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	51	0.5	---
15	Atherosclerosis (I70)	42	0.4	---
...	All other causes (Residual)	1,741	18.2	---
Non-Hispanic, black, male, 90-94 years				
...	All causes	4,760	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,307	27.5	---
2	Malignant neoplasms (C00-C97)	790	16.6	---
3	Cerebrovascular diseases (I60-I69)	279	5.9	---
4	Alzheimer's disease (G30)	248	5.2	---
5	Chronic lower respiratory diseases (J40-J47)	189	4.0	---
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	166	3.5	---
7	Diabetes mellitus (E10-E14)	154	3.2	---
8	Influenza and pneumonia (J09-J18)	145	3.0	---
9	Essential hypertension and hypertensive renal disease (I10,I12,I15)	102	2.1	---
10	Septicemia (A40-A41)	100	2.1	---
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	82	1.7	---
12	Pneumonitis due to solids and liquids (J69)	70	1.5	---
13	Parkinson's disease (G20-G21)	32	0.7	---
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	20	0.4	---
15	Atherosclerosis (I70)	16	0.3	---
...	All other causes (Residual)	1,060	22.3	---

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, male, 95-99 years				
...	All causes	1,488	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	436	29.3	---
2	Malignant neoplasms (C00-C97)	201	13.5	---
3	Alzheimer's disease (G30)	78	5.2	---
3	Cerebrovascular diseases (I60-I69)	78	5.2	---
5	Influenza and pneumonia (J09-J18)	61	4.1	---
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	52	3.5	---
7	Chronic lower respiratory diseases (J40-J47)	49	3.3	---
8	Diabetes mellitus (E10-E14)	43	2.9	---
9	Septicemia (A40-A41)	37	2.5	---
9	Essential hypertension and hypertensive renal disease (I10,I12,I15)	37	2.5	---
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	24	1.6	---
12	Pneumonitis due to solids and liquids (J69)	12	0.8	---
13	Parkinson's disease (G20-G21)	11	0.7	---
13	Atherosclerosis (I70)	11	0.7	---
15	Hyperplasia of prostate (N40)	7	0.5	---
...	All other causes (Residual)	351	23.6	---
Non-Hispanic, black, male, 100 years and over				
...	All causes	309	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	94	30.4	---
2	Malignant neoplasms (C00-C97)	19	6.1	---
2	Chronic lower respiratory diseases (J40-J47)	19	6.1	---
4	Influenza and pneumonia (J09-J18)	18	5.8	---
5	Alzheimer's disease (G30)	16	5.2	---
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	15	4.9	---
7	Essential hypertension and hypertensive renal disease (I10,I12,I15)	12	3.9	---
8	Cerebrovascular diseases (I60-I69)	11	3.6	---
9	Septicemia (A40-A41)	6	1.9	---
10	Diabetes mellitus (E10-E14)	5	1.6	---
10	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	5	1.6	---
12	Pneumonitis due to solids and liquids (J69)	4	1.3	---
13	Atherosclerosis (I70)	3	1.0	---
14	Anemias (D50-D64)	2	0.6	---
14	Nutritional deficiencies (E40-E64)	2	0.6	---
14	Parkinson's disease (G20-G21)	2	0.6	---
...	All other causes (Residual)	76	24.6	---

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, female, all ages				
...	All causes	141,745	100.0	679.2
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	33,119	23.4	158.7
2	Malignant neoplasms (C00-C97)	32,530	22.9	155.9
3	Cerebrovascular diseases (I60-I69)	8,743	6.2	41.9
4	Diabetes mellitus (E10-E14)	6,785	4.8	32.5
5	Chronic lower respiratory diseases (J40-J47)	4,346	3.1	20.8
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	4,310	3.0	20.7
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	4,080	2.9	19.5
8	Alzheimer's disease (G30)	3,889	2.7	18.6
9	Septicemia (A40-A41)	3,297	2.3	15.8
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	2,785	2.0	13.3
11	Influenza and pneumonia (J09-J18)	2,712	1.9	13.0
12	Certain conditions originating in the perinatal period (P00-P96)	1,719	1.2	8.2
13	Human immunodeficiency virus (HIV) disease (B20-B24)	1,445	1.0	6.9
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,106	0.8	5.3
15	Chronic liver disease and cirrhosis (K70,K73-K74)	1,033	0.7	4.9
...	All other causes (Residual)	29,846	21.1	143.0
Non-Hispanic, black, female, 1-4 years				
...	All causes	436	100.0	35.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	119	27.3	9.8
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	70	16.1	5.7
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	48	11.0	3.9
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	24	5.5	2.0
5	Malignant neoplasms (C00-C97)	23	5.3	1.9
6	Influenza and pneumonia (J09-J18)	14	3.2	*
7	Chronic lower respiratory diseases (J40-J47)	8	1.8	*
8	Septicemia (A40-A41)	5	1.1	*
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	4	0.9	*
9	Certain conditions originating in the perinatal period (P00-P96)	4	0.9	*
11	Anemias (D50-D64)	3	0.7	*
11	Cerebrovascular diseases (I60-I69)	3	0.7	*
13	Complications of medical and surgical care (Y40-Y84,Y88)	2	0.5	*
14	Meningitis (G00,G03)	1	0.2	*
14	Atherosclerosis (I70)	1	0.2	*
14	Acute bronchitis and bronchiolitis (J20-J21)	1	0.2	*
14	Pneumonitis due to solids and liquids (J69)	1	0.2	*
14	Peptic ulcer (K25-K28)	1	0.2	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1	0.2	*
14	Enterocolitis due to <i>Clostridium difficile</i> (A04.7)	1	0.2	*
...	All other causes (Residual)	102	23.4	8.4

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, female, 5-9 years				
...	All causes	233	100.0	15.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	65	27.9	4.4
2	Malignant neoplasms (C00-C97)	27	11.6	1.8
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	16	6.9	*
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	15	6.4	*
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	9	3.9	*
6	Chronic lower respiratory diseases (J40-J47)	8	3.4	*
7	Cerebrovascular diseases (I60-I69)	7	3.0	*
8	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	6	2.6	*
8	Anemias (D50-D64)	6	2.6	*
10	Septicemia (A40-A41)	4	1.7	*
10	Influenza and pneumonia (J09-J18)	4	1.7	*
12	Diseases of appendix (K35-K38)	2	0.9	*
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2	0.9	*
14	Meningitis (G00,G03)	1	0.4	*
14	Acute bronchitis and bronchiolitis (J20-J21)	1	0.4	*
14	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1	0.4	*
14	Complications of medical and surgical care (Y40-Y84,Y88)	1	0.4	*
...	All other causes (Residual)	58	24.9	3.9
Non-Hispanic, black, female, 10-14 years				
...	All causes	233	100.0	15.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	52	22.3	3.4
2	Malignant neoplasms (C00-C97)	27	11.6	1.7
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	19	8.2	*
4	Chronic lower respiratory diseases (J40-J47)	16	6.9	*
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	15	6.4	*
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	13	5.6	*
7	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	10	4.3	*
8	Influenza and pneumonia (J09-J18)	6	2.6	*
9	Septicemia (A40-A41)	3	1.3	*
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	3	1.3	*
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3	1.3	*
12	Anemias (D50-D64)	2	0.9	*
12	Meningitis (G00,G03)	2	0.9	*
12	Cerebrovascular diseases (I60-I69)	2	0.9	*
15	Diabetes mellitus (E10-E14)	1	0.4	*
15	Certain conditions originating in the perinatal period (P00-P96)	1	0.4	*
...	All other causes (Residual)	58	24.9	3.8

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, female, 15-19 years				
...	All causes	554	100.0	33.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	138	24.9	8.2
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	119	21.5	7.1
3	Malignant neoplasms (C00-C97)	39	7.0	2.3
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	34	6.1	2.0
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	26	4.7	1.6
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	23	4.2	1.4
7	Pregnancy, childbirth and the puerperium (O00-O99)	11	2.0	*
8	Human immunodeficiency virus (HIV) disease (B20-B24)	10	1.8	*
8	Anemias (D50-D64)	10	1.8	*
8	Chronic lower respiratory diseases (J40-J47)	10	1.8	*
11	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	5	0.9	*
11	Diabetes mellitus (E10-E14)	5	0.9	*
11	Cerebrovascular diseases (I60-I69)	5	0.9	*
14	Influenza and pneumonia (J09-J18)	4	0.7	*
14	Pneumonitis due to solids and liquids (J69)	4	0.7	*
...	All other causes (Residual)	111	20.0	6.6
Non-Hispanic, black, female, 20-24 years				
...	All causes	982	100.0	58.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	207	21.1	12.4
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	170	17.3	10.2
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	80	8.1	4.8
4	Malignant neoplasms (C00-C97)	67	6.8	4.0
5	Pregnancy, childbirth and the puerperium (O00-O99)	51	5.2	3.1
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	45	4.6	2.7
7	Human immunodeficiency virus (HIV) disease (B20-B24)	28	2.9	1.7
8	Septicemia (A40-A41)	20	2.0	1.2
8	Influenza and pneumonia (J09-J18)	20	2.0	1.2
10	Diabetes mellitus (E10-E14)	19	1.9	*
11	Anemias (D50-D64)	18	1.8	*
12	Chronic lower respiratory diseases (J40-J47)	17	1.7	*
12	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	17	1.7	*
14	Cerebrovascular diseases (I60-I69)	10	1.0	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	8	0.8	*
...	All other causes (Residual)	205	20.9	12.3

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, female, 25-29 years				
...	All causes	1,208	100.0	82.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	215	17.8	14.7
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	144	11.9	9.8
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	124	10.3	8.5
4	Malignant neoplasms (C00-C97)	123	10.2	8.4
5	Pregnancy, childbirth and the puerperium (O00-O99)	61	5.0	4.2
6	Human immunodeficiency virus (HIV) disease (B20-B24)	54	4.5	3.7
7	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	50	4.1	3.4
8	Diabetes mellitus (E10-E14)	45	3.7	3.1
9	Septicemia (A40-A41)	26	2.2	1.8
10	Anemias (D50-D64)	25	2.1	1.7
11	Influenza and pneumonia (J09-J18)	22	1.8	1.5
12	Cerebrovascular diseases (I60-I69)	17	1.4	*
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	16	1.3	*
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	13	1.1	*
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	13	1.1	*
...	All other causes (Residual)	260	21.5	17.7
Non-Hispanic, black, female, 30-34 years				
...	All causes	1,731	100.0	121.5
1	Malignant neoplasms (C00-C97)	257	14.8	18.0
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	250	14.4	17.5
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	244	14.1	17.1
4	Human immunodeficiency virus (HIV) disease (B20-B24)	118	6.8	8.3
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	97	5.6	6.8
6	Diabetes mellitus (E10-E14)	69	4.0	4.8
7	Pregnancy, childbirth and the puerperium (O00-O99)	63	3.6	4.4
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	52	3.0	3.6
9	Cerebrovascular diseases (I60-I69)	51	2.9	3.6
10	Anemias (D50-D64)	33	1.9	2.3
11	Septicemia (A40-A41)	31	1.8	2.2
12	Influenza and pneumonia (J09-J18)	29	1.7	2.0
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	26	1.5	1.8
14	Chronic lower respiratory diseases (J40-J47)	22	1.3	1.5
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	18	1.0	*
...	All other causes (Residual)	371	21.4	26.0

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, female, 35-39 years				
...	All causes	2,258	100.0	167.9
1	Malignant neoplasms (C00-C97)	458	20.3	34.1
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	376	16.7	28.0
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	213	9.4	15.8
4	Human immunodeficiency virus (HIV) disease (B20-B24)	165	7.3	12.3
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	99	4.4	7.4
6	Cerebrovascular diseases (I60-I69)	96	4.3	7.1
7	Diabetes mellitus (E10-E14)	74	3.3	5.5
8	Pregnancy, childbirth and the puerperium (O00-O99)	54	2.4	4.0
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	51	2.3	3.8
10	Chronic lower respiratory diseases (J40-J47)	39	1.7	2.9
11	Septicemia (A40-A41)	34	1.5	2.5
12	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	32	1.4	2.4
13	Chronic liver disease and cirrhosis (K70,K73-K74)	30	1.3	2.2
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	29	1.3	2.2
15	Anemias (D50-D64)	24	1.1	1.8
15	Influenza and pneumonia (J09-J18)	24	1.1	1.8
...	All other causes (Residual)	460	20.4	34.2
Non-Hispanic, black, female, 40-44 years				
...	All causes	3,484	100.0	247.1
1	Malignant neoplasms (C00-C97)	884	25.4	62.7
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	592	17.0	42.0
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	269	7.7	19.1
4	Human immunodeficiency virus (HIV) disease (B20-B24)	222	6.4	15.7
5	Cerebrovascular diseases (I60-I69)	151	4.3	10.7
6	Diabetes mellitus (E10-E14)	139	4.0	9.9
7	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	79	2.3	5.6
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	70	2.0	5.0
9	Chronic lower respiratory diseases (J40-J47)	66	1.9	4.7
10	Septicemia (A40-A41)	65	1.9	4.6
11	Influenza and pneumonia (J09-J18)	62	1.8	4.4
12	Chronic liver disease and cirrhosis (K70,K73-K74)	56	1.6	4.0
13	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	49	1.4	3.5
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	45	1.3	3.2
15	Anemias (D50-D64)	32	0.9	2.3
...	All other causes (Residual)	703	20.2	49.9

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, female, 45-49 years				
...	All causes	5,727	100.0	388.3
1	Malignant neoplasms (C00-C97)	1,655	28.9	112.2
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,060	18.5	71.9
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	351	6.1	23.8
4	Cerebrovascular diseases (I60-I69)	301	5.3	20.4
5	Diabetes mellitus (E10-E14)	238	4.2	16.1
6	Human immunodeficiency virus (HIV) disease (B20-B24)	236	4.1	16.0
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	129	2.3	8.7
8	Septicemia (A40-A41)	128	2.2	8.7
9	Chronic lower respiratory diseases (J40-J47)	125	2.2	8.5
10	Chronic liver disease and cirrhosis (K70,K73-K74)	123	2.1	8.3
11	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	94	1.6	6.4
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	82	1.4	5.6
13	Influenza and pneumonia (J09-J18)	74	1.3	5.0
14	Anemias (D50-D64)	52	0.9	3.5
15	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	36	0.6	2.4
...	All other causes (Residual)	1,043	18.2	70.7
Non-Hispanic, black, female, 50-54 years				
...	All causes	8,486	100.0	583.5
1	Malignant neoplasms (C00-C97)	2,750	32.4	189.1
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,730	20.4	119.0
3	Cerebrovascular diseases (I60-I69)	419	4.9	28.8
4	Diabetes mellitus (E10-E14)	406	4.8	27.9
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	394	4.6	27.1
6	Chronic lower respiratory diseases (J40-J47)	253	3.0	17.4
7	Human immunodeficiency virus (HIV) disease (B20-B24)	233	2.7	16.0
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	197	2.3	13.5
9	Chronic liver disease and cirrhosis (K70,K73-K74)	187	2.2	12.9
10	Septicemia (A40-A41)	181	2.1	12.4
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	116	1.4	8.0
12	Influenza and pneumonia (J09-J18)	107	1.3	7.4
13	Viral hepatitis (B15-B19)	68	0.8	4.7
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	63	0.7	4.3
15	Aortic aneurysm and dissection (I71)	33	0.4	2.3
...	All other causes (Residual)	1,349	15.9	92.8

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, female, 55-59 years				
...	All causes	10,390	100.0	836.1
1	Malignant neoplasms (C00-C97)	3,513	33.8	282.7
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,269	21.8	182.6
3	Cerebrovascular diseases (I60-I69)	517	5.0	41.6
4	Diabetes mellitus (E10-E14)	516	5.0	41.5
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	352	3.4	28.3
6	Chronic lower respiratory diseases (J40-J47)	305	2.9	24.5
7	Septicemia (A40-A41)	262	2.5	21.1
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	254	2.4	20.4
9	Chronic liver disease and cirrhosis (K70,K73-K74)	202	1.9	16.3
10	Human immunodeficiency virus (HIV) disease (B20-B24)	182	1.8	14.6
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	177	1.7	14.2
12	Influenza and pneumonia (J09-J18)	136	1.3	10.9
13	Viral hepatitis (B15-B19)	101	1.0	8.1
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	45	0.4	3.6
15	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	39	0.4	3.1
...	All other causes (Residual)	1,520	14.6	122.3
Non-Hispanic, black, female, 60-64 years				
...	All causes	11,635	100.0	1,156.0
1	Malignant neoplasms (C00-C97)	4,062	34.9	403.6
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,576	22.1	255.9
3	Diabetes mellitus (E10-E14)	695	6.0	69.1
4	Cerebrovascular diseases (I60-I69)	602	5.2	59.8
5	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	379	3.3	37.7
6	Chronic lower respiratory diseases (J40-J47)	363	3.1	36.1
7	Septicemia (A40-A41)	249	2.1	24.7
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	230	2.0	22.9
9	Essential hypertension and hypertensive renal disease (I10,I12,I15)	191	1.6	19.0
10	Influenza and pneumonia (J09-J18)	168	1.4	16.7
11	Chronic liver disease and cirrhosis (K70,K73-K74)	162	1.4	16.1
12	Human immunodeficiency virus (HIV) disease (B20-B24)	100	0.9	9.9
13	Viral hepatitis (B15-B19)	87	0.7	8.6
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	60	0.5	6.0
15	Pneumonitis due to solids and liquids (J69)	42	0.4	4.2
...	All other causes (Residual)	1,669	14.3	165.8

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, female, 65-69 years				
...	All causes	11,521	100.0	1,692.5
1	Malignant neoplasms (C00-C97)	3,846	33.4	565.0
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,641	22.9	388.0
3	Diabetes mellitus (E10-E14)	768	6.7	112.8
4	Cerebrovascular diseases (I60-I69)	606	5.3	89.0
5	Chronic lower respiratory diseases (J40-J47)	427	3.7	62.7
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	389	3.4	57.1
7	Septicemia (A40-A41)	299	2.6	43.9
8	Essential hypertension and hypertensive renal disease (I10,I12,I15)	188	1.6	27.6
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	182	1.6	26.7
10	Influenza and pneumonia (J09-J18)	179	1.6	26.3
11	Chronic liver disease and cirrhosis (K70,K73-K74)	90	0.8	13.2
12	Alzheimer's disease (G30)	60	0.5	8.8
13	Pneumonitis due to solids and liquids (J69)	54	0.5	7.9
14	Human immunodeficiency virus (HIV) disease (B20-B24)	52	0.5	7.6
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	51	0.4	7.5
...	All other causes (Residual)	1,689	14.7	248.1
Non-Hispanic, black, female, 70-74 years				
...	All causes	12,626	100.0	2,447.6
1	Malignant neoplasms (C00-C97)	3,781	29.9	733.0
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,027	24.0	586.8
3	Cerebrovascular diseases (I60-I69)	822	6.5	159.3
4	Diabetes mellitus (E10-E14)	735	5.8	142.5
5	Chronic lower respiratory diseases (J40-J47)	530	4.2	102.7
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	471	3.7	91.3
7	Septicemia (A40-A41)	357	2.8	69.2
8	Essential hypertension and hypertensive renal disease (I10,I12,I15)	244	1.9	47.3
9	Influenza and pneumonia (J09-J18)	226	1.8	43.8
10	Alzheimer's disease (G30)	162	1.3	31.4
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	155	1.2	30.0
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	68	0.5	13.2
13	Chronic liver disease and cirrhosis (K70,K73-K74)	53	0.4	10.3
14	Pneumonitis due to solids and liquids (J69)	51	0.4	9.9
15	Aortic aneurysm and dissection (I71)	49	0.4	9.5
...	All other causes (Residual)	1,895	15.0	367.4

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, female, 75-79 years				
...	All causes	14,626	100.0	3,720.0
1	Malignant neoplasms (C00-C97)	3,674	25.1	934.5
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,616	24.7	919.7
3	Cerebrovascular diseases (I60-I69)	1,044	7.1	265.5
4	Diabetes mellitus (E10-E14)	864	5.9	219.8
5	Chronic lower respiratory diseases (J40-J47)	583	4.0	148.3
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	572	3.9	145.5
7	Septicemia (A40-A41)	407	2.8	103.5
8	Alzheimer's disease (G30)	405	2.8	103.0
9	Influenza and pneumonia (J09-J18)	299	2.0	76.0
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	276	1.9	70.2
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	167	1.1	42.5
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	97	0.7	24.7
13	Pneumonitis due to solids and liquids (J69)	86	0.6	21.9
14	Parkinson's disease (G20-G21)	74	0.5	18.8
15	Aortic aneurysm and dissection (I71)	59	0.4	15.0
15	Enterocolitis due to <i>Clostridium difficile</i> (A04.7)	59	0.4	15.0
...	All other causes (Residual)	2,344	16.0	596.2
Non-Hispanic, black, female, 80-84 years				
...	All causes	16,680	100.0	5,917.2
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	4,364	26.2	1,548.1
2	Malignant neoplasms (C00-C97)	3,303	19.8	1,171.7
3	Cerebrovascular diseases (I60-I69)	1,311	7.9	465.1
4	Diabetes mellitus (E10-E14)	878	5.3	311.5
5	Alzheimer's disease (G30)	689	4.1	244.4
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	628	3.8	222.8
7	Chronic lower respiratory diseases (J40-J47)	612	3.7	217.1
8	Septicemia (A40-A41)	400	2.4	141.9
9	Essential hypertension and hypertensive renal disease (I10,I12,I15)	365	2.2	129.5
10	Influenza and pneumonia (J09-J18)	344	2.1	122.0
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	192	1.2	68.1
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	103	0.6	36.5
13	Pneumonitis due to solids and liquids (J69)	99	0.6	35.1
14	Parkinson's disease (G20-G21)	89	0.5	31.6
15	Enterocolitis due to <i>Clostridium difficile</i> (A04.7)	61	0.4	21.6
...	All other causes (Residual)	3,242	19.4	1,150.1

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, female, 85-89 years				
...	All causes	16,455	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	4,586	27.9	---
2	Malignant neoplasms (C00-C97)	2,372	14.4	---
3	Cerebrovascular diseases (I60-I69)	1,270	7.7	---
4	Alzheimer's disease (G30)	1,006	6.1	---
5	Diabetes mellitus (E10-E14)	698	4.2	---
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	554	3.4	---
7	Chronic lower respiratory diseases (J40-J47)	528	3.2	---
8	Essential hypertension and hypertensive renal disease (I10,I12,I15)	419	2.5	---
9	Septicemia (A40-A41)	417	2.5	---
10	Influenza and pneumonia (J09-J18)	414	2.5	---
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	157	1.0	---
12	Pneumonitis due to solids and liquids (J69)	129	0.8	---
13	Parkinson's disease (G20-G21)	89	0.5	---
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	81	0.5	---
15	Atherosclerosis (I70)	66	0.4	---
...	All other causes (Residual)	3,669	22.3	---
Non-Hispanic, black, female, 90-94 years				
...	All causes	11,789	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,418	29.0	---
2	Malignant neoplasms (C00-C97)	1,187	10.1	---
3	Cerebrovascular diseases (I60-I69)	955	8.1	---
4	Alzheimer's disease (G30)	911	7.7	---
5	Diabetes mellitus (E10-E14)	425	3.6	---
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	369	3.1	---
7	Essential hypertension and hypertensive renal disease (I10,I12,I15)	354	3.0	---
8	Influenza and pneumonia (J09-J18)	308	2.6	---
9	Chronic lower respiratory diseases (J40-J47)	284	2.4	---
10	Septicemia (A40-A41)	248	2.1	---
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	131	1.1	---
12	Pneumonitis due to solids and liquids (J69)	105	0.9	---
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	63	0.5	---
14	Atherosclerosis (I70)	54	0.5	---
15	Anemias (D50-D64)	48	0.4	---
...	All other causes (Residual)	2,929	24.8	---

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, female, 95-99 years				
...	All causes	5,719	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,700	29.7	---
2	Alzheimer's disease (G30)	476	8.3	---
3	Cerebrovascular diseases (I60-I69)	409	7.2	---
4	Malignant neoplasms (C00-C97)	387	6.8	---
5	Essential hypertension and hypertensive renal disease (I10,I12,I15)	200	3.5	---
6	Diabetes mellitus (E10-E14)	174	3.0	---
7	Influenza and pneumonia (J09-J18)	169	3.0	---
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	143	2.5	---
9	Chronic lower respiratory diseases (J40-J47)	112	2.0	---
10	Septicemia (A40-A41)	105	1.8	---
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	59	1.0	---
12	Pneumonitis due to solids and liquids (J69)	51	0.9	---
13	Atherosclerosis (I70)	27	0.5	---
14	Anemias (D50-D64)	19	0.3	---
14	Nutritional deficiencies (E40-E64)	19	0.3	---
...	All other causes (Residual)	1,669	29.2	---
Non-Hispanic, black, female, 100 years and over				
...	All causes	1,872	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	588	31.4	---
2	Alzheimer's disease (G30)	142	7.6	---
3	Cerebrovascular diseases (I60-I69)	125	6.7	---
4	Malignant neoplasms (C00-C97)	87	4.6	---
5	Essential hypertension and hypertensive renal disease (I10,I12,I15)	72	3.8	---
6	Influenza and pneumonia (J09-J18)	68	3.6	---
7	Diabetes mellitus (E10-E14)	36	1.9	---
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	35	1.9	---
9	Septicemia (A40-A41)	30	1.6	---
10	Chronic lower respiratory diseases (J40-J47)	24	1.3	---
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	23	1.2	---
12	Atherosclerosis (I70)	19	1.0	---
13	Pneumonitis due to solids and liquids (J69)	18	1.0	---
14	Anemias (D50-D64)	10	0.5	---
15	Nutritional deficiencies (E40-E64)	7	0.4	---
...	All other causes (Residual)	588	31.4	---

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011