

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> , Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, both sexes, all ages				
...	All causes	149,635	100.0	287.5
1	Malignant neoplasms (C00-C97)	32,381	21.6	62.2
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	30,385	20.3	58.4
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	11,166	7.5	21.5
4	Cerebrovascular diseases (I60-I69)	7,552	5.0	14.5
5	Diabetes mellitus (E10-E14)	7,054	4.7	13.6
6	Chronic liver disease and cirrhosis (K70,K73-K74)	4,774	3.2	9.2
7	Chronic lower respiratory diseases (J40-J47)	4,355	2.9	8.4
8	Alzheimer's disease (G30)	3,599	2.4	6.9
9	Influenza and pneumonia (J09-J18)	3,256	2.2	6.3
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2,799	1.9	5.4
11	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	2,759	1.8	5.3
12	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,720	1.8	5.2
13	Certain conditions originating in the perinatal period (P00-P96)	2,387	1.6	4.6
14	Septicemia (A40-A41)	2,168	1.4	4.2
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	1,803	1.2	3.5
...	All other causes (Residual)	30,477	20.4	58.6
Hispanic, both sexes, 1-4 years				
...	All causes	972	100.0	23.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	317	32.6	7.7
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	126	13.0	3.0
3	Malignant neoplasms (C00-C97)	97	10.0	2.3
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	94	9.7	2.3
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	35	3.6	0.8
6	Influenza and pneumonia (J09-J18)	21	2.2	0.5
7	Septicemia (A40-A41)	16	1.6	*
8	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	10	1.0	*
8	Cerebrovascular diseases (I60-I69)	10	1.0	*
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	7	0.7	*
11	Complications of medical and surgical care (Y40-Y84,Y88)	6	0.6	*
12	Certain conditions originating in the perinatal period (P00-P96)	5	0.5	*
13	Acute bronchitis and bronchiolitis (J20-J21)	4	0.4	*
13	Chronic lower respiratory diseases (J40-J47)	4	0.4	*
15	Meningitis (G00,G03)	3	0.3	*
...	All other causes (Residual)	217	22.3	5.2

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> , Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, both sexes, 5-14 years				
...	All causes	1,052	100.0	11.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	278	26.4	2.9
2	Malignant neoplasms (C00-C97)	200	19.0	2.1
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	92	8.7	1.0
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	58	5.5	0.6
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	54	5.1	0.6
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	45	4.3	0.5
7	Chronic lower respiratory diseases (J40-J47)	23	2.2	0.2
8	Cerebrovascular diseases (I60-I69)	22	2.1	0.2
8	Influenza and pneumonia (J09-J18)	22	2.1	0.2
10	Septicemia (A40-A41)	10	1.0	*
11	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	7	0.7	*
12	Anemias (D50-D64)	6	0.6	*
12	Diabetes mellitus (E10-E14)	6	0.6	*
14	Certain conditions originating in the perinatal period (P00-P96)	5	0.5	*
15	Meningitis (G00,G03)	4	0.4	*
...	All other causes (Residual)	220	20.9	2.3
Hispanic, both sexes, 15-24 years				
...	All causes	4,826	100.0	53.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,911	39.6	21.1
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	933	19.3	10.3
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	635	13.2	7.0
4	Malignant neoplasms (C00-C97)	383	7.9	4.2
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	156	3.2	1.7
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	56	1.2	0.6
7	Cerebrovascular diseases (I60-I69)	53	1.1	0.6
8	Legal intervention (Y35,Y89.0)	34	0.7	0.4
9	Pregnancy, childbirth and the puerperium (O00-O99)	31	0.6	0.3
10	Influenza and pneumonia (J09-J18)	28	0.6	0.3
11	Human immunodeficiency virus (HIV) disease (B20-B24)	21	0.4	0.2
12	Diabetes mellitus (E10-E14)	19	0.4	*
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	18	0.4	*
14	Chronic lower respiratory diseases (J40-J47)	17	0.4	*
15	Septicemia (A40-A41)	16	0.3	*
...	All other causes (Residual)	515	10.7	5.7

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> , Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, both sexes, 25-34 years				
...	All causes	6,061	100.0	70.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,096	34.6	24.4
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	782	12.9	9.1
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	654	10.8	7.6
4	Malignant neoplasms (C00-C97)	595	9.8	6.9
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	386	6.4	4.5
6	Human immunodeficiency virus (HIV) disease (B20-B24)	119	2.0	1.4
7	Cerebrovascular diseases (I60-I69)	110	1.8	1.3
8	Chronic liver disease and cirrhosis (K70,K73-K74)	106	1.7	1.2
9	Influenza and pneumonia (J09-J18)	79	1.3	0.9
10	Diabetes mellitus (E10-E14)	78	1.3	0.9
11	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	56	0.9	0.7
12	Pregnancy, childbirth and the puerperium (O00-O99)	55	0.9	0.6
13	Septicemia (A40-A41)	51	0.8	0.6
14	Legal intervention (Y35,Y89.0)	34	0.6	0.4
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	25	0.4	0.3
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	25	0.4	0.3
...	All other causes (Residual)	810	13.4	9.4
Hispanic, both sexes, 35-44 years				
...	All causes	8,192	100.0	109.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,807	22.1	24.1
2	Malignant neoplasms (C00-C97)	1,509	18.4	20.1
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	945	11.5	12.6
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	523	6.4	7.0
5	Chronic liver disease and cirrhosis (K70,K73-K74)	469	5.7	6.2
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	444	5.4	5.9
7	Human immunodeficiency virus (HIV) disease (B20-B24)	271	3.3	3.6
8	Cerebrovascular diseases (I60-I69)	255	3.1	3.4
9	Diabetes mellitus (E10-E14)	227	2.8	3.0
10	Influenza and pneumonia (J09-J18)	108	1.3	1.4
11	Septicemia (A40-A41)	94	1.1	1.3
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	69	0.8	0.9
13	Viral hepatitis (B15-B19)	65	0.8	0.9
14	Chronic lower respiratory diseases (J40-J47)	51	0.6	0.7
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	46	0.6	0.6
...	All other causes (Residual)	1,309	16.0	17.4

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> , Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, both sexes, 45-54 years				
...	All causes	15,549	100.0	272.0
1	Malignant neoplasms (C00-C97)	3,871	24.9	67.7
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,531	16.3	44.3
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,761	11.3	30.8
4	Chronic liver disease and cirrhosis (K70,K73-K74)	1,366	8.8	23.9
5	Diabetes mellitus (E10-E14)	671	4.3	11.7
6	Cerebrovascular diseases (I60-I69)	669	4.3	11.7
7	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	423	2.7	7.4
8	Human immunodeficiency virus (HIV) disease (B20-B24)	396	2.5	6.9
9	Viral hepatitis (B15-B19)	368	2.4	6.4
10	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	235	1.5	4.1
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	231	1.5	4.0
12	Septicemia (A40-A41)	217	1.4	3.8
13	Influenza and pneumonia (J09-J18)	204	1.3	3.6
14	Chronic lower respiratory diseases (J40-J47)	152	1.0	2.7
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	131	0.8	2.3
...	All other causes (Residual)	2,323	14.9	40.6
Hispanic, both sexes, 55-64 years				
...	All causes	20,958	100.0	606.1
1	Malignant neoplasms (C00-C97)	6,626	31.6	191.6
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	4,139	19.7	119.7
3	Chronic liver disease and cirrhosis (K70,K73-K74)	1,347	6.4	39.0
4	Diabetes mellitus (E10-E14)	1,267	6.0	36.6
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,017	4.9	29.4
6	Cerebrovascular diseases (I60-I69)	961	4.6	27.8
7	Viral hepatitis (B15-B19)	473	2.3	13.7
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	412	2.0	11.9
9	Chronic lower respiratory diseases (J40-J47)	390	1.9	11.3
10	Septicemia (A40-A41)	363	1.7	10.5
11	Influenza and pneumonia (J09-J18)	272	1.3	7.9
12	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	226	1.1	6.5
13	Human immunodeficiency virus (HIV) disease (B20-B24)	215	1.0	6.2
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	176	0.8	5.1
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	104	0.5	3.0
15	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	104	0.5	3.0
...	All other causes (Residual)	2,866	13.7	82.9

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> , Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, both sexes, 65-74 years				
...	All causes	23,984	100.0	1,366.2
1	Malignant neoplasms (C00-C97)	7,527	31.4	428.8
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	5,206	21.7	296.6
3	Diabetes mellitus (E10-E14)	1,701	7.1	96.9
4	Cerebrovascular diseases (I60-I69)	1,206	5.0	68.7
5	Chronic liver disease and cirrhosis (K70,K73-K74)	829	3.5	47.2
6	Chronic lower respiratory diseases (J40-J47)	799	3.3	45.5
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	607	2.5	34.6
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	591	2.5	33.7
9	Influenza and pneumonia (J09-J18)	449	1.9	25.6
10	Septicemia (A40-A41)	417	1.7	23.8
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	272	1.1	15.5
12	Alzheimer's disease (G30)	219	0.9	12.5
13	Viral hepatitis (B15-B19)	169	0.7	9.6
14	Parkinson's disease (G20-G21)	161	0.7	9.2
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	138	0.6	7.9
...	All other causes (Residual)	3,693	15.4	210.4
Hispanic, both sexes, 75-84 years				
...	All causes	32,621	100.0	3,585.5
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	7,945	24.4	873.3
2	Malignant neoplasms (C00-C97)	7,730	23.7	849.6
3	Cerebrovascular diseases (I60-I69)	2,171	6.7	238.6
4	Diabetes mellitus (E10-E14)	1,917	5.9	210.7
5	Chronic lower respiratory diseases (J40-J47)	1,460	4.5	160.5
6	Alzheimer's disease (G30)	1,160	3.6	127.5
7	Influenza and pneumonia (J09-J18)	883	2.7	97.1
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	770	2.4	84.6
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	650	2.0	71.4
10	Septicemia (A40-A41)	535	1.6	58.8
11	Chronic liver disease and cirrhosis (K70,K73-K74)	510	1.6	56.1
12	Parkinson's disease (G20-G21)	476	1.5	52.3
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	465	1.4	51.1
14	Pneumonitis due to solids and liquids (J69)	226	0.7	24.8
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	216	0.7	23.7
...	All other causes (Residual)	5,507	16.9	605.3

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> , Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, both sexes, 85 years and over				
...	All causes	30,588	100.0	10,072.2
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	8,932	29.2	2,941.2
2	Malignant neoplasms (C00-C97)	3,828	12.5	1,260.5
3	Alzheimer's disease (G30)	2,145	7.0	706.3
4	Cerebrovascular diseases (I60-I69)	2,074	6.8	682.9
5	Chronic lower respiratory diseases (J40-J47)	1,432	4.7	471.5
6	Diabetes mellitus (E10-E14)	1,166	3.8	383.9
7	Influenza and pneumonia (J09-J18)	1,160	3.8	382.0
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	651	2.1	214.4
9	Essential hypertension and hypertensive renal disease (I10,I12,I15)	597	2.0	196.6
10	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	576	1.9	189.7
11	Septicemia (A40-A41)	416	1.4	137.0
12	Parkinson's disease (G20-G21)	388	1.3	127.8
13	Pneumonitis due to solids and liquids (J69)	304	1.0	100.1
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	175	0.6	57.6
15	Enterocolitis due to <i>Clostridium difficile</i> (A04.7)	136	0.4	44.8
...	All other causes (Residual)	6,608	21.6	2,175.9
Hispanic, male, all ages				
...	All causes	81,887	100.0	309.7
1	Malignant neoplasms (C00-C97)	17,185	21.0	65.0
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	16,526	20.2	62.5
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	8,098	9.9	30.6
4	Diabetes mellitus (E10-E14)	3,726	4.6	14.1
5	Cerebrovascular diseases (I60-I69)	3,528	4.3	13.3
6	Chronic liver disease and cirrhosis (K70,K73-K74)	3,311	4.0	12.5
7	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	2,279	2.8	8.6
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,224	2.7	8.4
9	Chronic lower respiratory diseases (J40-J47)	2,222	2.7	8.4
10	Influenza and pneumonia (J09-J18)	1,606	2.0	6.1
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,388	1.7	5.2
12	Certain conditions originating in the perinatal period (P00-P96)	1,297	1.6	4.9
13	Alzheimer's disease (G30)	1,151	1.4	4.4
14	Septicemia (A40-A41)	1,048	1.3	4.0
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	958	1.2	3.6
...	All other causes (Residual)	15,340	18.7	58.0

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> , Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, male, 1-4 years				
...	All causes	539	100.0	25.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	187	34.7	8.8
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	70	13.0	3.3
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	55	10.2	2.6
4	Malignant neoplasms (C00-C97)	51	9.5	2.4
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	20	3.7	0.9
6	Influenza and pneumonia (J09-J18)	10	1.9	*
7	Septicemia (A40-A41)	8	1.5	*
8	Cerebrovascular diseases (I60-I69)	7	1.3	*
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	4	0.7	*
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	3	0.6	*
10	Meningitis (G00,G03)	3	0.6	*
10	Acute bronchitis and bronchiolitis (J20-J21)	3	0.6	*
10	Certain conditions originating in the perinatal period (P00-P96)	3	0.6	*
10	Complications of medical and surgical care (Y40-Y84,Y88)	3	0.6	*
15	Chronic lower respiratory diseases (J40-J47)	2	0.4	*
...	All other causes (Residual)	110	20.4	5.2
Hispanic, male, 5-14 years				
...	All causes	608	100.0	12.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	182	29.9	3.7
2	Malignant neoplasms (C00-C97)	108	17.8	2.2
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	48	7.9	1.0
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	38	6.3	0.8
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	36	5.9	0.7
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	17	2.8	*
7	Cerebrovascular diseases (I60-I69)	12	2.0	*
7	Chronic lower respiratory diseases (J40-J47)	12	2.0	*
9	Influenza and pneumonia (J09-J18)	9	1.5	*
10	Septicemia (A40-A41)	5	0.8	*
10	Anemias (D50-D64)	5	0.8	*
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	4	0.7	*
13	Meningitis (G00,G03)	3	0.5	*
14	Diabetes mellitus (E10-E14)	2	0.3	*
14	Diseases of appendix (K35-K38)	2	0.3	*
...	All other causes (Residual)	125	20.6	2.6

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> , Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, male, 15-24 years				
...	All causes	3,723	100.0	78.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,517	40.7	32.0
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	835	22.4	17.6
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	510	13.7	10.7
4	Malignant neoplasms (C00-C97)	250	6.7	5.3
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	98	2.6	2.1
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	37	1.0	0.8
7	Legal intervention (Y35,Y89.0)	33	0.9	0.7
8	Cerebrovascular diseases (I60-I69)	32	0.9	0.7
9	Influenza and pneumonia (J09-J18)	17	0.5	*
10	Chronic lower respiratory diseases (J40-J47)	13	0.3	*
11	Human immunodeficiency virus (HIV) disease (B20-B24)	11	0.3	*
12	Anemias (D50-D64)	9	0.2	*
12	Diabetes mellitus (E10-E14)	9	0.2	*
14	Septicemia (A40-A41)	8	0.2	*
14	Chronic liver disease and cirrhosis (K70,K73-K74)	8	0.2	*
...	All other causes (Residual)	336	9.0	7.1
Hispanic, male, 25-34 years				
...	All causes	4,467	100.0	98.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,691	37.9	37.3
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	664	14.9	14.6
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	546	12.2	12.0
4	Malignant neoplasms (C00-C97)	304	6.8	6.7
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	283	6.3	6.2
6	Human immunodeficiency virus (HIV) disease (B20-B24)	96	2.1	2.1
7	Chronic liver disease and cirrhosis (K70,K73-K74)	76	1.7	1.7
8	Cerebrovascular diseases (I60-I69)	71	1.6	1.6
9	Influenza and pneumonia (J09-J18)	52	1.2	1.1
10	Diabetes mellitus (E10-E14)	44	1.0	1.0
11	Legal intervention (Y35,Y89.0)	34	0.8	0.7
12	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	31	0.7	0.7
13	Septicemia (A40-A41)	30	0.7	0.7
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	15	0.3	*
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	13	0.3	*
...	All other causes (Residual)	517	11.6	11.4

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> , Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, male, 35-44 years				
...	All causes	5,478	100.0	142.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,386	25.3	36.0
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	701	12.8	18.2
3	Malignant neoplasms (C00-C97)	631	11.5	16.4
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	433	7.9	11.3
5	Chronic liver disease and cirrhosis (K70,K73-K74)	376	6.9	9.8
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	345	6.3	9.0
7	Human immunodeficiency virus (HIV) disease (B20-B24)	211	3.9	5.5
8	Cerebrovascular diseases (I60-I69)	156	2.8	4.1
9	Diabetes mellitus (E10-E14)	147	2.7	3.8
10	Influenza and pneumonia (J09-J18)	64	1.2	1.7
11	Septicemia (A40-A41)	51	0.9	1.3
12	Viral hepatitis (B15-B19)	50	0.9	1.3
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	45	0.8	1.2
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	34	0.6	0.9
15	Aortic aneurysm and dissection (I71)	29	0.5	0.8
...	All other causes (Residual)	819	15.0	21.3
Hispanic, male, 45-54 years				
...	All causes	10,271	100.0	357.3
1	Malignant neoplasms (C00-C97)	1,989	19.4	69.2
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,897	18.5	66.0
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,347	13.1	46.9
4	Chronic liver disease and cirrhosis (K70,K73-K74)	1,048	10.2	36.5
5	Diabetes mellitus (E10-E14)	439	4.3	15.3
6	Cerebrovascular diseases (I60-I69)	399	3.9	13.9
7	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	335	3.3	11.7
8	Human immunodeficiency virus (HIV) disease (B20-B24)	295	2.9	10.3
9	Viral hepatitis (B15-B19)	287	2.8	10.0
10	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	191	1.9	6.6
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	145	1.4	5.0
12	Influenza and pneumonia (J09-J18)	134	1.3	4.7
13	Septicemia (A40-A41)	113	1.1	3.9
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	85	0.8	3.0
15	Chronic lower respiratory diseases (J40-J47)	76	0.7	2.6
...	All other causes (Residual)	1,491	14.5	51.9

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> , Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, male, 55-64 years				
...	All causes	13,112	100.0	791.0
1	Malignant neoplasms (C00-C97)	3,718	28.4	224.3
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,856	21.8	172.3
3	Chronic liver disease and cirrhosis (K70,K73-K74)	1,000	7.6	60.3
4	Diabetes mellitus (E10-E14)	762	5.8	46.0
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	739	5.6	44.6
6	Cerebrovascular diseases (I60-I69)	583	4.4	35.2
7	Viral hepatitis (B15-B19)	342	2.6	20.6
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	243	1.9	14.7
9	Septicemia (A40-A41)	219	1.7	13.2
10	Chronic lower respiratory diseases (J40-J47)	218	1.7	13.2
11	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	189	1.4	11.4
12	Human immunodeficiency virus (HIV) disease (B20-B24)	161	1.2	9.7
12	Influenza and pneumonia (J09-J18)	161	1.2	9.7
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	108	0.8	6.5
15	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	78	0.6	4.7
...	All other causes (Residual)	1,735	13.2	104.7
Hispanic, male, 65-74 years				
...	All causes	13,540	100.0	1,722.4
1	Malignant neoplasms (C00-C97)	4,152	30.7	528.2
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,246	24.0	412.9
3	Diabetes mellitus (E10-E14)	962	7.1	122.4
4	Cerebrovascular diseases (I60-I69)	632	4.7	80.4
5	Chronic liver disease and cirrhosis (K70,K73-K74)	500	3.7	63.6
6	Chronic lower respiratory diseases (J40-J47)	453	3.3	57.6
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	364	2.7	46.3
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	279	2.1	35.5
9	Influenza and pneumonia (J09-J18)	246	1.8	31.3
10	Septicemia (A40-A41)	220	1.6	28.0
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	147	1.1	18.7
12	Parkinson's disease (G20-G21)	116	0.9	14.8
13	Alzheimer's disease (G30)	97	0.7	12.3
14	Viral hepatitis (B15-B19)	92	0.7	11.7
15	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	86	0.6	10.9
...	All other causes (Residual)	1,948	14.4	247.8

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> , Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, male, 75-84 years				
...	All causes	16,081	100.0	4,303.5
1	Malignant neoplasms (C00-C97)	4,127	25.7	1,104.5
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	4,083	25.4	1,092.7
3	Diabetes mellitus (E10-E14)	925	5.8	247.5
4	Cerebrovascular diseases (I60-I69)	917	5.7	245.4
5	Chronic lower respiratory diseases (J40-J47)	788	4.9	210.9
6	Alzheimer's disease (G30)	431	2.7	115.3
7	Influenza and pneumonia (J09-J18)	429	2.7	114.8
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	359	2.2	96.1
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	358	2.2	95.8
10	Parkinson's disease (G20-G21)	279	1.7	74.7
11	Chronic liver disease and cirrhosis (K70,K73-K74)	248	1.5	66.4
12	Septicemia (A40-A41)	225	1.4	60.2
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	206	1.3	55.1
14	Pneumonitis due to solids and liquids (J69)	137	0.9	36.7
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	124	0.8	33.2
...	All other causes (Residual)	2,445	15.2	654.3
Hispanic, male, 85 years and over				
...	All causes	11,442	100.0	10,707.8
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,287	28.7	3,076.1
2	Malignant neoplasms (C00-C97)	1,847	16.1	1,728.5
3	Cerebrovascular diseases (I60-I69)	711	6.2	665.4
4	Chronic lower respiratory diseases (J40-J47)	625	5.5	584.9
5	Alzheimer's disease (G30)	593	5.2	554.9
6	Influenza and pneumonia (J09-J18)	463	4.0	433.3
7	Diabetes mellitus (E10-E14)	435	3.8	407.1
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	284	2.5	265.8
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	235	2.1	219.9
10	Parkinson's disease (G20-G21)	199	1.7	186.2
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	191	1.7	178.7
12	Septicemia (A40-A41)	154	1.3	144.1
13	Pneumonitis due to solids and liquids (J69)	132	1.2	123.5
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	82	0.7	76.7
15	Chronic liver disease and cirrhosis (K70,K73-K74)	53	0.5	49.6
...	All other causes (Residual)	2,151	18.8	2,013.0

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> , Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, female, all ages				
...	All causes	67,748	100.0	264.6
1	Malignant neoplasms (C00-C97)	15,196	22.4	59.4
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	13,859	20.5	54.1
3	Cerebrovascular diseases (I60-I69)	4,024	5.9	15.7
4	Diabetes mellitus (E10-E14)	3,328	4.9	13.0
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,068	4.5	12.0
6	Alzheimer's disease (G30)	2,448	3.6	9.6
7	Chronic lower respiratory diseases (J40-J47)	2,133	3.1	8.3
8	Influenza and pneumonia (J09-J18)	1,650	2.4	6.4
9	Chronic liver disease and cirrhosis (K70,K73-K74)	1,463	2.2	5.7
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,411	2.1	5.5
11	Septicemia (A40-A41)	1,120	1.7	4.4
12	Certain conditions originating in the perinatal period (P00-P96)	1,090	1.6	4.3
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	927	1.4	3.6
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	845	1.2	3.3
15	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	496	0.7	1.9
...	All other causes (Residual)	14,690	21.7	57.4
Hispanic, female, 1-4 years				
...	All causes	433	100.0	21.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	130	30.0	6.4
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	56	12.9	2.8
3	Malignant neoplasms (C00-C97)	46	10.6	2.3
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	39	9.0	1.9
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	15	3.5	*
6	Influenza and pneumonia (J09-J18)	11	2.5	*
7	Septicemia (A40-A41)	8	1.8	*
8	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	7	1.6	*
9	Cerebrovascular diseases (I60-I69)	3	0.7	*
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3	0.7	*
9	Complications of medical and surgical care (Y40-Y84,Y88)	3	0.7	*
12	Chronic lower respiratory diseases (J40-J47)	2	0.5	*
12	Diseases of appendix (K35-K38)	2	0.5	*
12	Hernia (K40-K46)	2	0.5	*
12	Certain conditions originating in the perinatal period (P00-P96)	2	0.5	*
...	All other causes (Residual)	104	24.0	5.1

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> , Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, female, 5-14 years				
...	All causes	444	100.0	9.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	96	21.6	2.1
2	Malignant neoplasms (C00-C97)	92	20.7	2.0
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	44	9.9	0.9
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	28	6.3	0.6
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	20	4.5	0.4
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	18	4.1	*
7	Influenza and pneumonia (J09-J18)	13	2.9	*
8	Chronic lower respiratory diseases (J40-J47)	11	2.5	*
9	Cerebrovascular diseases (I60-I69)	10	2.3	*
10	Septicemia (A40-A41)	5	1.1	*
11	Diabetes mellitus (E10-E14)	4	0.9	*
11	Certain conditions originating in the perinatal period (P00-P96)	4	0.9	*
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	3	0.7	*
14	Meningococcal infection (A39)	2	0.5	*
14	Pneumonitis due to solids and liquids (J69)	2	0.5	*
14	Complications of medical and surgical care (Y40-Y84,Y88)	2	0.5	*
...	All other causes (Residual)	90	20.3	1.9
Hispanic, female, 15-24 years				
...	All causes	1,103	100.0	25.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	394	35.7	9.1
2	Malignant neoplasms (C00-C97)	133	12.1	3.1
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	125	11.3	2.9
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	98	8.9	2.3
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	58	5.3	1.3
6	Pregnancy, childbirth and the puerperium (O00-O99)	31	2.8	0.7
7	Cerebrovascular diseases (I60-I69)	21	1.9	0.5
8	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	19	1.7	*
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	13	1.2	*
10	Influenza and pneumonia (J09-J18)	11	1.0	*
11	Human immunodeficiency virus (HIV) disease (B20-B24)	10	0.9	*
11	Diabetes mellitus (E10-E14)	10	0.9	*
13	Septicemia (A40-A41)	8	0.7	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	6	0.5	*
15	Anemias (D50-D64)	4	0.4	*
15	Chronic lower respiratory diseases (J40-J47)	4	0.4	*
...	All other causes (Residual)	158	14.3	3.7

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> , Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, female, 25-34 years				
...	All causes	1,594	100.0	39.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	405	25.4	10.0
2	Malignant neoplasms (C00-C97)	291	18.3	7.2
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	118	7.4	2.9
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	108	6.8	2.7
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	103	6.5	2.5
6	Pregnancy, childbirth and the puerperium (O00-O99)	55	3.5	1.4
7	Cerebrovascular diseases (I60-I69)	39	2.4	1.0
8	Diabetes mellitus (E10-E14)	34	2.1	0.8
9	Chronic liver disease and cirrhosis (K70,K73-K74)	30	1.9	0.7
10	Influenza and pneumonia (J09-J18)	27	1.7	0.7
11	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	25	1.6	0.6
12	Human immunodeficiency virus (HIV) disease (B20-B24)	23	1.4	0.6
13	Septicemia (A40-A41)	21	1.3	0.5
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	12	0.8	*
15	Chronic lower respiratory diseases (J40-J47)	10	0.6	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	10	0.6	*
...	All other causes (Residual)	283	17.8	7.0
Hispanic, female, 35-44 years				
...	All causes	2,714	100.0	74.1
1	Malignant neoplasms (C00-C97)	878	32.4	24.0
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	421	15.5	11.5
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	244	9.0	6.7
4	Cerebrovascular diseases (I60-I69)	99	3.6	2.7
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	99	3.6	2.7
6	Chronic liver disease and cirrhosis (K70,K73-K74)	93	3.4	2.5
7	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	90	3.3	2.5
8	Diabetes mellitus (E10-E14)	80	2.9	2.2
9	Human immunodeficiency virus (HIV) disease (B20-B24)	60	2.2	1.6
10	Influenza and pneumonia (J09-J18)	44	1.6	1.2
11	Septicemia (A40-A41)	43	1.6	1.2
12	Pregnancy, childbirth and the puerperium (O00-O99)	32	1.2	0.9
13	Chronic lower respiratory diseases (J40-J47)	29	1.1	0.8
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	24	0.9	0.7
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	17	0.6	*
...	All other causes (Residual)	461	17.0	12.6

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> , Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, female, 45-54 years				
...	All causes	5,278	100.0	185.7
1	Malignant neoplasms (C00-C97)	1,882	35.7	66.2
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	634	12.0	22.3
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	414	7.8	14.6
4	Chronic liver disease and cirrhosis (K70,K73-K74)	318	6.0	11.2
5	Cerebrovascular diseases (I60-I69)	270	5.1	9.5
6	Diabetes mellitus (E10-E14)	232	4.4	8.2
7	Septicemia (A40-A41)	104	2.0	3.7
8	Human immunodeficiency virus (HIV) disease (B20-B24)	101	1.9	3.6
9	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	88	1.7	3.1
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	86	1.6	3.0
11	Viral hepatitis (B15-B19)	81	1.5	2.9
12	Chronic lower respiratory diseases (J40-J47)	76	1.4	2.7
13	Influenza and pneumonia (J09-J18)	70	1.3	2.5
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	46	0.9	1.6
15	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	44	0.8	1.5
...	All other causes (Residual)	832	15.8	29.3
Hispanic, female, 55-64 years				
...	All causes	7,846	100.0	435.9
1	Malignant neoplasms (C00-C97)	2,908	37.1	161.6
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,283	16.4	71.3
3	Diabetes mellitus (E10-E14)	505	6.4	28.1
4	Cerebrovascular diseases (I60-I69)	378	4.8	21.0
5	Chronic liver disease and cirrhosis (K70,K73-K74)	347	4.4	19.3
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	278	3.5	15.4
7	Chronic lower respiratory diseases (J40-J47)	172	2.2	9.6
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	169	2.2	9.4
9	Septicemia (A40-A41)	144	1.8	8.0
10	Viral hepatitis (B15-B19)	131	1.7	7.3
11	Influenza and pneumonia (J09-J18)	111	1.4	6.2
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	68	0.9	3.8
13	Human immunodeficiency virus (HIV) disease (B20-B24)	54	0.7	3.0
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	45	0.6	2.5
15	Alzheimer's disease (G30)	40	0.5	2.2
...	All other causes (Residual)	1,213	15.5	67.4

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> , Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, female, 65-74 years				
...	All causes	10,444	100.0	1,077.4
1	Malignant neoplasms (C00-C97)	3,375	32.3	348.2
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,960	18.8	202.2
3	Diabetes mellitus (E10-E14)	739	7.1	76.2
4	Cerebrovascular diseases (I60-I69)	574	5.5	59.2
5	Chronic lower respiratory diseases (J40-J47)	346	3.3	35.7
6	Chronic liver disease and cirrhosis (K70,K73-K74)	329	3.2	33.9
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	328	3.1	33.8
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	227	2.2	23.4
9	Influenza and pneumonia (J09-J18)	203	1.9	20.9
10	Septicemia (A40-A41)	197	1.9	20.3
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	125	1.2	12.9
12	Alzheimer's disease (G30)	122	1.2	12.6
13	Viral hepatitis (B15-B19)	77	0.7	7.9
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	70	0.7	7.2
15	Enterocolitis due to <i>Clostridium difficile</i> (A04.7)	53	0.5	5.5
...	All other causes (Residual)	1,719	16.5	177.3
Hispanic, female, 75-84 years				
...	All causes	16,540	100.0	3,085.1
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,862	23.3	720.3
2	Malignant neoplasms (C00-C97)	3,603	21.8	672.0
3	Cerebrovascular diseases (I60-I69)	1,254	7.6	233.9
4	Diabetes mellitus (E10-E14)	992	6.0	185.0
5	Alzheimer's disease (G30)	729	4.4	136.0
6	Chronic lower respiratory diseases (J40-J47)	672	4.1	125.3
7	Influenza and pneumonia (J09-J18)	454	2.7	84.7
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	411	2.5	76.7
9	Septicemia (A40-A41)	310	1.9	57.8
10	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	292	1.8	54.5
11	Chronic liver disease and cirrhosis (K70,K73-K74)	262	1.6	48.9
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	259	1.6	48.3
13	Parkinson's disease (G20-G21)	197	1.2	36.7
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	92	0.6	17.2
15	Pneumonitis due to solids and liquids (J69)	89	0.5	16.6
...	All other causes (Residual)	3,062	18.5	571.1

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> , Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, female, 85 years and over				
...	All causes	19,146	100.0	9,727.2
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	5,645	29.5	2,868.0
2	Malignant neoplasms (C00-C97)	1,981	10.3	1,006.5
3	Alzheimer's disease (G30)	1,552	8.1	788.5
4	Cerebrovascular diseases (I60-I69)	1,363	7.1	692.5
5	Chronic lower respiratory diseases (J40-J47)	807	4.2	410.0
6	Diabetes mellitus (E10-E14)	731	3.8	371.4
7	Influenza and pneumonia (J09-J18)	697	3.6	354.1
8	Essential hypertension and hypertensive renal disease (I10,I12,I15)	406	2.1	206.3
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	367	1.9	186.5
10	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	341	1.8	173.2
11	Septicemia (A40-A41)	262	1.4	133.1
12	Parkinson's disease (G20-G21)	189	1.0	96.0
13	Pneumonitis due to solids and liquids (J69)	172	0.9	87.4
14	Enterocolitis due to <i>Clostridium difficile</i> (A04.7)	97	0.5	49.3
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	93	0.5	47.2
...	All other causes (Residual)	4,443	23.2	2,257.3
Non-Hispanic, white, both sexes, all ages				
...	All causes	2,006,319	100.0	1,001.0
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	482,979	24.1	241.0
2	Malignant neoplasms (C00-C97)	460,026	22.9	229.5
3	Chronic lower respiratory diseases (J40-J47)	127,035	6.3	63.4
4	Cerebrovascular diseases (I60-I69)	101,043	5.0	50.4
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	98,605	4.9	49.2
6	Alzheimer's disease (G30)	74,448	3.7	37.1
7	Diabetes mellitus (E10-E14)	50,964	2.5	25.4
8	Influenza and pneumonia (J09-J18)	43,350	2.2	21.6
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	33,492	1.7	16.7
10	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	33,031	1.6	16.5
11	Septicemia (A40-A41)	26,669	1.3	13.3
12	Chronic liver disease and cirrhosis (K70,K73-K74)	24,752	1.2	12.3
13	Parkinson's disease (G20-G21)	20,625	1.0	10.3
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	20,153	1.0	10.1
15	Pneumonitis due to solids and liquids (J69)	15,513	0.8	7.7
...	All other causes (Residual)	393,634	19.6	196.4

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> , Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, both sexes, 1-4 years				
...	All causes	2,055	100.0	24.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	724	35.2	8.5
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	236	11.5	2.8
3	Malignant neoplasms (C00-C97)	183	8.9	2.1
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	150	7.3	1.8
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	74	3.6	0.9
6	Influenza and pneumonia (J09-J18)	45	2.2	0.5
7	Septicemia (A40-A41)	30	1.5	0.4
8	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	24	1.2	0.3
9	Chronic lower respiratory diseases (J40-J47)	19	0.9	*
10	Certain conditions originating in the perinatal period (P00-P96)	17	0.8	*
11	Cerebrovascular diseases (I60-I69)	14	0.7	*
12	Acute bronchitis and bronchiolitis (J20-J21)	8	0.4	*
13	Complications of medical and surgical care (Y40-Y84,Y88)	7	0.3	*
14	Anemias (D50-D64)	6	0.3	*
15	Meningitis (G00,G03)	5	0.2	*
...	All other causes (Residual)	513	25.0	6.0
Non-Hispanic, white, both sexes, 5-14 years				
...	All causes	2,908	100.0	12.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	973	33.5	4.3
2	Malignant neoplasms (C00-C97)	481	16.5	2.1
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	168	5.8	0.7
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	168	5.8	0.7
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	125	4.3	0.5
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	97	3.3	0.4
7	Influenza and pneumonia (J09-J18)	61	2.1	0.3
8	Septicemia (A40-A41)	46	1.6	0.2
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	43	1.5	0.2
10	Cerebrovascular diseases (I60-I69)	38	1.3	0.2
11	Chronic lower respiratory diseases (J40-J47)	29	1.0	0.1
12	Certain conditions originating in the perinatal period (P00-P96)	16	0.6	*
13	Diabetes mellitus (E10-E14)	15	0.5	*
14	Pneumonitis due to solids and liquids (J69)	10	0.3	*
15	Anemias (D50-D64)	9	0.3	*
...	All other causes (Residual)	629	21.6	2.8

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> , Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, both sexes, 15-24 years				
...	All causes	16,961	100.0	67.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	8,528	50.3	33.8
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	3,413	20.1	13.5
3	Malignant neoplasms (C00-C97)	891	5.3	3.5
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	701	4.1	2.8
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	493	2.9	2.0
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	252	1.5	1.0
7	Influenza and pneumonia (J09-J18)	138	0.8	0.5
8	Diabetes mellitus (E10-E14)	97	0.6	0.4
9	Cerebrovascular diseases (I60-I69)	90	0.5	0.4
10	Septicemia (A40-A41)	77	0.5	0.3
11	Pregnancy, childbirth and the puerperium (O00-O99)	73	0.4	0.3
12	Chronic lower respiratory diseases (J40-J47)	64	0.4	0.3
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	54	0.3	0.2
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	30	0.2	0.1
15	Pneumonitis due to solids and liquids (J69)	28	0.2	0.1
15	Complications of medical and surgical care (Y40-Y84,Y88)	28	0.2	0.1
...	All other causes (Residual)	2,004	11.8	7.9
Non-Hispanic, white, both sexes, 25-34 years				
...	All causes	26,754	100.0	109.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	11,182	41.8	45.6
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	4,581	17.1	18.7
3	Malignant neoplasms (C00-C97)	2,057	7.7	8.4
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,816	6.8	7.4
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	947	3.5	3.9
6	Diabetes mellitus (E10-E14)	358	1.3	1.5
7	Influenza and pneumonia (J09-J18)	310	1.2	1.3
8	Chronic liver disease and cirrhosis (K70,K73-K74)	289	1.1	1.2
9	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	281	1.1	1.1
10	Cerebrovascular diseases (I60-I69)	253	0.9	1.0
11	Septicemia (A40-A41)	186	0.7	0.8
12	Pregnancy, childbirth and the puerperium (O00-O99)	176	0.7	0.7
13	Chronic lower respiratory diseases (J40-J47)	142	0.5	0.6
14	Human immunodeficiency virus (HIV) disease (B20-B24)	114	0.4	0.5
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	103	0.4	0.4
...	All other causes (Residual)	3,959	14.8	16.1

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> , Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, both sexes, 35-44 years				
...	All causes	44,891	100.0	180.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	11,202	25.0	45.0
2	Malignant neoplasms (C00-C97)	7,463	16.6	30.0
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	6,579	14.7	26.4
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	5,364	11.9	21.6
5	Chronic liver disease and cirrhosis (K70,K73-K74)	1,591	3.5	6.4
6	Diabetes mellitus (E10-E14)	1,035	2.3	4.2
7	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	836	1.9	3.4
8	Cerebrovascular diseases (I60-I69)	824	1.8	3.3
9	Influenza and pneumonia (J09-J18)	521	1.2	2.1
10	Chronic lower respiratory diseases (J40-J47)	476	1.1	1.9
11	Septicemia (A40-A41)	451	1.0	1.8
12	Human immunodeficiency virus (HIV) disease (B20-B24)	425	0.9	1.7
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	300	0.7	1.2
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	296	0.7	1.2
15	Viral hepatitis (B15-B19)	180	0.4	0.7
...	All other causes (Residual)	7,348	16.4	29.5
Non-Hispanic, white, both sexes, 45-54 years				
...	All causes	127,836	100.0	414.4
1	Malignant neoplasms (C00-C97)	34,675	27.1	112.4
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	24,634	19.3	79.8
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	15,961	12.5	51.7
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	7,821	6.1	25.4
5	Chronic liver disease and cirrhosis (K70,K73-K74)	6,326	4.9	20.5
6	Chronic lower respiratory diseases (J40-J47)	3,698	2.9	12.0
7	Diabetes mellitus (E10-E14)	3,641	2.8	11.8
8	Cerebrovascular diseases (I60-I69)	3,086	2.4	10.0
9	Viral hepatitis (B15-B19)	1,575	1.2	5.1
10	Septicemia (A40-A41)	1,541	1.2	5.0
11	Influenza and pneumonia (J09-J18)	1,526	1.2	4.9
12	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	989	0.8	3.2
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	973	0.8	3.2
14	Human immunodeficiency virus (HIV) disease (B20-B24)	894	0.7	2.9
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	701	0.5	2.3
...	All other causes (Residual)	19,795	15.5	64.2

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> , Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, both sexes, 55-64 years				
...	All causes	238,484	100.0	837.8
1	Malignant neoplasms (C00-C97)	85,241	35.7	299.5
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	50,464	21.2	177.3
3	Chronic lower respiratory diseases (J40-J47)	12,904	5.4	45.3
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	11,607	4.9	40.8
5	Diabetes mellitus (E10-E14)	8,156	3.4	28.7
6	Chronic liver disease and cirrhosis (K70,K73-K74)	7,930	3.3	27.9
7	Cerebrovascular diseases (I60-I69)	6,975	2.9	24.5
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	5,892	2.5	20.7
9	Septicemia (A40-A41)	3,302	1.4	11.6
10	Influenza and pneumonia (J09-J18)	3,057	1.3	10.7
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2,771	1.2	9.7
12	Viral hepatitis (B15-B19)	2,311	1.0	8.1
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,570	0.7	5.5
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1,122	0.5	3.9
15	Aortic aneurysm and dissection (I71)	984	0.4	3.5
...	All other causes (Residual)	34,198	14.3	120.1
Non-Hispanic, white, both sexes, 65-74 years				
...	All causes	325,930	100.0	1,856.1
1	Malignant neoplasms (C00-C97)	116,814	35.8	665.2
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	68,657	21.1	391.0
3	Chronic lower respiratory diseases (J40-J47)	28,933	8.9	164.8
4	Cerebrovascular diseases (I60-I69)	12,619	3.9	71.9
5	Diabetes mellitus (E10-E14)	10,810	3.3	61.6
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	8,105	2.5	46.2
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	5,062	1.6	28.8
8	Influenza and pneumonia (J09-J18)	4,960	1.5	28.2
9	Septicemia (A40-A41)	4,818	1.5	27.4
10	Chronic liver disease and cirrhosis (K70,K73-K74)	4,503	1.4	25.6
11	Alzheimer's disease (G30)	3,629	1.1	20.7
12	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,881	0.9	16.4
13	Parkinson's disease (G20-G21)	2,457	0.8	14.0
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	2,408	0.7	13.7
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2,130	0.7	12.1
...	All other causes (Residual)	47,144	14.5	268.5

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> , Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, both sexes, 75-84 years				
...	All causes	519,160	100.0	4,871.3
1	Malignant neoplasms (C00-C97)	129,241	24.9	1,212.7
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	122,673	23.6	1,151.1
3	Chronic lower respiratory diseases (J40-J47)	44,596	8.6	418.4
4	Cerebrovascular diseases (I60-I69)	30,104	5.8	282.5
5	Alzheimer's disease (G30)	20,942	4.0	196.5
6	Diabetes mellitus (E10-E14)	13,983	2.7	131.2
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	12,214	2.4	114.6
8	Influenza and pneumonia (J09-J18)	11,023	2.1	103.4
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	10,163	2.0	95.4
10	Parkinson's disease (G20-G21)	8,919	1.7	83.7
11	Septicemia (A40-A41)	7,613	1.5	71.4
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	4,947	1.0	46.4
13	Pneumonitis due to solids and liquids (J69)	4,382	0.8	41.1
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	3,954	0.8	37.1
15	Chronic liver disease and cirrhosis (K70,K73-K74)	3,061	0.6	28.7
...	All other causes (Residual)	91,345	17.6	857.1
Non-Hispanic, white, both sexes, 85 years and over				
...	All causes	690,440	100.0	14,215.3
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	207,352	30.0	4,269.1
2	Malignant neoplasms (C00-C97)	82,940	12.0	1,707.6
3	Alzheimer's disease (G30)	49,103	7.1	1,011.0
4	Cerebrovascular diseases (I60-I69)	46,973	6.8	967.1
5	Chronic lower respiratory diseases (J40-J47)	36,158	5.2	744.4
6	Influenza and pneumonia (J09-J18)	21,618	3.1	445.1
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	17,521	2.5	360.7
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	14,066	2.0	289.6
9	Diabetes mellitus (E10-E14)	12,864	1.9	264.9
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	10,327	1.5	212.6
11	Parkinson's disease (G20-G21)	8,785	1.3	180.9
12	Septicemia (A40-A41)	8,538	1.2	175.8
13	Pneumonitis due to solids and liquids (J69)	8,069	1.2	166.1
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	4,404	0.6	90.7
15	Atherosclerosis (I70)	3,530	0.5	72.7
...	All other causes (Residual)	158,192	22.9	3,257.0

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> , Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, male, all ages				
...	All causes	989,835	100.0	1,004.1
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	248,787	25.1	252.4
2	Malignant neoplasms (C00-C97)	242,140	24.5	245.6
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	60,043	6.1	60.9
4	Chronic lower respiratory diseases (J40-J47)	59,223	6.0	60.1
5	Cerebrovascular diseases (I60-I69)	39,753	4.0	40.3
6	Diabetes mellitus (E10-E14)	27,060	2.7	27.4
7	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	25,858	2.6	26.2
8	Alzheimer's disease (G30)	22,482	2.3	22.8
9	Influenza and pneumonia (J09-J18)	20,316	2.1	20.6
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	16,898	1.7	17.1
11	Chronic liver disease and cirrhosis (K70,K73-K74)	15,764	1.6	16.0
12	Septicemia (A40-A41)	12,319	1.2	12.5
13	Parkinson's disease (G20-G21)	12,134	1.2	12.3
14	Pneumonitis due to solids and liquids (J69)	8,242	0.8	8.4
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	7,926	0.8	8.0
...	All other causes (Residual)	170,890	17.3	173.4
Non-Hispanic, white, male, 1-4 years				
...	All causes	1,177	100.0	27.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	454	38.6	10.4
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	124	10.5	2.8
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	93	7.9	2.1
4	Malignant neoplasms (C00-C97)	89	7.6	2.0
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	35	3.0	0.8
6	Influenza and pneumonia (J09-J18)	22	1.9	0.5
7	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	16	1.4	*
8	Chronic lower respiratory diseases (J40-J47)	15	1.3	*
9	Septicemia (A40-A41)	12	1.0	*
10	Certain conditions originating in the perinatal period (P00-P96)	9	0.8	*
11	Cerebrovascular diseases (I60-I69)	8	0.7	*
12	Anemias (D50-D64)	4	0.3	*
12	Acute bronchitis and bronchiolitis (J20-J21)	4	0.3	*
12	Complications of medical and surgical care (Y40-Y84,Y88)	4	0.3	*
15	Meningitis (G00,G03)	3	0.3	*
...	All other causes (Residual)	285	24.2	6.5

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> , Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, male, 5-14 years				
...	All causes	1,727	100.0	14.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	613	35.5	5.2
2	Malignant neoplasms (C00-C97)	274	15.9	2.3
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	121	7.0	1.0
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	94	5.4	0.8
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	72	4.2	0.6
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	48	2.8	0.4
7	Influenza and pneumonia (J09-J18)	35	2.0	0.3
8	Septicemia (A40-A41)	24	1.4	0.2
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	23	1.3	0.2
10	Cerebrovascular diseases (I60-I69)	21	1.2	0.2
11	Chronic lower respiratory diseases (J40-J47)	17	1.0	*
12	Pneumonitis due to solids and liquids (J69)	8	0.5	*
12	Certain conditions originating in the perinatal period (P00-P96)	8	0.5	*
14	Anemias (D50-D64)	6	0.3	*
15	Diabetes mellitus (E10-E14)	5	0.3	*
...	All other causes (Residual)	358	20.7	3.1
Non-Hispanic, white, male, 15-24 years				
...	All causes	12,222	100.0	94.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	6,249	51.1	48.5
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,817	23.0	21.9
3	Malignant neoplasms (C00-C97)	517	4.2	4.0
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	497	4.1	3.9
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	356	2.9	2.8
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	164	1.3	1.3
7	Influenza and pneumonia (J09-J18)	81	0.7	0.6
8	Diabetes mellitus (E10-E14)	67	0.5	0.5
9	Cerebrovascular diseases (I60-I69)	51	0.4	0.4
10	Septicemia (A40-A41)	48	0.4	0.4
11	Chronic lower respiratory diseases (J40-J47)	47	0.4	0.4
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	27	0.2	0.2
13	Pneumonitis due to solids and liquids (J69)	20	0.2	0.2
14	Legal intervention (Y35,Y89.0)	18	0.1	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	17	0.1	*
...	All other causes (Residual)	1,246	10.2	9.7

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> , Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, male, 25-34 years				
...	All causes	18,291	100.0	147.9
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	8,117	44.4	65.6
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	3,642	19.9	29.5
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,241	6.8	10.0
4	Malignant neoplasms (C00-C97)	1,038	5.7	8.4
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	662	3.6	5.4
6	Diabetes mellitus (E10-E14)	215	1.2	1.7
7	Influenza and pneumonia (J09-J18)	174	1.0	1.4
8	Chronic liver disease and cirrhosis (K70,K73-K74)	172	0.9	1.4
9	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	160	0.9	1.3
10	Cerebrovascular diseases (I60-I69)	128	0.7	1.0
11	Septicemia (A40-A41)	109	0.6	0.9
12	Human immunodeficiency virus (HIV) disease (B20-B24)	76	0.4	0.6
12	Chronic lower respiratory diseases (J40-J47)	76	0.4	0.6
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	63	0.3	0.5
15	Legal intervention (Y35,Y89.0)	62	0.3	0.5
...	All other causes (Residual)	2,356	12.9	19.1
Non-Hispanic, white, male, 35-44 years				
...	All causes	27,963	100.0	223.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	7,497	26.8	60.0
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	4,664	16.7	37.3
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	4,067	14.5	32.6
4	Malignant neoplasms (C00-C97)	3,401	12.2	27.2
5	Chronic liver disease and cirrhosis (K70,K73-K74)	980	3.5	7.8
6	Diabetes mellitus (E10-E14)	640	2.3	5.1
7	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	561	2.0	4.5
8	Cerebrovascular diseases (I60-I69)	422	1.5	3.4
9	Human immunodeficiency virus (HIV) disease (B20-B24)	326	1.2	2.6
10	Influenza and pneumonia (J09-J18)	271	1.0	2.2
11	Chronic lower respiratory diseases (J40-J47)	224	0.8	1.8
12	Septicemia (A40-A41)	217	0.8	1.7
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	183	0.7	1.5
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	166	0.6	1.3
15	Aortic aneurysm and dissection (I71)	129	0.5	1.0
...	All other causes (Residual)	4,215	15.1	33.7

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> , Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, male, 45-54 years				
...	All causes	78,465	100.0	512.3
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	17,931	22.9	117.1
2	Malignant neoplasms (C00-C97)	17,521	22.3	114.4
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	10,487	13.4	68.5
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	5,839	7.4	38.1
5	Chronic liver disease and cirrhosis (K70,K73-K74)	4,108	5.2	26.8
6	Diabetes mellitus (E10-E14)	2,349	3.0	15.3
7	Chronic lower respiratory diseases (J40-J47)	1,733	2.2	11.3
8	Cerebrovascular diseases (I60-I69)	1,717	2.2	11.2
9	Viral hepatitis (B15-B19)	1,105	1.4	7.2
10	Influenza and pneumonia (J09-J18)	876	1.1	5.7
11	Septicemia (A40-A41)	830	1.1	5.4
12	Human immunodeficiency virus (HIV) disease (B20-B24)	748	1.0	4.9
13	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	661	0.8	4.3
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	580	0.7	3.8
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	461	0.6	3.0
...	All other causes (Residual)	11,519	14.7	75.2
Non-Hispanic, white, male, 55-64 years				
...	All causes	146,023	100.0	1,050.0
1	Malignant neoplasms (C00-C97)	47,535	32.6	341.8
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	35,881	24.6	258.0
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	7,913	5.4	56.9
4	Chronic lower respiratory diseases (J40-J47)	6,717	4.6	48.3
5	Chronic liver disease and cirrhosis (K70,K73-K74)	5,582	3.8	40.1
6	Diabetes mellitus (E10-E14)	5,011	3.4	36.0
7	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	4,523	3.1	32.5
8	Cerebrovascular diseases (I60-I69)	3,934	2.7	28.3
9	Influenza and pneumonia (J09-J18)	1,782	1.2	12.8
10	Septicemia (A40-A41)	1,733	1.2	12.5
11	Viral hepatitis (B15-B19)	1,725	1.2	12.4
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,588	1.1	11.4
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,016	0.7	7.3
14	Aortic aneurysm and dissection (I71)	752	0.5	5.4
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	649	0.4	4.7
...	All other causes (Residual)	19,682	13.5	141.5

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> , Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, male, 65-74 years				
...	All causes	185,021	100.0	2,227.4
1	Malignant neoplasms (C00-C97)	64,437	34.8	775.7
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	44,104	23.8	530.9
3	Chronic lower respiratory diseases (J40-J47)	14,508	7.8	174.7
4	Cerebrovascular diseases (I60-I69)	6,625	3.6	79.8
5	Diabetes mellitus (E10-E14)	6,361	3.4	76.6
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	5,081	2.7	61.2
7	Chronic liver disease and cirrhosis (K70,K73-K74)	2,802	1.5	33.7
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2,776	1.5	33.4
9	Influenza and pneumonia (J09-J18)	2,759	1.5	33.2
10	Septicemia (A40-A41)	2,489	1.3	30.0
11	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,317	1.3	27.9
12	Parkinson's disease (G20-G21)	1,638	0.9	19.7
13	Alzheimer's disease (G30)	1,566	0.8	18.9
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1,276	0.7	15.4
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,260	0.7	15.2
...	All other causes (Residual)	25,022	13.5	301.2
Non-Hispanic, white, male, 75-84 years				
...	All causes	262,132	100.0	5,727.3
1	Malignant neoplasms (C00-C97)	69,027	26.3	1,508.2
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	66,499	25.4	1,452.9
3	Chronic lower respiratory diseases (J40-J47)	21,167	8.1	462.5
4	Cerebrovascular diseases (I60-I69)	13,041	5.0	284.9
5	Alzheimer's disease (G30)	7,922	3.0	173.1
6	Diabetes mellitus (E10-E14)	7,224	2.8	157.8
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	6,561	2.5	143.4
8	Influenza and pneumonia (J09-J18)	5,622	2.1	122.8
9	Parkinson's disease (G20-G21)	5,518	2.1	120.6
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	5,446	2.1	119.0
11	Septicemia (A40-A41)	3,688	1.4	80.6
12	Pneumonitis due to solids and liquids (J69)	2,474	0.9	54.1
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2,196	0.8	48.0
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	2,147	0.8	46.9
15	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,739	0.7	38.0
...	All other causes (Residual)	41,861	16.0	914.6

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> , Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, male, 85 years and over				
...	All causes	250,688	100.0	15,568.2
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	77,953	31.1	4,841.0
2	Malignant neoplasms (C00-C97)	38,278	15.3	2,377.1
3	Chronic lower respiratory diseases (J40-J47)	14,708	5.9	913.4
4	Cerebrovascular diseases (I60-I69)	13,764	5.5	854.8
5	Alzheimer's disease (G30)	12,651	5.0	785.7
6	Influenza and pneumonia (J09-J18)	8,639	3.4	536.5
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	6,739	2.7	418.5
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	6,238	2.5	387.4
9	Diabetes mellitus (E10-E14)	5,186	2.1	322.1
10	Parkinson's disease (G20-G21)	4,678	1.9	290.5
11	Pneumonitis due to solids and liquids (J69)	3,933	1.6	244.2
12	Septicemia (A40-A41)	3,136	1.3	194.8
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	2,909	1.2	180.7
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1,985	0.8	123.3
15	Aortic aneurysm and dissection (I71)	1,096	0.4	68.1
...	All other causes (Residual)	48,795	19.5	3,030.3
Non-Hispanic, white, female, all ages				
...	All causes	1,016,484	100.0	998.1
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	234,192	23.0	230.0
2	Malignant neoplasms (C00-C97)	217,886	21.4	213.9
3	Chronic lower respiratory diseases (J40-J47)	67,812	6.7	66.6
4	Cerebrovascular diseases (I60-I69)	61,290	6.0	60.2
5	Alzheimer's disease (G30)	51,966	5.1	51.0
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	38,562	3.8	37.9
7	Diabetes mellitus (E10-E14)	23,904	2.4	23.5
8	Influenza and pneumonia (J09-J18)	23,034	2.3	22.6
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	16,594	1.6	16.3
10	Septicemia (A40-A41)	14,350	1.4	14.1
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	12,227	1.2	12.0
12	Chronic liver disease and cirrhosis (K70,K73-K74)	8,988	0.9	8.8
13	Parkinson's disease (G20-G21)	8,491	0.8	8.3
14	Pneumonitis due to solids and liquids (J69)	7,271	0.7	7.1
15	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	7,173	0.7	7.0
...	All other causes (Residual)	222,744	21.9	218.7

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> , Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, female, 1-4 years				
...	All causes	878	100.0	21.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	270	30.8	6.5
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	112	12.8	2.7
3	Malignant neoplasms (C00-C97)	94	10.7	2.3
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	57	6.5	1.4
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	39	4.4	0.9
6	Influenza and pneumonia (J09-J18)	23	2.6	0.6
7	Septicemia (A40-A41)	18	2.1	*
8	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	8	0.9	*
8	Certain conditions originating in the perinatal period (P00-P96)	8	0.9	*
10	Cerebrovascular diseases (I60-I69)	6	0.7	*
11	Acute bronchitis and bronchiolitis (J20-J21)	4	0.5	*
11	Chronic lower respiratory diseases (J40-J47)	4	0.5	*
13	Diabetes mellitus (E10-E14)	3	0.3	*
13	Complications of medical and surgical care (Y40-Y84,Y88)	3	0.3	*
15	Anemias (D50-D64)	2	0.2	*
15	Meningitis (G00,G03)	2	0.2	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2	0.2	*
...	All other causes (Residual)	223	25.4	5.4
Non-Hispanic, white, female, 5-14 years				
...	All causes	1,181	100.0	10.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	360	30.5	3.2
2	Malignant neoplasms (C00-C97)	207	17.5	1.9
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	74	6.3	0.7
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	53	4.5	0.5
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	49	4.1	0.4
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	47	4.0	0.4
7	Influenza and pneumonia (J09-J18)	26	2.2	0.2
8	Septicemia (A40-A41)	22	1.9	0.2
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	20	1.7	0.2
10	Cerebrovascular diseases (I60-I69)	17	1.4	*
11	Chronic lower respiratory diseases (J40-J47)	12	1.0	*
12	Diabetes mellitus (E10-E14)	10	0.8	*
13	Certain conditions originating in the perinatal period (P00-P96)	8	0.7	*
14	Meningitis (G00,G03)	4	0.3	*
15	Anemias (D50-D64)	3	0.3	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3	0.3	*
15	Complications of medical and surgical care (Y40-Y84,Y88)	3	0.3	*
...	All other causes (Residual)	263	22.3	2.4

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> , Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, female, 15-24 years				
...	All causes	4,739	100.0	38.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,279	48.1	18.5
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	596	12.6	4.8
3	Malignant neoplasms (C00-C97)	374	7.9	3.0
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	204	4.3	1.7
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	137	2.9	1.1
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	88	1.9	0.7
7	Pregnancy, childbirth and the puerperium (O00-O99)	73	1.5	0.6
8	Influenza and pneumonia (J09-J18)	57	1.2	0.5
9	Cerebrovascular diseases (I60-I69)	39	0.8	0.3
10	Diabetes mellitus (E10-E14)	30	0.6	0.2
11	Septicemia (A40-A41)	29	0.6	0.2
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	27	0.6	0.2
13	Chronic lower respiratory diseases (J40-J47)	17	0.4	*
13	Complications of medical and surgical care (Y40-Y84,Y88)	17	0.4	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	13	0.3	*
...	All other causes (Residual)	759	16.0	6.1
Non-Hispanic, white, female, 25-34 years				
...	All causes	8,463	100.0	69.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,065	36.2	25.2
2	Malignant neoplasms (C00-C97)	1,019	12.0	8.4
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	939	11.1	7.7
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	575	6.8	4.7
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	285	3.4	2.3
6	Pregnancy, childbirth and the puerperium (O00-O99)	176	2.1	1.4
7	Diabetes mellitus (E10-E14)	143	1.7	1.2
8	Influenza and pneumonia (J09-J18)	136	1.6	1.1
9	Cerebrovascular diseases (I60-I69)	125	1.5	1.0
10	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	121	1.4	1.0
11	Chronic liver disease and cirrhosis (K70,K73-K74)	117	1.4	1.0
12	Septicemia (A40-A41)	77	0.9	0.6
13	Chronic lower respiratory diseases (J40-J47)	66	0.8	0.5
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	40	0.5	0.3
15	Human immunodeficiency virus (HIV) disease (B20-B24)	38	0.4	0.3
...	All other causes (Residual)	1,541	18.2	12.7

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> , Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, female, 35-44 years				
...	All causes	16,928	100.0	136.6
1	Malignant neoplasms (C00-C97)	4,062	24.0	32.8
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,705	21.9	29.9
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,915	11.3	15.5
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,297	7.7	10.5
5	Chronic liver disease and cirrhosis (K70,K73-K74)	611	3.6	4.9
6	Cerebrovascular diseases (I60-I69)	402	2.4	3.2
7	Diabetes mellitus (E10-E14)	395	2.3	3.2
8	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	275	1.6	2.2
9	Chronic lower respiratory diseases (J40-J47)	252	1.5	2.0
10	Influenza and pneumonia (J09-J18)	250	1.5	2.0
11	Septicemia (A40-A41)	234	1.4	1.9
12	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	130	0.8	1.0
13	Pregnancy, childbirth and the puerperium (O00-O99)	123	0.7	1.0
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	117	0.7	0.9
15	Human immunodeficiency virus (HIV) disease (B20-B24)	99	0.6	0.8
...	All other causes (Residual)	3,061	18.1	24.7
Non-Hispanic, white, female, 45-54 years				
...	All causes	49,371	100.0	317.8
1	Malignant neoplasms (C00-C97)	17,154	34.7	110.4
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	6,703	13.6	43.1
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	5,474	11.1	35.2
4	Chronic liver disease and cirrhosis (K70,K73-K74)	2,218	4.5	14.3
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,982	4.0	12.8
6	Chronic lower respiratory diseases (J40-J47)	1,965	4.0	12.6
7	Cerebrovascular diseases (I60-I69)	1,369	2.8	8.8
8	Diabetes mellitus (E10-E14)	1,292	2.6	8.3
9	Septicemia (A40-A41)	711	1.4	4.6
10	Influenza and pneumonia (J09-J18)	650	1.3	4.2
11	Viral hepatitis (B15-B19)	470	1.0	3.0
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	393	0.8	2.5
13	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	328	0.7	2.1
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	272	0.6	1.8
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	240	0.5	1.5
...	All other causes (Residual)	8,150	16.5	52.5

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> , Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, female, 55-64 years				
...	All causes	92,461	100.0	635.2
1	Malignant neoplasms (C00-C97)	37,706	40.8	259.0
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	14,583	15.8	100.2
3	Chronic lower respiratory diseases (J40-J47)	6,187	6.7	42.5
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,694	4.0	25.4
5	Diabetes mellitus (E10-E14)	3,145	3.4	21.6
6	Cerebrovascular diseases (I60-I69)	3,041	3.3	20.9
7	Chronic liver disease and cirrhosis (K70,K73-K74)	2,348	2.5	16.1
8	Septicemia (A40-A41)	1,569	1.7	10.8
9	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,369	1.5	9.4
10	Influenza and pneumonia (J09-J18)	1,275	1.4	8.8
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,183	1.3	8.1
12	Viral hepatitis (B15-B19)	586	0.6	4.0
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	554	0.6	3.8
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	473	0.5	3.2
15	Alzheimer's disease (G30)	379	0.4	2.6
...	All other causes (Residual)	14,369	15.5	98.7
Non-Hispanic, white, female, 65-74 years				
...	All causes	140,909	100.0	1,522.8
1	Malignant neoplasms (C00-C97)	52,377	37.2	566.0
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	24,553	17.4	265.3
3	Chronic lower respiratory diseases (J40-J47)	14,425	10.2	155.9
4	Cerebrovascular diseases (I60-I69)	5,994	4.3	64.8
5	Diabetes mellitus (E10-E14)	4,449	3.2	48.1
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,024	2.1	32.7
7	Septicemia (A40-A41)	2,329	1.7	25.2
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2,286	1.6	24.7
9	Influenza and pneumonia (J09-J18)	2,201	1.6	23.8
10	Alzheimer's disease (G30)	2,063	1.5	22.3
11	Chronic liver disease and cirrhosis (K70,K73-K74)	1,701	1.2	18.4
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,148	0.8	12.4
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	854	0.6	9.2
14	Parkinson's disease (G20-G21)	819	0.6	8.9
15	Pneumonitis due to solids and liquids (J69)	633	0.4	6.8
...	All other causes (Residual)	22,053	15.7	238.3

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> , Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, female, 75-84 years				
...	All causes	257,028	100.0	4,227.0
1	Malignant neoplasms (C00-C97)	60,214	23.4	990.3
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	56,174	21.9	923.8
3	Chronic lower respiratory diseases (J40-J47)	23,429	9.1	385.3
4	Cerebrovascular diseases (I60-I69)	17,063	6.6	280.6
5	Alzheimer's disease (G30)	13,020	5.1	214.1
6	Diabetes mellitus (E10-E14)	6,759	2.6	111.2
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	5,653	2.2	93.0
8	Influenza and pneumonia (J09-J18)	5,401	2.1	88.8
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	4,717	1.8	77.6
10	Septicemia (A40-A41)	3,925	1.5	64.5
11	Parkinson's disease (G20-G21)	3,401	1.3	55.9
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	2,800	1.1	46.0
13	Pneumonitis due to solids and liquids (J69)	1,908	0.7	31.4
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1,758	0.7	28.9
15	Chronic liver disease and cirrhosis (K70,K73-K74)	1,416	0.6	23.3
...	All other causes (Residual)	49,390	19.2	812.3
Non-Hispanic, white, female, 85 years and over				
...	All causes	439,752	100.0	13,544.3
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	129,399	29.4	3,985.5
2	Malignant neoplasms (C00-C97)	44,662	10.2	1,375.6
3	Alzheimer's disease (G30)	36,452	8.3	1,122.7
4	Cerebrovascular diseases (I60-I69)	33,209	7.6	1,022.8
5	Chronic lower respiratory diseases (J40-J47)	21,450	4.9	660.7
6	Influenza and pneumonia (J09-J18)	12,979	3.0	399.8
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	10,782	2.5	332.1
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	7,828	1.8	241.1
9	Diabetes mellitus (E10-E14)	7,678	1.7	236.5
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	7,418	1.7	228.5
11	Septicemia (A40-A41)	5,402	1.2	166.4
12	Pneumonitis due to solids and liquids (J69)	4,136	0.9	127.4
13	Parkinson's disease (G20-G21)	4,107	0.9	126.5
14	Atherosclerosis (I70)	2,447	0.6	75.4
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2,419	0.6	74.5
...	All other causes (Residual)	109,384	24.9	3,369.0

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> , Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, both sexes, all ages				
...	All causes	286,797	100.0	718.0
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	67,595	23.6	169.2
2	Malignant neoplasms (C00-C97)	66,158	23.1	165.6
3	Cerebrovascular diseases (I60-I69)	15,722	5.5	39.4
4	Diabetes mellitus (E10-E14)	12,771	4.5	32.0
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	12,299	4.3	30.8
6	Chronic lower respiratory diseases (J40-J47)	8,994	3.1	22.5
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	8,044	2.8	20.1
8	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	7,764	2.7	19.4
9	Septicemia (A40-A41)	5,963	2.1	14.9
10	Alzheimer's disease (G30)	5,446	1.9	13.6
11	Influenza and pneumonia (J09-J18)	5,098	1.8	12.8
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	4,965	1.7	12.4
13	Human immunodeficiency virus (HIV) disease (B20-B24)	4,138	1.4	10.4
14	Certain conditions originating in the perinatal period (P00-P96)	3,890	1.4	9.7
15	Chronic liver disease and cirrhosis (K70,K73-K74)	2,783	1.0	7.0
...	All other causes (Residual)	55,167	19.2	138.1
Non-Hispanic, black, both sexes, 1-4 years				
...	All causes	1,011	100.0	40.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	286	28.3	11.5
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	154	15.2	6.2
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	105	10.4	4.2
4	Malignant neoplasms (C00-C97)	53	5.2	2.1
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	52	5.1	2.1
6	Influenza and pneumonia (J09-J18)	32	3.2	1.3
7	Chronic lower respiratory diseases (J40-J47)	24	2.4	1.0
8	Cerebrovascular diseases (I60-I69)	17	1.7	*
9	Certain conditions originating in the perinatal period (P00-P96)	12	1.2	*
10	Septicemia (A40-A41)	11	1.1	*
11	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	10	1.0	*
12	Complications of medical and surgical care (Y40-Y84,Y88)	8	0.8	*
13	Anemias (D50-D64)	6	0.6	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3	0.3	*
15	Meningitis (G00,G03)	2	0.2	*
15	Acute bronchitis and bronchiolitis (J20-J21)	2	0.2	*
15	Pneumonitis due to solids and liquids (J69)	2	0.2	*
...	All other causes (Residual)	232	22.9	9.4

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> , Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, both sexes, 5-14 years				
...	All causes	1,150	100.0	18.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	313	27.2	5.1
2	Malignant neoplasms (C00-C97)	127	11.0	2.1
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	89	7.7	1.4
4	Chronic lower respiratory diseases (J40-J47)	81	7.0	1.3
5	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	78	6.8	1.3
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	49	4.3	0.8
7	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	43	3.7	0.7
8	Influenza and pneumonia (J09-J18)	25	2.2	0.4
9	Anemias (D50-D64)	22	1.9	0.4
10	Cerebrovascular diseases (I60-I69)	20	1.7	0.3
11	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	19	1.7	*
12	Septicemia (A40-A41)	11	1.0	*
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	9	0.8	*
14	Pneumonitis due to solids and liquids (J69)	5	0.4	*
15	Diabetes mellitus (E10-E14)	4	0.3	*
15	Meningitis (G00,G03)	4	0.3	*
15	Certain conditions originating in the perinatal period (P00-P96)	4	0.3	*
15	Complications of medical and surgical care (Y40-Y84,Y88)	4	0.3	*
...	All other causes (Residual)	243	21.1	3.9
Non-Hispanic, black, both sexes, 15-24 years				
...	All causes	6,576	100.0	97.6
1	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	2,797	42.5	41.5
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,357	20.6	20.1
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	487	7.4	7.2
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	303	4.6	4.5
5	Malignant neoplasms (C00-C97)	257	3.9	3.8
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	102	1.6	1.5
7	Human immunodeficiency virus (HIV) disease (B20-B24)	100	1.5	1.5
8	Chronic lower respiratory diseases (J40-J47)	81	1.2	1.2
9	Anemias (D50-D64)	71	1.1	1.1
10	Pregnancy, childbirth and the puerperium (O00-O99)	62	0.9	0.9
11	Diabetes mellitus (E10-E14)	47	0.7	0.7
12	Legal intervention (Y35,Y89.0)	46	0.7	0.7
13	Influenza and pneumonia (J09-J18)	44	0.7	0.7
14	Septicemia (A40-A41)	35	0.5	0.5
15	Cerebrovascular diseases (I60-I69)	31	0.5	0.5
...	All other causes (Residual)	756	11.5	11.2

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> , Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, both sexes, 25-34 years				
...	All causes	8,957	100.0	161.5
1	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	2,302	25.7	41.5
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,632	18.2	29.4
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	941	10.5	17.0
4	Malignant neoplasms (C00-C97)	640	7.1	11.5
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	548	6.1	9.9
6	Human immunodeficiency virus (HIV) disease (B20-B24)	412	4.6	7.4
7	Diabetes mellitus (E10-E14)	226	2.5	4.1
8	Cerebrovascular diseases (I60-I69)	135	1.5	2.4
9	Anemias (D50-D64)	133	1.5	2.4
10	Pregnancy, childbirth and the puerperium (O00-O99)	124	1.4	2.2
11	Influenza and pneumonia (J09-J18)	106	1.2	1.9
12	Septicemia (A40-A41)	97	1.1	1.7
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	94	1.0	1.7
14	Chronic lower respiratory diseases (J40-J47)	90	1.0	1.6
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	59	0.7	1.1
...	All other causes (Residual)	1,418	15.8	25.6
Non-Hispanic, black, both sexes, 35-44 years				
...	All causes	13,611	100.0	261.5
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,681	19.7	51.5
2	Malignant neoplasms (C00-C97)	2,089	15.3	40.1
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,659	12.2	31.9
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,123	8.3	21.6
5	Human immunodeficiency virus (HIV) disease (B20-B24)	889	6.5	17.1
6	Cerebrovascular diseases (I60-I69)	509	3.7	9.8
7	Diabetes mellitus (E10-E14)	487	3.6	9.4
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	417	3.1	8.0
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	270	2.0	5.2
10	Septicemia (A40-A41)	215	1.6	4.1
11	Chronic lower respiratory diseases (J40-J47)	191	1.4	3.7
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	186	1.4	3.6
13	Influenza and pneumonia (J09-J18)	179	1.3	3.4
13	Chronic liver disease and cirrhosis (K70,K73-K74)	179	1.3	3.4
15	Anemias (D50-D64)	110	0.8	2.1
...	All other causes (Residual)	2,427	17.8	46.6

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> , Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, both sexes, 45-54 years				
...	All causes	33,151	100.0	600.2
1	Malignant neoplasms (C00-C97)	8,310	25.1	150.5
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	7,749	23.4	140.3
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,358	7.1	42.7
4	Cerebrovascular diseases (I60-I69)	1,631	4.9	29.5
5	Diabetes mellitus (E10-E14)	1,436	4.3	26.0
6	Human immunodeficiency virus (HIV) disease (B20-B24)	1,429	4.3	25.9
7	Chronic liver disease and cirrhosis (K70,K73-K74)	785	2.4	14.2
8	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	727	2.2	13.2
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	725	2.2	13.1
10	Chronic lower respiratory diseases (J40-J47)	693	2.1	12.5
11	Septicemia (A40-A41)	623	1.9	11.3
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	495	1.5	9.0
13	Influenza and pneumonia (J09-J18)	396	1.2	7.2
14	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	349	1.1	6.3
15	Viral hepatitis (B15-B19)	275	0.8	5.0
...	All other causes (Residual)	5,170	15.6	93.6
Non-Hispanic, black, both sexes, 55-64 years				
...	All causes	53,296	100.0	1,292.2
1	Malignant neoplasms (C00-C97)	16,807	31.5	407.5
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	13,022	24.4	315.7
3	Diabetes mellitus (E10-E14)	2,745	5.2	66.6
4	Cerebrovascular diseases (I60-I69)	2,690	5.0	65.2
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,001	3.8	48.5
6	Chronic lower respiratory diseases (J40-J47)	1,469	2.8	35.6
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,417	2.7	34.4
8	Chronic liver disease and cirrhosis (K70,K73-K74)	1,149	2.2	27.9
9	Septicemia (A40-A41)	1,119	2.1	27.1
10	Human immunodeficiency virus (HIV) disease (B20-B24)	979	1.8	23.7
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	844	1.6	20.5
12	Influenza and pneumonia (J09-J18)	700	1.3	17.0
13	Viral hepatitis (B15-B19)	597	1.1	14.5
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	331	0.6	8.0
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	199	0.4	4.8
...	All other causes (Residual)	7,227	13.6	175.2

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> , Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, both sexes, 65-74 years				
...	All causes	52,744	100.0	2,524.5
1	Malignant neoplasms (C00-C97)	16,808	31.9	804.5
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	13,195	25.0	631.6
3	Cerebrovascular diseases (I60-I69)	3,004	5.7	143.8
4	Diabetes mellitus (E10-E14)	2,983	5.7	142.8
5	Chronic lower respiratory diseases (J40-J47)	2,206	4.2	105.6
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,745	3.3	83.5
7	Septicemia (A40-A41)	1,239	2.3	59.3
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	943	1.8	45.1
9	Essential hypertension and hypertensive renal disease (I10,I12,I15)	905	1.7	43.3
10	Influenza and pneumonia (J09-J18)	843	1.6	40.3
11	Alzheimer's disease (G30)	405	0.8	19.4
11	Chronic liver disease and cirrhosis (K70,K73-K74)	405	0.8	19.4
13	Pneumonitis due to solids and liquids (J69)	253	0.5	12.1
14	Human immunodeficiency virus (HIV) disease (B20-B24)	251	0.5	12.0
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	223	0.4	10.7
...	All other causes (Residual)	7,336	13.9	351.1
Non-Hispanic, black, both sexes, 75-84 years				
...	All causes	57,367	100.0	5,331.0
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	14,842	25.9	1,379.3
2	Malignant neoplasms (C00-C97)	13,981	24.4	1,299.2
3	Cerebrovascular diseases (I60-I69)	3,939	6.9	366.0
4	Diabetes mellitus (E10-E14)	2,910	5.1	270.4
5	Chronic lower respiratory diseases (J40-J47)	2,491	4.3	231.5
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2,072	3.6	192.5
7	Alzheimer's disease (G30)	1,710	3.0	158.9
8	Septicemia (A40-A41)	1,390	2.4	129.2
9	Influenza and pneumonia (J09-J18)	1,219	2.1	113.3
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,103	1.9	102.5
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	760	1.3	70.6
12	Pneumonitis due to solids and liquids (J69)	406	0.7	37.7
13	Parkinson's disease (G20-G21)	382	0.7	35.5
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	330	0.6	30.7
15	Enterocolitis due to <i>Clostridium difficile</i> (A04.7)	197	0.3	18.3
...	All other causes (Residual)	9,635	16.8	895.4

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> , Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, both sexes, 85 years and over				
...	All causes	51,933	100.0	13,131.3
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	14,661	28.2	3,707.0
2	Malignant neoplasms (C00-C97)	7,070	13.6	1,787.7
3	Cerebrovascular diseases (I60-I69)	3,701	7.1	935.8
4	Alzheimer's disease (G30)	3,254	6.3	822.8
5	Diabetes mellitus (E10-E14)	1,933	3.7	488.8
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,658	3.2	419.2
7	Chronic lower respiratory diseases (J40-J47)	1,656	3.2	418.7
8	Influenza and pneumonia (J09-J18)	1,474	2.8	372.7
9	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,373	2.6	347.2
10	Septicemia (A40-A41)	1,162	2.2	293.8
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	623	1.2	157.5
12	Pneumonitis due to solids and liquids (J69)	505	1.0	127.7
13	Parkinson's disease (G20-G21)	269	0.5	68.0
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	238	0.5	60.2
14	Atherosclerosis (I70)	238	0.5	60.2
...	All other causes (Residual)	12,118	23.3	3,064.0
Non-Hispanic, black, male, all ages				
...	All causes	145,052	100.0	760.4
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	34,476	23.8	180.7
2	Malignant neoplasms (C00-C97)	33,628	23.2	176.3
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	8,219	5.7	43.1
4	Cerebrovascular diseases (I60-I69)	6,979	4.8	36.6
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	6,658	4.6	34.9
6	Diabetes mellitus (E10-E14)	5,986	4.1	31.4
7	Chronic lower respiratory diseases (J40-J47)	4,648	3.2	24.4
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3,734	2.6	19.6
9	Human immunodeficiency virus (HIV) disease (B20-B24)	2,693	1.9	14.1
10	Septicemia (A40-A41)	2,666	1.8	14.0
11	Influenza and pneumonia (J09-J18)	2,386	1.6	12.5
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	2,180	1.5	11.4
13	Certain conditions originating in the perinatal period (P00-P96)	2,171	1.5	11.4
14	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,790	1.2	9.4
15	Chronic liver disease and cirrhosis (K70,K73-K74)	1,750	1.2	9.2
...	All other causes (Residual)	25,088	17.3	131.5

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> , Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, male, 1-4 years				
...	All causes	575	100.0	45.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	167	29.0	13.3
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	84	14.6	6.7
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	57	9.9	4.5
4	Malignant neoplasms (C00-C97)	30	5.2	2.4
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	28	4.9	2.2
6	Influenza and pneumonia (J09-J18)	18	3.1	*
7	Chronic lower respiratory diseases (J40-J47)	16	2.8	*
8	Cerebrovascular diseases (I60-I69)	14	2.4	*
9	Certain conditions originating in the perinatal period (P00-P96)	8	1.4	*
10	Septicemia (A40-A41)	6	1.0	*
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	6	1.0	*
10	Complications of medical and surgical care (Y40-Y84,Y88)	6	1.0	*
13	Anemias (D50-D64)	3	0.5	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2	0.3	*
15	Meningitis (G00,G03)	1	0.2	*
15	Acute bronchitis and bronchiolitis (J20-J21)	1	0.2	*
15	Pneumonitis due to solids and liquids (J69)	1	0.2	*
...	All other causes (Residual)	127	22.1	10.1
Non-Hispanic, black, male, 5-14 years				
...	All causes	684	100.0	21.9
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	196	28.7	6.3
2	Malignant neoplasms (C00-C97)	73	10.7	2.3
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	61	8.9	1.9
4	Chronic lower respiratory diseases (J40-J47)	57	8.3	1.8
5	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	43	6.3	1.4
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	32	4.7	1.0
7	Diseases of heart (I00-I09,I11,I13,I20-I51)	25	3.7	0.8
8	Influenza and pneumonia (J09-J18)	15	2.2	*
9	Anemias (D50-D64)	14	2.0	*
10	Cerebrovascular diseases (I60-I69)	11	1.6	*
11	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	10	1.5	*
12	Pneumonitis due to solids and liquids (J69)	5	0.7	*
13	Septicemia (A40-A41)	4	0.6	*
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	4	0.6	*
15	Diabetes mellitus (E10-E14)	3	0.4	*
15	Certain conditions originating in the perinatal period (P00-P96)	3	0.4	*
15	Complications of medical and surgical care (Y40-Y84,Y88)	3	0.4	*
...	All other causes (Residual)	125	18.3	4.0

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> , Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, male, 15-24 years				
...	All causes	5,040	100.0	148.6
1	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	2,508	49.8	73.9
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,012	20.1	29.8
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	408	8.1	12.0
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	197	3.9	5.8
5	Malignant neoplasms (C00-C97)	151	3.0	4.5
6	Human immunodeficiency virus (HIV) disease (B20-B24)	62	1.2	1.8
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	62	1.2	1.8
8	Chronic lower respiratory diseases (J40-J47)	54	1.1	1.6
9	Legal intervention (Y35,Y89.0)	45	0.9	1.3
10	Anemias (D50-D64)	43	0.9	1.3
11	Diabetes mellitus (E10-E14)	23	0.5	0.7
12	Influenza and pneumonia (J09-J18)	20	0.4	0.6
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	17	0.3	*
14	Cerebrovascular diseases (I60-I69)	16	0.3	*
15	Septicemia (A40-A41)	12	0.2	*
...	All other causes (Residual)	410	8.1	12.1
Non-Hispanic, black, male, 25-34 years				
...	All causes	6,018	100.0	226.7
1	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	2,061	34.2	77.6
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,173	19.5	44.2
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	567	9.4	21.4
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	446	7.4	16.8
5	Malignant neoplasms (C00-C97)	260	4.3	9.8
6	Human immunodeficiency virus (HIV) disease (B20-B24)	240	4.0	9.0
7	Diabetes mellitus (E10-E14)	112	1.9	4.2
8	Anemias (D50-D64)	75	1.2	2.8
9	Cerebrovascular diseases (I60-I69)	67	1.1	2.5
10	Chronic lower respiratory diseases (J40-J47)	57	0.9	2.1
11	Influenza and pneumonia (J09-J18)	55	0.9	2.1
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	52	0.9	2.0
13	Septicemia (A40-A41)	40	0.7	1.5
14	Legal intervention (Y35,Y89.0)	31	0.5	1.2
15	Chronic liver disease and cirrhosis (K70,K73-K74)	28	0.5	1.1
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	28	0.5	1.1
...	All other causes (Residual)	726	12.1	27.4

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> , Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, male, 35-44 years				
...	All causes	7,869	100.0	321.1
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,713	21.8	69.9
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,177	15.0	48.0
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	945	12.0	38.6
4	Malignant neoplasms (C00-C97)	747	9.5	30.5
5	Human immunodeficiency virus (HIV) disease (B20-B24)	502	6.4	20.5
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	336	4.3	13.7
7	Diabetes mellitus (E10-E14)	274	3.5	11.2
8	Cerebrovascular diseases (I60-I69)	262	3.3	10.7
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	149	1.9	6.1
10	Septicemia (A40-A41)	116	1.5	4.7
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	112	1.4	4.6
12	Influenza and pneumonia (J09-J18)	93	1.2	3.8
12	Chronic liver disease and cirrhosis (K70,K73-K74)	93	1.2	3.8
14	Chronic lower respiratory diseases (J40-J47)	86	1.1	3.5
15	Aortic aneurysm and dissection (I71)	67	0.9	2.7
...	All other causes (Residual)	1,197	15.2	48.8
Non-Hispanic, black, male, 45-54 years				
...	All causes	18,938	100.0	730.1
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	4,959	26.2	191.2
2	Malignant neoplasms (C00-C97)	3,905	20.6	150.5
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,613	8.5	62.2
4	Human immunodeficiency virus (HIV) disease (B20-B24)	960	5.1	37.0
5	Cerebrovascular diseases (I60-I69)	911	4.8	35.1
6	Diabetes mellitus (E10-E14)	792	4.2	30.5
7	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	570	3.0	22.0
8	Chronic liver disease and cirrhosis (K70,K73-K74)	475	2.5	18.3
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	399	2.1	15.4
10	Chronic lower respiratory diseases (J40-J47)	315	1.7	12.1
11	Septicemia (A40-A41)	314	1.7	12.1
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	297	1.6	11.4
13	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	284	1.5	10.9
14	Influenza and pneumonia (J09-J18)	215	1.1	8.3
15	Viral hepatitis (B15-B19)	179	0.9	6.9
...	All other causes (Residual)	2,750	14.5	106.0

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> , Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, male, 55-64 years				
...	All causes	31,271	100.0	1,667.7
1	Malignant neoplasms (C00-C97)	9,232	29.5	492.3
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	8,177	26.1	436.1
3	Cerebrovascular diseases (I60-I69)	1,571	5.0	83.8
4	Diabetes mellitus (E10-E14)	1,534	4.9	81.8
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,419	4.5	75.7
6	Chronic lower respiratory diseases (J40-J47)	801	2.6	42.7
7	Chronic liver disease and cirrhosis (K70,K73-K74)	785	2.5	41.9
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	784	2.5	41.8
9	Human immunodeficiency virus (HIV) disease (B20-B24)	697	2.2	37.2
10	Septicemia (A40-A41)	608	1.9	32.4
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	476	1.5	25.4
12	Viral hepatitis (B15-B19)	409	1.3	21.8
13	Influenza and pneumonia (J09-J18)	396	1.3	21.1
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	277	0.9	14.8
15	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	153	0.5	8.2
...	All other causes (Residual)	3,952	12.6	210.8
Non-Hispanic, black, male, 65-74 years				
...	All causes	28,597	100.0	3,203.3
1	Malignant neoplasms (C00-C97)	9,181	32.1	1,028.4
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	7,527	26.3	843.1
3	Cerebrovascular diseases (I60-I69)	1,576	5.5	176.5
4	Diabetes mellitus (E10-E14)	1,480	5.2	165.8
5	Chronic lower respiratory diseases (J40-J47)	1,249	4.4	139.9
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	885	3.1	99.1
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	606	2.1	67.9
8	Septicemia (A40-A41)	583	2.0	65.3
9	Essential hypertension and hypertensive renal disease (I10,I12,I15)	473	1.7	53.0
10	Influenza and pneumonia (J09-J18)	438	1.5	49.1
11	Chronic liver disease and cirrhosis (K70,K73-K74)	262	0.9	29.3
12	Alzheimer's disease (G30)	183	0.6	20.5
13	Human immunodeficiency virus (HIV) disease (B20-B24)	178	0.6	19.9
14	Pneumonitis due to solids and liquids (J69)	148	0.5	16.6
15	Parkinson's disease (G20-G21)	119	0.4	13.3
...	All other causes (Residual)	3,709	13.0	415.5

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> , Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, male, 75-84 years				
...	All causes	26,061	100.0	6,498.5
1	Malignant neoplasms (C00-C97)	7,004	26.9	1,746.5
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	6,862	26.3	1,711.1
3	Cerebrovascular diseases (I60-I69)	1,584	6.1	395.0
4	Chronic lower respiratory diseases (J40-J47)	1,296	5.0	323.2
5	Diabetes mellitus (E10-E14)	1,168	4.5	291.2
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	872	3.3	217.4
7	Alzheimer's disease (G30)	616	2.4	153.6
8	Septicemia (A40-A41)	583	2.2	145.4
9	Influenza and pneumonia (J09-J18)	576	2.2	143.6
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	462	1.8	115.2
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	401	1.5	100.0
12	Pneumonitis due to solids and liquids (J69)	221	0.8	55.1
13	Parkinson's disease (G20-G21)	219	0.8	54.6
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	130	0.5	32.4
15	Chronic liver disease and cirrhosis (K70,K73-K74)	90	0.3	22.4
...	All other causes (Residual)	3,977	15.3	991.7
Non-Hispanic, black, male, 85 years and over				
...	All causes	16,098	100.0	14,516.7
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	4,369	27.1	3,939.8
2	Malignant neoplasms (C00-C97)	3,037	18.9	2,738.7
3	Cerebrovascular diseases (I60-I69)	942	5.9	849.5
4	Alzheimer's disease (G30)	719	4.5	648.4
5	Chronic lower respiratory diseases (J40-J47)	708	4.4	638.5
6	Diabetes mellitus (E10-E14)	600	3.7	541.1
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	557	3.5	502.3
8	Influenza and pneumonia (J09-J18)	515	3.2	464.4
9	Septicemia (A40-A41)	362	2.2	326.4
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	328	2.0	295.8
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	253	1.6	228.1
12	Pneumonitis due to solids and liquids (J69)	202	1.3	182.2
13	Parkinson's disease (G20-G21)	124	0.8	111.8
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	72	0.4	64.9
14	Atherosclerosis (I70)	72	0.4	64.9
...	All other causes (Residual)	3,238	20.1	2,919.9

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, female, all ages				
...	All causes	141,745	100.0	679.2
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	33,119	23.4	158.7
2	Malignant neoplasms (C00-C97)	32,530	22.9	155.9
3	Cerebrovascular diseases (I60-I69)	8,743	6.2	41.9
4	Diabetes mellitus (E10-E14)	6,785	4.8	32.5
5	Chronic lower respiratory diseases (J40-J47)	4,346	3.1	20.8
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	4,310	3.0	20.7
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	4,080	2.9	19.5
8	Alzheimer's disease (G30)	3,889	2.7	18.6
9	Septicemia (A40-A41)	3,297	2.3	15.8
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	2,785	2.0	13.3
11	Influenza and pneumonia (J09-J18)	2,712	1.9	13.0
12	Certain conditions originating in the perinatal period (P00-P96)	1,719	1.2	8.2
13	Human immunodeficiency virus (HIV) disease (B20-B24)	1,445	1.0	6.9
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,106	0.8	5.3
15	Chronic liver disease and cirrhosis (K70,K73-K74)	1,033	0.7	4.9
...	All other causes (Residual)	29,846	21.1	143.0
Non-Hispanic, black, female, 1-4 years				
...	All causes	436	100.0	35.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	119	27.3	9.8
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	70	16.1	5.7
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	48	11.0	3.9
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	24	5.5	2.0
5	Malignant neoplasms (C00-C97)	23	5.3	1.9
6	Influenza and pneumonia (J09-J18)	14	3.2	*
7	Chronic lower respiratory diseases (J40-J47)	8	1.8	*
8	Septicemia (A40-A41)	5	1.1	*
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	4	0.9	*
9	Certain conditions originating in the perinatal period (P00-P96)	4	0.9	*
11	Anemias (D50-D64)	3	0.7	*
11	Cerebrovascular diseases (I60-I69)	3	0.7	*
13	Complications of medical and surgical care (Y40-Y84,Y88)	2	0.5	*
14	Meningitis (G00,G03)	1	0.2	*
14	Atherosclerosis (I70)	1	0.2	*
14	Acute bronchitis and bronchiolitis (J20-J21)	1	0.2	*
14	Pneumonitis due to solids and liquids (J69)	1	0.2	*
14	Peptic ulcer (K25-K28)	1	0.2	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1	0.2	*
14	Enterocolitis due to <i>Clostridium difficile</i> (A04.7)	1	0.2	*
...	All other causes (Residual)	102	23.4	8.4

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> , Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, female, 5-14 years				
...	All causes	466	100.0	15.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	117	25.1	3.9
2	Malignant neoplasms (C00-C97)	54	11.6	1.8
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	35	7.5	1.2
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	28	6.0	0.9
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	24	5.2	0.8
5	Chronic lower respiratory diseases (J40-J47)	24	5.2	0.8
7	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	11	2.4	*
8	Influenza and pneumonia (J09-J18)	10	2.1	*
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	9	1.9	*
9	Cerebrovascular diseases (I60-I69)	9	1.9	*
11	Anemias (D50-D64)	8	1.7	*
12	Septicemia (A40-A41)	7	1.5	*
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	5	1.1	*
14	Meningitis (G00,G03)	3	0.6	*
15	Diseases of appendix (K35-K38)	2	0.4	*
...	All other causes (Residual)	120	25.8	4.0
Non-Hispanic, black, female, 15-24 years				
...	All causes	1,536	100.0	45.9
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	345	22.5	10.3
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	289	18.8	8.6
2	Malignant neoplasms (C00-C97)	106	6.9	3.2
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	106	6.9	3.2
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	79	5.1	2.4
6	Pregnancy, childbirth and the puerperium (O00-O99)	62	4.0	1.9
7	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	40	2.6	1.2
8	Human immunodeficiency virus (HIV) disease (B20-B24)	38	2.5	1.1
9	Anemias (D50-D64)	28	1.8	0.8
10	Chronic lower respiratory diseases (J40-J47)	27	1.8	0.8
11	Diabetes mellitus (E10-E14)	24	1.6	0.7
11	Influenza and pneumonia (J09-J18)	24	1.6	0.7
13	Septicemia (A40-A41)	23	1.5	0.7
14	Cerebrovascular diseases (I60-I69)	15	1.0	*
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	11	0.7	*
...	All other causes (Residual)	319	20.8	9.5

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> , Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, female, 25-34 years				
...	All causes	2,939	100.0	101.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	459	15.6	15.9
2	Malignant neoplasms (C00-C97)	380	12.9	13.1
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	374	12.7	12.9
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	241	8.2	8.3
5	Human immunodeficiency virus (HIV) disease (B20-B24)	172	5.9	6.0
6	Pregnancy, childbirth and the puerperium (O00-O99)	124	4.2	4.3
7	Diabetes mellitus (E10-E14)	114	3.9	3.9
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	102	3.5	3.5
9	Cerebrovascular diseases (I60-I69)	68	2.3	2.4
10	Anemias (D50-D64)	58	2.0	2.0
11	Septicemia (A40-A41)	57	1.9	2.0
12	Influenza and pneumonia (J09-J18)	51	1.7	1.8
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	42	1.4	1.5
14	Chronic lower respiratory diseases (J40-J47)	33	1.1	1.1
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	31	1.1	1.1
...	All other causes (Residual)	633	21.5	21.9
Non-Hispanic, black, female, 35-44 years				
...	All causes	5,742	100.0	208.4
1	Malignant neoplasms (C00-C97)	1,342	23.4	48.7
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	968	16.9	35.1
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	482	8.4	17.5
4	Human immunodeficiency virus (HIV) disease (B20-B24)	387	6.7	14.0
5	Cerebrovascular diseases (I60-I69)	247	4.3	9.0
6	Diabetes mellitus (E10-E14)	213	3.7	7.7
7	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	178	3.1	6.5
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	121	2.1	4.4
9	Chronic lower respiratory diseases (J40-J47)	105	1.8	3.8
10	Septicemia (A40-A41)	99	1.7	3.6
11	Influenza and pneumonia (J09-J18)	86	1.5	3.1
11	Chronic liver disease and cirrhosis (K70,K73-K74)	86	1.5	3.1
13	Pregnancy, childbirth and the puerperium (O00-O99)	85	1.5	3.1
14	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	81	1.4	2.9
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	74	1.3	2.7
...	All other causes (Residual)	1,188	20.7	43.1

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> , Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, female, 45-54 years				
...	All causes	14,213	100.0	485.2
1	Malignant neoplasms (C00-C97)	4,405	31.0	150.4
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,790	19.6	95.2
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	745	5.2	25.4
4	Cerebrovascular diseases (I60-I69)	720	5.1	24.6
5	Diabetes mellitus (E10-E14)	644	4.5	22.0
6	Human immunodeficiency virus (HIV) disease (B20-B24)	469	3.3	16.0
7	Chronic lower respiratory diseases (J40-J47)	378	2.7	12.9
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	326	2.3	11.1
9	Chronic liver disease and cirrhosis (K70,K73-K74)	310	2.2	10.6
10	Septicemia (A40-A41)	309	2.2	10.5
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	198	1.4	6.8
12	Influenza and pneumonia (J09-J18)	181	1.3	6.2
13	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	157	1.1	5.4
14	Viral hepatitis (B15-B19)	96	0.7	3.3
15	Anemias (D50-D64)	83	0.6	2.8
...	All other causes (Residual)	2,402	16.9	82.0
Non-Hispanic, black, female, 55-64 years				
...	All causes	22,025	100.0	979.2
1	Malignant neoplasms (C00-C97)	7,575	34.4	336.8
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	4,845	22.0	215.4
3	Diabetes mellitus (E10-E14)	1,211	5.5	53.8
4	Cerebrovascular diseases (I60-I69)	1,119	5.1	49.8
5	Chronic lower respiratory diseases (J40-J47)	668	3.0	29.7
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	633	2.9	28.1
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	582	2.6	25.9
8	Septicemia (A40-A41)	511	2.3	22.7
9	Essential hypertension and hypertensive renal disease (I10,I12,I15)	368	1.7	16.4
10	Chronic liver disease and cirrhosis (K70,K73-K74)	364	1.7	16.2
11	Influenza and pneumonia (J09-J18)	304	1.4	13.5
12	Human immunodeficiency virus (HIV) disease (B20-B24)	282	1.3	12.5
13	Viral hepatitis (B15-B19)	188	0.9	8.4
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	105	0.5	4.7
15	Pneumonitis due to solids and liquids (J69)	74	0.3	3.3
...	All other causes (Residual)	3,196	14.5	142.1

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> , Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, female, 65-74 years				
...	All causes	24,147	100.0	2,018.0
1	Malignant neoplasms (C00-C97)	7,627	31.6	637.4
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	5,668	23.5	473.7
3	Diabetes mellitus (E10-E14)	1,503	6.2	125.6
4	Cerebrovascular diseases (I60-I69)	1,428	5.9	119.3
5	Chronic lower respiratory diseases (J40-J47)	957	4.0	80.0
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	860	3.6	71.9
7	Septicemia (A40-A41)	656	2.7	54.8
8	Essential hypertension and hypertensive renal disease (I10,I12,I15)	432	1.8	36.1
9	Influenza and pneumonia (J09-J18)	405	1.7	33.8
10	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	337	1.4	28.2
11	Alzheimer's disease (G30)	222	0.9	18.6
12	Chronic liver disease and cirrhosis (K70,K73-K74)	143	0.6	12.0
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	119	0.5	9.9
14	Pneumonitis due to solids and liquids (J69)	105	0.4	8.8
15	Aortic aneurysm and dissection (I71)	82	0.3	6.9
...	All other causes (Residual)	3,603	14.9	301.1
Non-Hispanic, black, female, 75-84 years				
...	All causes	31,306	100.0	4,637.5
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	7,980	25.5	1,182.1
2	Malignant neoplasms (C00-C97)	6,977	22.3	1,033.5
3	Cerebrovascular diseases (I60-I69)	2,355	7.5	348.9
4	Diabetes mellitus (E10-E14)	1,742	5.6	258.1
5	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,200	3.8	177.8
6	Chronic lower respiratory diseases (J40-J47)	1,195	3.8	177.0
7	Alzheimer's disease (G30)	1,094	3.5	162.1
8	Septicemia (A40-A41)	807	2.6	119.5
9	Influenza and pneumonia (J09-J18)	643	2.1	95.3
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	641	2.0	95.0
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	359	1.1	53.2
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	200	0.6	29.6
13	Pneumonitis due to solids and liquids (J69)	185	0.6	27.4
14	Parkinson's disease (G20-G21)	163	0.5	24.1
15	Enterocolitis due to <i>Clostridium difficile</i> (A04.7)	120	0.4	17.8
...	All other causes (Residual)	5,645	18.0	836.2

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> , Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, female, 85 years and over				
...	All causes	35,835	100.0	12,591.4
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	10,292	28.7	3,616.3
2	Malignant neoplasms (C00-C97)	4,033	11.3	1,417.1
3	Cerebrovascular diseases (I60-I69)	2,759	7.7	969.4
4	Alzheimer's disease (G30)	2,535	7.1	890.7
5	Diabetes mellitus (E10-E14)	1,333	3.7	468.4
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,101	3.1	386.9
7	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,045	2.9	367.2
8	Influenza and pneumonia (J09-J18)	959	2.7	337.0
9	Chronic lower respiratory diseases (J40-J47)	948	2.6	333.1
10	Septicemia (A40-A41)	800	2.2	281.1
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	370	1.0	130.0
12	Pneumonitis due to solids and liquids (J69)	303	0.8	106.5
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	166	0.5	58.3
13	Atherosclerosis (I70)	166	0.5	58.3
15	Parkinson's disease (G20-G21)	145	0.4	50.9
...	All other causes (Residual)	8,880	24.8	3,120.2

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011