

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2000

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Population denominators for 2000 death rates are based on the 1990 census population.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, both sexes, all ages				
...	All causes	107,254	100.0	330.4
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	25,819	24.1	79.5
2	Malignant neoplasms (C00-C97)	21,160	19.7	65.2
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	8,830	8.2	27.2
4	Cerebrovascular diseases (I60-I69)	6,187	5.8	19.1
5	Diabetes mellitus (E10-E14)	5,328	5.0	16.4
6	Chronic liver disease and cirrhosis (K70,K73-K74)	3,187	3.0	9.8
7	Assault (homicide) (X85-Y09,Y87.1)	2,917	2.7	9.0
8	Chronic lower respiratory diseases (J40-J47)	2,689	2.5	8.3
9	Influenza and pneumonia (J10-J18)	2,625	2.4	8.1
10	Certain conditions originating in the perinatal period (P00-P96)	2,145	2.0	6.6
11	Human immunodeficiency virus (HIV) disease (B20-B24)	1,908	1.8	5.9
12	Intentional self-harm (suicide) (X60-X84,Y87.0)	1,787	1.7	5.5
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,646	1.5	5.1
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	1,612	1.5	5.0
15	Septicemia (A40-A41)	1,226	1.1	3.8
...	All other causes (Residual)	18,188	17.0	56.0
Hispanic, both sexes, 1-4 years				
...	All causes	872	100.0	31.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	306	35.1	10.9
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	111	12.7	3.9
3	Malignant neoplasms (C00-C97)	75	8.6	2.7
4	Assault (homicide) (X85-Y09,Y87.1)	57	6.5	2.0
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	29	3.3	1.0
6	Influenza and pneumonia (J10-J18)	17	1.9	*
7	Septicemia (A40-A41)	15	1.7	*
8	Certain conditions originating in the perinatal period (P00-P96)	12	1.4	*
9	Chronic lower respiratory diseases (J40-J47)	9	1.0	*
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	7	0.8	*
11	Cerebrovascular diseases (I60-I69)	6	0.7	*
12	Meningitis (G00,G03)	4	0.5	*
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	4	0.5	*
12	Complications of medical and surgical care (Y40-Y84,Y88)	4	0.5	*
15	Meningococcal infection (A39)	3	0.3	*
...	All other causes (Residual)	213	24.4	7.6

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2000

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Population denominators for 2000 death rates are based on the 1990 census population.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, both sexes, 5-14 years				
...	All causes	1,065	100.0	17.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	383	36.0	6.1
2	Malignant neoplasms (C00-C97)	198	18.6	3.2
3	Assault (homicide) (X85-Y09,Y87.1)	76	7.1	1.2
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	52	4.9	0.8
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	38	3.6	0.6
6	Intentional self-harm (suicide) (X60-X84,Y87.0)	29	2.7	0.5
7	Influenza and pneumonia (J10-J18)	21	2.0	0.3
8	Chronic lower respiratory diseases (J40-J47)	15	1.4	*
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	13	1.2	*
10	Certain conditions originating in the perinatal period (P00-P96)	10	0.9	*
11	Human immunodeficiency virus (HIV) disease (B20-B24)	9	0.8	*
12	Cerebrovascular diseases (I60-I69)	8	0.8	*
13	Septicemia (A40-A41)	7	0.7	*
13	Anemias (D50-D64)	7	0.7	*
15	Complications of medical and surgical care (Y40-Y84,Y88)	6	0.6	*
...	All other causes (Residual)	193	18.1	3.1
Hispanic, both sexes, 15-24 years				
...	All causes	4,793	100.0	85.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,095	43.7	37.2
2	Assault (homicide) (X85-Y09,Y87.1)	1,126	23.5	20.0
3	Intentional self-harm (suicide) (X60-X84,Y87.0)	447	9.3	7.9
4	Malignant neoplasms (C00-C97)	275	5.7	4.9
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	125	2.6	2.2
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	54	1.1	1.0
7	Cerebrovascular diseases (I60-I69)	31	0.6	0.5
8	Pregnancy, childbirth and the puerperium (O00-099)	24	0.5	0.4
9	Human immunodeficiency virus (HIV) disease (B20-B24)	23	0.5	0.4
10	Influenza and pneumonia (J10-J18)	21	0.4	0.4
11	Legal intervention (Y35,Y89.0)	18	0.4	*
12	Chronic lower respiratory diseases (J40-J47)	16	0.3	*
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	14	0.3	*
14	Diabetes mellitus (E10-E14)	13	0.3	*
15	Septicemia (A40-A41)	12	0.3	*
...	All other causes (Residual)	499	10.4	8.9

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2000

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Population denominators for 2000 death rates are based on the 1990 census population.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
	Hispanic, both sexes, 25-44 years			
...	All causes	13,594	100.0	133.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,394	25.0	33.2
2	Malignant neoplasms (C00-C97)	1,771	13.0	17.3
3	Assault (homicide) (X85-Y09,Y87.1)	1,257	9.2	12.3
4	Human immunodeficiency virus (HIV) disease (B20-B24)	1,127	8.3	11.0
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,040	7.7	10.2
6	Intentional self-harm (suicide) (X60-X84,Y87.0)	805	5.9	7.9
7	Chronic liver disease and cirrhosis (K70,K73-K74)	593	4.4	5.8
8	Cerebrovascular diseases (I60-I69)	375	2.8	3.7
9	Diabetes mellitus (E10-E14)	218	1.6	2.1
10	Viral hepatitis (B15-B19)	140	1.0	1.4
11	Influenza and pneumonia (J10-J18)	130	1.0	1.3
12	Septicemia (A40-A41)	110	0.8	1.1
13	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	89	0.7	0.9
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	79	0.6	0.8
15	Chronic lower respiratory diseases (J40-J47)	75	0.6	0.7
...	All other causes (Residual)	2,391	17.6	23.4
	Hispanic, both sexes, 45-64 years			
...	All causes	23,089	100.0	473.7
1	Malignant neoplasms (C00-C97)	6,355	27.5	130.4
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	4,890	21.2	100.3
3	Chronic liver disease and cirrhosis (K70,K73-K74)	1,562	6.8	32.0
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,445	6.3	29.6
5	Diabetes mellitus (E10-E14)	1,389	6.0	28.5
6	Cerebrovascular diseases (I60-I69)	1,158	5.0	23.8
7	Human immunodeficiency virus (HIV) disease (B20-B24)	694	3.0	14.2
8	Viral hepatitis (B15-B19)	365	1.6	7.5
9	Chronic lower respiratory diseases (J40-J47)	349	1.5	7.2
10	Intentional self-harm (suicide) (X60-X84,Y87.0)	346	1.5	7.1
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	341	1.5	7.0
12	Influenza and pneumonia (J10-J18)	311	1.3	6.4
13	Assault (homicide) (X85-Y09,Y87.1)	288	1.2	5.9
14	Septicemia (A40-A41)	267	1.2	5.5
15	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	164	0.7	3.4
...	All other causes (Residual)	3,165	13.7	64.9

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2000

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Population denominators for 2000 death rates are based on the 1990 census population.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
	Hispanic, both sexes, 65 years and over			
...	All causes	59,189	100.0	3,034.3
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	19,606	33.1	1,005.1
2	Malignant neoplasms (C00-C97)	12,469	21.1	639.2
3	Cerebrovascular diseases (I60-I69)	4,593	7.8	235.5
4	Diabetes mellitus (E10-E14)	3,703	6.3	189.8
5	Chronic lower respiratory diseases (J40-J47)	2,218	3.7	113.7
6	Influenza and pneumonia (J10-J18)	2,065	3.5	105.9
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,179	2.0	60.4
8	Alzheimer's disease (G30)	1,102	1.9	56.5
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,061	1.8	54.4
10	Chronic liver disease and cirrhosis (K70,K73-K74)	1,024	1.7	52.5
11	Septicemia (A40-A41)	753	1.3	38.6
12	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	617	1.0	31.6
13	Pneumonitis due to solids and liquids (J69)	412	0.7	21.1
14	Parkinson's disease (G20-G21)	405	0.7	20.8
15	Atherosclerosis (I70)	387	0.7	19.8
...	All other causes (Residual)	7,595	12.8	389.4
	Hispanic, male, all ages			
...	All causes	60,172	100.0	368.9
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	13,566	22.5	83.2
2	Malignant neoplasms (C00-C97)	11,138	18.5	68.3
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	6,696	11.1	41.1
4	Cerebrovascular diseases (I60-I69)	2,865	4.8	17.6
5	Diabetes mellitus (E10-E14)	2,507	4.2	15.4
6	Assault (homicide) (X85-Y09,Y87.1)	2,431	4.0	14.9
7	Chronic liver disease and cirrhosis (K70,K73-K74)	2,312	3.8	14.2
8	Intentional self-harm (suicide) (X60-X84,Y87.0)	1,525	2.5	9.3
9	Human immunodeficiency virus (HIV) disease (B20-B24)	1,493	2.5	9.2
10	Chronic lower respiratory diseases (J40-J47)	1,451	2.4	8.9
11	Influenza and pneumonia (J10-J18)	1,303	2.2	8.0
12	Certain conditions originating in the perinatal period (P00-P96)	1,194	2.0	7.3
13	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	857	1.4	5.3
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	805	1.3	4.9
15	Septicemia (A40-A41)	625	1.0	3.8
...	All other causes (Residual)	9,404	15.6	57.7

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2000

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Population denominators for 2000 death rates are based on the 1990 census population.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, male, 1-4 years				
...	All causes	475	100.0	33.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	174	36.6	12.1
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	60	12.6	4.2
3	Malignant neoplasms (C00-C97)	43	9.1	3.0
4	Assault (homicide) (X85-Y09,Y87.1)	30	6.3	2.1
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	21	4.4	1.5
6	Septicemia (A40-A41)	9	1.9	*
7	Certain conditions originating in the perinatal period (P00-P96)	7	1.5	*
8	Influenza and pneumonia (J10-J18)	5	1.1	*
8	Chronic lower respiratory diseases (J40-J47)	5	1.1	*
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3	0.6	*
11	Meningococcal infection (A39)	2	0.4	*
11	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2	0.4	*
11	Meningitis (G00,G03)	2	0.4	*
11	Complications of medical and surgical care (Y40-Y84,Y88)	2	0.4	*
15	Acute bronchitis and bronchiolitis (J20-J21)	1	0.2	*
15	Diseases of appendix (K35-K38)	1	0.2	*
...	All other causes (Residual)	108	22.7	7.5
Hispanic, male, 5-14 years				
...	All causes	622	100.0	19.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	210	33.8	6.6
2	Malignant neoplasms (C00-C97)	122	19.6	3.8
3	Assault (homicide) (X85-Y09,Y87.1)	56	9.0	1.8
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	28	4.5	0.9
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	26	4.2	0.8
6	Intentional self-harm (suicide) (X60-X84,Y87.0)	22	3.5	0.7
7	Influenza and pneumonia (J10-J18)	10	1.6	*
8	Chronic lower respiratory diseases (J40-J47)	9	1.4	*
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	8	1.3	*
10	Certain conditions originating in the perinatal period (P00-P96)	6	1.0	*
11	Septicemia (A40-A41)	5	0.8	*
11	Human immunodeficiency virus (HIV) disease (B20-B24)	5	0.8	*
13	Complications of medical and surgical care (Y40-Y84,Y88)	4	0.6	*
14	Anemias (D50-D64)	3	0.5	*
14	Cerebrovascular diseases (I60-I69)	3	0.5	*
14	Diseases of appendix (K35-K38)	3	0.5	*
...	All other causes (Residual)	102	16.4	3.2

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2000

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Population denominators for 2000 death rates are based on the 1990 census population.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
	Hispanic, male, 15-24 years			
...	All causes	3,837	100.0	131.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,717	44.7	58.7
2	Assault (homicide) (X85-Y09,Y87.1)	1,014	26.4	34.7
3	Intentional self-harm (suicide) (X60-X84,Y87.0)	387	10.1	13.2
4	Malignant neoplasms (C00-C97)	162	4.2	5.5
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	87	2.3	3.0
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	39	1.0	1.3
7	Human immunodeficiency virus (HIV) disease (B20-B24)	18	0.5	*
8	Cerebrovascular diseases (I60-I69)	17	0.4	*
8	Legal intervention (Y35,Y89.0)	17	0.4	*
10	Chronic lower respiratory diseases (J40-J47)	13	0.3	*
11	Influenza and pneumonia (J10-J18)	10	0.3	*
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	8	0.2	*
13	Diabetes mellitus (E10-E14)	7	0.2	*
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	7	0.2	*
15	Meningococcal infection (A39)	6	0.2	*
15	Septicemia (A40-A41)	6	0.2	*
15	Complications of medical and surgical care (Y40-Y84,Y88)	6	0.2	*
...	All other causes (Residual)	316	8.2	10.8
	Hispanic, male, 25-44 years			
...	All causes	9,797	100.0	187.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,816	28.7	53.8
2	Assault (homicide) (X85-Y09,Y87.1)	1,056	10.8	20.2
3	Human immunodeficiency virus (HIV) disease (B20-B24)	877	9.0	16.8
4	Malignant neoplasms (C00-C97)	775	7.9	14.8
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	749	7.6	14.3
6	Intentional self-harm (suicide) (X60-X84,Y87.0)	690	7.0	13.2
7	Chronic liver disease and cirrhosis (K70,K73-K74)	481	4.9	9.2
8	Cerebrovascular diseases (I60-I69)	210	2.1	4.0
9	Diabetes mellitus (E10-E14)	143	1.5	2.7
10	Viral hepatitis (B15-B19)	111	1.1	2.1
11	Influenza and pneumonia (J10-J18)	83	0.8	1.6
12	Septicemia (A40-A41)	67	0.7	1.3
13	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	53	0.5	1.0
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	47	0.5	0.9
15	Chronic lower respiratory diseases (J40-J47)	36	0.4	0.7
...	All other causes (Residual)	1,603	16.4	30.7

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2000

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Population denominators for 2000 death rates are based on the 1990 census population.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
	Hispanic, male, 45-64 years			
...	All causes	14,568	100.0	622.6
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,435	23.6	146.8
2	Malignant neoplasms (C00-C97)	3,266	22.4	139.6
3	Chronic liver disease and cirrhosis (K70,K73-K74)	1,238	8.5	52.9
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,097	7.5	46.9
5	Diabetes mellitus (E10-E14)	760	5.2	32.5
6	Cerebrovascular diseases (I60-I69)	672	4.6	28.7
7	Human immunodeficiency virus (HIV) disease (B20-B24)	549	3.8	23.5
8	Intentional self-harm (suicide) (X60-X84,Y87.0)	283	1.9	12.1
9	Viral hepatitis (B15-B19)	259	1.8	11.1
10	Assault (homicide) (X85-Y09,Y87.1)	215	1.5	9.2
11	Influenza and pneumonia (J10-J18)	201	1.4	8.6
12	Chronic lower respiratory diseases (J40-J47)	199	1.4	8.5
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	184	1.3	7.9
14	Septicemia (A40-A41)	158	1.1	6.8
15	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	109	0.7	4.7
...	All other causes (Residual)	1,943	13.3	83.0
	Hispanic, male, 65 years and over			
...	All causes	28,308	100.0	3,434.5
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	9,200	32.5	1,116.2
2	Malignant neoplasms (C00-C97)	6,760	23.9	820.2
3	Cerebrovascular diseases (I60-I69)	1,955	6.9	237.2
4	Diabetes mellitus (E10-E14)	1,594	5.6	193.4
5	Chronic lower respiratory diseases (J40-J47)	1,185	4.2	143.8
6	Influenza and pneumonia (J10-J18)	964	3.4	117.0
7	Chronic liver disease and cirrhosis (K70,K73-K74)	588	2.1	71.3
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	587	2.1	71.2
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	545	1.9	66.1
10	Alzheimer's disease (G30)	370	1.3	44.9
11	Septicemia (A40-A41)	347	1.2	42.1
12	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	249	0.9	30.2
13	Parkinson's disease (G20-G21)	229	0.8	27.8
14	Pneumonitis due to solids and liquids (J69)	206	0.7	25.0
15	Aortic aneurysm and dissection (I71)	194	0.7	23.5
...	All other causes (Residual)	3,335	11.8	404.6

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2000

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Population denominators for 2000 death rates are based on the 1990 census population.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
	Hispanic, female, all ages			
...	All causes	47,082	100.0	291.5
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	12,253	26.0	75.9
2	Malignant neoplasms (C00-C97)	10,022	21.3	62.0
3	Cerebrovascular diseases (I60-I69)	3,322	7.1	20.6
4	Diabetes mellitus (E10-E14)	2,821	6.0	17.5
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,134	4.5	13.2
6	Influenza and pneumonia (J10-J18)	1,322	2.8	8.2
7	Chronic lower respiratory diseases (J40-J47)	1,238	2.6	7.7
8	Certain conditions originating in the perinatal period (P00-P96)	951	2.0	5.9
9	Chronic liver disease and cirrhosis (K70,K73-K74)	875	1.9	5.4
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	841	1.8	5.2
11	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	755	1.6	4.7
12	Alzheimer's disease (G30)	744	1.6	4.6
13	Septicemia (A40-A41)	601	1.3	3.7
14	Assault (homicide) (X85-Y09,Y87.1)	486	1.0	3.0
15	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	435	0.9	2.7
...	All other causes (Residual)	8,282	17.6	51.3
	Hispanic, female, 1-4 years			
...	All causes	397	100.0	28.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	132	33.2	9.6
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	51	12.8	3.7
3	Malignant neoplasms (C00-C97)	32	8.1	2.3
4	Assault (homicide) (X85-Y09,Y87.1)	27	6.8	2.0
5	Influenza and pneumonia (J10-J18)	12	3.0	*
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	8	2.0	*
7	Septicemia (A40-A41)	6	1.5	*
7	Cerebrovascular diseases (I60-I69)	6	1.5	*
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	5	1.3	*
9	Certain conditions originating in the perinatal period (P00-P96)	5	1.3	*
11	Chronic lower respiratory diseases (J40-J47)	4	1.0	*
12	Tuberculosis (A16-A19)	2	0.5	*
12	Meningitis (G00,G03)	2	0.5	*
12	Pneumonitis due to solids and liquids (J69)	2	0.5	*
12	Complications of medical and surgical care (Y40-Y84,Y88)	2	0.5	*
...	All other causes (Residual)	101	25.4	7.3

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2000

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Population denominators for 2000 death rates are based on the 1990 census population.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
	Hispanic, female, 5-14 years			
...	All causes	443	100.0	14.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	173	39.1	5.7
2	Malignant neoplasms (C00-C97)	76	17.2	2.5
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	24	5.4	0.8
4	Assault (homicide) (X85-Y09,Y87.1)	20	4.5	0.7
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	12	2.7	*
6	Influenza and pneumonia (J10-J18)	11	2.5	*
7	Intentional self-harm (suicide) (X60-X84,Y87.0)	7	1.6	*
8	Chronic lower respiratory diseases (J40-J47)	6	1.4	*
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	5	1.1	*
9	Cerebrovascular diseases (I60-I69)	5	1.1	*
11	Human immunodeficiency virus (HIV) disease (B20-B24)	4	0.9	*
11	Anemias (D50-D64)	4	0.9	*
11	Certain conditions originating in the perinatal period (P00-P96)	4	0.9	*
14	Meningococcal infection (A39)	2	0.5	*
14	Septicemia (A40-A41)	2	0.5	*
14	Diabetes mellitus (E10-E14)	2	0.5	*
14	Meningitis (G00,G03)	2	0.5	*
14	Complications of medical and surgical care (Y40-Y84,Y88)	2	0.5	*
...	All other causes (Residual)	82	18.5	2.7
	Hispanic, female, 15-24 years			
...	All causes	956	100.0	35.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	378	39.5	13.9
2	Malignant neoplasms (C00-C97)	113	11.8	4.2
3	Assault (homicide) (X85-Y09,Y87.1)	112	11.7	4.1
4	Intentional self-harm (suicide) (X60-X84,Y87.0)	60	6.3	2.2
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	38	4.0	1.4
6	Pregnancy, childbirth and the puerperium (O00-099)	24	2.5	0.9
7	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	15	1.6	*
8	Cerebrovascular diseases (I60-I69)	14	1.5	*
9	Influenza and pneumonia (J10-J18)	11	1.2	*
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	7	0.7	*
11	Septicemia (A40-A41)	6	0.6	*
11	Diabetes mellitus (E10-E14)	6	0.6	*
13	Human immunodeficiency virus (HIV) disease (B20-B24)	5	0.5	*
14	Anemias (D50-D64)	4	0.4	*
14	Aortic aneurysm and dissection (I71)	4	0.4	*
...	All other causes (Residual)	159	16.6	5.9

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2000

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Population denominators for 2000 death rates are based on the 1990 census population.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
	Hispanic, female, 25-44 years			
...	All causes	3,797	100.0	76.1
1	Malignant neoplasms (C00-C97)	996	26.2	20.0
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	578	15.2	11.6
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	291	7.7	5.8
4	Human immunodeficiency virus (HIV) disease (B20-B24)	250	6.6	5.0
5	Assault (homicide) (X85-Y09,Y87.1)	201	5.3	4.0
6	Cerebrovascular diseases (I60-I69)	165	4.3	3.3
7	Intentional self-harm (suicide) (X60-X84,Y87.0)	115	3.0	2.3
8	Chronic liver disease and cirrhosis (K70,K73-K74)	112	2.9	2.2
9	Diabetes mellitus (E10-E14)	75	2.0	1.5
10	Pregnancy, childbirth and the puerperium (O00-099)	59	1.6	1.2
11	Influenza and pneumonia (J10-J18)	47	1.2	0.9
12	Septicemia (A40-A41)	43	1.1	0.9
13	Chronic lower respiratory diseases (J40-J47)	39	1.0	0.8
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	36	0.9	0.7
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	33	0.9	0.7
...	All other causes (Residual)	757	19.9	15.2
	Hispanic, female, 45-64 years			
...	All causes	8,521	100.0	336.2
1	Malignant neoplasms (C00-C97)	3,089	36.3	121.9
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,455	17.1	57.4
3	Diabetes mellitus (E10-E14)	629	7.4	24.8
4	Cerebrovascular diseases (I60-I69)	486	5.7	19.2
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	348	4.1	13.7
6	Chronic liver disease and cirrhosis (K70,K73-K74)	324	3.8	12.8
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	157	1.8	6.2
8	Chronic lower respiratory diseases (J40-J47)	150	1.8	5.9
9	Human immunodeficiency virus (HIV) disease (B20-B24)	145	1.7	5.7
10	Influenza and pneumonia (J10-J18)	110	1.3	4.3
11	Septicemia (A40-A41)	109	1.3	4.3
12	Viral hepatitis (B15-B19)	106	1.2	4.2
13	Assault (homicide) (X85-Y09,Y87.1)	73	0.9	2.9
14	Intentional self-harm (suicide) (X60-X84,Y87.0)	63	0.7	2.5
15	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	55	0.6	2.2
...	All other causes (Residual)	1,222	14.3	48.2

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2000

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Population denominators for 2000 death rates are based on the 1990 census population.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
	Hispanic, female, 65 years and over			
...	All causes	30,881	100.0	2,741.5
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	10,406	33.7	923.8
2	Malignant neoplasms (C00-C97)	5,709	18.5	506.8
3	Cerebrovascular diseases (I60-I69)	2,638	8.5	234.2
4	Diabetes mellitus (E10-E14)	2,109	6.8	187.2
5	Influenza and pneumonia (J10-J18)	1,101	3.6	97.7
6	Chronic lower respiratory diseases (J40-J47)	1,033	3.3	91.7
7	Alzheimer's disease (G30)	732	2.4	65.0
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	634	2.1	56.3
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	474	1.5	42.1
10	Chronic liver disease and cirrhosis (K70,K73-K74)	436	1.4	38.7
11	Septicemia (A40-A41)	406	1.3	36.0
12	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	368	1.2	32.7
13	Atherosclerosis (I70)	231	0.7	20.5
14	Pneumonitis due to solids and liquids (J69)	206	0.7	18.3
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	184	0.6	16.3
...	All other causes (Residual)	4,214	13.6	374.1
	White, non-Hispanic, both sexes, all ages			
...	All causes	1,959,919	100.0	996.6
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	594,465	30.3	302.3
2	Malignant neoplasms (C00-C97)	458,200	23.4	233.0
3	Cerebrovascular diseases (I60-I69)	138,210	7.1	70.3
4	Chronic lower respiratory diseases (J40-J47)	109,901	5.6	55.9
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	73,536	3.8	37.4
6	Influenza and pneumonia (J10-J18)	55,135	2.8	28.0
7	Diabetes mellitus (E10-E14)	50,163	2.6	25.5
8	Alzheimer's disease (G30)	45,246	2.3	23.0
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	27,924	1.4	14.2
10	Intentional self-harm (suicide) (X60-X84,Y87.0)	24,616	1.3	12.5
11	Septicemia (A40-A41)	23,517	1.2	12.0
12	Chronic liver disease and cirrhosis (K70,K73-K74)	19,739	1.0	10.0
13	Parkinson's disease (G20-G21)	14,497	0.7	7.4
14	Pneumonitis due to solids and liquids (J69)	14,343	0.7	7.3
15	Aortic aneurysm and dissection (I71)	13,722	0.7	7.0
...	All other causes (Residual)	296,705	15.1	150.9

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2000

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Population denominators for 2000 death rates are based on the 1990 census population.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
White, non-Hispanic, both sexes, 1-4 years				
...	All causes	2,641	100.0	27.9
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,033	39.1	10.9
2	Malignant neoplasms (C00-C97)	256	9.7	2.7
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	240	9.1	2.5
4	Assault (homicide) (X85-Y09,Y87.1)	144	5.5	1.5
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	98	3.7	1.0
6	Septicemia (A40-A41)	51	1.9	0.5
6	Influenza and pneumonia (J10-J18)	51	1.9	0.5
8	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	37	1.4	0.4
9	Certain conditions originating in the perinatal period (P00-P96)	34	1.3	0.4
10	Cerebrovascular diseases (I60-I69)	25	0.9	0.3
11	Meningitis (G00,G03)	19	0.7	*
11	Chronic lower respiratory diseases (J40-J47)	19	0.7	*
13	Anemias (D50-D64)	14	0.5	*
14	Meningococcal infection (A39)	11	0.4	*
15	Complications of medical and surgical care (Y40-Y84,Y88)	8	0.3	*
...	All other causes (Residual)	601	22.8	6.4
White, non-Hispanic, both sexes, 5-14 years				
...	All causes	4,397	100.0	17.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,843	41.9	7.2
2	Malignant neoplasms (C00-C97)	627	14.3	2.5
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	240	5.5	0.9
4	Intentional self-harm (suicide) (X60-X84,Y87.0)	225	5.1	0.9
5	Assault (homicide) (X85-Y09,Y87.1)	164	3.7	0.6
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	140	3.2	0.5
7	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	63	1.4	0.2
8	Influenza and pneumonia (J10-J18)	49	1.1	0.2
9	Chronic lower respiratory diseases (J40-J47)	47	1.1	0.2
10	Cerebrovascular diseases (I60-I69)	46	1.0	0.2
11	Septicemia (A40-A41)	42	1.0	0.2
12	Certain conditions originating in the perinatal period (P00-P96)	20	0.5	0.1
13	Meningitis (G00,G03)	18	0.4	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	17	0.4	*
14	Complications of medical and surgical care (Y40-Y84,Y88)	17	0.4	*
...	All other causes (Residual)	839	19.1	3.3

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2000

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Population denominators for 2000 death rates are based on the 1990 census population.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
White, non-Hispanic, both sexes, 15-24 years				
...	All causes	18,261	100.0	72.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	9,735	53.3	38.4
2	Intentional self-harm (suicide) (X60-X84,Y87.0)	2,868	15.7	11.3
3	Malignant neoplasms (C00-C97)	1,059	5.8	4.2
4	Assault (homicide) (X85-Y09,Y87.1)	874	4.8	3.5
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	531	2.9	2.1
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	279	1.5	1.1
7	Cerebrovascular diseases (I60-I69)	118	0.6	0.5
8	Influenza and pneumonia (J10-J18)	116	0.6	0.5
9	Diabetes mellitus (E10-E14)	92	0.5	0.4
10	Chronic lower respiratory diseases (J40-J47)	87	0.5	0.3
11	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	63	0.3	0.2
12	Septicemia (A40-A41)	48	0.3	0.2
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	48	0.3	0.2
14	Meningococcal infection (A39)	37	0.2	0.1
15	Pregnancy, childbirth and the puerperium (O00-099)	33	0.2	0.1
...	All other causes (Residual)	2,273	12.4	9.0
White, non-Hispanic, both sexes, 25-44 years				
...	All causes	81,422	100.0	141.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	18,738	23.0	32.6
2	Malignant neoplasms (C00-C97)	14,057	17.3	24.5
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	10,396	12.8	18.1
4	Intentional self-harm (suicide) (X60-X84,Y87.0)	9,198	11.3	16.0
5	Human immunodeficiency virus (HIV) disease (B20-B24)	2,619	3.2	4.6
6	Chronic liver disease and cirrhosis (K70,K73-K74)	2,457	3.0	4.3
7	Assault (homicide) (X85-Y09,Y87.1)	2,296	2.8	4.0
8	Cerebrovascular diseases (I60-I69)	1,691	2.1	2.9
9	Diabetes mellitus (E10-E14)	1,617	2.0	2.8
10	Influenza and pneumonia (J10-J18)	891	1.1	1.6
11	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	763	0.9	1.3
12	Chronic lower respiratory diseases (J40-J47)	714	0.9	1.2
13	Septicemia (A40-A41)	632	0.8	1.1
14	Viral hepatitis (B15-B19)	538	0.7	0.9
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	469	0.6	0.8
...	All other causes (Residual)	14,346	17.6	25.0

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2000

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Population denominators for 2000 death rates are based on the 1990 census population.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
White, non-Hispanic, both sexes, 45-64 years				
...	All causes	294,312	100.0	619.9
1	Malignant neoplasms (C00-C97)	106,726	36.3	224.8
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	72,445	24.6	152.6
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	14,635	5.0	30.8
4	Chronic lower respiratory diseases (J40-J47)	11,716	4.0	24.7
5	Cerebrovascular diseases (I60-I69)	9,988	3.4	21.0
6	Diabetes mellitus (E10-E14)	9,001	3.1	19.0
7	Chronic liver disease and cirrhosis (K70,K73-K74)	8,872	3.0	18.7
8	Intentional self-harm (suicide) (X60-X84,Y87.0)	7,433	2.5	15.7
9	Influenza and pneumonia (J10-J18)	3,226	1.1	6.8
10	Septicemia (A40-A41)	3,011	1.0	6.3
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2,605	0.9	5.5
12	Viral hepatitis (B15-B19)	1,763	0.6	3.7
13	Human immunodeficiency virus (HIV) disease (B20-B24)	1,745	0.6	3.7
14	Aortic aneurysm and dissection (I71)	1,655	0.6	3.5
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1,520	0.5	3.2
...	All other causes (Residual)	37,971	12.9	80.0
White, non-Hispanic, both sexes, 65 years and over				
...	All causes	1,545,284	100.0	5,318.5
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	510,568	33.0	1,757.2
2	Malignant neoplasms (C00-C97)	335,408	21.7	1,154.4
3	Cerebrovascular diseases (I60-I69)	126,283	8.2	434.6
4	Chronic lower respiratory diseases (J40-J47)	97,301	6.3	334.9
5	Influenza and pneumonia (J10-J18)	50,687	3.3	174.5
6	Alzheimer's disease (G30)	44,760	2.9	154.1
7	Diabetes mellitus (E10-E14)	39,432	2.6	135.7
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	27,089	1.8	93.2
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	24,720	1.6	85.1
10	Septicemia (A40-A41)	19,624	1.3	67.5
11	Parkinson's disease (G20-G21)	14,217	0.9	48.9
12	Pneumonitis due to solids and liquids (J69)	13,419	0.9	46.2
13	Atherosclerosis (I70)	12,233	0.8	42.1
14	Aortic aneurysm and dissection (I71)	11,805	0.8	40.6
15	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	11,637	0.8	40.1
...	All other causes (Residual)	206,101	13.3	709.3

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2000

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Population denominators for 2000 death rates are based on the 1990 census population.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
	White, non-Hispanic, male, all ages			
...	All causes	944,781	100.0	980.9
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	287,210	30.4	298.2
2	Malignant neoplasms (C00-C97)	235,932	25.0	245.0
3	Chronic lower respiratory diseases (J40-J47)	53,231	5.6	55.3
4	Cerebrovascular diseases (I60-I69)	52,000	5.5	54.0
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	46,470	4.9	48.2
6	Influenza and pneumonia (J10-J18)	23,609	2.5	24.5
7	Diabetes mellitus (E10-E14)	23,465	2.5	24.4
8	Intentional self-harm (suicide) (X60-X84,Y87.0)	19,711	2.1	20.5
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	13,567	1.4	14.1
10	Alzheimer's disease (G30)	13,123	1.4	13.6
11	Chronic liver disease and cirrhosis (K70,K73-K74)	12,640	1.3	13.1
12	Septicemia (A40-A41)	10,056	1.1	10.4
13	Aortic aneurysm and dissection (I71)	8,488	0.9	8.8
14	Parkinson's disease (G20-G21)	8,116	0.9	8.4
15	Pneumonitis due to solids and liquids (J69)	7,214	0.8	7.5
...	All other causes (Residual)	129,949	13.8	134.9
	White, non-Hispanic, male, 1-4 years			
...	All causes	1,541	100.0	31.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	638	41.4	13.2
2	Malignant neoplasms (C00-C97)	146	9.5	3.0
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	124	8.0	2.6
4	Assault (homicide) (X85-Y09,Y87.1)	86	5.6	1.8
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	59	3.8	1.2
6	Septicemia (A40-A41)	26	1.7	0.5
7	Influenza and pneumonia (J10-J18)	24	1.6	0.5
8	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	23	1.5	0.5
8	Certain conditions originating in the perinatal period (P00-P96)	23	1.5	0.5
10	Cerebrovascular diseases (I60-I69)	14	0.9	*
11	Meningitis (G00,G03)	12	0.8	*
12	Anemias (D50-D64)	8	0.5	*
12	Chronic lower respiratory diseases (J40-J47)	8	0.5	*
14	Meningococcal infection (A39)	6	0.4	*
14	Complications of medical and surgical care (Y40-Y84,Y88)	6	0.4	*
...	All other causes (Residual)	338	21.9	7.0

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2000

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Population denominators for 2000 death rates are based on the 1990 census population.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
White, non-Hispanic, male, 5-14 years				
...	All causes	2,652	100.0	20.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,176	44.3	9.0
2	Malignant neoplasms (C00-C97)	338	12.7	2.6
3	Intentional self-harm (suicide) (X60-X84,Y87.0)	179	6.7	1.4
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	119	4.5	0.9
5	Assault (homicide) (X85-Y09,Y87.1)	93	3.5	0.7
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	87	3.3	0.7
7	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	31	1.2	0.2
8	Influenza and pneumonia (J10-J18)	29	1.1	0.2
9	Septicemia (A40-A41)	25	0.9	0.2
10	Chronic lower respiratory diseases (J40-J47)	24	0.9	0.2
11	Cerebrovascular diseases (I60-I69)	23	0.9	0.2
12	Certain conditions originating in the perinatal period (P00-P96)	14	0.5	*
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	12	0.5	*
14	Anemias (D50-D64)	9	0.3	*
14	Meningitis (G00,G03)	9	0.3	*
...	All other causes (Residual)	484	18.3	3.7
White, non-Hispanic, male, 15-24 years				
...	All causes	13,066	100.0	100.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	7,065	54.1	54.4
2	Intentional self-harm (suicide) (X60-X84,Y87.0)	2,467	18.9	19.0
3	Malignant neoplasms (C00-C97)	636	4.9	4.9
4	Assault (homicide) (X85-Y09,Y87.1)	588	4.5	4.5
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	326	2.5	2.5
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	163	1.2	1.3
7	Influenza and pneumonia (J10-J18)	68	0.5	0.5
8	Cerebrovascular diseases (I60-I69)	67	0.5	0.5
9	Diabetes mellitus (E10-E14)	57	0.4	0.4
10	Chronic lower respiratory diseases (J40-J47)	52	0.4	0.4
11	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	40	0.3	0.3
12	Septicemia (A40-A41)	33	0.3	0.3
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	25	0.2	0.2
13	Legal intervention (Y35,Y89.0)	25	0.2	0.2
15	Meningococcal infection (A39)	22	0.2	0.2
...	All other causes (Residual)	1,432	11.0	11.0

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2000

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Population denominators for 2000 death rates are based on the 1990 census population.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
	White, non-Hispanic, male, 25-44 years			
...	All causes	53,556	100.0	186.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	13,974	26.1	48.6
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	7,561	14.1	26.3
3	Intentional self-harm (suicide) (X60-X84,Y87.0)	7,276	13.6	25.3
4	Malignant neoplasms (C00-C97)	6,460	12.1	22.5
5	Human immunodeficiency virus (HIV) disease (B20-B24)	2,153	4.0	7.5
6	Chronic liver disease and cirrhosis (K70,K73-K74)	1,658	3.1	5.8
7	Assault (homicide) (X85-Y09,Y87.1)	1,513	2.8	5.3
8	Diabetes mellitus (E10-E14)	986	1.8	3.4
9	Cerebrovascular diseases (I60-I69)	855	1.6	3.0
10	Influenza and pneumonia (J10-J18)	505	0.9	1.8
11	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	436	0.8	1.5
12	Viral hepatitis (B15-B19)	361	0.7	1.3
13	Chronic lower respiratory diseases (J40-J47)	338	0.6	1.2
14	Septicemia (A40-A41)	336	0.6	1.2
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	268	0.5	0.9
...	All other causes (Residual)	8,876	16.6	30.9
	White, non-Hispanic, male, 45-64 years			
...	All causes	179,819	100.0	771.7
1	Malignant neoplasms (C00-C97)	56,460	31.4	242.3
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	52,280	29.1	224.4
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	10,392	5.8	44.6
4	Chronic liver disease and cirrhosis (K70,K73-K74)	6,412	3.6	27.5
5	Chronic lower respiratory diseases (J40-J47)	5,987	3.3	25.7
6	Intentional self-harm (suicide) (X60-X84,Y87.0)	5,658	3.1	24.3
7	Cerebrovascular diseases (I60-I69)	5,461	3.0	23.4
8	Diabetes mellitus (E10-E14)	5,140	2.9	22.1
9	Influenza and pneumonia (J10-J18)	1,843	1.0	7.9
10	Septicemia (A40-A41)	1,628	0.9	7.0
11	Human immunodeficiency virus (HIV) disease (B20-B24)	1,530	0.9	6.6
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,481	0.8	6.4
13	Viral hepatitis (B15-B19)	1,297	0.7	5.6
14	Aortic aneurysm and dissection (I71)	1,269	0.7	5.4
15	Assault (homicide) (X85-Y09,Y87.1)	848	0.5	3.6
...	All other causes (Residual)	22,133	12.3	95.0

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2000

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Population denominators for 2000 death rates are based on the 1990 census population.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
	White, non-Hispanic, male, 65 years and over			
...	All causes	686,417	100.0	5,660.3
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	226,738	33.0	1,869.7
2	Malignant neoplasms (C00-C97)	171,851	25.0	1,417.1
3	Chronic lower respiratory diseases (J40-J47)	46,809	6.8	386.0
4	Cerebrovascular diseases (I60-I69)	45,542	6.6	375.5
5	Influenza and pneumonia (J10-J18)	21,071	3.1	173.8
6	Diabetes mellitus (E10-E14)	17,271	2.5	142.4
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	12,952	1.9	106.8
8	Alzheimer's disease (G30)	12,910	1.9	106.5
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	11,738	1.7	96.8
10	Septicemia (A40-A41)	7,952	1.2	65.6
11	Parkinson's disease (G20-G21)	7,938	1.2	65.5
12	Aortic aneurysm and dissection (I71)	7,025	1.0	57.9
13	Pneumonitis due to solids and liquids (J69)	6,652	1.0	54.9
14	Chronic liver disease and cirrhosis (K70,K73-K74)	4,561	0.7	37.6
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	4,511	0.7	37.2
...	All other causes (Residual)	80,896	11.8	667.1
	White, non-Hispanic, female, all ages			
...	All causes	1,015,138	100.0	1,011.7
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	307,255	30.3	306.2
2	Malignant neoplasms (C00-C97)	222,268	21.9	221.5
3	Cerebrovascular diseases (I60-I69)	86,210	8.5	85.9
4	Chronic lower respiratory diseases (J40-J47)	56,670	5.6	56.5
5	Alzheimer's disease (G30)	32,123	3.2	32.0
6	Influenza and pneumonia (J10-J18)	31,526	3.1	31.4
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	27,066	2.7	27.0
8	Diabetes mellitus (E10-E14)	26,698	2.6	26.6
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	14,357	1.4	14.3
10	Septicemia (A40-A41)	13,461	1.3	13.4
11	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	8,196	0.8	8.2
12	Atherosclerosis (I70)	8,052	0.8	8.0
13	Pneumonitis due to solids and liquids (J69)	7,129	0.7	7.1
14	Chronic liver disease and cirrhosis (K70,K73-K74)	7,099	0.7	7.1
15	Parkinson's disease (G20-G21)	6,381	0.6	6.4
...	All other causes (Residual)	160,647	15.8	160.1

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2000

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Population denominators for 2000 death rates are based on the 1990 census population.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
	White, non-Hispanic, female, 1-4 years			
...	All causes	1,100	100.0	23.9
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	395	35.9	8.6
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	116	10.5	2.5
3	Malignant neoplasms (C00-C97)	110	10.0	2.4
4	Assault (homicide) (X85-Y09,Y87.1)	58	5.3	1.3
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	39	3.5	0.8
6	Influenza and pneumonia (J10-J18)	27	2.5	0.6
7	Septicemia (A40-A41)	25	2.3	0.5
8	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	14	1.3	*
9	Cerebrovascular diseases (I60-I69)	11	1.0	*
9	Chronic lower respiratory diseases (J40-J47)	11	1.0	*
9	Certain conditions originating in the perinatal period (P00-P96)	11	1.0	*
12	Meningitis (G00,G03)	7	0.6	*
13	Anemias (D50-D64)	6	0.5	*
14	Meningococcal infection (A39)	5	0.5	*
15	Diabetes mellitus (E10-E14)	4	0.4	*
...	All other causes (Residual)	261	23.7	5.7
	White, non-Hispanic, female, 5-14 years			
...	All causes	1,745	100.0	14.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	667	38.2	5.4
2	Malignant neoplasms (C00-C97)	289	16.6	2.3
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	121	6.9	1.0
4	Assault (homicide) (X85-Y09,Y87.1)	71	4.1	0.6
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	53	3.0	0.4
6	Intentional self-harm (suicide) (X60-X84,Y87.0)	46	2.6	0.4
7	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	32	1.8	0.3
8	Cerebrovascular diseases (I60-I69)	23	1.3	0.2
8	Chronic lower respiratory diseases (J40-J47)	23	1.3	0.2
10	Influenza and pneumonia (J10-J18)	20	1.1	0.2
11	Septicemia (A40-A41)	17	1.0	*
12	Complications of medical and surgical care (Y40-Y84,Y88)	10	0.6	*
13	Meningitis (G00,G03)	9	0.5	*
14	Human immunodeficiency virus (HIV) disease (B20-B24)	8	0.5	*
14	Diseases of appendix (K35-K38)	8	0.5	*
...	All other causes (Residual)	348	19.9	2.8

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2000

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Population denominators for 2000 death rates are based on the 1990 census population.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
White, non-Hispanic, female, 15-24 years				
...	All causes	5,195	100.0	42.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,670	51.4	21.7
2	Malignant neoplasms (C00-C97)	423	8.1	3.4
3	Intentional self-harm (suicide) (X60-X84,Y87.0)	401	7.7	3.3
4	Assault (homicide) (X85-Y09,Y87.1)	286	5.5	2.3
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	205	3.9	1.7
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	116	2.2	0.9
7	Cerebrovascular diseases (I60-I69)	51	1.0	0.4
8	Influenza and pneumonia (J10-J18)	48	0.9	0.4
9	Diabetes mellitus (E10-E14)	35	0.7	0.3
9	Chronic lower respiratory diseases (J40-J47)	35	0.7	0.3
11	Pregnancy, childbirth and the puerperium (O00-099)	33	0.6	0.3
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	23	0.4	0.2
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	23	0.4	0.2
14	Meningococcal infection (A39)	15	0.3	*
14	Septicemia (A40-A41)	15	0.3	*
...	All other causes (Residual)	816	15.7	6.6
White, non-Hispanic, female, 25-44 years				
...	All causes	27,866	100.0	97.0
1	Malignant neoplasms (C00-C97)	7,597	27.3	26.5
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	4,764	17.1	16.6
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,835	10.2	9.9
4	Intentional self-harm (suicide) (X60-X84,Y87.0)	1,922	6.9	6.7
5	Cerebrovascular diseases (I60-I69)	836	3.0	2.9
6	Chronic liver disease and cirrhosis (K70,K73-K74)	799	2.9	2.8
7	Assault (homicide) (X85-Y09,Y87.1)	783	2.8	2.7
8	Diabetes mellitus (E10-E14)	631	2.3	2.2
9	Human immunodeficiency virus (HIV) disease (B20-B24)	466	1.7	1.6
10	Influenza and pneumonia (J10-J18)	386	1.4	1.3
11	Chronic lower respiratory diseases (J40-J47)	376	1.3	1.3
12	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	327	1.2	1.1
13	Septicemia (A40-A41)	296	1.1	1.0
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	201	0.7	0.7
15	Viral hepatitis (B15-B19)	177	0.6	0.6
...	All other causes (Residual)	5,470	19.6	19.0

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2000

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Population denominators for 2000 death rates are based on the 1990 census population.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
White, non-Hispanic, female, 45-64 years				
...	All causes	114,493	100.0	473.5
1	Malignant neoplasms (C00-C97)	50,266	43.9	207.9
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	20,165	17.6	83.4
3	Chronic lower respiratory diseases (J40-J47)	5,729	5.0	23.7
4	Cerebrovascular diseases (I60-I69)	4,527	4.0	18.7
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	4,243	3.7	17.5
6	Diabetes mellitus (E10-E14)	3,861	3.4	16.0
7	Chronic liver disease and cirrhosis (K70,K73-K74)	2,460	2.1	10.2
8	Intentional self-harm (suicide) (X60-X84,Y87.0)	1,775	1.6	7.3
9	Septicemia (A40-A41)	1,383	1.2	5.7
9	Influenza and pneumonia (J10-J18)	1,383	1.2	5.7
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,124	1.0	4.6
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	680	0.6	2.8
13	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	468	0.4	1.9
14	Viral hepatitis (B15-B19)	466	0.4	1.9
15	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	428	0.4	1.8
...	All other causes (Residual)	15,535	13.6	64.3
White, non-Hispanic, female, 65 years and over				
...	All causes	858,867	100.0	5,073.6
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	283,830	33.0	1,676.7
2	Malignant neoplasms (C00-C97)	163,557	19.0	966.2
3	Cerebrovascular diseases (I60-I69)	80,741	9.4	477.0
4	Chronic lower respiratory diseases (J40-J47)	50,492	5.9	298.3
5	Alzheimer's disease (G30)	31,850	3.7	188.1
6	Influenza and pneumonia (J10-J18)	29,616	3.4	175.0
7	Diabetes mellitus (E10-E14)	22,161	2.6	130.9
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	14,137	1.6	83.5
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	12,982	1.5	76.7
10	Septicemia (A40-A41)	11,672	1.4	69.0
11	Atherosclerosis (I70)	7,895	0.9	46.6
12	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	7,715	0.9	45.6
13	Pneumonitis due to solids and liquids (J69)	6,767	0.8	40.0
14	Parkinson's disease (G20-G21)	6,279	0.7	37.1
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	5,041	0.6	29.8
...	All other causes (Residual)	124,132	14.5	733.3

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2000

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Population denominators for 2000 death rates are based on the 1990 census population.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
	Black, non-Hispanic, both sexes, all ages			
...	All causes	282,676	100.0	844.4
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	76,706	27.1	229.1
2	Malignant neoplasms (C00-C97)	61,390	21.7	183.4
3	Cerebrovascular diseases (I60-I69)	19,053	6.7	56.9
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	12,087	4.3	36.1
5	Diabetes mellitus (E10-E14)	11,918	4.2	35.6
6	Assault (homicide) (X85-Y09,Y87.1)	7,748	2.7	23.1
7	Human immunodeficiency virus (HIV) disease (B20-B24)	7,708	2.7	23.0
8	Chronic lower respiratory diseases (J40-J47)	7,528	2.7	22.5
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	6,853	2.4	20.5
10	Influenza and pneumonia (J10-J18)	5,920	2.1	17.7
11	Septicemia (A40-A41)	5,863	2.1	17.5
12	Certain conditions originating in the perinatal period (P00-P96)	4,878	1.7	14.6
13	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	3,862	1.4	11.5
14	Alzheimer's disease (G30)	2,719	1.0	8.1
15	Chronic liver disease and cirrhosis (K70,K73-K74)	2,705	1.0	8.1
...	All other causes (Residual)	45,738	16.2	136.6
	Black, non-Hispanic, both sexes, 1-4 years			
...	All causes	1,217	100.0	58.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	396	32.5	19.1
2	Assault (homicide) (X85-Y09,Y87.1)	143	11.8	6.9
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	123	10.1	5.9
4	Malignant neoplasms (C00-C97)	66	5.4	3.2
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	47	3.9	2.3
6	Influenza and pneumonia (J10-J18)	29	2.4	1.4
7	Certain conditions originating in the perinatal period (P00-P96)	27	2.2	1.3
8	Septicemia (A40-A41)	25	2.1	1.2
9	Chronic lower respiratory diseases (J40-J47)	21	1.7	1.0
10	Anemias (D50-D64)	19	1.6	*
11	Meningitis (G00,G03)	13	1.1	*
11	Cerebrovascular diseases (I60-I69)	13	1.1	*
13	Human immunodeficiency virus (HIV) disease (B20-B24)	12	1.0	*
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	9	0.7	*
15	Complications of medical and surgical care (Y40-Y84,Y88)	8	0.7	*
...	All other causes (Residual)	266	21.9	12.8

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2000

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Population denominators for 2000 death rates are based on the 1990 census population.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
	Black, non-Hispanic, both sexes, 5-14 years			
...	All causes	1,620	100.0	27.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	619	38.2	10.5
2	Malignant neoplasms (C00-C97)	147	9.1	2.5
3	Assault (homicide) (X85-Y09,Y87.1)	122	7.5	2.1
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	89	5.5	1.5
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	75	4.6	1.3
6	Chronic lower respiratory diseases (J40-J47)	70	4.3	1.2
7	Intentional self-harm (suicide) (X60-X84,Y87.0)	39	2.4	0.7
8	Anemias (D50-D64)	30	1.9	0.5
9	Human immunodeficiency virus (HIV) disease (B20-B24)	23	1.4	0.4
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	18	1.1	*
11	Cerebrovascular diseases (I60-I69)	15	0.9	*
12	Septicemia (A40-A41)	11	0.7	*
12	Influenza and pneumonia (J10-J18)	11	0.7	*
14	Certain conditions originating in the perinatal period (P00-P96)	10	0.6	*
15	Diabetes mellitus (E10-E14)	9	0.6	*
15	Meningitis (G00,G03)	9	0.6	*
...	All other causes (Residual)	323	19.9	5.5
	Black, non-Hispanic, both sexes, 15-24 years			
...	All causes	6,926	100.0	125.2
1	Assault (homicide) (X85-Y09,Y87.1)	2,771	40.0	50.1
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,662	24.0	30.0
3	Intentional self-harm (suicide) (X60-X84,Y87.0)	471	6.8	8.5
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	331	4.8	6.0
5	Malignant neoplasms (C00-C97)	297	4.3	5.4
6	Human immunodeficiency virus (HIV) disease (B20-B24)	125	1.8	2.3
7	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	91	1.3	1.6
8	Chronic lower respiratory diseases (J40-J47)	80	1.2	1.4
9	Anemias (D50-D64)	61	0.9	1.1
10	Diabetes mellitus (E10-E14)	55	0.8	1.0
11	Influenza and pneumonia (J10-J18)	44	0.6	0.8
12	Cerebrovascular diseases (I60-I69)	42	0.6	0.8
13	Pregnancy, childbirth and the puerperium (O00-099)	34	0.5	0.6
14	Septicemia (A40-A41)	33	0.5	0.6
15	Legal intervention (Y35,Y89.0)	27	0.4	0.5
...	All other causes (Residual)	802	11.6	14.5

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2000

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Population denominators for 2000 death rates are based on the 1990 census population.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Black, non-Hispanic, both sexes, 25-44 years				
...	All causes	30,382	100.0	293.3
1	Human immunodeficiency virus (HIV) disease (B20-B24)	4,442	14.6	42.9
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	4,188	13.8	40.4
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,993	13.1	38.5
4	Malignant neoplasms (C00-C97)	3,713	12.2	35.8
5	Assault (homicide) (X85-Y09,Y87.1)	3,552	11.7	34.3
6	Cerebrovascular diseases (I60-I69)	987	3.2	9.5
7	Intentional self-harm (suicide) (X60-X84,Y87.0)	896	2.9	8.6
8	Diabetes mellitus (E10-E14)	645	2.1	6.2
9	Chronic liver disease and cirrhosis (K70,K73-K74)	502	1.7	4.8
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	404	1.3	3.9
11	Chronic lower respiratory diseases (J40-J47)	386	1.3	3.7
12	Septicemia (A40-A41)	376	1.2	3.6
13	Influenza and pneumonia (J10-J18)	346	1.1	3.3
14	Anemias (D50-D64)	263	0.9	2.5
15	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	211	0.7	2.0
...	All other causes (Residual)	5,478	18.0	52.9
Black, non-Hispanic, both sexes, 45-64 years				
...	All causes	71,778	100.0	1,147.3
1	Malignant neoplasms (C00-C97)	20,087	28.0	321.1
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	18,863	26.3	301.5
3	Cerebrovascular diseases (I60-I69)	4,084	5.7	65.3
4	Diabetes mellitus (E10-E14)	3,265	4.5	52.2
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,944	4.1	47.1
6	Human immunodeficiency virus (HIV) disease (B20-B24)	2,836	4.0	45.3
7	Chronic lower respiratory diseases (J40-J47)	1,656	2.3	26.5
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,644	2.3	26.3
9	Chronic liver disease and cirrhosis (K70,K73-K74)	1,551	2.2	24.8
10	Septicemia (A40-A41)	1,328	1.9	21.2
11	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	960	1.3	15.3
12	Influenza and pneumonia (J10-J18)	954	1.3	15.2
13	Assault (homicide) (X85-Y09,Y87.1)	816	1.1	13.0
14	Viral hepatitis (B15-B19)	467	0.7	7.5
15	Intentional self-harm (suicide) (X60-X84,Y87.0)	367	0.5	5.9
...	All other causes (Residual)	9,956	13.9	159.1

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2000

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Population denominators for 2000 death rates are based on the 1990 census population.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
	Black, non-Hispanic, both sexes, 65 years and over			
...	All causes	162,179	100.0	5,759.3
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	53,072	32.7	1,884.7
2	Malignant neoplasms (C00-C97)	37,066	22.9	1,316.3
3	Cerebrovascular diseases (I60-I69)	13,865	8.5	492.4
4	Diabetes mellitus (E10-E14)	7,943	4.9	282.1
5	Chronic lower respiratory diseases (J40-J47)	5,304	3.3	188.4
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	4,703	2.9	167.0
7	Influenza and pneumonia (J10-J18)	4,435	2.7	157.5
8	Septicemia (A40-A41)	3,994	2.5	141.8
9	Alzheimer's disease (G30)	2,683	1.7	95.3
10	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	2,680	1.7	95.2
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,201	1.4	78.2
12	Pneumonitis due to solids and liquids (J69)	1,262	0.8	44.8
13	Atherosclerosis (I70)	908	0.6	32.2
14	Aortic aneurysm and dissection (I71)	808	0.5	28.7
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	729	0.4	25.9
...	All other causes (Residual)	20,526	12.7	728.9
	Black, non-Hispanic, male, all ages			
...	All causes	143,297	100.0	903.3
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	36,282	25.3	228.7
2	Malignant neoplasms (C00-C97)	32,493	22.7	204.8
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	8,380	5.8	52.8
4	Cerebrovascular diseases (I60-I69)	7,945	5.5	50.1
5	Assault (homicide) (X85-Y09,Y87.1)	6,382	4.5	40.2
6	Human immunodeficiency virus (HIV) disease (B20-B24)	5,291	3.7	33.4
7	Diabetes mellitus (E10-E14)	4,720	3.3	29.8
8	Chronic lower respiratory diseases (J40-J47)	4,184	2.9	26.4
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3,045	2.1	19.2
10	Influenza and pneumonia (J10-J18)	2,880	2.0	18.2
11	Certain conditions originating in the perinatal period (P00-P96)	2,760	1.9	17.4
12	Septicemia (A40-A41)	2,559	1.8	16.1
13	Chronic liver disease and cirrhosis (K70,K73-K74)	1,709	1.2	10.8
14	Intentional self-harm (suicide) (X60-X84,Y87.0)	1,601	1.1	10.1
15	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	1,576	1.1	9.9
...	All other causes (Residual)	21,490	15.0	135.5

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2000

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Population denominators for 2000 death rates are based on the 1990 census population.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Black, non-Hispanic, male, 1-4 years				
...	All causes	672	100.0	63.9
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	232	34.5	22.1
2	Assault (homicide) (X85-Y09,Y87.1)	73	10.9	6.9
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	67	10.0	6.4
4	Malignant neoplasms (C00-C97)	31	4.6	2.9
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	24	3.6	2.3
6	Influenza and pneumonia (J10-J18)	20	3.0	1.9
7	Certain conditions originating in the perinatal period (P00-P96)	18	2.7	*
8	Chronic lower respiratory diseases (J40-J47)	14	2.1	*
9	Septicemia (A40-A41)	13	1.9	*
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	8	1.2	*
10	Anemias (D50-D64)	8	1.2	*
12	Human immunodeficiency virus (HIV) disease (B20-B24)	6	0.9	*
12	Meningitis (G00,G03)	6	0.9	*
14	Meningococcal infection (A39)	4	0.6	*
15	Cerebrovascular diseases (I60-I69)	3	0.4	*
...	All other causes (Residual)	145	21.6	13.8
Black, non-Hispanic, male, 5-14 years				
...	All causes	959	100.0	32.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	406	42.3	13.6
2	Malignant neoplasms (C00-C97)	86	9.0	2.9
3	Assault (homicide) (X85-Y09,Y87.1)	70	7.3	2.3
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	50	5.2	1.7
5	Chronic lower respiratory diseases (J40-J47)	43	4.5	1.4
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	41	4.3	1.4
7	Intentional self-harm (suicide) (X60-X84,Y87.0)	34	3.5	1.1
8	Anemias (D50-D64)	17	1.8	*
9	Human immunodeficiency virus (HIV) disease (B20-B24)	12	1.3	*
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	8	0.8	*
11	Influenza and pneumonia (J10-J18)	7	0.7	*
12	Septicemia (A40-A41)	6	0.6	*
12	Meningitis (G00,G03)	6	0.6	*
14	Cerebrovascular diseases (I60-I69)	5	0.5	*
15	Diabetes mellitus (E10-E14)	4	0.4	*
...	All other causes (Residual)	164	17.1	5.5

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2000

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Population denominators for 2000 death rates are based on the 1990 census population.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Black, non-Hispanic, male, 15-24 years				
...	All causes	5,224	100.0	188.2
1	Assault (homicide) (X85-Y09,Y87.1)	2,459	47.1	88.6
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,233	23.6	44.4
3	Intentional self-harm (suicide) (X60-X84,Y87.0)	407	7.8	14.7
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	192	3.7	6.9
5	Malignant neoplasms (C00-C97)	178	3.4	6.4
6	Human immunodeficiency virus (HIV) disease (B20-B24)	60	1.1	2.2
7	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	58	1.1	2.1
8	Chronic lower respiratory diseases (J40-J47)	51	1.0	1.8
9	Anemias (D50-D64)	29	0.6	1.0
10	Diabetes mellitus (E10-E14)	28	0.5	1.0
11	Legal intervention (Y35,Y89.0)	27	0.5	1.0
12	Influenza and pneumonia (J10-J18)	21	0.4	0.8
13	Cerebrovascular diseases (I60-I69)	19	0.4	*
14	Septicemia (A40-A41)	18	0.3	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	10	0.2	*
...	All other causes (Residual)	434	8.3	15.6
Black, non-Hispanic, male, 25-44 years				
...	All causes	18,710	100.0	384.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,965	15.8	60.9
2	Assault (homicide) (X85-Y09,Y87.1)	2,910	15.6	59.7
3	Human immunodeficiency virus (HIV) disease (B20-B24)	2,882	15.4	59.1
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,515	13.4	51.6
5	Malignant neoplasms (C00-C97)	1,551	8.3	31.8
6	Intentional self-harm (suicide) (X60-X84,Y87.0)	743	4.0	15.2
7	Cerebrovascular diseases (I60-I69)	464	2.5	9.5
8	Diabetes mellitus (E10-E14)	350	1.9	7.2
9	Chronic liver disease and cirrhosis (K70,K73-K74)	305	1.6	6.3
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	225	1.2	4.6
11	Influenza and pneumonia (J10-J18)	209	1.1	4.3
12	Chronic lower respiratory diseases (J40-J47)	199	1.1	4.1
13	Septicemia (A40-A41)	184	1.0	3.8
14	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	114	0.6	2.3
15	Anemias (D50-D64)	110	0.6	2.3
...	All other causes (Residual)	2,984	15.9	61.2

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2000

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Population denominators for 2000 death rates are based on the 1990 census population.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
	Black, non-Hispanic, male, 45-64 years			
...	All causes	42,049	100.0	1,509.7
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	11,573	27.5	415.5
2	Malignant neoplasms (C00-C97)	10,915	26.0	391.9
3	Cerebrovascular diseases (I60-I69)	2,195	5.2	78.8
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,190	5.2	78.6
5	Human immunodeficiency virus (HIV) disease (B20-B24)	2,134	5.1	76.6
6	Diabetes mellitus (E10-E14)	1,600	3.8	57.4
7	Chronic liver disease and cirrhosis (K70,K73-K74)	1,041	2.5	37.4
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	878	2.1	31.5
9	Chronic lower respiratory diseases (J40-J47)	876	2.1	31.5
10	Septicemia (A40-A41)	700	1.7	25.1
11	Assault (homicide) (X85-Y09,Y87.1)	657	1.6	23.6
12	Influenza and pneumonia (J10-J18)	593	1.4	21.3
13	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	530	1.3	19.0
14	Viral hepatitis (B15-B19)	320	0.8	11.5
15	Intentional self-harm (suicide) (X60-X84,Y87.0)	290	0.7	10.4
...	All other causes (Residual)	5,557	13.2	199.5
	Black, non-Hispanic, male, 65 years and over			
...	All causes	70,891	100.0	6,400.7
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	21,871	30.9	1,974.7
2	Malignant neoplasms (C00-C97)	19,726	27.8	1,781.0
3	Cerebrovascular diseases (I60-I69)	5,232	7.4	472.4
4	Chronic lower respiratory diseases (J40-J47)	2,995	4.2	270.4
5	Diabetes mellitus (E10-E14)	2,737	3.9	247.1
6	Influenza and pneumonia (J10-J18)	1,975	2.8	178.3
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,888	2.7	170.5
8	Septicemia (A40-A41)	1,590	2.2	143.6
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,205	1.7	108.8
10	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	931	1.3	84.1
11	Alzheimer's disease (G30)	755	1.1	68.2
12	Pneumonitis due to solids and liquids (J69)	612	0.9	55.3
13	Aortic aneurysm and dissection (I71)	364	0.5	32.9
14	Chronic liver disease and cirrhosis (K70,K73-K74)	353	0.5	31.9
15	Atherosclerosis (I70)	333	0.5	30.1
...	All other causes (Residual)	8,324	11.7	751.6

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2000

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Population denominators for 2000 death rates are based on the 1990 census population.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Black, non-Hispanic, female, all ages				
...	All causes	139,379	100.0	791.4
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	40,424	29.0	229.5
2	Malignant neoplasms (C00-C97)	28,897	20.7	164.1
3	Cerebrovascular diseases (I60-I69)	11,108	8.0	63.1
4	Diabetes mellitus (E10-E14)	7,198	5.2	40.9
5	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3,808	2.7	21.6
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,707	2.7	21.0
7	Chronic lower respiratory diseases (J40-J47)	3,344	2.4	19.0
8	Septicemia (A40-A41)	3,304	2.4	18.8
9	Influenza and pneumonia (J10-J18)	3,040	2.2	17.3
10	Human immunodeficiency virus (HIV) disease (B20-B24)	2,417	1.7	13.7
11	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	2,286	1.6	13.0
12	Certain conditions originating in the perinatal period (P00-P96)	2,118	1.5	12.0
13	Alzheimer's disease (G30)	1,945	1.4	11.0
14	Assault (homicide) (X85-Y09,Y87.1)	1,366	1.0	7.8
15	Chronic liver disease and cirrhosis (K70,K73-K74)	996	0.7	5.7
...	All other causes (Residual)	23,421	16.8	133.0
Black, non-Hispanic, female, 1-4 years				
...	All causes	545	100.0	53.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	164	30.1	16.1
2	Assault (homicide) (X85-Y09,Y87.1)	70	12.8	6.9
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	56	10.3	5.5
4	Malignant neoplasms (C00-C97)	35	6.4	3.4
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	23	4.2	2.3
6	Septicemia (A40-A41)	12	2.2	*
7	Anemias (D50-D64)	11	2.0	*
8	Cerebrovascular diseases (I60-I69)	10	1.8	*
9	Influenza and pneumonia (J10-J18)	9	1.7	*
9	Certain conditions originating in the perinatal period (P00-P96)	9	1.7	*
11	Meningitis (G00,G03)	7	1.3	*
11	Chronic lower respiratory diseases (J40-J47)	7	1.3	*
13	Human immunodeficiency virus (HIV) disease (B20-B24)	6	1.1	*
13	Complications of medical and surgical care (Y40-Y84,Y88)	6	1.1	*
15	Acute bronchitis and bronchiolitis (J20-J21)	3	0.6	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3	0.6	*
...	All other causes (Residual)	114	20.9	11.2

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2000

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Population denominators for 2000 death rates are based on the 1990 census population.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Black, non-Hispanic, female, 5-14 years				
...	All causes	661	100.0	22.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	213	32.2	7.3
2	Malignant neoplasms (C00-C97)	61	9.2	2.1
3	Assault (homicide) (X85-Y09,Y87.1)	52	7.9	1.8
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	39	5.9	1.3
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	34	5.1	1.2
6	Chronic lower respiratory diseases (J40-J47)	27	4.1	0.9
7	Anemias (D50-D64)	13	2.0	*
8	Human immunodeficiency virus (HIV) disease (B20-B24)	11	1.7	*
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	10	1.5	*
9	Cerebrovascular diseases (I60-I69)	10	1.5	*
11	Certain conditions originating in the perinatal period (P00-P96)	7	1.1	*
12	Meningococcal infection (A39)	5	0.8	*
12	Septicemia (A40-A41)	5	0.8	*
12	Diabetes mellitus (E10-E14)	5	0.8	*
12	Intentional self-harm (suicide) (X60-X84,Y87.0)	5	0.8	*
...	All other causes (Residual)	164	24.8	5.6
Black, non-Hispanic, female, 15-24 years				
...	All causes	1,702	100.0	61.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	429	25.2	15.6
2	Assault (homicide) (X85-Y09,Y87.1)	312	18.3	11.3
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	139	8.2	5.0
4	Malignant neoplasms (C00-C97)	119	7.0	4.3
5	Human immunodeficiency virus (HIV) disease (B20-B24)	65	3.8	2.4
6	Intentional self-harm (suicide) (X60-X84,Y87.0)	64	3.8	2.3
7	Pregnancy, childbirth and the puerperium (O00-099)	34	2.0	1.2
8	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	33	1.9	1.2
9	Anemias (D50-D64)	32	1.9	1.2
10	Chronic lower respiratory diseases (J40-J47)	29	1.7	1.1
11	Diabetes mellitus (E10-E14)	27	1.6	1.0
12	Cerebrovascular diseases (I60-I69)	23	1.4	0.8
12	Influenza and pneumonia (J10-J18)	23	1.4	0.8
14	Septicemia (A40-A41)	15	0.9	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	14	0.8	*
...	All other causes (Residual)	344	20.2	12.5

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2000

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Population denominators for 2000 death rates are based on the 1990 census population.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
	Black, non-Hispanic, female, 25-44 years			
...	All causes	11,672	100.0	212.8
1	Malignant neoplasms (C00-C97)	2,162	18.5	39.4
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,673	14.3	30.5
3	Human immunodeficiency virus (HIV) disease (B20-B24)	1,560	13.4	28.4
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,028	8.8	18.7
5	Assault (homicide) (X85-Y09,Y87.1)	642	5.5	11.7
6	Cerebrovascular diseases (I60-I69)	523	4.5	9.5
7	Diabetes mellitus (E10-E14)	295	2.5	5.4
8	Chronic liver disease and cirrhosis (K70,K73-K74)	197	1.7	3.6
9	Septicemia (A40-A41)	192	1.6	3.5
10	Chronic lower respiratory diseases (J40-J47)	187	1.6	3.4
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	179	1.5	3.3
12	Anemias (D50-D64)	153	1.3	2.8
12	Intentional self-harm (suicide) (X60-X84,Y87.0)	153	1.3	2.8
14	Influenza and pneumonia (J10-J18)	137	1.2	2.5
15	Pregnancy, childbirth and the puerperium (O00-099)	102	0.9	1.9
...	All other causes (Residual)	2,489	21.3	45.4
	Black, non-Hispanic, female, 45-64 years			
...	All causes	29,729	100.0	856.4
1	Malignant neoplasms (C00-C97)	9,172	30.9	264.2
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	7,290	24.5	210.0
3	Cerebrovascular diseases (I60-I69)	1,889	6.4	54.4
4	Diabetes mellitus (E10-E14)	1,665	5.6	48.0
5	Chronic lower respiratory diseases (J40-J47)	780	2.6	22.5
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	766	2.6	22.1
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	754	2.5	21.7
8	Human immunodeficiency virus (HIV) disease (B20-B24)	702	2.4	20.2
9	Septicemia (A40-A41)	628	2.1	18.1
10	Chronic liver disease and cirrhosis (K70,K73-K74)	510	1.7	14.7
11	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	430	1.4	12.4
12	Influenza and pneumonia (J10-J18)	361	1.2	10.4
13	Assault (homicide) (X85-Y09,Y87.1)	159	0.5	4.6
14	Viral hepatitis (B15-B19)	147	0.5	4.2
15	Aortic aneurysm and dissection (I71)	119	0.4	3.4
...	All other causes (Residual)	4,357	14.7	125.5

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2000

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Population denominators for 2000 death rates are based on the 1990 census population.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
	Black, non-Hispanic, female, 65 years and over			
...	All causes	91,288	100.0	5,343.5
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	31,201	34.2	1,826.3
2	Malignant neoplasms (C00-C97)	17,340	19.0	1,015.0
3	Cerebrovascular diseases (I60-I69)	8,633	9.5	505.3
4	Diabetes mellitus (E10-E14)	5,206	5.7	304.7
5	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2,815	3.1	164.8
6	Influenza and pneumonia (J10-J18)	2,460	2.7	144.0
7	Septicemia (A40-A41)	2,404	2.6	140.7
8	Chronic lower respiratory diseases (J40-J47)	2,309	2.5	135.2
9	Alzheimer's disease (G30)	1,928	2.1	112.9
10	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	1,749	1.9	102.4
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	996	1.1	58.3
12	Pneumonitis due to solids and liquids (J69)	650	0.7	38.0
13	Atherosclerosis (I70)	575	0.6	33.7
14	Aortic aneurysm and dissection (I71)	444	0.5	26.0
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	416	0.5	24.4
...	All other causes (Residual)	12,162	13.3	711.9
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,167	1.3	69.0
12	Pneumonitis due to solids and liquids (J69)	632	0.7	37.3
13	Atherosclerosis (I70)	582	0.6	34.4
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	439	0.5	25.9
15	Aortic aneurysm and dissection (I71)	427	0.5	25.2
...	All other causes (Residual)	11,751	13.0	694.4

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable