

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2006

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. The title for Essential hypertension and hypertensive renal disease was changed in 2006 to reflect the addition of Secondary hypertension (ICD code I15); see the "Technical Notes" of *Deaths: Final Data for 2006*]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, both sexes, all ages				
...	All causes	133,004	100.0	300.1
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	28,921	21.7	65.3
2	Malignant neoplasms (C00-C97)	26,633	20.0	60.1
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	12,052	9.1	27.2
4	Cerebrovascular diseases (I60-I69)	7,005	5.3	15.8
5	Diabetes mellitus (E10-E14)	6,287	4.7	14.2
6	Chronic liver disease and cirrhosis (K70,K73-K74)	3,592	2.7	8.1
7	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	3,524	2.6	8.0
8	Chronic lower respiratory diseases (J40-J47)	3,310	2.5	7.5
9	Influenza and pneumonia (J10-J18)	2,966	2.2	6.7
10	Certain conditions originating in the perinatal period (P00-P96)	2,804	2.1	6.3
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2,592	1.9	5.8
12	Alzheimer's disease (G30)	2,399	1.8	5.4
13	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,177	1.6	4.9
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	1,952	1.5	4.4
15	Septicemia (A40-A41)	1,829	1.4	4.1
...	All other causes (Residual)	24,961	18.8	56.3
Hispanic, both sexes, 1-4 years				
...	All causes	987	100.0	26.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	361	36.6	9.7
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	125	12.7	3.3
3	Malignant neoplasms (C00-C97)	82	8.3	2.2
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	62	6.3	1.7
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	31	3.1	0.8
6	Influenza and pneumonia (J10-J18)	28	2.8	0.7
7	Septicemia (A40-A41)	18	1.8	*
8	Certain conditions originating in the perinatal period (P00-P96)	12	1.2	*
9	Chronic lower respiratory diseases (J40-J47)	11	1.1	*
10	Cerebrovascular diseases (I60-I69)	9	0.9	*
11	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	7	0.7	*
12	Meningitis (G00,G03)	6	0.6	*
13	Anemias (D50-D64)	4	0.4	*
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	4	0.4	*
13	Complications of medical and surgical care (Y40-Y84,Y88)	4	0.4	*
...	All other causes (Residual)	223	22.6	6.0

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2006

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. The title for Essential hypertension and hypertensive renal disease was changed in 2006 to reflect the addition of Secondary hypertension (ICD code I15); see the "Technical Notes" of *Deaths: Final Data for 2006*]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, both sexes, 5-14 years				
...	All causes	1,137	100.0	14.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	412	36.2	5.1
2	Malignant neoplasms (C00-C97)	183	16.1	2.3
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	95	8.4	1.2
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	61	5.4	0.8
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	45	4.0	0.6
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	25	2.2	0.3
7	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	18	1.6	*
8	Cerebrovascular diseases (I60-I69)	17	1.5	*
9	Septicemia (A40-A41)	16	1.4	*
9	Chronic lower respiratory diseases (J40-J47)	16	1.4	*
11	Influenza and pneumonia (J10-J18)	15	1.3	*
12	Certain conditions originating in the perinatal period (P00-P96)	10	0.9	*
13	Diabetes mellitus (E10-E14)	6	0.5	*
14	Anemias (D50-D64)	3	0.3	*
14	Pneumonitis due to solids and liquids (J69)	3	0.3	*
14	Diseases of appendix (K35-K38)	3	0.3	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3	0.3	*
...	All other causes (Residual)	206	18.1	2.6
Hispanic, both sexes, 15-24 years				
...	All causes	5,932	100.0	80.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,662	44.9	36.1
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,343	22.6	18.2
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	543	9.2	7.4
4	Malignant neoplasms (C00-C97)	321	5.4	4.4
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	130	2.2	1.8
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	71	1.2	1.0
7	Cerebrovascular diseases (I60-I69)	52	0.9	0.7
8	Pregnancy, childbirth and the puerperium (O00-099)	39	0.7	0.5
9	Human immunodeficiency virus (HIV) disease (B20-B24)	31	0.5	0.4
10	Legal intervention (Y35,Y89.0)	25	0.4	0.3
11	Influenza and pneumonia (J10-J18)	24	0.4	0.3
12	Septicemia (A40-A41)	22	0.4	0.3
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	22	0.4	0.3
14	Diabetes mellitus (E10-E14)	18	0.3	*
15	Chronic lower respiratory diseases (J40-J47)	15	0.3	*
...	All other causes (Residual)	614	10.4	8.3

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2006

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. The title for Essential hypertension and hypertensive renal disease was changed in 2006 to reflect the addition of Secondary hypertension (ICD code I15); see the "Technical Notes" of *Deaths: Final Data for 2006*]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, both sexes, 25-44 years				
...	All causes	15,543	100.0	105.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	4,502	29.0	30.5
2	Malignant neoplasms (C00-C97)	2,019	13.0	13.7
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,532	9.9	10.4
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,304	8.4	8.8
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,015	6.5	6.9
6	Human immunodeficiency virus (HIV) disease (B20-B24)	779	5.0	5.3
7	Chronic liver disease and cirrhosis (K70,K73-K74)	529	3.4	3.6
8	Cerebrovascular diseases (I60-I69)	402	2.6	2.7
9	Diabetes mellitus (E10-E14)	259	1.7	1.8
10	Influenza and pneumonia (J10-J18)	126	0.8	0.9
11	Septicemia (A40-A41)	120	0.8	0.8
12	Viral hepatitis (B15-B19)	118	0.8	0.8
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	104	0.7	0.7
13	Pregnancy, childbirth and the puerperium (O00-099)	104	0.7	0.7
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	98	0.6	0.7
...	All other causes (Residual)	2,532	16.3	17.2
Hispanic, both sexes, 45-64 years				
...	All causes	30,740	100.0	435.6
1	Malignant neoplasms (C00-C97)	8,418	27.4	119.3
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	5,949	19.4	84.3
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,453	8.0	34.8
4	Chronic liver disease and cirrhosis (K70,K73-K74)	1,942	6.3	27.5
5	Diabetes mellitus (E10-E14)	1,676	5.5	23.7
6	Cerebrovascular diseases (I60-I69)	1,521	4.9	21.6
7	Viral hepatitis (B15-B19)	752	2.4	10.7
8	Human immunodeficiency virus (HIV) disease (B20-B24)	721	2.3	10.2
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	588	1.9	8.3
10	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	447	1.5	6.3
11	Septicemia (A40-A41)	430	1.4	6.1
12	Chronic lower respiratory diseases (J40-J47)	423	1.4	6.0
13	Influenza and pneumonia (J10-J18)	414	1.3	5.9
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	357	1.2	5.1
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	287	0.9	4.1
...	All other causes (Residual)	4,362	14.2	61.8

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2006

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. The title for Essential hypertension and hypertensive renal disease was changed in 2006 to reflect the addition of Secondary hypertension (ICD code I15); see the "Technical Notes" of *Deaths: Final Data for 2006*]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, both sexes, 65 years and over				
...	All causes	72,894	100.0	3,038.1
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	21,396	29.4	891.8
2	Malignant neoplasms (C00-C97)	15,591	21.4	649.8
3	Cerebrovascular diseases (I60-I69)	4,970	6.8	207.1
4	Diabetes mellitus (E10-E14)	4,326	5.9	180.3
5	Chronic lower respiratory diseases (J40-J47)	2,762	3.8	115.1
6	Alzheimer's disease (G30)	2,335	3.2	97.3
7	Influenza and pneumonia (J10-J18)	2,299	3.2	95.8
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,845	2.5	76.9
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,457	2.0	60.7
10	Septicemia (A40-A41)	1,156	1.6	48.2
11	Chronic liver disease and cirrhosis (K70,K73-K74)	1,112	1.5	46.3
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	917	1.3	38.2
13	Parkinson's disease (G20-G21)	681	0.9	28.4
14	Pneumonitis due to solids and liquids (J69)	475	0.7	19.8
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	438	0.6	18.3
...	All other causes (Residual)	11,134	15.3	464.0
Hispanic, male, all ages				
...	All causes	74,250	100.0	323.9
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	15,518	20.9	67.7
2	Malignant neoplasms (C00-C97)	13,856	18.7	60.4
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	9,102	12.3	39.7
4	Cerebrovascular diseases (I60-I69)	3,269	4.4	14.3
5	Diabetes mellitus (E10-E14)	3,140	4.2	13.7
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	3,004	4.0	13.1
7	Chronic liver disease and cirrhosis (K70,K73-K74)	2,527	3.4	11.0
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,813	2.4	7.9
9	Chronic lower respiratory diseases (J40-J47)	1,708	2.3	7.5
10	Certain conditions originating in the perinatal period (P00-P96)	1,565	2.1	6.8
11	Influenza and pneumonia (J10-J18)	1,439	1.9	6.3
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,295	1.7	5.6
13	Human immunodeficiency virus (HIV) disease (B20-B24)	1,275	1.7	5.6
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	1,046	1.4	4.6
15	Septicemia (A40-A41)	932	1.3	4.1
...	All other causes (Residual)	12,761	17.2	55.7

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2006

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. The title for Essential hypertension and hypertensive renal disease was changed in 2006 to reflect the addition of Secondary hypertension (ICD code I15); see the "Technical Notes" of *Deaths: Final Data for 2006*]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, male, 1-4 years				
...	All causes	549	100.0	28.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	215	39.2	11.3
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	69	12.6	3.6
3	Malignant neoplasms (C00-C97)	49	8.9	2.6
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	35	6.4	1.8
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	19	3.5	*
6	Influenza and pneumonia (J10-J18)	13	2.4	*
7	Chronic lower respiratory diseases (J40-J47)	7	1.3	*
8	Septicemia (A40-A41)	6	1.1	*
8	Cerebrovascular diseases (I60-I69)	6	1.1	*
8	Certain conditions originating in the perinatal period (P00-P96)	6	1.1	*
11	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	3	0.5	*
12	Meningococcal infection (A39)	2	0.4	*
12	Anemias (D50-D64)	2	0.4	*
12	Meningitis (G00,G03)	2	0.4	*
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2	0.4	*
12	Complications of medical and surgical care (Y40-Y84,Y88)	2	0.4	*
...	All other causes (Residual)	111	20.2	5.8
Hispanic, male, 5-14 years				
...	All causes	673	100.0	16.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	259	38.5	6.3
2	Malignant neoplasms (C00-C97)	95	14.1	2.3
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	63	9.4	1.5
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	34	5.1	0.8
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	27	4.0	0.7
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	18	2.7	*
7	Chronic lower respiratory diseases (J40-J47)	13	1.9	*
8	Cerebrovascular diseases (I60-I69)	12	1.8	*
9	Septicemia (A40-A41)	10	1.5	*
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	9	1.3	*
11	Certain conditions originating in the perinatal period (P00-P96)	7	1.0	*
12	Influenza and pneumonia (J10-J18)	6	0.9	*
13	Diabetes mellitus (E10-E14)	3	0.4	*
13	Pneumonitis due to solids and liquids (J69)	3	0.4	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2	0.3	*
...	All other causes (Residual)	112	16.6	2.7

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2006

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. The title for Essential hypertension and hypertensive renal disease was changed in 2006 to reflect the addition of Secondary hypertension (ICD code I15); see the "Technical Notes" of *Deaths: Final Data for 2006*]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, male, 15-24 years				
...	All causes	4,712	100.0	120.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,169	46.0	55.5
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,210	25.7	31.0
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	452	9.6	11.6
4	Malignant neoplasms (C00-C97)	201	4.3	5.1
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	99	2.1	2.5
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	48	1.0	1.2
7	Cerebrovascular diseases (I60-I69)	31	0.7	0.8
8	Legal intervention (Y35,Y89.0)	25	0.5	0.6
9	Human immunodeficiency virus (HIV) disease (B20-B24)	19	0.4	*
10	Diabetes mellitus (E10-E14)	14	0.3	*
11	Influenza and pneumonia (J10-J18)	13	0.3	*
12	Septicemia (A40-A41)	11	0.2	*
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	10	0.2	*
13	Chronic lower respiratory diseases (J40-J47)	10	0.2	*
15	Meningitis (G00,G03)	6	0.1	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	6	0.1	*
...	All other causes (Residual)	388	8.2	9.9
Hispanic, male, 25-44 years				
...	All causes	11,247	100.0	140.9
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,691	32.8	46.2
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,320	11.7	16.5
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,000	8.9	12.5
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	859	7.6	10.8
5	Malignant neoplasms (C00-C97)	835	7.4	10.5
6	Human immunodeficiency virus (HIV) disease (B20-B24)	608	5.4	7.6
7	Chronic liver disease and cirrhosis (K70,K73-K74)	414	3.7	5.2
8	Cerebrovascular diseases (I60-I69)	254	2.3	3.2
9	Diabetes mellitus (E10-E14)	154	1.4	1.9
10	Viral hepatitis (B15-B19)	98	0.9	1.2
11	Influenza and pneumonia (J10-J18)	83	0.7	1.0
12	Septicemia (A40-A41)	65	0.6	0.8
13	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	63	0.6	0.8
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	62	0.6	0.8
15	Legal intervention (Y35,Y89.0)	55	0.5	0.7
...	All other causes (Residual)	1,686	15.0	21.1

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2006

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. The title for Essential hypertension and hypertensive renal disease was changed in 2006 to reflect the addition of Secondary hypertension (ICD code I15); see the "Technical Notes" of *Deaths: Final Data for 2006*]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, male, 45-64 years				
...	All causes	19,505	100.0	556.6
1	Malignant neoplasms (C00-C97)	4,370	22.4	124.7
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	4,192	21.5	119.6
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,859	9.5	53.0
4	Chronic liver disease and cirrhosis (K70,K73-K74)	1,514	7.8	43.2
5	Diabetes mellitus (E10-E14)	978	5.0	27.9
6	Cerebrovascular diseases (I60-I69)	890	4.6	25.4
7	Human immunodeficiency virus (HIV) disease (B20-B24)	583	3.0	16.6
8	Viral hepatitis (B15-B19)	533	2.7	15.2
9	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	360	1.8	10.3
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	333	1.7	9.5
11	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	291	1.5	8.3
12	Influenza and pneumonia (J10-J18)	255	1.3	7.3
13	Septicemia (A40-A41)	250	1.3	7.1
14	Chronic lower respiratory diseases (J40-J47)	234	1.2	6.7
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	165	0.8	4.7
...	All other causes (Residual)	2,698	13.8	77.0
Hispanic, male, 65 years and over				
...	All causes	34,356	100.0	3,363.3
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	10,144	29.5	993.1
2	Malignant neoplasms (C00-C97)	8,295	24.1	812.0
3	Cerebrovascular diseases (I60-I69)	2,059	6.0	201.6
4	Diabetes mellitus (E10-E14)	1,989	5.8	194.7
5	Chronic lower respiratory diseases (J40-J47)	1,400	4.1	137.1
6	Influenza and pneumonia (J10-J18)	1,033	3.0	101.1
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	871	2.5	85.3
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	792	2.3	77.5
9	Alzheimer's disease (G30)	779	2.3	76.3
10	Chronic liver disease and cirrhosis (K70,K73-K74)	590	1.7	57.8
11	Septicemia (A40-A41)	553	1.6	54.1
12	Parkinson's disease (G20-G21)	408	1.2	39.9
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	357	1.0	34.9
14	Pneumonitis due to solids and liquids (J69)	223	0.6	21.8
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	212	0.6	20.8
...	All other causes (Residual)	4,651	13.5	455.3

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2006

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. The title for Essential hypertension and hypertensive renal disease was changed in 2006 to reflect the addition of Secondary hypertension (ICD code I15); see the "Technical Notes" of *Deaths: Final Data for 2006*]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, female, all ages				
...	All causes	58,754	100.0	274.6
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	13,403	22.8	62.6
2	Malignant neoplasms (C00-C97)	12,777	21.7	59.7
3	Cerebrovascular diseases (I60-I69)	3,736	6.4	17.5
4	Diabetes mellitus (E10-E14)	3,147	5.4	14.7
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,950	5.0	13.8
6	Chronic lower respiratory diseases (J40-J47)	1,602	2.7	7.5
7	Alzheimer's disease (G30)	1,587	2.7	7.4
8	Influenza and pneumonia (J10-J18)	1,527	2.6	7.1
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,297	2.2	6.1
10	Certain conditions originating in the perinatal period (P00-P96)	1,239	2.1	5.8
11	Chronic liver disease and cirrhosis (K70,K73-K74)	1,065	1.8	5.0
12	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	906	1.5	4.2
13	Septicemia (A40-A41)	897	1.5	4.2
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	697	1.2	3.3
15	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	520	0.9	2.4
...	All other causes (Residual)	11,404	19.4	53.3
Hispanic, female, 1-4 years				
...	All causes	438	100.0	24.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	146	33.3	8.0
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	56	12.8	3.1
3	Malignant neoplasms (C00-C97)	33	7.5	1.8
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	27	6.2	1.5
5	Influenza and pneumonia (J10-J18)	15	3.4	*
6	Septicemia (A40-A41)	12	2.7	*
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	12	2.7	*
8	Certain conditions originating in the perinatal period (P00-P96)	6	1.4	*
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	4	0.9	*
9	Meningitis (G00,G03)	4	0.9	*
9	Chronic lower respiratory diseases (J40-J47)	4	0.9	*
12	Cerebrovascular diseases (I60-I69)	3	0.7	*
13	Anemias (D50-D64)	2	0.5	*
13	Acute bronchitis and bronchiolitis (J20-J21)	2	0.5	*
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2	0.5	*
13	Complications of medical and surgical care (Y40-Y84,Y88)	2	0.5	*
...	All other causes (Residual)	108	24.7	5.9

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2006

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. The title for Essential hypertension and hypertensive renal disease was changed in 2006 to reflect the addition of Secondary hypertension (ICD code I15); see the "Technical Notes" of *Deaths: Final Data for 2006*]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, female, 5-14 years				
...	All causes	464	100.0	11.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	153	33.0	3.9
2	Malignant neoplasms (C00-C97)	88	19.0	2.2
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	32	6.9	0.8
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	27	5.8	0.7
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	18	3.9	*
6	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	9	1.9	*
6	Influenza and pneumonia (J10-J18)	9	1.9	*
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	7	1.5	*
9	Septicemia (A40-A41)	6	1.3	*
10	Cerebrovascular diseases (I60-I69)	5	1.1	*
11	Diabetes mellitus (E10-E14)	3	0.6	*
11	Chronic lower respiratory diseases (J40-J47)	3	0.6	*
11	Certain conditions originating in the perinatal period (P00-P96)	3	0.6	*
14	Meningococcal infection (A39)	2	0.4	*
14	Human immunodeficiency virus (HIV) disease (B20-B24)	2	0.4	*
14	Anemias (D50-D64)	2	0.4	*
14	Diseases of appendix (K35-K38)	2	0.4	*
...	All other causes (Residual)	93	20.0	2.4
Hispanic, female, 15-24 years				
...	All causes	1,220	100.0	35.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	493	40.4	14.2
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	133	10.9	3.8
3	Malignant neoplasms (C00-C97)	120	9.8	3.5
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	91	7.5	2.6
5	Pregnancy, childbirth and the puerperium (O00-099)	39	3.2	1.1
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	31	2.5	0.9
7	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	23	1.9	0.7
8	Cerebrovascular diseases (I60-I69)	21	1.7	0.6
9	Human immunodeficiency virus (HIV) disease (B20-B24)	12	1.0	*
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	12	1.0	*
11	Septicemia (A40-A41)	11	0.9	*
11	Influenza and pneumonia (J10-J18)	11	0.9	*
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	6	0.5	*
14	Chronic lower respiratory diseases (J40-J47)	5	0.4	*
14	Complications of medical and surgical care (Y40-Y84,Y88)	5	0.4	*
...	All other causes (Residual)	207	17.0	6.0

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2006

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. The title for Essential hypertension and hypertensive renal disease was changed in 2006 to reflect the addition of Secondary hypertension (ICD code I15); see the "Technical Notes" of *Deaths: Final Data for 2006*]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, female, 25-44 years				
...	All causes	4,296	100.0	63.5
1	Malignant neoplasms (C00-C97)	1,184	27.6	17.5
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	811	18.9	12.0
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	304	7.1	4.5
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	212	4.9	3.1
5	Human immunodeficiency virus (HIV) disease (B20-B24)	171	4.0	2.5
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	156	3.6	2.3
7	Cerebrovascular diseases (I60-I69)	148	3.4	2.2
8	Chronic liver disease and cirrhosis (K70,K73-K74)	115	2.7	1.7
9	Diabetes mellitus (E10-E14)	105	2.4	1.6
10	Pregnancy, childbirth and the puerperium (O00-099)	104	2.4	1.5
11	Septicemia (A40-A41)	55	1.3	0.8
12	Influenza and pneumonia (J10-J18)	43	1.0	0.6
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	42	1.0	0.6
14	Chronic lower respiratory diseases (J40-J47)	37	0.9	0.5
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	35	0.8	0.5
...	All other causes (Residual)	774	18.0	11.4
Hispanic, female, 45-64 years				
...	All causes	11,235	100.0	316.2
1	Malignant neoplasms (C00-C97)	4,048	36.0	113.9
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,757	15.6	49.5
3	Diabetes mellitus (E10-E14)	698	6.2	19.6
4	Cerebrovascular diseases (I60-I69)	631	5.6	17.8
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	594	5.3	16.7
6	Chronic liver disease and cirrhosis (K70,K73-K74)	428	3.8	12.0
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	255	2.3	7.2
8	Viral hepatitis (B15-B19)	219	1.9	6.2
9	Chronic lower respiratory diseases (J40-J47)	189	1.7	5.3
10	Septicemia (A40-A41)	180	1.6	5.1
11	Influenza and pneumonia (J10-J18)	159	1.4	4.5
12	Human immunodeficiency virus (HIV) disease (B20-B24)	138	1.2	3.9
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	122	1.1	3.4
14	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	87	0.8	2.4
15	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	66	0.6	1.9
...	All other causes (Residual)	1,664	14.8	46.8

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2006

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. The title for Essential hypertension and hypertensive renal disease was changed in 2006 to reflect the addition of Secondary hypertension (ICD code I15); see the "Technical Notes" of *Deaths: Final Data for 2006*]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, female, 65 years and over				
...	All causes	38,538	100.0	2,797.0
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	11,252	29.2	816.6
2	Malignant neoplasms (C00-C97)	7,296	18.9	529.5
3	Cerebrovascular diseases (I60-I69)	2,911	7.6	211.3
4	Diabetes mellitus (E10-E14)	2,337	6.1	169.6
5	Alzheimer's disease (G30)	1,556	4.0	112.9
6	Chronic lower respiratory diseases (J40-J47)	1,362	3.5	98.9
7	Influenza and pneumonia (J10-J18)	1,266	3.3	91.9
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	974	2.5	70.7
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	665	1.7	48.3
10	Septicemia (A40-A41)	603	1.6	43.8
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	560	1.5	40.6
12	Chronic liver disease and cirrhosis (K70,K73-K74)	522	1.4	37.9
13	Parkinson's disease (G20-G21)	273	0.7	19.8
14	Pneumonitis due to solids and liquids (J69)	252	0.7	18.3
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	226	0.6	16.4
...	All other causes (Residual)	6,483	16.8	470.5
White, non-Hispanic, both sexes, all ages				
...	All causes	1,944,617	100.0	968.5
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	516,883	26.6	257.4
2	Malignant neoplasms (C00-C97)	455,978	23.4	227.1
3	Chronic lower respiratory diseases (J40-J47)	111,559	5.7	55.6
4	Cerebrovascular diseases (I60-I69)	108,886	5.6	54.2
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	91,830	4.7	45.7
6	Alzheimer's disease (G30)	64,660	3.3	32.2
7	Diabetes mellitus (E10-E14)	50,950	2.6	25.4
8	Influenza and pneumonia (J10-J18)	46,419	2.4	23.1
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	33,208	1.7	16.5
10	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	27,952	1.4	13.9
11	Septicemia (A40-A41)	25,556	1.3	12.7
12	Chronic liver disease and cirrhosis (K70,K73-K74)	20,605	1.1	10.3
13	Parkinson's disease (G20-G21)	17,806	0.9	8.9
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	17,034	0.9	8.5
15	Pneumonitis due to solids and liquids (J69)	14,563	0.7	7.3
...	All other causes (Residual)	340,728	17.5	169.7

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2006

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. The title for Essential hypertension and hypertensive renal disease was changed in 2006 to reflect the addition of Secondary hypertension (ICD code I15); see the "Technical Notes" of *Deaths: Final Data for 2006*]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
White, non-Hispanic, both sexes, 1-4 years				
...	All causes	2,279	100.0	25.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	814	35.7	8.9
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	253	11.1	2.8
3	Malignant neoplasms (C00-C97)	209	9.2	2.3
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	133	5.8	1.5
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	79	3.5	0.9
6	Influenza and pneumonia (J10-J18)	54	2.4	0.6
7	Septicemia (A40-A41)	44	1.9	0.5
7	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	44	1.9	0.5
9	Cerebrovascular diseases (I60-I69)	32	1.4	0.4
10	Certain conditions originating in the perinatal period (P00-P96)	29	1.3	0.3
11	Chronic lower respiratory diseases (J40-J47)	15	0.7	*
12	Complications of medical and surgical care (Y40-Y84,Y88)	13	0.6	*
13	Meningococcal infection (A39)	10	0.4	*
13	Meningitis (G00,G03)	10	0.4	*
15	Anemias (D50-D64)	9	0.4	*
...	All other causes (Residual)	531	23.3	5.8
White, non-Hispanic, both sexes, 5-14 years				
...	All causes	3,353	100.0	14.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,267	37.8	5.3
2	Malignant neoplasms (C00-C97)	540	16.1	2.3
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	198	5.9	0.8
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	143	4.3	0.6
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	134	4.0	0.6
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	128	3.8	0.5
7	Cerebrovascular diseases (I60-I69)	51	1.5	0.2
8	Septicemia (A40-A41)	47	1.4	0.2
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	41	1.2	0.2
10	Influenza and pneumonia (J10-J18)	39	1.2	0.2
11	Chronic lower respiratory diseases (J40-J47)	30	0.9	0.1
12	Diabetes mellitus (E10-E14)	13	0.4	*
12	Complications of medical and surgical care (Y40-Y84,Y88)	13	0.4	*
14	Anemias (D50-D64)	11	0.3	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	10	0.3	*
...	All other causes (Residual)	688	20.5	2.9

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2006

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. The title for Essential hypertension and hypertensive renal disease was changed in 2006 to reflect the addition of Secondary hypertension (ICD code I15); see the "Technical Notes" of *Deaths: Final Data for 2006*]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
White, non-Hispanic, both sexes, 15-24 years				
...	All causes	20,055	100.0	76.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	11,203	55.9	42.5
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,944	14.7	11.2
3	Malignant neoplasms (C00-C97)	980	4.9	3.7
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	900	4.5	3.4
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	608	3.0	2.3
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	272	1.4	1.0
7	Influenza and pneumonia (J10-J18)	114	0.6	0.4
8	Cerebrovascular diseases (I60-I69)	105	0.5	0.4
9	Diabetes mellitus (E10-E14)	89	0.4	0.3
10	Pregnancy, childbirth and the puerperium (O00-099)	78	0.4	0.3
11	Septicemia (A40-A41)	69	0.3	0.3
12	Chronic lower respiratory diseases (J40-J47)	54	0.3	0.2
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	54	0.3	0.2
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	49	0.2	0.2
15	Legal intervention (Y35,Y89.0)	30	0.1	0.1
...	All other causes (Residual)	2,506	12.5	9.5
White, non-Hispanic, both sexes, 25-44 years				
...	All causes	77,804	100.0	147.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	22,644	29.1	42.8
2	Malignant neoplasms (C00-C97)	11,553	14.8	21.8
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	9,734	12.5	18.4
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	9,197	11.8	17.4
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,970	2.5	3.7
6	Chronic liver disease and cirrhosis (K70,K73-K74)	1,859	2.4	3.5
7	Diabetes mellitus (E10-E14)	1,591	2.0	3.0
8	Human immunodeficiency virus (HIV) disease (B20-B24)	1,402	1.8	2.7
9	Cerebrovascular diseases (I60-I69)	1,337	1.7	2.5
10	Influenza and pneumonia (J10-J18)	738	0.9	1.4
11	Chronic lower respiratory diseases (J40-J47)	654	0.8	1.2
12	Septicemia (A40-A41)	635	0.8	1.2
13	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	632	0.8	1.2
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	470	0.6	0.9
15	Viral hepatitis (B15-B19)	352	0.5	0.7
...	All other causes (Residual)	13,036	16.8	24.6

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2006

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. The title for Essential hypertension and hypertensive renal disease was changed in 2006 to reflect the addition of Secondary hypertension (ICD code I15); see the "Technical Notes" of *Deaths: Final Data for 2006*]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
White, non-Hispanic, both sexes, 45-64 years				
...	All causes	338,287	100.0	605.2
1	Malignant neoplasms (C00-C97)	115,839	34.2	207.2
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	74,191	21.9	132.7
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	23,272	6.9	41.6
4	Chronic lower respiratory diseases (J40-J47)	13,779	4.1	24.7
5	Chronic liver disease and cirrhosis (K70,K73-K74)	10,949	3.2	19.6
6	Diabetes mellitus (E10-E14)	10,863	3.2	19.4
7	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	10,781	3.2	19.3
8	Cerebrovascular diseases (I60-I69)	10,050	3.0	18.0
9	Septicemia (A40-A41)	4,016	1.2	7.2
10	Influenza and pneumonia (J10-J18)	3,640	1.1	6.5
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3,625	1.1	6.5
12	Viral hepatitis (B15-B19)	3,296	1.0	5.9
13	Human immunodeficiency virus (HIV) disease (B20-B24)	1,899	0.6	3.4
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,776	0.5	3.2
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1,600	0.5	2.9
...	All other causes (Residual)	48,711	14.4	87.1
White, non-Hispanic, both sexes, 65 years and over				
...	All causes	1,489,841	100.0	4,914.3
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	431,970	29.0	1,424.9
2	Malignant neoplasms (C00-C97)	326,804	21.9	1,078.0
3	Cerebrovascular diseases (I60-I69)	97,255	6.5	320.8
4	Chronic lower respiratory diseases (J40-J47)	97,016	6.5	320.0
5	Alzheimer's disease (G30)	64,020	4.3	211.2
6	Influenza and pneumonia (J10-J18)	41,731	2.8	137.7
7	Diabetes mellitus (E10-E14)	38,385	2.6	126.6
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	32,051	2.2	105.7
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	28,964	1.9	95.5
10	Septicemia (A40-A41)	20,654	1.4	68.1
11	Parkinson's disease (G20-G21)	17,408	1.2	57.4
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	15,079	1.0	49.7
13	Pneumonitis due to solids and liquids (J69)	13,347	0.9	44.0
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	9,792	0.7	32.3
15	Aortic aneurysm and dissection (I71)	9,269	0.6	30.6
...	All other causes (Residual)	246,096	16.5	811.8

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2006

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. The title for Essential hypertension and hypertensive renal disease was changed in 2006 to reflect the addition of Secondary hypertension (ICD code I15); see the "Technical Notes" of *Deaths: Final Data for 2006*]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
White, non-Hispanic, male, all ages				
...	All causes	947,966	100.0	962.0
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	256,459	27.1	260.3
2	Malignant neoplasms (C00-C97)	236,393	24.9	239.9
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	57,758	6.1	58.6
4	Chronic lower respiratory diseases (J40-J47)	52,270	5.5	53.0
5	Cerebrovascular diseases (I60-I69)	41,941	4.4	42.6
6	Diabetes mellitus (E10-E14)	25,931	2.7	26.3
7	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	21,952	2.3	22.3
8	Influenza and pneumonia (J10-J18)	20,863	2.2	21.2
9	Alzheimer's disease (G30)	18,844	2.0	19.1
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	16,415	1.7	16.7
11	Chronic liver disease and cirrhosis (K70,K73-K74)	13,212	1.4	13.4
12	Septicemia (A40-A41)	11,526	1.2	11.7
13	Parkinson's disease (G20-G21)	10,277	1.1	10.4
14	Pneumonitis due to solids and liquids (J69)	7,694	0.8	7.8
15	Aortic aneurysm and dissection (I71)	6,512	0.7	6.6
...	All other causes (Residual)	149,919	15.8	152.1
White, non-Hispanic, male, 1-4 years				
...	All causes	1,247	100.0	26.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	458	36.7	9.8
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	122	9.8	2.6
3	Malignant neoplasms (C00-C97)	116	9.3	2.5
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	77	6.2	1.6
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	41	3.3	0.9
6	Influenza and pneumonia (J10-J18)	29	2.3	0.6
7	Septicemia (A40-A41)	24	1.9	0.5
8	Certain conditions originating in the perinatal period (P00-P96)	21	1.7	0.4
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	16	1.3	*
10	Cerebrovascular diseases (I60-I69)	12	1.0	*
11	Chronic lower respiratory diseases (J40-J47)	10	0.8	*
12	Meningococcal infection (A39)	7	0.6	*
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	6	0.5	*
13	Complications of medical and surgical care (Y40-Y84,Y88)	6	0.5	*
15	Anemias (D50-D64)	3	0.2	*
15	Meningitis (G00,G03)	3	0.2	*
15	Acute bronchitis and bronchiolitis (J20-J21)	3	0.2	*
...	All other causes (Residual)	293	23.5	6.3

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2006

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. The title for Essential hypertension and hypertensive renal disease was changed in 2006 to reflect the addition of Secondary hypertension (ICD code I15); see the "Technical Notes" of *Deaths: Final Data for 2006*]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
White, non-Hispanic, male, 5-14 years				
...	All causes	1,983	100.0	16.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	766	38.6	6.2
2	Malignant neoplasms (C00-C97)	309	15.6	2.5
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	113	5.7	0.9
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	100	5.0	0.8
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	73	3.7	0.6
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	68	3.4	0.6
7	Cerebrovascular diseases (I60-I69)	34	1.7	0.3
8	Septicemia (A40-A41)	28	1.4	0.2
9	Influenza and pneumonia (J10-J18)	25	1.3	0.2
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	24	1.2	0.2
11	Chronic lower respiratory diseases (J40-J47)	22	1.1	0.2
12	Diabetes mellitus (E10-E14)	8	0.4	*
12	Complications of medical and surgical care (Y40-Y84,Y88)	8	0.4	*
14	Aortic aneurysm and dissection (I71)	7	0.4	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	6	0.3	*
...	All other causes (Residual)	392	19.8	3.2
White, non-Hispanic, male, 15-24 years				
...	All causes	14,553	100.0	107.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	8,290	57.0	61.3
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,496	17.2	18.5
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	638	4.4	4.7
4	Malignant neoplasms (C00-C97)	608	4.2	4.5
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	410	2.8	3.0
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	158	1.1	1.2
7	Influenza and pneumonia (J10-J18)	62	0.4	0.5
8	Cerebrovascular diseases (I60-I69)	60	0.4	0.4
9	Diabetes mellitus (E10-E14)	47	0.3	0.3
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	37	0.3	0.3
11	Septicemia (A40-A41)	33	0.2	0.2
12	Chronic lower respiratory diseases (J40-J47)	30	0.2	0.2
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	29	0.2	0.2
14	Legal intervention (Y35,Y89.0)	28	0.2	0.2
15	Aortic aneurysm and dissection (I71)	22	0.2	0.2
...	All other causes (Residual)	1,605	11.0	11.9

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2006

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. The title for Essential hypertension and hypertensive renal disease was changed in 2006 to reflect the addition of Secondary hypertension (ICD code I15); see the "Technical Notes" of *Deaths: Final Data for 2006*]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
White, non-Hispanic, male, 25-44 years				
...	All causes	50,724	100.0	191.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	16,292	32.1	61.4
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	7,139	14.1	26.9
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	6,988	13.8	26.3
4	Malignant neoplasms (C00-C97)	5,291	10.4	19.9
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,364	2.7	5.1
6	Chronic liver disease and cirrhosis (K70,K73-K74)	1,182	2.3	4.5
7	Human immunodeficiency virus (HIV) disease (B20-B24)	1,133	2.2	4.3
8	Diabetes mellitus (E10-E14)	966	1.9	3.6
9	Cerebrovascular diseases (I60-I69)	692	1.4	2.6
10	Influenza and pneumonia (J10-J18)	393	0.8	1.5
11	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	368	0.7	1.4
12	Septicemia (A40-A41)	339	0.7	1.3
13	Chronic lower respiratory diseases (J40-J47)	310	0.6	1.2
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	288	0.6	1.1
15	Viral hepatitis (B15-B19)	224	0.4	0.8
...	All other causes (Residual)	7,755	15.3	29.2
White, non-Hispanic, male, 45-64 years				
...	All causes	207,206	100.0	752.6
1	Malignant neoplasms (C00-C97)	61,832	29.8	224.6
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	53,340	25.7	193.7
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	15,971	7.7	58.0
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	8,078	3.9	29.3
5	Chronic liver disease and cirrhosis (K70,K73-K74)	7,723	3.7	28.1
6	Chronic lower respiratory diseases (J40-J47)	6,945	3.4	25.2
7	Diabetes mellitus (E10-E14)	6,697	3.2	24.3
8	Cerebrovascular diseases (I60-I69)	5,505	2.7	20.0
9	Viral hepatitis (B15-B19)	2,397	1.2	8.7
10	Septicemia (A40-A41)	2,158	1.0	7.8
11	Influenza and pneumonia (J10-J18)	2,144	1.0	7.8
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2,144	1.0	7.8
13	Human immunodeficiency virus (HIV) disease (B20-B24)	1,630	0.8	5.9
14	Aortic aneurysm and dissection (I71)	1,205	0.6	4.4
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,128	0.5	4.1
...	All other causes (Residual)	28,309	13.7	102.8

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2006

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. The title for Essential hypertension and hypertensive renal disease was changed in 2006 to reflect the addition of Secondary hypertension (ICD code I15); see the "Technical Notes" of *Deaths: Final Data for 2006*]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
White, non-Hispanic, male, 65 years and over				
...	All causes	664,897	100.0	5,184.7
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	195,509	29.4	1,524.5
2	Malignant neoplasms (C00-C97)	168,211	25.3	1,311.7
3	Chronic lower respiratory diseases (J40-J47)	44,946	6.8	350.5
4	Cerebrovascular diseases (I60-I69)	35,607	5.4	277.7
5	Alzheimer's disease (G30)	18,540	2.8	144.6
6	Diabetes mellitus (E10-E14)	18,207	2.7	142.0
7	Influenza and pneumonia (J10-J18)	18,142	2.7	141.5
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	15,636	2.4	121.9
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	13,880	2.1	108.2
10	Parkinson's disease (G20-G21)	10,009	1.5	78.0
11	Septicemia (A40-A41)	8,896	1.3	69.4
12	Pneumonitis due to solids and liquids (J69)	6,940	1.0	54.1
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	5,208	0.8	40.6
14	Aortic aneurysm and dissection (I71)	5,072	0.8	39.6
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	4,748	0.7	37.0
...	All other causes (Residual)	95,346	14.3	743.5
White, non-Hispanic, female, all ages				
...	All causes	996,651	100.0	974.7
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	260,424	26.1	254.7
2	Malignant neoplasms (C00-C97)	219,585	22.0	214.7
3	Cerebrovascular diseases (I60-I69)	66,945	6.7	65.5
4	Chronic lower respiratory diseases (J40-J47)	59,289	5.9	58.0
5	Alzheimer's disease (G30)	45,816	4.6	44.8
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	34,072	3.4	33.3
7	Influenza and pneumonia (J10-J18)	25,556	2.6	25.0
8	Diabetes mellitus (E10-E14)	25,019	2.5	24.5
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	16,793	1.7	16.4
10	Septicemia (A40-A41)	14,030	1.4	13.7
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	10,583	1.1	10.3
12	Parkinson's disease (G20-G21)	7,529	0.8	7.4
13	Chronic liver disease and cirrhosis (K70,K73-K74)	7,393	0.7	7.2
14	Pneumonitis due to solids and liquids (J69)	6,869	0.7	6.7
15	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	6,000	0.6	5.9
...	All other causes (Residual)	190,748	19.1	186.5

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2006

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. The title for Essential hypertension and hypertensive renal disease was changed in 2006 to reflect the addition of Secondary hypertension (ICD code I15); see the "Technical Notes" of *Deaths: Final Data for 2006*]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
White, non-Hispanic, female, 1-4 years				
...	All causes	1,032	100.0	23.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	356	34.5	8.0
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	131	12.7	2.9
3	Malignant neoplasms (C00-C97)	93	9.0	2.1
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	56	5.4	1.3
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	38	3.7	0.9
6	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	28	2.7	0.6
7	Influenza and pneumonia (J10-J18)	25	2.4	0.6
8	Septicemia (A40-A41)	20	1.9	0.4
8	Cerebrovascular diseases (I60-I69)	20	1.9	0.4
10	Certain conditions originating in the perinatal period (P00-P96)	8	0.8	*
11	Meningitis (G00,G03)	7	0.7	*
11	Complications of medical and surgical care (Y40-Y84,Y88)	7	0.7	*
13	Anemias (D50-D64)	6	0.6	*
14	Chronic lower respiratory diseases (J40-J47)	5	0.5	*
15	Pneumonitis due to solids and liquids (J69)	4	0.4	*
...	All other causes (Residual)	228	22.1	5.1
White, non-Hispanic, female, 5-14 years				
...	All causes	1,370	100.0	11.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	501	36.6	4.3
2	Malignant neoplasms (C00-C97)	231	16.9	2.0
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	85	6.2	0.7
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	66	4.8	0.6
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	55	4.0	0.5
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	43	3.1	0.4
7	Septicemia (A40-A41)	19	1.4	*
8	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	17	1.2	*
8	Cerebrovascular diseases (I60-I69)	17	1.2	*
10	Influenza and pneumonia (J10-J18)	14	1.0	*
11	Chronic lower respiratory diseases (J40-J47)	8	0.6	*
12	Anemias (D50-D64)	6	0.4	*
13	Diabetes mellitus (E10-E14)	5	0.4	*
13	Complications of medical and surgical care (Y40-Y84,Y88)	5	0.4	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	4	0.3	*
...	All other causes (Residual)	294	21.5	2.5

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2006

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. The title for Essential hypertension and hypertensive renal disease was changed in 2006 to reflect the addition of Secondary hypertension (ICD code I15); see the "Technical Notes" of *Deaths: Final Data for 2006*]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
White, non-Hispanic, female, 15-24 years				
...	All causes	5,502	100.0	42.9
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,913	52.9	22.7
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	448	8.1	3.5
3	Malignant neoplasms (C00-C97)	372	6.8	2.9
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	262	4.8	2.0
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	198	3.6	1.5
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	114	2.1	0.9
7	Pregnancy, childbirth and the puerperium (O00-099)	78	1.4	0.6
8	Influenza and pneumonia (J10-J18)	52	0.9	0.4
9	Cerebrovascular diseases (I60-I69)	45	0.8	0.4
10	Diabetes mellitus (E10-E14)	42	0.8	0.3
11	Septicemia (A40-A41)	36	0.7	0.3
12	Chronic lower respiratory diseases (J40-J47)	24	0.4	0.2
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	20	0.4	0.2
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	17	0.3	*
15	Complications of medical and surgical care (Y40-Y84,Y88)	12	0.2	*
...	All other causes (Residual)	869	15.8	6.8
White, non-Hispanic, female, 25-44 years				
...	All causes	27,080	100.0	102.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	6,352	23.5	24.1
2	Malignant neoplasms (C00-C97)	6,262	23.1	23.8
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,746	10.1	10.4
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,058	7.6	7.8
5	Chronic liver disease and cirrhosis (K70,K73-K74)	677	2.5	2.6
6	Cerebrovascular diseases (I60-I69)	645	2.4	2.4
7	Diabetes mellitus (E10-E14)	625	2.3	2.4
8	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	606	2.2	2.3
9	Influenza and pneumonia (J10-J18)	345	1.3	1.3
10	Chronic lower respiratory diseases (J40-J47)	344	1.3	1.3
11	Septicemia (A40-A41)	296	1.1	1.1
12	Human immunodeficiency virus (HIV) disease (B20-B24)	269	1.0	1.0
13	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	264	1.0	1.0
14	Pregnancy, childbirth and the puerperium (O00-099)	192	0.7	0.7
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	182	0.7	0.7
...	All other causes (Residual)	5,217	19.3	19.8

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2006

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. The title for Essential hypertension and hypertensive renal disease was changed in 2006 to reflect the addition of Secondary hypertension (ICD code I15); see the "Technical Notes" of *Deaths: Final Data for 2006*]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
White, non-Hispanic, female, 45-64 years				
...	All causes	131,081	100.0	462.2
1	Malignant neoplasms (C00-C97)	54,007	41.2	190.4
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	20,851	15.9	73.5
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	7,301	5.6	25.7
4	Chronic lower respiratory diseases (J40-J47)	6,834	5.2	24.1
5	Cerebrovascular diseases (I60-I69)	4,545	3.5	16.0
6	Diabetes mellitus (E10-E14)	4,166	3.2	14.7
7	Chronic liver disease and cirrhosis (K70,K73-K74)	3,226	2.5	11.4
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,703	2.1	9.5
9	Septicemia (A40-A41)	1,858	1.4	6.6
10	Influenza and pneumonia (J10-J18)	1,496	1.1	5.3
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,481	1.1	5.2
12	Viral hepatitis (B15-B19)	899	0.7	3.2
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	698	0.5	2.5
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	648	0.5	2.3
15	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	504	0.4	1.8
...	All other causes (Residual)	19,864	15.2	70.0
White, non-Hispanic, female, 65 years and over				
...	All causes	824,944	100.0	4,716.0
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	236,461	28.7	1,351.8
2	Malignant neoplasms (C00-C97)	158,593	19.2	906.6
3	Cerebrovascular diseases (I60-I69)	61,648	7.5	352.4
4	Chronic lower respiratory diseases (J40-J47)	52,070	6.3	297.7
5	Alzheimer's disease (G30)	45,480	5.5	260.0
6	Influenza and pneumonia (J10-J18)	23,589	2.9	134.9
7	Diabetes mellitus (E10-E14)	20,178	2.4	115.4
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	16,415	2.0	93.8
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	15,084	1.8	86.2
10	Septicemia (A40-A41)	11,758	1.4	67.2
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	9,871	1.2	56.4
12	Parkinson's disease (G20-G21)	7,399	0.9	42.3
13	Pneumonitis due to solids and liquids (J69)	6,407	0.8	36.6
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	5,044	0.6	28.8
15	Atherosclerosis (I70)	4,517	0.5	25.8
...	All other causes (Residual)	150,430	18.2	860.0

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2006

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. The title for Essential hypertension and hypertensive renal disease was changed in 2006 to reflect the addition of Secondary hypertension (ICD code I15); see the "Technical Notes" of *Deaths: Final Data for 2006*]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Black, non-Hispanic, both sexes, all ages				
...	All causes	286,581	100.0	759.1
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	71,461	24.9	189.3
2	Malignant neoplasms (C00-C97)	62,475	21.8	165.5
3	Cerebrovascular diseases (I60-I69)	16,882	5.9	44.7
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	13,684	4.8	36.2
5	Diabetes mellitus (E10-E14)	12,671	4.4	33.6
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	8,902	3.1	23.6
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	8,334	2.9	22.1
8	Chronic lower respiratory diseases (J40-J47)	7,657	2.7	20.3
9	Human immunodeficiency virus (HIV) disease (B20-B24)	6,767	2.4	17.9
10	Septicemia (A40-A41)	6,045	2.1	16.0
11	Influenza and pneumonia (J10-J18)	5,242	1.8	13.9
12	Certain conditions originating in the perinatal period (P00-P96)	4,906	1.7	13.0
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	4,851	1.7	12.8
14	Alzheimer's disease (G30)	4,422	1.5	11.7
15	Chronic liver disease and cirrhosis (K70,K73-K74)	2,310	0.8	6.1
...	All other causes (Residual)	49,972	17.4	132.4
Black, non-Hispanic, both sexes, 1-4 years				
...	All causes	1,108	100.0	44.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	355	32.0	14.2
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	150	13.5	6.0
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	106	9.6	4.2
4	Malignant neoplasms (C00-C97)	63	5.7	2.5
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	44	4.0	1.8
6	Influenza and pneumonia (J10-J18)	34	3.1	1.4
7	Certain conditions originating in the perinatal period (P00-P96)	22	2.0	0.9
8	Septicemia (A40-A41)	19	1.7	*
9	Chronic lower respiratory diseases (J40-J47)	15	1.4	*
10	Anemias (D50-D64)	12	1.1	*
11	Cerebrovascular diseases (I60-I69)	11	1.0	*
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	9	0.8	*
13	Acute bronchitis and bronchiolitis (J20-J21)	5	0.5	*
14	Complications of medical and surgical care (Y40-Y84,Y88)	4	0.4	*
15	Meningitis (G00,G03)	3	0.3	*
...	All other causes (Residual)	256	23.1	10.2

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2006

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. The title for Essential hypertension and hypertensive renal disease was changed in 2006 to reflect the addition of Secondary hypertension (ICD code I15); see the "Technical Notes" of *Deaths: Final Data for 2006*]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Black, non-Hispanic, both sexes, 5-14 years				
...	All causes	1,361	100.0	21.9
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	460	33.8	7.4
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	147	10.8	2.4
3	Malignant neoplasms (C00-C97)	140	10.3	2.2
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	67	4.9	1.1
5	Chronic lower respiratory diseases (J40-J47)	66	4.8	1.1
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	63	4.6	1.0
7	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	35	2.6	0.6
8	Septicemia (A40-A41)	20	1.5	0.3
8	Cerebrovascular diseases (I60-I69)	20	1.5	0.3
10	Anemias (D50-D64)	19	1.4	*
11	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	15	1.1	*
12	Human immunodeficiency virus (HIV) disease (B20-B24)	11	0.8	*
13	Diabetes mellitus (E10-E14)	9	0.7	*
13	Influenza and pneumonia (J10-J18)	9	0.7	*
15	Meningitis (G00,G03)	8	0.6	*
...	All other causes (Residual)	272	20.0	4.4
Black, non-Hispanic, both sexes, 15-24 years				
...	All causes	7,347	100.0	115.6
1	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	3,253	44.3	51.2
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,703	23.2	26.8
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	407	5.5	6.4
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	302	4.1	4.8
5	Malignant neoplasms (C00-C97)	259	3.5	4.1
6	Human immunodeficiency virus (HIV) disease (B20-B24)	143	1.9	2.2
7	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	87	1.2	1.4
8	Chronic lower respiratory diseases (J40-J47)	74	1.0	1.2
9	Anemias (D50-D64)	64	0.9	1.0
10	Pregnancy, childbirth and the puerperium (O00-099)	57	0.8	0.9
11	Diabetes mellitus (E10-E14)	54	0.7	0.8
12	Cerebrovascular diseases (I60-I69)	44	0.6	0.7
13	Septicemia (A40-A41)	41	0.6	0.6
13	Influenza and pneumonia (J10-J18)	41	0.6	0.6
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	30	0.4	0.5
...	All other causes (Residual)	788	10.7	12.4

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2006

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. The title for Essential hypertension and hypertensive renal disease was changed in 2006 to reflect the addition of Secondary hypertension (ICD code I15); see the "Technical Notes" of *Deaths: Final Data for 2006*]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Black, non-Hispanic, both sexes, 25-44 years				
...	All causes	27,620	100.0	256.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	4,210	15.2	39.0
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	4,057	14.7	37.6
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	3,966	14.4	36.8
4	Malignant neoplasms (C00-C97)	3,131	11.3	29.0
5	Human immunodeficiency virus (HIV) disease (B20-B24)	2,924	10.6	27.1
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	872	3.2	8.1
7	Cerebrovascular diseases (I60-I69)	840	3.0	7.8
8	Diabetes mellitus (E10-E14)	815	3.0	7.6
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	439	1.6	4.1
10	Septicemia (A40-A41)	358	1.3	3.3
11	Chronic lower respiratory diseases (J40-J47)	330	1.2	3.1
12	Chronic liver disease and cirrhosis (K70,K73-K74)	273	1.0	2.5
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	272	1.0	2.5
14	Influenza and pneumonia (J10-J18)	260	0.9	2.4
15	Anemias (D50-D64)	254	0.9	2.4
...	All other causes (Residual)	4,619	16.7	42.8
Black, non-Hispanic, both sexes, 45-64 years				
...	All causes	82,951	100.0	1,020.3
1	Malignant neoplasms (C00-C97)	22,761	27.4	280.0
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	20,462	24.7	251.7
3	Cerebrovascular diseases (I60-I69)	4,490	5.4	55.2
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	4,361	5.3	53.6
5	Diabetes mellitus (E10-E14)	3,877	4.7	47.7
6	Human immunodeficiency virus (HIV) disease (B20-B24)	3,379	4.1	41.6
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2,170	2.6	26.7
8	Chronic lower respiratory diseases (J40-J47)	1,804	2.2	22.2
9	Septicemia (A40-A41)	1,659	2.0	20.4
10	Chronic liver disease and cirrhosis (K70,K73-K74)	1,492	1.8	18.4
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,295	1.6	15.9
12	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,116	1.3	13.7
13	Influenza and pneumonia (J10-J18)	956	1.2	11.8
14	Viral hepatitis (B15-B19)	782	0.9	9.6
15	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	456	0.5	5.6
...	All other causes (Residual)	11,891	14.3	146.3

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2006

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. The title for Essential hypertension and hypertensive renal disease was changed in 2006 to reflect the addition of Secondary hypertension (ICD code I15); see the "Technical Notes" of *Deaths: Final Data for 2006*]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Black, non-Hispanic, both sexes, 65 years and over				
...	All causes	157,668	100.0	5,051.0
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	46,413	29.4	1,486.9
2	Malignant neoplasms (C00-C97)	36,113	22.9	1,156.9
3	Cerebrovascular diseases (I60-I69)	11,428	7.2	366.1
4	Diabetes mellitus (E10-E14)	7,913	5.0	253.5
5	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	5,648	3.6	180.9
6	Chronic lower respiratory diseases (J40-J47)	5,357	3.4	171.6
7	Alzheimer's disease (G30)	4,358	2.8	139.6
8	Influenza and pneumonia (J10-J18)	3,855	2.4	123.5
9	Septicemia (A40-A41)	3,849	2.4	123.3
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	3,275	2.1	104.9
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,245	1.4	71.9
12	Pneumonitis due to solids and liquids (J69)	1,166	0.7	37.4
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	767	0.5	24.6
14	Aortic aneurysm and dissection (I71)	716	0.5	22.9
15	Parkinson's disease (G20-G21)	631	0.4	20.2
...	All other causes (Residual)	23,934	15.2	766.7
Black, non-Hispanic, male, all ages				
...	All causes	146,729	100.0	815.3
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	35,787	24.4	198.8
2	Malignant neoplasms (C00-C97)	32,253	22.0	179.2
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	9,443	6.4	52.5
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	7,566	5.2	42.0
5	Cerebrovascular diseases (I60-I69)	7,349	5.0	40.8
6	Diabetes mellitus (E10-E14)	5,701	3.9	31.7
7	Human immunodeficiency virus (HIV) disease (B20-B24)	4,384	3.0	24.4
8	Chronic lower respiratory diseases (J40-J47)	4,091	2.8	22.7
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3,782	2.6	21.0
10	Certain conditions originating in the perinatal period (P00-P96)	2,706	1.8	15.0
11	Septicemia (A40-A41)	2,688	1.8	14.9
12	Influenza and pneumonia (J10-J18)	2,449	1.7	13.6
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	2,100	1.4	11.7
14	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,630	1.1	9.1
15	Chronic liver disease and cirrhosis (K70,K73-K74)	1,485	1.0	8.3
...	All other causes (Residual)	23,315	15.9	129.5

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2006

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. The title for Essential hypertension and hypertensive renal disease was changed in 2006 to reflect the addition of Secondary hypertension (ICD code I15); see the "Technical Notes" of *Deaths: Final Data for 2006*]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Black, non-Hispanic, male, 1-4 years				
...	All causes	612	100.0	48.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	207	33.8	16.3
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	81	13.2	6.4
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	41	6.7	3.2
4	Malignant neoplasms (C00-C97)	34	5.6	2.7
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	27	4.4	2.1
6	Influenza and pneumonia (J10-J18)	16	2.6	*
7	Chronic lower respiratory diseases (J40-J47)	12	2.0	*
8	Septicemia (A40-A41)	11	1.8	*
9	Anemias (D50-D64)	9	1.5	*
9	Certain conditions originating in the perinatal period (P00-P96)	9	1.5	*
11	Cerebrovascular diseases (I60-I69)	7	1.1	*
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	5	0.8	*
13	Acute bronchitis and bronchiolitis (J20-J21)	3	0.5	*
13	Complications of medical and surgical care (Y40-Y84,Y88)	3	0.5	*
15	Meningococcal infection (A39)	1	0.2	*
15	Syphilis (A50-A53)	1	0.2	*
15	Chronic liver disease and cirrhosis (K70,K73-K74)	1	0.2	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1	0.2	*
...	All other causes (Residual)	143	23.4	11.3
Black, non-Hispanic, male, 5-14 years				
...	All causes	814	100.0	25.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	288	35.4	9.1
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	104	12.8	3.3
3	Malignant neoplasms (C00-C97)	80	9.8	2.5
4	Chronic lower respiratory diseases (J40-J47)	41	5.0	1.3
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	34	4.2	1.1
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	29	3.6	0.9
7	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	28	3.4	0.9
8	Septicemia (A40-A41)	13	1.6	*
9	Anemias (D50-D64)	9	1.1	*
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	7	0.9	*
10	Cerebrovascular diseases (I60-I69)	7	0.9	*
12	Human immunodeficiency virus (HIV) disease (B20-B24)	6	0.7	*
13	Meningitis (G00,G03)	3	0.4	*
13	Influenza and pneumonia (J10-J18)	3	0.4	*
13	Pneumonitis due to solids and liquids (J69)	3	0.4	*
...	All other causes (Residual)	159	19.5	5.0

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2006

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. The title for Essential hypertension and hypertensive renal disease was changed in 2006 to reflect the addition of Secondary hypertension (ICD code I15); see the "Technical Notes" of *Deaths: Final Data for 2006*]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Black, non-Hispanic, male, 15-24 years				
...	All causes	5,691	100.0	176.9
1	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	2,946	51.8	91.6
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,293	22.7	40.2
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	351	6.2	10.9
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	180	3.2	5.6
5	Malignant neoplasms (C00-C97)	151	2.7	4.7
6	Human immunodeficiency virus (HIV) disease (B20-B24)	83	1.5	2.6
7	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	52	0.9	1.6
8	Chronic lower respiratory diseases (J40-J47)	46	0.8	1.4
9	Anemias (D50-D64)	37	0.7	1.1
10	Diabetes mellitus (E10-E14)	31	0.5	1.0
11	Legal intervention (Y35,Y89.0)	24	0.4	0.7
12	Influenza and pneumonia (J10-J18)	23	0.4	0.7
13	Septicemia (A40-A41)	21	0.4	0.7
14	Cerebrovascular diseases (I60-I69)	20	0.4	0.6
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	11	0.2	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	11	0.2	*
...	All other causes (Residual)	411	7.2	12.8
Black, non-Hispanic, male, 25-44 years				
...	All causes	17,191	100.0	335.6
1	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	3,367	19.6	65.7
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,117	18.1	60.8
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,517	14.6	49.1
4	Human immunodeficiency virus (HIV) disease (B20-B24)	1,752	10.2	34.2
5	Malignant neoplasms (C00-C97)	1,244	7.2	24.3
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	756	4.4	14.8
7	Diabetes mellitus (E10-E14)	478	2.8	9.3
8	Cerebrovascular diseases (I60-I69)	414	2.4	8.1
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	237	1.4	4.6
10	Septicemia (A40-A41)	176	1.0	3.4
11	Chronic lower respiratory diseases (J40-J47)	163	0.9	3.2
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	162	0.9	3.2
13	Chronic liver disease and cirrhosis (K70,K73-K74)	160	0.9	3.1
14	Influenza and pneumonia (J10-J18)	132	0.8	2.6
15	Anemias (D50-D64)	119	0.7	2.3
...	All other causes (Residual)	2,397	13.9	46.8

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2006

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. The title for Essential hypertension and hypertensive renal disease was changed in 2006 to reflect the addition of Secondary hypertension (ICD code I15); see the "Technical Notes" of *Deaths: Final Data for 2006*]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Black, non-Hispanic, male, 45-64 years				
...	All causes	48,758	100.0	1,316.5
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	12,919	26.5	348.8
2	Malignant neoplasms (C00-C97)	12,076	24.8	326.1
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,147	6.5	85.0
4	Cerebrovascular diseases (I60-I69)	2,543	5.2	68.7
5	Human immunodeficiency virus (HIV) disease (B20-B24)	2,325	4.8	62.8
6	Diabetes mellitus (E10-E14)	2,081	4.3	56.2
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,187	2.4	32.1
8	Chronic liver disease and cirrhosis (K70,K73-K74)	1,002	2.1	27.1
9	Chronic lower respiratory diseases (J40-J47)	993	2.0	26.8
10	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	880	1.8	23.8
11	Septicemia (A40-A41)	877	1.8	23.7
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	727	1.5	19.6
13	Influenza and pneumonia (J10-J18)	574	1.2	15.5
14	Viral hepatitis (B15-B19)	535	1.1	14.4
15	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	370	0.8	10.0
...	All other causes (Residual)	6,522	13.4	176.1
Black, non-Hispanic, male, 65 years and over				
...	All causes	68,938	100.0	5,752.5
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	20,044	29.1	1,672.6
2	Malignant neoplasms (C00-C97)	18,664	27.1	1,557.4
3	Cerebrovascular diseases (I60-I69)	4,325	6.3	360.9
4	Diabetes mellitus (E10-E14)	3,109	4.5	259.4
5	Chronic lower respiratory diseases (J40-J47)	2,830	4.1	236.1
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2,320	3.4	193.6
7	Influenza and pneumonia (J10-J18)	1,648	2.4	137.5
8	Septicemia (A40-A41)	1,537	2.2	128.3
9	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,206	1.7	100.6
10	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,182	1.7	98.6
11	Alzheimer's disease (G30)	1,148	1.7	95.8
12	Pneumonitis due to solids and liquids (J69)	595	0.9	49.6
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	352	0.5	29.4
14	Parkinson's disease (G20-G21)	351	0.5	29.3
15	Chronic liver disease and cirrhosis (K70,K73-K74)	319	0.5	26.6
...	All other causes (Residual)	9,308	13.5	776.7

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2006

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. The title for Essential hypertension and hypertensive renal disease was changed in 2006 to reflect the addition of Secondary hypertension (ICD code I15); see the "Technical Notes" of *Deaths: Final Data for 2006*]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Black, non-Hispanic, female, all ages				
...	All causes	139,852	100.0	708.0
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	35,674	25.5	180.6
2	Malignant neoplasms (C00-C97)	30,222	21.6	153.0
3	Cerebrovascular diseases (I60-I69)	9,533	6.8	48.3
4	Diabetes mellitus (E10-E14)	6,970	5.0	35.3
5	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	4,552	3.3	23.0
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	4,241	3.0	21.5
7	Chronic lower respiratory diseases (J40-J47)	3,566	2.5	18.1
8	Septicemia (A40-A41)	3,357	2.4	17.0
9	Alzheimer's disease (G30)	3,246	2.3	16.4
10	Influenza and pneumonia (J10-J18)	2,793	2.0	14.1
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	2,751	2.0	13.9
12	Human immunodeficiency virus (HIV) disease (B20-B24)	2,383	1.7	12.1
13	Certain conditions originating in the perinatal period (P00-P96)	2,200	1.6	11.1
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,336	1.0	6.8
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	828	0.6	4.2
...	All other causes (Residual)	26,200	18.7	132.6
Black, non-Hispanic, female, 1-4 years				
...	All causes	496	100.0	40.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	148	29.8	12.0
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	69	13.9	5.6
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	65	13.1	5.3
4	Malignant neoplasms (C00-C97)	29	5.8	2.4
5	Influenza and pneumonia (J10-J18)	18	3.6	*
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	17	3.4	*
7	Certain conditions originating in the perinatal period (P00-P96)	13	2.6	*
8	Septicemia (A40-A41)	8	1.6	*
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	4	0.8	*
9	Cerebrovascular diseases (I60-I69)	4	0.8	*
11	Anemias (D50-D64)	3	0.6	*
11	Meningitis (G00,G03)	3	0.6	*
11	Chronic lower respiratory diseases (J40-J47)	3	0.6	*
14	Diabetes mellitus (E10-E14)	2	0.4	*
14	Acute bronchitis and bronchiolitis (J20-J21)	2	0.4	*
...	All other causes (Residual)	108	21.8	8.8

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2006

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. The title for Essential hypertension and hypertensive renal disease was changed in 2006 to reflect the addition of Secondary hypertension (ICD code I15); see the "Technical Notes" of *Deaths: Final Data for 2006*]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Black, non-Hispanic, female, 5-14 years				
...	All causes	547	100.0	17.9
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	172	31.4	5.6
2	Malignant neoplasms (C00-C97)	60	11.0	2.0
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	43	7.9	1.4
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	38	6.9	1.2
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	29	5.3	0.9
6	Chronic lower respiratory diseases (J40-J47)	25	4.6	0.8
7	Cerebrovascular diseases (I60-I69)	13	2.4	*
8	Anemias (D50-D64)	10	1.8	*
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	8	1.5	*
10	Septicemia (A40-A41)	7	1.3	*
10	Diabetes mellitus (E10-E14)	7	1.3	*
10	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	7	1.3	*
13	Influenza and pneumonia (J10-J18)	6	1.1	*
14	Human immunodeficiency virus (HIV) disease (B20-B24)	5	0.9	*
14	Meningitis (G00,G03)	5	0.9	*
...	All other causes (Residual)	112	20.5	3.7
Black, non-Hispanic, female, 15-24 years				
...	All causes	1,656	100.0	52.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	410	24.8	13.1
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	307	18.5	9.8
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	122	7.4	3.9
4	Malignant neoplasms (C00-C97)	108	6.5	3.4
5	Human immunodeficiency virus (HIV) disease (B20-B24)	60	3.6	1.9
6	Pregnancy, childbirth and the puerperium (O00-099)	57	3.4	1.8
7	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	56	3.4	1.8
8	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	35	2.1	1.1
9	Chronic lower respiratory diseases (J40-J47)	28	1.7	0.9
10	Anemias (D50-D64)	27	1.6	0.9
11	Cerebrovascular diseases (I60-I69)	24	1.4	0.8
12	Diabetes mellitus (E10-E14)	23	1.4	0.7
13	Septicemia (A40-A41)	20	1.2	0.6
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	19	1.1	*
15	Influenza and pneumonia (J10-J18)	18	1.1	*
...	All other causes (Residual)	342	20.7	10.9

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2006

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. The title for Essential hypertension and hypertensive renal disease was changed in 2006 to reflect the addition of Secondary hypertension (ICD code I15); see the "Technical Notes" of *Deaths: Final Data for 2006*]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Black, non-Hispanic, female, 25-44 years				
...	All causes	10,429	100.0	184.2
1	Malignant neoplasms (C00-C97)	1,887	18.1	33.3
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,540	14.8	27.2
3	Human immunodeficiency virus (HIV) disease (B20-B24)	1,172	11.2	20.7
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,093	10.5	19.3
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	599	5.7	10.6
6	Cerebrovascular diseases (I60-I69)	426	4.1	7.5
7	Diabetes mellitus (E10-E14)	337	3.2	6.0
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	202	1.9	3.6
9	Septicemia (A40-A41)	182	1.7	3.2
10	Pregnancy, childbirth and the puerperium (O00-099)	176	1.7	3.1
11	Chronic lower respiratory diseases (J40-J47)	167	1.6	2.9
12	Anemias (D50-D64)	135	1.3	2.4
13	Influenza and pneumonia (J10-J18)	128	1.2	2.3
14	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	116	1.1	2.0
15	Chronic liver disease and cirrhosis (K70,K73-K74)	113	1.1	2.0
...	All other causes (Residual)	2,156	20.7	38.1
Black, non-Hispanic, female, 45-64 years				
...	All causes	34,193	100.0	772.4
1	Malignant neoplasms (C00-C97)	10,685	31.2	241.4
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	7,543	22.1	170.4
3	Cerebrovascular diseases (I60-I69)	1,947	5.7	44.0
4	Diabetes mellitus (E10-E14)	1,796	5.3	40.6
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,214	3.6	27.4
6	Human immunodeficiency virus (HIV) disease (B20-B24)	1,054	3.1	23.8
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	983	2.9	22.2
8	Chronic lower respiratory diseases (J40-J47)	811	2.4	18.3
9	Septicemia (A40-A41)	782	2.3	17.7
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	568	1.7	12.8
11	Chronic liver disease and cirrhosis (K70,K73-K74)	490	1.4	11.1
12	Influenza and pneumonia (J10-J18)	382	1.1	8.6
13	Viral hepatitis (B15-B19)	247	0.7	5.6
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	236	0.7	5.3
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	156	0.5	3.5
...	All other causes (Residual)	5,299	15.5	119.7

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2006

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. The title for Essential hypertension and hypertensive renal disease was changed in 2006 to reflect the addition of Secondary hypertension (ICD code I15); see the "Technical Notes" of *Deaths: Final Data for 2006*]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Black, non-Hispanic, female, 65 years and over				
...	All causes	88,730	100.0	4,613.8
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	26,369	29.7	1,371.1
2	Malignant neoplasms (C00-C97)	17,449	19.7	907.3
3	Cerebrovascular diseases (I60-I69)	7,103	8.0	369.3
4	Diabetes mellitus (E10-E14)	4,804	5.4	249.8
5	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3,328	3.8	173.1
6	Alzheimer's disease (G30)	3,210	3.6	166.9
7	Chronic lower respiratory diseases (J40-J47)	2,527	2.8	131.4
8	Septicemia (A40-A41)	2,312	2.6	120.2
9	Influenza and pneumonia (J10-J18)	2,207	2.5	114.8
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	2,069	2.3	107.6
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,063	1.2	55.3
12	Pneumonitis due to solids and liquids (J69)	571	0.6	29.7
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	415	0.5	21.6
14	Aortic aneurysm and dissection (I71)	409	0.5	21.3
15	Atherosclerosis (I70)	355	0.4	18.5
...	All other causes (Residual)	14,539	16.4	756.0

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable