

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2007

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007; there were no deaths due to J09 in 2007]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, both sexes, all ages				
...	All causes	135,519	100.0	297.8
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	29,021	21.4	63.8
2	Malignant neoplasms (C00-C97)	27,660	20.4	60.8
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	11,723	8.7	25.8
4	Cerebrovascular diseases (I60-I69)	7,078	5.2	15.6
5	Diabetes mellitus (E10-E14)	6,417	4.7	14.1
6	Chronic liver disease and cirrhosis (K70,K73-K74)	3,913	2.9	8.6
7	Chronic lower respiratory diseases (J40-J47)	3,531	2.6	7.8
8	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	3,466	2.6	7.6
9	Certain conditions originating in the perinatal period (P00-P96)	2,946	2.2	6.5
10	Influenza and pneumonia (J09-J18)	2,735	2.0	6.0
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2,691	2.0	5.9
12	Alzheimer's disease (G30)	2,471	1.8	5.4
13	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,465	1.8	5.4
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	1,991	1.5	4.4
15	Septicemia (A40-A41)	1,894	1.4	4.2
...	All other causes (Residual)	25,517	18.8	56.1
Hispanic, both sexes, 1-4 years				
...	All causes	1,009	100.0	26.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	348	34.5	9.0
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	132	13.1	3.4
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	76	7.5	2.0
4	Malignant neoplasms (C00-C97)	68	6.7	1.8
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	39	3.9	1.0
6	Influenza and pneumonia (J09-J18)	24	2.4	0.6
7	Septicemia (A40-A41)	18	1.8	*
8	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	17	1.7	*
9	Certain conditions originating in the perinatal period (P00-P96)	16	1.6	*
10	Cerebrovascular diseases (I60-I69)	8	0.8	*
11	Chronic lower respiratory diseases (J40-J47)	6	0.6	*
12	Pneumonitis due to solids and liquids (J69)	5	0.5	*
13	Acute bronchitis and bronchiolitis (J20-J21)	4	0.4	*
13	Complications of medical and surgical care (Y40-Y84,Y88)	4	0.4	*
15	Anemias (D50-D64)	2	0.2	*
15	Meningitis (G00,G03)	2	0.2	*
15	Diseases of appendix (K35-K38)	2	0.2	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2	0.2	*
...	All other causes (Residual)	236	23.4	6.1

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2007

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007; there were no deaths due to J09 in 2007]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, both sexes, 5-14 years				
...	All causes	1,157	100.0	14.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	387	33.4	4.7
2	Malignant neoplasms (C00-C97)	228	19.7	2.8
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	76	6.6	0.9
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	73	6.3	0.9
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	40	3.5	0.5
6	Influenza and pneumonia (J09-J18)	20	1.7	0.2
7	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	19	1.6	*
8	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	17	1.5	*
8	Chronic lower respiratory diseases (J40-J47)	17	1.5	*
10	Septicemia (A40-A41)	16	1.4	*
10	Cerebrovascular diseases (I60-I69)	16	1.4	*
12	Complications of medical and surgical care (Y40-Y84,Y88)	6	0.5	*
13	Anemias (D50-D64)	5	0.4	*
13	Certain conditions originating in the perinatal period (P00-P96)	5	0.4	*
15	Diabetes mellitus (E10-E14)	3	0.3	*
15	Meningitis (G00,G03)	3	0.3	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3	0.3	*
...	All other causes (Residual)	223	19.3	2.7
Hispanic, both sexes, 15-24 years				
...	All causes	5,686	100.0	76.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,511	44.2	33.8
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,295	22.8	17.4
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	527	9.3	7.1
4	Malignant neoplasms (C00-C97)	345	6.1	4.6
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	145	2.6	1.9
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	55	1.0	0.7
7	Cerebrovascular diseases (I60-I69)	37	0.7	0.5
8	Human immunodeficiency virus (HIV) disease (B20-B24)	32	0.6	0.4
9	Pregnancy, childbirth and the puerperium (O00-O99)	28	0.5	0.4
10	Legal intervention (Y35,Y89.0)	24	0.4	0.3
11	Chronic lower respiratory diseases (J40-J47)	20	0.4	0.3
12	Diabetes mellitus (E10-E14)	18	0.3	*
13	Septicemia (A40-A41)	17	0.3	*
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	17	0.3	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	16	0.3	*
...	All other causes (Residual)	599	10.5	8.1

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2007

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007; there were no deaths due to J09 in 2007]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, both sexes, 25-44 years				
...	All causes	15,241	100.0	101.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	4,279	28.1	28.5
2	Malignant neoplasms (C00-C97)	2,094	13.7	14.0
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,530	10.0	10.2
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,340	8.8	8.9
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,148	7.5	7.6
6	Human immunodeficiency virus (HIV) disease (B20-B24)	687	4.5	4.6
7	Chronic liver disease and cirrhosis (K70,K73-K74)	565	3.7	3.8
8	Cerebrovascular diseases (I60-I69)	401	2.6	2.7
9	Diabetes mellitus (E10-E14)	287	1.9	1.9
10	Influenza and pneumonia (J09-J18)	123	0.8	0.8
11	Septicemia (A40-A41)	122	0.8	0.8
12	Viral hepatitis (B15-B19)	118	0.8	0.8
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	103	0.7	0.7
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	100	0.7	0.7
15	Pregnancy, childbirth and the puerperium (O00-O99)	97	0.6	0.6
...	All other causes (Residual)	2,247	14.7	15.0
Hispanic, both sexes, 45-64 years				
...	All causes	31,824	100.0	427.7
1	Malignant neoplasms (C00-C97)	8,880	27.9	119.3
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	6,068	19.1	81.5
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,454	7.7	33.0
4	Chronic liver disease and cirrhosis (K70,K73-K74)	2,121	6.7	28.5
5	Diabetes mellitus (E10-E14)	1,762	5.5	23.7
6	Cerebrovascular diseases (I60-I69)	1,567	4.9	21.1
7	Viral hepatitis (B15-B19)	750	2.4	10.1
8	Human immunodeficiency virus (HIV) disease (B20-B24)	707	2.2	9.5
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	601	1.9	8.1
10	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	582	1.8	7.8
11	Chronic lower respiratory diseases (J40-J47)	438	1.4	5.9
12	Septicemia (A40-A41)	435	1.4	5.8
13	Influenza and pneumonia (J09-J18)	356	1.1	4.8
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	348	1.1	4.7
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	282	0.9	3.8
...	All other causes (Residual)	4,473	14.1	60.1

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2007

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007; there were no deaths due to J09 in 2007]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, both sexes, 65 years and over				
...	All causes	74,507	100.0	2,966.4
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	21,296	28.6	847.9
2	Malignant neoplasms (C00-C97)	16,026	21.5	638.1
3	Cerebrovascular diseases (I60-I69)	5,020	6.7	199.9
4	Diabetes mellitus (E10-E14)	4,343	5.8	172.9
5	Chronic lower respiratory diseases (J40-J47)	2,965	4.0	118.0
6	Alzheimer's disease (G30)	2,420	3.2	96.3
7	Influenza and pneumonia (J09-J18)	2,151	2.9	85.6
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,930	2.6	76.8
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,568	2.1	62.4
10	Septicemia (A40-A41)	1,220	1.6	48.6
10	Chronic liver disease and cirrhosis (K70,K73-K74)	1,220	1.6	48.6
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,063	1.4	42.3
13	Parkinson's disease (G20-G21)	743	1.0	29.6
14	Pneumonitis due to solids and liquids (J69)	440	0.6	17.5
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	434	0.6	17.3
...	All other causes (Residual)	11,668	15.7	464.5
Hispanic, male, all ages				
...	All causes	75,708	100.0	321.8
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	15,657	20.7	66.6
2	Malignant neoplasms (C00-C97)	14,493	19.1	61.6
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	8,844	11.7	37.6
4	Cerebrovascular diseases (I60-I69)	3,319	4.4	14.1
5	Diabetes mellitus (E10-E14)	3,199	4.2	13.6
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	2,926	3.9	12.4
7	Chronic liver disease and cirrhosis (K70,K73-K74)	2,799	3.7	11.9
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,078	2.7	8.8
9	Chronic lower respiratory diseases (J40-J47)	1,894	2.5	8.1
10	Certain conditions originating in the perinatal period (P00-P96)	1,643	2.2	7.0
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,363	1.8	5.8
12	Influenza and pneumonia (J09-J18)	1,361	1.8	5.8
13	Human immunodeficiency virus (HIV) disease (B20-B24)	1,180	1.6	5.0
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	1,063	1.4	4.5
15	Septicemia (A40-A41)	944	1.2	4.0
...	All other causes (Residual)	12,945	17.1	55.0

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2007

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007; there were no deaths due to J09 in 2007]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, male, 1-4 years				
...	All causes	556	100.0	28.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	225	40.5	11.3
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	76	13.7	3.8
3	Malignant neoplasms (C00-C97)	38	6.8	1.9
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	30	5.4	1.5
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	17	3.1	*
6	Influenza and pneumonia (J09-J18)	13	2.3	*
7	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	7	1.3	*
8	Septicemia (A40-A41)	6	1.1	*
9	Certain conditions originating in the perinatal period (P00-P96)	5	0.9	*
10	Cerebrovascular diseases (I60-I69)	4	0.7	*
10	Pneumonitis due to solids and liquids (J69)	4	0.7	*
12	Chronic lower respiratory diseases (J40-J47)	3	0.5	*
13	Meningococcal infection (A39)	1	0.2	*
13	Anemias (D50-D64)	1	0.2	*
13	Diabetes mellitus (E10-E14)	1	0.2	*
13	Meningitis (G00,G03)	1	0.2	*
13	Diseases of appendix (K35-K38)	1	0.2	*
13	Hernia (K40-K46)	1	0.2	*
13	Chronic liver disease and cirrhosis (K70,K73-K74)	1	0.2	*
13	Cholelithiasis and other disorders of gallbladder (K80-K82)	1	0.2	*
13	Infections of kidney (N10-N12,N13.6,N15.1)	1	0.2	*
13	Complications of medical and surgical care (Y40-Y84,Y88)	1	0.2	*
...	All other causes (Residual)	118	21.2	5.9
Hispanic, male, 5-14 years				
...	All causes	663	100.0	15.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	244	36.8	5.8
2	Malignant neoplasms (C00-C97)	114	17.2	2.7
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	48	7.2	1.1
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	35	5.3	0.8
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	23	3.5	0.5
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	13	2.0	*
7	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	11	1.7	*
7	Influenza and pneumonia (J09-J18)	11	1.7	*
9	Septicemia (A40-A41)	10	1.5	*
9	Chronic lower respiratory diseases (J40-J47)	10	1.5	*
11	Cerebrovascular diseases (I60-I69)	8	1.2	*
12	Anemias (D50-D64)	4	0.6	*
13	Certain conditions originating in the perinatal period (P00-P96)	3	0.5	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2	0.3	*
14	Complications of medical and surgical care (Y40-Y84,Y88)	2	0.3	*
...	All other causes (Residual)	125	18.9	3.0

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2007

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007; there were no deaths due to J09 in 2007]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, male, 15-24 years				
...	All causes	4,506	100.0	115.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,035	45.2	52.1
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,171	26.0	30.0
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	451	10.0	11.5
4	Malignant neoplasms (C00-C97)	209	4.6	5.3
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	104	2.3	2.7
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	39	0.9	1.0
7	Legal intervention (Y35,Y89.0)	24	0.5	0.6
8	Human immunodeficiency virus (HIV) disease (B20-B24)	23	0.5	0.6
9	Cerebrovascular diseases (I60-I69)	20	0.4	0.5
10	Diabetes mellitus (E10-E14)	14	0.3	*
11	Chronic lower respiratory diseases (J40-J47)	11	0.2	*
12	Septicemia (A40-A41)	10	0.2	*
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	10	0.2	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	7	0.2	*
15	Aortic aneurysm and dissection (I71)	6	0.1	*
15	Influenza and pneumonia (J09-J18)	6	0.1	*
...	All other causes (Residual)	366	8.1	9.4
Hispanic, male, 25-44 years				
...	All causes	10,993	100.0	135.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,497	31.8	43.0
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,304	11.9	16.0
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,020	9.3	12.5
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	965	8.8	11.9
5	Malignant neoplasms (C00-C97)	915	8.3	11.2
6	Human immunodeficiency virus (HIV) disease (B20-B24)	531	4.8	6.5
7	Chronic liver disease and cirrhosis (K70,K73-K74)	463	4.2	5.7
8	Cerebrovascular diseases (I60-I69)	254	2.3	3.1
9	Diabetes mellitus (E10-E14)	191	1.7	2.3
10	Viral hepatitis (B15-B19)	92	0.8	1.1
11	Influenza and pneumonia (J09-J18)	81	0.7	1.0
12	Septicemia (A40-A41)	68	0.6	0.8
13	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	63	0.6	0.8
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	58	0.5	0.7
15	Legal intervention (Y35,Y89.0)	55	0.5	0.7
...	All other causes (Residual)	1,436	13.1	17.7

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2007

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007; there were no deaths due to J09 in 2007]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, male, 45-64 years				
...	All causes	20,398	100.0	550.1
1	Malignant neoplasms (C00-C97)	4,664	22.9	125.8
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	4,384	21.5	118.2
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,871	9.2	50.5
4	Chronic liver disease and cirrhosis (K70,K73-K74)	1,652	8.1	44.6
5	Diabetes mellitus (E10-E14)	1,044	5.1	28.2
6	Cerebrovascular diseases (I60-I69)	955	4.7	25.8
7	Human immunodeficiency virus (HIV) disease (B20-B24)	553	2.7	14.9
8	Viral hepatitis (B15-B19)	548	2.7	14.8
9	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	479	2.3	12.9
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	351	1.7	9.5
11	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	284	1.4	7.7
12	Chronic lower respiratory diseases (J40-J47)	257	1.3	6.9
13	Septicemia (A40-A41)	256	1.3	6.9
14	Influenza and pneumonia (J09-J18)	222	1.1	6.0
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	168	0.8	4.5
...	All other causes (Residual)	2,710	13.3	73.1
Hispanic, male, 65 years and over				
...	All causes	35,225	100.0	3,285.3
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	10,060	28.6	938.3
2	Malignant neoplasms (C00-C97)	8,546	24.3	797.0
3	Cerebrovascular diseases (I60-I69)	2,065	5.9	192.6
4	Diabetes mellitus (E10-E14)	1,947	5.5	181.6
5	Chronic lower respiratory diseases (J40-J47)	1,559	4.4	145.4
6	Influenza and pneumonia (J09-J18)	999	2.8	93.2
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	922	2.6	86.0
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	871	2.5	81.2
9	Alzheimer's disease (G30)	784	2.2	73.1
10	Chronic liver disease and cirrhosis (K70,K73-K74)	679	1.9	63.3
11	Septicemia (A40-A41)	558	1.6	52.0
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	439	1.2	40.9
13	Parkinson's disease (G20-G21)	420	1.2	39.2
14	Pneumonitis due to solids and liquids (J69)	231	0.7	21.5
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	182	0.5	17.0
...	All other causes (Residual)	4,963	14.1	462.9

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2007

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007; there were no deaths due to J09 in 2007]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, female, all ages				
...	All causes	59,811	100.0	272.1
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	13,364	22.3	60.8
2	Malignant neoplasms (C00-C97)	13,167	22.0	59.9
3	Cerebrovascular diseases (I60-I69)	3,759	6.3	17.1
4	Diabetes mellitus (E10-E14)	3,218	5.4	14.6
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,879	4.8	13.1
6	Alzheimer's disease (G30)	1,670	2.8	7.6
7	Chronic lower respiratory diseases (J40-J47)	1,637	2.7	7.4
8	Influenza and pneumonia (J09-J18)	1,374	2.3	6.3
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,328	2.2	6.0
10	Certain conditions originating in the perinatal period (P00-P96)	1,303	2.2	5.9
11	Chronic liver disease and cirrhosis (K70,K73-K74)	1,114	1.9	5.1
12	Septicemia (A40-A41)	950	1.6	4.3
13	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	928	1.6	4.2
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	756	1.3	3.4
15	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	540	0.9	2.5
...	All other causes (Residual)	11,824	19.8	53.8
Hispanic, female, 1-4 years				
...	All causes	453	100.0	23.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	123	27.2	6.5
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	56	12.4	2.9
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	46	10.2	2.4
4	Malignant neoplasms (C00-C97)	30	6.6	1.6
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	22	4.9	1.2
6	Septicemia (A40-A41)	12	2.6	*
7	Influenza and pneumonia (J09-J18)	11	2.4	*
7	Certain conditions originating in the perinatal period (P00-P96)	11	2.4	*
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	10	2.2	*
10	Cerebrovascular diseases (I60-I69)	4	0.9	*
10	Acute bronchitis and bronchiolitis (J20-J21)	4	0.9	*
12	Chronic lower respiratory diseases (J40-J47)	3	0.7	*
12	Complications of medical and surgical care (Y40-Y84,Y88)	3	0.7	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2	0.4	*
15	Viral hepatitis (B15-B19)	1	0.2	*
15	Anemias (D50-D64)	1	0.2	*
15	Nutritional deficiencies (E40-E64)	1	0.2	*
15	Meningitis (G00,G03)	1	0.2	*
15	Pneumonitis due to solids and liquids (J69)	1	0.2	*
15	Diseases of appendix (K35-K38)	1	0.2	*
...	All other causes (Residual)	110	24.3	5.8

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2007

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007; there were no deaths due to J09 in 2007]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, female, 5-14 years				
...	All causes	494	100.0	12.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	143	28.9	3.6
2	Malignant neoplasms (C00-C97)	114	23.1	2.8
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	38	7.7	0.9
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	28	5.7	0.7
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	17	3.4	*
6	Influenza and pneumonia (J09-J18)	9	1.8	*
7	Cerebrovascular diseases (I60-I69)	8	1.6	*
8	Chronic lower respiratory diseases (J40-J47)	7	1.4	*
9	Septicemia (A40-A41)	6	1.2	*
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	6	1.2	*
9	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	6	1.2	*
12	Complications of medical and surgical care (Y40-Y84,Y88)	4	0.8	*
13	Diabetes mellitus (E10-E14)	2	0.4	*
13	Meningitis (G00,G03)	2	0.4	*
13	Certain conditions originating in the perinatal period (P00-P96)	2	0.4	*
...	All other causes (Residual)	102	20.6	2.5
Hispanic, female, 15-24 years				
...	All causes	1,180	100.0	33.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	476	40.3	13.5
2	Malignant neoplasms (C00-C97)	136	11.5	3.9
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	124	10.5	3.5
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	76	6.4	2.2
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	41	3.5	1.2
6	Pregnancy, childbirth and the puerperium (O00-O99)	28	2.4	0.8
7	Cerebrovascular diseases (I60-I69)	17	1.4	*
8	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	16	1.4	*
9	Human immunodeficiency virus (HIV) disease (B20-B24)	9	0.8	*
9	Chronic lower respiratory diseases (J40-J47)	9	0.8	*
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	9	0.8	*
12	Septicemia (A40-A41)	7	0.6	*
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	7	0.6	*
12	Influenza and pneumonia (J09-J18)	7	0.6	*
15	Anemias (D50-D64)	5	0.4	*
15	Pneumonitis due to solids and liquids (J69)	5	0.4	*
...	All other causes (Residual)	208	17.6	5.9

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2007

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007; there were no deaths due to J09 in 2007]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, female, 25-44 years				
...	All causes	4,248	100.0	61.8
1	Malignant neoplasms (C00-C97)	1,179	27.8	17.1
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	782	18.4	11.4
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	320	7.5	4.7
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	226	5.3	3.3
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	183	4.3	2.7
6	Human immunodeficiency virus (HIV) disease (B20-B24)	156	3.7	2.3
7	Cerebrovascular diseases (I60-I69)	147	3.5	2.1
8	Chronic liver disease and cirrhosis (K70,K73-K74)	102	2.4	1.5
9	Pregnancy, childbirth and the puerperium (O00-O99)	97	2.3	1.4
10	Diabetes mellitus (E10-E14)	96	2.3	1.4
11	Septicemia (A40-A41)	54	1.3	0.8
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	45	1.1	0.7
13	Influenza and pneumonia (J09-J18)	42	1.0	0.6
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	37	0.9	0.5
15	Chronic lower respiratory diseases (J40-J47)	28	0.7	0.4
...	All other causes (Residual)	754	17.7	11.0
Hispanic, female, 45-64 years				
...	All causes	11,426	100.0	306.1
1	Malignant neoplasms (C00-C97)	4,216	36.9	112.9
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,684	14.7	45.1
3	Diabetes mellitus (E10-E14)	718	6.3	19.2
4	Cerebrovascular diseases (I60-I69)	612	5.4	16.4
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	583	5.1	15.6
6	Chronic liver disease and cirrhosis (K70,K73-K74)	469	4.1	12.6
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	250	2.2	6.7
8	Viral hepatitis (B15-B19)	202	1.8	5.4
9	Chronic lower respiratory diseases (J40-J47)	181	1.6	4.8
10	Septicemia (A40-A41)	179	1.6	4.8
11	Human immunodeficiency virus (HIV) disease (B20-B24)	154	1.3	4.1
12	Influenza and pneumonia (J09-J18)	134	1.2	3.6
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	114	1.0	3.1
14	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	103	0.9	2.8
15	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	64	0.6	1.7
...	All other causes (Residual)	1,763	15.4	47.2

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2007

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007; there were no deaths due to J09 in 2007]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, female, 65 years and over				
...	All causes	39,282	100.0	2,728.9
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	11,236	28.6	780.6
2	Malignant neoplasms (C00-C97)	7,480	19.0	519.6
3	Cerebrovascular diseases (I60-I69)	2,955	7.5	205.3
4	Diabetes mellitus (E10-E14)	2,396	6.1	166.4
5	Alzheimer's disease (G30)	1,636	4.2	113.7
6	Chronic lower respiratory diseases (J40-J47)	1,406	3.6	97.7
7	Influenza and pneumonia (J09-J18)	1,152	2.9	80.0
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,008	2.6	70.0
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	697	1.8	48.4
10	Septicemia (A40-A41)	662	1.7	46.0
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	624	1.6	43.3
12	Chronic liver disease and cirrhosis (K70,K73-K74)	541	1.4	37.6
13	Parkinson's disease (G20-G21)	323	0.8	22.4
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	252	0.6	17.5
15	Pneumonitis due to solids and liquids (J69)	209	0.5	14.5
...	All other causes (Residual)	6,705	17.1	465.8
Non-Hispanic, white, both sexes, all ages				
...	All causes	1,939,606	100.0	964.1
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	502,683	25.9	249.9
2	Malignant neoplasms (C00-C97)	456,576	23.5	226.9
3	Chronic lower respiratory diseases (J40-J47)	114,480	5.9	56.9
4	Cerebrovascular diseases (I60-I69)	107,678	5.6	53.5
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	94,584	4.9	47.0
6	Alzheimer's disease (G30)	66,453	3.4	33.0
7	Diabetes mellitus (E10-E14)	50,046	2.6	24.9
8	Influenza and pneumonia (J09-J18)	43,219	2.2	21.5
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	34,219	1.8	17.0
10	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	28,897	1.5	14.4
11	Septicemia (A40-A41)	25,856	1.3	12.9
12	Chronic liver disease and cirrhosis (K70,K73-K74)	21,598	1.1	10.7
13	Parkinson's disease (G20-G21)	18,098	0.9	9.0
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	17,216	0.9	8.6
15	Pneumonitis due to solids and liquids (J69)	14,680	0.8	7.3
...	All other causes (Residual)	343,323	17.7	170.6

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2007

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007; there were no deaths due to J09 in 2007]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, both sexes, 1-4 years				
...	All causes	2,325	100.0	25.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	835	35.9	9.2
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	266	11.4	2.9
3	Malignant neoplasms (C00-C97)	224	9.6	2.5
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	147	6.3	1.6
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	75	3.2	0.8
6	Influenza and pneumonia (J09-J18)	50	2.2	0.5
7	Septicemia (A40-A41)	42	1.8	0.5
8	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	32	1.4	0.4
9	Cerebrovascular diseases (I60-I69)	26	1.1	0.3
10	Certain conditions originating in the perinatal period (P00-P96)	25	1.1	0.3
11	Chronic lower respiratory diseases (J40-J47)	18	0.8	*
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	13	0.6	*
12	Complications of medical and surgical care (Y40-Y84,Y88)	13	0.6	*
14	Meningitis (G00,G03)	10	0.4	*
15	Acute bronchitis and bronchiolitis (J20-J21)	5	0.2	*
...	All other causes (Residual)	544	23.4	6.0
Non-Hispanic, white, both sexes, 5-14 years				
...	All causes	3,362	100.0	14.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,250	37.2	5.3
2	Malignant neoplasms (C00-C97)	559	16.6	2.4
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	202	6.0	0.9
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	127	3.8	0.5
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	119	3.5	0.5
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	107	3.2	0.5
7	Influenza and pneumonia (J09-J18)	58	1.7	0.2
8	Cerebrovascular diseases (I60-I69)	47	1.4	0.2
9	Septicemia (A40-A41)	43	1.3	0.2
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	43	1.3	0.2
11	Chronic lower respiratory diseases (J40-J47)	39	1.2	0.2
12	Meningitis (G00,G03)	16	0.5	*
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	15	0.4	*
14	Anemias (D50-D64)	13	0.4	*
15	Diabetes mellitus (E10-E14)	12	0.4	*
...	All other causes (Residual)	712	21.2	3.0

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2007

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007; there were no deaths due to J09 in 2007]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, both sexes, 15-24 years				
...	All causes	19,607	100.0	74.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	11,032	56.3	41.9
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,932	15.0	11.1
3	Malignant neoplasms (C00-C97)	958	4.9	3.6
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	937	4.8	3.6
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	565	2.9	2.1
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	245	1.2	0.9
7	Influenza and pneumonia (J09-J18)	116	0.6	0.4
8	Cerebrovascular diseases (I60-I69)	102	0.5	0.4
9	Diabetes mellitus (E10-E14)	92	0.5	0.3
10	Septicemia (A40-A41)	89	0.5	0.3
11	Chronic lower respiratory diseases (J40-J47)	59	0.3	0.2
11	Pregnancy, childbirth and the puerperium (O00-O99)	59	0.3	0.2
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	44	0.2	0.2
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	41	0.2	0.2
15	Pneumonitis due to solids and liquids (J69)	30	0.2	0.1
...	All other causes (Residual)	2,306	11.8	8.8
Non-Hispanic, white, both sexes, 25-44 years				
...	All causes	75,723	100.0	144.9
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	22,617	29.9	43.3
2	Malignant neoplasms (C00-C97)	10,890	14.4	20.8
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	9,427	12.4	18.0
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	9,166	12.1	17.5
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	2,126	2.8	4.1
6	Chronic liver disease and cirrhosis (K70,K73-K74)	1,932	2.6	3.7
7	Diabetes mellitus (E10-E14)	1,439	1.9	2.8
8	Cerebrovascular diseases (I60-I69)	1,299	1.7	2.5
9	Human immunodeficiency virus (HIV) disease (B20-B24)	1,206	1.6	2.3
10	Influenza and pneumonia (J09-J18)	685	0.9	1.3
11	Septicemia (A40-A41)	678	0.9	1.3
12	Chronic lower respiratory diseases (J40-J47)	668	0.9	1.3
13	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	643	0.8	1.2
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	448	0.6	0.9
15	Viral hepatitis (B15-B19)	326	0.4	0.6
...	All other causes (Residual)	12,173	16.1	23.3

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2007

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007; there were no deaths due to J09 in 2007]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, both sexes, 45-64 years				
...	All causes	341,774	100.0	601.2
1	Malignant neoplasms (C00-C97)	116,155	34.0	204.3
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	73,626	21.5	129.5
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	24,747	7.2	43.5
4	Chronic lower respiratory diseases (J40-J47)	14,346	4.2	25.2
5	Chronic liver disease and cirrhosis (K70,K73-K74)	11,778	3.4	20.7
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	11,459	3.4	20.2
7	Diabetes mellitus (E10-E14)	10,777	3.2	19.0
8	Cerebrovascular diseases (I60-I69)	10,026	2.9	17.6
9	Septicemia (A40-A41)	4,313	1.3	7.6
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3,645	1.1	6.4
11	Influenza and pneumonia (J09-J18)	3,523	1.0	6.2
12	Viral hepatitis (B15-B19)	3,496	1.0	6.1
13	Human immunodeficiency virus (HIV) disease (B20-B24)	1,762	0.5	3.1
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,707	0.5	3.0
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1,617	0.5	2.8
...	All other causes (Residual)	48,797	14.3	85.8
Non-Hispanic, white, both sexes, 65 years and over				
...	All causes	1,483,736	100.0	4,832.8
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	418,941	28.2	1,364.6
2	Malignant neoplasms (C00-C97)	327,746	22.1	1,067.5
3	Chronic lower respiratory diseases (J40-J47)	99,330	6.7	323.5
4	Cerebrovascular diseases (I60-I69)	96,110	6.5	313.1
5	Alzheimer's disease (G30)	65,733	4.4	214.1
6	Influenza and pneumonia (J09-J18)	38,697	2.6	126.0
7	Diabetes mellitus (E10-E14)	37,720	2.5	122.9
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	33,452	2.3	109.0
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	29,995	2.0	97.7
10	Septicemia (A40-A41)	20,593	1.4	67.1
11	Parkinson's disease (G20-G21)	17,706	1.2	57.7
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	15,306	1.0	49.9
13	Pneumonitis due to solids and liquids (J69)	13,476	0.9	43.9
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	9,884	0.7	32.2
15	Aortic aneurysm and dissection (I71)	8,992	0.6	29.3
...	All other causes (Residual)	250,055	16.9	814.5

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2007

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007; there were no deaths due to J09 in 2007]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, male, all ages				
...	All causes	948,662	100.0	960.4
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	251,229	26.5	254.3
2	Malignant neoplasms (C00-C97)	237,672	25.1	240.6
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	59,274	6.2	60.0
4	Chronic lower respiratory diseases (J40-J47)	53,989	5.7	54.7
5	Cerebrovascular diseases (I60-I69)	41,416	4.4	41.9
6	Diabetes mellitus (E10-E14)	25,584	2.7	25.9
7	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	22,660	2.4	22.9
8	Alzheimer's disease (G30)	19,378	2.0	19.6
9	Influenza and pneumonia (J09-J18)	19,361	2.0	19.6
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	16,896	1.8	17.1
11	Chronic liver disease and cirrhosis (K70,K73-K74)	14,007	1.5	14.2
12	Septicemia (A40-A41)	11,653	1.2	11.8
13	Parkinson's disease (G20-G21)	10,457	1.1	10.6
14	Pneumonitis due to solids and liquids (J69)	7,713	0.8	7.8
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	6,480	0.7	6.6
...	All other causes (Residual)	150,893	15.9	152.8
Non-Hispanic, white, male, 1-4 years				
...	All causes	1,316	100.0	28.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	521	39.6	11.1
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	136	10.3	2.9
3	Malignant neoplasms (C00-C97)	111	8.4	2.4
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	87	6.6	1.9
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	30	2.3	0.6
6	Septicemia (A40-A41)	27	2.1	0.6
7	Influenza and pneumonia (J09-J18)	21	1.6	0.4
8	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	18	1.4	*
9	Certain conditions originating in the perinatal period (P00-P96)	14	1.1	*
10	Cerebrovascular diseases (I60-I69)	12	0.9	*
11	Chronic lower respiratory diseases (J40-J47)	11	0.8	*
12	Meningitis (G00,G03)	6	0.5	*
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	6	0.5	*
14	Complications of medical and surgical care (Y40-Y84,Y88)	5	0.4	*
15	Acute bronchitis and bronchiolitis (J20-J21)	4	0.3	*
...	All other causes (Residual)	307	23.3	6.6

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2007

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007; there were no deaths due to J09 in 2007]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, male, 5-14 years				
...	All causes	1,945	100.0	16.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	763	39.2	6.3
2	Malignant neoplasms (C00-C97)	293	15.1	2.4
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	111	5.7	0.9
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	78	4.0	0.6
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	68	3.5	0.6
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	57	2.9	0.5
7	Septicemia (A40-A41)	29	1.5	0.2
8	Chronic lower respiratory diseases (J40-J47)	26	1.3	0.2
9	Influenza and pneumonia (J09-J18)	25	1.3	0.2
10	Cerebrovascular diseases (I60-I69)	24	1.2	0.2
11	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	20	1.0	0.2
12	Anemias (D50-D64)	9	0.5	*
13	Diabetes mellitus (E10-E14)	8	0.4	*
13	Pneumonitis due to solids and liquids (J69)	8	0.4	*
13	Certain conditions originating in the perinatal period (P00-P96)	8	0.4	*
...	All other causes (Residual)	418	21.5	3.5
Non-Hispanic, white, male, 15-24 years				
...	All causes	14,129	100.0	104.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	8,137	57.6	60.2
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,460	17.4	18.2
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	658	4.7	4.9
4	Malignant neoplasms (C00-C97)	576	4.1	4.3
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	343	2.4	2.5
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	136	1.0	1.0
7	Influenza and pneumonia (J09-J18)	68	0.5	0.5
8	Diabetes mellitus (E10-E14)	60	0.4	0.4
9	Cerebrovascular diseases (I60-I69)	56	0.4	0.4
10	Septicemia (A40-A41)	49	0.3	0.4
11	Chronic lower respiratory diseases (J40-J47)	38	0.3	0.3
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	22	0.2	0.2
12	Pneumonitis due to solids and liquids (J69)	22	0.2	0.2
12	Legal intervention (Y35,Y89.0)	22	0.2	0.2
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	20	0.1	0.1
...	All other causes (Residual)	1,462	10.3	10.8

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2007

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007; there were no deaths due to J09 in 2007]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, male, 25-44 years				
...	All causes	49,277	100.0	187.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	16,246	33.0	61.9
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	7,352	14.9	28.0
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	6,481	13.2	24.7
4	Malignant neoplasms (C00-C97)	4,949	10.0	18.9
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,466	3.0	5.6
6	Chronic liver disease and cirrhosis (K70,K73-K74)	1,232	2.5	4.7
7	Human immunodeficiency virus (HIV) disease (B20-B24)	982	2.0	3.7
8	Diabetes mellitus (E10-E14)	905	1.8	3.4
9	Cerebrovascular diseases (I60-I69)	669	1.4	2.5
10	Influenza and pneumonia (J09-J18)	372	0.8	1.4
11	Septicemia (A40-A41)	353	0.7	1.3
12	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	336	0.7	1.3
13	Chronic lower respiratory diseases (J40-J47)	321	0.7	1.2
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	263	0.5	1.0
15	Viral hepatitis (B15-B19)	205	0.4	0.8
...	All other causes (Residual)	7,145	14.5	27.2
Non-Hispanic, white, male, 45-64 years				
...	All causes	209,908	100.0	749.5
1	Malignant neoplasms (C00-C97)	62,311	29.7	222.5
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	53,108	25.3	189.6
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	16,808	8.0	60.0
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	8,569	4.1	30.6
5	Chronic liver disease and cirrhosis (K70,K73-K74)	8,323	4.0	29.7
6	Chronic lower respiratory diseases (J40-J47)	7,457	3.6	26.6
7	Diabetes mellitus (E10-E14)	6,657	3.2	23.8
8	Cerebrovascular diseases (I60-I69)	5,628	2.7	20.1
9	Viral hepatitis (B15-B19)	2,540	1.2	9.1
10	Septicemia (A40-A41)	2,233	1.1	8.0
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2,096	1.0	7.5
12	Influenza and pneumonia (J09-J18)	2,068	1.0	7.4
13	Human immunodeficiency virus (HIV) disease (B20-B24)	1,469	0.7	5.2
14	Aortic aneurysm and dissection (I71)	1,240	0.6	4.4
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,096	0.5	3.9
...	All other causes (Residual)	28,305	13.5	101.1

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2007

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007; there were no deaths due to J09 in 2007]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, male, 65 years and over				
...	All causes	664,684	100.0	5,098.3
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	191,090	28.7	1,465.7
2	Malignant neoplasms (C00-C97)	169,406	25.5	1,299.4
3	Chronic lower respiratory diseases (J40-J47)	46,126	6.9	353.8
4	Cerebrovascular diseases (I60-I69)	34,987	5.3	268.4
5	Alzheimer's disease (G30)	19,050	2.9	146.1
6	Diabetes mellitus (E10-E14)	17,949	2.7	137.7
7	Influenza and pneumonia (J09-J18)	16,748	2.5	128.5
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	16,424	2.5	126.0
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	14,477	2.2	111.0
10	Parkinson's disease (G20-G21)	10,205	1.5	78.3
11	Septicemia (A40-A41)	8,910	1.3	68.3
12	Pneumonitis due to solids and liquids (J69)	6,950	1.0	53.3
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	5,238	0.8	40.2
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	4,951	0.7	38.0
15	Aortic aneurysm and dissection (I71)	4,872	0.7	37.4
...	All other causes (Residual)	97,301	14.6	746.3
Non-Hispanic, white, female, all ages				
...	All causes	990,944	100.0	967.6
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	251,454	25.4	245.5
2	Malignant neoplasms (C00-C97)	218,904	22.1	213.7
3	Cerebrovascular diseases (I60-I69)	66,262	6.7	64.7
4	Chronic lower respiratory diseases (J40-J47)	60,491	6.1	59.1
5	Alzheimer's disease (G30)	47,075	4.8	46.0
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	35,310	3.6	34.5
7	Diabetes mellitus (E10-E14)	24,462	2.5	23.9
8	Influenza and pneumonia (J09-J18)	23,858	2.4	23.3
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	17,323	1.7	16.9
10	Septicemia (A40-A41)	14,203	1.4	13.9
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	10,736	1.1	10.5
12	Parkinson's disease (G20-G21)	7,641	0.8	7.5
13	Chronic liver disease and cirrhosis (K70,K73-K74)	7,591	0.8	7.4
14	Pneumonitis due to solids and liquids (J69)	6,967	0.7	6.8
15	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	6,237	0.6	6.1
...	All other causes (Residual)	192,430	19.4	187.9

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2007

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007; there were no deaths due to J09 in 2007]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, female, 1-4 years				
...	All causes	1,009	100.0	22.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	314	31.1	7.1
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	130	12.9	2.9
3	Malignant neoplasms (C00-C97)	113	11.2	2.5
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	60	5.9	1.4
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	45	4.5	1.0
6	Influenza and pneumonia (J09-J18)	29	2.9	0.7
7	Septicemia (A40-A41)	15	1.5	*
8	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	14	1.4	*
8	Cerebrovascular diseases (I60-I69)	14	1.4	*
10	Certain conditions originating in the perinatal period (P00-P96)	11	1.1	*
11	Complications of medical and surgical care (Y40-Y84,Y88)	8	0.8	*
12	Chronic lower respiratory diseases (J40-J47)	7	0.7	*
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	7	0.7	*
14	Meningitis (G00,G03)	4	0.4	*
15	Anemias (D50-D64)	2	0.2	*
15	Chronic liver disease and cirrhosis (K70,K73-K74)	2	0.2	*
...	All other causes (Residual)	234	23.2	5.3
Non-Hispanic, white, female, 5-14 years				
...	All causes	1,417	100.0	12.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	487	34.4	4.2
2	Malignant neoplasms (C00-C97)	266	18.8	2.3
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	91	6.4	0.8
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	62	4.4	0.5
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	59	4.2	0.5
6	Influenza and pneumonia (J09-J18)	33	2.3	0.3
7	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	29	2.0	0.3
8	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	23	1.6	0.2
8	Cerebrovascular diseases (I60-I69)	23	1.6	0.2
10	Septicemia (A40-A41)	14	1.0	*
11	Chronic lower respiratory diseases (J40-J47)	13	0.9	*
12	Meningitis (G00,G03)	10	0.7	*
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	9	0.6	*
14	Anemias (D50-D64)	4	0.3	*
14	Diabetes mellitus (E10-E14)	4	0.3	*
...	All other causes (Residual)	290	20.5	2.5

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2007

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007; there were no deaths due to J09 in 2007]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, female, 15-24 years				
...	All causes	5,478	100.0	42.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,895	52.8	22.6
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	472	8.6	3.7
3	Malignant neoplasms (C00-C97)	382	7.0	3.0
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	279	5.1	2.2
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	222	4.1	1.7
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	109	2.0	0.9
7	Pregnancy, childbirth and the puerperium (O00-O99)	59	1.1	0.5
8	Influenza and pneumonia (J09-J18)	48	0.9	0.4
9	Cerebrovascular diseases (I60-I69)	46	0.8	0.4
10	Septicemia (A40-A41)	40	0.7	0.3
11	Diabetes mellitus (E10-E14)	32	0.6	0.2
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	22	0.4	0.2
13	Chronic lower respiratory diseases (J40-J47)	21	0.4	0.2
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	21	0.4	0.2
15	Chronic liver disease and cirrhosis (K70,K73-K74)	11	0.2	*
...	All other causes (Residual)	819	15.0	6.4
Non-Hispanic, white, female, 25-44 years				
...	All causes	26,446	100.0	101.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	6,371	24.1	24.5
2	Malignant neoplasms (C00-C97)	5,941	22.5	22.8
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,685	10.2	10.3
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,075	7.8	8.0
5	Chronic liver disease and cirrhosis (K70,K73-K74)	700	2.6	2.7
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	660	2.5	2.5
7	Cerebrovascular diseases (I60-I69)	630	2.4	2.4
8	Diabetes mellitus (E10-E14)	534	2.0	2.1
9	Chronic lower respiratory diseases (J40-J47)	347	1.3	1.3
10	Septicemia (A40-A41)	325	1.2	1.2
11	Influenza and pneumonia (J09-J18)	313	1.2	1.2
12	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	307	1.2	1.2
13	Pregnancy, childbirth and the puerperium (O00-O99)	235	0.9	0.9
14	Human immunodeficiency virus (HIV) disease (B20-B24)	224	0.8	0.9
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	185	0.7	0.7
...	All other causes (Residual)	4,914	18.6	18.9

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2007

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007; there were no deaths due to J09 in 2007]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, female, 45-64 years				
...	All causes	131,866	100.0	457.2
1	Malignant neoplasms (C00-C97)	53,844	40.8	186.7
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	20,518	15.6	71.1
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	7,939	6.0	27.5
4	Chronic lower respiratory diseases (J40-J47)	6,889	5.2	23.9
5	Cerebrovascular diseases (I60-I69)	4,398	3.3	15.2
6	Diabetes mellitus (E10-E14)	4,120	3.1	14.3
7	Chronic liver disease and cirrhosis (K70,K73-K74)	3,455	2.6	12.0
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,890	2.2	10.0
9	Septicemia (A40-A41)	2,080	1.6	7.2
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,549	1.2	5.4
11	Influenza and pneumonia (J09-J18)	1,455	1.1	5.0
12	Viral hepatitis (B15-B19)	956	0.7	3.3
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	675	0.5	2.3
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	611	0.5	2.1
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	516	0.4	1.8
...	All other causes (Residual)	19,971	15.1	69.2
Non-Hispanic, white, female, 65 years and over				
...	All causes	819,052	100.0	4,636.9
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	227,851	27.8	1,289.9
2	Malignant neoplasms (C00-C97)	158,340	19.3	896.4
3	Cerebrovascular diseases (I60-I69)	61,123	7.5	346.0
4	Chronic lower respiratory diseases (J40-J47)	53,204	6.5	301.2
5	Alzheimer's disease (G30)	46,683	5.7	264.3
6	Influenza and pneumonia (J09-J18)	21,949	2.7	124.3
7	Diabetes mellitus (E10-E14)	19,771	2.4	111.9
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	17,028	2.1	96.4
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	15,518	1.9	87.9
10	Septicemia (A40-A41)	11,683	1.4	66.1
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	10,068	1.2	57.0
12	Parkinson's disease (G20-G21)	7,501	0.9	42.5
13	Pneumonitis due to solids and liquids (J69)	6,526	0.8	36.9
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	4,933	0.6	27.9
15	Atherosclerosis (I70)	4,240	0.5	24.0
...	All other causes (Residual)	152,634	18.6	864.1

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2007

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007; there were no deaths due to J09 in 2007]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, both sexes, all ages				
...	All causes	286,366	100.0	750.7
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	70,443	24.6	184.7
2	Malignant neoplasms (C00-C97)	63,441	22.2	166.3
3	Cerebrovascular diseases (I60-I69)	16,934	5.9	44.4
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	13,332	4.7	34.9
5	Diabetes mellitus (E10-E14)	12,343	4.3	32.4
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	8,746	3.1	22.9
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	8,318	2.9	21.8
8	Chronic lower respiratory diseases (J40-J47)	7,830	2.7	20.5
9	Human immunodeficiency virus (HIV) disease (B20-B24)	6,398	2.2	16.8
10	Septicemia (A40-A41)	6,241	2.2	16.4
11	Influenza and pneumonia (J09-J18)	5,091	1.8	13.3
12	Certain conditions originating in the perinatal period (P00-P96)	4,868	1.7	12.8
13	Alzheimer's disease (G30)	4,729	1.7	12.4
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	4,597	1.6	12.1
15	Chronic liver disease and cirrhosis (K70,K73-K74)	2,525	0.9	6.6
...	All other causes (Residual)	50,530	17.6	132.5
Non-Hispanic, black, both sexes, 1-4 years				
...	All causes	1,102	100.0	43.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	318	28.9	12.6
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	160	14.5	6.3
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	116	10.5	4.6
4	Malignant neoplasms (C00-C97)	51	4.6	2.0
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	42	3.8	1.7
6	Chronic lower respiratory diseases (J40-J47)	29	2.6	1.2
7	Influenza and pneumonia (J09-J18)	28	2.5	1.1
8	Certain conditions originating in the perinatal period (P00-P96)	26	2.4	1.0
9	Septicemia (A40-A41)	16	1.5	*
9	Cerebrovascular diseases (I60-I69)	16	1.5	*
11	Anemias (D50-D64)	12	1.1	*
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	9	0.8	*
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	6	0.5	*
13	Complications of medical and surgical care (Y40-Y84,Y88)	6	0.5	*
15	Acute bronchitis and bronchiolitis (J20-J21)	5	0.5	*
...	All other causes (Residual)	262	23.8	10.4

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2007

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007; there were no deaths due to J09 in 2007]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, both sexes, 5-14 years				
...	All causes	1,335	100.0	21.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	452	33.9	7.3
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	138	10.3	2.2
3	Malignant neoplasms (C00-C97)	134	10.0	2.2
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	77	5.8	1.2
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	63	4.7	1.0
6	Chronic lower respiratory diseases (J40-J47)	59	4.4	1.0
7	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	41	3.1	0.7
8	Influenza and pneumonia (J09-J18)	21	1.6	0.3
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	20	1.5	0.3
10	Anemias (D50-D64)	13	1.0	*
10	Cerebrovascular diseases (I60-I69)	13	1.0	*
12	Septicemia (A40-A41)	10	0.7	*
13	Human immunodeficiency virus (HIV) disease (B20-B24)	8	0.6	*
14	Diabetes mellitus (E10-E14)	6	0.4	*
14	Certain conditions originating in the perinatal period (P00-P96)	6	0.4	*
14	Complications of medical and surgical care (Y40-Y84,Y88)	6	0.4	*
...	All other causes (Residual)	268	20.1	4.3
Non-Hispanic, black, both sexes, 15-24 years				
...	All causes	7,237	100.0	112.6
1	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	3,162	43.7	49.2
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,699	23.5	26.4
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	396	5.5	6.2
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	344	4.8	5.4
5	Malignant neoplasms (C00-C97)	258	3.6	4.0
6	Human immunodeficiency virus (HIV) disease (B20-B24)	109	1.5	1.7
7	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	81	1.1	1.3
8	Pregnancy, childbirth and the puerperium (O00-O99)	68	0.9	1.1
9	Anemias (D50-D64)	66	0.9	1.0
10	Chronic lower respiratory diseases (J40-J47)	65	0.9	1.0
11	Diabetes mellitus (E10-E14)	55	0.8	0.9
12	Septicemia (A40-A41)	47	0.6	0.7
12	Legal intervention (Y35,Y89.0)	47	0.6	0.7
14	Cerebrovascular diseases (I60-I69)	45	0.6	0.7
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	28	0.4	0.4
...	All other causes (Residual)	767	10.6	11.9

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2007

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007; there were no deaths due to J09 in 2007]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, both sexes, 25-44 years				
...	All causes	26,318	100.0	243.5
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,947	15.0	36.5
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	3,879	14.7	35.9
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,866	14.7	35.8
4	Malignant neoplasms (C00-C97)	2,944	11.2	27.2
5	Human immunodeficiency virus (HIV) disease (B20-B24)	2,691	10.2	24.9
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	886	3.4	8.2
7	Cerebrovascular diseases (I60-I69)	796	3.0	7.4
8	Diabetes mellitus (E10-E14)	766	2.9	7.1
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	439	1.7	4.1
10	Septicemia (A40-A41)	362	1.4	3.3
11	Chronic lower respiratory diseases (J40-J47)	286	1.1	2.6
12	Influenza and pneumonia (J09-J18)	267	1.0	2.5
13	Anemias (D50-D64)	240	0.9	2.2
14	Chronic liver disease and cirrhosis (K70,K73-K74)	226	0.9	2.1
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	223	0.8	2.1
...	All other causes (Residual)	4,500	17.1	41.6
Non-Hispanic, black, both sexes, 45-64 years				
...	All causes	83,664	100.0	998.8
1	Malignant neoplasms (C00-C97)	23,376	27.9	279.1
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	20,441	24.4	244.0
3	Cerebrovascular diseases (I60-I69)	4,498	5.4	53.7
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	4,308	5.1	51.4
5	Diabetes mellitus (E10-E14)	3,872	4.6	46.2
6	Human immunodeficiency virus (HIV) disease (B20-B24)	3,298	3.9	39.4
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2,192	2.6	26.2
8	Chronic lower respiratory diseases (J40-J47)	1,835	2.2	21.9
9	Septicemia (A40-A41)	1,711	2.0	20.4
10	Chronic liver disease and cirrhosis (K70,K73-K74)	1,700	2.0	20.3
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,276	1.5	15.2
12	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,112	1.3	13.3
13	Influenza and pneumonia (J09-J18)	1,001	1.2	11.9
14	Viral hepatitis (B15-B19)	774	0.9	9.2
15	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	470	0.6	5.6
...	All other causes (Residual)	11,800	14.1	140.9

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2007

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007; there were no deaths due to J09 in 2007]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, both sexes, 65 years and over				
...	All causes	158,050	100.0	4,968.6
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	45,467	28.8	1,429.3
2	Malignant neoplasms (C00-C97)	36,665	23.2	1,152.6
3	Cerebrovascular diseases (I60-I69)	11,531	7.3	362.5
4	Diabetes mellitus (E10-E14)	7,641	4.8	240.2
5	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	5,606	3.5	176.2
6	Chronic lower respiratory diseases (J40-J47)	5,542	3.5	174.2
7	Alzheimer's disease (G30)	4,680	3.0	147.1
8	Septicemia (A40-A41)	3,985	2.5	125.3
9	Influenza and pneumonia (J09-J18)	3,677	2.3	115.6
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	3,087	2.0	97.0
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,267	1.4	71.3
12	Pneumonitis due to solids and liquids (J69)	1,140	0.7	35.8
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	753	0.5	23.7
14	Parkinson's disease (G20-G21)	748	0.5	23.5
15	Aortic aneurysm and dissection (I71)	663	0.4	20.8
...	All other causes (Residual)	24,598	15.6	773.3
Non-Hispanic, black, male, all ages				
...	All causes	146,474	100.0	805.1
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	35,243	24.1	193.7
2	Malignant neoplasms (C00-C97)	32,766	22.4	180.1
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	9,093	6.2	50.0
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	7,477	5.1	41.1
5	Cerebrovascular diseases (I60-I69)	7,470	5.1	41.1
6	Diabetes mellitus (E10-E14)	5,434	3.7	29.9
7	Chronic lower respiratory diseases (J40-J47)	4,167	2.8	22.9
8	Human immunodeficiency virus (HIV) disease (B20-B24)	4,127	2.8	22.7
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3,730	2.5	20.5
10	Septicemia (A40-A41)	2,807	1.9	15.4
11	Certain conditions originating in the perinatal period (P00-P96)	2,734	1.9	15.0
12	Influenza and pneumonia (J09-J18)	2,464	1.7	13.5
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,963	1.3	10.8
14	Chronic liver disease and cirrhosis (K70,K73-K74)	1,662	1.1	9.1
15	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,571	1.1	8.6
...	All other causes (Residual)	23,766	16.2	130.6

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2007

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007; there were no deaths due to J09 in 2007]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, male, 1-4 years				
...	All causes	609	100.0	47.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	186	30.5	14.5
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	88	14.4	6.9
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	57	9.4	4.4
4	Malignant neoplasms (C00-C97)	31	5.1	2.4
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	28	4.6	2.2
6	Influenza and pneumonia (J09-J18)	15	2.5	*
7	Chronic lower respiratory diseases (J40-J47)	13	2.1	*
8	Certain conditions originating in the perinatal period (P00-P96)	12	2.0	*
9	Septicemia (A40-A41)	11	1.8	*
10	Anemias (D50-D64)	4	0.7	*
10	Cerebrovascular diseases (I60-I69)	4	0.7	*
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	4	0.7	*
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	3	0.5	*
13	Meningitis (G00,G03)	3	0.5	*
13	Acute bronchitis and bronchiolitis (J20-J21)	3	0.5	*
13	Complications of medical and surgical care (Y40-Y84,Y88)	3	0.5	*
...	All other causes (Residual)	144	23.6	11.2
Non-Hispanic, black, male, 5-14 years				
...	All causes	801	100.0	25.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	275	34.3	8.8
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	94	11.7	3.0
3	Malignant neoplasms (C00-C97)	70	8.7	2.2
4	Chronic lower respiratory diseases (J40-J47)	40	5.0	1.3
5	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	37	4.6	1.2
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	35	4.4	1.1
7	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	29	3.6	0.9
8	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	14	1.7	*
9	Influenza and pneumonia (J09-J18)	13	1.6	*
10	Anemias (D50-D64)	8	1.0	*
11	Septicemia (A40-A41)	4	0.5	*
11	Meningitis (G00,G03)	4	0.5	*
13	Cerebrovascular diseases (I60-I69)	3	0.4	*
13	Diseases of appendix (K35-K38)	3	0.4	*
13	Certain conditions originating in the perinatal period (P00-P96)	3	0.4	*
...	All other causes (Residual)	169	21.1	5.4

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2007

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007; there were no deaths due to J09 in 2007]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, male, 15-24 years				
...	All causes	5,635	100.0	173.1
1	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	2,871	50.9	88.2
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,267	22.5	38.9
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	343	6.1	10.5
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	233	4.1	7.2
5	Malignant neoplasms (C00-C97)	142	2.5	4.4
6	Human immunodeficiency virus (HIV) disease (B20-B24)	56	1.0	1.7
7	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	52	0.9	1.6
8	Legal intervention (Y35,Y89.0)	45	0.8	1.4
9	Chronic lower respiratory diseases (J40-J47)	44	0.8	1.4
10	Anemias (D50-D64)	36	0.6	1.1
11	Cerebrovascular diseases (I60-I69)	30	0.5	0.9
12	Septicemia (A40-A41)	22	0.4	0.7
13	Diabetes mellitus (E10-E14)	21	0.4	0.6
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	14	0.2	*
15	Influenza and pneumonia (J09-J18)	12	0.2	*
...	All other causes (Residual)	447	7.9	13.7
Non-Hispanic, black, male, 25-44 years				
...	All causes	16,519	100.0	321.0
1	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	3,339	20.2	64.9
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,873	17.4	55.8
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,552	15.4	49.6
4	Human immunodeficiency virus (HIV) disease (B20-B24)	1,589	9.6	30.9
5	Malignant neoplasms (C00-C97)	1,155	7.0	22.4
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	728	4.4	14.1
7	Diabetes mellitus (E10-E14)	425	2.6	8.3
8	Cerebrovascular diseases (I60-I69)	401	2.4	7.8
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	235	1.4	4.6
10	Septicemia (A40-A41)	166	1.0	3.2
11	Chronic lower respiratory diseases (J40-J47)	155	0.9	3.0
12	Influenza and pneumonia (J09-J18)	151	0.9	2.9
13	Chronic liver disease and cirrhosis (K70,K73-K74)	135	0.8	2.6
14	Anemias (D50-D64)	130	0.8	2.5
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	123	0.7	2.4
...	All other causes (Residual)	2,362	14.3	45.9

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2007

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007; there were no deaths due to J09 in 2007]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, male, 45-64 years				
...	All causes	49,083	100.0	1,286.6
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	12,924	26.3	338.8
2	Malignant neoplasms (C00-C97)	12,447	25.4	326.3
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,057	6.2	80.1
4	Cerebrovascular diseases (I60-I69)	2,480	5.1	65.0
5	Human immunodeficiency virus (HIV) disease (B20-B24)	2,264	4.6	59.3
6	Diabetes mellitus (E10-E14)	2,060	4.2	54.0
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,153	2.3	30.2
8	Chronic liver disease and cirrhosis (K70,K73-K74)	1,150	2.3	30.1
9	Chronic lower respiratory diseases (J40-J47)	966	2.0	25.3
10	Septicemia (A40-A41)	900	1.8	23.6
11	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	883	1.8	23.1
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	707	1.4	18.5
13	Influenza and pneumonia (J09-J18)	595	1.2	15.6
14	Viral hepatitis (B15-B19)	536	1.1	14.0
15	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	363	0.7	9.5
...	All other causes (Residual)	6,598	13.4	172.9
Non-Hispanic, black, male, 65 years and over				
...	All causes	69,015	100.0	5,645.6
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	19,390	28.1	1,586.1
2	Malignant neoplasms (C00-C97)	18,913	27.4	1,547.1
3	Cerebrovascular diseases (I60-I69)	4,527	6.6	370.3
4	Chronic lower respiratory diseases (J40-J47)	2,937	4.3	240.3
5	Diabetes mellitus (E10-E14)	2,924	4.2	239.2
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2,296	3.3	187.8
7	Septicemia (A40-A41)	1,644	2.4	134.5
8	Influenza and pneumonia (J09-J18)	1,636	2.4	133.8
9	Alzheimer's disease (G30)	1,272	1.8	104.1
10	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,216	1.8	99.5
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,127	1.6	92.2
12	Pneumonitis due to solids and liquids (J69)	558	0.8	45.6
13	Parkinson's disease (G20-G21)	407	0.6	33.3
14	Chronic liver disease and cirrhosis (K70,K73-K74)	372	0.5	30.4
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	348	0.5	28.5
...	All other causes (Residual)	9,448	13.7	772.9

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2007

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007; there were no deaths due to J09 in 2007]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, female, all ages				
...	All causes	139,892	100.0	701.0
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	35,200	25.2	176.4
2	Malignant neoplasms (C00-C97)	30,675	21.9	153.7
3	Cerebrovascular diseases (I60-I69)	9,464	6.8	47.4
4	Diabetes mellitus (E10-E14)	6,909	4.9	34.6
5	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	4,588	3.3	23.0
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	4,239	3.0	21.2
7	Chronic lower respiratory diseases (J40-J47)	3,663	2.6	18.4
8	Alzheimer's disease (G30)	3,436	2.5	17.2
9	Septicemia (A40-A41)	3,434	2.5	17.2
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	2,634	1.9	13.2
11	Influenza and pneumonia (J09-J18)	2,627	1.9	13.2
12	Human immunodeficiency virus (HIV) disease (B20-B24)	2,271	1.6	11.4
13	Certain conditions originating in the perinatal period (P00-P96)	2,134	1.5	10.7
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,269	0.9	6.4
15	Chronic liver disease and cirrhosis (K70,K73-K74)	863	0.6	4.3
...	All other causes (Residual)	26,486	18.9	132.7
Non-Hispanic, black, female, 1-4 years				
...	All causes	493	100.0	39.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	132	26.8	10.6
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	72	14.6	5.8
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	59	12.0	4.8
4	Malignant neoplasms (C00-C97)	20	4.1	1.6
5	Chronic lower respiratory diseases (J40-J47)	16	3.2	*
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	14	2.8	*
6	Certain conditions originating in the perinatal period (P00-P96)	14	2.8	*
8	Influenza and pneumonia (J09-J18)	13	2.6	*
9	Cerebrovascular diseases (I60-I69)	12	2.4	*
10	Anemias (D50-D64)	8	1.6	*
11	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	6	1.2	*
12	Septicemia (A40-A41)	5	1.0	*
13	Complications of medical and surgical care (Y40-Y84,Y88)	3	0.6	*
14	Acute bronchitis and bronchiolitis (J20-J21)	2	0.4	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2	0.4	*
...	All other causes (Residual)	115	23.3	9.3

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2007

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007; there were no deaths due to J09 in 2007]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, female, 5-14 years				
...	All causes	534	100.0	17.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	177	33.1	5.8
2	Malignant neoplasms (C00-C97)	64	12.0	2.1
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	44	8.2	1.4
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	40	7.5	1.3
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	28	5.2	0.9
6	Chronic lower respiratory diseases (J40-J47)	19	3.6	*
7	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	12	2.2	*
8	Cerebrovascular diseases (I60-I69)	10	1.9	*
9	Influenza and pneumonia (J09-J18)	8	1.5	*
10	Septicemia (A40-A41)	6	1.1	*
10	Human immunodeficiency virus (HIV) disease (B20-B24)	6	1.1	*
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	6	1.1	*
13	Anemias (D50-D64)	5	0.9	*
14	Diabetes mellitus (E10-E14)	4	0.7	*
14	Complications of medical and surgical care (Y40-Y84,Y88)	4	0.7	*
...	All other causes (Residual)	101	18.9	3.3
Non-Hispanic, black, female, 15-24 years				
...	All causes	1,602	100.0	50.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	432	27.0	13.6
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	291	18.2	9.2
3	Malignant neoplasms (C00-C97)	116	7.2	3.7
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	111	6.9	3.5
5	Pregnancy, childbirth and the puerperium (O00-O99)	68	4.2	2.1
6	Human immunodeficiency virus (HIV) disease (B20-B24)	53	3.3	1.7
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	53	3.3	1.7
8	Diabetes mellitus (E10-E14)	34	2.1	1.1
9	Anemias (D50-D64)	30	1.9	0.9
10	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	29	1.8	0.9
11	Septicemia (A40-A41)	25	1.6	0.8
12	Chronic lower respiratory diseases (J40-J47)	21	1.3	0.7
13	Cerebrovascular diseases (I60-I69)	15	0.9	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	14	0.9	*
15	Influenza and pneumonia (J09-J18)	11	0.7	*
...	All other causes (Residual)	299	18.7	9.4

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2007

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007; there were no deaths due to J09 in 2007]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, female, 25-44 years				
...	All causes	9,799	100.0	173.1
1	Malignant neoplasms (C00-C97)	1,789	18.3	31.6
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,395	14.2	24.6
3	Human immunodeficiency virus (HIV) disease (B20-B24)	1,102	11.2	19.5
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	993	10.1	17.5
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	540	5.5	9.5
6	Cerebrovascular diseases (I60-I69)	395	4.0	7.0
7	Diabetes mellitus (E10-E14)	341	3.5	6.0
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	204	2.1	3.6
9	Septicemia (A40-A41)	196	2.0	3.5
10	Pregnancy, childbirth and the puerperium (O00-O99)	167	1.7	3.0
11	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	158	1.6	2.8
12	Chronic lower respiratory diseases (J40-J47)	131	1.3	2.3
13	Influenza and pneumonia (J09-J18)	116	1.2	2.0
14	Anemias (D50-D64)	110	1.1	1.9
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	100	1.0	1.8
...	All other causes (Residual)	2,062	21.0	36.4
Non-Hispanic, black, female, 45-64 years				
...	All causes	34,581	100.0	758.1
1	Malignant neoplasms (C00-C97)	10,929	31.6	239.6
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	7,517	21.7	164.8
3	Cerebrovascular diseases (I60-I69)	2,018	5.8	44.2
4	Diabetes mellitus (E10-E14)	1,812	5.2	39.7
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,251	3.6	27.4
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,039	3.0	22.8
7	Human immunodeficiency virus (HIV) disease (B20-B24)	1,034	3.0	22.7
8	Chronic lower respiratory diseases (J40-J47)	869	2.5	19.0
9	Septicemia (A40-A41)	811	2.3	17.8
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	569	1.6	12.5
11	Chronic liver disease and cirrhosis (K70,K73-K74)	550	1.6	12.1
12	Influenza and pneumonia (J09-J18)	406	1.2	8.9
13	Viral hepatitis (B15-B19)	238	0.7	5.2
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	229	0.7	5.0
15	Aortic aneurysm and dissection (I71)	141	0.4	3.1
...	All other causes (Residual)	5,168	14.9	113.3

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2007

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007; there were no deaths due to J09 in 2007]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, female, 65 years and over				
...	All causes	89,035	100.0	4,546.0
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	26,077	29.3	1,331.5
2	Malignant neoplasms (C00-C97)	17,752	19.9	906.4
3	Cerebrovascular diseases (I60-I69)	7,004	7.9	357.6
4	Diabetes mellitus (E10-E14)	4,717	5.3	240.8
5	Alzheimer's disease (G30)	3,408	3.8	174.0
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3,310	3.7	169.0
7	Chronic lower respiratory diseases (J40-J47)	2,605	2.9	133.0
8	Septicemia (A40-A41)	2,341	2.6	119.5
9	Influenza and pneumonia (J09-J18)	2,041	2.3	104.2
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,960	2.2	100.1
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,051	1.2	53.7
12	Pneumonitis due to solids and liquids (J69)	582	0.7	29.7
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	405	0.5	20.7
14	Aortic aneurysm and dissection (I71)	371	0.4	18.9
15	Atherosclerosis (I70)	362	0.4	18.5
...	All other causes (Residual)	15,049	16.9	768.4

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable