

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, both sexes, all ages				
...	All causes	163,241	100.0	301.9
1	Malignant neoplasms (C00-C97)	35,147	21.5	65.0
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	33,243	20.4	61.5
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	12,015	7.4	22.2
4	Cerebrovascular diseases (I60-I69)	8,127	5.0	15.0
5	Diabetes mellitus (E10-E14)	7,632	4.7	14.1
6	Chronic liver disease and cirrhosis (K70,K73-K74)	5,141	3.1	9.5
7	Chronic lower respiratory diseases (J40-J47)	4,827	3.0	8.9
8	Alzheimer's disease (G30)	4,127	2.5	7.6
9	Influenza and pneumonia (J09-J18)	3,592	2.2	6.6
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3,083	1.9	5.7
11	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,865	1.8	5.3
12	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	2,571	1.6	4.8
13	Septicemia (A40-A41)	2,435	1.5	4.5
14	Certain conditions originating in the perinatal period (P00-P96)	2,428	1.5	4.5
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	2,092	1.3	3.9
...	All other causes (Residual)	33,916	20.8	62.7
Hispanic, both sexes, 1-4 years				
...	All causes	854	100.0	20.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	278	32.6	6.8
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	122	14.3	3.0
3	Malignant neoplasms (C00-C97)	90	10.5	2.2
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	53	6.2	1.3
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	23	2.7	0.6
6	Influenza and pneumonia (J09-J18)	19	2.2	*
7	Septicemia (A40-A41)	9	1.1	*
7	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	9	1.1	*
7	Chronic lower respiratory diseases (J40-J47)	9	1.1	*
7	Certain conditions originating in the perinatal period (P00-P96)	9	1.1	*
11	Anemias (D50-D64)	5	0.6	*
11	Meningitis (G00,G03)	5	0.6	*
13	Cerebrovascular diseases (I60-I69)	4	0.5	*
14	Diseases of appendix (K35-K38)	3	0.4	*
14	Complications of medical and surgical care (Y40-Y84,Y88)	3	0.4	*
...	All other causes (Residual)	213	24.9	5.2

... Category not applicable

\* Figure does not meet standards of reliability or precision

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality, 2013

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, both sexes, 5-14 years				
...	All causes	1,073	100.0	10.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	279	26.0	2.8
2	Malignant neoplasms (C00-C97)	209	19.5	2.1
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	79	7.4	0.8
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	71	6.6	0.7
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	59	5.5	0.6
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	31	2.9	0.3
7	Cerebrovascular diseases (I60-I69)	26	2.4	0.3
8	Influenza and pneumonia (J09-J18)	22	2.1	0.2
8	Chronic lower respiratory diseases (J40-J47)	22	2.1	0.2
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	12	1.1	*
11	Septicemia (A40-A41)	10	0.9	*
12	Complications of medical and surgical care (Y40-Y84,Y88)	7	0.7	*
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	5	0.5	*
13	Certain conditions originating in the perinatal period (P00-P96)	5	0.5	*
15	Diabetes mellitus (E10-E14)	4	0.4	*
...	All other causes (Residual)	232	21.6	2.3
Hispanic, both sexes, 15-24 years				
...	All causes	4,657	100.0	50.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,944	41.7	21.0
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	849	18.2	9.2
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	635	13.6	6.8
4	Malignant neoplasms (C00-C97)	339	7.3	3.7
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	120	2.6	1.3
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	55	1.2	0.6
7	Cerebrovascular diseases (I60-I69)	32	0.7	0.3
8	Influenza and pneumonia (J09-J18)	27	0.6	0.3
8	Pregnancy, childbirth and the puerperium (O00-O99)	27	0.6	0.3
10	Septicemia (A40-A41)	23	0.5	0.2
11	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	21	0.5	0.2
12	Diabetes mellitus (E10-E14)	18	0.4	*
13	Legal intervention (Y35,Y89.0)	17	0.4	*
14	Human immunodeficiency virus (HIV) disease (B20-B24)	16	0.3	*
15	Chronic lower respiratory diseases (J40-J47)	14	0.3	*
...	All other causes (Residual)	520	11.2	5.6

... Category not applicable

\* Figure does not meet standards of reliability or precision

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality, 2013

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, both sexes, 25-44 years				
...	All causes	15,039	100.0	90.9
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	4,117	27.4	24.9
2	Malignant neoplasms (C00-C97)	2,343	15.6	14.2
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,367	9.1	8.3
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,212	8.1	7.3
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,138	7.6	6.9
6	Chronic liver disease and cirrhosis (K70,K73-K74)	611	4.1	3.7
7	Cerebrovascular diseases (I60-I69)	414	2.8	2.5
8	Diabetes mellitus (E10-E14)	354	2.4	2.1
9	Human immunodeficiency virus (HIV) disease (B20-B24)	295	2.0	1.8
10	Influenza and pneumonia (J09-J18)	193	1.3	1.2
11	Septicemia (A40-A41)	142	0.9	0.9
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	121	0.8	0.7
13	Pregnancy, childbirth and the puerperium (O00-O99)	109	0.7	0.7
14	Chronic lower respiratory diseases (J40-J47)	95	0.6	0.6
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	91	0.6	0.5
...	All other causes (Residual)	2,437	16.2	14.7
Hispanic, both sexes, 45-64 years				
...	All causes	39,720	100.0	401.2
1	Malignant neoplasms (C00-C97)	11,380	28.7	114.9
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	7,145	18.0	72.2
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,122	7.9	31.5
4	Chronic liver disease and cirrhosis (K70,K73-K74)	2,869	7.2	29.0
5	Diabetes mellitus (E10-E14)	2,159	5.4	21.8
6	Cerebrovascular diseases (I60-I69)	1,708	4.3	17.3
7	Viral hepatitis (B15-B19)	865	2.2	8.7
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	720	1.8	7.3
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	678	1.7	6.8
10	Chronic lower respiratory diseases (J40-J47)	629	1.6	6.4
11	Influenza and pneumonia (J09-J18)	602	1.5	6.1
12	Septicemia (A40-A41)	600	1.5	6.1
13	Human immunodeficiency virus (HIV) disease (B20-B24)	521	1.3	5.3
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	358	0.9	3.6
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	355	0.9	3.6
...	All other causes (Residual)	6,009	15.1	60.7

... Category not applicable

\* Figure does not meet standards of reliability or precision

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality, 2013

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, both sexes, 65 years and over				
...	All causes	97,140	100.0	2,916.3
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	24,499	25.2	735.5
2	Malignant neoplasms (C00-C97)	20,774	21.4	623.7
3	Cerebrovascular diseases (I60-I69)	5,928	6.1	178.0
4	Diabetes mellitus (E10-E14)	5,096	5.2	153.0
5	Alzheimer's disease (G30)	4,061	4.2	121.9
6	Chronic lower respiratory diseases (J40-J47)	4,053	4.2	121.7
7	Influenza and pneumonia (J09-J18)	2,694	2.8	80.9
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2,246	2.3	67.4
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,106	2.2	63.2
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,672	1.7	50.2
11	Chronic liver disease and cirrhosis (K70,K73-K74)	1,652	1.7	49.6
12	Septicemia (A40-A41)	1,616	1.7	48.5
13	Parkinson's disease (G20-G21)	1,251	1.3	37.6
14	Pneumonitis due to solids and liquids (J69)	677	0.7	20.3
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	564	0.6	16.9
...	All other causes (Residual)	18,251	18.8	547.9
Hispanic, male, all ages				
...	All causes	88,880	100.0	323.7
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	18,377	20.7	66.9
2	Malignant neoplasms (C00-C97)	18,371	20.7	66.9
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	8,760	9.9	31.9
4	Diabetes mellitus (E10-E14)	3,934	4.4	14.3
5	Cerebrovascular diseases (I60-I69)	3,841	4.3	14.0
6	Chronic liver disease and cirrhosis (K70,K73-K74)	3,552	4.0	12.9
7	Chronic lower respiratory diseases (J40-J47)	2,536	2.9	9.2
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,279	2.6	8.3
9	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	2,132	2.4	7.8
10	Influenza and pneumonia (J09-J18)	1,816	2.0	6.6
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,583	1.8	5.8
12	Certain conditions originating in the perinatal period (P00-P96)	1,320	1.5	4.8
13	Alzheimer's disease (G30)	1,298	1.5	4.7
14	Septicemia (A40-A41)	1,226	1.4	4.5
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	972	1.1	3.5
...	All other causes (Residual)	16,883	19.0	61.5

... Category not applicable

\* Figure does not meet standards of reliability or precision

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality, 2013

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, male, 1-4 years				
...	All causes	483	100.0	23.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	182	37.7	8.7
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	63	13.0	3.0
3	Malignant neoplasms (C00-C97)	48	9.9	2.3
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	31	6.4	1.5
5	Influenza and pneumonia (J09-J18)	9	1.9	*
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	8	1.7	*
7	Chronic lower respiratory diseases (J40-J47)	6	1.2	*
8	Septicemia (A40-A41)	4	0.8	*
8	Certain conditions originating in the perinatal period (P00-P96)	4	0.8	*
10	Meningitis (G00,G03)	3	0.6	*
11	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2	0.4	*
11	Anemias (D50-D64)	2	0.4	*
11	Diseases of appendix (K35-K38)	2	0.4	*
11	Complications of medical and surgical care (Y40-Y84,Y88)	2	0.4	*
15	Tuberculosis (A16-A19)	1	0.2	*
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1	0.2	*
15	Cerebrovascular diseases (I60-I69)	1	0.2	*
15	Acute bronchitis and bronchiolitis (J20-J21)	1	0.2	*
...	All other causes (Residual)	113	23.4	5.4
Hispanic, male, 5-14 years				
...	All causes	591	100.0	11.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	173	29.3	3.4
2	Malignant neoplasms (C00-C97)	106	17.9	2.1
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	38	6.4	0.8
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	36	6.1	0.7
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	33	5.6	0.7
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	18	3.0	*
7	Chronic lower respiratory diseases (J40-J47)	15	2.5	*
8	Cerebrovascular diseases (I60-I69)	10	1.7	*
9	Influenza and pneumonia (J09-J18)	7	1.2	*
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	6	1.0	*
11	Septicemia (A40-A41)	5	0.8	*
11	Complications of medical and surgical care (Y40-Y84,Y88)	5	0.8	*
13	Anemias (D50-D64)	2	0.3	*
13	Meningitis (G00,G03)	2	0.3	*
13	Pneumonitis due to solids and liquids (J69)	2	0.3	*
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2	0.3	*
...	All other causes (Residual)	131	22.2	2.6

... Category not applicable

\* Figure does not meet standards of reliability or precision

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality, 2013

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, male, 15-24 years				
...	All causes	3,516	100.0	72.9
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,513	43.0	31.4
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	740	21.0	15.3
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	487	13.9	10.1
4	Malignant neoplasms (C00-C97)	209	5.9	4.3
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	86	2.4	1.8
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	32	0.9	0.7
7	Legal intervention (Y35,Y89.0)	17	0.5	*
8	Human immunodeficiency virus (HIV) disease (B20-B24)	16	0.5	*
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	15	0.4	*
9	Cerebrovascular diseases (I60-I69)	15	0.4	*
11	Influenza and pneumonia (J09-J18)	14	0.4	*
12	Septicemia (A40-A41)	13	0.4	*
13	Chronic lower respiratory diseases (J40-J47)	10	0.3	*
14	Diabetes mellitus (E10-E14)	9	0.3	*
15	Chronic liver disease and cirrhosis (K70,K73-K74)	8	0.2	*
...	All other causes (Residual)	332	9.4	6.9
Hispanic, male, 25-44 years				
...	All causes	10,360	100.0	120.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,250	31.4	37.6
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,019	9.8	11.8
3	Malignant neoplasms (C00-C97)	1,018	9.8	11.8
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	993	9.6	11.5
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	951	9.2	11.0
6	Chronic liver disease and cirrhosis (K70,K73-K74)	468	4.5	5.4
7	Cerebrovascular diseases (I60-I69)	252	2.4	2.9
8	Human immunodeficiency virus (HIV) disease (B20-B24)	241	2.3	2.8
9	Diabetes mellitus (E10-E14)	223	2.2	2.6
10	Influenza and pneumonia (J09-J18)	120	1.2	1.4
11	Septicemia (A40-A41)	84	0.8	1.0
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	74	0.7	0.9
13	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	59	0.6	0.7
14	Legal intervention (Y35,Y89.0)	57	0.6	0.7
15	Viral hepatitis (B15-B19)	51	0.5	0.6
...	All other causes (Residual)	1,500	14.5	17.4

... Category not applicable

\* Figure does not meet standards of reliability or precision

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality, 2013

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, male, 45-64 years				
...	All causes	25,357	100.0	516.4
1	Malignant neoplasms (C00-C97)	6,040	23.8	123.0
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	5,226	20.6	106.4
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,374	9.4	48.3
4	Chronic liver disease and cirrhosis (K70,K73-K74)	2,180	8.6	44.4
5	Diabetes mellitus (E10-E14)	1,303	5.1	26.5
6	Cerebrovascular diseases (I60-I69)	1,046	4.1	21.3
7	Viral hepatitis (B15-B19)	630	2.5	12.8
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	559	2.2	11.4
9	Human immunodeficiency virus (HIV) disease (B20-B24)	417	1.6	8.5
10	Influenza and pneumonia (J09-J18)	384	1.5	7.8
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	381	1.5	7.8
12	Chronic lower respiratory diseases (J40-J47)	363	1.4	7.4
13	Septicemia (A40-A41)	331	1.3	6.7
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	299	1.2	6.1
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	219	0.9	4.5
...	All other causes (Residual)	3,605	14.2	73.4
Hispanic, male, 65 years and over				
...	All causes	45,969	100.0	3,211.2
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	11,986	26.1	837.3
2	Malignant neoplasms (C00-C97)	10,946	23.8	764.6
3	Cerebrovascular diseases (I60-I69)	2,510	5.5	175.3
4	Diabetes mellitus (E10-E14)	2,397	5.2	167.4
5	Chronic lower respiratory diseases (J40-J47)	2,090	4.5	146.0
6	Alzheimer's disease (G30)	1,268	2.8	88.6
7	Influenza and pneumonia (J09-J18)	1,259	2.7	87.9
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,175	2.6	82.1
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,105	2.4	77.2
10	Chronic liver disease and cirrhosis (K70,K73-K74)	896	1.9	62.6
11	Septicemia (A40-A41)	767	1.7	53.6
12	Parkinson's disease (G20-G21)	721	1.6	50.4
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	705	1.5	49.2
14	Pneumonitis due to solids and liquids (J69)	357	0.8	24.9
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	278	0.6	19.4
...	All other causes (Residual)	7,509	16.3	524.5

... Category not applicable

\* Figure does not meet standards of reliability or precision

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality, 2013

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, female, all ages				
...	All causes	74,361	100.0	279.4
1	Malignant neoplasms (C00-C97)	16,776	22.6	63.0
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	14,866	20.0	55.9
3	Cerebrovascular diseases (I60-I69)	4,286	5.8	16.1
4	Diabetes mellitus (E10-E14)	3,698	5.0	13.9
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,255	4.4	12.2
6	Alzheimer's disease (G30)	2,829	3.8	10.6
7	Chronic lower respiratory diseases (J40-J47)	2,291	3.1	8.6
8	Influenza and pneumonia (J09-J18)	1,776	2.4	6.7
9	Chronic liver disease and cirrhosis (K70,K73-K74)	1,589	2.1	6.0
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,500	2.0	5.6
11	Septicemia (A40-A41)	1,209	1.6	4.5
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,120	1.5	4.2
13	Certain conditions originating in the perinatal period (P00-P96)	1,108	1.5	4.2
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	813	1.1	3.1
15	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	586	0.8	2.2
...	All other causes (Residual)	16,659	22.4	62.6
Hispanic, female, 1-4 years				
...	All causes	371	100.0	18.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	96	25.9	4.8
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	59	15.9	2.9
3	Malignant neoplasms (C00-C97)	42	11.3	2.1
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	22	5.9	1.1
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	15	4.0	*
6	Influenza and pneumonia (J09-J18)	10	2.7	*
7	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	7	1.9	*
8	Septicemia (A40-A41)	5	1.3	*
8	Certain conditions originating in the perinatal period (P00-P96)	5	1.3	*
10	Anemias (D50-D64)	3	0.8	*
10	Cerebrovascular diseases (I60-I69)	3	0.8	*
10	Chronic lower respiratory diseases (J40-J47)	3	0.8	*
13	Meningitis (G00,G03)	2	0.5	*
14	Shigellosis and amebiasis (A03,A06)	1	0.3	*
14	Tuberculosis (A16-A19)	1	0.3	*
14	Meningococcal infection (A39)	1	0.3	*
14	Acute bronchitis and bronchiolitis (J20-J21)	1	0.3	*
14	Diseases of appendix (K35-K38)	1	0.3	*
14	Complications of medical and surgical care (Y40-Y84,Y88)	1	0.3	*
...	All other causes (Residual)	93	25.1	4.6

... Category not applicable

\* Figure does not meet standards of reliability or precision

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality, 2013



LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, female, 5-14 years				
...	All causes	482	100.0	9.9
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	106	22.0	2.2
2	Malignant neoplasms (C00-C97)	103	21.4	2.1
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	41	8.5	0.8
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	35	7.3	0.7
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	26	5.4	0.5
6	Cerebrovascular diseases (I60-I69)	16	3.3	*
7	Influenza and pneumonia (J09-J18)	15	3.1	*
8	Diseases of heart (I00-I09,I11,I13,I20-I51)	13	2.7	*
9	Chronic lower respiratory diseases (J40-J47)	7	1.5	*
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	6	1.2	*
11	Septicemia (A40-A41)	5	1.0	*
12	Certain conditions originating in the perinatal period (P00-P96)	4	0.8	*
13	Diabetes mellitus (E10-E14)	3	0.6	*
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3	0.6	*
15	Complications of medical and surgical care (Y40-Y84,Y88)	2	0.4	*
...	All other causes (Residual)	97	20.1	2.0
Hispanic, female, 15-24 years				
...	All causes	1,141	100.0	25.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	431	37.8	9.7
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	148	13.0	3.3
3	Malignant neoplasms (C00-C97)	130	11.4	2.9
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	109	9.6	2.5
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	34	3.0	0.8
6	Pregnancy, childbirth and the puerperium (O00-O99)	27	2.4	0.6
7	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	23	2.0	0.5
8	Cerebrovascular diseases (I60-I69)	17	1.5	*
9	Influenza and pneumonia (J09-J18)	13	1.1	*
10	Septicemia (A40-A41)	10	0.9	*
11	Diabetes mellitus (E10-E14)	9	0.8	*
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	6	0.5	*
13	Anemias (D50-D64)	5	0.4	*
14	Chronic lower respiratory diseases (J40-J47)	4	0.4	*
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	2	0.2	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2	0.2	*
15	Complications of medical and surgical care (Y40-Y84,Y88)	2	0.2	*
...	All other causes (Residual)	169	14.8	3.8

... Category not applicable

\* Figure does not meet standards of reliability or precision

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality, 2013

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, female, 25-44 years				
...	All causes	4,679	100.0	59.1
1	Malignant neoplasms (C00-C97)	1,325	28.3	16.7
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	867	18.5	10.9
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	348	7.4	4.4
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	219	4.7	2.8
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	187	4.0	2.4
6	Cerebrovascular diseases (I60-I69)	162	3.5	2.0
7	Chronic liver disease and cirrhosis (K70,K73-K74)	143	3.1	1.8
8	Diabetes mellitus (E10-E14)	131	2.8	1.7
9	Pregnancy, childbirth and the puerperium (O00-O99)	109	2.3	1.4
10	Influenza and pneumonia (J09-J18)	73	1.6	0.9
11	Septicemia (A40-A41)	58	1.2	0.7
12	Human immunodeficiency virus (HIV) disease (B20-B24)	54	1.2	0.7
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	47	1.0	0.6
14	Chronic lower respiratory diseases (J40-J47)	45	1.0	0.6
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	32	0.7	0.4
...	All other causes (Residual)	879	18.8	11.1
Hispanic, female, 45-64 years				
...	All causes	14,363	100.0	287.8
1	Malignant neoplasms (C00-C97)	5,340	37.2	107.0
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,919	13.4	38.5
3	Diabetes mellitus (E10-E14)	856	6.0	17.2
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	748	5.2	15.0
5	Chronic liver disease and cirrhosis (K70,K73-K74)	689	4.8	13.8
6	Cerebrovascular diseases (I60-I69)	662	4.6	13.3
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	297	2.1	6.0
8	Septicemia (A40-A41)	269	1.9	5.4
9	Chronic lower respiratory diseases (J40-J47)	266	1.9	5.3
10	Viral hepatitis (B15-B19)	235	1.6	4.7
11	Influenza and pneumonia (J09-J18)	218	1.5	4.4
12	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	161	1.1	3.2
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	136	0.9	2.7
14	Human immunodeficiency virus (HIV) disease (B20-B24)	104	0.7	2.1
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	64	0.4	1.3
...	All other causes (Residual)	2,399	16.7	48.1

... Category not applicable

\* Figure does not meet standards of reliability or precision

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality, 2013

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, female, 65 years and over				
...	All causes	51,171	100.0	2,694.1
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	12,513	24.5	658.8
2	Malignant neoplasms (C00-C97)	9,828	19.2	517.4
3	Cerebrovascular diseases (I60-I69)	3,418	6.7	180.0
4	Alzheimer's disease (G30)	2,793	5.5	147.0
5	Diabetes mellitus (E10-E14)	2,699	5.3	142.1
6	Chronic lower respiratory diseases (J40-J47)	1,963	3.8	103.3
7	Influenza and pneumonia (J09-J18)	1,435	2.8	75.6
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,141	2.2	60.1
9	Essential hypertension and hypertensive renal disease (I10,I12,I15)	967	1.9	50.9
10	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	931	1.8	49.0
11	Septicemia (A40-A41)	849	1.7	44.7
12	Chronic liver disease and cirrhosis (K70,K73-K74)	756	1.5	39.8
13	Parkinson's disease (G20-G21)	530	1.0	27.9
14	Pneumonitis due to solids and liquids (J69)	320	0.6	16.8
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	286	0.6	15.1
...	All other causes (Residual)	10,742	21.0	565.6
Non-Hispanic, white, both sexes, all ages				
...	All causes	2,052,660	100.0	1,021.6
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	488,817	23.8	243.3
2	Malignant neoplasms (C00-C97)	462,850	22.5	230.4
3	Chronic lower respiratory diseases (J40-J47)	131,667	6.4	65.5
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	100,773	4.9	50.2
5	Cerebrovascular diseases (I60-I69)	99,759	4.9	49.7
6	Alzheimer's disease (G30)	73,234	3.6	36.4
7	Diabetes mellitus (E10-E14)	51,297	2.5	25.5
8	Influenza and pneumonia (J09-J18)	45,350	2.2	22.6
9	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	34,253	1.7	17.0
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	34,173	1.7	17.0
11	Septicemia (A40-A41)	28,431	1.4	14.2
12	Chronic liver disease and cirrhosis (K70,K73-K74)	26,692	1.3	13.3
13	Parkinson's disease (G20-G21)	22,241	1.1	11.1
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	22,043	1.1	11.0
15	Pneumonitis due to solids and liquids (J69)	15,630	0.8	7.8
...	All other causes (Residual)	415,450	20.2	206.8

... Category not applicable

\* Figure does not meet standards of reliability or precision

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality, 2013

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, both sexes, 1-4 years				
...	All causes	1,977	100.0	23.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	675	34.1	8.1
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	227	11.5	2.7
3	Malignant neoplasms (C00-C97)	160	8.1	1.9
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	128	6.5	1.5
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	82	4.1	1.0
6	Influenza and pneumonia (J09-J18)	53	2.7	0.6
7	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	30	1.5	0.4
8	Chronic lower respiratory diseases (J40-J47)	24	1.2	0.3
9	Septicemia (A40-A41)	22	1.1	0.3
9	Certain conditions originating in the perinatal period (P00-P96)	22	1.1	0.3
11	Cerebrovascular diseases (I60-I69)	15	0.8	*
12	Meningitis (G00,G03)	9	0.5	*
12	Acute bronchitis and bronchiolitis (J20-J21)	9	0.5	*
14	Complications of medical and surgical care (Y40-Y84,Y88)	7	0.4	*
15	Anemias (D50-D64)	6	0.3	*
...	All other causes (Residual)	508	25.7	6.1
Non-Hispanic, white, both sexes, 5-14 years				
...	All causes	2,824	100.0	12.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	838	29.7	3.7
2	Malignant neoplasms (C00-C97)	512	18.1	2.3
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	244	8.6	1.1
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	167	5.9	0.7
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	121	4.3	0.5
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	76	2.7	0.3
7	Influenza and pneumonia (J09-J18)	71	2.5	0.3
8	Cerebrovascular diseases (I60-I69)	42	1.5	0.2
9	Chronic lower respiratory diseases (J40-J47)	38	1.3	0.2
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	35	1.2	0.2
11	Septicemia (A40-A41)	28	1.0	0.1
12	Certain conditions originating in the perinatal period (P00-P96)	19	0.7	*
13	Diabetes mellitus (E10-E14)	18	0.6	*
14	Complications of medical and surgical care (Y40-Y84,Y88)	10	0.4	*
15	Meningitis (G00,G03)	7	0.2	*
15	Pneumonitis due to solids and liquids (J69)	7	0.2	*
...	All other causes (Residual)	591	20.9	2.6

... Category not applicable

\* Figure does not meet standards of reliability or precision

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality, 2013

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, both sexes, 15-24 years				
...	All causes	16,092	100.0	64.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	7,820	48.6	31.3
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	3,388	21.1	13.6
3	Malignant neoplasms (C00-C97)	804	5.0	3.2
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	687	4.3	2.7
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	485	3.0	1.9
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	227	1.4	0.9
7	Influenza and pneumonia (J09-J18)	132	0.8	0.5
8	Diabetes mellitus (E10-E14)	95	0.6	0.4
9	Pregnancy, childbirth and the puerperium (O00-O99)	76	0.5	0.3
10	Cerebrovascular diseases (I60-I69)	75	0.5	0.3
11	Septicemia (A40-A41)	70	0.4	0.3
12	Chronic lower respiratory diseases (J40-J47)	54	0.3	0.2
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	43	0.3	0.2
14	Legal intervention (Y35,Y89.0)	34	0.2	0.1
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	32	0.2	0.1
...	All other causes (Residual)	2,070	12.9	8.3
Non-Hispanic, white, both sexes, 25-44 years				
...	All causes	71,514	100.0	145.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	22,714	31.8	46.1
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	10,023	14.0	20.4
3	Malignant neoplasms (C00-C97)	9,172	12.8	18.6
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	7,914	11.1	16.1
5	Chronic liver disease and cirrhosis (K70,K73-K74)	2,016	2.8	4.1
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,756	2.5	3.6
7	Diabetes mellitus (E10-E14)	1,404	2.0	2.9
8	Cerebrovascular diseases (I60-I69)	995	1.4	2.0
9	Influenza and pneumonia (J09-J18)	824	1.2	1.7
10	Septicemia (A40-A41)	645	0.9	1.3
11	Chronic lower respiratory diseases (J40-J47)	620	0.9	1.3
12	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	601	0.8	1.2
13	Human immunodeficiency virus (HIV) disease (B20-B24)	461	0.6	0.9
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	386	0.5	0.8
15	Pregnancy, childbirth and the puerperium (O00-O99)	349	0.5	0.7
...	All other causes (Residual)	11,634	16.3	23.6

... Category not applicable

\* Figure does not meet standards of reliability or precision

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality, 2013

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, both sexes, 45-64 years				
...	All causes	368,049	100.0	630.3
1	Malignant neoplasms (C00-C97)	116,845	31.7	200.1
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	75,063	20.4	128.6
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	28,278	7.7	48.4
4	Chronic lower respiratory diseases (J40-J47)	17,096	4.6	29.3
5	Chronic liver disease and cirrhosis (K70,K73-K74)	15,032	4.1	25.7
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	13,938	3.8	23.9
7	Diabetes mellitus (E10-E14)	11,662	3.2	20.0
8	Cerebrovascular diseases (I60-I69)	9,908	2.7	17.0
9	Septicemia (A40-A41)	5,142	1.4	8.8
10	Influenza and pneumonia (J09-J18)	4,814	1.3	8.2
11	Viral hepatitis (B15-B19)	3,884	1.1	6.7
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3,825	1.0	6.6
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	2,676	0.7	4.6
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1,584	0.4	2.7
15	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,561	0.4	2.7
...	All other causes (Residual)	56,741	15.4	97.2
Non-Hispanic, white, both sexes, 65 years and over				
...	All causes	1,581,591	100.0	4,465.5
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	405,042	25.6	1,143.6
2	Malignant neoplasms (C00-C97)	335,309	21.2	946.7
3	Chronic lower respiratory diseases (J40-J47)	113,828	7.2	321.4
4	Cerebrovascular diseases (I60-I69)	88,674	5.6	250.4
5	Alzheimer's disease (G30)	72,426	4.6	204.5
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	39,898	2.5	112.6
7	Influenza and pneumonia (J09-J18)	39,390	2.5	111.2
8	Diabetes mellitus (E10-E14)	38,110	2.4	107.6
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	29,884	1.9	84.4
10	Septicemia (A40-A41)	22,466	1.4	63.4
11	Parkinson's disease (G20-G21)	21,706	1.4	61.3
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	19,142	1.2	54.0
13	Pneumonitis due to solids and liquids (J69)	14,041	0.9	39.6
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	10,754	0.7	30.4
15	Chronic liver disease and cirrhosis (K70,K73-K74)	9,621	0.6	27.2
...	All other causes (Residual)	321,300	20.3	907.2

... Category not applicable

\* Figure does not meet standards of reliability or precision

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality, 2013

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, male, all ages				
...	All causes	1,021,135	100.0	1,032.1
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	256,294	25.1	259.0
2	Malignant neoplasms (C00-C97)	244,646	24.0	247.3
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	61,391	6.0	62.1
4	Chronic lower respiratory diseases (J40-J47)	61,106	6.0	61.8
5	Cerebrovascular diseases (I60-I69)	40,350	4.0	40.8
6	Diabetes mellitus (E10-E14)	27,807	2.7	28.1
7	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	26,621	2.6	26.9
8	Alzheimer's disease (G30)	22,345	2.2	22.6
9	Influenza and pneumonia (J09-J18)	21,059	2.1	21.3
10	Chronic liver disease and cirrhosis (K70,K73-K74)	17,290	1.7	17.5
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	17,213	1.7	17.4
12	Parkinson's disease (G20-G21)	13,370	1.3	13.5
13	Septicemia (A40-A41)	13,364	1.3	13.5
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	8,955	0.9	9.1
15	Pneumonitis due to solids and liquids (J69)	8,534	0.8	8.6
...	All other causes (Residual)	180,790	17.7	182.7
Non-Hispanic, white, male, 1-4 years				
...	All causes	1,149	100.0	26.9
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	437	38.0	10.2
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	101	8.8	2.4
3	Malignant neoplasms (C00-C97)	96	8.4	2.2
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	67	5.8	1.6
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	48	4.2	1.1
6	Influenza and pneumonia (J09-J18)	31	2.7	0.7
7	Certain conditions originating in the perinatal period (P00-P96)	18	1.6	*
8	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	16	1.4	*
9	Septicemia (A40-A41)	13	1.1	*
9	Chronic lower respiratory diseases (J40-J47)	13	1.1	*
11	Cerebrovascular diseases (I60-I69)	8	0.7	*
12	Acute bronchitis and bronchiolitis (J20-J21)	5	0.4	*
13	Anemias (D50-D64)	4	0.3	*
13	Meningitis (G00,G03)	4	0.3	*
15	Complications of medical and surgical care (Y40-Y84,Y88)	3	0.3	*
...	All other causes (Residual)	285	24.8	6.7

... Category not applicable

\* Figure does not meet standards of reliability or precision

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality, 2013

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, male, 5-14 years				
...	All causes	1,657	100.0	14.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	519	31.3	4.5
2	Malignant neoplasms (C00-C97)	266	16.1	2.3
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	169	10.2	1.5
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	95	5.7	0.8
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	72	4.3	0.6
6	Influenza and pneumonia (J09-J18)	41	2.5	0.4
7	Diseases of heart (I00-I09,I11,I13,I20-I51)	37	2.2	0.3
8	Chronic lower respiratory diseases (J40-J47)	24	1.4	0.2
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	22	1.3	0.2
10	Cerebrovascular diseases (I60-I69)	21	1.3	0.2
11	Septicemia (A40-A41)	16	1.0	*
12	Certain conditions originating in the perinatal period (P00-P96)	11	0.7	*
13	Diabetes mellitus (E10-E14)	10	0.6	*
14	Meningitis (G00,G03)	6	0.4	*
14	Aortic aneurysm and dissection (I71)	6	0.4	*
...	All other causes (Residual)	342	20.6	3.0
Non-Hispanic, white, male, 15-24 years				
...	All causes	11,461	100.0	89.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	5,695	49.7	44.5
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,732	23.8	21.4
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	496	4.3	3.9
4	Malignant neoplasms (C00-C97)	456	4.0	3.6
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	315	2.7	2.5
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	142	1.2	1.1
7	Influenza and pneumonia (J09-J18)	79	0.7	0.6
8	Diabetes mellitus (E10-E14)	58	0.5	0.5
9	Cerebrovascular diseases (I60-I69)	46	0.4	0.4
10	Septicemia (A40-A41)	37	0.3	0.3
11	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	33	0.3	0.3
12	Chronic lower respiratory diseases (J40-J47)	32	0.3	0.3
13	Legal intervention (Y35,Y89.0)	30	0.3	0.2
14	Aortic aneurysm and dissection (I71)	16	0.1	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	16	0.1	*
...	All other causes (Residual)	1,278	11.2	10.0

... Category not applicable

\* Figure does not meet standards of reliability or precision

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality, 2013



LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, male, 25-44 years				
...	All causes	46,101	100.0	185.9
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	15,952	34.6	64.3
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	7,742	16.8	31.2
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	5,495	11.9	22.2
4	Malignant neoplasms (C00-C97)	4,198	9.1	16.9
5	Chronic liver disease and cirrhosis (K70,K73-K74)	1,282	2.8	5.2
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,213	2.6	4.9
7	Diabetes mellitus (E10-E14)	879	1.9	3.5
8	Cerebrovascular diseases (I60-I69)	554	1.2	2.2
9	Influenza and pneumonia (J09-J18)	431	0.9	1.7
10	Human immunodeficiency virus (HIV) disease (B20-B24)	365	0.8	1.5
11	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	335	0.7	1.4
12	Septicemia (A40-A41)	320	0.7	1.3
13	Chronic lower respiratory diseases (J40-J47)	282	0.6	1.1
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	202	0.4	0.8
15	Aortic aneurysm and dissection (I71)	171	0.4	0.7
...	All other causes (Residual)	6,680	14.5	26.9
Non-Hispanic, white, male, 45-64 years				
...	All causes	225,780	100.0	785.0
1	Malignant neoplasms (C00-C97)	63,365	28.1	220.3
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	53,888	23.9	187.4
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	18,711	8.3	65.1
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	10,384	4.6	36.1
5	Chronic liver disease and cirrhosis (K70,K73-K74)	10,322	4.6	35.9
6	Chronic lower respiratory diseases (J40-J47)	8,635	3.8	30.0
7	Diabetes mellitus (E10-E14)	7,361	3.3	25.6
8	Cerebrovascular diseases (I60-I69)	5,704	2.5	19.8
9	Septicemia (A40-A41)	2,781	1.2	9.7
10	Viral hepatitis (B15-B19)	2,769	1.2	9.6
11	Influenza and pneumonia (J09-J18)	2,721	1.2	9.5
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2,196	1.0	7.6
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,793	0.8	6.2
14	Human immunodeficiency virus (HIV) disease (B20-B24)	1,185	0.5	4.1
15	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,027	0.5	3.6
...	All other causes (Residual)	32,938	14.6	114.5

... Category not applicable

\* Figure does not meet standards of reliability or precision

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality, 2013

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, male, 65 years and over				
...	All causes	728,907	100.0	4,637.1
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	196,413	26.9	1,249.5
2	Malignant neoplasms (C00-C97)	176,236	24.2	1,121.2
3	Chronic lower respiratory diseases (J40-J47)	52,115	7.1	331.5
4	Cerebrovascular diseases (I60-I69)	33,989	4.7	216.2
5	Alzheimer's disease (G30)	22,020	3.0	140.1
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	19,735	2.7	125.5
7	Diabetes mellitus (E10-E14)	19,493	2.7	124.0
8	Influenza and pneumonia (J09-J18)	17,714	2.4	112.7
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	14,774	2.0	94.0
10	Parkinson's disease (G20-G21)	13,010	1.8	82.8
11	Septicemia (A40-A41)	10,165	1.4	64.7
12	Pneumonitis due to solids and liquids (J69)	7,595	1.0	48.3
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	7,011	1.0	44.6
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	5,677	0.8	36.1
15	Chronic liver disease and cirrhosis (K70,K73-K74)	5,673	0.8	36.1
...	All other causes (Residual)	127,287	17.5	809.8
Non-Hispanic, white, female, all ages				
...	All causes	1,031,525	100.0	1,011.5
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	232,523	22.5	228.0
2	Malignant neoplasms (C00-C97)	218,204	21.2	214.0
3	Chronic lower respiratory diseases (J40-J47)	70,561	6.8	69.2
4	Cerebrovascular diseases (I60-I69)	59,409	5.8	58.3
5	Alzheimer's disease (G30)	50,889	4.9	49.9
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	39,382	3.8	38.6
7	Influenza and pneumonia (J09-J18)	24,291	2.4	23.8
8	Diabetes mellitus (E10-E14)	23,490	2.3	23.0
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	16,960	1.6	16.6
10	Septicemia (A40-A41)	15,067	1.5	14.8
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	13,088	1.3	12.8
12	Chronic liver disease and cirrhosis (K70,K73-K74)	9,402	0.9	9.2
13	Parkinson's disease (G20-G21)	8,871	0.9	8.7
14	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	7,632	0.7	7.5
15	Pneumonitis due to solids and liquids (J69)	7,096	0.7	7.0
...	All other causes (Residual)	234,660	22.7	230.1

... Category not applicable

\* Figure does not meet standards of reliability or precision

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality, 2013

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, female, 1-4 years				
...	All causes	828	100.0	20.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	238	28.7	5.9
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	126	15.2	3.1
3	Malignant neoplasms (C00-C97)	64	7.7	1.6
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	61	7.4	1.5
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	34	4.1	0.8
6	Influenza and pneumonia (J09-J18)	22	2.7	0.5
7	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	14	1.7	*
8	Chronic lower respiratory diseases (J40-J47)	11	1.3	*
9	Septicemia (A40-A41)	9	1.1	*
10	Cerebrovascular diseases (I60-I69)	7	0.8	*
11	Meningitis (G00,G03)	5	0.6	*
12	Acute bronchitis and bronchiolitis (J20-J21)	4	0.5	*
12	Certain conditions originating in the perinatal period (P00-P96)	4	0.5	*
12	Complications of medical and surgical care (Y40-Y84,Y88)	4	0.5	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3	0.4	*
...	All other causes (Residual)	222	26.8	5.5
Non-Hispanic, white, female, 5-14 years				
...	All causes	1,167	100.0	10.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	319	27.3	2.9
2	Malignant neoplasms (C00-C97)	246	21.1	2.2
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	75	6.4	0.7
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	72	6.2	0.7
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	49	4.2	0.4
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	39	3.3	0.4
7	Influenza and pneumonia (J09-J18)	30	2.6	0.3
8	Cerebrovascular diseases (I60-I69)	21	1.8	0.2
9	Chronic lower respiratory diseases (J40-J47)	14	1.2	*
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	13	1.1	*
11	Septicemia (A40-A41)	12	1.0	*
12	Diabetes mellitus (E10-E14)	8	0.7	*
12	Certain conditions originating in the perinatal period (P00-P96)	8	0.7	*
14	Complications of medical and surgical care (Y40-Y84,Y88)	5	0.4	*
15	Pneumonitis due to solids and liquids (J69)	3	0.3	*
...	All other causes (Residual)	253	21.7	2.3

... Category not applicable

\* Figure does not meet standards of reliability or precision

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality, 2013

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, female, 15-24 years				
...	All causes	4,631	100.0	38.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,125	45.9	17.4
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	656	14.2	5.4
3	Malignant neoplasms (C00-C97)	348	7.5	2.9
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	191	4.1	1.6
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	170	3.7	1.4
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	85	1.8	0.7
7	Pregnancy, childbirth and the puerperium (O00-O99)	76	1.6	0.6
8	Influenza and pneumonia (J09-J18)	53	1.1	0.4
9	Diabetes mellitus (E10-E14)	37	0.8	0.3
10	Septicemia (A40-A41)	33	0.7	0.3
11	Cerebrovascular diseases (I60-I69)	29	0.6	0.2
12	Chronic lower respiratory diseases (J40-J47)	22	0.5	0.2
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	16	0.3	*
14	Complications of medical and surgical care (Y40-Y84,Y88)	15	0.3	*
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	10	0.2	*
...	All other causes (Residual)	765	16.5	6.3
Non-Hispanic, white, female, 25-44 years				
...	All causes	25,413	100.0	104.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	6,762	26.6	27.7
2	Malignant neoplasms (C00-C97)	4,974	19.6	20.4
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,419	9.5	9.9
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,281	9.0	9.3
5	Chronic liver disease and cirrhosis (K70,K73-K74)	734	2.9	3.0
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	543	2.1	2.2
7	Diabetes mellitus (E10-E14)	525	2.1	2.1
8	Cerebrovascular diseases (I60-I69)	441	1.7	1.8
9	Influenza and pneumonia (J09-J18)	393	1.5	1.6
10	Pregnancy, childbirth and the puerperium (O00-O99)	349	1.4	1.4
11	Chronic lower respiratory diseases (J40-J47)	338	1.3	1.4
12	Septicemia (A40-A41)	325	1.3	1.3
13	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	266	1.0	1.1
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	184	0.7	0.8
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	116	0.5	0.5
...	All other causes (Residual)	4,763	18.7	19.5

... Category not applicable

\* Figure does not meet standards of reliability or precision

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality, 2013

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, female, 45-64 years				
...	All causes	142,269	100.0	480.2
1	Malignant neoplasms (C00-C97)	53,480	37.6	180.5
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	21,175	14.9	71.5
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	9,567	6.7	32.3
4	Chronic lower respiratory diseases (J40-J47)	8,461	5.9	28.6
5	Chronic liver disease and cirrhosis (K70,K73-K74)	4,710	3.3	15.9
6	Diabetes mellitus (E10-E14)	4,301	3.0	14.5
7	Cerebrovascular diseases (I60-I69)	4,204	3.0	14.2
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	3,554	2.5	12.0
9	Septicemia (A40-A41)	2,361	1.7	8.0
10	Influenza and pneumonia (J09-J18)	2,093	1.5	7.1
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,629	1.1	5.5
12	Viral hepatitis (B15-B19)	1,115	0.8	3.8
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	883	0.6	3.0
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	667	0.5	2.3
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	613	0.4	2.1
...	All other causes (Residual)	23,456	16.5	79.2
Non-Hispanic, white, female, 65 years and over				
...	All causes	852,684	100.0	4,328.5
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	208,629	24.5	1,059.1
2	Malignant neoplasms (C00-C97)	159,073	18.7	807.5
3	Chronic lower respiratory diseases (J40-J47)	61,713	7.2	313.3
4	Cerebrovascular diseases (I60-I69)	54,685	6.4	277.6
5	Alzheimer's disease (G30)	50,406	5.9	255.9
6	Influenza and pneumonia (J09-J18)	21,676	2.5	110.0
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	20,163	2.4	102.4
8	Diabetes mellitus (E10-E14)	18,617	2.2	94.5
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	15,110	1.8	76.7
10	Septicemia (A40-A41)	12,301	1.4	62.4
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	12,131	1.4	61.6
12	Parkinson's disease (G20-G21)	8,696	1.0	44.1
13	Pneumonitis due to solids and liquids (J69)	6,446	0.8	32.7
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	5,077	0.6	25.8
15	Chronic liver disease and cirrhosis (K70,K73-K74)	3,948	0.5	20.0
...	All other causes (Residual)	194,013	22.8	984.9

... Category not applicable

\* Figure does not meet standards of reliability or precision

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality, 2013

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, both sexes, all ages				
...	All causes	299,227	100.0	733.4
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	71,102	23.8	174.3
2	Malignant neoplasms (C00-C97)	67,272	22.5	164.9
3	Cerebrovascular diseases (I60-I69)	16,111	5.4	39.5
4	Diabetes mellitus (E10-E14)	13,239	4.4	32.4
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	13,169	4.4	32.3
6	Chronic lower respiratory diseases (J40-J47)	9,816	3.3	24.1
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	8,340	2.8	20.4
8	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	7,950	2.7	19.5
9	Septicemia (A40-A41)	6,193	2.1	15.2
10	Alzheimer's disease (G30)	5,664	1.9	13.9
11	Influenza and pneumonia (J09-J18)	5,494	1.8	13.5
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	5,400	1.8	13.2
13	Certain conditions originating in the perinatal period (P00-P96)	3,854	1.3	9.4
14	Human immunodeficiency virus (HIV) disease (B20-B24)	3,673	1.2	9.0
15	Chronic liver disease and cirrhosis (K70,K73-K74)	2,998	1.0	7.3
...	All other causes (Residual)	58,952	19.7	144.5
Non-Hispanic, black, both sexes, 1-4 years				
...	All causes	963	100.0	39.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	285	29.6	11.7
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	133	13.8	5.5
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	88	9.1	3.6
4	Malignant neoplasms (C00-C97)	62	6.4	2.5
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	58	6.0	2.4
6	Chronic lower respiratory diseases (J40-J47)	26	2.7	1.1
7	Influenza and pneumonia (J09-J18)	21	2.2	0.9
8	Septicemia (A40-A41)	18	1.9	*
9	Cerebrovascular diseases (I60-I69)	15	1.6	*
10	Certain conditions originating in the perinatal period (P00-P96)	12	1.2	*
11	Anemias (D50-D64)	8	0.8	*
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	6	0.6	*
13	Meningitis (G00,G03)	5	0.5	*
13	Complications of medical and surgical care (Y40-Y84,Y88)	5	0.5	*
15	Acute bronchitis and bronchiolitis (J20-J21)	2	0.2	*
...	All other causes (Residual)	219	22.7	9.0

... Category not applicable

\* Figure does not meet standards of reliability or precision

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality, 2013

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, both sexes, 5-14 years				
...	All causes	1,132	100.0	18.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	324	28.6	5.2
2	Malignant neoplasms (C00-C97)	119	10.5	1.9
3	Chronic lower respiratory diseases (J40-J47)	86	7.6	1.4
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	81	7.2	1.3
5	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	80	7.1	1.3
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	55	4.9	0.9
7	Diseases of heart (I00-I09,I11,I13,I20-I51)	49	4.3	0.8
8	Influenza and pneumonia (J09-J18)	27	2.4	0.4
9	Cerebrovascular diseases (I60-I69)	18	1.6	*
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	14	1.2	*
11	Septicemia (A40-A41)	13	1.1	*
12	Anemias (D50-D64)	11	1.0	*
13	Certain conditions originating in the perinatal period (P00-P96)	6	0.5	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	5	0.4	*
14	Complications of medical and surgical care (Y40-Y84,Y88)	5	0.4	*
...	All other causes (Residual)	239	21.1	3.9
Non-Hispanic, black, both sexes, 15-24 years				
...	All causes	6,423	100.0	94.1
1	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	2,693	41.9	39.5
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,376	21.4	20.2
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	513	8.0	7.5
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	287	4.5	4.2
5	Malignant neoplasms (C00-C97)	239	3.7	3.5
6	Human immunodeficiency virus (HIV) disease (B20-B24)	85	1.3	1.2
7	Chronic lower respiratory diseases (J40-J47)	80	1.2	1.2
8	Diabetes mellitus (E10-E14)	73	1.1	1.1
9	Anemias (D50-D64)	70	1.1	1.0
10	Pregnancy, childbirth and the puerperium (O00-O99)	69	1.1	1.0
11	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	57	0.9	0.8
12	Legal intervention (Y35,Y89.0)	43	0.7	0.6
13	Cerebrovascular diseases (I60-I69)	38	0.6	0.6
14	Septicemia (A40-A41)	36	0.6	0.5
15	Influenza and pneumonia (J09-J18)	31	0.5	0.5
...	All other causes (Residual)	733	11.4	10.7

... Category not applicable

\* Figure does not meet standards of reliability or precision

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality, 2013

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, both sexes, 25-44 years				
...	All causes	22,961	100.0	209.2
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,741	16.3	34.1
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	3,670	16.0	33.4
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,511	15.3	32.0
4	Malignant neoplasms (C00-C97)	2,531	11.0	23.1
5	Human immunodeficiency virus (HIV) disease (B20-B24)	1,066	4.6	9.7
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	970	4.2	8.8
7	Diabetes mellitus (E10-E14)	752	3.3	6.9
8	Cerebrovascular diseases (I60-I69)	627	2.7	5.7
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	342	1.5	3.1
10	Septicemia (A40-A41)	314	1.4	2.9
11	Chronic lower respiratory diseases (J40-J47)	300	1.3	2.7
12	Influenza and pneumonia (J09-J18)	253	1.1	2.3
13	Pregnancy, childbirth and the puerperium (O00-O99)	235	1.0	2.1
14	Chronic liver disease and cirrhosis (K70,K73-K74)	233	1.0	2.1
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	232	1.0	2.1
...	All other causes (Residual)	4,184	18.2	38.1
Non-Hispanic, black, both sexes, 45-64 years				
...	All causes	89,240	100.0	903.7
1	Malignant neoplasms (C00-C97)	25,120	28.1	254.4
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	21,739	24.4	220.1
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	4,705	5.3	47.6
4	Diabetes mellitus (E10-E14)	4,264	4.8	43.2
5	Cerebrovascular diseases (I60-I69)	4,260	4.8	43.1
6	Chronic lower respiratory diseases (J40-J47)	2,436	2.7	24.7
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2,207	2.5	22.3
8	Human immunodeficiency virus (HIV) disease (B20-B24)	2,127	2.4	21.5
9	Chronic liver disease and cirrhosis (K70,K73-K74)	1,992	2.2	20.2
10	Septicemia (A40-A41)	1,803	2.0	18.3
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,453	1.6	14.7
12	Influenza and pneumonia (J09-J18)	1,302	1.5	13.2
13	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,111	1.2	11.3
14	Viral hepatitis (B15-B19)	848	1.0	8.6
15	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	605	0.7	6.1
...	All other causes (Residual)	13,268	14.9	134.4

... Category not applicable

\* Figure does not meet standards of reliability or precision

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality, 2013



LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, both sexes, 65 years and over				
...	All causes	171,707	100.0	4,412.8
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	45,131	26.3	1,159.9
2	Malignant neoplasms (C00-C97)	39,192	22.8	1,007.2
3	Cerebrovascular diseases (I60-I69)	11,113	6.5	285.6
4	Diabetes mellitus (E10-E14)	8,144	4.7	209.3
5	Chronic lower respiratory diseases (J40-J47)	6,877	4.0	176.7
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	5,747	3.3	147.7
7	Alzheimer's disease (G30)	5,577	3.2	143.3
8	Septicemia (A40-A41)	3,955	2.3	101.6
9	Influenza and pneumonia (J09-J18)	3,790	2.2	97.4
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	3,708	2.2	95.3
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,555	1.5	65.7
12	Pneumonitis due to solids and liquids (J69)	1,239	0.7	31.8
13	Parkinson's disease (G20-G21)	970	0.6	24.9
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	790	0.5	20.3
15	Chronic liver disease and cirrhosis (K70,K73-K74)	772	0.4	19.8
...	All other causes (Residual)	32,147	18.7	826.2
Non-Hispanic, black, male, all ages				
...	All causes	152,661	100.0	782.5
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	36,562	23.9	187.4
2	Malignant neoplasms (C00-C97)	34,319	22.5	175.9
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	8,845	5.8	45.3
4	Cerebrovascular diseases (I60-I69)	7,266	4.8	37.2
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	6,851	4.5	35.1
6	Diabetes mellitus (E10-E14)	6,298	4.1	32.3
7	Chronic lower respiratory diseases (J40-J47)	5,019	3.3	25.7
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3,944	2.6	20.2
9	Septicemia (A40-A41)	2,854	1.9	14.6
10	Influenza and pneumonia (J09-J18)	2,657	1.7	13.6
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	2,467	1.6	12.6
12	Human immunodeficiency virus (HIV) disease (B20-B24)	2,429	1.6	12.4
13	Certain conditions originating in the perinatal period (P00-P96)	2,102	1.4	10.8
14	Chronic liver disease and cirrhosis (K70,K73-K74)	1,915	1.3	9.8
15	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,850	1.2	9.5
...	All other causes (Residual)	27,283	17.9	139.8

... Category not applicable

\* Figure does not meet standards of reliability or precision

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality, 2013

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, male, 1-4 years				
...	All causes	538	100.0	43.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	165	30.7	13.3
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	68	12.6	5.5
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	50	9.3	4.0
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	36	6.7	2.9
5	Malignant neoplasms (C00-C97)	33	6.1	2.7
6	Chronic lower respiratory diseases (J40-J47)	15	2.8	*
7	Septicemia (A40-A41)	12	2.2	*
7	Cerebrovascular diseases (I60-I69)	12	2.2	*
7	Influenza and pneumonia (J09-J18)	12	2.2	*
10	Certain conditions originating in the perinatal period (P00-P96)	6	1.1	*
11	Anemias (D50-D64)	4	0.7	*
11	Meningitis (G00,G03)	4	0.7	*
13	Complications of medical and surgical care (Y40-Y84,Y88)	2	0.4	*
14	Acute bronchitis and bronchiolitis (J20-J21)	1	0.2	*
14	Pneumoconioses and chemical effects (J60-J66,J68)	1	0.2	*
14	Pneumonitis due to solids and liquids (J69)	1	0.2	*
14	Enterocolitis due to <i>Clostridium difficile</i> (A04.7)	1	0.2	*
...	All other causes (Residual)	115	21.4	9.3
Non-Hispanic, black, male, 5-14 years				
...	All causes	653	100.0	20.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	210	32.2	6.7
2	Malignant neoplasms (C00-C97)	65	10.0	2.1
3	Chronic lower respiratory diseases (J40-J47)	54	8.3	1.7
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	53	8.1	1.7
5	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	37	5.7	1.2
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	33	5.1	1.0
7	Diseases of heart (I00-I09,I11,I13,I20-I51)	29	4.4	0.9
8	Influenza and pneumonia (J09-J18)	18	2.8	*
9	Cerebrovascular diseases (I60-I69)	10	1.5	*
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	9	1.4	*
11	Septicemia (A40-A41)	8	1.2	*
12	Anemias (D50-D64)	5	0.8	*
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3	0.5	*
13	Certain conditions originating in the perinatal period (P00-P96)	3	0.5	*
13	Complications of medical and surgical care (Y40-Y84,Y88)	3	0.5	*
...	All other causes (Residual)	113	17.3	3.6

... Category not applicable

\* Figure does not meet standards of reliability or precision

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality, 2013

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, male, 15-24 years				
...	All causes	4,957	100.0	143.7
1	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	2,450	49.4	71.0
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	995	20.1	28.8
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	421	8.5	12.2
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	191	3.9	5.5
5	Malignant neoplasms (C00-C97)	128	2.6	3.7
6	Human immunodeficiency virus (HIV) disease (B20-B24)	61	1.2	1.8
7	Chronic lower respiratory diseases (J40-J47)	55	1.1	1.6
8	Diabetes mellitus (E10-E14)	48	1.0	1.4
9	Anemias (D50-D64)	44	0.9	1.3
10	Legal intervention (Y35,Y89.0)	43	0.9	1.2
11	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	35	0.7	1.0
12	Cerebrovascular diseases (I60-I69)	21	0.4	0.6
13	Septicemia (A40-A41)	19	0.4	*
14	Influenza and pneumonia (J09-J18)	16	0.3	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	9	0.2	*
...	All other causes (Residual)	421	8.5	12.2
Non-Hispanic, black, male, 25-44 years				
...	All causes	14,594	100.0	278.5
1	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	3,230	22.1	61.6
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,531	17.3	48.3
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,422	16.6	46.2
4	Malignant neoplasms (C00-C97)	1,015	7.0	19.4
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	800	5.5	15.3
6	Human immunodeficiency virus (HIV) disease (B20-B24)	654	4.5	12.5
7	Diabetes mellitus (E10-E14)	436	3.0	8.3
8	Cerebrovascular diseases (I60-I69)	337	2.3	6.4
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	188	1.3	3.6
10	Chronic lower respiratory diseases (J40-J47)	170	1.2	3.2
11	Septicemia (A40-A41)	151	1.0	2.9
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	140	1.0	2.7
13	Influenza and pneumonia (J09-J18)	133	0.9	2.5
14	Chronic liver disease and cirrhosis (K70,K73-K74)	122	0.8	2.3
15	Anemias (D50-D64)	111	0.8	2.1
...	All other causes (Residual)	2,154	14.8	41.1

... Category not applicable

\* Figure does not meet standards of reliability or precision

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality, 2013

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, male, 45-64 years				
...	All causes	51,716	100.0	1,130.9
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	13,624	26.3	297.9
2	Malignant neoplasms (C00-C97)	13,137	25.4	287.3
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,298	6.4	72.1
4	Cerebrovascular diseases (I60-I69)	2,436	4.7	53.3
5	Diabetes mellitus (E10-E14)	2,365	4.6	51.7
6	Human immunodeficiency virus (HIV) disease (B20-B24)	1,441	2.8	31.5
7	Chronic liver disease and cirrhosis (K70,K73-K74)	1,314	2.5	28.7
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,273	2.5	27.8
9	Chronic lower respiratory diseases (J40-J47)	1,212	2.3	26.5
10	Septicemia (A40-A41)	933	1.8	20.4
11	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	875	1.7	19.1
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	859	1.7	18.8
13	Influenza and pneumonia (J09-J18)	743	1.4	16.2
14	Viral hepatitis (B15-B19)	570	1.1	12.5
15	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	463	0.9	10.1
...	All other causes (Residual)	7,173	13.9	156.9
Non-Hispanic, black, male, 65 years and over				
...	All causes	76,492	100.0	4,930.5
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	20,207	26.4	1,302.5
2	Malignant neoplasms (C00-C97)	19,939	26.1	1,285.2
3	Cerebrovascular diseases (I60-I69)	4,427	5.8	285.4
4	Chronic lower respiratory diseases (J40-J47)	3,507	4.6	226.1
5	Diabetes mellitus (E10-E14)	3,446	4.5	222.1
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2,457	3.2	158.4
7	Septicemia (A40-A41)	1,701	2.2	109.6
8	Influenza and pneumonia (J09-J18)	1,695	2.2	109.3
9	Alzheimer's disease (G30)	1,595	2.1	102.8
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,464	1.9	94.4
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,432	1.9	92.3
12	Pneumonitis due to solids and liquids (J69)	642	0.8	41.4
13	Parkinson's disease (G20-G21)	573	0.7	36.9
14	Chronic liver disease and cirrhosis (K70,K73-K74)	478	0.6	30.8
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	384	0.5	24.8
...	All other causes (Residual)	12,545	16.4	808.6

... Category not applicable

\* Figure does not meet standards of reliability or precision

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality, 2013

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, female, all ages				
...	All causes	146,566	100.0	688.4
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	34,540	23.6	162.2
2	Malignant neoplasms (C00-C97)	32,953	22.5	154.8
3	Cerebrovascular diseases (I60-I69)	8,845	6.0	41.5
4	Diabetes mellitus (E10-E14)	6,941	4.7	32.6
5	Chronic lower respiratory diseases (J40-J47)	4,797	3.3	22.5
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	4,396	3.0	20.6
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	4,324	3.0	20.3
8	Alzheimer's disease (G30)	4,018	2.7	18.9
9	Septicemia (A40-A41)	3,339	2.3	15.7
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	2,933	2.0	13.8
11	Influenza and pneumonia (J09-J18)	2,837	1.9	13.3
12	Certain conditions originating in the perinatal period (P00-P96)	1,752	1.2	8.2
13	Human immunodeficiency virus (HIV) disease (B20-B24)	1,244	0.8	5.8
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,099	0.7	5.2
15	Chronic liver disease and cirrhosis (K70,K73-K74)	1,083	0.7	5.1
...	All other causes (Residual)	31,465	21.5	147.8
Non-Hispanic, black, female, 1-4 years				
...	All causes	425	100.0	35.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	120	28.2	10.0
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	65	15.3	5.4
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	38	8.9	3.2
4	Malignant neoplasms (C00-C97)	29	6.8	2.4
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	22	5.2	1.8
6	Chronic lower respiratory diseases (J40-J47)	11	2.6	*
7	Influenza and pneumonia (J09-J18)	9	2.1	*
8	Septicemia (A40-A41)	6	1.4	*
8	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	6	1.4	*
8	Certain conditions originating in the perinatal period (P00-P96)	6	1.4	*
11	Anemias (D50-D64)	4	0.9	*
12	Cerebrovascular diseases (I60-I69)	3	0.7	*
12	Complications of medical and surgical care (Y40-Y84,Y88)	3	0.7	*
14	Diabetes mellitus (E10-E14)	1	0.2	*
14	Meningitis (G00,G03)	1	0.2	*
14	Acute bronchitis and bronchiolitis (J20-J21)	1	0.2	*
14	Diseases of appendix (K35-K38)	1	0.2	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1	0.2	*
...	All other causes (Residual)	98	23.1	8.2

... Category not applicable

\* Figure does not meet standards of reliability or precision

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality, 2013

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, female, 5-14 years				
...	All causes	479	100.0	15.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	114	23.8	3.7
2	Malignant neoplasms (C00-C97)	54	11.3	1.8
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	43	9.0	1.4
4	Chronic lower respiratory diseases (J40-J47)	32	6.7	1.1
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	28	5.8	0.9
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	22	4.6	0.7
7	Diseases of heart (I00-I09,I11,I13,I20-I51)	20	4.2	0.7
8	Influenza and pneumonia (J09-J18)	9	1.9	*
9	Cerebrovascular diseases (I60-I69)	8	1.7	*
10	Anemias (D50-D64)	6	1.3	*
11	Septicemia (A40-A41)	5	1.0	*
11	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	5	1.0	*
13	Certain conditions originating in the perinatal period (P00-P96)	3	0.6	*
14	Meningitis (G00,G03)	2	0.4	*
14	Pneumonitis due to solids and liquids (J69)	2	0.4	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2	0.4	*
14	Complications of medical and surgical care (Y40-Y84,Y88)	2	0.4	*
...	All other causes (Residual)	122	25.5	4.0
Non-Hispanic, black, female, 15-24 years				
...	All causes	1,466	100.0	43.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	381	26.0	11.3
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	243	16.6	7.2
3	Malignant neoplasms (C00-C97)	111	7.6	3.3
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	96	6.5	2.8
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	92	6.3	2.7
6	Pregnancy, childbirth and the puerperium (O00-O99)	69	4.7	2.0
7	Anemias (D50-D64)	26	1.8	0.8
8	Diabetes mellitus (E10-E14)	25	1.7	0.7
8	Chronic lower respiratory diseases (J40-J47)	25	1.7	0.7
10	Human immunodeficiency virus (HIV) disease (B20-B24)	24	1.6	0.7
11	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	22	1.5	0.7
12	Septicemia (A40-A41)	17	1.2	*
12	Cerebrovascular diseases (I60-I69)	17	1.2	*
14	Influenza and pneumonia (J09-J18)	15	1.0	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	7	0.5	*
...	All other causes (Residual)	296	20.2	8.8

... Category not applicable

\* Figure does not meet standards of reliability or precision

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality, 2013

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, female, 25-44 years				
...	All causes	8,367	100.0	145.8
1	Malignant neoplasms (C00-C97)	1,516	18.1	26.4
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,319	15.8	23.0
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	980	11.7	17.1
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	440	5.3	7.7
5	Human immunodeficiency virus (HIV) disease (B20-B24)	412	4.9	7.2
6	Diabetes mellitus (E10-E14)	316	3.8	5.5
7	Cerebrovascular diseases (I60-I69)	290	3.5	5.1
8	Pregnancy, childbirth and the puerperium (O00-O99)	235	2.8	4.1
9	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	170	2.0	3.0
10	Septicemia (A40-A41)	163	1.9	2.8
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	154	1.8	2.7
12	Chronic lower respiratory diseases (J40-J47)	130	1.6	2.3
13	Influenza and pneumonia (J09-J18)	120	1.4	2.1
14	Chronic liver disease and cirrhosis (K70,K73-K74)	111	1.3	1.9
15	Anemias (D50-D64)	102	1.2	1.8
...	All other causes (Residual)	1,909	22.8	33.3
Non-Hispanic, black, female, 45-64 years				
...	All causes	37,524	100.0	707.7
1	Malignant neoplasms (C00-C97)	11,983	31.9	226.0
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	8,115	21.6	153.1
3	Diabetes mellitus (E10-E14)	1,899	5.1	35.8
4	Cerebrovascular diseases (I60-I69)	1,824	4.9	34.4
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,407	3.7	26.5
6	Chronic lower respiratory diseases (J40-J47)	1,224	3.3	23.1
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	934	2.5	17.6
8	Septicemia (A40-A41)	870	2.3	16.4
9	Human immunodeficiency virus (HIV) disease (B20-B24)	686	1.8	12.9
10	Chronic liver disease and cirrhosis (K70,K73-K74)	678	1.8	12.8
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	594	1.6	11.2
12	Influenza and pneumonia (J09-J18)	559	1.5	10.5
13	Viral hepatitis (B15-B19)	278	0.7	5.2
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	236	0.6	4.5
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	146	0.4	2.8
15	Anemias (D50-D64)	146	0.4	2.8
...	All other causes (Residual)	5,945	15.8	112.1

... Category not applicable

\* Figure does not meet standards of reliability or precision

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality, 2013

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, female, 65 years and over				
...	All causes	95,215	100.0	4,069.6
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	24,924	26.2	1,065.3
2	Malignant neoplasms (C00-C97)	19,253	20.2	822.9
3	Cerebrovascular diseases (I60-I69)	6,686	7.0	285.8
4	Diabetes mellitus (E10-E14)	4,698	4.9	200.8
5	Alzheimer's disease (G30)	3,982	4.2	170.2
6	Chronic lower respiratory diseases (J40-J47)	3,370	3.5	144.0
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3,290	3.5	140.6
8	Septicemia (A40-A41)	2,254	2.4	96.3
9	Essential hypertension and hypertensive renal disease (I10,I12,I15)	2,244	2.4	95.9
10	Influenza and pneumonia (J09-J18)	2,095	2.2	89.5
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,123	1.2	48.0
12	Pneumonitis due to solids and liquids (J69)	597	0.6	25.5
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	406	0.4	17.4
14	Parkinson's disease (G20-G21)	397	0.4	17.0
15	Chronic liver disease and cirrhosis (K70,K73-K74)	294	0.3	12.6
...	All other causes (Residual)	19,602	20.6	837.8

... Category not applicable

\* Figure does not meet standards of reliability or precision

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality, 2013