

LCWK8. Infant, neonatal, and postneonatal deaths, percent of total deaths, and mortality rates for the 15 leading causes of infant death by Hispanic origin, races for non-Hispanic population, and sex: United States, 2010

[Rates are infant (under 1 year), neonatal (under 28 days), and postneonatal (28 days-11 months) deaths per 100,000 live births in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on birth and death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), Hispanic origin and sex	Number	Percent of total deaths	Mortality rate
Infants, Hispanic, both sexes				
...	All causes	5,170	100.0	547.0
1	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	1,336	25.8	141.3
2	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	834	16.1	88.2
3	Newborn affected by maternal complications of pregnancy (P01)	337	6.5	35.7
4	Sudden infant death syndrome (R95)	314	6.1	33.2
5	Newborn affected by complications of placenta, cord and membranes (P02)	201	3.9	21.3
6	Accidents (unintentional injuries) (V01-X59)	152	2.9	16.1
7	Bacterial sepsis of newborn (P36)	121	2.3	12.8
7	Neonatal hemorrhage (P50-P52,P54)	121	2.3	12.8
9	Necrotizing enterocolitis of newborn (P77)	111	2.1	11.7
10	Diseases of the circulatory system (I00-I99)	99	1.9	10.5
11	Respiratory distress of newborn (P22)	83	1.6	8.8
12	Diarrhea and gastroenteritis of infectious origin (A09)	73	1.4	7.7
13	Assault (homicide) (*U01,X85-Y09)	69	1.3	7.3
14	Atelectasis (P28.0-P28.1)	56	1.1	5.9
15	Intrauterine hypoxia and birth asphyxia (P20-P21)	53	1.0	5.6
...	All other causes (Residual)	1,210	23.4	128.0
Infants, Hispanic, male				
...	All causes	2,870	100.0	596.3
1	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	659	23.0	136.9
2	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	496	17.3	103.0
3	Sudden infant death syndrome (R95)	192	6.7	39.9
4	Newborn affected by maternal complications of pregnancy (P01)	177	6.2	36.8
5	Newborn affected by complications of placenta, cord and membranes (P02)	104	3.6	21.6
6	Accidents (unintentional injuries) (V01-X59)	87	3.0	18.1
7	Bacterial sepsis of newborn (P36)	78	2.7	16.2
8	Neonatal hemorrhage (P50-P52,P54)	73	2.5	15.2
9	Necrotizing enterocolitis of newborn (P77)	66	2.3	13.7
10	Diseases of the circulatory system (I00-I99)	59	2.1	12.3
11	Respiratory distress of newborn (P22)	57	2.0	11.8
12	Diarrhea and gastroenteritis of infectious origin (A09)	39	1.4	8.1
13	Assault (homicide) (*U01,X85-Y09)	36	1.3	7.5
14	Atelectasis (P28.0-P28.1)	35	1.2	7.3
15	Intrauterine hypoxia and birth asphyxia (P20-P21)	33	1.1	6.9
...	All other causes (Residual)	679	23.7	141.1

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK8. Infant, neonatal, and postneonatal deaths, percent of total deaths, and mortality rates for the 15 leading causes of infant death by Hispanic origin, races for non-Hispanic population, and sex: United States, 2010

[Rates are infant (under 1 year), neonatal (under 28 days), and postneonatal (28 days-11 months) deaths per 100,000 live births in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on birth and death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), Hispanic origin and sex	Number	Percent of total deaths	Mortality rate
Infants, Hispanic, female				
...	All causes	2,300	100.0	495.8
1	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	677	29.4	146.0
2	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	338	14.7	72.9
3	Newborn affected by maternal complications of pregnancy (P01)	160	7.0	34.5
4	Sudden infant death syndrome (R95)	122	5.3	26.3
5	Newborn affected by complications of placenta, cord and membranes (P02)	97	4.2	20.9
6	Accidents (unintentional injuries) (V01-X59)	65	2.8	14.0
7	Neonatal hemorrhage (P50-P52,P54)	48	2.1	10.3
8	Necrotizing enterocolitis of newborn (P77)	45	2.0	9.7
9	Bacterial sepsis of newborn (P36)	43	1.9	9.3
10	Diseases of the circulatory system (I00-I99)	40	1.7	8.6
11	Diarrhea and gastroenteritis of infectious origin (A09)	34	1.5	7.3
12	Assault (homicide) (*U01,X85-Y09)	33	1.4	7.1
13	Respiratory distress of newborn (P22)	26	1.1	5.6
14	Atelectasis (P28.0-P28.1)	21	0.9	4.5
15	Septicemia (A40-A41)	20	0.9	4.3
15	Intrauterine hypoxia and birth asphyxia (P20-P21)	20	0.9	4.3
...	All other causes (Residual)	511	22.2	110.2
Infants, non-Hispanic white, both sexes				
...	All causes	11,025	100.0	509.8
1	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	2,535	23.0	117.2
2	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	1,522	13.8	70.4
3	Sudden infant death syndrome (R95)	1,033	9.4	47.8
4	Newborn affected by maternal complications of pregnancy (P01)	657	6.0	30.4
5	Accidents (unintentional injuries) (V01-X59)	566	5.1	26.2
6	Newborn affected by complications of placenta, cord and membranes (P02)	449	4.1	20.8
7	Bacterial sepsis of newborn (P36)	270	2.4	12.5
8	Diseases of the circulatory system (I00-I99)	237	2.1	11.0
8	Respiratory distress of newborn (P22)	237	2.1	11.0
10	Neonatal hemorrhage (P50-P52,P54)	220	2.0	10.2
11	Necrotizing enterocolitis of newborn (P77)	172	1.6	8.0
12	Intrauterine hypoxia and birth asphyxia (P20-P21)	157	1.4	7.3
13	Assault (homicide) (*U01,X85-Y09)	147	1.3	6.8
14	Atelectasis (P28.0-P28.1)	123	1.1	5.7
15	Diarrhea and gastroenteritis of infectious origin (A09)	107	1.0	4.9
...	All other causes (Residual)	2,593	23.5	119.9

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK8. Infant, neonatal, and postneonatal deaths, percent of total deaths, and mortality rates for the 15 leading causes of infant death by Hispanic origin, races for non-Hispanic population, and sex: United States, 2010

[Rates are infant (under 1 year), neonatal (under 28 days), and postneonatal (28 days-11 months) deaths per 100,000 live births in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on birth and death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), Hispanic origin and sex	Number	Percent of total deaths	Mortality rate
Infants, non-Hispanic white, male				
...	All causes	6,144	100.0	553.8
1	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	1,307	21.3	117.8
2	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	824	13.4	74.3
3	Sudden infant death syndrome (R95)	605	9.8	54.5
4	Newborn affected by maternal complications of pregnancy (P01)	376	6.1	33.9
5	Accidents (unintentional injuries) (V01-X59)	325	5.3	29.3
6	Newborn affected by complications of placenta, cord and membranes (P02)	254	4.1	22.9
7	Bacterial sepsis of newborn (P36)	142	2.3	12.8
8	Respiratory distress of newborn (P22)	140	2.3	12.6
9	Neonatal hemorrhage (P50-P52,P54)	133	2.2	12.0
10	Diseases of the circulatory system (I00-I99)	129	2.1	11.6
11	Necrotizing enterocolitis of newborn (P77)	103	1.7	9.3
12	Assault (homicide) (*U01,X85-Y09)	93	1.5	8.4
13	Intrauterine hypoxia and birth asphyxia (P20-P21)	86	1.4	7.8
14	Atelectasis (P28.0-P28.1)	70	1.1	6.3
15	Diarrhea and gastroenteritis of infectious origin (A09)	65	1.1	5.9
...	All other causes (Residual)	1,492	24.3	134.5
Infants, non-Hispanic white, female				
...	All causes	4,881	100.0	463.5
1	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	1,228	25.2	116.6
2	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	698	14.3	66.3
3	Sudden infant death syndrome (R95)	428	8.8	40.6
4	Newborn affected by maternal complications of pregnancy (P01)	281	5.8	26.7
5	Accidents (unintentional injuries) (V01-X59)	241	4.9	22.9
6	Newborn affected by complications of placenta, cord and membranes (P02)	195	4.0	18.5
7	Bacterial sepsis of newborn (P36)	128	2.6	12.2
8	Diseases of the circulatory system (I00-I99)	108	2.2	10.3
9	Respiratory distress of newborn (P22)	97	2.0	9.2
10	Neonatal hemorrhage (P50-P52,P54)	87	1.8	8.3
11	Intrauterine hypoxia and birth asphyxia (P20-P21)	71	1.5	6.7
12	Necrotizing enterocolitis of newborn (P77)	69	1.4	6.6
13	Assault (homicide) (*U01,X85-Y09)	54	1.1	5.1
14	Atelectasis (P28.0-P28.1)	53	1.1	5.0
15	Hydrops fetalis not due to hemolytic disease (P83.2)	45	0.9	4.3
...	All other causes (Residual)	1,098	22.5	104.3

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK8. Infant, neonatal, and postneonatal deaths, percent of total deaths, and mortality rates for the 15 leading causes of infant death by Hispanic origin, races for non-Hispanic population, and sex: United States, 2010

[Rates are infant (under 1 year), neonatal (under 28 days), and postneonatal (28 days-11 months) deaths per 100,000 live births in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on birth and death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), Hispanic origin and sex	Number	Percent of total deaths	Mortality rate
Infants, non-Hispanic black, both sexes				
...	All causes	7,071	100.0	1,198.9
1	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	1,550	21.9	262.8
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	969	13.7	164.3
3	Sudden infant death syndrome (R95)	639	9.0	108.3
4	Newborn affected by maternal complications of pregnancy (P01)	465	6.6	78.8
5	Accidents (unintentional injuries) (V01-X59)	339	4.8	57.5
6	Newborn affected by complications of placenta, cord and membranes (P02)	312	4.4	52.9
7	Respiratory distress of newborn (P22)	170	2.4	28.8
8	Necrotizing enterocolitis of newborn (P77)	168	2.4	28.5
9	Bacterial sepsis of newborn (P36)	166	2.3	28.1
10	Diseases of the circulatory system (I00-I99)	144	2.0	24.4
11	Diarrhea and gastroenteritis of infectious origin (A09)	123	1.7	20.9
12	Neonatal hemorrhage (P50-P52,P54)	112	1.6	19.0
13	Septicemia (A40-A41)	87	1.2	14.8
14	Assault (homicide) (*U01,X85-Y09)	85	1.2	14.4
15	Intrauterine hypoxia and birth asphyxia (P20-P21)	72	1.0	12.2
...	All other causes (Residual)	1,670	23.6	283.1
Infants, non-Hispanic black, male				
...	All causes	3,931	100.0	1,308.2
1	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	861	21.9	286.5
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	511	13.0	170.1
3	Sudden infant death syndrome (R95)	372	9.5	123.8
4	Newborn affected by maternal complications of pregnancy (P01)	245	6.2	81.5
5	Accidents (unintentional injuries) (V01-X59)	194	4.9	64.6
6	Newborn affected by complications of placenta, cord and membranes (P02)	148	3.8	49.3
7	Respiratory distress of newborn (P22)	102	2.6	33.9
8	Necrotizing enterocolitis of newborn (P77)	96	2.4	31.9
9	Bacterial sepsis of newborn (P36)	93	2.4	30.9
10	Diseases of the circulatory system (I00-I99)	83	2.1	27.6
11	Diarrhea and gastroenteritis of infectious origin (A09)	75	1.9	25.0
12	Neonatal hemorrhage (P50-P52,P54)	70	1.8	23.3
13	Septicemia (A40-A41)	45	1.1	15.0
14	Influenza and pneumonia (J09-J18)	42	1.1	14.0
15	Intrauterine hypoxia and birth asphyxia (P20-P21)	40	1.0	13.3
15	Assault (homicide) (*U01,X85-Y09)	40	1.0	13.3
...	All other causes (Residual)	914	23.3	304.2

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK8. Infant, neonatal, and postneonatal deaths, percent of total deaths, and mortality rates for the 15 leading causes of infant death by Hispanic origin, races for non-Hispanic population, and sex: United States, 2010

[Rates are infant (under 1 year), neonatal (under 28 days), and postneonatal (28 days-11 months) deaths per 100,000 live births in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on birth and death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), Hispanic origin and sex	Number	Percent of total deaths	Mortality rate
Infants, non-Hispanic black, female				
...	All causes	3,140	100.0	1,085.3
1	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	689	21.9	238.1
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	458	14.6	158.3
3	Sudden infant death syndrome (R95)	267	8.5	92.3
4	Newborn affected by maternal complications of pregnancy (P01)	220	7.0	76.0
5	Newborn affected by complications of placenta, cord and membranes (P02)	164	5.2	56.7
6	Accidents (unintentional injuries) (V01-X59)	145	4.6	50.1
7	Bacterial sepsis of newborn (P36)	73	2.3	25.2
8	Necrotizing enterocolitis of newborn (P77)	72	2.3	24.9
9	Respiratory distress of newborn (P22)	68	2.2	23.5
10	Diseases of the circulatory system (I00-I99)	61	1.9	21.1
11	Diarrhea and gastroenteritis of infectious origin (A09)	48	1.5	16.6
12	Assault (homicide) (*U01,X85-Y09)	45	1.4	15.6
13	Septicemia (A40-A41)	42	1.3	14.5
13	Neonatal hemorrhage (P50-P52,P54)	42	1.3	14.5
15	Intrauterine hypoxia and birth asphyxia (P20-P21)	32	1.0	11.1
...	All other causes (Residual)	714	22.7	246.8
Neonates, Hispanic, both sexes				
...	All causes	3,524	100.0	372.8
1	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	941	26.7	99.6
2	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	818	23.2	86.5
3	Newborn affected by maternal complications of pregnancy (P01)	334	9.5	35.3
4	Newborn affected by complications of placenta, cord and membranes (P02)	197	5.6	20.8
5	Neonatal hemorrhage (P50-P52,P54)	121	3.4	12.8
6	Bacterial sepsis of newborn (P36)	112	3.2	11.8
7	Necrotizing enterocolitis of newborn (P77)	93	2.6	9.8
8	Respiratory distress of newborn (P22)	83	2.4	8.8
9	Atelectasis (P28.0-P28.1)	54	1.5	5.7
10	Intrauterine hypoxia and birth asphyxia (P20-P21)	49	1.4	5.2
11	Pulmonary hemorrhage originating in the perinatal period (P26)	33	0.9	3.5
12	Newborn affected by other complications of labor and delivery (P03)	28	0.8	3.0
12	Hydrops fetalis not due to hemolytic disease (P83.2)	28	0.8	3.0
14	Newborn affected by other maternal conditions which may be unrelated to present pregnancy (P00.1-P00.9)	26	0.7	2.8
15	Interstitial emphysema and related conditions originating in the perinatal period (P25)	24	0.7	2.5
15	Sudden infant death syndrome (R95)	24	0.7	2.5
...	All other causes (Residual)	559	15.9	59.1

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK8. Infant, neonatal, and postneonatal deaths, percent of total deaths, and mortality rates for the 15 leading causes of infant death by Hispanic origin, races for non-Hispanic population, and sex: United States, 2010

[Rates are infant (under 1 year), neonatal (under 28 days), and postneonatal (28 days-11 months) deaths per 100,000 live births in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on birth and death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), Hispanic origin and sex	Number	Percent of total deaths	Mortality rate
Neonates, Hispanic, male				
...	All causes	1,960	100.0	407.2
1	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	484	24.7	100.6
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	477	24.3	99.1
3	Newborn affected by maternal complications of pregnancy (P01)	177	9.0	36.8
4	Newborn affected by complications of placenta, cord and membranes (P02)	100	5.1	20.8
5	Bacterial sepsis of newborn (P36)	74	3.8	15.4
6	Neonatal hemorrhage (P50-P52,P54)	73	3.7	15.2
7	Necrotizing enterocolitis of newborn (P77)	58	3.0	12.0
8	Respiratory distress of newborn (P22)	57	2.9	11.8
9	Atelectasis (P28.0-P28.1)	33	1.7	6.9
10	Intrauterine hypoxia and birth asphyxia (P20-P21)	30	1.5	6.2
11	Pulmonary hemorrhage originating in the perinatal period (P26)	23	1.2	4.8
12	Diseases of the circulatory system (I00-I99)	16	0.8	*
12	Sudden infant death syndrome (R95)	16	0.8	*
14	Hydrops fetalis not due to hemolytic disease (P83.2)	14	0.7	*
15	Newborn affected by other maternal conditions which may be unrelated to present pregnancy (P00.1-P00.9)	12	0.6	*
...	All other causes (Residual)	316	16.1	65.7
Neonates, Hispanic, female				
...	All causes	1,564	100.0	337.2
1	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	464	29.7	100.0
2	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	334	21.4	72.0
3	Newborn affected by maternal complications of pregnancy (P01)	157	10.0	33.8
4	Newborn affected by complications of placenta, cord and membranes (P02)	97	6.2	20.9
5	Neonatal hemorrhage (P50-P52,P54)	48	3.1	10.3
6	Bacterial sepsis of newborn (P36)	38	2.4	8.2
7	Necrotizing enterocolitis of newborn (P77)	35	2.2	7.5
8	Respiratory distress of newborn (P22)	26	1.7	5.6
9	Atelectasis (P28.0-P28.1)	21	1.3	4.5
10	Intrauterine hypoxia and birth asphyxia (P20-P21)	19	1.2	*
11	Newborn affected by other complications of labor and delivery (P03)	18	1.2	*
12	Newborn affected by other maternal conditions which may be unrelated to present pregnancy (P00.1-P00.9)	14	0.9	*
12	Interstitial emphysema and related conditions originating in the perinatal period (P25)	14	0.9	*
12	Hydrops fetalis not due to hemolytic disease (P83.2)	14	0.9	*
15	Newborn affected by maternal hypertensive disorders (P00.0)	13	0.8	*
...	All other causes (Residual)	252	16.1	54.3

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK8. Infant, neonatal, and postneonatal deaths, percent of total deaths, and mortality rates for the 15 leading causes of infant death by Hispanic origin, races for non-Hispanic population, and sex: United States, 2010

[Rates are infant (under 1 year), neonatal (under 28 days), and postneonatal (28 days-11 months) deaths per 100,000 live births in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on birth and death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), Hispanic origin and sex	Number	Percent of total deaths	Mortality rate
Neonates, non-Hispanic white, both sexes				
...	All causes	7,212	100.0	333.5
1	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	1,851	25.7	85.6
2	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	1,490	20.7	68.9
3	Newborn affected by maternal complications of pregnancy (P01)	649	9.0	30.0
4	Newborn affected by complications of placenta, cord and membranes (P02)	443	6.1	20.5
5	Bacterial sepsis of newborn (P36)	257	3.6	11.9
6	Respiratory distress of newborn (P22)	227	3.1	10.5
7	Neonatal hemorrhage (P50-P52,P54)	216	3.0	10.0
8	Necrotizing enterocolitis of newborn (P77)	159	2.2	7.4
9	Intrauterine hypoxia and birth asphyxia (P20-P21)	155	2.1	7.2
10	Atelectasis (P28.0-P28.1)	114	1.6	5.3
11	Sudden infant death syndrome (R95)	107	1.5	4.9
12	Hydrops fetalis not due to hemolytic disease (P83.2)	87	1.2	4.0
13	Accidents (unintentional injuries) (V01-X59)	74	1.0	3.4
14	Pulmonary hemorrhage originating in the perinatal period (P26)	63	0.9	2.9
15	Interstitial emphysema and related conditions originating in the perinatal period (P25)	59	0.8	2.7
...	All other causes (Residual)	1,261	17.5	58.3
Neonates, non-Hispanic white, male				
...	All causes	3,974	100.0	358.2
1	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	967	24.3	87.2
2	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	809	20.4	72.9
3	Newborn affected by maternal complications of pregnancy (P01)	373	9.4	33.6
4	Newborn affected by complications of placenta, cord and membranes (P02)	250	6.3	22.5
5	Bacterial sepsis of newborn (P36)	135	3.4	12.2
6	Respiratory distress of newborn (P22)	132	3.3	11.9
7	Neonatal hemorrhage (P50-P52,P54)	131	3.3	11.8
8	Necrotizing enterocolitis of newborn (P77)	97	2.4	8.7
9	Intrauterine hypoxia and birth asphyxia (P20-P21)	85	2.1	7.7
10	Atelectasis (P28.0-P28.1)	63	1.6	5.7
11	Sudden infant death syndrome (R95)	60	1.5	5.4
12	Hydrops fetalis not due to hemolytic disease (P83.2)	42	1.1	3.8
12	Accidents (unintentional injuries) (V01-X59)	42	1.1	3.8
14	Pulmonary hemorrhage originating in the perinatal period (P26)	37	0.9	3.3
15	Interstitial emphysema and related conditions originating in the perinatal period (P25)	36	0.9	3.2
...	All other causes (Residual)	715	18.0	64.5

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK8. Infant, neonatal, and postneonatal deaths, percent of total deaths, and mortality rates for the 15 leading causes of infant death by Hispanic origin, races for non-Hispanic population, and sex: United States, 2010

[Rates are infant (under 1 year), neonatal (under 28 days), and postneonatal (28 days-11 months) deaths per 100,000 live births in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on birth and death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), Hispanic origin and sex	Number	Percent of total deaths	Mortality rate
Neonates, non-Hispanic white, female				
...	All causes	3,238	100.0	307.5
1	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	884	27.3	83.9
2	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	681	21.0	64.7
3	Newborn affected by maternal complications of pregnancy (P01)	276	8.5	26.2
4	Newborn affected by complications of placenta, cord and membranes (P02)	193	6.0	18.3
5	Bacterial sepsis of newborn (P36)	122	3.8	11.6
6	Respiratory distress of newborn (P22)	95	2.9	9.0
7	Neonatal hemorrhage (P50-P52,P54)	85	2.6	8.1
8	Intrauterine hypoxia and birth asphyxia (P20-P21)	70	2.2	6.6
9	Necrotizing enterocolitis of newborn (P77)	62	1.9	5.9
10	Atelectasis (P28.0-P28.1)	51	1.6	4.8
11	Sudden infant death syndrome (R95)	47	1.5	4.5
12	Hydrops fetalis not due to hemolytic disease (P83.2)	45	1.4	4.3
13	Accidents (unintentional injuries) (V01-X59)	32	1.0	3.0
14	Pulmonary hemorrhage originating in the perinatal period (P26)	26	0.8	2.5
15	Interstitial emphysema and related conditions originating in the perinatal period (P25)	23	0.7	2.2
...	All other causes (Residual)	546	16.9	51.9
Neonates, non-Hispanic black, both sexes				
...	All causes	4,549	100.0	771.3
1	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	1,518	33.4	257.4
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	629	13.8	106.6
3	Newborn affected by maternal complications of pregnancy (P01)	463	10.2	78.5
4	Newborn affected by complications of placenta, cord and membranes (P02)	307	6.7	52.1
5	Respiratory distress of newborn (P22)	164	3.6	27.8
6	Bacterial sepsis of newborn (P36)	157	3.5	26.6
7	Necrotizing enterocolitis of newborn (P77)	138	3.0	23.4
8	Neonatal hemorrhage (P50-P52,P54)	106	2.3	18.0
9	Intrauterine hypoxia and birth asphyxia (P20-P21)	64	1.4	10.9
10	Pulmonary hemorrhage originating in the perinatal period (P26)	60	1.3	10.2
11	Atelectasis (P28.0-P28.1)	54	1.2	9.2
11	Sudden infant death syndrome (R95)	54	1.2	9.2
13	Accidents (unintentional injuries) (V01-X59)	34	0.7	5.8
14	Newborn affected by maternal hypertensive disorders (P00.0)	30	0.7	5.1
15	Newborn affected by other complications of labor and delivery (P03)	29	0.6	4.9
15	Slow fetal growth and fetal malnutrition (P05)	29	0.6	4.9
...	All other causes (Residual)	713	15.7	120.9

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK8. Infant, neonatal, and postneonatal deaths, percent of total deaths, and mortality rates for the 15 leading causes of infant death by Hispanic origin, races for non-Hispanic population, and sex: United States, 2010

[Rates are infant (under 1 year), neonatal (under 28 days), and postneonatal (28 days-11 months) deaths per 100,000 live births in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on birth and death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), Hispanic origin and sex	Number	Percent of total deaths	Mortality rate
Neonates, non-Hispanic black, male				
...	All causes	2,499	100.0	831.6
1	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	852	34.1	283.5
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	332	13.3	110.5
3	Newborn affected by maternal complications of pregnancy (P01)	243	9.7	80.9
4	Newborn affected by complications of placenta, cord and membranes (P02)	146	5.8	48.6
5	Respiratory distress of newborn (P22)	96	3.8	31.9
6	Bacterial sepsis of newborn (P36)	89	3.6	29.6
7	Necrotizing enterocolitis of newborn (P77)	80	3.2	26.6
8	Neonatal hemorrhage (P50-P52,P54)	67	2.7	22.3
9	Intrauterine hypoxia and birth asphyxia (P20-P21)	36	1.4	12.0
10	Pulmonary hemorrhage originating in the perinatal period (P26)	33	1.3	11.0
11	Atelectasis (P28.0-P28.1)	30	1.2	10.0
12	Sudden infant death syndrome (R95)	26	1.0	8.7
13	Accidents (unintentional injuries) (V01-X59)	19	0.8	*
14	Newborn affected by maternal hypertensive disorders (P00.0)	17	0.7	*
15	Slow fetal growth and fetal malnutrition (P05)	16	0.6	*
...	All other causes (Residual)	417	16.7	138.8
Neonates, non-Hispanic black, female				
...	All causes	2,050	100.0	708.6
1	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	666	32.5	230.2
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	297	14.5	102.7
3	Newborn affected by maternal complications of pregnancy (P01)	220	10.7	76.0
4	Newborn affected by complications of placenta, cord and membranes (P02)	161	7.9	55.6
5	Respiratory distress of newborn (P22)	68	3.3	23.5
5	Bacterial sepsis of newborn (P36)	68	3.3	23.5
7	Necrotizing enterocolitis of newborn (P77)	58	2.8	20.0
8	Neonatal hemorrhage (P50-P52,P54)	39	1.9	13.5
9	Intrauterine hypoxia and birth asphyxia (P20-P21)	28	1.4	9.7
9	Sudden infant death syndrome (R95)	28	1.4	9.7
11	Pulmonary hemorrhage originating in the perinatal period (P26)	27	1.3	9.3
12	Atelectasis (P28.0-P28.1)	24	1.2	8.3
13	Accidents (unintentional injuries) (V01-X59)	15	0.7	*
14	Newborn affected by other complications of labor and delivery (P03)	14	0.7	*
15	Newborn affected by maternal hypertensive disorders (P00.0)	13	0.6	*
15	Slow fetal growth and fetal malnutrition (P05)	13	0.6	*
...	All other causes (Residual)	311	15.2	107.5

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK8. Infant, neonatal, and postneonatal deaths, percent of total deaths, and mortality rates for the 15 leading causes of infant death by Hispanic origin, races for non-Hispanic population, and sex: United States, 2010

[Rates are infant (under 1 year), neonatal (under 28 days), and postneonatal (28 days-11 months) deaths per 100,000 live births in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on birth and death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), Hispanic origin and sex	Number	Percent of total deaths	Mortality rate
Postneonates, Hispanic, both sexes				
...	All causes	1,646	100.0	174.1
1	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	395	24.0	41.8
2	Sudden infant death syndrome (R95)	290	17.6	30.7
3	Accidents (unintentional injuries) (V01-X59)	135	8.2	14.3
4	Diseases of the circulatory system (I00-I99)	77	4.7	8.1
5	Diarrhea and gastroenteritis of infectious origin (A09)	70	4.3	7.4
6	Assault (homicide) (*U01,X85-Y09)	58	3.5	6.1
7	Septicemia (A40-A41)	44	2.7	4.7
8	Influenza and pneumonia (J09-J18)	32	1.9	3.4
9	Diseases of the blood and blood-forming organs and certain disorders involving the immune mechanism (D50-D89)	22	1.3	2.3
10	Necrotizing enterocolitis of newborn (P77)	18	1.1	*
11	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	16	1.0	*
12	Renal failure and other disorders of kidney (N17-N19,N25,N27)	15	0.9	*
13	Chronic respiratory disease originating in the perinatal period (P27)	14	0.9	*
14	Malignant neoplasms (C00-C97)	12	0.7	*
15	Whooping cough (A37)	11	0.7	*
15	Meningitis (G00,G03)	11	0.7	*
...	All other causes (Residual)	426	25.9	45.1
Postneonates, Hispanic, male				
...	All causes	910	100.0	189.1
1	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	182	20.0	37.8
2	Sudden infant death syndrome (R95)	176	19.3	36.6
3	Accidents (unintentional injuries) (V01-X59)	77	8.5	16.0
4	Diseases of the circulatory system (I00-I99)	43	4.7	8.9
5	Diarrhea and gastroenteritis of infectious origin (A09)	37	4.1	7.7
6	Assault (homicide) (*U01,X85-Y09)	35	3.8	7.3
7	Septicemia (A40-A41)	24	2.6	5.0
8	Influenza and pneumonia (J09-J18)	17	1.9	*
9	Diseases of the blood and blood-forming organs and certain disorders involving the immune mechanism (D50-D89)	12	1.3	*
9	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	12	1.3	*
11	Renal failure and other disorders of kidney (N17-N19,N25,N27)	9	1.0	*
12	Necrotizing enterocolitis of newborn (P77)	8	0.9	*
13	Malignant neoplasms (C00-C97)	7	0.8	*
13	Meningitis (G00,G03)	7	0.8	*
15	Anoxic brain damage, not elsewhere classified (G93.1)	6	0.7	*
15	Chronic respiratory disease originating in the perinatal period (P27)	6	0.7	*
...	All other causes (Residual)	252	27.7	52.4

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK8. Infant, neonatal, and postneonatal deaths, percent of total deaths, and mortality rates for the 15 leading causes of infant death by Hispanic origin, races for non-Hispanic population, and sex: United States, 2010

[Rates are infant (under 1 year), neonatal (under 28 days), and postneonatal (28 days-11 months) deaths per 100,000 live births in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on birth and death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), Hispanic origin and sex	Number	Percent of total deaths	Mortality rate
Postneonates, Hispanic, female				
...	All causes	736	100.0	158.7
1	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	213	28.9	45.9
2	Sudden infant death syndrome (R95)	114	15.5	24.6
3	Accidents (unintentional injuries) (V01-X59)	58	7.9	12.5
4	Diseases of the circulatory system (I00-I99)	34	4.6	7.3
5	Diarrhea and gastroenteritis of infectious origin (A09)	33	4.5	7.1
6	Assault (homicide) (*U01,X85-Y09)	23	3.1	5.0
7	Septicemia (A40-A41)	20	2.7	4.3
8	Influenza and pneumonia (J09-J18)	15	2.0	*
9	Diseases of the blood and blood-forming organs and certain disorders involving the immune mechanism (D50-D89)	10	1.4	*
9	Necrotizing enterocolitis of newborn (P77)	10	1.4	*
11	Chronic respiratory disease originating in the perinatal period (P27)	8	1.1	*
12	Whooping cough (A37)	6	0.8	*
12	Renal failure and other disorders of kidney (N17-N19,N25,N27)	6	0.8	*
14	Malignant neoplasms (C00-C97)	5	0.7	*
14	Hernia of abdominal cavity and intestinal obstruction without hernia (K40-K46,K56)	5	0.7	*
14	Bacterial sepsis of newborn (P36)	5	0.7	*
...	All other causes (Residual)	171	23.2	36.9
Postneonates, non-Hispanic white, both sexes				
...	All causes	3,813	100.0	176.3
1	Sudden infant death syndrome (R95)	926	24.3	42.8
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	684	17.9	31.6
3	Accidents (unintentional injuries) (V01-X59)	492	12.9	22.8
4	Diseases of the circulatory system (I00-I99)	180	4.7	8.3
5	Assault (homicide) (*U01,X85-Y09)	136	3.6	6.3
6	Diarrhea and gastroenteritis of infectious origin (A09)	103	2.7	4.8
7	Septicemia (A40-A41)	73	1.9	3.4
8	Influenza and pneumonia (J09-J18)	69	1.8	3.2
9	Chronic respiratory disease originating in the perinatal period (P27)	33	0.9	1.5
10	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	32	0.8	1.5
11	Malignant neoplasms (C00-C97)	28	0.7	1.3
12	Diseases of the blood and blood-forming organs and certain disorders involving the immune mechanism (D50-D89)	19	0.5	*
13	Anoxic brain damage, not elsewhere classified (G93.1)	16	0.4	*
14	Meningitis (G00,G03)	15	0.4	*
15	Bacterial sepsis of newborn (P36)	13	0.3	*
15	Necrotizing enterocolitis of newborn (P77)	13	0.3	*
...	All other causes (Residual)	981	25.7	45.4

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK8. Infant, neonatal, and postneonatal deaths, percent of total deaths, and mortality rates for the 15 leading causes of infant death by Hispanic origin, races for non-Hispanic population, and sex: United States, 2010

[Rates are infant (under 1 year), neonatal (under 28 days), and postneonatal (28 days-11 months) deaths per 100,000 live births in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on birth and death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), Hispanic origin and sex	Number	Percent of total deaths	Mortality rate
Postneonates, non-Hispanic white, male				
...	All causes	2,170	100.0	195.6
1	Sudden infant death syndrome (R95)	545	25.1	49.1
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	340	15.7	30.6
3	Accidents (unintentional injuries) (V01-X59)	283	13.0	25.5
4	Diseases of the circulatory system (I00-I99)	94	4.3	8.5
5	Assault (homicide) (*U01,X85-Y09)	86	4.0	7.8
6	Diarrhea and gastroenteritis of infectious origin (A09)	64	2.9	5.8
7	Septicemia (A40-A41)	46	2.1	4.1
8	Influenza and pneumonia (J09-J18)	39	1.8	3.5
9	Chronic respiratory disease originating in the perinatal period (P27)	17	0.8	*
10	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	15	0.7	*
11	Malignant neoplasms (C00-C97)	14	0.6	*
12	Acute bronchitis and acute bronchiolitis (J20-J21)	11	0.5	*
13	Diseases of the blood and blood-forming organs and certain disorders involving the immune mechanism (D50-D89)	8	0.4	*
13	Respiratory distress of newborn (P22)	8	0.4	*
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	7	0.3	*
15	Atelectasis (P28.0-P28.1)	7	0.3	*
15	Bacterial sepsis of newborn (P36)	7	0.3	*
...	All other causes (Residual)	579	26.7	52.2
Postneonates, non-Hispanic white, female				
...	All causes	1,643	100.0	156.0
1	Sudden infant death syndrome (R95)	381	23.2	36.2
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	344	20.9	32.7
3	Accidents (unintentional injuries) (V01-X59)	209	12.7	19.8
4	Diseases of the circulatory system (I00-I99)	86	5.2	8.2
5	Assault (homicide) (*U01,X85-Y09)	50	3.0	4.7
6	Diarrhea and gastroenteritis of infectious origin (A09)	39	2.4	3.7
7	Influenza and pneumonia (J09-J18)	30	1.8	2.8
8	Septicemia (A40-A41)	27	1.6	2.6
9	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	17	1.0	*
10	Chronic respiratory disease originating in the perinatal period (P27)	16	1.0	*
11	Malignant neoplasms (C00-C97)	14	0.9	*
12	Diseases of the blood and blood-forming organs and certain disorders involving the immune mechanism (D50-D89) (D50-D89)	11	0.7	*
13	Meningitis (G00,G03)	10	0.6	*
13	Anoxic brain damage, not elsewhere classified (G93.1)	10	0.6	*
15	Volume depletion, disorders of fluid, electrolyte and acid-base balance (E86-E87)	7	0.4	*
15	Bronchitis, chronic and unspecified (J40-J42)	7	0.4	*
15	Necrotizing enterocolitis of newborn (P77)	7	0.4	*
...	All other causes (Residual)	378	23.0	35.9

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK8. Infant, neonatal, and postneonatal deaths, percent of total deaths, and mortality rates for the 15 leading causes of infant death by Hispanic origin, races for non-Hispanic population, and sex: United States, 2010

[Rates are infant (under 1 year), neonatal (under 28 days), and postneonatal (28 days-11 months) deaths per 100,000 live births in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on birth and death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), Hispanic origin and sex	Number	Percent of total deaths	Mortality rate
Postneonates, non-Hispanic black, both sexes				
...	All causes	2,522	100.0	427.6
1	Sudden infant death syndrome (R95)	585	23.2	99.2
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	340	13.5	57.6
3	Accidents (unintentional injuries) (V01-X59)	305	12.1	51.7
4	Diseases of the circulatory system (I00-I99)	123	4.9	20.9
5	Diarrhea and gastroenteritis of infectious origin (A09)	116	4.6	19.7
6	Septicemia (A40-A41)	84	3.3	14.2
7	Assault (homicide) (*U01,X85-Y09)	78	3.1	13.2
8	Influenza and pneumonia (J09-J18)	71	2.8	12.0
9	Chronic respiratory disease originating in the perinatal period (P27)	37	1.5	6.3
10	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	32	1.3	5.4
11	Necrotizing enterocolitis of newborn (P77)	30	1.2	5.1
12	Meningitis (G00,G03)	17	0.7	*
13	Renal failure and other disorders of kidney (N17-N19,N25,N27)	16	0.6	*
14	Diseases of the blood and blood-forming organs and certain disorders involving the immune mechanism (D50-D89)	15	0.6	*
15	Volume depletion, disorders of fluid, electrolyte and acid-base balance (E86-E87)	11	0.4	*
...	All other causes (Residual)	662	26.2	112.2
Postneonates, non-Hispanic black, male				
...	All causes	1,432	100.0	476.6
1	Sudden infant death syndrome (R95)	346	24.2	115.1
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	179	12.5	59.6
3	Accidents (unintentional injuries) (V01-X59)	175	12.2	58.2
4	Diseases of the circulatory system (I00-I99)	70	4.9	23.3
5	Diarrhea and gastroenteritis of infectious origin (A09)	69	4.8	23.0
6	Septicemia (A40-A41)	44	3.1	14.6
7	Influenza and pneumonia (J09-J18)	42	2.9	14.0
8	Assault (homicide) (*U01,X85-Y09)	36	2.5	12.0
9	Chronic respiratory disease originating in the perinatal period (P27)	22	1.5	7.3
10	Necrotizing enterocolitis of newborn (P77)	16	1.1	*
11	Renal failure and other disorders of kidney (N17-N19,N25,N27)	10	0.7	*
12	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	9	0.6	*
13	Diseases of the blood and blood-forming organs and certain disorders involving the immune mechanism (D50-D89)	8	0.6	*
13	Hernia of abdominal cavity and intestinal obstruction without hernia (K40-K46,K56)	8	0.6	*
15	Volume depletion, disorders of fluid, electrolyte and acid-base balance (E86-E87)	7	0.5	*
15	Meningitis (G00,G03)	7	0.5	*
...	All other causes (Residual)	384	26.8	127.8

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK8. Infant, neonatal, and postneonatal deaths, percent of total deaths, and mortality rates for the 15 leading causes of infant death by Hispanic origin, races for non-Hispanic population, and sex: United States, 2010

[Rates are infant (under 1 year), neonatal (under 28 days), and postneonatal (28 days-11 months) deaths per 100,000 live births in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on birth and death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), Hispanic origin and sex	Number	Percent of total deaths	Mortality rate
Postneonates, non-Hispanic black, female				
...	All causes	1,090	100.0	376.7
1	Sudden infant death syndrome (R95)	239	21.9	82.6
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	161	14.8	55.6
3	Accidents (unintentional injuries) (V01-X59)	130	11.9	44.9
4	Diseases of the circulatory system (I00-I99)	53	4.9	18.3
5	Diarrhea and gastroenteritis of infectious origin (A09)	47	4.3	16.2
6	Assault (homicide) (*U01,X85-Y09)	42	3.9	14.5
7	Septicemia (A40-A41)	40	3.7	13.8
8	Influenza and pneumonia (J09-J18)	29	2.7	10.0
9	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	23	2.1	7.9
10	Chronic respiratory disease originating in the perinatal period (P27)	15	1.4	*
11	Necrotizing enterocolitis of newborn (P77)	14	1.3	*
12	Meningitis (G00,G03)	10	0.9	*
13	Diseases of the blood and blood-forming organs and certain disorders involving the immune mechanism (D50-D89)	7	0.6	*
14	Renal failure and other disorders of kidney (N17-N19,N25,N27)	6	0.6	*
15	Bacterial sepsis of newborn (P36)	5	0.5	*
...	All other causes (Residual)	269	24.7	93.0

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable