

LCWK9. Deaths, percent of total deaths, and death rates for the 15 leading causes of death: United States and each State, 2001

[Rates per 100,000 population.]

| Rank ¹ | Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992) and State | Number | Percent of total deaths | Rate |
|-------------------|--|-----------|-------------------------|---------|
| United States | | | | |
| ... | All causes | 2,416,425 | 100.0 | 848.5 |
| 1 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 700,142 | 29.0 | 245.8 |
| 2 | Malignant neoplasms (C00-C97) | 553,768 | 22.9 | 194.4 |
| 3 | Cerebrovascular diseases (I60-I69) | 163,538 | 6.8 | 57.4 |
| 4 | Chronic lower respiratory diseases (J40-J47) | 123,013 | 5.1 | 43.2 |
| 5 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 101,537 | 4.2 | 35.7 |
| 6 | Diabetes mellitus (E10-E14) | 71,372 | 3.0 | 25.1 |
| 7 | Influenza and pneumonia (J10-J18) | 62,034 | 2.6 | 21.8 |
| 8 | Alzheimer's disease (G30) | 53,852 | 2.2 | 18.9 |
| 9 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 39,480 | 1.6 | 13.9 |
| 10 | Septicemia (A40-A41) | 32,238 | 1.3 | 11.3 |
| 11 | Intentional self-harm (suicide) (*U03,X60-X84,Y87.0) | 30,622 | 1.3 | 10.8 |
| 12 | Chronic liver disease and cirrhosis (K70,K73-K74) | 27,035 | 1.1 | 9.5 |
| 13 | Assault (homicide) (*U01-*U02,X85-Y09,Y87.1) | 20,308 | 0.8 | 7.1 |
| 14 | Essential (primary) hypertension and hypertensive renal disease (I10,I12) | 19,250 | 0.8 | 6.8 |
| 15 | Pneumonitis due to solids and liquids (J69) | 17,301 | 0.7 | 6.1 |
| ... | All other causes (Residual) | 400,935 | 16.6 | 140.8 |
| Alabama | | | | |
| ... | All causes | 45,316 | 100.0 | 1,014.0 |
| 1 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 13,207 | 29.1 | 295.5 |
| 2 | Malignant neoplasms (C00-C97) | 9,801 | 21.6 | 219.3 |
| 3 | Cerebrovascular diseases (I60-I69) | 2,998 | 6.6 | 67.1 |
| 4 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 2,211 | 4.9 | 49.5 |
| 5 | Chronic lower respiratory diseases (J40-J47) | 2,204 | 4.9 | 49.3 |
| 6 | Diabetes mellitus (E10-E14) | 1,344 | 3.0 | 30.1 |
| 7 | Influenza and pneumonia (J10-J18) | 1,105 | 2.4 | 24.7 |
| 8 | Alzheimer's disease (G30) | 1,103 | 2.4 | 24.7 |
| 9 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 974 | 2.1 | 21.8 |
| 10 | Septicemia (A40-A41) | 771 | 1.7 | 17.3 |
| 11 | Intentional self-harm (suicide) (*U03,X60-X84,Y87.0) | 512 | 1.1 | 11.5 |
| 12 | Chronic liver disease and cirrhosis (K70,K73-K74) | 445 | 1.0 | 10.0 |
| 13 | Assault (homicide) (*U01-*U02,X85-Y09,Y87.1) | 425 | 0.9 | 9.5 |
| 14 | Essential (primary) hypertension and hypertensive renal disease (I10,I12) | 369 | 0.8 | 8.3 |
| 15 | Pneumonitis due to solids and liquids (J69) | 290 | 0.6 | 6.5 |
| ... | All other causes (Residual) | 7,557 | 16.7 | 169.1 |

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

... Category not applicable

LCWK9. Deaths, percent of total deaths, and death rates for the 15 leading causes of death: United States and each State, 2001

[Rates per 100,000 population.]

| Rank ¹ | Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992) and State | Number | Percent of total deaths | Rate |
|-------------------|--|--------|-------------------------|-------|
| Alaska | | | | |
| ... | All causes | 2,974 | 100.0 | 469.4 |
| 1 | Malignant neoplasms (C00-C97) | 689 | 23.2 | 108.7 |
| 2 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 603 | 20.3 | 95.2 |
| 3 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 345 | 11.6 | 54.4 |
| 4 | Cerebrovascular diseases (I60-I69) | 158 | 5.3 | 24.9 |
| 5 | Chronic lower respiratory diseases (J40-J47) | 147 | 4.9 | 23.2 |
| 6 | Intentional self-harm (suicide) (*U03,X60-X84,Y87.0) | 102 | 3.4 | 16.1 |
| 7 | Diabetes mellitus (E10-E14) | 80 | 2.7 | 12.6 |
| 8 | Atherosclerosis (I70) | 60 | 2.0 | 9.5 |
| 9 | Chronic liver disease and cirrhosis (K70,K73-K74) | 57 | 1.9 | 9.0 |
| 10 | Alzheimer's disease (G30) | 44 | 1.5 | 6.9 |
| 11 | Assault (homicide) (*U01-*U02,X85-Y09,Y87.1) | 38 | 1.3 | 6.0 |
| 12 | Influenza and pneumonia (J10-J18) | 36 | 1.2 | 5.7 |
| 13 | Septicemia (A40-A41) | 25 | 0.8 | 3.9 |
| 13 | Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99) | 25 | 0.8 | 3.9 |
| 15 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 24 | 0.8 | 3.8 |
| ... | All other causes (Residual) | 541 | 18.2 | 85.4 |
| Arizona | | | | |
| ... | All causes | 41,058 | 100.0 | 773.7 |
| 1 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 10,588 | 25.8 | 199.5 |
| 2 | Malignant neoplasms (C00-C97) | 9,143 | 22.3 | 172.3 |
| 3 | Chronic lower respiratory diseases (J40-J47) | 2,499 | 6.1 | 47.1 |
| 4 | Cerebrovascular diseases (I60-I69) | 2,480 | 6.0 | 46.7 |
| 5 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 2,475 | 6.0 | 46.6 |
| 6 | Influenza and pneumonia (J10-J18) | 1,133 | 2.8 | 21.3 |
| 7 | Alzheimer's disease (G30) | 1,108 | 2.7 | 20.9 |
| 8 | Diabetes mellitus (E10-E14) | 1,058 | 2.6 | 19.9 |
| 9 | Intentional self-harm (suicide) (*U03,X60-X84,Y87.0) | 767 | 1.9 | 14.5 |
| 10 | Chronic liver disease and cirrhosis (K70,K73-K74) | 661 | 1.6 | 12.5 |
| 11 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 644 | 1.6 | 12.1 |
| 12 | Assault (homicide) (*U01-*U02,X85-Y09,Y87.1) | 494 | 1.2 | 9.3 |
| 13 | Septicemia (A40-A41) | 383 | 0.9 | 7.2 |
| 14 | Parkinson's disease (G20-G21) | 370 | 0.9 | 7.0 |
| 15 | Essential (primary) hypertension and hypertensive renal disease (I10,I12) | 304 | 0.7 | 5.7 |
| ... | All other causes (Residual) | 6,951 | 16.9 | 131.0 |

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

... Category not applicable

LCWK9. Deaths, percent of total deaths, and death rates for the 15 leading causes of death: United States and each State, 2001

[Rates per 100,000 population.]

| Rank ¹ | Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992) and State | Number | Percent of total deaths | Rate |
|-------------------|--|---------|-------------------------|---------|
| Arkansas | | | | |
| ... | All causes | 27,759 | 100.0 | 1,030.1 |
| 1 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 8,263 | 29.8 | 306.6 |
| 2 | Malignant neoplasms (C00-C97) | 6,092 | 21.9 | 226.1 |
| 3 | Cerebrovascular diseases (I60-I69) | 2,256 | 8.1 | 83.7 |
| 4 | Chronic lower respiratory diseases (J40-J47) | 1,368 | 4.9 | 50.8 |
| 5 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 1,277 | 4.6 | 47.4 |
| 6 | Diabetes mellitus (E10-E14) | 756 | 2.7 | 28.1 |
| 7 | Influenza and pneumonia (J10-J18) | 735 | 2.6 | 27.3 |
| 8 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 521 | 1.9 | 19.3 |
| 9 | Septicemia (A40-A41) | 499 | 1.8 | 18.5 |
| 10 | Alzheimer's disease (G30) | 452 | 1.6 | 16.8 |
| 11 | Intentional self-harm (suicide) (*U03,X60-X84,Y87.0) | 382 | 1.4 | 14.2 |
| 12 | Chronic liver disease and cirrhosis (K70,K73-K74) | 233 | 0.8 | 8.6 |
| 13 | Assault (homicide) (*U01-*U02,X85-Y09,Y87.1) | 183 | 0.7 | 6.8 |
| 14 | Aortic aneurysm and dissection (I71) | 175 | 0.6 | 6.5 |
| 15 | Parkinson's disease (G20-G21) | 174 | 0.6 | 6.5 |
| ... | All other causes (Residual) | 4,393 | 15.8 | 163.0 |
| California | | | | |
| ... | All causes | 234,044 | 100.0 | 676.4 |
| 1 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 68,234 | 29.2 | 197.2 |
| 2 | Malignant neoplasms (C00-C97) | 53,924 | 23.0 | 155.8 |
| 3 | Cerebrovascular diseases (I60-I69) | 18,088 | 7.7 | 52.3 |
| 4 | Chronic lower respiratory diseases (J40-J47) | 12,965 | 5.5 | 37.5 |
| 5 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 8,132 | 3.5 | 23.5 |
| 6 | Influenza and pneumonia (J10-J18) | 8,129 | 3.5 | 23.5 |
| 7 | Diabetes mellitus (E10-E14) | 6,395 | 2.7 | 18.5 |
| 8 | Alzheimer's disease (G30) | 4,935 | 2.1 | 14.3 |
| 9 | Chronic liver disease and cirrhosis (K70,K73-K74) | 3,692 | 1.6 | 10.7 |
| 10 | Intentional self-harm (suicide) (*U03,X60-X84,Y87.0) | 2,831 | 1.2 | 8.2 |
| 11 | Essential (primary) hypertension and hypertensive renal disease (I10,I12) | 2,339 | 1.0 | 6.8 |
| 12 | Assault (homicide) (*U01-*U02,X85-Y09,Y87.1) | 2,223 | 0.9 | 6.4 |
| 13 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 1,994 | 0.9 | 5.8 |
| 14 | Atherosclerosis (I70) | 1,601 | 0.7 | 4.6 |
| 15 | Parkinson's disease (G20-G21) | 1,512 | 0.6 | 4.4 |
| ... | All other causes (Residual) | 37,050 | 15.8 | 107.1 |

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

... Category not applicable

LCWK9. Deaths, percent of total deaths, and death rates for the 15 leading causes of death: United States and each State, 2001

[Rates per 100,000 population.]

| Rank ¹ | Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992) and State | Number | Percent of total deaths | Rate |
|-------------------|--|--------|-------------------------|-------|
| Colorado | | | | |
| ... | All causes | 28,294 | 100.0 | 638.5 |
| 1 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 6,293 | 22.2 | 142.0 |
| 2 | Malignant neoplasms (C00-C97) | 6,145 | 21.7 | 138.7 |
| 3 | Chronic lower respiratory diseases (J40-J47) | 1,836 | 6.5 | 41.4 |
| 4 | Cerebrovascular diseases (I60-I69) | 1,825 | 6.5 | 41.2 |
| 5 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 1,720 | 6.1 | 38.8 |
| 6 | Alzheimer's disease (G30) | 852 | 3.0 | 19.2 |
| 7 | Intentional self-harm (suicide) (*U03,X60-X84,Y87.0) | 722 | 2.6 | 16.3 |
| 8 | Diabetes mellitus (E10-E14) | 667 | 2.4 | 15.1 |
| 9 | Influenza and pneumonia (J10-J18) | 650 | 2.3 | 14.7 |
| 10 | Atherosclerosis (I70) | 542 | 1.9 | 12.2 |
| 11 | Chronic liver disease and cirrhosis (K70,K73-K74) | 420 | 1.5 | 9.5 |
| 12 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 391 | 1.4 | 8.8 |
| 13 | Septicemia (A40-A41) | 297 | 1.0 | 6.7 |
| 14 | Pneumonitis due to solids and liquids (J69) | 241 | 0.9 | 5.4 |
| 15 | Aortic aneurysm and dissection (I71) | 220 | 0.8 | 5.0 |
| 15 | Certain conditions originating in the perinatal period (P00-P96) | 220 | 0.8 | 5.0 |
| ... | All other causes (Residual) | 5,253 | 18.6 | 118.6 |
| Connecticut | | | | |
| ... | All causes | 29,827 | 100.0 | 868.4 |
| 1 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 8,582 | 28.8 | 249.9 |
| 2 | Malignant neoplasms (C00-C97) | 7,093 | 23.8 | 206.5 |
| 3 | Cerebrovascular diseases (I60-I69) | 2,003 | 6.7 | 58.3 |
| 4 | Chronic lower respiratory diseases (J40-J47) | 1,486 | 5.0 | 43.3 |
| 5 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 1,056 | 3.5 | 30.7 |
| 6 | Influenza and pneumonia (J10-J18) | 854 | 2.9 | 24.9 |
| 7 | Diabetes mellitus (E10-E14) | 759 | 2.5 | 22.1 |
| 8 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 585 | 2.0 | 17.0 |
| 9 | Alzheimer's disease (G30) | 573 | 1.9 | 16.7 |
| 10 | Septicemia (A40-A41) | 552 | 1.9 | 16.1 |
| 11 | Pneumonitis due to solids and liquids (J69) | 392 | 1.3 | 11.4 |
| 12 | Chronic liver disease and cirrhosis (K70,K73-K74) | 294 | 1.0 | 8.6 |
| 13 | Intentional self-harm (suicide) (*U03,X60-X84,Y87.0) | 283 | 0.9 | 8.2 |
| 14 | Essential (primary) hypertension and hypertensive renal disease (I10,I12) | 252 | 0.8 | 7.3 |
| 15 | Aortic aneurysm and dissection (I71) | 223 | 0.7 | 6.5 |
| ... | All other causes (Residual) | 4,840 | 16.2 | 140.9 |

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

... Category not applicable

LCWK9. Deaths, percent of total deaths, and death rates for the 15 leading causes of death: United States and each State, 2001

[Rates per 100,000 population.]

| Rank ¹ | Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992) and State | Number | Percent of total deaths | Rate |
|----------------------|--|--------|-------------------------|---------|
| Delaware | | | | |
| ... | All causes | 7,112 | 100.0 | 892.8 |
| 1 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 2,033 | 28.6 | 255.2 |
| 2 | Malignant neoplasms (C00-C97) | 1,786 | 25.1 | 224.2 |
| 3 | Cerebrovascular diseases (I60-I69) | 383 | 5.4 | 48.1 |
| 4 | Chronic lower respiratory diseases (J40-J47) | 306 | 4.3 | 38.4 |
| 5 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 290 | 4.1 | 36.4 |
| 6 | Diabetes mellitus (E10-E14) | 218 | 3.1 | 27.4 |
| 7 | Influenza and pneumonia (J10-J18) | 181 | 2.5 | 22.7 |
| 8 | Alzheimer's disease (G30) | 135 | 1.9 | 16.9 |
| 9 | Septicemia (A40-A41) | 128 | 1.8 | 16.1 |
| 10 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 108 | 1.5 | 13.6 |
| 10 | Intentional self-harm (suicide) (*U03,X60-X84,Y87.0) | 108 | 1.5 | 13.6 |
| 12 | Pneumonitis due to solids and liquids (J69) | 84 | 1.2 | 10.5 |
| 13 | Certain conditions originating in the perinatal period (P00-P96) | 82 | 1.2 | 10.3 |
| 14 | Chronic liver disease and cirrhosis (K70,K73-K74) | 75 | 1.1 | 9.4 |
| 15 | Human immunodeficiency virus (HIV) disease (B20-B24) | 67 | 0.9 | 8.4 |
| ... | All other causes (Residual) | 1,128 | 15.9 | 141.6 |
| District of Columbia | | | | |
| ... | All causes | 5,951 | 100.0 | 1,037.1 |
| 1 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 1,761 | 29.6 | 306.9 |
| 2 | Malignant neoplasms (C00-C97) | 1,327 | 22.3 | 231.3 |
| 3 | Cerebrovascular diseases (I60-I69) | 265 | 4.5 | 46.2 |
| 4 | Human immunodeficiency virus (HIV) disease (B20-B24) | 223 | 3.7 | 38.9 |
| 4 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 223 | 3.7 | 38.9 |
| 6 | Diabetes mellitus (E10-E14) | 209 | 3.5 | 36.4 |
| 7 | Assault (homicide) (*U01-*U02,X85-Y09,Y87.1) | 197 | 3.3 | 34.3 |
| 8 | Chronic lower respiratory diseases (J40-J47) | 150 | 2.5 | 26.1 |
| 9 | Septicemia (A40-A41) | 100 | 1.7 | 17.4 |
| 10 | Alzheimer's disease (G30) | 99 | 1.7 | 17.3 |
| 11 | Influenza and pneumonia (J10-J18) | 86 | 1.4 | 15.0 |
| 12 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 79 | 1.3 | 13.8 |
| 13 | Chronic liver disease and cirrhosis (K70,K73-K74) | 63 | 1.1 | 11.0 |
| 14 | Certain conditions originating in the perinatal period (P00-P96) | 49 | 0.8 | 8.5 |
| 15 | Essential (primary) hypertension and hypertensive renal disease (I10,I12) | 47 | 0.8 | 8.2 |
| ... | All other causes (Residual) | 1,073 | 18.0 | 187.0 |

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

... Category not applicable

LCWK9. Deaths, percent of total deaths, and death rates for the 15 leading causes of death: United States and each State, 2001

[Rates per 100,000 population.]

| Rank ¹ | Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992) and State | Number | Percent of total deaths | Rate |
|-------------------|--|---------|-------------------------|---------|
| Florida | | | | |
| ... | All causes | 167,269 | 100.0 | 1,021.6 |
| 1 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 50,629 | 30.3 | 309.2 |
| 2 | Malignant neoplasms (C00-C97) | 39,090 | 23.4 | 238.7 |
| 3 | Cerebrovascular diseases (I60-I69) | 10,414 | 6.2 | 63.6 |
| 4 | Chronic lower respiratory diseases (J40-J47) | 8,954 | 5.4 | 54.7 |
| 5 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 6,969 | 4.2 | 42.6 |
| 6 | Diabetes mellitus (E10-E14) | 4,631 | 2.8 | 28.3 |
| 7 | Alzheimer's disease (G30) | 3,643 | 2.2 | 22.2 |
| 8 | Influenza and pneumonia (J10-J18) | 3,317 | 2.0 | 20.3 |
| 9 | Intentional self-harm (suicide) (*U03,X60-X84,Y87.0) | 2,314 | 1.4 | 14.1 |
| 10 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 2,172 | 1.3 | 13.3 |
| 11 | Chronic liver disease and cirrhosis (K70,K73-K74) | 2,013 | 1.2 | 12.3 |
| 12 | Septicemia (A40-A41) | 1,729 | 1.0 | 10.6 |
| 13 | Human immunodeficiency virus (HIV) disease (B20-B24) | 1,664 | 1.0 | 10.2 |
| 14 | Parkinson's disease (G20-G21) | 1,322 | 0.8 | 8.1 |
| 15 | Essential (primary) hypertension and hypertensive renal disease (I10,I12) | 1,277 | 0.8 | 7.8 |
| ... | All other causes (Residual) | 27,131 | 16.2 | 165.7 |
| Georgia | | | | |
| ... | All causes | 64,485 | 100.0 | 767.2 |
| 1 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 17,478 | 27.1 | 207.9 |
| 2 | Malignant neoplasms (C00-C97) | 13,818 | 21.4 | 164.4 |
| 3 | Cerebrovascular diseases (I60-I69) | 4,312 | 6.7 | 51.3 |
| 4 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 3,396 | 5.3 | 40.4 |
| 5 | Chronic lower respiratory diseases (J40-J47) | 3,097 | 4.8 | 36.8 |
| 6 | Influenza and pneumonia (J10-J18) | 1,515 | 2.3 | 18.0 |
| 7 | Diabetes mellitus (E10-E14) | 1,475 | 2.3 | 17.5 |
| 8 | Alzheimer's disease (G30) | 1,419 | 2.2 | 16.9 |
| 9 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 1,362 | 2.1 | 16.2 |
| 10 | Septicemia (A40-A41) | 1,313 | 2.0 | 15.6 |
| 11 | Intentional self-harm (suicide) (*U03,X60-X84,Y87.0) | 935 | 1.4 | 11.1 |
| 12 | Human immunodeficiency virus (HIV) disease (B20-B24) | 722 | 1.1 | 8.6 |
| 13 | Assault (homicide) (*U01-*U02,X85-Y09,Y87.1) | 681 | 1.1 | 8.1 |
| 14 | Essential (primary) hypertension and hypertensive renal disease (I10,I12) | 660 | 1.0 | 7.9 |
| 15 | Chronic liver disease and cirrhosis (K70,K73-K74) | 654 | 1.0 | 7.8 |
| ... | All other causes (Residual) | 11,648 | 18.1 | 138.6 |

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

... Category not applicable

LCWK9. Deaths, percent of total deaths, and death rates for the 15 leading causes of death: United States and each State, 2001

[Rates per 100,000 population.]

| Rank ¹ | Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992) and State | Number | Percent of total deaths | Rate |
|-------------------|--|--------|-------------------------|-------|
| Hawaii | | | | |
| ... | All causes | 8,394 | 100.0 | 684.1 |
| 1 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 2,310 | 27.5 | 188.3 |
| 2 | Malignant neoplasms (C00-C97) | 2,024 | 24.1 | 165.0 |
| 3 | Cerebrovascular diseases (I60-I69) | 766 | 9.1 | 62.4 |
| 4 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 372 | 4.4 | 30.3 |
| 5 | Chronic lower respiratory diseases (J40-J47) | 280 | 3.3 | 22.8 |
| 6 | Influenza and pneumonia (J10-J18) | 207 | 2.5 | 16.9 |
| 7 | Diabetes mellitus (E10-E14) | 173 | 2.1 | 14.1 |
| 8 | Septicemia (A40-A41) | 164 | 2.0 | 13.4 |
| 9 | Intentional self-harm (suicide) (*U03,X60-X84,Y87.0) | 136 | 1.6 | 11.1 |
| 10 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 123 | 1.5 | 10.0 |
| 11 | Alzheimer's disease (G30) | 122 | 1.5 | 9.9 |
| 12 | Pneumonitis due to solids and liquids (J69) | 107 | 1.3 | 8.7 |
| 13 | Parkinson's disease (G20-G21) | 84 | 1.0 | 6.8 |
| 13 | Chronic liver disease and cirrhosis (K70,K73-K74) | 84 | 1.0 | 6.8 |
| 15 | Essential (primary) hypertension and hypertensive renal disease (I10,I12) | 73 | 0.9 | 5.9 |
| ... | All other causes (Residual) | 1,369 | 16.3 | 111.6 |
| Idaho | | | | |
| ... | All causes | 9,753 | 100.0 | 738.5 |
| 1 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 2,489 | 25.5 | 188.5 |
| 2 | Malignant neoplasms (C00-C97) | 2,093 | 21.5 | 158.5 |
| 3 | Cerebrovascular diseases (I60-I69) | 781 | 8.0 | 59.1 |
| 4 | Chronic lower respiratory diseases (J40-J47) | 581 | 6.0 | 44.0 |
| 5 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 568 | 5.8 | 43.0 |
| 6 | Diabetes mellitus (E10-E14) | 318 | 3.3 | 24.1 |
| 7 | Alzheimer's disease (G30) | 294 | 3.0 | 22.3 |
| 8 | Influenza and pneumonia (J10-J18) | 239 | 2.5 | 18.1 |
| 9 | Intentional self-harm (suicide) (*U03,X60-X84,Y87.0) | 210 | 2.2 | 15.9 |
| 10 | Chronic liver disease and cirrhosis (K70,K73-K74) | 105 | 1.1 | 8.0 |
| 11 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 95 | 1.0 | 7.2 |
| 12 | Parkinson's disease (G20-G21) | 91 | 0.9 | 6.9 |
| 13 | Pneumonitis due to solids and liquids (J69) | 76 | 0.8 | 5.8 |
| 14 | Aortic aneurysm and dissection (I71) | 66 | 0.7 | 5.0 |
| 15 | Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99) | 64 | 0.7 | 4.8 |
| ... | All other causes (Residual) | 1,683 | 17.3 | 127.4 |

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

... Category not applicable

LCWK9. Deaths, percent of total deaths, and death rates for the 15 leading causes of death: United States and each State, 2001

[Rates per 100,000 population.]

| Rank ¹ | Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992) and State | Number | Percent of total deaths | Rate |
|-------------------|--|---------|-------------------------|-------|
| Illinois | | | | |
| ... | All causes | 105,430 | 100.0 | 842.1 |
| 1 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 30,990 | 29.4 | 247.5 |
| 2 | Malignant neoplasms (C00-C97) | 24,778 | 23.5 | 197.9 |
| 3 | Cerebrovascular diseases (I60-I69) | 7,230 | 6.9 | 57.7 |
| 4 | Chronic lower respiratory diseases (J40-J47) | 4,776 | 4.5 | 38.1 |
| 5 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 4,077 | 3.9 | 32.6 |
| 6 | Diabetes mellitus (E10-E14) | 3,092 | 2.9 | 24.7 |
| 7 | Influenza and pneumonia (J10-J18) | 2,659 | 2.5 | 21.2 |
| 8 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 2,178 | 2.1 | 17.4 |
| 9 | Alzheimer's disease (G30) | 2,163 | 2.1 | 17.3 |
| 10 | Septicemia (A40-A41) | 1,968 | 1.9 | 15.7 |
| 11 | Intentional self-harm (suicide) (*U03,X60-X84,Y87.0) | 1,139 | 1.1 | 9.1 |
| 12 | Chronic liver disease and cirrhosis (K70,K73-K74) | 1,103 | 1.0 | 8.8 |
| 13 | Assault (homicide) (*U01-*U02,X85-Y09,Y87.1) | 1,087 | 1.0 | 8.7 |
| 14 | Pneumonitis due to solids and liquids (J69) | 848 | 0.8 | 6.8 |
| 15 | Essential (primary) hypertension and hypertensive renal disease (I10,I12) | 791 | 0.8 | 6.3 |
| ... | All other causes (Residual) | 16,551 | 15.7 | 132.2 |
| Indiana | | | | |
| ... | All causes | 55,198 | 100.0 | 900.9 |
| 1 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 15,682 | 28.4 | 256.0 |
| 2 | Malignant neoplasms (C00-C97) | 12,831 | 23.2 | 209.4 |
| 3 | Cerebrovascular diseases (I60-I69) | 3,877 | 7.0 | 63.3 |
| 4 | Chronic lower respiratory diseases (J40-J47) | 3,136 | 5.7 | 51.2 |
| 5 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 2,185 | 4.0 | 35.7 |
| 6 | Diabetes mellitus (E10-E14) | 1,677 | 3.0 | 27.4 |
| 7 | Alzheimer's disease (G30) | 1,231 | 2.2 | 20.1 |
| 8 | Influenza and pneumonia (J10-J18) | 1,184 | 2.1 | 19.3 |
| 9 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 1,149 | 2.1 | 18.8 |
| 10 | Septicemia (A40-A41) | 795 | 1.4 | 13.0 |
| 11 | Intentional self-harm (suicide) (*U03,X60-X84,Y87.0) | 715 | 1.3 | 11.7 |
| 12 | Chronic liver disease and cirrhosis (K70,K73-K74) | 459 | 0.8 | 7.5 |
| 13 | Assault (homicide) (*U01-*U02,X85-Y09,Y87.1) | 437 | 0.8 | 7.1 |
| 14 | Pneumonitis due to solids and liquids (J69) | 419 | 0.8 | 6.8 |
| 15 | Parkinson's disease (G20-G21) | 407 | 0.7 | 6.6 |
| ... | All other causes (Residual) | 9,014 | 16.3 | 147.1 |

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

... Category not applicable

LCWK9. Deaths, percent of total deaths, and death rates for the 15 leading causes of death: United States and each State, 2001

[Rates per 100,000 population.]

| Rank ¹ | Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992) and State | Number | Percent of total deaths | Rate |
|-------------------|--|--------|-------------------------|-------|
| Iowa | | | | |
| ... | All causes | 27,791 | 100.0 | 947.9 |
| 1 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 8,250 | 29.7 | 281.4 |
| 2 | Malignant neoplasms (C00-C97) | 6,404 | 23.0 | 218.4 |
| 3 | Cerebrovascular diseases (I60-I69) | 2,218 | 8.0 | 75.6 |
| 4 | Chronic lower respiratory diseases (J40-J47) | 1,552 | 5.6 | 52.9 |
| 5 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 1,051 | 3.8 | 35.8 |
| 6 | Influenza and pneumonia (J10-J18) | 877 | 3.2 | 29.9 |
| 7 | Alzheimer's disease (G30) | 863 | 3.1 | 29.4 |
| 8 | Diabetes mellitus (E10-E14) | 709 | 2.6 | 24.2 |
| 9 | Intentional self-harm (suicide) (*U03,X60-X84,Y87.0) | 304 | 1.1 | 10.4 |
| 10 | Septicemia (A40-A41) | 279 | 1.0 | 9.5 |
| 11 | Atherosclerosis (I70) | 272 | 1.0 | 9.3 |
| 12 | Aortic aneurysm and dissection (I71) | 227 | 0.8 | 7.7 |
| 13 | Pneumonitis due to solids and liquids (J69) | 225 | 0.8 | 7.7 |
| 14 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 220 | 0.8 | 7.5 |
| 15 | Parkinson's disease (G20-G21) | 218 | 0.8 | 7.4 |
| ... | All other causes (Residual) | 4,122 | 14.8 | 140.6 |
| Kansas | | | | |
| ... | All causes | 24,647 | 100.0 | 912.1 |
| 1 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 6,716 | 27.2 | 248.5 |
| 2 | Malignant neoplasms (C00-C97) | 5,448 | 22.1 | 201.6 |
| 3 | Cerebrovascular diseases (I60-I69) | 1,848 | 7.5 | 68.4 |
| 4 | Chronic lower respiratory diseases (J40-J47) | 1,444 | 5.9 | 53.4 |
| 5 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 1,156 | 4.7 | 42.8 |
| 6 | Diabetes mellitus (E10-E14) | 721 | 2.9 | 26.7 |
| 7 | Alzheimer's disease (G30) | 642 | 2.6 | 23.8 |
| 8 | Influenza and pneumonia (J10-J18) | 561 | 2.3 | 20.8 |
| 9 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 498 | 2.0 | 18.4 |
| 10 | Atherosclerosis (I70) | 305 | 1.2 | 11.3 |
| 11 | Intentional self-harm (suicide) (*U03,X60-X84,Y87.0) | 293 | 1.2 | 10.8 |
| 12 | Septicemia (A40-A41) | 283 | 1.1 | 10.5 |
| 13 | Parkinson's disease (G20-G21) | 214 | 0.9 | 7.9 |
| 14 | Chronic liver disease and cirrhosis (K70,K73-K74) | 208 | 0.8 | 7.7 |
| 15 | Pneumonitis due to solids and liquids (J69) | 184 | 0.7 | 6.8 |
| ... | All other causes (Residual) | 4,126 | 16.7 | 152.7 |

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

... Category not applicable

LCWK9. Deaths, percent of total deaths, and death rates for the 15 leading causes of death: United States and each State, 2001

[Rates per 100,000 population.]

| Rank ¹ | Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992) and State | Number | Percent of total deaths | Rate |
|-------------------|--|--------|-------------------------|-------|
| Kentucky | | | | |
| ... | All causes | 39,861 | 100.0 | 979.7 |
| 1 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 11,808 | 29.6 | 290.2 |
| 2 | Malignant neoplasms (C00-C97) | 9,358 | 23.5 | 230.0 |
| 3 | Cerebrovascular diseases (I60-I69) | 2,557 | 6.4 | 62.8 |
| 4 | Chronic lower respiratory diseases (J40-J47) | 2,265 | 5.7 | 55.7 |
| 5 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 1,990 | 5.0 | 48.9 |
| 6 | Influenza and pneumonia (J10-J18) | 1,114 | 2.8 | 27.4 |
| 7 | Diabetes mellitus (E10-E14) | 1,099 | 2.8 | 27.0 |
| 8 | Alzheimer's disease (G30) | 966 | 2.4 | 23.7 |
| 9 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 807 | 2.0 | 19.8 |
| 10 | Septicemia (A40-A41) | 606 | 1.5 | 14.9 |
| 11 | Intentional self-harm (suicide) (*U03,X60-X84,Y87.0) | 495 | 1.2 | 12.2 |
| 12 | Chronic liver disease and cirrhosis (K70,K73-K74) | 387 | 1.0 | 9.5 |
| 13 | Pneumonitis due to solids and liquids (J69) | 303 | 0.8 | 7.4 |
| 14 | Aortic aneurysm and dissection (I71) | 238 | 0.6 | 5.8 |
| 15 | In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48) | 237 | 0.6 | 5.8 |
| ... | All other causes (Residual) | 5,631 | 14.1 | 138.4 |
| Louisiana | | | | |
| ... | All causes | 41,757 | 100.0 | 934.1 |
| 1 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 11,474 | 27.5 | 256.7 |
| 2 | Malignant neoplasms (C00-C97) | 9,517 | 22.8 | 212.9 |
| 3 | Cerebrovascular diseases (I60-I69) | 2,638 | 6.3 | 59.0 |
| 4 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 2,029 | 4.9 | 45.4 |
| 5 | Chronic lower respiratory diseases (J40-J47) | 1,763 | 4.2 | 39.4 |
| 6 | Diabetes mellitus (E10-E14) | 1,734 | 4.2 | 38.8 |
| 7 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 993 | 2.4 | 22.2 |
| 8 | Alzheimer's disease (G30) | 957 | 2.3 | 21.4 |
| 9 | Influenza and pneumonia (J10-J18) | 941 | 2.3 | 21.0 |
| 10 | Septicemia (A40-A41) | 758 | 1.8 | 17.0 |
| 11 | Assault (homicide) (*U01-*U02,X85-Y09,Y87.1) | 541 | 1.3 | 12.1 |
| 12 | Intentional self-harm (suicide) (*U03,X60-X84,Y87.0) | 493 | 1.2 | 11.0 |
| 13 | Human immunodeficiency virus (HIV) disease (B20-B24) | 404 | 1.0 | 9.0 |
| 14 | Chronic liver disease and cirrhosis (K70,K73-K74) | 327 | 0.8 | 7.3 |
| 15 | Certain conditions originating in the perinatal period (P00-P96) | 326 | 0.8 | 7.3 |
| ... | All other causes (Residual) | 6,862 | 16.4 | 153.5 |

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

... Category not applicable

LCWK9. Deaths, percent of total deaths, and death rates for the 15 leading causes of death: United States and each State, 2001

[Rates per 100,000 population.]

| Rank ¹ | Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992) and State | Number | Percent of total deaths | Rate |
|-------------------|--|--------|-------------------------|-------|
| Maine | | | | |
| ... | All causes | 12,421 | 100.0 | 967.0 |
| 1 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 3,272 | 26.3 | 254.7 |
| 2 | Malignant neoplasms (C00-C97) | 3,045 | 24.5 | 237.1 |
| 3 | Cerebrovascular diseases (I60-I69) | 822 | 6.6 | 64.0 |
| 4 | Chronic lower respiratory diseases (J40-J47) | 800 | 6.4 | 62.3 |
| 5 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 490 | 3.9 | 38.1 |
| 6 | Alzheimer's disease (G30) | 471 | 3.8 | 36.7 |
| 7 | Diabetes mellitus (E10-E14) | 398 | 3.2 | 31.0 |
| 8 | Influenza and pneumonia (J10-J18) | 269 | 2.2 | 20.9 |
| 9 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 208 | 1.7 | 16.2 |
| 10 | Intentional self-harm (suicide) (*U03,X60-X84,Y87.0) | 161 | 1.3 | 12.5 |
| 11 | Septicemia (A40-A41) | 122 | 1.0 | 9.5 |
| 12 | Aortic aneurysm and dissection (I71) | 119 | 1.0 | 9.3 |
| 13 | Chronic liver disease and cirrhosis (K70,K73-K74) | 111 | 0.9 | 8.6 |
| 14 | Parkinson's disease (G20-G21) | 103 | 0.8 | 8.0 |
| 15 | Pneumonitis due to solids and liquids (J69) | 85 | 0.7 | 6.6 |
| ... | All other causes (Residual) | 1,945 | 15.7 | 151.4 |
| Maryland | | | | |
| ... | All causes | 43,839 | 100.0 | 813.9 |
| 1 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 12,310 | 28.1 | 228.6 |
| 2 | Malignant neoplasms (C00-C97) | 10,322 | 23.5 | 191.6 |
| 3 | Cerebrovascular diseases (I60-I69) | 2,882 | 6.6 | 53.5 |
| 4 | Chronic lower respiratory diseases (J40-J47) | 1,907 | 4.4 | 35.4 |
| 5 | Diabetes mellitus (E10-E14) | 1,458 | 3.3 | 27.1 |
| 6 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 1,339 | 3.1 | 24.9 |
| 7 | Septicemia (A40-A41) | 986 | 2.2 | 18.3 |
| 8 | Alzheimer's disease (G30) | 948 | 2.2 | 17.6 |
| 9 | Influenza and pneumonia (J10-J18) | 928 | 2.1 | 17.2 |
| 10 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 593 | 1.4 | 11.0 |
| 11 | Human immunodeficiency virus (HIV) disease (B20-B24) | 572 | 1.3 | 10.6 |
| 12 | Assault (homicide) (*U01-*U02,X85-Y09,Y87.1) | 555 | 1.3 | 10.3 |
| 13 | Chronic liver disease and cirrhosis (K70,K73-K74) | 481 | 1.1 | 8.9 |
| 14 | Intentional self-harm (suicide) (*U03,X60-X84,Y87.0) | 454 | 1.0 | 8.4 |
| 15 | Pneumonitis due to solids and liquids (J69) | 418 | 1.0 | 7.8 |
| ... | All other causes (Residual) | 7,686 | 17.5 | 142.7 |

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

... Category not applicable

LCWK9. Deaths, percent of total deaths, and death rates for the 15 leading causes of death: United States and each State, 2001

[Rates per 100,000 population.]

| Rank ¹ | Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992) and State | Number | Percent of total deaths | Rate |
|-------------------|--|--------|-------------------------|-------|
| Massachusetts | | | | |
| ... | All causes | 56,754 | 100.0 | 886.6 |
| 1 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 15,144 | 26.7 | 236.6 |
| 2 | Malignant neoplasms (C00-C97) | 13,750 | 24.2 | 214.8 |
| 3 | Cerebrovascular diseases (I60-I69) | 3,535 | 6.2 | 55.2 |
| 4 | Chronic lower respiratory diseases (J40-J47) | 2,810 | 5.0 | 43.9 |
| 5 | Influenza and pneumonia (J10-J18) | 1,770 | 3.1 | 27.7 |
| 6 | Alzheimer's disease (G30) | 1,533 | 2.7 | 23.9 |
| 7 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 1,506 | 2.7 | 23.5 |
| 8 | Diabetes mellitus (E10-E14) | 1,422 | 2.5 | 22.2 |
| 9 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 1,207 | 2.1 | 18.9 |
| 10 | Septicemia (A40-A41) | 956 | 1.7 | 14.9 |
| 11 | Pneumonitis due to solids and liquids (J69) | 633 | 1.1 | 9.9 |
| 12 | Chronic liver disease and cirrhosis (K70,K73-K74) | 556 | 1.0 | 8.7 |
| 13 | Parkinson's disease (G20-G21) | 451 | 0.8 | 7.0 |
| 14 | Intentional self-harm (suicide) (*U03,X60-X84,Y87.0) | 426 | 0.8 | 6.7 |
| 15 | Aortic aneurysm and dissection (I71) | 399 | 0.7 | 6.2 |
| ... | All other causes (Residual) | 10,656 | 18.8 | 166.5 |
| Michigan | | | | |
| ... | All causes | 86,424 | 100.0 | 863.7 |
| 1 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 26,896 | 31.1 | 268.8 |
| 2 | Malignant neoplasms (C00-C97) | 19,689 | 22.8 | 196.8 |
| 3 | Cerebrovascular diseases (I60-I69) | 5,701 | 6.6 | 57.0 |
| 4 | Chronic lower respiratory diseases (J40-J47) | 4,148 | 4.8 | 41.5 |
| 5 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 3,291 | 3.8 | 32.9 |
| 6 | Diabetes mellitus (E10-E14) | 2,655 | 3.1 | 26.5 |
| 7 | Influenza and pneumonia (J10-J18) | 2,076 | 2.4 | 20.7 |
| 8 | Alzheimer's disease (G30) | 1,803 | 2.1 | 18.0 |
| 9 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 1,533 | 1.8 | 15.3 |
| 10 | Chronic liver disease and cirrhosis (K70,K73-K74) | 1,058 | 1.2 | 10.6 |
| 11 | Intentional self-harm (suicide) (*U03,X60-X84,Y87.0) | 1,051 | 1.2 | 10.5 |
| 12 | Septicemia (A40-A41) | 885 | 1.0 | 8.8 |
| 13 | Assault (homicide) (*U01-*U02,X85-Y09,Y87.1) | 689 | 0.8 | 6.9 |
| 14 | Essential (primary) hypertension and hypertensive renal disease (I10,I12) | 672 | 0.8 | 6.7 |
| 15 | Certain conditions originating in the perinatal period (P00-P96) | 622 | 0.7 | 6.2 |
| ... | All other causes (Residual) | 13,655 | 15.8 | 136.5 |

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

... Category not applicable

LCWK9. Deaths, percent of total deaths, and death rates for the 15 leading causes of death: United States and each State, 2001

[Rates per 100,000 population.]

| Rank ¹ | Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992) and State | Number | Percent of total deaths | Rate |
|-------------------|--|--------|-------------------------|-------|
| Minnesota | | | | |
| ... | All causes | 37,735 | 100.0 | 757.0 |
| 1 | Malignant neoplasms (C00-C97) | 8,967 | 23.8 | 179.9 |
| 2 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 8,760 | 23.2 | 175.7 |
| 3 | Cerebrovascular diseases (I60-I69) | 2,727 | 7.2 | 54.7 |
| 4 | Chronic lower respiratory diseases (J40-J47) | 1,909 | 5.1 | 38.3 |
| 5 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 1,797 | 4.8 | 36.1 |
| 6 | Diabetes mellitus (E10-E14) | 1,213 | 3.2 | 24.3 |
| 7 | Alzheimer's disease (G30) | 1,173 | 3.1 | 23.5 |
| 8 | Influenza and pneumonia (J10-J18) | 828 | 2.2 | 16.6 |
| 9 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 729 | 1.9 | 14.6 |
| 10 | Intentional self-harm (suicide) (*U03,X60-X84,Y87.0) | 480 | 1.3 | 9.6 |
| 11 | Essential (primary) hypertension and hypertensive renal disease (I10,I12) | 463 | 1.2 | 9.3 |
| 12 | Parkinson's disease (G20-G21) | 453 | 1.2 | 9.1 |
| 13 | Chronic liver disease and cirrhosis (K70,K73-K74) | 359 | 1.0 | 7.2 |
| 14 | Aortic aneurysm and dissection (I71) | 348 | 0.9 | 7.0 |
| 15 | In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48) | 276 | 0.7 | 5.5 |
| ... | All other causes (Residual) | 7,253 | 19.2 | 145.5 |
| Mississippi | | | | |
| ... | All causes | 28,259 | 100.0 | 988.2 |
| 1 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 9,050 | 32.0 | 316.5 |
| 2 | Malignant neoplasms (C00-C97) | 5,958 | 21.1 | 208.3 |
| 3 | Cerebrovascular diseases (I60-I69) | 1,935 | 6.8 | 67.7 |
| 4 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 1,571 | 5.6 | 54.9 |
| 5 | Chronic lower respiratory diseases (J40-J47) | 1,333 | 4.7 | 46.6 |
| 6 | Influenza and pneumonia (J10-J18) | 754 | 2.7 | 26.4 |
| 7 | Diabetes mellitus (E10-E14) | 661 | 2.3 | 23.1 |
| 8 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 602 | 2.1 | 21.1 |
| 9 | Septicemia (A40-A41) | 491 | 1.7 | 17.2 |
| 10 | Alzheimer's disease (G30) | 449 | 1.6 | 15.7 |
| 11 | Intentional self-harm (suicide) (*U03,X60-X84,Y87.0) | 328 | 1.2 | 11.5 |
| 12 | Assault (homicide) (*U01-*U02,X85-Y09,Y87.1) | 326 | 1.2 | 11.4 |
| 13 | Essential (primary) hypertension and hypertensive renal disease (I10,I12) | 287 | 1.0 | 10.0 |
| 14 | Chronic liver disease and cirrhosis (K70,K73-K74) | 265 | 0.9 | 9.3 |
| 15 | Certain conditions originating in the perinatal period (P00-P96) | 212 | 0.8 | 7.4 |
| ... | All other causes (Residual) | 4,037 | 14.3 | 141.2 |

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

... Category not applicable

LCWK9. Deaths, percent of total deaths, and death rates for the 15 leading causes of death: United States and each State, 2001

[Rates per 100,000 population.]

| Rank ¹ | Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992) and State | Number | Percent of total deaths | Rate |
|-------------------|--|--------|-------------------------|-------|
| Missouri | | | | |
| ... | All causes | 54,982 | 100.0 | 975.3 |
| 1 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 16,633 | 30.3 | 295.1 |
| 2 | Malignant neoplasms (C00-C97) | 12,365 | 22.5 | 219.3 |
| 3 | Cerebrovascular diseases (I60-I69) | 3,796 | 6.9 | 67.3 |
| 4 | Chronic lower respiratory diseases (J40-J47) | 2,882 | 5.2 | 51.1 |
| 5 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 2,455 | 4.5 | 43.5 |
| 6 | Influenza and pneumonia (J10-J18) | 1,604 | 2.9 | 28.5 |
| 7 | Diabetes mellitus (E10-E14) | 1,535 | 2.8 | 27.2 |
| 8 | Alzheimer's disease (G30) | 1,106 | 2.0 | 19.6 |
| 9 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 991 | 1.8 | 17.6 |
| 10 | Intentional self-harm (suicide) (*U03,X60-X84,Y87.0) | 725 | 1.3 | 12.9 |
| 11 | Septicemia (A40-A41) | 602 | 1.1 | 10.7 |
| 12 | Chronic liver disease and cirrhosis (K70,K73-K74) | 439 | 0.8 | 7.8 |
| 13 | Assault (homicide) (*U01-*U02,X85-Y09,Y87.1) | 437 | 0.8 | 7.8 |
| 14 | Pneumonitis due to solids and liquids (J69) | 433 | 0.8 | 7.7 |
| 15 | Parkinson's disease (G20-G21) | 371 | 0.7 | 6.6 |
| ... | All other causes (Residual) | 8,608 | 15.7 | 152.7 |
| Montana | | | | |
| ... | All causes | 8,265 | 100.0 | 912.9 |
| 1 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 1,970 | 23.8 | 217.6 |
| 2 | Malignant neoplasms (C00-C97) | 1,956 | 23.7 | 216.0 |
| 3 | Chronic lower respiratory diseases (J40-J47) | 583 | 7.1 | 64.4 |
| 4 | Cerebrovascular diseases (I60-I69) | 578 | 7.0 | 63.8 |
| 5 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 468 | 5.7 | 51.7 |
| 6 | Alzheimer's disease (G30) | 249 | 3.0 | 27.5 |
| 7 | Diabetes mellitus (E10-E14) | 229 | 2.8 | 25.3 |
| 8 | Influenza and pneumonia (J10-J18) | 193 | 2.3 | 21.3 |
| 9 | Intentional self-harm (suicide) (*U03,X60-X84,Y87.0) | 175 | 2.1 | 19.3 |
| 10 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 128 | 1.5 | 14.1 |
| 11 | Chronic liver disease and cirrhosis (K70,K73-K74) | 104 | 1.3 | 11.5 |
| 12 | Septicemia (A40-A41) | 71 | 0.9 | 7.8 |
| 13 | Parkinson's disease (G20-G21) | 69 | 0.8 | 7.6 |
| 14 | Pneumonitis due to solids and liquids (J69) | 62 | 0.8 | 6.8 |
| 15 | In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48) | 54 | 0.7 | 6.0 |
| ... | All other causes (Residual) | 1,376 | 16.6 | 152.0 |

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

... Category not applicable

LCWK9. Deaths, percent of total deaths, and death rates for the 15 leading causes of death: United States and each State, 2001

[Rates per 100,000 population.]

| Rank ¹ | Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992) and State | Number | Percent of total deaths | Rate |
|-------------------|--|--------|-------------------------|-------|
| Nebraska | | | | |
| ... | All causes | 15,174 | 100.0 | 882.2 |
| 1 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 4,150 | 27.3 | 241.3 |
| 2 | Malignant neoplasms (C00-C97) | 3,389 | 22.3 | 197.0 |
| 3 | Cerebrovascular diseases (I60-I69) | 1,130 | 7.4 | 65.7 |
| 4 | Chronic lower respiratory diseases (J40-J47) | 882 | 5.8 | 51.3 |
| 5 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 633 | 4.2 | 36.8 |
| 6 | Alzheimer's disease (G30) | 430 | 2.8 | 25.0 |
| 7 | Diabetes mellitus (E10-E14) | 400 | 2.6 | 23.3 |
| 8 | Influenza and pneumonia (J10-J18) | 296 | 2.0 | 17.2 |
| 9 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 289 | 1.9 | 16.8 |
| 10 | Intentional self-harm (suicide) (*U03,X60-X84,Y87.0) | 187 | 1.2 | 10.9 |
| 11 | Atherosclerosis (I70) | 148 | 1.0 | 8.6 |
| 12 | Essential (primary) hypertension and hypertensive renal disease (I10,I12) | 140 | 0.9 | 8.1 |
| 13 | Parkinson's disease (G20-G21) | 137 | 0.9 | 8.0 |
| 14 | Septicemia (A40-A41) | 132 | 0.9 | 7.7 |
| 15 | Pneumonitis due to solids and liquids (J69) | 126 | 0.8 | 7.3 |
| ... | All other causes (Residual) | 2,705 | 17.8 | 157.3 |
| Nevada | | | | |
| ... | All causes | 16,285 | 100.0 | 776.3 |
| 1 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 4,393 | 27.0 | 209.4 |
| 2 | Malignant neoplasms (C00-C97) | 3,815 | 23.4 | 181.9 |
| 3 | Chronic lower respiratory diseases (J40-J47) | 1,136 | 7.0 | 54.2 |
| 4 | Cerebrovascular diseases (I60-I69) | 913 | 5.6 | 43.5 |
| 5 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 739 | 4.5 | 35.2 |
| 6 | Septicemia (A40-A41) | 391 | 2.4 | 18.6 |
| 7 | Intentional self-harm (suicide) (*U03,X60-X84,Y87.0) | 387 | 2.4 | 18.4 |
| 8 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 376 | 2.3 | 17.9 |
| 9 | Influenza and pneumonia (J10-J18) | 352 | 2.2 | 16.8 |
| 10 | Diabetes mellitus (E10-E14) | 322 | 2.0 | 15.3 |
| 11 | Chronic liver disease and cirrhosis (K70,K73-K74) | 297 | 1.8 | 14.2 |
| 12 | Alzheimer's disease (G30) | 211 | 1.3 | 10.1 |
| 13 | Assault (homicide) (*U01-*U02,X85-Y09,Y87.1) | 175 | 1.1 | 8.3 |
| 14 | Essential (primary) hypertension and hypertensive renal disease (I10,I12) | 125 | 0.8 | 6.0 |
| 14 | Aortic aneurysm and dissection (I71) | 125 | 0.8 | 6.0 |
| ... | All other causes (Residual) | 2,528 | 15.5 | 120.5 |

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

... Category not applicable

LCWK9. Deaths, percent of total deaths, and death rates for the 15 leading causes of death: United States and each State, 2001

[Rates per 100,000 population.]

| Rank ¹ | Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992) and State | Number | Percent of total deaths | Rate |
|-------------------|--|--------|-------------------------|-------|
| New Hampshire | | | | |
| ... | All causes | 9,815 | 100.0 | 779.4 |
| 1 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 2,835 | 28.9 | 225.1 |
| 2 | Malignant neoplasms (C00-C97) | 2,398 | 24.4 | 190.4 |
| 3 | Cerebrovascular diseases (I60-I69) | 633 | 6.4 | 50.3 |
| 4 | Chronic lower respiratory diseases (J40-J47) | 616 | 6.3 | 48.9 |
| 5 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 374 | 3.8 | 29.7 |
| 6 | Alzheimer's disease (G30) | 305 | 3.1 | 24.2 |
| 7 | Diabetes mellitus (E10-E14) | 291 | 3.0 | 23.1 |
| 8 | Influenza and pneumonia (J10-J18) | 216 | 2.2 | 17.2 |
| 9 | Intentional self-harm (suicide) (*U03,X60-X84,Y87.0) | 167 | 1.7 | 13.3 |
| 10 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 136 | 1.4 | 10.8 |
| 11 | Chronic liver disease and cirrhosis (K70,K73-K74) | 121 | 1.2 | 9.6 |
| 12 | Aortic aneurysm and dissection (I71) | 90 | 0.9 | 7.1 |
| 13 | Parkinson's disease (G20-G21) | 79 | 0.8 | 6.3 |
| 14 | Atherosclerosis (I70) | 72 | 0.7 | 5.7 |
| 15 | Essential (primary) hypertension and hypertensive renal disease (I10,I12) | 67 | 0.7 | 5.3 |
| ... | All other causes (Residual) | 1,415 | 14.4 | 112.4 |
| New Jersey | | | | |
| ... | All causes | 74,710 | 100.0 | 877.8 |
| 1 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 22,704 | 30.4 | 266.8 |
| 2 | Malignant neoplasms (C00-C97) | 18,165 | 24.3 | 213.4 |
| 3 | Cerebrovascular diseases (I60-I69) | 4,007 | 5.4 | 47.1 |
| 4 | Chronic lower respiratory diseases (J40-J47) | 2,911 | 3.9 | 34.2 |
| 5 | Diabetes mellitus (E10-E14) | 2,556 | 3.4 | 30.0 |
| 6 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 2,405 | 3.2 | 28.3 |
| 7 | Influenza and pneumonia (J10-J18) | 1,893 | 2.5 | 22.2 |
| 8 | Septicemia (A40-A41) | 1,830 | 2.4 | 21.5 |
| 9 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 1,495 | 2.0 | 17.6 |
| 10 | Alzheimer's disease (G30) | 1,376 | 1.8 | 16.2 |
| 11 | Assault (homicide) (*U01-*U02,X85-Y09,Y87.1) | 1,051 | 1.4 | 12.3 |
| 12 | Human immunodeficiency virus (HIV) disease (B20-B24) | 785 | 1.1 | 9.2 |
| 13 | Chronic liver disease and cirrhosis (K70,K73-K74) | 778 | 1.0 | 9.1 |
| 14 | Intentional self-harm (suicide) (*U03,X60-X84,Y87.0) | 588 | 0.8 | 6.9 |
| 15 | Pneumonitis due to solids and liquids (J69) | 569 | 0.8 | 6.7 |
| ... | All other causes (Residual) | 11,597 | 15.5 | 136.3 |

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

... Category not applicable

LCWK9. Deaths, percent of total deaths, and death rates for the 15 leading causes of death: United States and each State, 2001

[Rates per 100,000 population.]

| Rank ¹ | Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992) and State | Number | Percent of total deaths | Rate |
|-------------------|--|---------|-------------------------|-------|
| New Mexico | | | | |
| ... | All causes | 14,129 | 100.0 | 771.7 |
| 1 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 3,423 | 24.2 | 187.0 |
| 2 | Malignant neoplasms (C00-C97) | 2,900 | 20.5 | 158.4 |
| 3 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 1,020 | 7.2 | 55.7 |
| 4 | Cerebrovascular diseases (I60-I69) | 824 | 5.8 | 45.0 |
| 5 | Chronic lower respiratory diseases (J40-J47) | 775 | 5.5 | 42.3 |
| 6 | Diabetes mellitus (E10-E14) | 538 | 3.8 | 29.4 |
| 7 | Influenza and pneumonia (J10-J18) | 398 | 2.8 | 21.7 |
| 8 | Intentional self-harm (suicide) (*U03,X60-X84,Y87.0) | 362 | 2.6 | 19.8 |
| 9 | Chronic liver disease and cirrhosis (K70,K73-K74) | 302 | 2.1 | 16.5 |
| 10 | Alzheimer's disease (G30) | 297 | 2.1 | 16.2 |
| 11 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 172 | 1.2 | 9.4 |
| 12 | Assault (homicide) (*U01-*U02,X85-Y09,Y87.1) | 130 | 0.9 | 7.1 |
| 13 | Parkinson's disease (G20-G21) | 126 | 0.9 | 6.9 |
| 14 | Septicemia (A40-A41) | 110 | 0.8 | 6.0 |
| 15 | In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48) | 101 | 0.7 | 5.5 |
| ... | All other causes (Residual) | 2,651 | 18.8 | 144.8 |
| New York | | | | |
| ... | All causes | 159,240 | 100.0 | 834.4 |
| 1 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 56,643 | 35.6 | 296.8 |
| 2 | Malignant neoplasms (C00-C97) | 36,975 | 23.2 | 193.7 |
| 3 | Cerebrovascular diseases (I60-I69) | 7,706 | 4.8 | 40.4 |
| 4 | Chronic lower respiratory diseases (J40-J47) | 6,906 | 4.3 | 36.2 |
| 5 | Influenza and pneumonia (J10-J18) | 5,085 | 3.2 | 26.6 |
| 6 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 4,982 | 3.1 | 26.1 |
| 7 | Diabetes mellitus (E10-E14) | 3,844 | 2.4 | 20.1 |
| 8 | Assault (homicide) (*U01-*U02,X85-Y09,Y87.1) | 2,818 | 1.8 | 14.8 |
| 9 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 2,421 | 1.5 | 12.7 |
| 10 | Septicemia (A40-A41) | 2,308 | 1.4 | 12.1 |
| 11 | Human immunodeficiency virus (HIV) disease (B20-B24) | 2,085 | 1.3 | 10.9 |
| 12 | Alzheimer's disease (G30) | 1,705 | 1.1 | 8.9 |
| 13 | Essential (primary) hypertension and hypertensive renal disease (I10,I12) | 1,517 | 1.0 | 7.9 |
| 14 | Chronic liver disease and cirrhosis (K70,K73-K74) | 1,488 | 0.9 | 7.8 |
| 15 | Intentional self-harm (suicide) (*U03,X60-X84,Y87.0) | 1,253 | 0.8 | 6.6 |
| ... | All other causes (Residual) | 21,504 | 13.5 | 112.7 |

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

... Category not applicable

LCWK9. Deaths, percent of total deaths, and death rates for the 15 leading causes of death: United States and each State, 2001

[Rates per 100,000 population.]

| Rank ¹ | Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992) and State | Number | Percent of total deaths | Rate |
|-------------------|--|--------|-------------------------|-------|
| North Carolina | | | | |
| ... | All causes | 70,934 | 100.0 | 864.4 |
| 1 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 18,792 | 26.5 | 229.0 |
| 2 | Malignant neoplasms (C00-C97) | 16,065 | 22.6 | 195.8 |
| 3 | Cerebrovascular diseases (I60-I69) | 5,401 | 7.6 | 65.8 |
| 4 | Chronic lower respiratory diseases (J40-J47) | 3,514 | 5.0 | 42.8 |
| 5 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 3,438 | 4.8 | 41.9 |
| 6 | Diabetes mellitus (E10-E14) | 2,181 | 3.1 | 26.6 |
| 7 | Alzheimer's disease (G30) | 1,786 | 2.5 | 21.8 |
| 8 | Influenza and pneumonia (J10-J18) | 1,780 | 2.5 | 21.7 |
| 9 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 1,332 | 1.9 | 16.2 |
| 10 | Septicemia (A40-A41) | 1,075 | 1.5 | 13.1 |
| 11 | Intentional self-harm (suicide) (*U03,X60-X84,Y87.0) | 997 | 1.4 | 12.1 |
| 12 | Chronic liver disease and cirrhosis (K70,K73-K74) | 725 | 1.0 | 8.8 |
| 13 | Pneumonitis due to solids and liquids (J69) | 663 | 0.9 | 8.1 |
| 14 | Essential (primary) hypertension and hypertensive renal disease (I10,I12) | 627 | 0.9 | 7.6 |
| 15 | Assault (homicide) (*U01-*U02,X85-Y09,Y87.1) | 599 | 0.8 | 7.3 |
| ... | All other causes (Residual) | 11,959 | 16.9 | 145.7 |
| North Dakota | | | | |
| ... | All causes | 6,048 | 100.0 | 950.1 |
| 1 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 1,700 | 28.1 | 267.1 |
| 2 | Malignant neoplasms (C00-C97) | 1,391 | 23.0 | 218.5 |
| 3 | Cerebrovascular diseases (I60-I69) | 507 | 8.4 | 79.6 |
| 4 | Chronic lower respiratory diseases (J40-J47) | 308 | 5.1 | 48.4 |
| 5 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 238 | 3.9 | 37.4 |
| 6 | Alzheimer's disease (G30) | 224 | 3.7 | 35.2 |
| 7 | Diabetes mellitus (E10-E14) | 196 | 3.2 | 30.8 |
| 8 | Influenza and pneumonia (J10-J18) | 162 | 2.7 | 25.4 |
| 9 | Intentional self-harm (suicide) (*U03,X60-X84,Y87.0) | 79 | 1.3 | 12.4 |
| 10 | Essential (primary) hypertension and hypertensive renal disease (I10,I12) | 67 | 1.1 | 10.5 |
| 11 | Chronic liver disease and cirrhosis (K70,K73-K74) | 62 | 1.0 | 9.7 |
| 12 | Atherosclerosis (I70) | 56 | 0.9 | 8.8 |
| 13 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 50 | 0.8 | 7.9 |
| 14 | In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48) | 48 | 0.8 | 7.5 |
| 15 | Parkinson's disease (G20-G21) | 44 | 0.7 | 6.9 |
| ... | All other causes (Residual) | 916 | 15.1 | 143.9 |

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

... Category not applicable

LCWK9. Deaths, percent of total deaths, and death rates for the 15 leading causes of death: United States and each State, 2001

[Rates per 100,000 population.]

| Rank ¹ | Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992) and State | Number | Percent of total deaths | Rate |
|-------------------|--|---------|-------------------------|-------|
| Ohio | | | | |
| ... | All causes | 108,027 | 100.0 | 948.5 |
| 1 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 32,453 | 30.0 | 284.9 |
| 2 | Malignant neoplasms (C00-C97) | 24,804 | 23.0 | 217.8 |
| 3 | Cerebrovascular diseases (I60-I69) | 6,891 | 6.4 | 60.5 |
| 4 | Chronic lower respiratory diseases (J40-J47) | 5,881 | 5.4 | 51.6 |
| 5 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 3,857 | 3.6 | 33.9 |
| 6 | Diabetes mellitus (E10-E14) | 3,750 | 3.5 | 32.9 |
| 7 | Influenza and pneumonia (J10-J18) | 2,401 | 2.2 | 21.1 |
| 8 | Alzheimer's disease (G30) | 2,367 | 2.2 | 20.8 |
| 9 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 1,912 | 1.8 | 16.8 |
| 10 | Septicemia (A40-A41) | 1,452 | 1.3 | 12.7 |
| 11 | Intentional self-harm (suicide) (*U03,X60-X84,Y87.0) | 1,219 | 1.1 | 10.7 |
| 12 | Chronic liver disease and cirrhosis (K70,K73-K74) | 1,027 | 1.0 | 9.0 |
| 13 | Essential (primary) hypertension and hypertensive renal disease (I10,I12) | 903 | 0.8 | 7.9 |
| 14 | Pneumonitis due to solids and liquids (J69) | 693 | 0.6 | 6.1 |
| 15 | Parkinson's disease (G20-G21) | 679 | 0.6 | 6.0 |
| ... | All other causes (Residual) | 17,738 | 16.4 | 155.7 |
| Oklahoma | | | | |
| ... | All causes | 34,682 | 100.0 | 999.6 |
| 1 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 10,840 | 31.3 | 312.4 |
| 2 | Malignant neoplasms (C00-C97) | 7,411 | 21.4 | 213.6 |
| 3 | Cerebrovascular diseases (I60-I69) | 2,384 | 6.9 | 68.7 |
| 4 | Chronic lower respiratory diseases (J40-J47) | 1,922 | 5.5 | 55.4 |
| 5 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 1,700 | 4.9 | 49.0 |
| 6 | Diabetes mellitus (E10-E14) | 1,065 | 3.1 | 30.7 |
| 7 | Influenza and pneumonia (J10-J18) | 913 | 2.6 | 26.3 |
| 8 | Alzheimer's disease (G30) | 682 | 2.0 | 19.7 |
| 9 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 567 | 1.6 | 16.3 |
| 10 | Intentional self-harm (suicide) (*U03,X60-X84,Y87.0) | 515 | 1.5 | 14.8 |
| 11 | Septicemia (A40-A41) | 425 | 1.2 | 12.2 |
| 12 | Atherosclerosis (I70) | 372 | 1.1 | 10.7 |
| 13 | Chronic liver disease and cirrhosis (K70,K73-K74) | 363 | 1.0 | 10.5 |
| 14 | Pneumonitis due to solids and liquids (J69) | 252 | 0.7 | 7.3 |
| 15 | Essential (primary) hypertension and hypertensive renal disease (I10,I12) | 240 | 0.7 | 6.9 |
| ... | All other causes (Residual) | 5,031 | 14.5 | 145.0 |

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

... Category not applicable

LCWK9. Deaths, percent of total deaths, and death rates for the 15 leading causes of death: United States and each State, 2001

[Rates per 100,000 population.]

| Rank ¹ | Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992) and State | Number | Percent of total deaths | Rate |
|-------------------|--|---------|-------------------------|---------|
| Oregon | | | | |
| ... | All causes | 30,158 | 100.0 | 868.2 |
| 1 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 7,075 | 23.5 | 203.7 |
| 2 | Malignant neoplasms (C00-C97) | 7,057 | 23.4 | 203.2 |
| 3 | Cerebrovascular diseases (I60-I69) | 2,588 | 8.6 | 74.5 |
| 4 | Chronic lower respiratory diseases (J40-J47) | 1,726 | 5.7 | 49.7 |
| 5 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 1,313 | 4.4 | 37.8 |
| 6 | Alzheimer's disease (G30) | 1,030 | 3.4 | 29.7 |
| 7 | Diabetes mellitus (E10-E14) | 1,011 | 3.4 | 29.1 |
| 8 | Influenza and pneumonia (J10-J18) | 575 | 1.9 | 16.6 |
| 9 | Intentional self-harm (suicide) (*U03,X60-X84,Y87.0) | 505 | 1.7 | 14.5 |
| 10 | Chronic liver disease and cirrhosis (K70,K73-K74) | 339 | 1.1 | 9.8 |
| 11 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 331 | 1.1 | 9.5 |
| 12 | Essential (primary) hypertension and hypertensive renal disease (I10,I12) | 297 | 1.0 | 8.6 |
| 13 | Parkinson's disease (G20-G21) | 291 | 1.0 | 8.4 |
| 14 | Aortic aneurysm and dissection (I71) | 229 | 0.8 | 6.6 |
| 15 | Septicemia (A40-A41) | 207 | 0.7 | 6.0 |
| ... | All other causes (Residual) | 5,584 | 18.5 | 160.8 |
| Pennsylvania | | | | |
| ... | All causes | 129,729 | 100.0 | 1,054.4 |
| 1 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 39,438 | 30.4 | 320.6 |
| 2 | Malignant neoplasms (C00-C97) | 29,914 | 23.1 | 243.1 |
| 3 | Cerebrovascular diseases (I60-I69) | 8,619 | 6.6 | 70.1 |
| 4 | Chronic lower respiratory diseases (J40-J47) | 5,845 | 4.5 | 47.5 |
| 5 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 4,552 | 3.5 | 37.0 |
| 6 | Diabetes mellitus (E10-E14) | 3,826 | 2.9 | 31.1 |
| 7 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 2,838 | 2.2 | 23.1 |
| 8 | Alzheimer's disease (G30) | 2,753 | 2.1 | 22.4 |
| 9 | Influenza and pneumonia (J10-J18) | 2,735 | 2.1 | 22.2 |
| 10 | Septicemia (A40-A41) | 2,649 | 2.0 | 21.5 |
| 11 | Intentional self-harm (suicide) (*U03,X60-X84,Y87.0) | 1,276 | 1.0 | 10.4 |
| 12 | Pneumonitis due to solids and liquids (J69) | 1,259 | 1.0 | 10.2 |
| 13 | Chronic liver disease and cirrhosis (K70,K73-K74) | 1,134 | 0.9 | 9.2 |
| 14 | Parkinson's disease (G20-G21) | 1,022 | 0.8 | 8.3 |
| 15 | Aortic aneurysm and dissection (I71) | 869 | 0.7 | 7.1 |
| ... | All other causes (Residual) | 21,000 | 16.2 | 170.7 |

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

... Category not applicable

LCWK9. Deaths, percent of total deaths, and death rates for the 15 leading causes of death: United States and each State, 2001

[Rates per 100,000 population.]

| Rank ¹ | Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992) and State | Number | Percent of total deaths | Rate |
|-------------------|--|--------|-------------------------|-------|
| Rhode Island | | | | |
| ... | All causes | 10,021 | 100.0 | 945.7 |
| 1 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 3,076 | 30.7 | 290.3 |
| 2 | Malignant neoplasms (C00-C97) | 2,398 | 23.9 | 226.3 |
| 3 | Cerebrovascular diseases (I60-I69) | 614 | 6.1 | 57.9 |
| 4 | Chronic lower respiratory diseases (J40-J47) | 511 | 5.1 | 48.2 |
| 5 | Influenza and pneumonia (J10-J18) | 295 | 2.9 | 27.8 |
| 6 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 293 | 2.9 | 27.7 |
| 7 | Diabetes mellitus (E10-E14) | 265 | 2.6 | 25.0 |
| 8 | Alzheimer's disease (G30) | 251 | 2.5 | 23.7 |
| 9 | Septicemia (A40-A41) | 136 | 1.4 | 12.8 |
| 10 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 118 | 1.2 | 11.1 |
| 11 | Chronic liver disease and cirrhosis (K70,K73-K74) | 96 | 1.0 | 9.1 |
| 12 | Atherosclerosis (I70) | 91 | 0.9 | 8.6 |
| 13 | Intentional self-harm (suicide) (*U03,X60-X84,Y87.0) | 88 | 0.9 | 8.3 |
| 14 | Pneumonitis due to solids and liquids (J69) | 87 | 0.9 | 8.2 |
| 15 | Parkinson's disease (G20-G21) | 74 | 0.7 | 7.0 |
| ... | All other causes (Residual) | 1,628 | 16.2 | 153.6 |
| South Carolina | | | | |
| ... | All causes | 36,612 | 100.0 | 901.3 |
| 1 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 9,471 | 25.9 | 233.2 |
| 2 | Malignant neoplasms (C00-C97) | 8,272 | 22.6 | 203.6 |
| 3 | Cerebrovascular diseases (I60-I69) | 2,832 | 7.7 | 69.7 |
| 4 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 1,960 | 5.4 | 48.3 |
| 5 | Chronic lower respiratory diseases (J40-J47) | 1,731 | 4.7 | 42.6 |
| 6 | Diabetes mellitus (E10-E14) | 1,088 | 3.0 | 26.8 |
| 7 | Alzheimer's disease (G30) | 881 | 2.4 | 21.7 |
| 8 | Influenza and pneumonia (J10-J18) | 754 | 2.1 | 18.6 |
| 9 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 632 | 1.7 | 15.6 |
| 10 | Septicemia (A40-A41) | 543 | 1.5 | 13.4 |
| 11 | Intentional self-harm (suicide) (*U03,X60-X84,Y87.0) | 467 | 1.3 | 11.5 |
| 12 | Chronic liver disease and cirrhosis (K70,K73-K74) | 461 | 1.3 | 11.3 |
| 13 | Assault (homicide) (*U01-*U02,X85-Y09,Y87.1) | 350 | 1.0 | 8.6 |
| 14 | Essential (primary) hypertension and hypertensive renal disease (I10,I12) | 338 | 0.9 | 8.3 |
| 15 | Pneumonitis due to solids and liquids (J69) | 317 | 0.9 | 7.8 |
| ... | All other causes (Residual) | 6,515 | 17.8 | 160.4 |

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

... Category not applicable

LCWK9. Deaths, percent of total deaths, and death rates for the 15 leading causes of death: United States and each State, 2001

[Rates per 100,000 population.]

| Rank ¹ | Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992) and State | Number | Percent of total deaths | Rate |
|-------------------|--|--------|-------------------------|-------|
| South Dakota | | | | |
| ... | All causes | 6,923 | 100.0 | 912.9 |
| 1 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 1,985 | 28.7 | 261.8 |
| 2 | Malignant neoplasms (C00-C97) | 1,590 | 23.0 | 209.7 |
| 3 | Cerebrovascular diseases (I60-I69) | 490 | 7.1 | 64.6 |
| 4 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 382 | 5.5 | 50.4 |
| 5 | Chronic lower respiratory diseases (J40-J47) | 362 | 5.2 | 47.7 |
| 6 | Diabetes mellitus (E10-E14) | 212 | 3.1 | 28.0 |
| 7 | Influenza and pneumonia (J10-J18) | 188 | 2.7 | 24.8 |
| 8 | Alzheimer's disease (G30) | 156 | 2.3 | 20.6 |
| 9 | Intentional self-harm (suicide) (*U03,X60-X84,Y87.0) | 105 | 1.5 | 13.8 |
| 10 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 103 | 1.5 | 13.6 |
| 11 | Chronic liver disease and cirrhosis (K70,K73-K74) | 77 | 1.1 | 10.2 |
| 12 | Septicemia (A40-A41) | 70 | 1.0 | 9.2 |
| 13 | Aortic aneurysm and dissection (I71) | 56 | 0.8 | 7.4 |
| 14 | Parkinson's disease (G20-G21) | 55 | 0.8 | 7.3 |
| 15 | Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99) | 42 | 0.6 | 5.5 |
| ... | All other causes (Residual) | 1,050 | 15.2 | 138.5 |
| Tennessee | | | | |
| ... | All causes | 55,151 | 100.0 | 959.2 |
| 1 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 15,688 | 28.4 | 272.9 |
| 2 | Malignant neoplasms (C00-C97) | 12,229 | 22.2 | 212.7 |
| 3 | Cerebrovascular diseases (I60-I69) | 4,037 | 7.3 | 70.2 |
| 4 | Chronic lower respiratory diseases (J40-J47) | 2,941 | 5.3 | 51.2 |
| 5 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 2,709 | 4.9 | 47.1 |
| 6 | Diabetes mellitus (E10-E14) | 1,746 | 3.2 | 30.4 |
| 7 | Influenza and pneumonia (J10-J18) | 1,617 | 2.9 | 28.1 |
| 8 | Alzheimer's disease (G30) | 1,148 | 2.1 | 20.0 |
| 9 | Intentional self-harm (suicide) (*U03,X60-X84,Y87.0) | 711 | 1.3 | 12.4 |
| 10 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 624 | 1.1 | 10.9 |
| 11 | Chronic liver disease and cirrhosis (K70,K73-K74) | 613 | 1.1 | 10.7 |
| 12 | Essential (primary) hypertension and hypertensive renal disease (I10,I12) | 539 | 1.0 | 9.4 |
| 13 | Septicemia (A40-A41) | 495 | 0.9 | 8.6 |
| 14 | Assault (homicide) (*U01-*U02,X85-Y09,Y87.1) | 454 | 0.8 | 7.9 |
| 15 | Pneumonitis due to solids and liquids (J69) | 431 | 0.8 | 7.5 |
| ... | All other causes (Residual) | 9,169 | 16.6 | 159.5 |

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

... Category not applicable

LCWK9. Deaths, percent of total deaths, and death rates for the 15 leading causes of death: United States and each State, 2001

[Rates per 100,000 population.]

| Rank ¹ | Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992) and State | Number | Percent of total deaths | Rate |
|-------------------|--|---------|-------------------------|-------|
| Texas | | | | |
| ... | All causes | 152,779 | 100.0 | 714.9 |
| 1 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 43,199 | 28.3 | 202.1 |
| 2 | Malignant neoplasms (C00-C97) | 33,488 | 21.9 | 156.7 |
| 3 | Cerebrovascular diseases (I60-I69) | 10,612 | 6.9 | 49.7 |
| 4 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 7,920 | 5.2 | 37.1 |
| 5 | Chronic lower respiratory diseases (J40-J47) | 7,739 | 5.1 | 36.2 |
| 6 | Diabetes mellitus (E10-E14) | 5,456 | 3.6 | 25.5 |
| 7 | Influenza and pneumonia (J10-J18) | 3,608 | 2.4 | 16.9 |
| 8 | Alzheimer's disease (G30) | 3,439 | 2.3 | 16.1 |
| 9 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 2,269 | 1.5 | 10.6 |
| 10 | Intentional self-harm (suicide) (*U03,X60-X84,Y87.0) | 2,225 | 1.5 | 10.4 |
| 11 | Chronic liver disease and cirrhosis (K70,K73-K74) | 2,213 | 1.4 | 10.4 |
| 12 | Septicemia (A40-A41) | 2,003 | 1.3 | 9.4 |
| 13 | Assault (homicide) (*U01-*U02,X85-Y09,Y87.1) | 1,415 | 0.9 | 6.6 |
| 14 | Essential (primary) hypertension and hypertensive renal disease (I10,I12) | 1,173 | 0.8 | 5.5 |
| 15 | Human immunodeficiency virus (HIV) disease (B20-B24) | 1,055 | 0.7 | 4.9 |
| ... | All other causes (Residual) | 24,965 | 16.3 | 116.8 |
| Utah | | | | |
| ... | All causes | 12,662 | 100.0 | 555.7 |
| 1 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 2,896 | 22.9 | 127.1 |
| 2 | Malignant neoplasms (C00-C97) | 2,313 | 18.3 | 101.5 |
| 3 | Cerebrovascular diseases (I60-I69) | 870 | 6.9 | 38.2 |
| 4 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 643 | 5.1 | 28.2 |
| 5 | Chronic lower respiratory diseases (J40-J47) | 525 | 4.1 | 23.0 |
| 6 | Diabetes mellitus (E10-E14) | 509 | 4.0 | 22.3 |
| 7 | Influenza and pneumonia (J10-J18) | 411 | 3.2 | 18.0 |
| 8 | Intentional self-harm (suicide) (*U03,X60-X84,Y87.0) | 321 | 2.5 | 14.1 |
| 9 | Alzheimer's disease (G30) | 314 | 2.5 | 13.8 |
| 10 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 178 | 1.4 | 7.8 |
| 11 | Chronic liver disease and cirrhosis (K70,K73-K74) | 138 | 1.1 | 6.1 |
| 12 | Parkinson's disease (G20-G21) | 134 | 1.1 | 5.9 |
| 13 | Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99) | 120 | 0.9 | 5.3 |
| 14 | Septicemia (A40-A41) | 112 | 0.9 | 4.9 |
| 15 | Certain conditions originating in the perinatal period (P00-P96) | 97 | 0.8 | 4.3 |
| ... | All other causes (Residual) | 3,081 | 24.3 | 135.2 |

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

... Category not applicable

LCWK9. Deaths, percent of total deaths, and death rates for the 15 leading causes of death: United States and each State, 2001

[Rates per 100,000 population.]

| Rank ¹ | Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992) and State | Number | Percent of total deaths | Rate |
|-------------------|--|--------|-------------------------|-------|
| Vermont | | | | |
| ... | All causes | 5,201 | 100.0 | 848.5 |
| 1 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 1,429 | 27.5 | 233.1 |
| 2 | Malignant neoplasms (C00-C97) | 1,249 | 24.0 | 203.8 |
| 3 | Cerebrovascular diseases (I60-I69) | 323 | 6.2 | 52.7 |
| 4 | Chronic lower respiratory diseases (J40-J47) | 305 | 5.9 | 49.8 |
| 5 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 230 | 4.4 | 37.5 |
| 6 | Diabetes mellitus (E10-E14) | 155 | 3.0 | 25.3 |
| 7 | Alzheimer's disease (G30) | 149 | 2.9 | 24.3 |
| 8 | Influenza and pneumonia (J10-J18) | 133 | 2.6 | 21.7 |
| 9 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 86 | 1.7 | 14.0 |
| 10 | Intentional self-harm (suicide) (*U03,X60-X84,Y87.0) | 72 | 1.4 | 11.7 |
| 11 | Parkinson's disease (G20-G21) | 64 | 1.2 | 10.4 |
| 12 | Septicemia (A40-A41) | 49 | 0.9 | 8.0 |
| 13 | Aortic aneurysm and dissection (I71) | 48 | 0.9 | 7.8 |
| 14 | In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48) | 40 | 0.8 | 6.5 |
| 15 | Pneumonitis due to solids and liquids (J69) | 37 | 0.7 | 6.0 |
| 15 | Chronic liver disease and cirrhosis (K70,K73-K74) | 37 | 0.7 | 6.0 |
| ... | All other causes (Residual) | 795 | 15.3 | 129.7 |
| Virginia | | | | |
| ... | All causes | 56,280 | 100.0 | 782.0 |
| 1 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 14,913 | 26.5 | 207.2 |
| 2 | Malignant neoplasms (C00-C97) | 13,345 | 23.7 | 185.4 |
| 3 | Cerebrovascular diseases (I60-I69) | 4,129 | 7.3 | 57.4 |
| 4 | Chronic lower respiratory diseases (J40-J47) | 2,744 | 4.9 | 38.1 |
| 5 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 2,432 | 4.3 | 33.8 |
| 6 | Diabetes mellitus (E10-E14) | 1,613 | 2.9 | 22.4 |
| 7 | Influenza and pneumonia (J10-J18) | 1,481 | 2.6 | 20.6 |
| 8 | Alzheimer's disease (G30) | 1,184 | 2.1 | 16.5 |
| 9 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 1,128 | 2.0 | 15.7 |
| 10 | Septicemia (A40-A41) | 1,098 | 2.0 | 15.3 |
| 11 | Intentional self-harm (suicide) (*U03,X60-X84,Y87.0) | 797 | 1.4 | 11.1 |
| 12 | Pneumonitis due to solids and liquids (J69) | 590 | 1.0 | 8.2 |
| 13 | Chronic liver disease and cirrhosis (K70,K73-K74) | 531 | 0.9 | 7.4 |
| 14 | Assault (homicide) (*U01-*U02,X85-Y09,Y87.1) | 509 | 0.9 | 7.1 |
| 15 | Essential (primary) hypertension and hypertensive renal disease (I10,I12) | 399 | 0.7 | 5.5 |
| ... | All other causes (Residual) | 9,387 | 16.7 | 130.4 |

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

... Category not applicable

LCWK9. Deaths, percent of total deaths, and death rates for the 15 leading causes of death: United States and each State, 2001

[Rates per 100,000 population.]

| Rank ¹ | Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992) and State | Number | Percent of total deaths | Rate |
|-------------------|--|--------|-------------------------|---------|
| Washington | | | | |
| ... | All causes | 44,642 | 100.0 | 744.9 |
| 1 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 11,281 | 25.3 | 188.2 |
| 2 | Malignant neoplasms (C00-C97) | 10,802 | 24.2 | 180.2 |
| 3 | Cerebrovascular diseases (I60-I69) | 3,765 | 8.4 | 62.8 |
| 4 | Chronic lower respiratory diseases (J40-J47) | 2,634 | 5.9 | 43.9 |
| 5 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 2,072 | 4.6 | 34.6 |
| 6 | Alzheimer's disease (G30) | 2,050 | 4.6 | 34.2 |
| 7 | Diabetes mellitus (E10-E14) | 1,403 | 3.1 | 23.4 |
| 8 | Influenza and pneumonia (J10-J18) | 965 | 2.2 | 16.1 |
| 9 | Intentional self-harm (suicide) (*U03,X60-X84,Y87.0) | 712 | 1.6 | 11.9 |
| 10 | Chronic liver disease and cirrhosis (K70,K73-K74) | 571 | 1.3 | 9.5 |
| 11 | Pneumonitis due to solids and liquids (J69) | 448 | 1.0 | 7.5 |
| 12 | Parkinson's disease (G20-G21) | 434 | 1.0 | 7.2 |
| 13 | Septicemia (A40-A41) | 350 | 0.8 | 5.8 |
| 14 | Atherosclerosis (I70) | 347 | 0.8 | 5.8 |
| 15 | Aortic aneurysm and dissection (I71) | 321 | 0.7 | 5.4 |
| ... | All other causes (Residual) | 6,487 | 14.5 | 108.2 |
| West Virginia | | | | |
| ... | All causes | 20,967 | 100.0 | 1,164.2 |
| 1 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 6,325 | 30.2 | 351.2 |
| 2 | Malignant neoplasms (C00-C97) | 4,685 | 22.3 | 260.1 |
| 3 | Chronic lower respiratory diseases (J40-J47) | 1,273 | 6.1 | 70.7 |
| 4 | Cerebrovascular diseases (I60-I69) | 1,272 | 6.1 | 70.6 |
| 5 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 834 | 4.0 | 46.3 |
| 6 | Diabetes mellitus (E10-E14) | 802 | 3.8 | 44.5 |
| 7 | Influenza and pneumonia (J10-J18) | 483 | 2.3 | 26.8 |
| 8 | Alzheimer's disease (G30) | 398 | 1.9 | 22.1 |
| 9 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 391 | 1.9 | 21.7 |
| 10 | Septicemia (A40-A41) | 334 | 1.6 | 18.5 |
| 11 | Intentional self-harm (suicide) (*U03,X60-X84,Y87.0) | 286 | 1.4 | 15.9 |
| 12 | Chronic liver disease and cirrhosis (K70,K73-K74) | 212 | 1.0 | 11.8 |
| 13 | Essential (primary) hypertension and hypertensive renal disease (I10,I12) | 164 | 0.8 | 9.1 |
| 14 | Atherosclerosis (I70) | 140 | 0.7 | 7.8 |
| 15 | Pneumonitis due to solids and liquids (J69) | 137 | 0.7 | 7.6 |
| ... | All other causes (Residual) | 3,231 | 15.4 | 179.4 |

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

... Category not applicable

LCWK9. Deaths, percent of total deaths, and death rates for the 15 leading causes of death: United States and each State, 2001

[Rates per 100,000 population.]

| Rank ¹ | Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992) and State | Number | Percent of total deaths | Rate |
|-------------------|--|--------|-------------------------|-------|
| Wisconsin | | | | |
| ... | All causes | 46,628 | 100.0 | 862.5 |
| 1 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 13,023 | 27.9 | 240.9 |
| 2 | Malignant neoplasms (C00-C97) | 10,777 | 23.1 | 199.4 |
| 3 | Cerebrovascular diseases (I60-I69) | 3,658 | 7.8 | 67.7 |
| 4 | Chronic lower respiratory diseases (J40-J47) | 2,375 | 5.1 | 43.9 |
| 5 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 2,100 | 4.5 | 38.8 |
| 6 | Diabetes mellitus (E10-E14) | 1,337 | 2.9 | 24.7 |
| 7 | Alzheimer's disease (G30) | 1,269 | 2.7 | 23.5 |
| 8 | Influenza and pneumonia (J10-J18) | 1,236 | 2.7 | 22.9 |
| 9 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 791 | 1.7 | 14.6 |
| 10 | Intentional self-harm (suicide) (*U03,X60-X84,Y87.0) | 639 | 1.4 | 11.8 |
| 11 | Chronic liver disease and cirrhosis (K70,K73-K74) | 431 | 0.9 | 8.0 |
| 12 | Septicemia (A40-A41) | 416 | 0.9 | 7.7 |
| 13 | Parkinson's disease (G20-G21) | 383 | 0.8 | 7.1 |
| 14 | Aortic aneurysm and dissection (I71) | 377 | 0.8 | 7.0 |
| 15 | In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48) | 356 | 0.8 | 6.6 |
| ... | All other causes (Residual) | 7,460 | 16.0 | 138.0 |
| Wyoming | | | | |
| ... | All causes | 4,029 | 100.0 | 816.0 |
| 1 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 985 | 24.4 | 199.5 |
| 2 | Malignant neoplasms (C00-C97) | 923 | 22.9 | 186.9 |
| 3 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 272 | 6.8 | 55.1 |
| 4 | Chronic lower respiratory diseases (J40-J47) | 270 | 6.7 | 54.7 |
| 5 | Cerebrovascular diseases (I60-I69) | 260 | 6.5 | 52.7 |
| 6 | Diabetes mellitus (E10-E14) | 120 | 3.0 | 24.3 |
| 7 | Alzheimer's disease (G30) | 114 | 2.8 | 23.1 |
| 8 | Influenza and pneumonia (J10-J18) | 112 | 2.8 | 22.7 |
| 9 | Intentional self-harm (suicide) (*U03,X60-X84,Y87.0) | 83 | 2.1 | 16.8 |
| 10 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 54 | 1.3 | 10.9 |
| 11 | Chronic liver disease and cirrhosis (K70,K73-K74) | 49 | 1.2 | 9.9 |
| 12 | Septicemia (A40-A41) | 39 | 1.0 | 7.9 |
| 13 | In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48) | 31 | 0.8 | 6.3 |
| 14 | Parkinson's disease (G20-G21) | 26 | 0.6 | 5.3 |
| 15 | Aortic aneurysm and dissection (I71) | 23 | 0.6 | 4.7 |
| ... | All other causes (Residual) | 668 | 16.6 | 135.3 |

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

... Category not applicable