Person with disabilities experiencing problems accessing NCHS birth and death tables should contact **nchsed@cdc.gov**, or call 301-458-4688

Worktable 13. Age-adjusted death rates for 358 selected causes, by race and sex: United States, 1999-2005

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992) and year	All races 1/			White 2/			Black 2/		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
All causes									
2005	798.8	951.1	677.6	785.3	933.2	666.5	1,016.5	1,252.9	845.7
2004	800.8	955.7	679.2	786.3	936.9	666.9	1,027.3	1,269.4	855.3
2003	832.7	994.3	706.2	817.0	973.9	693.1	1,065.9	1,319.1	885.6
2002	845.3	1,013.7	715.2	829.0	992.9	701.3	1,083.3	1,341.4	901.8
2001	854.5	1,029.1	721.8	836.5	1,006.1	706.7	1,101.2	1,375.0	912.5
2000	869.0	1,053.8	731.4	849.8	1,029.4	715.3	1,121.4	1,403.5	927.6
1999	875.6	1,067.0	734.0	854.6	1,040.0	716.6	1,135.7	1,432.6	933.6
I. Certain infectious and parasitic diseases (A00-B99)									
2005	21.8	26.3	17.9	18.7	22.4	15.4	49.6	63.7	38.6
2004	21.7	26.3	17.6	18.4	22.2	15.0	50.5	65.0	39.1
2003	22.1	26.7	17.9	18.6	22.4	15.2	52.5	68.3	40.0
2002	22.2	27.2	17.7	18.5	22.5	14.9	54.2	71.0	41.0
2001	21.3	26.3	16.9	17.6	21.7	14.1	53.3	70.1	40.2
2000	21.3	26.3	16.8	17.4	21.4	13.9	55.0	72.9	41.0
1999	21.8	27.2	17.2	17.8	22.0	14.1	56.9	75.9	42.0
Intestinal infectious diseases (A00-A09)									
2005	1.9	1.9	1.8	2.0	2.0	1.9	1.2	1.3	1.2
2004	1.5	1.4	1.5	1.5	1.5	1.6	0.9	1.0	0.9
2003	1.1	1.0	1.1	1.1	1.0	1.1	0.8	0.7	0.8
2002	0.9	0.9	0.9	0.9	0.9	0.9	0.6	0.7	0.6
2001	0.5	0.5	0.6	0.6	0.6	0.6	0.4	0.5	0.4
2000	0.5	0.5	0.5	0.5	0.5	0.5	0.3	0.4	0.3
1999	0.4	0.4	0.4	0.4	0.4	0.4	0.3	0.3	0.3

Worktable 13. Age-adjusted death rates for 358 selected causes, by race and sex: United States, 1999-2005

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992) and year	All races 1/			White 2/			Black 2/		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
Cholera (A00)									
2005	*	*	*	*	*	*	*	*	*
2004	*	*	*	*	*	*	*	*	*
2003	*	*	*	*	*	*	*	*	*
2002	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*
Other intestinal infectious diseases (A01-A08)									
2005	1.8	1.8	1.8	1.9	1.9	1.9	1.2	1.3	1.2
2004	1.4	1.4	1.4	1.5	1.5	1.5	0.9	0.9	0.9
2003	1.0	1.0	1.0	1.1	1.0	1.1	0.7	0.7	0.8
2002	0.8	0.8	0.8	0.9	0.9	0.9	0.6	0.7	0.6
2001	0.5	0.5	0.5	0.5	0.5	0.6	0.4	0.5	0.4
2000	0.5	0.5	0.5	0.5	0.5	0.5	0.3	0.4	0.3
1999	0.4	0.4	0.4	0.4	0.4	0.4	0.3	0.3	0.3
Typhoid fever (A01.0)									
2005	*	*	*	*	*	*	*	*	*
2004	*	*	*	*	*	*	*	*	*
2003	*	*	*	*	*	*	*	*	*
2002	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*

Worktable 13. Age-adjusted death rates for 358 selected causes, by race and sex: United States, 1999-2005

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992) and year	All races 1/			White 2/			Black 2/		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
Paratyphoid fevers and other salmonella infections (A01.1-A01.4,A02)									
2005	0.0	*	*	0.0	*	*	*	*	*
2004	0.0	0.0	*	0.0	*	*	*	*	*
2003	0.0	0.0	0.0	0.0	*	*	*	*	*
2002	0.0	*	*	*	*	*	*	*	*
2001	0.0	0.0	*	0.0	*	*	*	*	*
2000	0.0	*	*	0.0	*	*	*	*	*
1999	0.0	0.0	*	0.0	*	*	*	*	*
Shigellosis (A03)									
2005	*	*	*	*	*	*	*	*	*
2004	*	*	*	*	*	*	*	*	*
2003	*	*	*	*	*	*	*	*	*
2002	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*
Other bacterial food-borne intoxications (A05)									
2005	*	*	*	*	*	*	*	*	*
2004	*	*	*	*	*	*	*	*	*
2003	*	*	*	*	*	*	*	*	*
2002	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*

Worktable 13. Age-adjusted death rates for 358 selected causes, by race and sex: United States, 1999-2005

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992) and year	All races 1/			White 2/			Black 2/		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
Amebiasis (A06)									
2005	*	*	*	*	*	*	*	*	*
2004	*	*	*	*	*	*	*	*	*
2003	*	*	*	*	*	*	*	*	*
2002	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*
Intestinal infections due to other specified organisms (A04,A07-A08)									
2005	1.8	1.8	1.8	1.9	1.9	1.9	1.2	1.2	1.2
2004	1.4	1.4	1.4	1.5	1.4	1.5	0.9	0.9	0.9
2003	1.0	1.0	1.0	1.0	1.0	1.1	0.7	0.7	0.7
2002	0.8	0.8	0.8	0.8	0.9	0.8	0.6	0.7	0.6
2001	0.5	0.5	0.5	0.5	0.5	0.5	0.4	0.4	0.4
2000	0.5	0.5	0.4	0.5	0.5	0.5	0.3	0.4	0.3
1999	0.4	0.4	0.4	0.4	0.4	0.4	0.3	0.3	0.3
Diarrhea and gastroenteritis of infectious origin (A09)									
2005	0.0	0.0	0.0	0.0	0.0	0.0	*	*	*
2004	0.0	0.0	0.0	0.0	0.0	0.0	*	*	*
2003	0.0	0.0	0.0	0.0	0.0	0.0	*	*	*
2002	0.0	0.0	0.0	0.0	0.0	0.0	*	*	*
2001	0.0	0.0	0.0	0.0	0.0	0.0	*	*	*
2000	0.0	0.0	0.0	0.0	0.0	0.0	*	*	*
1999	0.0	0.0	0.0	0.0	0.0	0.0	*	*	*

Worktable 13. Age-adjusted death rates for 358 selected causes, by race and sex: United States, 1999-2005

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992) and year	All races 1/			White 2/			Black 2/		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
Tuberculosis (A16-A19)									
2005	0.2	0.3	0.1	0.1	0.2	0.1	0.5	0.8	0.3
2004	0.2	0.3	0.1	0.2	0.2	0.1	0.7	1.0	0.4
2003	0.2	0.3	0.1	0.2	0.2	0.1	0.6	1.0	0.4
2002	0.3	0.4	0.2	0.2	0.3	0.1	0.7	1.1	0.5
2001	0.3	0.4	0.2	0.2	0.2	0.1	0.8	1.2	0.5
2000	0.3	0.4	0.2	0.2	0.2	0.1	0.9	1.5	0.6
1999	0.3	0.5	0.2	0.2	0.3	0.1	0.9	1.5	0.6
Respiratory tuberculosis (A16)									
2005	0.1	0.2	0.1	0.1	0.1	0.1	0.4	0.6	0.2
2004	0.2	0.2	0.1	0.1	0.2	0.1	0.5	0.8	0.3
2003	0.2	0.3	0.1	0.1	0.2	0.1	0.5	0.8	0.3
2002	0.2	0.3	0.1	0.2	0.2	0.1	0.6	0.9	0.4
2001	0.2	0.3	0.1	0.1	0.2	0.1	0.6	1.0	0.3
2000	0.2	0.3	0.1	0.1	0.2	0.1	0.7	1.3	0.4
1999	0.3	0.4	0.2	0.2	0.3	0.1	0.8	1.2	0.5
Other tuberculosis (A17-A19)									
2005	0.0	0.1	0.0	0.0	0.0	0.0	0.1	0.2	*
2004	0.1	0.1	0.0	0.0	0.0	0.0	0.2	0.2	*
2003	0.0	0.1	0.0	0.0	0.0	0.0	0.1	0.1	*
2002	0.0	0.1	0.0	0.0	0.0	0.0	0.1	0.2	*
2001	0.1	0.1	0.0	0.0	0.0	0.0	0.2	0.2	0.1
2000	0.1	0.1	0.0	0.0	0.0	0.0	0.2	0.3	0.2
1999	0.1	0.1	0.0	0.0	0.1	0.0	0.2	0.3	*

Worktable 13. Age-adjusted death rates for 358 selected causes, by race and sex: United States, 1999-2005

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992) and year	All races 1/			White 2/			Black 2/		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
Zoonotic and other bacterial diseases (A20-A49)									
2005	11.8	13.0	10.9	10.7	11.9	9.9	23.4	26.9	21.2
2004	11.8	13.0	10.8	10.7	12.0	9.9	23.4	26.9	21.1
2003	12.2	13.3	11.4	11.1	12.1	10.3	24.6	28.5	22.2
2002	12.3	13.5	11.4	11.2	12.4	10.4	25.1	28.8	22.8
2001	12.0	13.4	11.0	10.9	12.2	10.0	24.5	28.8	22.0
2000	11.9	13.1	11.0	10.7	11.8	9.9	25.5	29.2	23.2
1999	12.0	13.4	11.0	10.7	11.9	9.9	26.0	30.6	23.0
Plague (A20)									
2005	*	*	*	*	*	*	*	*	*
2004	*	*	*	*	*	*	*	*	*
2003	*	*	*	*	*	*	*	*	*
2002	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*
Brucellosis (A23)									
2005	*	*	*	*	*	*	*	*	*
2004	*	*	*	*	*	*	*	*	*
2003	*	*	*	*	*	*	*	*	*
2002	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*

Worktable 13. Age-adjusted death rates for 358 selected causes, by race and sex: United States, 1999-2005

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992) and year	All races 1/			White 2/			Black 2/		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
Tetanus (A33-A35)									
2005	*	*	*	*	*	*	*	*	*
2004	*	*	*	*	*	*	*	*	*
2003	*	*	*	*	*	*	*	*	*
2002	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*
Diphtheria (A36)									
2005	*	*	*	*	*	*	*	*	*
2004	*	*	*	*	*	*	*	*	*
2003	*	*	*	*	*	*	*	*	*
2002	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*
Whooping cough (A37)									
2005	0.0	*	*	0.0	*	*	*	*	*
2004	*	*	*	*	*	*	*	*	*
2003	*	*	*	*	*	*	*	*	*
2002	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*

Worktable 13. Age-adjusted death rates for 358 selected causes, by race and sex: United States, 1999-2005

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992) and year	All races 1/			White 2/			Black 2/		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
Scarlet fever and erysipelas (A38,A46)									
2005	*	*	*	*	*	*	*	*	*
2004	*	*	*	*	*	*	*	*	*
2003	*	*	*	*	*	*	*	*	*
2002	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*
Meningococcal infection (A39)									
2005	0.0	0.0	0.0	0.0	0.0	0.0	*	*	*
2004	0.0	0.0	0.0	0.0	0.0	0.0	0.1	*	*
2003	0.0	0.0	0.0	0.0	0.0	0.0	0.1	*	*
2002	0.0	0.1	0.0	0.0	0.1	0.0	0.1	*	*
2001	0.1	0.1	0.1	0.1	0.0	0.1	0.1	*	*
2000	0.0	0.1	0.1	0.0	0.1	0.1	0.1	0.1	*
1999	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	*
Septicemia (A40-A41)									
2005	11.2	12.3	10.4	10.2	11.2	9.4	22.6	25.8	20.5
2004	11.2	12.3	10.3	10.2	11.3	9.4	22.5	25.7	20.4
2003	11.6	12.7	10.9	10.5	11.5	9.9	23.8	27.3	21.6
2002	11.7	12.9	10.9	10.7	11.7	9.9	24.2	27.8	22.1
2001	11.4	12.7	10.5	10.3	11.5	9.5	23.6	27.7	21.2
2000	11.3	12.4	10.5	10.2	11.2	9.4	24.6	28.0	22.5
1999	11.3	12.5	10.4	10.0	11.1	9.3	24.6	28.9	21.9

Worktable 13. Age-adjusted death rates for 358 selected causes, by race and sex: United States, 1999-2005

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992) and year	All races 1/			White 2/			Black 2/		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
Other zoonotic and bacterial diseases (A21-A22,A24-A32,A42-A44,A48-A49)									
2005	0.5	0.7	0.4	0.5	0.6	0.4	0.8	1.0	0.6
2004	0.5	0.6	0.5	0.5	0.6	0.4	0.8	1.1	0.7
2003	0.5	0.6	0.4	0.5	0.6	0.4	0.8	1.1	0.5
2002	0.5	0.6	0.4	0.5	0.6	0.4	0.8	1.0	0.7
2001	0.5	0.6	0.4	0.5	0.6	0.4	0.8	1.0	0.7
2000	0.5	0.6	0.4	0.4	0.5	0.4	0.7	1.0	0.6
1999	0.7	0.8	0.6	0.6	0.7	0.5	1.2	1.6	1.0
Infections with a predominately sexual mode of transmission (A50-A64)									
2005	0.0	0.0	*	0.0	*	*	0.1	0.2	*
2004	0.0	0.0	0.0	0.0	*	*	0.1	*	*
2003	0.0	0.0	0.0	0.0	*	*	*	*	*
2002	0.0	0.0	0.0	0.0	*	*	0.1	*	*
2001	0.0	0.0	0.0	0.0	*	*	*	*	*
2000	0.0	0.0	0.0	0.0	*	0.0	0.1	0.2	*
1999	0.0	*	0.0	0.0	*	*	*	*	*
Syphilis (A50-A53)									
2005	0.0	0.0	*	*	*	*	0.1	0.2	*
2004	0.0	0.0	0.0	*	*	*	0.1	*	*
2003	0.0	0.0	*	*	*	*	*	*	*
2002	0.0	0.0	0.0	*	*	*	0.1	*	*
2001	0.0	*	*	0.0	*	*	*	*	*
2000	0.0	0.0	*	0.0	*	*	0.1	*	*
1999	0.0	*	0.0	0.0	*	*	*	*	*

Worktable 13. Age-adjusted death rates for 358 selected causes, by race and sex: United States, 1999-2005

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992) and year	All races 1/			White 2/			Black 2/		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
Cardiovascular syphilis (A52.0)									
2005	*	*	*	*	*	*	*	*	*
2004	*	*	*	*	*	*	*	*	*
2003	*	*	*	*	*	*	*	*	*
2002	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*
Neurosyphilis (A52.1-A52.3)									
2005	0.0	0.0	*	*	*	*	*	*	*
2004	0.0	*	*	*	*	*	*	*	*
2003	*	*	*	*	*	*	*	*	*
2002	0.0	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*
2000	0.0	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*
Other and unspecified syphilis (A50-A51,A52.7-A52.9,A53)									
2005	*	*	*	*	*	*	*	*	*
2004	*	*	*	*	*	*	*	*	*
2003	*	*	*	*	*	*	*	*	*
2002	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*

Worktable 13. Age-adjusted death rates for 358 selected causes, by race and sex: United States, 1999-2005

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992) and year	All races 1/			White 2/			Black 2/		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
Gonococcal infection (A54)									
2005	*	*	*	*	*	*	*	*	*
2004	*	*	*	*	*	*	*	*	*
2003	*	*	*	*	*	*	*	*	*
2002	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*
Other infections with a predominately sexual mode of transmission (A55-A64)									
2005	*	*	*	*	*	*	*	*	*
2004	*	*	*	*	*	*	*	*	*
2003	*	*	*	*	*	*	*	*	*
2002	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*
Other infections caused by spirochetes, chlamydia or rickettsia (A65-A79)									
2005	*	*	*	*	*	*	*	*	*
2004	*	*	*	*	*	*	*	*	*
2003	*	*	*	*	*	*	*	*	*
2002	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*
1999	0.0	*	*	*	*	*	*	*	*

Worktable 13. Age-adjusted death rates for 358 selected causes, by race and sex: United States, 1999-2005

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992) and year	All races 1/			White 2/			Black 2/		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
Lyme disease (A69.2)									
2005	*	*	*	*	*	*	*	*	*
2004	*	*	*	*	*	*	*	*	*
2003	*	*	*	*	*	*	*	*	*
2002	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*
Typhus fever (A75)									
2005	*	*	*	*	*	*	*	*	*
2004	*	*	*	*	*	*	*	*	*
2003	*	*	*	*	*	*	*	*	*
2002	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*
All other infections caused by spirochetes, chlamydia or rickettsia (A65-A68,A69.0-A69.1,A69.8-A69.9,A70-A74,A77-A79)									
2005	*	*	*	*	*	*	*	*	*
2004	*	*	*	*	*	*	*	*	*
2003	*	*	*	*	*	*	*	*	*
2002	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*

Worktable 13. Age-adjusted death rates for 358 selected causes, by race and sex: United States, 1999-2005

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992) and year	All races 1/			White 2/			Black 2/		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
Viral diseases (A80-B34)									
2005	6.5	9.2	3.9	4.4	6.5	2.3	22.4	32.1	14.2
2004	6.7	9.5	4.0	4.5	6.6	2.3	23.4	33.4	15.0
2003	7.1	10.2	4.1	4.8	7.2	2.5	24.5	35.6	15.1
2002	7.4	10.7	4.3	5.0	7.4	2.6	25.8	37.8	15.7
2001	7.4	10.6	4.2	4.9	7.4	2.5	25.9	37.5	15.9
2000	7.5	11.0	4.3	5.0	7.5	2.6	26.5	39.3	15.6
1999	7.6	11.1	4.2	5.1	7.7	2.6	26.7	40.1	15.4
Acute poliomyelitis (A80)									
2005	*	*	*	*	*	*	*	*	*
2004	*	*	*	*	*	*	*	*	*
2003	*	*	*	*	*	*	*	*	*
2002	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*
Rabies (A82)									
2005	*	*	*	*	*	*	*	*	*
2004	*	*	*	*	*	*	*	*	*
2003	*	*	*	*	*	*	*	*	*
2002	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*

Worktable 13. Age-adjusted death rates for 358 selected causes, by race and sex: United States, 1999-2005

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992) and year	All races 1/			White 2/			Black 2/		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
Arthropod-borne viral encephalitis (A83-A84,A85.2)									
2005	*	*	*	*	*	*	*	*	*
2004	*	*	*	*	*	*	*	*	*
2003	*	*	*	*	*	*	*	*	*
2002	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*
Yellow fever (A95)									
2005	*	*	*	*	*	*	*	*	*
2004	*	*	*	*	*	*	*	*	*
2003	*	*	*	*	*	*	*	*	*
2002	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*
Other and unspecified arthropod-borne viral and hemorrhagic fevers (A90-A94,A96-A99)									
2005	0.0	0.0	0.0	0.0	0.0	0.0	*	*	*
2004	0.0	0.0	0.0	0.0	0.0	0.0	*	*	*
2003	0.1	0.1	0.0	0.1	0.1	0.0	*	*	*
2002	0.0	0.1	0.0	0.0	0.1	0.0	0.1	*	*
2001	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*

Worktable 13. Age-adjusted death rates for 358 selected causes, by race and sex: United States, 1999-2005

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992) and year	All races 1/			White 2/			Black 2/		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
Herpesviral (herpes simplex) infections (B00)									
2005	0.1	0.1	0.0	0.0	0.1	0.0	0.1	*	*
2004	0.1	0.1	0.1	0.1	0.1	0.1	*	*	*
2003	0.1	0.0	0.1	0.1	0.1	0.1	*	*	*
2002	0.0	0.1	0.1	0.1	0.0	0.1	0.1	*	*
2001	0.1	0.1	0.1	0.1	0.1	0.1	0.0	*	*
2000	0.0	0.0	0.0	0.0	0.0	0.0	0.1	*	*
1999	0.0	0.0	0.0	0.0	0.0	0.0	0.1	*	*
Zoster (herpes zoster)(B02)									
2005	0.0	0.0	0.1	0.0	0.0	0.1	*	*	*
2004	0.0	0.1	0.0	0.1	0.0	0.1	*	*	*
2003	0.0	0.0	0.0	0.0	0.0	0.0	*	*	*
2002	0.0	0.0	0.0	0.0	0.0	0.0	*	*	*
2001	0.0	0.0	0.0	0.0	0.0	0.1	*	*	*
2000	0.0	0.0	0.0	0.0	0.0	0.0	*	*	*
1999	0.1	0.0	0.1	0.1	0.0	0.1	*	*	*
Smallpox (B03)									
2005	*	*	*	*	*	*	*	*	*
2004	*	*	*	*	*	*	*	*	*
2003	*	*	*	*	*	*	*	*	*
2002	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*

Worktable 13. Age-adjusted death rates for 358 selected causes, by race and sex: United States, 1999-2005

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992) and year	All races 1/			White 2/			Black 2/		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
Measles (B05)									
2005	*	*	*	*	*	*	*	*	*
2004	*	*	*	*	*	*	*	*	*
2003	*	*	*	*	*	*	*	*	*
2002	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*
Rubella (German measles) (B06)									
2005	*	*	*	*	*	*	*	*	*
2004	*	*	*	*	*	*	*	*	*
2003	*	*	*	*	*	*	*	*	*
2002	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*
Viral hepatitis (B15-B19)									
2005	1.8	2.4	1.2	1.7	2.3	1.1	2.6	3.4	1.9
2004	1.8	2.4	1.2	1.7	2.3	1.1	2.6	3.8	1.7
2003	1.8	2.5	1.2	1.7	2.4	1.1	2.7	3.8	1.8
2002	2.0	2.7	1.3	1.9	2.6	1.2	2.8	3.8	1.9
2001	2.0	2.6	1.3	1.8	2.5	1.2	2.7	3.4	2.1
2000	1.9	2.6	1.3	1.8	2.4	1.2	2.7	3.8	1.8
1999	1.8	2.4	1.2	1.6	2.2	1.1	2.6	3.4	1.9

Worktable 13. Age-adjusted death rates for 358 selected causes, by race and sex: United States, 1999-2005

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992) and year	All races 1/			White 2/			Black 2/		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
Human immunodeficiency virus (HIV) disease (B20-B24)									
2005	4.2	6.2	2.3	2.2	3.6	0.8	19.4	28.2	12.0
2004	4.5	6.6	2.4	2.3	3.8	0.9	20.4	29.2	13.0
2003	4.7	7.1	2.4	2.5	4.2	0.9	21.3	31.3	12.8
2002	4.9	7.4	2.5	2.6	4.3	0.9	22.5	33.3	13.4
2001	5.0	7.5	2.5	2.6	4.4	0.9	22.8	33.8	13.4
2000	5.2	7.9	2.5	2.8	4.6	1.0	23.3	35.1	13.2
1999	5.3	8.2	2.5	2.9	4.9	1.0	23.6	36.1	13.1
Human immunodeficiency virus (HIV) disease resulting in infectious and parasitic diseases (B20)									
2005	1.6	2.4	0.9	0.9	1.4	0.3	7.2	10.4	4.6
2004	1.8	2.5	1.0	0.9	1.5	0.4	7.8	10.9	5.1
2003	1.8	2.7	0.9	1.0	1.7	0.4	7.7	11.5	4.5
2002	1.9	2.9	1.0	1.1	1.7	0.4	8.6	12.8	5.1
2001	1.9	2.9	1.0	1.1	1.8	0.4	8.6	12.6	5.1
2000	2.0	3.1	1.0	1.1	1.8	0.4	9.0	13.4	5.2
1999	2.1	3.3	1.1	1.2	1.9	0.4	9.3	14.1	5.2
Human immunodeficiency virus (HIV) disease resulting in malignant neoplasms (B21)									
2005	0.3	0.5	0.1	0.2	0.4	0.1	1.2	1.9	0.6
2004	0.3	0.5	0.1	0.2	0.4	0.0	1.1	1.8	0.5
2003	0.3	0.5	0.1	0.2	0.4	0.0	1.1	1.6	0.6
2002	0.3	0.5	0.1	0.2	0.4	0.0	1.2	1.8	0.6
2001	0.3	0.6	0.1	0.2	0.4	0.1	1.2	2.0	0.6
2000	0.4	0.6	0.1	0.3	0.5	0.0	1.3	2.1	0.5
1999	0.4	0.7	0.1	0.3	0.5	0.1	1.3	2.2	0.5

Worktable 13. Age-adjusted death rates for 358 selected causes, by race and sex: United States, 1999-2005

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992) and year	All races 1/			White 2/			Black 2/		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
Human immunodeficiency virus (HIV) disease resulting in other specified diseases (B22)									
2005	0.4	0.6	0.2	0.2	0.4	0.1	1.8	2.6	1.2
2004	0.4	0.7	0.2	0.3	0.4	0.1	1.8	2.7	1.1
2003	0.4	0.7	0.2	0.3	0.5	0.1	1.8	2.9	0.9
2002	0.5	0.7	0.2	0.3	0.5	0.1	1.8	2.8	1.0
2001	0.5	0.8	0.2	0.3	0.5	0.1	2.1	3.1	1.2
2000	0.6	0.9	0.2	0.3	0.6	0.1	2.3	3.6	1.2
1999	0.6	0.9	0.2	0.4	0.6	0.1	2.3	3.6	1.2
Human immunodeficiency virus (HIV) disease resulting in other conditions (B23)									
2005	0.7	1.0	0.4	0.3	0.5	0.1	3.7	5.4	2.2
2004	0.7	1.1	0.4	0.3	0.5	0.1	3.8	5.6	2.4
2003	1.1	1.6	0.6	0.5	0.8	0.2	5.8	8.2	3.8
2002	1.2	1.7	0.7	0.5	0.8	0.2	6.2	9.1	3.8
2001	1.2	1.8	0.7	0.6	0.9	0.2	6.2	9.1	3.8
2000	1.2	1.8	0.7	0.6	0.9	0.2	6.0	8.9	3.7
1999	1.2	1.8	0.7	0.6	0.9	0.2	6.4	9.5	3.7
Unspecified human immunodeficiency virus (HIV) disease (B24)									
2005	1.2	1.7	0.6	0.6	1.0	0.2	5.5	7.9	3.4
2004	1.2	1.7	0.7	0.6	1.0	0.2	5.8	8.2	3.9
2003	1.0	1.5	0.6	0.5	0.8	0.2	4.9	7.1	3.0
2002	1.0	1.4	0.5	0.5	0.8	0.2	4.6	6.8	2.9
2001	1.0	1.5	0.5	0.5	0.8	0.1	4.7	7.0	2.8
2000	1.0	1.5	0.5	0.5	0.8	0.2	4.7	7.1	2.6
1999	1.0	1.5	0.5	0.5	0.9	0.2	4.4	6.6	2.5

Worktable 13. Age-adjusted death rates for 358 selected causes, by race and sex: United States, 1999-2005

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992) and year	All races 1/			White 2/			Black 2/		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
All other and unspecified viral diseases (A81,A85.0-A85.1,A85.8,A86-A89,B01,B04,B07-B09,B25-B34)									
2005	0.3	0.4	0.3	0.3	0.4	0.3	0.3	0.3	0.3
2004	0.3	0.4	0.3	0.3	0.4	0.3	0.3	0.3	0.3
2003	0.3	0.4	0.3	0.3	0.4	0.3	0.3	0.3	0.3
2002	0.4	0.4	0.3	0.4	0.4	0.3	0.3	0.4	0.3
2001	0.3	0.3	0.3	0.3	0.3	0.3	0.3	0.3	0.3
2000	0.4	0.4	0.3	0.4	0.4	0.3	0.4	0.4	0.4
1999	0.4	0.4	0.4	0.4	0.4	0.4	0.4	0.4	0.4
Other and unspecified infectious and parasitic diseases and their sequelae (B35-B99)									
2005	1.5	1.9	1.2	1.4	1.8	1.1	1.9	2.4	1.5
2004	1.5	1.9	1.1	1.5	1.9	1.1	2.0	2.5	1.5
2003	1.5	1.8	1.2	1.4	1.7	1.1	1.9	2.4	1.6
2002	1.3	1.7	0.9	1.2	1.6	0.9	1.8	2.5	1.3
2001	1.1	1.3	0.9	1.0	1.2	0.8	1.7	2.1	1.4
2000	1.1	1.3	0.8	1.0	1.2	0.8	1.7	2.2	1.3
1999	1.5	1.8	1.2	1.3	1.6	1.1	2.9	3.3	2.6
Mycoses (B35-B49)									
2005	0.3	0.4	0.3	0.3	0.4	0.2	0.7	0.8	0.6
2004	0.4	0.5	0.3	0.3	0.4	0.3	0.7	0.9	0.5
2003	0.4	0.4	0.3	0.3	0.4	0.3	0.7	0.8	0.6
2002	0.4	0.6	0.3	0.4	0.5	0.3	0.8	1.2	0.5
2001	0.4	0.5	0.3	0.3	0.4	0.3	0.8	1.1	0.6
2000	0.4	0.6	0.3	0.4	0.5	0.3	0.8	1.1	0.7
1999	0.7	0.9	0.5	0.6	0.8	0.4	1.3	1.6	1.1

Worktable 13. Age-adjusted death rates for 358 selected causes, by race and sex: United States, 1999-2005

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992) and year	All races 1/			White 2/			Black 2/		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
Protozoal diseases (B50-B64)									
2005	0.1	0.1	0.0	0.0	0.1	0.0	0.1	0.1	*
2004	0.1	0.1	0.0	0.0	0.1	0.0	0.1	0.2	*
2003	0.1	0.1	0.1	0.0	0.1	0.0	0.1	0.2	0.1
2002	0.1	0.1	0.1	0.1	0.1	0.0	0.2	0.2	0.2
2001	0.1	0.1	0.1	0.1	0.1	0.1	0.2	0.2	0.2
2000	0.1	0.1	0.1	0.1	0.1	0.0	0.2	0.3	0.2
1999	0.1	0.1	0.1	0.1	0.1	0.1	0.2	0.2	0.2
Malaria (B50-B54)									
2005	*	*	*	*	*	*	*	*	*
2004	*	*	*	*	*	*	*	*	*
2003	*	*	*	*	*	*	*	*	*
2002	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*
Leishmaniasis (B55)									
2005	*	*	*	*	*	*	*	*	*
2004	*	*	*	*	*	*	*	*	*
2003	*	*	*	*	*	*	*	*	*
2002	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*

Worktable 13. Age-adjusted death rates for 358 selected causes, by race and sex: United States, 1999-2005

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992) and year	All races 1/			White 2/			Black 2/		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
Trypanosomiasis (B56-B57)									
2005	*	*	*	*	*	*	*	*	*
2004	*	*	*	*	*	*	*	*	*
2003	*	*	*	*	*	*	*	*	*
2002	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*
Pneumocystosis (B59)									
2005	0.0	0.0	0.0	0.0	0.0	0.0	0.1	*	*
2004	0.1	0.1	0.0	0.0	0.1	0.0	0.1	0.1	*
2003	0.1	0.1	0.0	0.0	0.0	0.0	0.1	0.1	*
2002	0.1	0.1	0.0	0.1	0.1	0.0	0.1	0.2	0.1
2001	0.1	0.1	0.1	0.1	0.1	0.1	0.2	0.2	0.1
2000	0.1	0.1	0.1	0.1	0.1	0.0	0.2	0.2	0.1
1999	0.1	0.1	0.1	0.1	0.1	0.1	0.2	0.2	0.2
Other and unspecified protozoal diseases (B58,B60-B64)									
2005	0.0	*	*	*	*	*	*	*	*
2004	*	*	*	*	*	*	*	*	*
2003	*	*	*	*	*	*	*	*	*
2002	0.0	*	*	*	*	*	*	*	*
2001	0.0	*	*	*	*	*	*	*	*
2000	0.0	*	*	*	*	*	*	*	*
1999	0.0	0.0	*	*	*	*	*	*	*

Worktable 13. Age-adjusted death rates for 358 selected causes, by race and sex: United States, 1999-2005

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992) and year	All races 1/			White 2/			Black 2/		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
Helminthiases (B65-B83)									
2005	*	*	*	*	*	*	*	*	*
2004	0.0	0.0	*	0.0	*	*	*	*	*
2003	0.0	*	*	*	*	*	*	*	*
2002	*	*	*	*	*	*	*	*	*
2001	0.0	*	*	0.0	*	*	*	*	*
2000	0.0	0.0	*	0.0	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*
Schistosomiasis (bilharziasis) (B65)									
2005	*	*	*	*	*	*	*	*	*
2004	*	*	*	*	*	*	*	*	*
2003	*	*	*	*	*	*	*	*	*
2002	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*
Other and unspecified helminthiases (B66-B83)									
2005	*	*	*	*	*	*	*	*	*
2004	0.0	*	*	0.0	*	*	*	*	*
2003	0.0	*	*	*	*	*	*	*	*
2002	*	*	*	*	*	*	*	*	*
2001	0.0	*	*	0.0	*	*	*	*	*
2000	0.0	0.0	*	0.0	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*

Worktable 13. Age-adjusted death rates for 358 selected causes, by race and sex: United States, 1999-2005

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992) and year	All races 1/			White 2/			Black 2/		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
Sequelae of tuberculosis (B90)									
2005	0.0	0.0	0.0	0.0	0.0	0.0	*	*	*
2004	0.0	0.0	0.0	0.0	0.0	0.0	*	*	*
2003	0.0	0.0	0.0	0.0	0.0	0.0	*	*	*
2002	0.0	0.0	0.0	0.0	0.0	0.0	*	*	*
2001	0.0	0.0	0.0	0.0	0.0	0.0	*	*	*
2000	0.0	0.1	0.0	0.0	0.0	0.0	*	*	*
1999	0.1	0.1	0.1	0.1	0.1	0.1	0.1	*	*
Sequelae of poliomyelitis (B91)									
2005	0.1	0.1	0.1	0.1	0.1	0.1	*	*	*
2004	0.1	0.1	0.1	0.1	0.1	0.1	*	*	*
2003	0.1	0.1	0.1	0.1	0.1	0.1	*	*	*
2002	0.1	0.1	0.1	0.1	0.1	0.1	*	*	*
2001	0.1	0.1	0.1	0.1	0.1	0.1	*	*	*
2000	0.1	0.1	0.1	0.1	0.1	0.1	*	*	*
1999	0.1	0.1	0.1	0.1	0.1	0.1	*	*	*
All other and unspecified infectious and parasitic diseases and their sequelae (B85-B89,B92-B99)									
2005	0.9	1.2	0.7	0.9	1.2	0.7	1.1	1.4	0.9
2004	0.9	1.2	0.7	0.9	1.2	0.7	1.0	1.3	0.8
2003	0.9	1.1	0.7	0.9	1.1	0.6	1.0	1.4	0.8
2002	0.6	0.8	0.4	0.6	0.8	0.4	0.7	1.0	0.6
2001	0.4	0.5	0.4	0.4	0.5	0.3	0.6	0.6	0.5
2000	0.3	0.4	0.3	0.3	0.4	0.3	0.5	0.7	0.4
1999	0.5	0.6	0.5	0.4	0.5	0.4	1.2	1.4	1.1

Worktable 13. Age-adjusted death rates for 358 selected causes, by race and sex: United States, 1999-2005

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992) and year	All races 1/			White 2/			Black 2/		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
II. Neoplasms (C00-D48)									
2005	188.3	230.6	159.4	187.2	228.0	159.1	226.9	298.3	183.4
2004	190.4	233.3	161.2	189.0	230.1	160.9	231.6	306.3	186.4
2003	194.7	239.1	164.8	193.2	235.9	164.2	237.6	313.9	191.4
2002	198.1	244.6	167.0	196.4	241.0	166.4	243.2	324.7	194.4
2001	200.9	249.8	168.8	198.8	245.4	168.1	247.8	336.5	195.5
2000	204.5	255.0	171.8	202.2	250.1	171.2	253.1	345.7	197.9
1999	205.6	258.0	171.7	202.9	252.7	170.5	257.3	351.4	201.3
Malignant neoplasms (C00-C97)									
2005	183.8	225.1	155.6	182.6	222.3	155.2	222.7	293.7	179.6
2004	185.8	227.7	157.4	184.4	224.4	157.0	227.2	301.2	182.5
2003	190.1	233.3	160.9	188.5	230.1	160.2	233.3	308.8	187.7
2002	193.5	238.9	163.1	191.7	235.2	162.4	238.8	319.6	190.3
2001	196.0	243.7	164.7	193.9	239.2	163.9	243.1	330.9	191.3
2000	199.6	248.9	167.6	197.2	243.9	166.9	248.5	340.3	193.8
1999	200.8	251.9	167.6	197.9	246.5	166.4	252.5	346.1	196.9
Malignant neoplasms of lip, oral cavity and pharynx (C00-C14)									
2005	2.5	3.9	1.4	2.4	3.6	1.4	3.5	6.5	1.4
2004	2.6	4.0	1.5	2.5	3.7	1.5	3.8	6.9	1.6
2003	2.6	4.0	1.5	2.5	3.8	1.4	3.8	6.9	1.6
2002	2.7	4.1	1.5	2.5	3.8	1.5	3.6	6.3	1.7
2001	2.7	4.1	1.6	2.6	3.9	1.5	3.9	6.7	1.8
2000	2.7	4.1	1.6	2.5	3.7	1.6	4.1	7.3	1.8
1999	2.7	4.1	1.6	2.5	3.8	1.5	4.3	7.5	2.1

Worktable 13. Age-adjusted death rates for 358 selected causes, by race and sex: United States, 1999-2005

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992) and year	All races 1/			White 2/			Black 2/		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
Of lip (C00)									
2005	0.0	0.0	0.0	0.0	0.0	0.0	*	*	*
2004	0.0	0.0	0.0	0.0	0.0	0.0	*	*	*
2003	0.0	0.0	0.0	0.0	0.0	*	*	*	*
2002	0.0	0.0	0.0	0.0	0.0	*	*	*	*
2001	0.0	0.0	*	0.0	0.0	*	*	*	*
2000	0.0	0.0	*	0.0	0.0	*	*	*	*
1999	0.0	0.0	*	0.0	0.0	*	*	*	*
Of tongue (C01-C02)									
2005	0.6	0.9	0.4	0.6	0.9	0.4	0.7	1.4	0.2
2004	0.6	0.9	0.4	0.6	0.9	0.4	0.7	1.2	0.3
2003	0.6	0.9	0.4	0.6	0.9	0.4	0.7	1.3	0.4
2002	0.7	0.9	0.4	0.6	0.9	0.4	0.7	1.3	0.3
2001	0.6	0.9	0.4	0.6	0.9	0.4	0.7	1.2	0.3
2000	0.6	0.9	0.4	0.6	0.9	0.4	0.8	1.4	0.3
1999	0.6	0.9	0.4	0.6	0.9	0.4	0.8	1.4	0.4
Of pharynx (C10-C13,C14.0)									
2005	1.0	1.7	0.5	0.9	1.5	0.5	1.9	3.6	0.6
2004	1.1	1.8	0.5	1.0	1.5	0.5	2.1	3.9	0.9
2003	1.1	1.8	0.5	1.0	1.5	0.5	2.0	3.8	0.7
2002	1.1	1.7	0.6	1.0	1.5	0.5	1.8	3.3	0.8
2001	1.1	1.8	0.6	1.0	1.6	0.5	2.0	3.6	0.9
2000	1.1	1.8	0.6	1.0	1.5	0.5	2.2	4.1	0.9
1999	1.1	1.8	0.6	1.0	1.6	0.5	2.3	4.0	1.0

Worktable 13. Age-adjusted death rates for 358 selected causes, by race and sex: United States, 1999-2005

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992) and year	All races 1/			White 2/			Black 2/		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
Of other and unspecified sites within the lip, oral cavity and pharynx (C03-C09,C14.2-C14.8)									
2005	0.8	1.2	0.5	0.8	1.2	0.5	0.9	1.5	0.5
2004	0.9	1.2	0.5	0.8	1.2	0.6	1.0	1.8	0.4
2003	0.9	1.3	0.5	0.9	1.2	0.6	1.1	1.8	0.5
2002	0.9	1.3	0.6	0.9	1.3	0.6	1.0	1.7	0.5
2001	0.9	1.3	0.6	0.9	1.3	0.6	1.2	1.9	0.6
2000	0.9	1.3	0.6	0.9	1.3	0.6	1.1	1.9	0.5
1999	0.9	1.3	0.6	0.9	1.3	0.6	1.2	2.0	0.7
Malignant neoplasms of digestive organs (C15-C26)									
2005	43.5	56.0	33.7	41.9	54.0	32.3	59.8	78.6	46.7
2004	43.9	56.4	34.2	42.4	54.4	32.8	60.4	78.4	47.7
2003	45.1	57.7	35.3	43.4	55.6	33.7	62.4	80.8	49.6
2002	45.7	58.8	35.6	44.0	56.6	34.1	63.4	82.9	50.1
2001	46.4	59.4	36.5	44.6	57.2	34.9	64.8	83.3	52.1
2000	47.2	60.5	37.2	45.3	58.1	35.6	66.2	86.2	52.5
1999	47.3	60.7	37.3	45.4	58.4	35.7	66.7	87.1	52.7
Of esophagus (C15)									
2005	4.4	7.9	1.7	4.5	8.0	1.6	5.0	8.8	2.3
2004	4.3	7.7	1.7	4.3	7.7	1.6	5.4	9.0	2.8
2003	4.4	7.7	1.7	4.3	7.7	1.7	5.7	9.7	2.9
2002	4.4	7.8	1.7	4.3	7.7	1.6	5.9	10.1	2.9
2001	4.4	7.8	1.8	4.3	7.6	1.7	6.4	11.3	3.0
2000	4.4	7.7	1.8	4.3	7.6	1.7	6.5	11.0	3.3
1999	4.4	7.7	1.8	4.2	7.4	1.6	6.5	11.5	3.1

Worktable 13. Age-adjusted death rates for 358 selected causes, by race and sex: United States, 1999-2005

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992) and year	All races 1/			White 2/			Black 2/		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
Of stomach (C16)									
2005	3.8	5.2	2.7	3.3	4.5	2.4	7.2	10.6	4.9
2004	4.0	5.5	2.8	3.5	4.9	2.5	7.5	11.1	5.2
2003	4.1	5.7	3.0	3.6	5.0	2.6	7.8	11.3	5.4
2002	4.2	5.9	3.0	3.7	5.2	2.5	8.4	11.9	6.1
2001	4.4	6.1	3.1	3.8	5.4	2.7	8.3	11.9	5.8
2000	4.6	6.4	3.3	4.0	5.6	2.8	9.0	12.7	6.5
1999	4.6	6.6	3.3	4.1	5.8	2.8	9.2	13.5	6.4
Of small intestine (C17)									
2005	0.4	0.4	0.3	0.3	0.4	0.3	0.5	0.6	0.5
2004	0.4	0.5	0.3	0.3	0.4	0.3	0.6	0.7	0.5
2003	0.4	0.4	0.3	0.3	0.4	0.3	0.5	0.7	0.4
2002	0.4	0.4	0.3	0.3	0.4	0.3	0.6	0.7	0.5
2001	0.4	0.5	0.3	0.4	0.4	0.3	0.6	0.8	0.5
2000	0.4	0.5	0.3	0.4	0.4	0.3	0.5	0.6	0.4
1999	0.4	0.5	0.3	0.4	0.5	0.3	0.6	0.8	0.4
Of colon, rectum and anus (C18-C21)									
2005	17.5	20.9	14.8	16.9	20.3	14.3	24.8	30.2	21.2
2004	18.0	21.5	15.3	17.6	21.0	14.9	24.7	30.0	21.2
2003	19.1	22.9	16.2	18.6	22.4	15.7	26.4	31.9	22.8
2002	19.7	23.7	16.7	19.2	23.1	16.2	26.8	33.1	22.8
2001	20.1	24.2	17.2	19.6	23.7	16.6	27.6	33.1	24.1
2000	20.8	25.1	17.7	20.3	24.5	17.2	28.2	34.9	24.0
1999	20.9	25.3	17.8	20.4	24.8	17.3	28.4	34.2	24.7

Worktable 13. Age-adjusted death rates for 358 selected causes, by race and sex: United States, 1999-2005

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992) and year	All races 1/			White 2/			Black 2/		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
Colon (C18)									
2005	14.4	17.1	12.4	13.9	16.6	11.9	21.3	25.5	18.5
2004	14.9	17.6	12.9	14.5	17.1	12.5	21.3	25.5	18.5
2003	16.0	19.0	13.8	15.5	18.5	13.2	22.7	27.1	19.9
2002	16.5	19.7	14.2	16.0	19.1	13.7	23.2	28.3	19.9
2001	16.9	20.1	14.7	16.4	19.6	14.1	24.0	28.4	21.3
2000	17.6	21.0	15.1	17.1	20.4	14.7	24.6	29.9	21.3
1999	17.7	21.2	15.3	17.3	20.8	14.7	24.8	29.2	22.0
Rectosigmoid junction and rectum (C19-C20)									
2005	2.9	3.7	2.2	2.8	3.6	2.2	3.3	4.5	2.5
2004	2.9	3.7	2.2	2.9	3.7	2.2	3.3	4.3	2.5
2003	2.9	3.8	2.2	2.9	3.7	2.2	3.4	4.5	2.7
2002	3.0	3.9	2.3	3.0	3.8	2.3	3.4	4.6	2.7
2001	3.0	4.0	2.3	3.0	4.0	2.3	3.4	4.5	2.7
2000	3.0	4.0	2.3	3.0	3.9	2.3	3.4	4.7	2.5
1999	3.0	3.9	2.3	3.0	3.9	2.4	3.4	4.8	2.5
Anus and anus canal (C21)									
2005	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2
2004	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.3	0.2
2003	0.2	0.1	0.2	0.2	0.1	0.2	0.2	0.2	0.2
2002	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.1
2001	0.2	0.2	0.2	0.2	0.2	0.2	0.2	*	0.2
2000	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2
1999	0.2	0.1	0.2	0.2	0.1	0.2	0.2	0.2	0.2

Worktable 13. Age-adjusted death rates for 358 selected causes, by race and sex: United States, 1999-2005

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992) and year	All races 1/			White 2/			Black 2/		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
Of liver and intrahepatic bile ducts (C22)									
2005	5.2	7.7	3.2	4.8	7.1	3.0	7.0	11.3	3.8
2004	5.1	7.4	3.2	4.7	6.7	3.0	6.9	10.7	4.1
2003	5.0	7.4	3.1	4.6	6.7	2.8	6.6	10.2	4.0
2002	4.9	7.1	3.0	4.5	6.5	2.8	6.3	9.9	3.6
2001	4.7	6.9	3.0	4.3	6.3	2.7	6.2	9.1	4.1
2000	4.7	6.8	3.0	4.2	6.2	2.7	6.1	9.5	3.6
1999	4.5	6.5	3.0	4.2	5.9	2.7	5.9	8.7	3.8
Liver (C22.0,C22.2-C22.9)									
2005	4.1	6.4	2.2	3.7	5.7	2.0	5.9	9.9	2.8
2004	3.9	6.1	2.2	3.5	5.4	1.9	5.9	9.6	3.1
2003	3.9	6.0	2.1	3.5	5.4	1.9	5.6	8.9	3.1
2002	3.8	5.9	2.1	3.5	5.3	1.9	5.4	8.8	2.8
2001	3.7	5.7	2.1	3.3	5.2	1.9	5.3	8.0	3.3
2000	3.7	5.7	2.1	3.3	5.0	1.9	5.3	8.6	2.8
1999	3.6	5.4	2.1	3.2	4.9	1.9	5.1	7.9	3.1
Intrahepatic bile duct carcinoma (C22.1)									
2005	1.1	1.4	1.0	1.1	1.3	1.0	1.1	1.4	0.9
2004	1.2	1.3	1.1	1.2	1.3	1.0	1.0	1.1	0.9
2003	1.1	1.3	1.0	1.1	1.3	1.0	1.1	1.3	0.9
2002	1.0	1.2	0.9	1.0	1.2	0.9	0.9	1.1	0.8
2001	1.0	1.2	0.9	1.0	1.2	0.9	0.9	1.1	0.8
2000	1.0	1.1	0.9	1.0	1.2	0.9	0.8	0.9	0.8
1999	0.9	1.0	0.9	0.9	1.1	0.9	0.8	0.8	0.7

Worktable 13. Age-adjusted death rates for 358 selected causes, by race and sex: United States, 1999-2005

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992) and year	All races 1/			White 2/			Black 2/		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
Of gallbladder and extrahepatic bile ducts (C23-C24)									
2005	1.1	1.0	1.2	1.1	1.0	1.2	1.3	1.2	1.3
2004	1.1	1.0	1.2	1.1	1.0	1.2	1.2	0.9	1.4
2003	1.2	1.1	1.2	1.1	1.1	1.2	1.1	0.9	1.2
2002	1.2	1.1	1.3	1.1	1.1	1.2	1.4	1.1	1.5
2001	1.3	1.2	1.4	1.3	1.1	1.4	1.3	1.2	1.5
2000	1.3	1.2	1.4	1.3	1.2	1.4	1.2	1.0	1.4
1999	1.3	1.2	1.4	1.3	1.2	1.4	1.3	1.0	1.4
Of pancreas (C25)									
2005	10.8	12.3	9.5	10.6	12.2	9.3	13.6	15.3	12.3
2004	10.6	12.3	9.3	10.4	12.2	9.1	13.6	15.4	12.1
2003	10.5	12.0	9.3	10.3	11.9	9.0	13.7	15.3	12.4
2002	10.5	12.1	9.2	10.3	12.0	9.0	13.6	15.3	12.3
2001	10.5	12.1	9.3	10.4	12.0	9.1	13.6	15.0	12.5
2000	10.6	12.2	9.3	10.4	12.0	9.0	14.1	15.8	12.7
1999	10.6	12.3	9.3	10.4	12.0	9.1	14.1	16.4	12.3
Of other and ill-defined digestive organs (C26)									
2005	0.4	0.4	0.3	0.3	0.4	0.3	0.4	0.6	0.3
2004	0.4	0.4	0.3	0.4	0.4	0.3	0.4	0.5	0.4
2003	0.5	0.5	0.4	0.5	0.5	0.4	0.6	0.8	0.5
2002	0.5	0.5	0.4	0.5	0.5	0.4	0.6	0.8	0.5
2001	0.5	0.6	0.4	0.5	0.6	0.4	0.7	0.8	0.6
2000	0.5	0.6	0.4	0.5	0.6	0.4	0.7	0.7	0.6
1999	0.6	0.7	0.5	0.6	0.7	0.5	0.7	0.9	0.5

Worktable 13. Age-adjusted death rates for 358 selected causes, by race and sex: United States, 1999-2005

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992) and year	All races 1/			White 2/			Black 2/		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
Malignant neoplasms of respiratory and intrathoracic organs (C30-C39)									
2005	54.2	71.6	41.2	54.6	71.1	42.2	61.1	91.4	41.0
2004	54.8	72.7	41.6	55.0	71.9	42.6	62.4	95.0	40.9
2003	55.7	74.5	42.0	56.0	73.6	43.0	63.7	98.1	41.1
2002	56.5	76.0	42.3	56.8	75.1	43.3	64.7	100.3	41.2
2001	57.0	78.1	41.8	57.1	76.9	42.8	65.4	104.7	39.9
2000	57.8	79.7	42.1	57.8	78.4	43.0	67.2	107.1	41.1
1999	57.2	79.8	41.1	57.0	78.4	41.8	67.7	108.5	41.1
Of nasal cavity, middle ear and accessory sinuses (C30-C31)									
2005	0.2	0.2	0.1	0.2	0.2	0.1	0.2	0.3	*
2004	0.2	0.2	0.1	0.2	0.2	0.1	0.2	0.3	*
2003	0.1	0.2	0.1	0.2	0.2	0.1	0.2	0.3	*
2002	0.2	0.2	0.1	0.2	0.2	0.1	0.1	0.2	*
2001	0.2	0.2	0.1	0.2	0.2	0.1	0.2	0.3	*
2000	0.1	0.2	0.1	0.1	0.2	0.1	0.2	0.3	0.1
1999	0.2	0.2	0.1	0.2	0.2	0.1	0.2	0.2	*
Of larynx (C32)									
2005	1.2	2.2	0.5	1.2	2.1	0.5	2.3	4.6	0.8
2004	1.2	2.2	0.4	1.1	2.1	0.4	2.2	4.6	0.7
2003	1.3	2.4	0.5	1.2	2.2	0.5	2.5	5.1	0.7
2002	1.3	2.3	0.5	1.2	2.1	0.5	2.5	4.8	0.8
2001	1.3	2.5	0.5	1.2	2.3	0.5	2.5	5.0	0.9
2000	1.4	2.5	0.5	1.3	2.3	0.5	2.8	5.5	0.9
1999	1.4	2.5	0.5	1.3	2.3	0.5	2.6	5.1	0.8

Worktable 13. Age-adjusted death rates for 358 selected causes, by race and sex: United States, 1999-2005

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992) and year	All races 1/			White 2/			Black 2/		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
Of trachea, bronchus and lung (C33-C34)									
2005	52.6	69.0	40.5	53.1	68.7	41.5	58.4	86.4	40.0
2004	53.2	70.1	40.9	53.6	69.4	41.9	59.8	90.0	39.9
2003	54.1	71.7	41.3	54.5	71.1	42.3	60.8	92.4	40.2
2002	54.9	73.2	41.6	55.3	72.5	42.6	61.9	95.0	40.1
2001	55.3	75.2	41.0	55.6	74.2	42.1	62.5	99.1	38.7
2000	56.1	76.7	41.3	56.2	75.7	42.3	64.0	101.1	39.8
1999	55.5	76.9	40.2	55.4	75.6	41.0	64.8	103.0	39.9
Of thymus, heart, mediastinum and pleura (C37-C38)									
2005	0.1	0.2	0.1	0.1	0.2	0.1	0.2	0.2	0.2
2004	0.2	0.2	0.1	0.1	0.2	0.1	0.2	0.2	0.2
2003	0.2	0.2	0.1	0.1	0.2	0.1	0.2	0.2	0.1
2002	0.2	0.2	0.1	0.2	0.2	0.1	0.2	0.2	0.1
2001	0.2	0.2	0.1	0.2	0.2	0.2	0.2	0.2	0.2
2000	0.1	0.2	0.1	0.1	0.2	0.1	0.2	0.3	0.2
1999	0.1	0.2	0.1	0.1	0.2	0.1	0.2	0.2	0.2
Of other and ill-defined sites in respiratory system and intrathoracic organs (C39)									
2005	*	*	*	*	*	*	*	*	*
2004	*	*	*	*	*	*	*	*	*
2003	0.0	0.0	0.0	0.0	0.0	*	*	*	*
2002	0.0	0.0	0.0	0.0	0.0	*	*	*	*
2001	0.0	0.0	*	0.0	0.0	*	*	*	*
2000	0.0	0.0	*	0.0	0.0	*	*	*	*
1999	0.0	0.0	*	0.0	0.0	*	*	*	*

Worktable 13. Age-adjusted death rates for 358 selected causes, by race and sex: United States, 1999-2005

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992) and year	All races 1/			White 2/			Black 2/		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
Malignant neoplasms of bone and articular cartilage (C40-C41)									
2005	0.5	0.6	0.4	0.5	0.6	0.4	0.4	0.5	0.4
2004	0.4	0.5	0.4	0.4	0.5	0.3	0.5	0.6	0.4
2003	0.4	0.5	0.3	0.4	0.5	0.4	0.4	0.5	0.3
2002	0.4	0.5	0.3	0.4	0.5	0.3	0.4	0.5	0.4
2001	0.5	0.6	0.4	0.5	0.6	0.3	0.5	0.6	0.4
2000	0.4	0.6	0.3	0.4	0.5	0.3	0.4	0.6	0.3
1999	0.4	0.5	0.4	0.4	0.5	0.4	0.4	0.5	0.3
Melanoma and other malignant neoplasms of skin (C43-C44)									
2005	3.5	5.4	2.2	4.0	5.9	2.5	0.9	1.3	0.7
2004	3.5	5.3	2.1	3.9	5.9	2.4	0.9	1.3	0.6
2003	3.5	5.2	2.2	3.9	5.8	2.4	0.9	1.1	0.7
2002	3.4	5.1	2.1	3.8	5.7	2.4	0.9	1.5	0.6
2001	3.5	5.3	2.2	3.9	5.9	2.5	1.0	1.4	0.7
2000	3.5	5.2	2.2	3.8	5.8	2.5	1.1	1.5	0.8
1999	3.5	5.2	2.2	3.9	5.8	2.5	0.9	1.6	0.6
Malignant melanoma of skin (C43)									
2005	2.7	4.0	1.8	3.1	4.5	2.1	0.4	0.5	0.4
2004	2.7	3.9	1.7	3.0	4.5	1.9	0.4	0.5	0.4
2003	2.7	3.9	1.8	3.0	4.4	2.0	0.4	0.4	0.4
2002	2.6	3.8	1.7	2.9	4.2	2.0	0.4	0.5	0.3
2001	2.7	3.9	1.8	3.0	4.3	2.0	0.4	0.3	0.4
2000	2.7	3.8	1.8	3.0	4.3	2.0	0.6	0.6	0.5
1999	2.6	3.8	1.7	3.0	4.2	2.0	0.4	0.5	0.3

Worktable 13. Age-adjusted death rates for 358 selected causes, by race and sex: United States, 1999-2005

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992) and year	All races 1/			White 2/			Black 2/		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
Other malignant neoplasms of skin (C44)									
2005	0.8	1.4	0.4	0.8	1.5	0.4	0.5	0.8	0.3
2004	0.8	1.3	0.4	0.8	1.4	0.4	0.4	0.8	0.2
2003	0.8	1.4	0.4	0.9	1.5	0.4	0.5	0.7	0.2
2002	0.8	1.4	0.4	0.9	1.5	0.5	0.6	1.0	0.3
2001	0.9	1.4	0.5	0.9	1.5	0.5	0.6	1.0	0.2
2000	0.8	1.4	0.4	0.9	1.5	0.4	0.6	0.9	0.3
1999	0.8	1.4	0.4	0.9	1.5	0.4	0.5	1.0	0.2
Malignant neoplasms of mesothelial and soft tissue (C45-C49)									
2005	2.4	3.2	1.9	2.5	3.3	1.9	2.0	2.2	1.8
2004	2.4	3.2	1.8	2.5	3.4	1.9	2.0	2.3	1.8
2003	2.4	3.2	1.8	2.5	3.3	1.8	2.0	2.5	1.7
2002	2.4	3.2	1.8	2.4	3.4	1.8	2.0	2.4	1.7
2001	2.4	3.2	1.8	2.5	3.4	1.8	1.9	2.2	1.7
2000	2.5	3.3	1.8	2.5	3.5	1.9	2.1	2.6	1.9
1999	2.4	3.3	1.8	2.5	3.4	1.8	2.2	2.6	1.9
Mesothelioma (C45)									
2005	0.9	1.6	0.3	0.9	1.7	0.3	0.4	0.8	0.1
2004	0.8	1.6	0.3	0.9	1.7	0.3	0.3	0.7	*
2003	0.8	1.6	0.3	0.9	1.8	0.3	0.4	0.9	*
2002	0.9	1.7	0.2	0.9	1.8	0.3	0.4	0.8	*
2001	0.8	1.6	0.3	0.9	1.8	0.3	0.2	0.5	*
2000	0.9	1.7	0.3	0.9	1.8	0.3	0.4	0.8	0.1
1999	0.9	1.6	0.3	0.9	1.8	0.3	0.4	0.8	0.2

Worktable 13. Age-adjusted death rates for 358 selected causes, by race and sex: United States, 1999-2005

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992) and year	All races 1/			White 2/			Black 2/		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
Kaposi's sarcoma (C46)									
2005	0.0	0.0	*	0.0	0.0	*	*	*	*
2004	0.0	0.0	*	0.0	0.0	*	*	*	*
2003	0.0	0.0	0.0	0.0	0.0	0.0	*	*	*
2002	0.0	0.0	0.0	0.0	0.0	0.0	*	*	*
2001	0.0	0.0	*	0.0	0.0	*	*	*	*
2000	0.0	0.0	*	0.0	0.0	*	*	*	*
1999	0.0	0.0	*	0.0	0.0	*	*	*	*
Of other mesothelial and soft tissue (C47-C49)									
2005	1.5	1.5	1.5	1.5	1.6	1.5	1.6	1.4	1.7
2004	1.5	1.6	1.5	1.6	1.6	1.5	1.6	1.5	1.7
2003	1.5	1.5	1.5	1.5	1.5	1.5	1.6	1.6	1.6
2002	1.5	1.5	1.5	1.5	1.5	1.5	1.6	1.6	1.6
2001	1.5	1.5	1.5	1.5	1.6	1.5	1.6	1.7	1.6
2000	1.6	1.6	1.5	1.6	1.7	1.6	1.6	1.6	1.7
1999	1.6	1.6	1.5	1.5	1.6	1.5	1.7	1.8	1.7
Malignant neoplasm of breast (C50)									
2005	13.5	0.3	24.1	13.1	0.3	23.4	19.3	0.5	32.8
2004	13.8	0.3	24.4	13.4	0.3	23.9	19.0	0.5	32.2
2003	14.2	0.3	25.3	13.8	0.3	24.7	20.1	0.5	34.0
2002	14.5	0.3	25.6	14.1	0.3	25.0	20.2	0.6	34.0
2001	14.8	0.3	26.0	14.4	0.3	25.5	20.4	0.6	34.4
2000	15.3	0.4	26.8	14.9	0.4	26.3	20.5	0.6	34.5
1999	15.2	0.3	26.6	14.8	0.3	26.0	20.8	0.7	35.1

Worktable 13. Age-adjusted death rates for 358 selected causes, by race and sex: United States, 1999-2005

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992) and year	All races 1/			White 2/			Black 2/		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
Malignant neoplasms of female genital organs (C51-C58)									
2005	8.9	...	16.0	8.8	...	15.9	11.5	...	19.4
2004	9.0	...	16.2	8.9	...	16.0	11.8	...	19.9
2003	9.2	...	16.4	9.0	...	16.2	11.9	...	20.0
2002	9.4	...	16.7	9.2	...	16.5	12.2	...	20.5
2001	9.5	...	16.8	9.4	...	16.6	12.1	...	20.4
2000	9.5	...	16.8	9.3	...	16.6	12.5	...	20.8
1999	9.4	...	16.7	9.3	...	16.5	12.3	...	20.6
Of vulva and vagina (C51-C52)									
2005	0.4	...	0.7	0.4	...	0.7	0.3	...	0.6
2004	0.4	...	0.7	0.4	...	0.7	0.4	...	0.7
2003	0.4	...	0.7	0.4	...	0.7	0.4	...	0.6
2002	0.4	...	0.7	0.4	...	0.7	0.4	...	0.5
2001	0.4	...	0.7	0.4	...	0.7	0.5	...	0.8
2000	0.4	...	0.7	0.4	...	0.7	0.4	...	0.7
1999	0.4	...	0.7	0.4	...	0.7	0.4	...	0.7
Of cervix uteri (C53)									
2005	1.3	...	2.4	1.1	...	2.2	2.5	...	4.4
2004	1.3	...	2.4	1.1	...	2.2	2.6	...	4.5
2003	1.3	...	2.5	1.2	...	2.2	2.7	...	4.7
2002	1.3	...	2.6	1.2	...	2.3	2.9	...	4.9
2001	1.4	...	2.7	1.3	...	2.5	2.8	...	4.8
2000	1.5	...	2.8	1.3	...	2.5	3.2	...	5.4
1999	1.5	...	2.8	1.4	...	2.6	3.1	...	5.4

Worktable 13. Age-adjusted death rates for 358 selected causes, by race and sex: United States, 1999-2005

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992) and year	All races 1/			White 2/			Black 2/		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
Of corpus uteri and uterus, part unspecified (C54-C55)									
2005	2.3	...	4.1	2.2	...	3.8	4.2	...	7.1
2004	2.3	...	4.1	2.2	...	3.9	4.1	...	6.9
2003	2.3	...	4.1	2.2	...	3.9	4.2	...	7.0
2002	2.4	...	4.2	2.2	...	3.9	4.4	...	7.3
2001	2.4	...	4.2	2.3	...	4.0	4.2	...	7.0
2000	2.4	...	4.1	2.2	...	3.9	4.4	...	7.2
1999	2.4	...	4.1	2.2	...	3.9	4.2	...	6.8
Of ovary (C56)									
2005	4.8	...	8.6	5.0	...	9.0	4.3	...	7.2
2004	4.9	...	8.8	5.1	...	9.1	4.5	...	7.6
2003	5.0	...	8.8	5.1	...	9.2	4.4	...	7.4
2002	5.1	...	9.0	5.3	...	9.4	4.4	...	7.4
2001	5.1	...	9.0	5.2	...	9.3	4.5	...	7.5
2000	5.1	...	8.9	5.3	...	9.3	4.3	...	7.2
1999	5.0	...	8.8	5.1	...	9.1	4.4	...	7.3
Of other and unspecified female genital organs (C57)									
2005	0.1	...	0.1	0.1	...	0.1	0.1	...	0.1
2004	0.1	...	0.1	0.1	...	0.1	0.1	...	0.1
2003	0.1	...	0.2	0.1	...	0.2	0.1	...	0.3
2002	0.1	...	0.3	0.1	...	0.3	0.2	...	0.3
2001	0.1	...	0.2	0.1	...	0.2	0.1	...	0.2
2000	0.1	...	0.2	0.1	...	0.2	0.2	...	0.3
1999	0.1	...	0.3	0.1	...	0.3	0.2	...	0.3

Worktable 13. Age-adjusted death rates for 358 selected causes, by race and sex: United States, 1999-2005

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992) and year	All races 1/			White 2/			Black 2/		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
Of placenta (C58)									
2005	*	...	*	*	...	*	*	...	*
2004	0.0	...	0.0	*	...	*	*	...	*
2003	*	...	*	*	...	*	*	...	*
2002	*	...	*	*	...	*	*	...	*
2001	*	...	*	*	...	*	*	...	*
2000	*	...	*	*	...	*	*	...	*
1999	*	...	*	*	...	*	*	...	*
Malignant neoplasms of male genital organs (C60-C63)									
2005	9.7	25.0	...	9.0	23.1	...	19.3	53.8	...
2004	10.0	25.9	...	9.2	23.9	...	19.9	55.9	...
2003	10.3	26.9	...	9.6	24.9	...	20.6	58.0	...
2002	10.8	28.4	...	10.0	26.2	...	22.1	62.4	...
2001	11.1	29.6	...	10.2	27.1	...	23.2	66.5	...
2000	11.5	30.8	...	10.5	28.3	...	24.1	68.6	...
1999	11.9	31.8	...	10.9	29.3	...	24.4	69.5	...
Of prostate (C61)									
2005	9.5	24.5	...	8.8	22.6	...	19.1	53.3	...
2004	9.8	25.4	...	9.0	23.4	...	19.7	55.5	...
2003	10.1	26.5	...	9.3	24.4	...	20.4	57.4	...
2002	10.6	27.9	...	9.8	25.7	...	22.0	62.0	...
2001	10.9	29.1	...	10.0	26.6	...	23.0	66.1	...
2000	11.3	30.4	...	10.3	27.8	...	23.9	68.1	...
1999	11.6	31.3	...	10.7	28.7	...	24.2	69.0	...

Worktable 13. Age-adjusted death rates for 358 selected causes, by race and sex: United States, 1999-2005

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992) and year	All races 1/			White 2/			Black 2/		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
Of testis (C62)									
2005	0.1	0.3	...	0.1	0.3	...	0.1	0.2	...
2004	0.1	0.3	...	0.1	0.3	...	0.1	0.2	...
2003	0.1	0.2	...	0.1	0.3	...	0.1	0.2	...
2002	0.1	0.3	...	0.2	0.3	...	0.1	0.2	...
2001	0.1	0.2	...	0.1	0.3	...	0.1	0.2	...
2000	0.1	0.3	...	0.1	0.3	...	0.1	0.2	...
1999	0.1	0.3	...	0.2	0.3	...	0.1	0.1	...
Of penis and other and unspecified male genital organs (C60,C63)									
2005	0.1	0.2	...	0.1	0.2	...	0.1	0.3	...
2004	0.1	0.2	...	0.1	0.2	...	0.1	0.2	...
2003	0.1	0.2	...	0.1	0.2	...	0.1	0.4	...
2002	0.1	0.2	...	0.1	0.2	...	0.1	0.2	...
2001	0.1	0.2	...	0.1	0.2	...	0.1	0.3	...
2000	0.1	0.2	...	0.1	0.2	...	0.1	0.3	...
1999	0.1	0.2	...	0.1	0.2	...	0.1	0.3	...
Malignant neoplasms of urinary tract (C64-C68)									
2005	8.7	13.7	5.1	8.9	14.2	5.2	7.9	11.9	5.3
2004	8.7	13.7	5.1	9.0	14.3	5.2	7.7	10.9	5.6
2003	8.7	13.7	5.1	8.9	14.2	5.2	8.0	11.6	5.7
2002	8.9	13.9	5.3	9.1	14.4	5.4	8.1	11.4	6.1
2001	8.8	14.1	5.2	9.0	14.5	5.2	8.2	11.7	5.9
2000	8.8	14.1	5.3	9.0	14.5	5.3	8.0	12.1	5.6
1999	8.7	13.8	5.2	8.8	14.2	5.1	8.3	12.0	6.0

Worktable 13. Age-adjusted death rates for 358 selected causes, by race and sex: United States, 1999-2005

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992) and year	All races 1/			White 2/			Black 2/		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
Of kidney and renal pelvis (C64-C65)									
2005	4.1	5.9	2.7	4.2	6.0	2.8	4.0	6.0	2.6
2004	4.1	5.9	2.7	4.2	6.0	2.8	4.0	5.8	2.7
2003	4.2	6.0	2.7	4.3	6.2	2.8	4.1	6.0	2.7
2002	4.2	6.1	2.8	4.3	6.3	2.8	4.0	5.8	2.8
2001	4.3	6.2	2.8	4.3	6.3	2.8	4.2	6.4	2.8
2000	4.2	6.2	2.8	4.3	6.3	2.8	4.1	6.3	2.8
1999	4.1	5.9	2.7	4.1	5.9	2.7	4.1	6.1	2.8
Of bladder (C67)									
2005	4.3	7.5	2.2	4.5	7.9	2.2	3.7	5.7	2.6
2004	4.4	7.6	2.2	4.5	8.0	2.3	3.6	4.9	2.7
2003	4.3	7.3	2.2	4.4	7.7	2.2	3.7	5.4	2.7
2002	4.4	7.5	2.4	4.5	7.8	2.4	3.9	5.5	3.0
2001	4.3	7.5	2.3	4.5	7.9	2.2	3.7	5.1	2.9
2000	4.3	7.6	2.3	4.5	7.9	2.3	3.7	5.6	2.7
1999	4.4	7.6	2.3	4.5	7.9	2.3	4.0	5.8	3.0
Of other and unspecified sites within the urinary tract (C66,C68)									
2005	0.2	0.3	0.2	0.2	0.3	0.2	0.2	*	0.2
2004	0.2	0.3	0.2	0.2	0.3	0.1	0.2	*	0.2
2003	0.2	0.3	0.2	0.2	0.3	0.2	0.2	*	0.2
2002	0.2	0.3	0.2	0.2	0.3	0.2	0.2	*	0.2
2001	0.2	0.3	0.2	0.2	0.3	0.2	0.2	0.2	0.2
2000	0.2	0.3	0.2	0.2	0.3	0.2	0.2	*	0.2
1999	0.2	0.3	0.2	0.2	0.3	0.2	0.2	*	0.2

Worktable 13. Age-adjusted death rates for 358 selected causes, by race and sex: United States, 1999-2005

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992) and year	All races 1/			White 2/			Black 2/		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
Malignant neoplasm of eye and adnexa (C69)									
2005	0.1	0.1	0.1	0.1	0.1	0.1	*	*	*
2004	0.1	0.1	0.0	0.1	0.1	0.1	*	*	*
2003	0.1	0.1	0.1	0.1	0.1	0.1	*	*	*
2002	0.1	0.1	0.1	0.1	0.1	0.1	*	*	*
2001	0.1	0.1	0.0	0.1	0.1	0.1	*	*	*
2000	0.1	0.1	0.0	0.1	0.1	0.1	*	*	*
1999	0.1	0.1	0.1	0.1	0.1	0.1	*	*	*
Malignant neoplasms of meninges, brain and other parts of central nervous system (C70-C72)									
2005	4.3	5.3	3.5	4.7	5.7	3.8	2.5	3.2	2.0
2004	4.3	5.2	3.5	4.7	5.6	3.8	2.5	3.0	2.0
2003	4.4	5.3	3.6	4.7	5.7	3.9	2.7	3.4	2.1
2002	4.4	5.5	3.6	4.8	5.8	3.9	2.6	3.2	2.2
2001	4.5	5.5	3.6	4.8	5.9	3.9	2.6	3.1	2.2
2000	4.5	5.6	3.7	4.9	6.0	3.9	2.7	3.3	2.3
1999	4.6	5.6	3.8	5.0	6.0	4.1	2.7	3.3	2.3
Of brain (C71)									
2005	4.2	5.2	3.5	4.6	5.6	3.8	2.4	3.1	1.9
2004	4.2	5.1	3.4	4.6	5.5	3.8	2.4	2.9	2.0
2003	4.3	5.3	3.5	4.7	5.6	3.8	2.6	3.3	2.0
2002	4.3	5.3	3.5	4.7	5.7	3.8	2.5	3.0	2.1
2001	4.4	5.4	3.5	4.7	5.8	3.8	2.5	3.1	2.2
2000	4.5	5.5	3.6	4.8	5.9	3.9	2.6	3.2	2.2
1999	4.5	5.5	3.7	4.9	5.9	4.0	2.6	3.1	2.2

Worktable 13. Age-adjusted death rates for 358 selected causes, by race and sex: United States, 1999-2005

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992) and year	All races 1/			White 2/			Black 2/		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
Of other parts of central nervous system (C70,C72)									
2005	0.1	0.1	0.0	0.1	0.1	0.0	0.1	*	*
2004	0.1	0.1	0.1	0.1	0.1	0.1	0.1	*	*
2003	0.1	0.1	0.1	0.1	0.1	0.1	0.1	*	*
2002	0.1	0.1	0.1	0.1	0.1	0.1	0.1	*	*
2001	0.1	0.1	0.1	0.1	0.1	0.1	*	*	*
2000	0.1	0.1	0.1	0.1	0.1	0.1	0.1	*	*
1999	0.1	0.1	0.1	0.1	0.1	0.1	0.1	*	*
Malignant neoplasms of thyroid and other endocrine glands (C73-C75)									
2005	0.7	0.7	0.7	0.7	0.8	0.7	0.8	0.6	0.9
2004	0.7	0.7	0.7	0.7	0.7	0.7	0.6	0.6	0.7
2003	0.7	0.7	0.7	0.7	0.7	0.7	0.6	0.5	0.6
2002	0.7	0.7	0.7	0.7	0.7	0.7	0.7	0.6	0.7
2001	0.8	0.8	0.7	0.8	0.8	0.7	0.7	0.6	0.8
2000	0.8	0.8	0.7	0.7	0.8	0.7	0.8	0.6	0.9
1999	0.7	0.8	0.7	0.7	0.8	0.7	0.7	0.7	0.7
Other malignant neoplasms of other and unspecified sites (C76-C80,C97)									
2005	13.1	15.9	11.1	13.0	15.7	11.1	16.0	20.8	12.8
2004	13.3	16.1	11.2	13.2	15.8	11.1	16.8	22.1	13.2
2003	13.7	16.6	11.6	13.6	16.4	11.6	17.2	22.0	14.0
2002	14.1	17.2	11.9	13.9	16.9	11.7	18.2	23.1	15.0
2001	14.2	17.2	12.0	13.9	16.8	11.9	18.4	24.3	14.5
2000	14.7	17.8	12.5	14.5	17.4	12.4	18.7	24.8	14.7
1999	16.0	19.5	13.6	15.7	19.1	13.4	20.6	26.5	16.8

Worktable 13. Age-adjusted death rates for 358 selected causes, by race and sex: United States, 1999-2005

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992) and year	All races 1/			White 2/			Black 2/		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
Malignant neoplasms of lymphoid, hematopoietic and related tissue (C81-C96)									
2005	18.2	23.5	14.3	18.5	24.0	14.5	17.6	22.5	14.3
2004	18.4	23.6	14.6	18.6	24.0	14.7	18.9	23.6	15.8
2003	19.1	24.5	15.1	19.4	25.1	15.2	18.8	22.9	16.1
2002	19.5	25.1	15.5	19.9	25.6	15.7	19.4	24.4	16.1
2001	20.0	25.5	16.0	20.2	25.9	16.2	20.0	25.3	16.5
2000	20.4	26.0	16.5	20.7	26.4	16.7	19.9	25.0	16.6
1999	20.6	26.1	16.7	20.8	26.5	16.8	20.3	25.6	16.8
Hodgkin's disease (C81)									
2005	0.4	0.5	0.4	0.4	0.5	0.4	0.4	0.5	0.3
2004	0.4	0.5	0.3	0.4	0.6	0.3	0.4	0.5	0.4
2003	0.5	0.5	0.4	0.5	0.6	0.4	0.4	0.5	0.3
2002	0.5	0.6	0.4	0.5	0.6	0.4	0.4	0.6	0.3
2001	0.5	0.6	0.4	0.5	0.6	0.4	0.4	0.5	0.4
2000	0.5	0.6	0.4	0.5	0.6	0.4	0.3	0.4	0.3
1999	0.5	0.6	0.4	0.5	0.7	0.4	0.4	0.5	0.3
Of other lymphoid, hematopoietic and related tissue (C82-C85,C96)									
2005	6.9	8.8	5.5	7.2	9.1	5.8	4.7	6.0	3.8
2004	7.1	8.9	5.7	7.4	9.2	6.0	5.0	6.2	4.1
2003	7.3	9.3	5.9	7.7	9.7	6.2	4.9	5.8	4.3
2002	7.6	9.6	6.2	7.9	10.0	6.5	5.4	6.8	4.4
2001	7.9	10.0	6.4	8.2	10.3	6.7	5.6	7.3	4.5
2000	8.3	10.3	6.8	8.6	10.7	7.1	5.2	6.7	4.2
1999	8.4	10.4	6.9	8.7	10.7	7.2	5.6	7.5	4.4

Worktable 13. Age-adjusted death rates for 358 selected causes, by race and sex: United States, 1999-2005

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992) and year	All races 1/			White 2/			Black 2/		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
Non-Hodgkin's lymphoma (C82-C85)									
2005	6.9	8.7	5.5	7.2	9.1	5.8	4.7	6.0	3.8
2004	7.0	8.8	5.7	7.3	9.2	6.0	5.0	6.2	4.1
2003	7.3	9.3	5.9	7.7	9.7	6.1	4.9	5.8	4.2
2002	7.6	9.6	6.2	7.9	10.0	6.5	5.4	6.8	4.4
2001	7.9	9.9	6.4	8.2	10.2	6.7	5.6	7.2	4.5
2000	8.2	10.2	6.7	8.6	10.7	7.1	5.2	6.6	4.2
1999	8.3	10.3	6.9	8.7	10.7	7.2	5.6	7.4	4.4
Of other and unspecified lymphoid, hematopoietic and related tissue (C96)									
2005	0.0	0.0	0.0	0.0	0.0	*	*	*	*
2004	0.0	0.0	0.0	0.0	0.0	0.0	*	*	*
2003	0.0	0.0	0.0	0.0	0.0	0.0	*	*	*
2002	0.0	0.0	0.0	0.0	0.0	0.0	*	*	*
2001	0.0	0.0	0.0	0.0	0.0	0.0	*	*	*
2000	0.0	0.0	0.0	0.0	0.0	0.0	*	*	*
1999	0.0	0.0	0.0	0.0	0.0	0.0	*	*	*
Malignant immunoproliferative diseases (C88)									
2005	0.1	0.2	0.1	0.1	0.2	0.1	*	*	*
2004	0.1	0.1	0.1	0.1	0.2	0.1	*	*	*
2003	0.1	0.2	0.1	0.1	0.2	0.1	*	*	*
2002	0.1	0.2	0.1	0.1	0.2	0.1	*	*	*
2001	0.1	0.2	0.1	0.1	0.2	0.1	*	*	*
2000	0.1	0.2	0.1	0.1	0.2	0.1	*	*	*
1999	0.2	0.2	0.1	0.2	0.2	0.1	0.1	*	*

Worktable 13. Age-adjusted death rates for 358 selected causes, by race and sex: United States, 1999-2005

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992) and year	All races 1/			White 2/			Black 2/		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
Multiple myeloma and malignant plasma cell neoplasms (C90)									
2005	3.6	4.5	2.9	3.3	4.2	2.7	6.5	8.0	5.5
2004	3.6	4.4	3.0	3.3	4.2	2.7	6.8	8.2	6.0
2003	3.7	4.6	3.1	3.5	4.3	2.8	7.1	8.3	6.3
2002	3.8	4.8	3.2	3.6	4.5	2.9	6.9	8.2	6.0
2001	3.8	4.7	3.2	3.6	4.4	2.9	7.3	8.7	6.3
2000	3.9	4.7	3.3	3.6	4.4	3.0	7.6	8.8	6.8
1999	3.9	4.7	3.3	3.5	4.4	2.9	7.5	8.8	6.6
Leukemia (C91-C95)									
2005	7.1	9.6	5.4	7.4	10.0	5.6	5.9	7.8	4.7
2004	7.2	9.6	5.5	7.4	9.9	5.6	6.7	8.7	5.3
2003	7.4	9.9	5.6	7.6	10.2	5.7	6.4	8.3	5.2
2002	7.5	10.0	5.7	7.7	10.3	5.8	6.6	8.6	5.3
2001	7.6	10.1	5.9	7.8	10.4	6.0	6.7	8.7	5.4
2000	7.7	10.3	5.9	7.9	10.5	6.1	6.7	9.0	5.2
1999	7.6	10.2	6.0	7.8	10.4	6.1	6.7	8.7	5.4
Lymphoid leukemia (C91)									
2005	2.1	2.9	1.5	2.2	3.0	1.5	1.8	2.6	1.2
2004	2.1	2.9	1.5	2.1	3.0	1.5	1.9	2.9	1.3
2003	2.1	3.1	1.5	2.2	3.2	1.5	1.8	2.6	1.3
2002	2.2	3.1	1.6	2.3	3.2	1.6	1.9	2.7	1.4
2001	2.2	3.1	1.6	2.3	3.1	1.6	2.1	2.9	1.5
2000	2.2	3.1	1.6	2.3	3.2	1.6	1.9	2.8	1.3
1999	2.3	3.2	1.6	2.4	3.3	1.7	2.0	2.9	1.4

Worktable 13. Age-adjusted death rates for 358 selected causes, by race and sex: United States, 1999-2005

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992) and year	All races 1/			White 2/			Black 2/		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
Myeloid leukemia (C92)									
2005	3.3	4.3	2.6	3.4	4.5	2.6	2.6	3.1	2.2
2004	3.4	4.3	2.7	3.4	4.5	2.7	2.9	3.4	2.5
2003	3.4	4.4	2.7	3.5	4.6	2.7	2.9	3.7	2.4
2002	3.4	4.5	2.7	3.5	4.6	2.8	2.8	3.7	2.3
2001	3.5	4.5	2.8	3.6	4.7	2.9	2.9	3.6	2.4
2000	3.5	4.6	2.8	3.6	4.7	2.8	3.0	3.7	2.6
1999	3.4	4.3	2.7	3.4	4.4	2.8	2.8	3.6	2.3
Monocytic leukemia (C93)									
2005	0.0	0.1	0.0	0.0	0.1	0.0	*	*	*
2004	0.0	0.1	0.0	0.0	0.1	0.0	*	*	*
2003	0.0	0.1	0.0	0.1	0.1	0.0	*	*	*
2002	0.0	0.1	0.0	0.0	0.1	0.0	*	*	*
2001	0.1	0.1	0.0	0.1	0.1	0.0	*	*	*
2000	0.1	0.1	0.0	0.1	0.1	0.0	*	*	*
1999	0.0	0.1	0.0	0.1	0.1	0.0	*	*	*
Other and unspecified leukemia (C94-C95)									
2005	1.7	2.3	1.3	1.8	2.4	1.4	1.6	2.1	1.2
2004	1.7	2.4	1.3	1.8	2.4	1.3	1.8	2.3	1.5
2003	1.8	2.3	1.4	1.8	2.4	1.4	1.7	2.0	1.4
2002	1.8	2.4	1.4	1.8	2.4	1.4	1.8	2.2	1.5
2001	1.8	2.4	1.4	1.9	2.5	1.4	1.7	2.2	1.4
2000	1.9	2.5	1.5	1.9	2.5	1.6	1.7	2.4	1.3
1999	1.9	2.5	1.5	2.0	2.6	1.5	1.8	2.1	1.6

Worktable 13. Age-adjusted death rates for 358 selected causes, by race and sex: United States, 1999-2005

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992) and year	All races 1/			White 2/			Black 2/		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
In situ neoplasms (D00-D09)									
2005	*	*	*	*	*	*	*	*	*
2004	*	*	*	*	*	*	*	*	*
2003	*	*	*	*	*	*	*	*	*
2002	*	*	*	*	*	*	*	*	*
2001	0.0	*	*	0.0	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*
Carcinoma in situ of breast and genitourinary system (D05-D07)									
2005	*	*	*	*	*	*	*	*	*
2004	*	*	*	*	*	*	*	*	*
2003	*	*	*	*	*	*	*	*	*
2002	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*
Carcinoma in situ of other and unspecified sites (D00-D04,D09)									
2005	*	*	*	*	*	*	*	*	*
2004	*	*	*	*	*	*	*	*	*
2003	*	*	*	*	*	*	*	*	*
2002	*	*	*	*	*	*	*	*	*
2001	0.0	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*

Worktable 13. Age-adjusted death rates for 358 selected causes, by race and sex: United States, 1999-2005

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992) and year	All races 1/			White 2/			Black 2/		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
Benign neoplasms (D10-D36)									
2005	0.5	0.4	0.5	0.4	0.4	0.5	0.8	0.6	0.9
2004	0.5	0.5	0.6	0.5	0.4	0.5	0.9	0.9	0.9
2003	0.5	0.5	0.5	0.5	0.5	0.5	0.8	0.8	0.7
2002	0.5	0.5	0.6	0.5	0.5	0.5	0.8	0.9	0.9
2001	0.6	0.6	0.6	0.5	0.5	0.5	1.0	1.1	0.9
2000	0.6	0.5	0.6	0.5	0.5	0.6	0.9	0.8	1.0
1999	0.6	0.6	0.6	0.5	0.5	0.5	1.0	1.0	0.9
Of female genital organs (D25-D28)									
2005	0.0	...	0.0	*	...	*	0.1	...	0.1
2004	0.0	...	0.0	0.0	...	0.0	*	...	*
2003	0.0	...	0.0	0.0	...	0.0	*	...	*
2002	0.0	...	0.0	0.0	...	0.0	0.1	...	0.1
2001	0.0	...	0.0	0.0	...	0.0	*	...	*
2000	0.0	...	0.0	0.0	...	0.0	0.1	...	0.2
1999	0.0	...	0.0	*	...	*	*	...	*
Of eye, brain and other parts of central nervous system (D31-D33)									
2005	0.3	0.2	0.3	0.2	0.2	0.3	0.5	0.3	0.5
2004	0.3	0.3	0.3	0.3	0.3	0.3	0.5	0.5	0.5
2003	0.3	0.3	0.3	0.3	0.3	0.3	0.5	0.4	0.4
2002	0.3	0.3	0.4	0.3	0.3	0.4	0.5	0.4	0.5
2001	0.3	0.3	0.3	0.3	0.3	0.3	0.6	0.6	0.6
2000	0.3	0.3	0.4	0.3	0.3	0.4	0.5	0.4	0.5
1999	0.3	0.3	0.3	0.3	0.3	0.3	0.6	0.5	0.6

Worktable 13. Age-adjusted death rates for 358 selected causes, by race and sex: United States, 1999-2005

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992) and year	All races 1/			White 2/			Black 2/		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
Of other and unspecified sites (D10-D24,D29-D30,D34-D36)									
2005	0.2	0.2	0.2	0.1	0.2	0.2	0.3	0.3	0.3
2004	0.2	0.2	0.2	0.2	0.2	0.2	0.3	0.3	0.2
2003	0.2	0.2	0.2	0.2	0.2	0.2	0.3	0.4	0.2
2002	0.2	0.2	0.2	0.2	0.2	0.2	0.3	0.4	0.3
2001	0.2	0.3	0.2	0.2	0.3	0.2	0.3	0.5	0.2
2000	0.2	0.2	0.2	0.2	0.2	0.2	0.3	0.4	0.3
1999	0.2	0.2	0.2	0.2	0.2	0.2	0.3	0.4	0.3
Neoplasms of uncertain or unknown behavior of specified sites (D37-D47)									
2005	3.8	4.9	3.0	3.9	5.0	3.1	3.1	3.7	2.7
2004	3.8	4.9	3.0	3.9	5.1	3.1	3.2	4.0	2.7
2003	3.9	5.0	3.2	4.0	5.2	3.2	3.3	4.0	2.8
2002	3.8	5.0	3.1	4.0	5.2	3.2	3.3	4.1	2.9
2001	4.1	5.3	3.3	4.2	5.4	3.4	3.4	4.3	2.9
2000	4.1	5.3	3.3	4.2	5.4	3.4	3.4	4.4	2.9
1999	4.1	5.3	3.3	4.2	5.5	3.4	3.6	4.1	3.2
Neoplasms of uncertain or unknown behavior of unspecified sites (D48)									
2005	0.3	0.3	0.2	0.3	0.3	0.3	0.3	0.3	0.2
2004	0.3	0.2	0.2	0.3	0.2	0.2	0.3	0.3	0.3
2003	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2
2002	0.2	0.2	0.3	0.2	0.2	0.2	0.2	0.2	0.2
2001	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2
2000	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.3	0.2
1999	0.2	0.2	0.2	0.2	0.1	0.2	0.3	0.3	0.3

Worktable 13. Age-adjusted death rates for 358 selected causes, by race and sex: United States, 1999-2005

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992) and year	All races 1/			White 2/			Black 2/		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
III. Diseases of the blood and blood-forming organs and certain disorders involving the immune mechanism (D50-D89)									
2005	3.1	3.2	3.1	2.7	2.7	2.6	7.1	7.2	7.0
2004	3.1	3.2	3.1	2.7	2.8	2.6	7.0	7.2	6.8
2003	3.3	3.3	3.2	2.8	2.9	2.6	7.5	7.7	7.3
2002	3.4	3.4	3.3	2.8	2.9	2.8	7.5	7.7	7.4
2001	3.4	3.4	3.3	2.9	3.0	2.8	7.4	7.2	7.5
2000	3.4	3.4	3.3	2.9	3.0	2.8	7.4	7.2	7.5
1999	3.3	3.4	3.2	2.8	3.0	2.7	7.3	7.3	7.2
Anemias (D50-D64)									
2005	1.5	1.5	1.5	1.3	1.2	1.3	3.3	3.4	3.2
2004	1.5	1.5	1.5	1.3	1.3	1.3	3.3	3.5	3.1
2003	1.6	1.6	1.5	1.3	1.3	1.3	3.5	3.9	3.2
2002	1.6	1.6	1.6	1.3	1.3	1.4	3.7	3.9	3.5
2001	1.6	1.6	1.6	1.4	1.4	1.4	3.5	3.6	3.4
2000	1.6	1.6	1.6	1.4	1.4	1.4	3.4	3.4	3.5
1999	1.6	1.7	1.6	1.4	1.4	1.4	3.6	3.7	3.5
Nutritional anemias (D50-D53)									
2005	0.1	0.1	0.1	0.1	0.1	0.1	0.1	*	*
2004	0.1	0.1	0.1	0.1	0.1	0.1	0.1	*	*
2003	0.1	0.1	0.1	0.1	0.1	0.1	0.1	*	0.1
2002	0.1	0.1	0.1	0.1	0.1	0.1	0.1	*	*
2001	0.1	0.1	0.1	0.1	0.1	0.1	0.1	*	*
2000	0.1	0.1	0.1	0.1	0.1	0.1	0.2	*	0.2
1999	0.1	0.1	0.1	0.1	0.1	0.1	0.2	*	0.2

Worktable 13. Age-adjusted death rates for 358 selected causes, by race and sex: United States, 1999-2005

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992) and year	All races 1/			White 2/			Black 2/		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
Hemolytic anemias (D55-D59)									
2005	0.3	0.3	0.3	0.1	0.1	0.1	1.4	1.5	1.4
2004	0.3	0.3	0.3	0.1	0.1	0.1	1.5	1.6	1.4
2003	0.3	0.3	0.3	0.1	0.1	0.1	1.4	1.5	1.3
2002	0.3	0.3	0.3	0.1	0.1	0.1	1.5	1.5	1.5
2001	0.3	0.3	0.3	0.1	0.1	0.1	1.4	1.5	1.3
2000	0.3	0.3	0.3	0.1	0.1	0.1	1.4	1.3	1.5
1999	0.3	0.3	0.3	0.1	0.1	0.1	1.5	1.7	1.4
Aplastic anemias (D60-D61)									
2005	0.3	0.3	0.3	0.3	0.3	0.3	0.3	0.4	0.2
2004	0.3	0.4	0.3	0.3	0.3	0.3	0.3	0.4	0.3
2003	0.3	0.4	0.3	0.3	0.4	0.3	0.4	0.4	0.4
2002	0.3	0.4	0.3	0.3	0.4	0.3	0.4	0.5	0.4
2001	0.4	0.4	0.4	0.4	0.4	0.4	0.4	0.4	0.3
2000	0.4	0.4	0.4	0.4	0.4	0.4	0.5	0.6	0.4
1999	0.4	0.5	0.4	0.4	0.4	0.4	0.5	0.5	0.4
Acute posthemorrhagic and other anemias (D62,D64)									
2005	0.8	0.8	0.8	0.8	0.7	0.8	1.5	1.5	1.5
2004	0.8	0.8	0.8	0.8	0.7	0.8	1.4	1.5	1.3
2003	0.8	0.8	0.8	0.8	0.8	0.8	1.5	1.8	1.3
2002	0.8	0.8	0.9	0.8	0.7	0.8	1.7	1.8	1.5
2001	0.9	0.8	0.9	0.8	0.7	0.8	1.5	1.5	1.6
2000	0.8	0.8	0.8	0.8	0.7	0.8	1.4	1.4	1.4
1999	0.8	0.8	0.9	0.8	0.7	0.8	1.5	1.3	1.5

Worktable 13. Age-adjusted death rates for 358 selected causes, by race and sex: United States, 1999-2005

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992) and year	All races 1/			White 2/			Black 2/		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
Coagulation defects, purpura and other hemorrhagic conditions (D65-D69)									
2005	0.9	1.0	0.8	0.8	0.9	0.8	1.4	1.5	1.3
2004	0.9	0.9	0.8	0.8	0.9	0.8	1.4	1.5	1.3
2003	0.9	1.0	0.9	0.9	0.9	0.8	1.5	1.5	1.4
2002	0.9	1.0	0.9	0.9	1.0	0.8	1.4	1.4	1.4
2001	0.9	1.0	0.9	0.9	0.9	0.8	1.5	1.4	1.5
2000	1.0	1.0	0.9	0.9	1.0	0.9	1.5	1.5	1.5
1999	1.0	1.0	0.9	0.9	0.9	0.8	1.6	1.8	1.5
Other diseases of blood and blood-forming organs (D70-D76)									
2005	0.3	0.3	0.2	0.3	0.3	0.2	0.3	0.3	0.3
2004	0.3	0.3	0.3	0.3	0.3	0.3	0.2	0.3	0.2
2003	0.3	0.3	0.3	0.3	0.3	0.3	0.3	0.3	0.3
2002	0.3	0.4	0.3	0.3	0.4	0.3	0.3	0.4	0.3
2001	0.3	0.3	0.3	0.3	0.3	0.3	0.3	0.3	0.3
2000	0.3	0.4	0.3	0.3	0.3	0.3	0.3	0.4	0.3
1999	0.3	0.3	0.3	0.3	0.3	0.2	0.3	0.4	0.3
Certain disorders involving the immune mechanism (D80-D89)									
2005	0.5	0.5	0.5	0.3	0.3	0.3	2.1	2.0	2.2
2004	0.5	0.5	0.5	0.3	0.3	0.3	2.0	1.9	2.1
2003	0.5	0.5	0.5	0.3	0.3	0.3	2.2	2.0	2.4
2002	0.5	0.5	0.5	0.3	0.3	0.3	2.2	2.0	2.3
2001	0.5	0.5	0.5	0.3	0.3	0.3	2.1	1.9	2.4
2000	0.5	0.5	0.5	0.3	0.3	0.3	2.1	1.9	2.2
1999	0.4	0.4	0.4	0.3	0.3	0.2	1.7	1.5	1.9

Worktable 13. Age-adjusted death rates for 358 selected causes, by race and sex: United States, 1999-2005

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992) and year	All races 1/			White 2/			Black 2/		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
IV. Endocrine, nutritional and metabolic diseases (E00-E88)									
2005	33.7	37.8	30.3	31.4	35.8	27.7	59.2	63.1	56.0
2004	33.1	37.2	29.8	30.7	35.1	27.2	59.8	63.6	56.8
2003	34.2	38.0	31.1	31.7	36.0	28.3	61.3	63.1	59.3
2002	34.4	37.7	31.7	31.9	35.7	28.8	62.3	62.3	61.2
2001	34.3	37.1	31.8	31.8	35.0	29.0	61.7	63.0	60.1
2000	34.1	36.9	31.8	31.6	34.9	28.9	62.1	61.4	61.8
1999	34.0	36.6	31.8	31.3	34.4	28.8	62.2	63.0	61.3
Endocrine diseases (E00-E34)									
2005	25.6	29.1	22.7	23.4	27.2	20.4	48.3	52.0	45.3
2004	25.4	28.9	22.7	23.2	26.8	20.2	49.4	52.6	46.7
2003	26.3	29.6	23.7	24.0	27.6	21.1	50.5	51.9	48.9
2002	26.4	29.3	24.1	24.1	27.4	21.5	50.8	50.4	50.0
2001	26.3	28.9	24.2	24.0	27.0	21.7	50.6	51.2	49.5
2000	26.0	28.4	24.1	23.7	26.6	21.5	51.0	49.9	51.0
1999	26.0	28.5	24.1	23.5	26.5	21.3	51.2	50.8	51.0
Disorders of thyroid gland (E00-E07)									
2005	0.6	0.4	0.8	0.6	0.3	0.8	0.7	0.4	0.9
2004	0.6	0.4	0.7	0.6	0.4	0.7	0.8	0.5	0.9
2003	0.7	0.4	0.8	0.7	0.3	0.8	0.7	0.5	0.9
2002	0.7	0.4	0.8	0.7	0.4	0.8	0.7	0.4	0.9
2001	0.7	0.4	0.8	0.7	0.4	0.8	0.8	0.6	0.9
2000	0.6	0.4	0.8	0.6	0.3	0.8	0.8	0.6	1.0
1999	0.7	0.4	0.8	0.6	0.4	0.8	0.9	0.6	1.0

Worktable 13. Age-adjusted death rates for 358 selected causes, by race and sex: United States, 1999-2005

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992) and year	All races 1/			White 2/			Black 2/		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
Diabetes mellitus (E10-E14)									
2005	24.6	28.4	21.6	22.5	26.5	19.3	46.9	50.8	43.8
2004	24.5	28.2	21.7	22.3	26.2	19.2	48.0	51.3	45.3
2003	25.3	28.9	22.5	23.0	27.0	19.9	49.2	50.7	47.5
2002	25.4	28.6	23.0	23.1	26.8	20.3	49.5	49.4	48.6
2001	25.3	28.1	23.1	23.0	26.2	20.5	49.2	49.9	48.1
2000	25.0	27.8	23.0	22.8	26.0	20.4	49.5	48.7	49.4
1999	25.0	27.8	23.0	22.6	25.8	20.2	49.7	49.4	49.5
Disorders of pituitary gland (E22-E23)									
2005	0.1	0.1	0.1	0.1	0.1	0.0	0.1	*	*
2004	0.1	0.1	0.1	0.1	0.1	0.1	0.1	*	*
2003	0.1	0.1	0.1	0.1	0.1	0.0	0.1	*	*
2002	0.1	0.1	0.1	0.1	0.1	0.1	0.1	*	*
2001	0.1	0.1	0.1	0.1	0.1	0.1	0.1	*	*
2000	0.1	0.1	0.1	0.1	0.1	0.1	0.1	*	*
1999	0.1	0.1	0.1	0.1	0.1	0.1	0.2	0.2	0.1
Disorders of adrenal glands (E24-E27)									
2005	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.2	*
2004	0.1	0.1	0.1	0.1	0.1	0.1	0.1	*	0.1
2003	0.1	0.1	0.1	0.1	0.1	0.1	0.1	*	*
2002	0.1	0.1	0.1	0.1	0.1	0.1	0.1	*	*
2001	0.1	0.1	0.1	0.1	0.1	0.1	0.1	*	*
2000	0.1	0.1	0.1	0.1	0.1	0.1	0.1	*	0.2
1999	0.1	0.1	0.1	0.1	0.1	0.1	0.1	*	*

Worktable 13. Age-adjusted death rates for 358 selected causes, by race and sex: United States, 1999-2005

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992) and year	All races 1/			White 2/			Black 2/		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
Diseases of thymus (E32)									
2005	*	*	*	*	*	*	*	*	*
2004	*	*	*	*	*	*	*	*	*
2003	*	*	*	*	*	*	*	*	*
2002	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*
Other endocrine diseases (E15-E21,E28-E31,E34)									
2005	0.1	0.1	0.1	0.1	0.1	0.1	0.4	0.5	0.4
2004	0.1	0.2	0.1	0.1	0.1	0.1	0.4	0.5	0.3
2003	0.2	0.1	0.1	0.1	0.1	0.1	0.4	0.4	0.3
2002	0.1	0.1	0.1	0.1	0.1	0.1	0.3	0.3	0.4
2001	0.2	0.2	0.1	0.1	0.1	0.1	0.3	0.4	0.3
2000	0.1	0.1	0.2	0.1	0.1	0.1	0.4	0.3	0.4
1999	0.2	0.2	0.1	0.1	0.1	0.1	0.3	0.5	0.3
Nutritional diseases (E40-E68)									
2005	2.5	2.5	2.6	2.4	2.4	2.4	4.3	3.7	4.7
2004	2.4	2.3	2.4	2.2	2.2	2.2	4.2	3.8	4.5
2003	2.5	2.4	2.5	2.4	2.3	2.4	4.0	3.8	4.2
2002	2.6	2.5	2.6	2.4	2.4	2.4	4.5	4.0	4.7
2001	2.4	2.2	2.5	2.3	2.1	2.3	4.1	4.0	4.3
2000	2.6	2.5	2.6	2.5	2.4	2.5	4.3	4.2	4.4
1999	2.5	2.5	2.5	2.4	2.3	2.4	4.4	4.5	4.3

Worktable 13. Age-adjusted death rates for 358 selected causes, by race and sex: United States, 1999-2005

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992) and year	All races 1/			White 2/			Black 2/		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
Malnutrition (E40-E46)									
2005	1.0	0.9	1.0	0.9	0.8	1.0	1.6	1.5	1.6
2004	0.9	0.9	1.0	0.9	0.8	0.9	1.6	1.8	1.5
2003	1.1	1.0	1.1	1.0	0.9	1.1	1.8	2.1	1.7
2002	1.2	1.1	1.2	1.1	1.1	1.2	2.0	2.0	2.0
2001	1.2	1.1	1.3	1.2	1.0	1.2	1.9	2.2	1.7
2000	1.4	1.3	1.5	1.4	1.2	1.4	2.2	2.5	2.1
1999	1.5	1.4	1.5	1.4	1.3	1.4	2.5	2.9	2.2
Nutritional marasmus (E41)									
2005	0.0	0.0	0.0	0.0	0.0	0.0	*	*	*
2004	0.0	0.0	0.0	0.0	0.0	0.0	*	*	*
2003	0.0	0.0	0.1	0.0	0.0	0.1	*	*	*
2002	0.1	0.0	0.1	0.1	0.0	0.1	0.1	*	*
2001	0.1	0.1	0.1	0.1	0.1	0.1	0.1	*	*
2000	0.1	0.1	0.1	0.1	0.1	0.1	0.1	*	*
1999	0.1	0.1	0.1	0.1	0.1	0.1	0.1	*	*
Other and unspecified malnutrition (E40,E42-E46)									
2005	0.9	0.9	1.0	0.9	0.8	0.9	1.6	1.5	1.6
2004	0.9	0.9	0.9	0.8	0.8	0.9	1.6	1.8	1.4
2003	1.0	0.9	1.0	1.0	0.9	1.0	1.8	2.0	1.6
2002	1.1	1.1	1.1	1.1	1.0	1.1	1.9	1.9	1.9
2001	1.1	1.0	1.2	1.1	1.0	1.2	1.8	2.1	1.6
2000	1.3	1.2	1.4	1.3	1.2	1.3	2.1	2.3	2.0
1999	1.4	1.3	1.4	1.3	1.2	1.3	2.3	2.8	2.0

Worktable 13. Age-adjusted death rates for 358 selected causes, by race and sex: United States, 1999-2005

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992) and year	All races 1/			White 2/			Black 2/		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
Other nutritional deficiencies (E50-E64)									
2005	0.1	0.0	0.1	0.1	0.0	0.1	*	*	*
2004	0.1	0.0	0.0	0.1	0.0	0.1	*	*	*
2003	0.1	0.1	0.0	0.1	0.1	0.1	*	*	*
2002	0.1	0.1	0.1	0.1	0.1	0.1	0.1	*	*
2001	0.1	0.1	0.1	0.1	0.1	0.1	0.1	*	*
2000	0.1	0.1	0.1	0.1	0.1	0.1	0.1	*	0.1
1999	0.1	0.1	0.1	0.1	0.1	0.1	0.2	*	0.2
Obesity and other hyperalimentation (E65-E68)									
2005	1.5	1.5	1.5	1.4	1.5	1.4	2.6	2.1	3.0
2004	1.4	1.4	1.4	1.3	1.3	1.3	2.6	2.0	3.0
2003	1.3	1.3	1.4	1.3	1.3	1.3	2.1	1.7	2.5
2002	1.3	1.2	1.3	1.2	1.2	1.1	2.3	1.8	2.7
2001	1.1	1.0	1.2	1.0	1.0	1.0	2.1	1.6	2.5
2000	1.1	1.1	1.1	1.0	1.0	0.9	1.9	1.6	2.2
1999	0.9	0.9	1.0	0.9	0.9	0.8	1.7	1.3	2.0
Metabolic disorders (E70-E88)									
2005	5.6	6.2	5.0	5.6	6.3	5.0	6.6	7.4	6.0
2004	5.3	6.0	4.7	5.3	6.0	4.7	6.2	7.1	5.5
2003	5.4	6.0	4.8	5.4	6.0	4.8	6.7	7.4	6.1
2002	5.4	5.9	5.0	5.4	5.9	4.9	7.1	8.0	6.5
2001	5.5	6.0	5.0	5.5	6.0	5.0	7.0	7.9	6.3
2000	5.5	5.9	5.0	5.5	5.9	5.0	6.8	7.3	6.3
1999	5.5	5.7	5.1	5.4	5.6	5.1	6.6	7.7	6.0

Worktable 13. Age-adjusted death rates for 358 selected causes, by race and sex: United States, 1999-2005

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992) and year	All races 1/			White 2/			Black 2/		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
Cystic fibrosis (E84)									
2005	0.1	0.1	0.1	0.2	0.2	0.2	*	*	*
2004	0.2	0.2	0.2	0.2	0.2	0.2	*	*	*
2003	0.2	0.2	0.1	0.2	0.2	0.2	0.0	*	*
2002	0.2	0.2	0.2	0.2	0.2	0.2	0.1	*	*
2001	0.2	0.2	0.2	0.2	0.2	0.2	*	*	*
2000	0.2	0.2	0.2	0.2	0.2	0.2	*	*	*
1999	0.2	0.2	0.2	0.2	0.2	0.2	*	*	*
Volume depletion, disorders of fluid, electrolyte and acid-base balance (E86-E87)									
2005	2.2	2.0	2.2	2.1	2.0	2.2	3.3	3.5	3.1
2004	2.2	2.0	2.2	2.1	1.9	2.2	3.2	3.5	2.9
2003	2.4	2.2	2.4	2.3	2.1	2.4	3.7	3.9	3.6
2002	2.5	2.3	2.6	2.4	2.1	2.6	4.0	4.2	3.8
2001	2.7	2.5	2.8	2.6	2.3	2.8	4.4	4.5	4.2
2000	2.8	2.7	2.8	2.7	2.6	2.7	4.2	4.5	4.0
1999	3.0	2.8	3.1	2.9	2.6	3.1	4.3	5.1	3.7
Other metabolic disorders (E70-E83,E85,E88)									
2005	3.3	4.1	2.6	3.3	4.2	2.6	3.3	3.9	2.8
2004	3.0	3.8	2.3	3.1	3.9	2.4	3.0	3.5	2.6
2003	2.9	3.7	2.3	2.9	3.8	2.3	2.9	3.4	2.5
2002	2.8	3.5	2.2	2.8	3.5	2.2	3.1	3.7	2.6
2001	2.6	3.4	2.0	2.7	3.4	2.1	2.6	3.3	2.1
2000	2.5	3.1	2.0	2.5	3.2	2.0	2.5	2.7	2.3
1999	2.3	2.8	1.9	2.3	2.8	1.9	2.4	2.5	2.2

Worktable 13. Age-adjusted death rates for 358 selected causes, by race and sex: United States, 1999-2005

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992) and year	All races 1/			White 2/			Black 2/		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
V. Mental and behavioral disorders (F01-F99)									
2005	22.8	22.0	22.3	23.1	22.0	22.7	24.6	26.4	22.5
2004	20.3	19.9	19.5	20.5	19.7	19.9	21.7	25.1	18.6
2003	20.2	19.6	19.6	20.4	19.5	20.0	21.3	24.0	18.5
2002	19.2	18.8	18.4	19.3	18.6	18.8	20.1	23.5	17.0
2001	18.0	17.8	17.1	18.1	17.6	17.4	18.9	22.7	15.6
2000	16.7	16.8	15.7	16.8	16.5	16.0	17.9	22.0	14.4
1999	15.5	16.0	14.3	15.6	15.8	14.5	15.9	20.4	12.3
Organic dementia (F01,F03)									
2005	17.8	15.0	19.2	18.1	15.1	19.5	18.6	17.1	19.0
2004	15.3	12.9	16.4	15.6	13.0	16.8	15.5	15.1	15.4
2003	15.3	12.8	16.4	15.6	12.9	16.8	15.0	14.3	15.0
2002	14.3	12.2	15.3	14.6	12.3	15.7	13.5	13.2	13.4
2001	13.2	11.4	14.0	13.5	11.5	14.3	12.0	11.7	12.0
2000	12.0	10.3	12.6	12.2	10.3	12.9	11.1	11.3	10.8
1999	10.6	9.3	11.1	10.9	9.5	11.5	8.7	8.9	8.4
Other and unspecified organic mental disorders (F04-F09)									
2005	0.5	0.5	0.6	0.6	0.5	0.6	0.6	0.5	0.5
2004	0.6	0.6	0.6	0.6	0.6	0.6	0.5	0.5	0.5
2003	0.6	0.6	0.6	0.6	0.6	0.6	0.6	0.7	0.5
2002	0.7	0.6	0.7	0.7	0.6	0.7	0.6	0.5	0.6
2001	0.7	0.7	0.8	0.8	0.7	0.8	0.8	0.8	0.7
2000	0.8	0.7	0.9	0.8	0.7	0.9	0.8	0.8	0.7
1999	0.9	0.8	0.9	0.9	0.8	1.0	0.9	1.1	0.8

Worktable 13. Age-adjusted death rates for 358 selected causes, by race and sex: United States, 1999-2005

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992) and year	All races 1/			White 2/			Black 2/		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
Mental and behavioral disorders due to psychoactive substance use (F10-F19)									
2005	3.7	5.7	1.8	3.6	5.6	1.8	4.8	7.9	2.3
2004	3.6	5.6	1.7	3.5	5.4	1.7	5.0	8.6	2.1
2003	3.4	5.4	1.6	3.3	5.1	1.6	4.9	8.3	2.2
2002	3.3	5.2	1.5	3.1	4.9	1.5	5.2	8.9	2.3
2001	3.2	5.0	1.5	3.0	4.6	1.4	5.4	9.4	2.3
2000	3.1	5.0	1.4	2.9	4.6	1.3	5.4	9.3	2.3
1999	3.1	5.1	1.4	2.9	4.6	1.2	5.7	9.7	2.6
Mental and behavioral disorders due to use of alcohol (F10)									
2005	2.4	3.9	1.0	2.4	3.9	1.0	2.7	4.9	1.1
2004	2.4	3.9	1.0	2.4	3.9	1.0	3.0	5.5	1.0
2003	2.4	3.8	1.0	2.3	3.7	1.0	3.0	5.5	1.1
2002	2.3	3.9	1.0	2.3	3.7	0.9	3.3	6.1	1.2
2001	2.3	3.8	1.0	2.2	3.5	0.9	3.7	6.7	1.3
2000	2.2	3.7	0.9	2.1	3.5	0.8	3.6	6.6	1.3
1999	2.2	3.7	0.9	2.1	3.4	0.8	3.8	6.8	1.5
Mental and behavioral disorders due to other psychoactive substance use (F11-F19)									
2005	1.3	1.8	0.8	1.2	1.7	0.8	2.0	3.0	1.3
2004	1.2	1.6	0.7	1.1	1.5	0.7	2.0	3.1	1.1
2003	1.1	1.5	0.6	1.0	1.4	0.6	1.9	2.8	1.2
2002	0.9	1.3	0.6	0.9	1.2	0.5	1.9	2.8	1.1
2001	0.9	1.2	0.5	0.8	1.1	0.5	1.8	2.6	1.1
2000	0.9	1.2	0.6	0.8	1.1	0.5	1.8	2.7	1.1
1999	0.9	1.3	0.5	0.8	1.1	0.4	1.9	2.9	1.1

Worktable 13. Age-adjusted death rates for 358 selected causes, by race and sex: United States, 1999-2005

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992) and year	All races 1/			White 2/			Black 2/		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
Schizophrenia, schizotypal and delusional disorders (F20-F29)									
2005	0.1	0.2	0.2	0.1	0.1	0.2	0.2	0.3	0.2
2004	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.4	0.2
2003	0.2	0.1	0.2	0.2	0.1	0.2	0.2	0.2	0.2
2002	0.2	0.2	0.2	0.2	0.1	0.2	0.3	0.3	0.2
2001	0.2	0.1	0.2	0.2	0.1	0.2	0.2	0.2	0.2
2000	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2
1999	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2
Mood (affective) disorders (F30-F39)									
2005	0.2	0.2	0.3	0.3	0.2	0.3	0.1	*	*
2004	0.3	0.2	0.3	0.3	0.2	0.3	0.1	*	0.1
2003	0.3	0.3	0.3	0.3	0.3	0.4	0.2	*	0.2
2002	0.3	0.2	0.3	0.3	0.2	0.4	0.2	*	0.1
2001	0.3	0.3	0.3	0.3	0.3	0.3	0.1	0.2	0.1
2000	0.3	0.3	0.3	0.3	0.3	0.3	0.1	*	0.1
1999	0.3	0.3	0.3	0.3	0.3	0.3	0.1	*	*
Neurotic, stress-related and somatoform disorders (F40-F48)									
2005	0.1	0.1	0.0	0.1	0.1	0.0	*	*	*
2004	0.0	0.0	0.0	0.0	0.0	0.0	*	*	*
2003	0.0	0.0	0.0	0.0	0.0	0.0	*	*	*
2002	0.0	0.0	0.0	0.0	0.0	0.0	*	*	*
2001	0.0	0.0	0.0	0.0	0.0	0.0	*	*	*
2000	0.0	0.0	0.0	0.0	0.0	0.0	*	*	*
1999	0.0	0.0	0.0	0.0	0.0	0.0	*	*	*

Worktable 13. Age-adjusted death rates for 358 selected causes, by race and sex: United States, 1999-2005

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992) and year	All races 1/			White 2/			Black 2/		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
Mental retardation (F70-F79)									
2005	0.2	0.2	0.1	0.2	0.2	0.1	0.2	0.2	0.1
2004	0.2	0.2	0.1	0.2	0.2	0.1	0.2	0.2	0.1
2003	0.2	0.2	0.1	0.2	0.2	0.1	0.2	0.2	0.1
2002	0.2	0.2	0.1	0.2	0.2	0.1	0.2	0.2	*
2001	0.1	0.1	0.1	0.1	0.1	0.1	0.2	0.2	0.1
2000	0.1	0.2	0.1	0.1	0.2	0.1	0.1	0.2	0.1
1999	0.2	0.2	0.1	0.2	0.2	0.1	0.2	0.2	0.2
Other and unspecified mental and behavioral disorders (F50-F69,F80-F99)									
2005	0.1	0.1	0.2	0.1	0.1	0.1	0.2	0.2	0.2
2004	0.2	0.1	0.2	0.2	0.1	0.2	0.1	*	0.1
2003	0.2	0.1	0.2	0.2	0.1	0.2	0.2	*	0.2
2002	0.2	0.1	0.2	0.2	0.1	0.2	0.1	*	0.1
2001	0.1	0.1	0.2	0.1	0.1	0.2	0.1	*	*
2000	0.1	0.1	0.2	0.1	0.1	0.2	0.1	*	*
1999	0.1	0.1	0.2	0.1	0.1	0.2	0.1	*	*
VI. Diseases of the nervous system (G00-G98)									
2005	39.0	39.3	38.2	40.2	40.4	39.3	34.4	34.5	33.7
2004	37.1	37.5	36.3	38.2	38.7	37.3	32.7	32.1	32.5
2003	37.2	37.8	36.1	38.3	38.9	37.3	31.1	32.2	30.0
2002	35.6	36.5	34.4	36.7	37.6	35.6	29.3	30.5	27.9
2001	34.2	35.8	32.7	35.2	36.8	33.7	28.6	30.2	27.3
2000	33.0	34.8	31.5	34.0	35.8	32.4	27.0	28.8	25.7
1999	30.9	33.2	29.1	31.8	34.2	29.9	25.0	26.6	23.7

Worktable 13. Age-adjusted death rates for 358 selected causes, by race and sex: United States, 1999-2005

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992) and year	All races 1/			White 2/			Black 2/		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
Meningitis (G00,G03)									
2005	0.2	0.3	0.2	0.2	0.2	0.2	0.4	0.5	0.4
2004	0.3	0.3	0.2	0.2	0.2	0.2	0.5	0.5	0.4
2003	0.3	0.3	0.2	0.2	0.3	0.2	0.6	0.5	0.6
2002	0.2	0.3	0.2	0.2	0.3	0.2	0.5	0.7	0.4
2001	0.3	0.3	0.2	0.2	0.3	0.2	0.5	0.8	0.4
2000	0.3	0.3	0.2	0.2	0.3	0.2	0.6	0.8	0.6
1999	0.3	0.3	0.3	0.3	0.3	0.2	0.7	0.9	0.6
Other inflammatory diseases of central nervous system (G04-G09)									
2005	0.2	0.3	0.2	0.2	0.2	0.2	0.4	0.5	0.3
2004	0.2	0.3	0.2	0.2	0.3	0.2	0.3	0.4	0.3
2003	0.3	0.3	0.2	0.2	0.3	0.2	0.3	0.4	0.3
2002	0.2	0.3	0.2	0.2	0.3	0.2	0.4	0.5	0.3
2001	0.2	0.2	0.2	0.2	0.2	0.2	0.4	0.4	0.4
2000	0.2	0.3	0.2	0.2	0.3	0.2	0.3	0.4	0.3
1999	0.2	0.3	0.2	0.2	0.2	0.2	0.3	0.4	0.3
Parkinson's disease (G20-G21)									
2005	6.4	9.8	4.4	6.8	10.4	4.6	3.0	4.9	1.9
2004	6.1	9.3	4.1	6.4	9.8	4.4	2.7	4.3	1.9
2003	6.2	9.3	4.2	6.5	9.8	4.5	2.7	4.4	1.8
2002	5.9	9.0	4.1	6.3	9.5	4.3	2.5	4.0	1.7
2001	5.9	9.1	4.0	6.2	9.7	4.3	2.5	3.9	1.8
2000	5.7	8.8	3.9	6.0	9.3	4.2	2.5	4.2	1.6
1999	5.4	8.4	3.7	5.7	8.9	3.9	2.4	3.7	1.7

Worktable 13. Age-adjusted death rates for 358 selected causes, by race and sex: United States, 1999-2005

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992) and year	All races 1/			White 2/			Black 2/		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
Alzheimer's disease (G30)									
2005	22.9	18.5	25.1	23.7	19.0	26.1	19.4	16.4	20.7
2004	21.8	17.7	23.8	22.5	18.3	24.7	18.5	15.2	19.9
2003	21.4	17.5	23.3	22.2	18.2	24.3	16.6	14.7	17.3
2002	20.2	16.6	22.0	21.1	17.3	22.9	15.2	13.2	15.9
2001	19.1	16.1	20.5	19.8	16.7	21.3	14.6	13.1	15.1
2000	18.1	15.2	19.3	18.8	15.8	20.1	13.0	11.4	13.7
1999	16.5	14.4	17.4	17.2	15.0	18.1	11.3	10.2	11.8
Multiple sclerosis (G35)									
2005	1.1	0.8	1.4	1.2	0.8	1.5	1.1	0.8	1.4
2004	1.1	0.8	1.4	1.1	0.8	1.4	1.0	0.7	1.3
2003	1.1	0.8	1.3	1.1	0.9	1.4	1.1	0.8	1.3
2002	1.1	0.8	1.3	1.1	0.8	1.4	1.1	0.6	1.5
2001	1.1	0.8	1.3	1.1	0.8	1.4	1.0	0.6	1.3
2000	1.0	0.8	1.2	1.1	0.8	1.3	1.0	0.7	1.3
1999	1.0	0.7	1.2	1.0	0.7	1.2	0.8	0.6	1.0
Epilepsy (G40-G41)									
2005	0.5	0.5	0.5	0.4	0.5	0.4	1.0	1.1	0.9
2004	0.5	0.5	0.4	0.4	0.5	0.3	0.9	1.1	0.7
2003	0.5	0.5	0.4	0.4	0.5	0.4	1.0	1.3	0.8
2002	0.5	0.6	0.4	0.4	0.5	0.4	1.0	1.3	0.8
2001	0.4	0.5	0.4	0.4	0.5	0.4	0.9	1.2	0.8
2000	0.5	0.5	0.4	0.4	0.5	0.4	1.0	1.3	0.9
1999	0.5	0.6	0.4	0.4	0.5	0.4	1.1	1.3	0.9

Worktable 13. Age-adjusted death rates for 358 selected causes, by race and sex: United States, 1999-2005

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992) and year	All races 1/			White 2/			Black 2/		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
Transient cerebral ischemic attacks and related syndromes (G45)									
2005	0.1	0.0	0.1	0.1	0.0	0.1	*	*	*
2004	0.1	0.0	0.1	0.1	0.0	0.1	*	*	*
2003	0.1	0.1	0.1	0.1	0.1	0.1	*	*	*
2002	0.1	0.1	0.1	0.1	0.1	0.1	*	*	*
2001	0.1	0.1	0.1	0.1	0.1	0.1	0.1	*	*
2000	0.1	0.1	0.1	0.1	0.1	0.1	*	*	*
1999	0.1	0.1	0.1	0.1	0.1	0.1	*	*	*
Infantile cerebral palsy (G80)									
2005	0.5	0.5	0.4	0.4	0.5	0.4	0.7	0.8	0.6
2004	0.5	0.5	0.4	0.5	0.5	0.4	0.7	0.8	0.6
2003	0.5	0.5	0.4	0.4	0.5	0.4	0.6	0.8	0.6
2002	0.4	0.5	0.4	0.4	0.5	0.3	0.6	0.7	0.5
2001	0.4	0.5	0.3	0.4	0.4	0.3	0.5	0.6	0.4
2000	0.4	0.4	0.3	0.4	0.4	0.3	0.6	0.6	0.5
1999	0.4	0.4	0.3	0.3	0.4	0.3	0.5	0.6	0.4
All other diseases of nervous system (G10-G12,G23-G25,G31,G36-G37,G43-G44,G47-G72,G81-G98)									
2005	7.1	8.6	6.0	7.1	8.7	5.9	8.4	9.6	7.5
2004	6.8	8.1	5.7	6.8	8.2	5.7	8.0	9.1	7.3
2003	7.0	8.4	5.9	7.0	8.5	5.9	8.0	9.1	7.2
2002	6.9	8.3	5.8	6.9	8.3	5.8	7.9	9.4	6.8
2001	6.7	8.1	5.7	6.7	8.1	5.6	8.1	9.6	7.2
2000	6.8	8.3	5.7	6.8	8.3	5.6	7.9	9.5	6.8
1999	6.6	8.1	5.6	6.6	8.1	5.5	7.7	8.8	7.0

Worktable 13. Age-adjusted death rates for 358 selected causes, by race and sex: United States, 1999-2005

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992) and year	All races 1/			White 2/			Black 2/		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
VII. Diseases of the eye and adnexa (H00-H57)									
2005	0.0	*	0.0	0.0	*	0.0	*	*	*
2004	0.0	*	0.0	0.0	*	0.0	*	*	*
2003	0.0	*	0.0	0.0	*	0.0	*	*	*
2002	0.0	*	0.0	0.0	*	0.0	*	*	*
2001	0.0	*	0.0	0.0	*	0.0	*	*	*
2000	0.0	0.0	0.0	0.0	0.0	0.0	*	*	*
1999	0.0	*	0.0	0.0	*	0.0	*	*	*
VIII. Diseases of the ear and mastoid process (H60-H93)									
2005	0.0	0.0	0.0	0.0	0.0	*	*	*	*
2004	0.0	0.0	0.0	0.0	0.0	0.0	*	*	*
2003	0.0	0.0	0.0	0.0	0.0	0.0	*	*	*
2002	0.0	0.0	0.0	0.0	0.0	0.0	*	*	*
2001	0.0	0.0	0.0	0.0	0.0	0.0	*	*	*
2000	0.0	0.0	0.0	0.0	0.0	0.0	*	*	*
1999	0.0	0.0	0.0	0.0	0.0	0.0	*	*	*
IX. Diseases of the circulatory system (I00-I99)									
2005	278.8	331.1	237.1	272.5	324.7	230.4	370.6	438.4	319.8
2004	288.0	341.7	245.3	281.2	335.1	238.0	384.6	454.0	333.6
2003	307.7	365.0	262.3	300.4	357.9	254.5	409.0	482.3	355.1
2002	319.0	378.8	272.1	312.0	372.2	264.3	418.9	490.6	366.4
2001	328.2	389.8	280.4	320.8	382.6	272.3	430.2	508.3	374.9
2000	341.3	409.0	290.0	334.0	402.3	281.7	441.6	520.7	386.1
1999	350.8	420.5	297.9	343.3	413.2	289.8	450.0	534.9	390.8

Worktable 13. Age-adjusted death rates for 358 selected causes, by race and sex: United States, 1999-2005

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992) and year	All races 1/			White 2/			Black 2/		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
Acute rheumatic fever and chronic rheumatic heart diseases (I00-I09)									
2005	1.1	0.8	1.3	1.1	0.8	1.3	0.8	0.7	0.8
2004	1.1	0.8	1.3	1.1	0.8	1.3	0.8	0.6	1.0
2003	1.2	1.0	1.4	1.3	1.0	1.4	1.1	0.9	1.2
2002	1.2	0.9	1.5	1.3	0.9	1.5	0.9	0.8	1.0
2001	1.2	0.9	1.5	1.2	0.9	1.5	1.1	0.7	1.3
2000	1.3	1.0	1.5	1.3	1.0	1.6	1.0	0.7	1.2
1999	1.3	0.9	1.6	1.4	0.9	1.6	1.0	0.8	1.2
Acute rheumatic fever (I00-I02)									
2005	0.0	*	*	0.0	*	*	*	*	*
2004	0.0	*	0.0	0.0	*	*	*	*	*
2003	0.0	*	0.0	0.0	*	*	*	*	*
2002	0.0	*	0.0	0.0	*	0.0	*	*	*
2001	0.0	0.0	0.0	0.0	0.0	0.0	*	*	*
2000	0.0	*	0.0	0.0	*	*	*	*	*
1999	0.0	*	*	0.0	*	*	*	*	*
Chronic rheumatic heart diseases (I05-I09)									
2005	1.1	0.8	1.3	1.1	0.8	1.3	0.7	0.7	0.8
2004	1.1	0.8	1.3	1.1	0.8	1.3	0.8	0.6	1.0
2003	1.2	1.0	1.4	1.2	1.0	1.4	1.0	0.9	1.2
2002	1.2	0.9	1.5	1.2	0.9	1.5	0.9	0.8	1.0
2001	1.2	0.9	1.4	1.2	0.9	1.4	1.0	0.7	1.3
2000	1.3	0.9	1.5	1.3	0.9	1.6	1.0	0.7	1.2
1999	1.3	0.9	1.6	1.4	0.9	1.6	1.0	0.7	1.2

Worktable 13. Age-adjusted death rates for 358 selected causes, by race and sex: United States, 1999-2005

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992) and year	All races 1/			White 2/			Black 2/		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
Rheumatic mitral valve diseases (I05)									
2005	0.5	0.4	0.6	0.5	0.4	0.6	0.4	0.3	0.4
2004	0.5	0.4	0.6	0.5	0.4	0.6	0.4	0.3	0.5
2003	0.6	0.5	0.7	0.6	0.5	0.7	0.6	0.5	0.6
2002	0.6	0.4	0.8	0.6	0.4	0.8	0.5	0.4	0.5
2001	0.7	0.4	0.8	0.7	0.5	0.8	0.6	0.4	0.8
2000	0.7	0.5	0.9	0.8	0.5	0.9	0.5	0.4	0.6
1999	0.7	0.5	0.9	0.8	0.5	0.9	0.6	0.5	0.7
Rheumatic aortic valve diseases (I06)									
2005	0.0	0.0	0.0	0.0	0.0	0.0	*	*	*
2004	0.0	0.0	0.0	0.0	0.0	0.0	*	*	*
2003	0.0	0.0	0.0	0.0	0.0	0.0	*	*	*
2002	0.0	0.0	0.1	0.0	0.0	0.0	*	*	*
2001	0.0	0.0	0.0	0.0	0.0	0.0	*	*	*
2000	0.0	*	*	0.0	*	*	*	*	*
1999	0.0	*	0.0	0.0	*	0.0	*	*	*
Disorders of both mitral and aortic valves (I08.0)									
2005	0.3	0.2	0.3	0.3	0.2	0.3	0.1	*	*
2004	0.2	0.2	0.2	0.3	0.2	0.3	0.1	0.2	0.1
2003	0.3	0.2	0.3	0.3	0.3	0.3	0.2	0.2	0.2
2002	0.2	0.2	0.2	0.2	0.2	0.2	0.1	*	*
2001	0.1	0.1	0.1	0.1	0.1	0.1	0.1	*	*
2000	0.1	0.1	0.1	0.1	0.1	0.1	0.1	*	*
1999	0.1	0.1	0.1	0.1	0.1	0.1	0.1	*	*

Worktable 13. Age-adjusted death rates for 358 selected causes, by race and sex: United States, 1999-2005

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992) and year	All races 1/			White 2/			Black 2/		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
Other chronic rheumatic heart diseases (I07,I08.1-I08.9,I09)									
2005	0.3	0.2	0.4	0.3	0.2	0.4	0.3	0.2	0.3
2004	0.3	0.2	0.4	0.3	0.2	0.4	0.2	*	0.3
2003	0.3	0.2	0.4	0.3	0.2	0.4	0.3	0.2	0.4
2002	0.4	0.2	0.4	0.4	0.3	0.4	0.3	0.2	0.3
2001	0.4	0.3	0.5	0.4	0.3	0.5	0.3	0.2	0.4
2000	0.4	0.3	0.5	0.4	0.3	0.5	0.4	0.2	0.5
1999	0.4	0.3	0.5	0.5	0.3	0.5	0.3	0.2	0.3
Hypertensive diseases (I10-I13)									
2005	18.5	18.9	17.4	15.8	15.8	15.0	45.7	52.2	40.3
2004	18.1	18.7	17.0	15.4	15.7	14.5	45.7	51.0	40.9
2003	18.1	18.3	17.2	15.3	15.3	14.6	46.0	51.1	41.7
2002	17.1	17.4	16.3	14.4	14.4	13.8	44.8	49.6	40.6
2001	16.5	16.6	15.8	13.9	13.7	13.4	43.0	47.8	38.9
2000	16.2	16.3	15.5	13.5	13.4	12.9	43.3	47.2	39.9
1999	15.8	15.9	15.2	13.1	12.9	12.6	43.1	47.5	39.4
Essential (primary) hypertension (I10)									
2005	4.8	4.1	5.0	4.4	3.7	4.7	8.6	8.9	8.2
2004	4.4	4.0	4.6	4.1	3.6	4.2	8.2	7.9	8.1
2003	4.4	3.7	4.6	4.0	3.3	4.3	8.4	8.6	8.1
2002	4.0	3.5	4.2	3.7	3.1	3.9	7.7	7.5	7.5
2001	3.9	3.5	4.0	3.6	3.1	3.7	7.6	8.2	7.1
2000	3.7	3.4	3.8	3.4	3.0	3.5	7.3	7.6	6.9
1999	3.6	3.3	3.7	3.3	2.9	3.4	7.2	7.4	7.0

Worktable 13. Age-adjusted death rates for 358 selected causes, by race and sex: United States, 1999-2005

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992) and year	All races 1/			White 2/			Black 2/		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
Hypertensive heart disease (I11)									
2005	9.4	10.1	8.5	7.9	8.3	7.2	24.1	28.4	20.4
2004	9.5	10.1	8.5	8.0	8.4	7.2	24.2	28.4	20.6
2003	9.6	10.1	8.8	8.0	8.3	7.3	25.0	28.7	21.9
2002	9.1	9.5	8.3	7.6	7.8	7.0	24.4	28.0	21.2
2001	8.7	8.9	8.1	7.3	7.3	6.8	23.2	26.3	20.5
2000	8.6	8.7	8.1	7.1	7.1	6.7	23.7	26.3	21.2
1999	8.3	8.3	7.9	6.9	6.7	6.6	22.9	25.7	20.5
Hypertensive renal disease (I12)									
2005	3.3	3.6	3.0	2.6	2.9	2.4	9.8	11.0	8.9
2004	3.2	3.5	3.0	2.5	2.8	2.3	10.0	11.0	9.2
2003	3.1	3.4	2.8	2.4	2.7	2.2	9.2	10.0	8.5
2002	3.0	3.3	2.8	2.4	2.7	2.2	9.3	10.2	8.6
2001	2.9	3.2	2.7	2.3	2.5	2.1	8.9	9.7	8.3
2000	2.8	3.1	2.6	2.2	2.5	2.0	9.0	9.5	8.7
1999	2.6	2.9	2.4	2.0	2.3	1.8	8.8	9.8	8.0
Hypertensive heart and renal disease (I13)									
2005	1.0	1.1	0.9	0.8	0.9	0.8	3.2	3.8	2.8
2004	1.0	1.1	0.9	0.8	0.8	0.7	3.2	3.6	2.9
2003	1.1	1.1	1.0	0.8	0.9	0.7	3.4	3.7	3.1
2002	1.0	1.0	0.9	0.7	0.8	0.7	3.5	3.9	3.2
2001	1.0	1.1	1.0	0.8	0.8	0.7	3.3	3.6	3.0
2000	1.0	1.1	0.9	0.8	0.9	0.7	3.3	3.8	3.1
1999	1.2	1.4	1.1	0.9	1.0	0.9	4.2	4.5	3.9

Worktable 13. Age-adjusted death rates for 358 selected causes, by race and sex: United States, 1999-2005

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992) and year	All races 1/			White 2/			Black 2/		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
Ischemic heart diseases (I20-I25)									
2005	144.4	187.4	111.7	143.8	187.7	110.0	171.3	213.9	140.9
2004	150.2	194.1	116.7	149.2	194.2	114.7	179.8	223.9	148.7
2003	162.9	209.9	127.2	161.7	209.6	125.0	195.0	243.3	160.9
2002	170.8	220.4	133.6	169.8	220.5	131.2	203.0	250.6	169.7
2001	177.8	228.5	139.9	176.5	228.4	137.4	211.6	262.0	176.7
2000	186.8	241.4	146.5	185.6	241.6	143.9	218.3	269.5	183.2
1999	194.6	251.2	152.9	193.4	251.2	150.5	224.9	278.7	188.0
Acute myocardial infarction (I21-I22)									
2005	49.1	63.6	37.8	49.1	64.1	37.2	57.6	69.7	48.8
2004	52.2	67.8	40.2	52.2	68.3	39.5	62.0	74.8	52.7
2003	57.9	75.0	44.8	57.8	75.3	44.0	68.6	84.3	57.6
2002	62.1	80.6	48.1	61.9	81.0	47.1	73.9	89.5	63.2
2001	65.4	84.9	50.9	65.1	85.4	49.9	76.8	93.1	66.0
2000	69.9	91.3	54.1	69.6	91.8	53.1	81.0	98.9	69.2
1999	73.2	96.1	56.3	73.1	96.8	55.4	83.1	102.1	70.5
Other acute ischemic heart diseases (I24)									
2005	1.2	1.4	0.9	1.1	1.3	0.9	2.0	2.6	1.5
2004	0.8	1.0	0.6	0.8	1.0	0.6	1.4	2.1	1.0
2003	1.1	1.3	0.9	1.0	1.2	0.8	1.6	2.4	1.1
2002	1.2	1.5	0.9	1.1	1.4	0.9	1.7	2.3	1.3
2001	1.2	1.5	0.9	1.1	1.4	0.9	1.8	2.3	1.3
2000	1.2	1.5	0.9	1.2	1.5	0.9	1.8	2.5	1.3
1999	1.2	1.5	0.9	1.1	1.4	0.9	1.8	2.4	1.3

Worktable 13. Age-adjusted death rates for 358 selected causes, by race and sex: United States, 1999-2005

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992) and year	All races 1/			White 2/			Black 2/		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
Other forms of chronic ischemic heart disease (I20,I25)									
2005	94.2	122.4	73.0	93.6	122.3	72.0	111.8	141.6	90.5
2004	97.1	125.3	75.8	96.3	125.0	74.6	116.4	146.9	95.0
2003	103.8	133.5	81.5	102.9	133.0	80.1	124.8	156.7	102.2
2002	107.6	138.3	84.5	106.8	138.0	83.2	127.5	158.9	105.2
2001	111.2	142.1	88.1	110.2	141.6	86.7	133.0	166.7	109.3
2000	115.7	148.6	91.5	114.8	148.3	90.0	135.4	168.1	112.6
1999	120.2	153.6	95.6	119.2	153.0	94.2	140.0	174.2	116.2
Atherosclerotic cardiovascular disease, so described (I25.0)									
2005	20.3	26.6	15.0	19.1	25.0	14.0	34.5	46.9	25.3
2004	21.1	27.5	15.7	19.9	25.9	14.7	36.1	48.9	27.0
2003	22.7	29.6	17.1	21.4	27.8	16.1	38.2	52.7	28.0
2002	23.5	30.4	17.8	22.1	28.5	16.7	40.3	54.8	30.0
2001	24.1	30.8	18.5	22.5	28.8	17.3	41.8	56.0	31.7
2000	25.2	32.3	19.4	23.6	30.2	18.2	43.5	58.7	32.9
1999	26.4	33.5	20.6	24.7	31.2	19.3	45.8	61.5	34.9
All other forms of chronic ischemic heart disease (I20,I25.1-I25.9)									
2005	73.9	95.8	58.0	74.5	97.3	57.9	77.3	94.7	65.2
2004	76.0	97.8	60.1	76.5	99.1	59.9	80.3	98.0	68.0
2003	81.1	104.0	64.4	81.5	105.2	64.0	86.6	104.0	74.2
2002	84.1	108.0	66.7	84.7	109.5	66.4	87.2	104.1	75.2
2001	87.1	111.3	69.6	87.6	112.8	69.4	91.1	110.6	77.6
2000	90.5	116.3	72.0	91.2	118.1	71.8	91.9	109.5	79.7
1999	93.8	120.1	75.0	94.5	121.8	74.9	94.1	112.7	81.3

Worktable 13. Age-adjusted death rates for 358 selected causes, by race and sex: United States, 1999-2005

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992) and year	All races 1/			White 2/			Black 2/		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
Other heart diseases (I26-I51)									
2005	55.2	61.5	49.9	54.1	60.2	49.0	71.9	83.0	63.4
2004	55.2	61.7	49.9	54.1	60.3	49.0	72.5	85.6	63.2
2003	57.6	64.5	52.0	56.4	63.1	50.9	75.7	87.7	66.7
2002	58.5	65.5	52.8	57.3	64.1	51.7	76.6	87.8	68.1
2001	59.1	66.1	53.4	57.7	64.4	52.3	77.8	91.8	68.2
2000	60.0	67.9	53.8	58.6	66.3	52.6	78.4	92.2	69.0
1999	60.9	69.2	54.5	59.3	67.2	53.1	81.2	97.5	70.1
Pulmonary heart disease and diseases of pulmonary circulation (I26-I28)									
2005	4.4	4.1	4.7	4.2	3.9	4.4	7.5	7.1	7.8
2004	4.4	4.0	4.7	4.1	3.8	4.4	7.6	7.3	7.8
2003	4.8	4.6	5.0	4.5	4.3	4.7	8.2	8.1	8.4
2002	4.8	4.4	5.0	4.5	4.2	4.7	8.1	7.2	8.8
2001	4.7	4.5	4.9	4.5	4.2	4.6	8.0	7.9	8.3
2000	4.7	4.5	4.9	4.5	4.3	4.6	8.0	7.6	8.3
1999	4.8	4.7	5.0	4.5	4.5	4.6	8.7	8.4	8.9
Pulmonary embolism (I26)									
2005	2.7	2.7	2.7	2.6	2.6	2.6	4.8	5.0	4.6
2004	2.7	2.7	2.7	2.6	2.5	2.6	4.9	5.2	4.7
2003	3.0	3.0	2.9	2.8	2.8	2.7	5.4	5.8	5.2
2002	3.0	2.9	3.1	2.8	2.7	2.9	5.2	5.0	5.4
2001	3.1	3.0	3.1	2.9	2.8	2.9	5.3	5.7	5.2
2000	3.0	3.0	3.0	2.9	2.9	2.8	5.4	5.7	5.4
1999	3.3	3.3	3.3	3.1	3.1	3.1	6.1	6.5	5.9

Worktable 13. Age-adjusted death rates for 358 selected causes, by race and sex: United States, 1999-2005

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992) and year	All races 1/			White 2/			Black 2/		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
Other pulmonary heart disease and diseases of pulmonary circulation (I27-I28)									
2005	1.7	1.4	1.9	1.6	1.4	1.8	2.7	2.1	3.2
2004	1.7	1.4	1.9	1.6	1.3	1.8	2.7	2.1	3.1
2003	1.8	1.6	2.0	1.7	1.5	1.9	2.8	2.2	3.2
2002	1.8	1.5	1.9	1.7	1.5	1.8	2.9	2.2	3.3
2001	1.7	1.5	1.8	1.6	1.4	1.7	2.7	2.3	3.0
2000	1.7	1.5	1.8	1.6	1.4	1.7	2.5	2.0	2.9
1999	1.5	1.4	1.6	1.5	1.4	1.5	2.6	1.9	3.0
Other non-pulmonary forms of heart disease (I30-I51)									
2005	50.7	57.3	45.2	49.9	56.3	44.6	64.4	75.9	55.6
2004	50.8	57.7	45.3	50.0	56.5	44.6	65.0	78.3	55.4
2003	52.8	60.0	47.0	51.9	58.8	46.2	67.4	79.7	58.3
2002	53.8	61.1	47.8	52.8	59.9	47.0	68.5	80.6	59.4
2001	54.4	61.6	48.5	53.3	60.2	47.7	69.8	83.9	60.0
2000	55.2	63.4	48.9	54.2	62.0	48.0	70.5	84.5	60.7
1999	56.1	64.5	49.6	54.9	62.7	48.6	72.5	89.1	61.2
Acute and subacute endocarditis (I33)									
2005	0.4	0.5	0.3	0.4	0.5	0.3	0.7	0.9	0.6
2004	0.4	0.5	0.3	0.4	0.5	0.3	0.7	0.8	0.6
2003	0.4	0.5	0.3	0.4	0.5	0.3	0.8	0.9	0.6
2002	0.4	0.5	0.3	0.4	0.5	0.3	0.7	0.8	0.6
2001	0.4	0.5	0.3	0.4	0.5	0.3	0.8	1.1	0.6
2000	0.4	0.5	0.3	0.4	0.5	0.3	0.7	0.9	0.7
1999	0.4	0.5	0.3	0.4	0.5	0.3	0.8	0.9	0.6

Worktable 13. Age-adjusted death rates for 358 selected causes, by race and sex: United States, 1999-2005

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992) and year	All races 1/			White 2/			Black 2/		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
Diseases of pericardium and acute myocarditis (I30-I31,I40)									
2005	0.3	0.3	0.3	0.3	0.3	0.2	0.5	0.5	0.5
2004	0.3	0.3	0.2	0.2	0.3	0.2	0.5	0.6	0.4
2003	0.3	0.3	0.3	0.3	0.3	0.2	0.5	0.5	0.5
2002	0.3	0.3	0.3	0.3	0.3	0.3	0.5	0.5	0.4
2001	0.3	0.3	0.3	0.3	0.3	0.3	0.5	0.5	0.4
2000	0.3	0.3	0.2	0.3	0.3	0.2	0.4	0.5	0.4
1999	0.3	0.4	0.3	0.3	0.3	0.3	0.5	0.5	0.5
Other diseases of endocardium (I34-I38)									
2005	6.7	6.8	6.6	7.0	7.1	6.8	4.5	4.5	4.4
2004	6.7	6.9	6.5	7.0	7.1	6.7	4.9	5.1	4.6
2003	6.8	7.0	6.6	7.0	7.3	6.8	4.9	5.0	4.7
2002	6.9	7.1	6.7	7.2	7.4	7.0	4.8	4.6	4.8
2001	6.9	7.2	6.6	7.2	7.5	6.9	4.9	5.3	4.6
2000	7.2	7.5	6.9	7.4	7.7	7.1	5.3	5.4	5.1
1999	7.2	7.6	6.9	7.4	7.8	7.1	5.3	5.4	5.1
Nonrheumatic mitral valve disorders (I34)									
2005	0.8	0.7	0.9	0.9	0.8	0.9	0.7	0.4	0.8
2004	0.8	0.8	0.9	0.9	0.8	0.9	0.7	0.6	0.8
2003	0.9	0.8	0.9	0.9	0.8	0.9	0.6	0.6	0.7
2002	1.0	0.8	1.0	1.0	0.9	1.1	0.6	0.6	0.7
2001	1.0	0.9	1.1	1.0	0.9	1.1	0.7	0.5	0.8
2000	1.0	1.0	1.1	1.1	1.0	1.1	0.8	0.6	0.9
1999	1.1	1.0	1.1	1.1	1.0	1.2	0.8	0.6	0.9

Worktable 13. Age-adjusted death rates for 358 selected causes, by race and sex: United States, 1999-2005

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992) and year	All races 1/			White 2/			Black 2/		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
Nonrheumatic aortic valve disorders (I35)									
2005	4.2	4.4	4.0	4.5	4.7	4.2	2.2	2.4	2.0
2004	4.2	4.4	4.0	4.4	4.6	4.2	2.4	2.6	2.2
2003	4.2	4.6	4.0	4.5	4.8	4.2	2.4	2.4	2.3
2002	4.3	4.6	4.0	4.5	4.9	4.2	2.4	2.4	2.4
2001	4.3	4.7	4.0	4.5	4.9	4.2	2.5	3.0	2.1
2000	4.5	4.9	4.2	4.7	5.1	4.4	2.7	2.9	2.5
1999	4.5	5.0	4.2	4.7	5.2	4.3	2.7	3.1	2.5
All other diseases of endocardium (I36-I38)									
2005	1.7	1.6	1.6	1.7	1.6	1.7	1.6	1.7	1.6
2004	1.7	1.7	1.6	1.7	1.7	1.6	1.8	1.9	1.7
2003	1.7	1.6	1.7	1.7	1.6	1.7	1.9	2.0	1.8
2002	1.6	1.6	1.7	1.6	1.6	1.7	1.7	1.6	1.7
2001	1.6	1.7	1.6	1.6	1.7	1.6	1.7	1.8	1.6
2000	1.6	1.6	1.6	1.6	1.6	1.6	1.8	2.0	1.7
1999	1.6	1.6	1.6	1.6	1.6	1.6	1.8	1.8	1.8
Cardiomyopathy (I42)									
2005	8.3	11.7	5.8	7.7	10.9	5.3	14.7	20.6	10.3
2004	8.5	11.9	6.1	7.9	11.1	5.6	14.9	20.6	10.7
2003	9.1	12.6	6.5	8.5	11.8	6.0	15.9	22.1	11.5
2002	9.6	13.4	6.8	8.9	12.5	6.2	16.6	23.0	12.0
2001	9.5	13.1	6.8	8.7	12.2	6.2	16.9	23.3	12.5
2000	9.8	13.8	7.0	9.0	12.7	6.4	17.8	24.9	12.8
1999	10.0	13.9	7.1	9.1	12.7	6.5	18.2	25.9	12.9

Worktable 13. Age-adjusted death rates for 358 selected causes, by race and sex: United States, 1999-2005

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992) and year	All races 1/			White 2/			Black 2/		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
Conduction disorders and cardiac dysrhythmias (I44-I49)									
2005	12.5	13.7	11.3	12.1	13.3	11.0	17.1	20.0	15.1
2004	12.5	13.8	11.3	12.1	13.3	10.9	17.6	21.1	15.3
2003	13.2	14.6	12.0	12.9	14.1	11.6	18.4	21.3	16.1
2002	13.6	15.1	12.2	13.2	14.6	11.9	18.8	21.7	16.7
2001	13.6	15.1	12.3	13.2	14.5	12.0	19.1	22.8	16.6
2000	13.9	15.6	12.4	13.6	15.2	12.1	18.8	21.8	16.6
1999	14.7	16.7	13.0	14.3	16.2	12.8	20.0	24.6	17.1
Heart failure (I50)									
2005	18.9	19.9	18.0	18.9	20.0	18.1	21.7	23.0	20.4
2004	18.9	19.9	18.0	19.0	20.0	18.1	20.9	23.4	19.3
2003	19.4	20.6	18.4	19.5	20.7	18.4	21.8	23.6	20.5
2002	19.5	20.4	18.6	19.5	20.5	18.6	21.8	23.3	20.5
2001	20.2	21.2	19.2	20.2	21.2	19.3	22.1	24.4	20.6
2000	20.3	21.5	19.2	20.3	21.6	19.3	22.0	24.2	20.7
1999	20.3	21.4	19.3	20.3	21.4	19.3	22.5	25.3	20.9
Congestive heart failure (I50.0)									
2005	17.6	18.5	16.8	17.7	18.6	17.0	19.2	20.2	18.2
2004	17.6	18.5	16.8	17.7	18.6	17.0	18.8	20.8	17.4
2003	18.1	19.1	17.3	18.2	19.3	17.4	19.5	20.9	18.5
2002	18.0	18.8	17.3	18.1	19.0	17.4	19.6	20.7	18.5
2001	18.7	19.5	18.0	18.8	19.6	18.1	19.9	21.7	18.8
2000	18.8	19.8	17.9	18.9	20.0	18.0	19.7	21.3	18.8
1999	18.8	19.8	17.9	18.8	19.7	18.0	20.3	22.7	18.9

Worktable 13. Age-adjusted death rates for 358 selected causes, by race and sex: United States, 1999-2005

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992) and year	All races 1/			White 2/			Black 2/		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
Other and unspecified heart failure (I50.1-I50.9)									
2005	1.3	1.5	1.2	1.2	1.4	1.1	2.5	2.8	2.2
2004	1.3	1.5	1.1	1.2	1.4	1.1	2.2	2.6	1.9
2003	1.3	1.5	1.1	1.2	1.4	1.1	2.3	2.7	2.0
2002	1.4	1.6	1.3	1.3	1.5	1.2	2.2	2.6	2.0
2001	1.5	1.7	1.3	1.4	1.6	1.2	2.2	2.7	1.8
2000	1.5	1.7	1.4	1.4	1.6	1.3	2.3	2.8	1.9
1999	1.5	1.7	1.3	1.4	1.6	1.3	2.2	2.5	2.0
Myocarditis, unspecified and myocardial degeneration (I51.4-I51.5)									
2005	0.2	0.3	0.2	0.2	0.2	0.2	0.4	0.5	0.4
2004	0.2	0.2	0.2	0.2	0.2	0.2	0.4	0.5	0.4
2003	0.2	0.2	0.2	0.2	0.2	0.2	0.4	0.4	0.3
2002	0.3	0.3	0.3	0.3	0.3	0.3	0.5	0.6	0.4
2001	0.3	0.3	0.2	0.2	0.2	0.2	0.5	0.6	0.4
2000	0.3	0.3	0.2	0.2	0.3	0.2	0.4	0.4	0.3
1999	0.3	0.3	0.2	0.2	0.3	0.2	0.4	0.5	0.3
All other and ill-defined forms of heart disease (I50.0-I51.3,I51.6-I51.9)									
2005	3.4	4.2	2.7	3.3	4.0	2.7	4.8	5.9	3.9
2004	3.3	4.0	2.7	3.2	3.9	2.6	5.1	6.4	4.1
2003	3.3	4.0	2.7	3.2	3.9	2.6	4.9	6.0	4.0
2002	3.2	4.0	2.6	3.1	3.8	2.5	4.9	6.1	3.9
2001	3.2	3.8	2.6	3.0	3.7	2.5	5.0	6.0	4.2
2000	3.2	3.9	2.6	3.0	3.7	2.4	5.0	6.5	4.0
1999	3.0	3.7	2.4	2.8	3.5	2.2	4.7	6.0	3.8

Worktable 13. Age-adjusted death rates for 358 selected causes, by race and sex: United States, 1999-2005

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992) and year	All races 1/			White 2/			Black 2/		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
Cerebrovascular diseases (I60-I69)									
2005	46.6	46.9	45.6	44.7	44.7	44.0	65.2	70.5	60.7
2004	50.0	50.4	48.9	48.0	48.1	47.2	69.9	74.9	65.5
2003	53.5	54.1	52.3	51.4	51.7	50.5	74.3	79.5	69.8
2002	56.2	56.5	55.2	54.2	54.2	53.4	76.3	81.7	71.8
2001	57.9	59.0	56.4	55.8	56.5	54.5	78.8	85.4	73.7
2000	60.9	62.4	59.1	58.8	59.8	57.3	81.9	89.6	76.2
1999	61.6	63.2	59.8	59.6	60.8	58.0	81.8	89.6	76.2
Subarachnoid hemorrhage (I60)									
2005	2.0	1.5	2.3	1.9	1.5	2.2	2.7	2.1	3.2
2004	2.0	1.6	2.4	1.9	1.5	2.3	2.7	2.0	3.2
2003	2.0	1.6	2.4	1.9	1.5	2.2	2.9	2.3	3.4
2002	2.1	1.6	2.6	2.0	1.5	2.4	2.9	2.1	3.5
2001	2.2	1.7	2.5	2.0	1.6	2.4	3.1	2.3	3.7
2000	2.4	1.8	2.8	2.2	1.7	2.6	3.4	2.3	4.2
1999	2.4	1.8	2.8	2.2	1.7	2.6	3.3	2.4	4.0
Intracerebral and other intracranial hemorrhage (I61-I62)									
2005	9.1	10.3	8.2	8.5	9.6	7.7	13.2	15.5	11.3
2004	9.2	10.3	8.4	8.6	9.6	7.8	13.5	15.8	11.8
2003	9.3	10.4	8.4	8.6	9.6	7.8	14.1	16.7	12.2
2002	9.4	10.4	8.5	8.8	9.7	8.0	13.2	15.3	11.6
2001	9.4	10.5	8.6	8.8	9.8	8.0	13.7	15.8	12.1
2000	9.9	11.1	9.0	9.2	10.3	8.5	14.3	17.0	12.3
1999	9.3	10.3	8.5	8.7	9.6	8.0	13.3	15.5	11.7

Worktable 13. Age-adjusted death rates for 358 selected causes, by race and sex: United States, 1999-2005

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992) and year	All races 1/			White 2/			Black 2/		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
Cerebral infarction (I63)									
2005	2.4	2.5	2.4	2.4	2.4	2.3	3.3	3.6	3.1
2004	3.6	3.8	3.5	3.5	3.6	3.4	5.1	5.8	4.5
2003	4.1	4.2	3.9	4.0	4.1	3.9	5.4	5.8	5.0
2002	4.3	4.4	4.1	4.1	4.3	4.0	5.5	6.1	5.0
2001	4.5	4.6	4.3	4.4	4.5	4.2	5.8	6.8	5.2
2000	5.0	5.3	4.7	4.9	5.1	4.6	6.6	7.5	5.9
1999	5.4	5.7	5.2	5.3	5.5	5.1	7.1	8.3	6.3
Stroke, not specified as hemorrhage or infarction (I64)									
2005	24.1	23.7	23.8	23.3	22.7	23.2	33.5	36.1	31.2
2004	25.4	24.9	25.3	24.6	23.9	24.5	35.8	37.0	34.3
2003	28.0	27.7	27.6	27.1	26.6	26.9	38.7	40.6	36.8
2002	30.0	29.6	29.8	29.1	28.4	29.0	41.2	44.1	38.8
2001	31.4	31.4	30.9	30.5	30.2	30.2	42.7	46.4	39.8
2000	32.8	33.0	32.2	31.9	31.8	31.4	44.2	48.5	41.1
1999	33.6	34.0	32.8	32.7	32.8	32.0	44.6	48.4	41.8
Other cerebrovascular diseases and their sequelae (I67,I69)									
2005	8.9	8.9	8.9	8.7	8.5	8.6	12.4	13.1	11.8
2004	9.6	9.7	9.4	9.3	9.4	9.2	12.8	14.2	11.7
2003	10.1	10.2	9.9	9.9	9.9	9.7	13.3	14.2	12.5
2002	10.4	10.6	10.2	10.2	10.3	10.0	13.4	14.0	12.9
2001	10.4	10.7	10.1	10.2	10.5	9.8	13.5	14.1	13.0
2000	10.8	11.3	10.3	10.5	11.0	10.1	13.4	14.4	12.7
1999	10.9	11.4	10.5	10.7	11.1	10.3	13.6	15.0	12.5

Worktable 13. Age-adjusted death rates for 358 selected causes, by race and sex: United States, 1999-2005

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992) and year	All races 1/			White 2/			Black 2/		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
Diseases of arteries, arterioles and capillaries (I70-I78)									
2005	11.5	13.9	9.7	11.5	14.0	9.7	13.0	15.0	11.3
2004	11.9	14.3	10.1	11.9	14.4	10.0	13.2	15.1	11.7
2003	12.8	15.7	10.7	12.9	15.8	10.6	14.0	16.6	12.1
2002	13.4	16.4	11.2	13.5	16.6	11.2	14.4	17.0	12.6
2001	14.0	17.1	11.7	14.0	17.2	11.7	15.1	17.4	13.4
2000	14.7	18.4	12.0	14.7	18.6	11.9	15.7	18.4	13.9
1999	15.0	18.5	12.5	15.1	18.7	12.5	15.5	18.2	13.7
Atherosclerosis (I70)									
2005	3.8	3.9	3.7	3.9	4.0	3.7	3.4	3.4	3.3
2004	3.9	4.1	3.7	4.0	4.2	3.8	3.5	3.5	3.3
2003	4.4	4.5	4.3	4.5	4.5	4.4	4.2	4.7	3.8
2002	4.7	4.9	4.6	4.8	5.0	4.7	4.4	4.9	4.0
2001	5.0	5.1	4.8	5.1	5.2	5.0	4.3	4.6	4.1
2000	5.2	5.5	5.0	5.3	5.6	5.1	4.8	5.4	4.4
1999	5.5	5.7	5.3	5.6	5.8	5.4	5.0	5.8	4.5
Aortic aneurysm and dissection (I71)									
2005	4.6	6.5	3.1	4.6	6.6	3.1	4.6	6.0	3.6
2004	4.6	6.7	3.1	4.7	6.8	3.2	4.5	6.1	3.3
2003	5.1	7.5	3.3	5.1	7.7	3.3	4.6	6.1	3.5
2002	5.2	7.7	3.3	5.2	7.9	3.4	4.6	6.1	3.5
2001	5.4	8.0	3.6	5.5	8.2	3.6	5.0	6.4	3.9
2000	5.7	8.7	3.7	5.8	9.0	3.6	5.2	6.8	4.1
1999	5.8	8.8	3.7	5.9	9.1	3.7	4.9	6.3	3.9

Worktable 13. Age-adjusted death rates for 358 selected causes, by race and sex: United States, 1999-2005

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992) and year	All races 1/			White 2/			Black 2/		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
Other diseases of arteries, arterioles and capillaries (I72-I78)									
2005	3.2	3.5	2.9	3.1	3.4	2.8	5.0	5.6	4.5
2004	3.3	3.5	3.1	3.2	3.4	3.0	5.2	5.6	5.0
2003	3.3	3.7	3.1	3.2	3.6	3.0	5.2	5.7	4.9
2002	3.5	3.8	3.3	3.4	3.7	3.1	5.4	6.0	5.0
2001	3.6	3.9	3.3	3.4	3.8	3.2	5.9	6.4	5.5
2000	3.7	4.2	3.4	3.6	4.0	3.2	5.7	6.2	5.3
1999	3.7	4.0	3.4	3.5	3.9	3.3	5.6	6.0	5.3
Other disorders of circulatory system (I80-I99)									
2005	1.6	1.6	1.5	1.5	1.5	1.4	2.7	3.1	2.3
2004	1.5	1.6	1.5	1.5	1.5	1.4	2.7	2.9	2.6
2003	1.6	1.6	1.6	1.5	1.4	1.5	2.9	3.2	2.8
2002	1.6	1.6	1.6	1.5	1.5	1.6	2.9	3.2	2.6
2001	1.6	1.7	1.6	1.5	1.6	1.5	2.7	3.1	2.5
2000	1.7	1.7	1.6	1.6	1.6	1.5	2.9	3.0	2.8
1999	1.5	1.5	1.4	1.4	1.4	1.3	2.4	2.6	2.2
Phlebitis, thrombophlebitis, venous embolism and thrombosis (I80-I82)									
2005	1.1	1.1	1.1	1.0	1.0	1.1	2.0	2.2	1.9
2004	1.2	1.2	1.2	1.1	1.1	1.0	2.2	2.2	2.1
2003	1.2	1.1	1.3	1.1	1.0	1.2	2.3	2.4	2.3
2002	1.2	1.1	1.3	1.2	1.0	1.2	2.3	2.4	2.2
2001	1.2	1.2	1.3	1.2	1.1	1.2	2.2	2.3	2.1
2000	1.2	1.2	1.2	1.1	1.1	1.2	2.2	2.1	2.3
1999	1.0	0.9	1.0	0.9	0.9	0.9	1.7	1.7	1.6

Worktable 13. Age-adjusted death rates for 358 selected causes, by race and sex: United States, 1999-2005

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992) and year	All races 1/			White 2/			Black 2/		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
All other and unspecified disorders of circulatory system (I83-I99)									
2005	0.4	0.5	0.4	0.4	0.5	0.3	0.6	0.9	0.4
2004	0.4	0.5	0.3	0.4	0.5	0.3	0.5	0.7	0.4
2003	0.4	0.4	0.3	0.4	0.4	0.3	0.6	0.7	0.5
2002	0.4	0.4	0.3	0.4	0.4	0.3	0.6	0.8	0.4
2001	0.4	0.5	0.3	0.4	0.5	0.3	0.5	0.8	0.4
2000	0.4	0.5	0.3	0.4	0.5	0.3	0.6	0.9	0.5
1999	0.5	0.6	0.4	0.5	0.6	0.4	0.7	1.0	0.6
X. Diseases of the respiratory system (J00-J98)									
2005	78.5	94.3	68.2	80.7	95.7	70.9	67.7	91.3	53.7
2004	76.1	93.0	65.5	78.2	94.4	68.0	65.4	89.7	51.1
2003	80.5	98.0	69.5	82.6	99.2	72.2	68.5	95.1	53.1
2002	81.4	100.4	69.7	83.4	101.7	72.1	70.5	97.1	55.3
2001	81.6	101.2	69.8	83.3	102.3	71.9	71.8	99.5	56.3
2000	83.7	105.7	70.9	85.4	106.8	73.0	73.5	102.9	56.9
1999	84.3	107.7	70.6	85.9	108.7	72.5	75.5	106.9	57.8
Acute upper respiratory infections (J00-J06)									
2005	0.1	0.0	0.1	0.1	0.0	0.1	0.1	*	*
2004	0.0	0.0	0.0	0.0	0.0	0.0	*	*	*
2003	0.1	0.1	0.0	0.1	0.1	0.0	0.1	*	*
2002	0.0	0.0	0.1	0.0	0.0	0.1	*	*	*
2001	0.0	0.0	0.1	0.0	0.0	0.1	0.1	*	*
2000	0.0	0.0	0.1	0.1	0.1	0.1	*	*	*
1999	0.1	0.1	0.1	0.1	0.1	0.1	0.1	*	*

Worktable 13. Age-adjusted death rates for 358 selected causes, by race and sex: United States, 1999-2005

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992) and year	All races 1/			White 2/			Black 2/		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
Acute pharyngitis and tonsillitis (J02-J03)									
2005	0.0	*	*	*	*	*	*	*	*
2004	0.0	*	*	0.0	*	*	*	*	*
2003	0.0	*	*	*	*	*	*	*	*
2002	0.0	*	*	*	*	*	*	*	*
2001	0.0	*	*	0.0	*	*	*	*	*
2000	0.0	*	*	0.0	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*
Acute laryngitis and tracheitis (J04)									
2005	0.0	*	*	*	*	*	*	*	*
2004	0.0	*	*	0.0	*	*	*	*	*
2003	0.0	*	*	*	*	*	*	*	*
2002	0.0	*	*	*	*	*	*	*	*
2001	0.0	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*
1999	0.0	*	*	0.0	*	*	*	*	*
Other and unspecified acute upper respiratory infections (J00-J01,J05-J06)									
2005	0.0	0.0	0.0	0.0	0.0	0.0	*	*	*
2004	0.0	0.0	0.0	0.0	0.0	0.0	*	*	*
2003	0.0	0.1	0.0	0.0	0.0	0.0	*	*	*
2002	0.0	0.0	0.0	0.0	0.0	0.0	*	*	*
2001	0.0	0.0	0.0	0.0	0.0	0.0	*	*	*
2000	0.0	0.0	0.0	0.0	0.0	0.1	*	*	*
1999	0.0	0.1	0.0	0.0	0.0	0.0	0.1	*	*

Worktable 13. Age-adjusted death rates for 358 selected causes, by race and sex: United States, 1999-2005

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992) and year	All races 1/			White 2/			Black 2/		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
Other diseases of the respiratory system (J10-J98)									
2005	78.4	94.2	68.2	80.6	95.6	70.9	67.6	91.2	53.6
2004	76.1	92.9	65.5	78.1	94.4	67.9	65.3	89.6	51.1
2003	80.4	97.9	69.5	82.5	99.1	72.1	68.4	94.9	53.0
2002	81.3	100.4	69.7	83.3	101.7	72.1	70.5	97.0	55.3
2001	81.5	101.2	69.7	83.2	102.2	71.8	71.8	99.4	56.2
2000	83.7	105.7	70.8	85.3	106.7	72.9	73.5	102.8	56.8
1999	84.3	107.6	70.5	85.8	108.6	72.4	75.4	106.8	57.8
Influenza (J10-J11)									
2005	0.6	0.6	0.6	0.6	0.6	0.6	0.2	0.3	0.2
2004	0.4	0.4	0.3	0.4	0.4	0.4	0.2	0.2	0.1
2003	0.6	0.6	0.6	0.6	0.6	0.6	0.3	0.3	0.3
2002	0.2	0.3	0.2	0.2	0.3	0.2	0.1	*	*
2001	0.1	0.1	0.1	0.1	0.1	0.1	0.1	*	*
2000	0.6	0.7	0.6	0.6	0.7	0.6	0.3	0.4	0.3
1999	0.6	0.6	0.6	0.6	0.6	0.6	0.3	0.3	0.3
Pneumonia (J12-J18)									
2005	19.7	23.3	17.3	19.6	23.0	17.3	21.5	26.6	18.2
2004	19.4	23.3	16.9	19.2	22.9	16.8	22.1	28.9	18.0
2003	21.4	25.5	18.8	21.3	25.0	18.9	23.0	30.6	18.4
2002	22.4	26.7	19.6	22.3	26.4	19.7	23.9	30.7	19.9
2001	21.9	26.5	19.1	21.6	25.9	19.0	24.1	32.2	19.4
2000	23.1	28.2	20.1	22.9	27.7	20.0	25.3	33.5	20.5
1999	22.9	27.9	20.0	22.7	27.5	19.9	25.1	33.2	20.5

Worktable 13. Age-adjusted death rates for 358 selected causes, by race and sex: United States, 1999-2005

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992) and year	All races 1/			White 2/			Black 2/		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
Viral pneumonia, not elsewhere classified (J12)									
2005	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	*
2004	0.1	0.1	0.1	0.1	0.1	0.1	0.1	*	*
2003	0.1	0.1	0.1	0.1	0.1	0.1	0.1	*	*
2002	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	*
2001	0.1	0.1	0.1	0.1	0.1	0.1	0.1	*	*
2000	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	*
1999	0.1	0.1	0.1	0.1	0.1	0.1	0.2	0.2	0.1
Bacterial pneumonia (J13-J15)									
2005	1.6	2.0	1.3	1.5	2.0	1.3	1.8	2.2	1.5
2004	1.5	1.9	1.2	1.4	1.8	1.2	2.0	2.6	1.6
2003	1.6	2.1	1.3	1.6	2.0	1.3	1.9	2.5	1.5
2002	1.6	2.0	1.3	1.6	2.0	1.3	1.7	2.3	1.4
2001	1.6	2.0	1.2	1.5	2.0	1.3	1.9	2.8	1.3
2000	1.6	2.2	1.3	1.6	2.1	1.2	2.0	2.9	1.5
1999	1.4	1.8	1.2	1.4	1.7	1.2	1.7	2.3	1.3
Pneumonia due to other or unspecified organisms (J16,J18)									
2005	18.1	21.2	16.0	18.0	20.9	16.0	19.6	24.3	16.7
2004	17.9	21.4	15.6	17.7	21.0	15.6	20.1	26.2	16.4
2003	19.7	23.3	17.4	19.6	22.9	17.5	21.0	27.9	16.8
2002	20.7	24.6	18.2	20.6	24.3	18.3	22.1	28.3	18.5
2001	20.2	24.3	17.7	20.0	23.8	17.7	22.2	29.3	18.0
2000	21.3	25.9	18.7	21.1	25.4	18.6	23.2	30.5	18.9
1999	21.3	26.0	18.6	21.2	25.6	18.6	23.3	30.7	19.0

Worktable 13. Age-adjusted death rates for 358 selected causes, by race and sex: United States, 1999-2005

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992) and year	All races 1/			White 2/			Black 2/		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
Other acute lower respiratory infections (J20-J22)									
2005	0.1	0.1	0.1	0.1	0.1	0.1	0.1	*	0.1
2004	0.1	0.1	0.1	0.1	0.1	0.1	0.1	*	*
2003	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	*
2002	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1
2001	0.1	0.1	0.1	0.1	0.1	0.1	0.1	*	*
2000	0.1	0.1	0.1	0.1	0.1	0.2	0.1	0.2	0.1
1999	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2
Acute bronchitis and bronchiolitis (J20-J21)									
2005	0.1	0.1	0.1	0.1	0.1	0.1	0.1	*	*
2004	0.1	0.1	0.1	0.1	0.1	0.1	0.1	*	*
2003	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	*
2002	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	*
2001	0.1	0.1	0.1	0.1	0.1	0.1	0.1	*	*
2000	0.1	0.1	0.1	0.1	0.1	0.1	0.1	*	0.1
1999	0.1	0.1	0.1	0.1	0.1	0.1	0.2	0.2	0.2
Unspecified acute lower respiratory infection (J22)									
2005	0.0	0.0	0.0	0.0	0.0	0.0	*	*	*
2004	0.0	0.0	0.0	0.0	0.0	0.0	*	*	*
2003	0.0	0.0	0.0	0.0	0.0	0.0	*	*	*
2002	0.0	0.0	0.0	0.0	0.0	0.0	*	*	*
2001	0.0	0.0	0.0	0.0	0.0	0.0	*	*	*
2000	0.0	0.0	0.0	0.1	0.0	0.0	*	*	*
1999	0.1	0.0	0.1	0.1	0.0	0.1	*	*	*

Worktable 13. Age-adjusted death rates for 358 selected causes, by race and sex: United States, 1999-2005

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992) and year	All races 1/			White 2/			Black 2/		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
Other diseases of upper respiratory tract (J30-J39)									
2005	0.1	0.2	0.1	0.1	0.2	0.1	0.2	0.3	0.2
2004	0.1	0.2	0.1	0.1	0.1	0.1	0.2	0.3	0.2
2003	0.1	0.2	0.1	0.1	0.2	0.1	0.2	0.2	0.2
2002	0.1	0.1	0.1	0.1	0.1	0.1	0.2	0.2	0.1
2001	0.2	0.2	0.1	0.1	0.2	0.1	0.3	0.4	0.2
2000	0.2	0.2	0.1	0.2	0.2	0.1	0.2	0.3	0.2
1999	0.2	0.2	0.1	0.1	0.2	0.1	0.3	0.3	0.2
Chronic lower respiratory diseases (J40-J47)									
2005	43.2	51.2	38.1	45.4	52.8	40.7	30.6	44.1	22.8
2004	41.1	49.5	36.0	43.2	51.1	38.4	28.2	40.9	20.9
2003	43.3	52.3	37.8	45.4	53.8	40.3	30.1	44.4	22.0
2002	43.5	53.5	37.4	45.4	54.9	39.7	31.2	46.3	22.6
2001	43.7	54.0	37.6	45.6	55.5	39.8	30.9	46.3	22.4
2000	44.2	55.8	37.4	46.0	57.2	39.5	31.6	47.5	22.7
1999	45.4	58.7	37.7	47.1	60.0	39.7	33.5	51.5	23.4
Bronchitis, chronic and unspecified, emphysema and asthma (J40-J46)									
2005	42.9	51.0	37.7	45.1	52.6	40.3	30.5	44.0	22.7
2004	40.8	49.2	35.6	42.9	50.8	38.0	28.1	40.8	20.8
2003	43.0	52.1	37.4	45.1	53.5	39.9	30.0	44.3	21.8
2002	43.1	53.2	37.0	45.1	54.7	39.3	31.0	46.2	22.4
2001	43.3	53.8	37.2	45.3	55.2	39.4	30.8	46.1	22.3
2000	43.8	55.5	37.0	45.6	56.9	39.1	31.4	47.3	22.6
1999	45.1	58.4	37.3	46.8	59.7	39.3	33.3	51.3	23.2

Worktable 13. Age-adjusted death rates for 358 selected causes, by race and sex: United States, 1999-2005

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992) and year	All races 1/			White 2/			Black 2/		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
Bronchitis, not specified as acute or chronic (J40)									
2005	0.1	0.2	0.1	0.1	0.2	0.1	0.1	*	*
2004	0.1	0.2	0.1	0.1	0.2	0.1	0.1	*	*
2003	0.1	0.1	0.1	0.2	0.1	0.1	0.1	*	0.2
2002	0.2	0.2	0.2	0.2	0.2	0.2	0.1	*	0.1
2001	0.2	0.2	0.2	0.2	0.2	0.2	0.1	0.2	*
2000	0.2	0.2	0.2	0.2	0.3	0.2	0.2	0.2	0.2
1999	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.3	0.2
Chronic bronchitis (J41-J42)									
2005	0.1	0.1	0.1	0.1	0.1	0.1	0.1	*	*
2004	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	*
2003	0.1	0.2	0.1	0.1	0.2	0.1	0.1	*	0.1
2002	0.2	0.2	0.1	0.2	0.2	0.1	0.1	0.2	*
2001	0.1	0.2	0.1	0.2	0.2	0.1	0.1	*	*
2000	0.2	0.2	0.2	0.2	0.2	0.2	0.1	*	*
1999	0.2	0.2	0.2	0.2	0.2	0.2	0.1	*	*
Emphysema (J43)									
2005	4.6	5.8	3.8	4.9	6.0	4.1	3.1	4.9	1.9
2004	4.6	5.8	3.8	4.9	6.1	4.1	2.7	4.6	1.6
2003	5.1	6.3	4.3	5.4	6.6	4.6	3.2	5.3	2.0
2002	5.4	7.0	4.4	5.7	7.2	4.8	3.4	5.8	2.0
2001	5.8	7.4	4.7	6.1	7.8	5.1	3.3	5.3	2.2
2000	6.0	7.8	4.9	6.4	8.1	5.3	3.8	6.2	2.4
1999	6.5	8.6	5.2	6.9	8.9	5.6	4.0	6.7	2.3

Worktable 13. Age-adjusted death rates for 358 selected causes, by race and sex: United States, 1999-2005

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992) and year	All races 1/			White 2/			Black 2/		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
Other chronic obstructive pulmonary disease (J44)									
2005	36.7	43.9	32.2	38.9	45.5	34.6	24.2	36.1	17.4
2004	34.7	42.0	30.1	36.7	43.7	32.4	22.1	32.9	15.9
2003	36.2	44.2	31.3	38.2	45.8	33.6	23.3	35.9	16.2
2002	35.9	44.7	30.7	37.8	46.1	32.8	24.0	36.8	16.8
2001	35.7	44.8	30.3	37.5	46.2	32.4	23.6	37.3	16.1
2000	35.8	45.9	29.9	37.5	47.3	31.9	23.5	37.3	15.8
1999	36.5	47.9	29.8	38.1	49.3	31.6	25.1	40.6	16.5
Asthma (J45-J46)									
2005	1.3	1.0	1.5	1.0	0.7	1.3	3.0	2.7	3.3
2004	1.3	1.0	1.4	1.0	0.8	1.2	3.0	3.0	3.1
2003	1.4	1.1	1.6	1.1	0.9	1.3	3.2	2.9	3.3
2002	1.5	1.2	1.7	1.2	0.9	1.4	3.4	3.3	3.4
2001	1.5	1.2	1.7	1.2	0.9	1.5	3.6	3.2	3.8
2000	1.6	1.3	1.8	1.3	1.0	1.5	3.9	3.5	4.1
1999	1.7	1.3	2.0	1.4	1.0	1.7	3.9	3.5	4.1
Bronchiectasis (J47)									
2005	0.3	0.2	0.4	0.3	0.2	0.4	0.1	*	*
2004	0.3	0.2	0.4	0.3	0.2	0.4	0.1	*	0.1
2003	0.3	0.2	0.4	0.3	0.2	0.4	0.1	*	*
2002	0.3	0.2	0.4	0.3	0.2	0.4	0.1	*	*
2001	0.3	0.2	0.4	0.4	0.2	0.4	0.1	*	*
2000	0.3	0.2	0.4	0.4	0.3	0.4	0.1	*	*
1999	0.3	0.3	0.4	0.4	0.3	0.4	0.2	*	0.2

Worktable 13. Age-adjusted death rates for 358 selected causes, by race and sex: United States, 1999-2005

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992) and year	All races 1/			White 2/			Black 2/		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
Lung diseases due to external agents (J60-J70)									
2005	5.9	8.4	4.4	6.0	8.5	4.4	6.6	9.5	4.9
2004	6.0	8.6	4.3	6.0	8.7	4.4	6.3	9.3	4.6
2003	6.3	9.0	4.7	6.3	9.1	4.6	6.9	9.7	5.4
2002	6.5	9.4	4.8	6.5	9.5	4.8	6.8	9.9	5.2
2001	6.6	9.5	4.9	6.6	9.6	4.8	7.3	10.1	5.8
2000	6.5	9.7	4.7	6.5	9.8	4.7	7.0	10.2	5.2
1999	6.1	9.2	4.4	6.0	9.1	4.3	7.1	10.7	5.1
Pneumoconioses and chemical effects (J60-J66,J68)									
2005	0.3	0.8	0.0	0.4	0.9	0.0	0.2	0.5	*
2004	0.3	0.9	0.0	0.4	0.9	0.0	0.2	0.6	*
2003	0.4	1.0	0.0	0.4	1.0	0.0	0.2	0.5	*
2002	0.4	1.0	0.0	0.4	1.0	0.0	0.2	0.6	*
2001	0.4	1.0	0.0	0.4	1.1	0.0	0.2	0.7	*
2000	0.4	1.1	0.0	0.4	1.1	0.0	0.2	0.7	*
1999	0.4	1.0	0.0	0.4	1.1	0.0	0.3	0.8	*
Pneumonitis due to solids and liquids (J69)									
2005	5.6	7.6	4.4	5.6	7.6	4.4	6.4	9.0	4.9
2004	5.6	7.7	4.3	5.6	7.7	4.3	6.0	8.6	4.6
2003	5.9	8.0	4.6	5.8	8.0	4.6	6.7	9.1	5.3
2002	6.1	8.4	4.7	6.1	8.5	4.7	6.6	9.2	5.1
2001	6.1	8.5	4.8	6.1	8.5	4.8	7.0	9.4	5.7
2000	6.1	8.6	4.6	6.0	8.6	4.6	6.8	9.4	5.2
1999	5.6	8.1	4.3	5.6	8.0	4.3	6.8	9.9	5.1

Worktable 13. Age-adjusted death rates for 358 selected causes, by race and sex: United States, 1999-2005

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992) and year	All races 1/			White 2/			Black 2/		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
Pneumonitis due to food and vomit (J69.0)									
2005	5.6	7.6	4.4	5.6	7.6	4.4	6.4	9.0	4.9
2004	5.6	7.7	4.3	5.6	7.7	4.3	6.0	8.6	4.6
2003	5.9	8.0	4.6	5.8	8.0	4.6	6.7	9.1	5.3
2002	6.1	8.4	4.7	6.1	8.5	4.7	6.6	9.2	5.1
2001	6.1	8.5	4.8	6.1	8.5	4.8	7.0	9.4	5.7
2000	6.1	8.6	4.6	6.0	8.6	4.6	6.8	9.4	5.2
1999	5.6	8.1	4.3	5.6	8.0	4.3	6.8	9.9	5.1
Other pneumonitis due to solids and liquids (J69.1-J69.8)									
2005	*	*	*	*	*	*	*	*	*
2004	*	*	*	*	*	*	*	*	*
2003	*	*	*	*	*	*	*	*	*
2002	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*
Other lung diseases due to external agents (J67,J70)									
2005	0.0	0.0	0.0	0.0	0.0	0.0	*	*	*
2004	0.0	0.0	0.0	0.0	0.0	0.0	*	*	*
2003	0.0	0.0	0.0	0.0	0.0	0.0	*	*	*
2002	0.0	0.0	0.0	0.0	0.0	0.0	*	*	*
2001	0.0	0.0	0.0	0.0	0.1	0.0	*	*	*
2000	0.0	0.0	0.0	0.0	0.0	0.0	*	*	*
1999	0.0	0.0	0.0	0.0	0.0	0.0	*	*	*

Worktable 13. Age-adjusted death rates for 358 selected causes, by race and sex: United States, 1999-2005

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992) and year	All races 1/			White 2/			Black 2/		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
Suppurative and necrotic conditions of lower respiratory tract (J85-J86)									
2005	0.2	0.3	0.1	0.2	0.3	0.2	0.3	0.5	0.1
2004	0.2	0.3	0.2	0.2	0.3	0.2	0.3	0.4	0.2
2003	0.2	0.3	0.2	0.2	0.3	0.2	0.4	0.6	0.2
2002	0.2	0.4	0.2	0.2	0.3	0.1	0.4	0.7	0.3
2001	0.3	0.4	0.2	0.2	0.3	0.2	0.4	0.6	0.2
2000	0.2	0.4	0.2	0.2	0.4	0.2	0.4	0.7	0.3
1999	0.3	0.4	0.2	0.3	0.4	0.2	0.5	0.8	0.3
Pleural effusion and plaque (J90,J92)									
2005	0.3	0.4	0.3	0.3	0.4	0.3	0.3	0.4	0.2
2004	0.3	0.4	0.3	0.3	0.4	0.3	0.3	0.4	0.3
2003	0.3	0.4	0.3	0.3	0.4	0.3	0.3	0.4	0.3
2002	0.3	0.4	0.3	0.3	0.4	0.3	0.4	0.4	0.3
2001	0.3	0.4	0.3	0.3	0.4	0.3	0.5	0.5	0.4
2000	0.3	0.4	0.3	0.3	0.4	0.3	0.4	0.5	0.3
1999	0.3	0.4	0.3	0.3	0.4	0.3	0.3	0.4	0.3
All other diseases of respiratory system (J80-J84,J93-J98)									
2005	8.1	9.7	7.0	8.2	9.7	7.1	7.8	9.4	6.8
2004	8.4	10.1	7.2	8.5	10.3	7.3	7.6	9.2	6.6
2003	7.9	9.5	6.9	8.0	9.6	6.9	7.1	8.6	6.1
2002	7.9	9.4	6.9	8.0	9.5	6.9	7.3	8.5	6.6
2001	8.4	10.0	7.3	8.5	10.1	7.3	8.2	9.2	7.7
2000	8.4	10.2	7.2	8.5	10.3	7.3	8.1	9.6	7.2
1999	8.3	10.0	7.2	8.3	10.1	7.1	8.2	9.5	7.5

Worktable 13. Age-adjusted death rates for 358 selected causes, by race and sex: United States, 1999-2005

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992) and year	All races 1/			White 2/			Black 2/		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
XI. Diseases of the digestive system (K00-K92)									
2005	28.4	33.3	24.0	28.5	33.2	24.1	30.7	38.0	25.0
2004	28.8	33.4	24.5	28.8	33.2	24.5	31.6	39.6	25.5
2003	29.8	34.8	25.1	29.8	34.7	25.2	32.7	41.1	26.2
2002	29.9	35.0	25.3	30.0	34.9	25.4	32.4	40.1	26.5
2001	30.3	35.5	25.7	30.1	34.9	25.6	34.6	44.0	27.5
2000	30.4	35.8	25.5	30.0	35.3	25.3	35.3	43.8	28.8
1999	30.3	36.0	25.3	30.0	35.5	25.0	35.3	44.1	28.7
Diseases of oral cavity, salivary glands and jaws (K00-K14)									
2005	0.1	0.1	0.1	0.1	0.1	0.1	0.1	*	*
2004	0.1	0.1	0.1	0.1	0.1	0.1	0.1	*	*
2003	0.0	0.0	0.0	0.0	0.0	0.0	*	*	*
2002	0.0	0.1	0.0	0.0	0.1	0.0	*	*	*
2001	0.0	0.0	0.0	0.0	0.0	0.0	*	*	*
2000	0.0	0.0	0.0	0.0	0.0	0.0	*	*	*
1999	0.0	0.0	0.0	0.0	0.0	0.0	*	*	*
Diseases of other parts of digestive system (K20-K92)									
2005	28.4	33.2	23.9	28.4	33.1	24.0	30.6	37.9	25.0
2004	28.7	33.3	24.4	28.7	33.1	24.4	31.6	39.6	25.4
2003	29.8	34.8	25.1	29.7	34.6	25.2	32.6	41.1	26.2
2002	29.9	35.0	25.3	29.9	34.8	25.3	32.3	40.1	26.4
2001	30.2	35.4	25.6	30.1	34.9	25.6	34.5	44.0	27.4
2000	30.3	35.7	25.5	30.0	35.2	25.2	35.2	43.8	28.7
1999	30.3	35.9	25.3	30.0	35.4	25.0	35.3	44.0	28.6

Worktable 13. Age-adjusted death rates for 358 selected causes, by race and sex: United States, 1999-2005

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992) and year	All races 1/			White 2/			Black 2/		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
Diseases of esophagus (K20-K22)									
2005	0.6	0.8	0.5	0.6	0.8	0.5	0.5	0.6	0.5
2004	0.6	0.7	0.6	0.6	0.7	0.6	0.6	0.8	0.4
2003	0.7	0.8	0.6	0.7	0.9	0.6	0.6	0.7	0.5
2002	0.7	0.9	0.6	0.8	0.9	0.6	0.7	0.8	0.5
2001	0.7	0.9	0.7	0.8	0.9	0.7	0.6	0.9	0.5
2000	0.8	1.0	0.7	0.8	1.0	0.7	0.7	1.0	0.5
1999	0.8	1.0	0.7	0.8	1.0	0.7	0.7	1.0	0.5
Peptic ulcer (K25-K28)									
2005	1.1	1.4	0.9	1.1	1.3	0.9	1.3	1.8	0.9
2004	1.2	1.4	1.1	1.2	1.4	1.1	1.3	1.7	1.1
2003	1.3	1.6	1.1	1.3	1.5	1.2	1.4	2.1	1.0
2002	1.4	1.7	1.2	1.4	1.7	1.2	1.4	2.0	1.1
2001	1.6	1.8	1.4	1.6	1.8	1.4	1.6	2.3	1.1
2000	1.6	2.0	1.4	1.6	2.0	1.4	1.7	2.3	1.3
1999	1.7	2.0	1.4	1.7	2.0	1.4	1.9	2.4	1.4
Gastric ulcer (K25)									
2005	0.3	0.4	0.3	0.3	0.4	0.3	0.5	0.6	0.4
2004	0.4	0.4	0.3	0.4	0.4	0.4	0.5	0.6	0.4
2003	0.4	0.4	0.4	0.4	0.4	0.4	0.5	0.7	0.4
2002	0.4	0.5	0.4	0.4	0.4	0.4	0.5	0.6	0.4
2001	0.5	0.4	0.4	0.4	0.4	0.4	0.6	0.8	0.4
2000	0.5	0.5	0.4	0.5	0.5	0.4	0.7	1.0	0.4
1999	0.5	0.5	0.4	0.5	0.5	0.4	0.7	1.0	0.5

Worktable 13. Age-adjusted death rates for 358 selected causes, by race and sex: United States, 1999-2005

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992) and year	All races 1/			White 2/			Black 2/		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
Duodenal ulcer (K26)									
2005	0.4	0.5	0.3	0.4	0.5	0.3	0.4	0.7	0.2
2004	0.4	0.5	0.3	0.4	0.5	0.3	0.4	0.5	0.2
2003	0.4	0.6	0.4	0.5	0.6	0.3	0.5	0.7	0.3
2002	0.5	0.6	0.4	0.5	0.6	0.4	0.4	0.6	0.3
2001	0.5	0.7	0.4	0.6	0.7	0.4	0.4	0.6	0.3
2000	0.5	0.7	0.4	0.6	0.7	0.4	0.5	0.6	0.3
1999	0.5	0.7	0.4	0.5	0.7	0.4	0.5	0.6	0.4
Peptic ulcer, site unspecified (K27)									
2005	0.4	0.4	0.3	0.4	0.4	0.3	0.4	0.5	0.3
2004	0.4	0.5	0.4	0.4	0.4	0.4	0.4	0.5	0.4
2003	0.5	0.6	0.4	0.5	0.6	0.4	0.5	0.7	0.3
2002	0.5	0.6	0.4	0.5	0.6	0.4	0.5	0.7	0.3
2001	0.6	0.7	0.5	0.6	0.7	0.5	0.6	0.9	0.4
2000	0.6	0.8	0.5	0.6	0.8	0.5	0.6	0.7	0.5
1999	0.6	0.8	0.6	0.6	0.8	0.6	0.6	0.8	0.5
Gastrojejunal ulcer (K28)									
2005	0.0	*	*	0.0	*	*	*	*	*
2004	0.0	0.0	0.0	0.0	*	0.0	*	*	*
2003	0.0	0.0	0.0	0.0	0.0	0.0	*	*	*
2002	0.0	0.0	0.0	0.0	0.0	*	*	*	*
2001	0.0	*	0.0	0.0	*	0.0	*	*	*
2000	0.0	0.0	0.0	0.0	0.0	0.0	*	*	*
1999	0.0	*	*	0.0	*	*	*	*	*

Worktable 13. Age-adjusted death rates for 358 selected causes, by race and sex: United States, 1999-2005

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992) and year	All races 1/			White 2/			Black 2/		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
Gastritis and duodenitis (K29)									
2005	0.1	0.1	0.1	0.1	0.1	0.1	0.1	*	*
2004	0.1	0.2	0.1	0.1	0.1	0.1	0.2	0.2	0.1
2003	0.1	0.2	0.1	0.1	0.2	0.1	0.2	0.2	0.1
2002	0.1	0.1	0.1	0.1	0.2	0.1	0.1	*	0.1
2001	0.1	0.1	0.1	0.1	0.1	0.1	0.1	*	*
2000	0.1	0.2	0.1	0.1	0.2	0.1	0.2	*	0.1
1999	0.1	0.2	0.1	0.1	0.2	0.1	0.2	0.3	0.2
Dyspepsia and other diseases of stomach and duodenum (K30-K31)									
2005	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.3	0.2
2004	0.2	0.2	0.2	0.2	0.2	0.2	0.3	0.4	0.2
2003	0.2	0.2	0.2	0.2	0.2	0.2	0.3	0.2	0.3
2002	0.2	0.2	0.2	0.2	0.2	0.2	0.3	0.3	0.3
2001	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2
2000	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.3	0.2
1999	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2
Diseases of appendix (K35-K38)									
2005	0.1	0.2	0.1	0.1	0.2	0.1	0.2	0.2	0.1
2004	0.1	0.2	0.1	0.1	0.2	0.1	0.2	0.3	0.1
2003	0.1	0.2	0.1	0.1	0.2	0.1	0.2	0.3	0.2
2002	0.2	0.2	0.1	0.1	0.2	0.1	0.2	0.4	0.1
2001	0.1	0.2	0.1	0.1	0.2	0.1	0.2	0.4	0.2
2000	0.1	0.2	0.1	0.1	0.2	0.1	0.3	0.4	0.2
1999	0.1	0.2	0.1	0.1	0.2	0.1	0.2	0.4	0.1

Worktable 13. Age-adjusted death rates for 358 selected causes, by race and sex: United States, 1999-2005

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992) and year	All races 1/			White 2/			Black 2/		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
Hernia (K40-K46)									
2005	0.5	0.5	0.5	0.5	0.5	0.5	0.5	0.6	0.5
2004	0.5	0.5	0.5	0.5	0.5	0.5	0.5	0.6	0.5
2003	0.6	0.6	0.5	0.6	0.6	0.5	0.6	0.7	0.6
2002	0.5	0.5	0.6	0.6	0.5	0.6	0.5	0.6	0.5
2001	0.5	0.5	0.5	0.5	0.5	0.5	0.6	0.8	0.5
2000	0.5	0.5	0.5	0.6	0.6	0.6	0.6	0.6	0.6
1999	0.5	0.6	0.5	0.5	0.6	0.5	0.6	0.6	0.6
Crohn's disease and ulcerative colitis (K50-K51)									
2005	0.3	0.3	0.3	0.3	0.3	0.3	0.2	*	0.2
2004	0.3	0.3	0.3	0.3	0.3	0.3	0.2	0.3	0.1
2003	0.3	0.3	0.3	0.3	0.3	0.3	0.2	0.2	0.2
2002	0.3	0.3	0.3	0.3	0.3	0.3	0.2	0.2	0.1
2001	0.3	0.3	0.3	0.3	0.3	0.3	0.2	0.2	0.2
2000	0.3	0.3	0.3	0.3	0.3	0.3	0.2	*	0.2
1999	0.3	0.3	0.3	0.3	0.3	0.3	0.2	0.2	0.2
Vascular disorders and obstruction of intestine without hernia (K55-K56)									
2005	4.7	4.3	5.0	4.7	4.1	5.0	5.7	6.1	5.3
2004	4.8	4.3	5.1	4.8	4.2	5.1	6.0	6.6	5.6
2003	5.1	4.6	5.3	5.0	4.4	5.4	6.0	6.9	5.5
2002	5.0	4.5	5.3	5.0	4.4	5.3	6.0	6.3	5.7
2001	5.1	4.6	5.3	5.0	4.5	5.3	6.2	7.0	5.8
2000	5.2	4.7	5.4	5.1	4.5	5.4	6.6	7.3	6.1
1999	5.1	4.6	5.4	5.1	4.4	5.4	6.4	6.6	6.2

Worktable 13. Age-adjusted death rates for 358 selected causes, by race and sex: United States, 1999-2005

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992) and year	All races 1/			White 2/			Black 2/		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
Diverticular disease of intestine (K57)									
2005	1.0	0.8	1.1	1.0	0.8	1.2	0.9	0.8	1.0
2004	1.1	0.9	1.3	1.2	0.9	1.3	1.0	0.9	1.0
2003	1.2	1.0	1.3	1.2	1.0	1.3	1.0	0.8	1.1
2002	1.2	0.9	1.3	1.2	1.0	1.3	1.1	1.0	1.1
2001	1.2	1.0	1.3	1.2	1.0	1.4	1.1	0.8	1.2
2000	1.2	1.0	1.3	1.2	1.0	1.3	1.0	0.8	1.2
1999	1.3	1.0	1.4	1.3	1.1	1.4	1.1	0.8	1.2
Other diseases of intestines and peritoneum (K52,K58-K63)									
2005	1.8	1.7	1.9	1.8	1.7	1.9	2.1	2.4	1.8
2004	1.7	1.5	1.7	1.7	1.5	1.8	1.7	1.9	1.5
2003	1.7	1.6	1.7	1.7	1.6	1.7	1.9	2.0	1.8
2002	1.7	1.6	1.7	1.7	1.6	1.7	1.7	1.9	1.5
2001	1.7	1.6	1.7	1.7	1.5	1.8	1.9	2.1	1.8
2000	1.6	1.5	1.7	1.6	1.5	1.7	1.8	1.9	1.7
1999	1.6	1.6	1.7	1.6	1.5	1.7	1.7	2.0	1.5
Diseases of peritoneum (K65-K66)									
2005	0.6	0.6	0.6	0.6	0.6	0.6	0.8	0.8	0.9
2004	0.6	0.6	0.6	0.6	0.6	0.5	0.8	0.8	0.8
2003	0.6	0.7	0.6	0.6	0.6	0.6	0.9	0.9	1.0
2002	0.7	0.6	0.7	0.6	0.6	0.7	1.0	0.9	1.0
2001	0.7	0.7	0.6	0.6	0.7	0.6	1.1	1.2	1.0
2000	0.7	0.7	0.7	0.7	0.7	0.6	1.1	1.3	1.0
1999	0.7	0.8	0.6	0.7	0.7	0.6	1.1	1.2	1.0

Worktable 13. Age-adjusted death rates for 358 selected causes, by race and sex: United States, 1999-2005

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992) and year	All races 1/			White 2/			Black 2/		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
Diseases of liver (K70-K76)									
2005	12.0	16.2	8.2	12.2	16.4	8.3	11.4	16.1	7.8
2004	12.0	16.2	8.2	12.1	16.4	8.2	11.8	16.6	7.9
2003	12.4	16.8	8.4	12.5	17.0	8.5	12.2	17.6	8.1
2002	12.3	16.7	8.4	12.4	16.9	8.4	12.3	16.9	8.6
2001	12.3	16.7	8.3	12.3	16.7	8.3	13.0	18.9	8.5
2000	12.2	16.7	8.2	12.1	16.7	8.0	13.2	18.4	9.1
1999	12.3	17.0	8.1	12.1	16.9	7.9	13.9	19.6	9.5
Alcoholic liver disease (K70)									
2005	4.2	6.4	2.2	4.3	6.6	2.3	3.5	5.4	1.9
2004	4.2	6.3	2.2	4.3	6.5	2.2	3.5	5.6	1.9
2003	4.2	6.4	2.1	4.2	6.6	2.1	3.8	6.0	2.0
2002	4.2	6.3	2.2	4.2	6.5	2.2	3.8	5.8	2.3
2001	4.3	6.6	2.2	4.3	6.6	2.2	4.4	6.9	2.4
2000	4.3	6.8	2.2	4.3	6.9	2.0	4.6	7.0	2.8
1999	4.3	6.9	2.1	4.3	6.8	2.0	5.1	8.1	2.8
Chronic hepatitis, not elsewhere classified (K73)									
2005	0.0	0.0	0.0	0.0	0.0	0.0	*	*	*
2004	0.0	0.0	0.0	0.0	0.0	0.0	*	*	*
2003	0.0	0.1	0.0	0.0	0.1	0.0	*	*	*
2002	0.2	0.1	0.2	0.2	0.1	0.3	0.2	*	0.2
2001	0.2	0.1	0.2	0.2	0.1	0.2	0.2	*	0.2
2000	0.2	0.1	0.2	0.1	0.1	0.2	0.2	*	0.2
1999	0.2	0.1	0.2	0.1	0.1	0.2	0.2	*	0.3

Worktable 13. Age-adjusted death rates for 358 selected causes, by race and sex: United States, 1999-2005

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992) and year	All races 1/			White 2/			Black 2/		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
Fibrosis and cirrhosis of liver (K74)									
2005	4.7	6.0	3.6	4.8	6.1	3.7	4.2	6.0	2.8
2004	4.8	6.1	3.6	4.9	6.2	3.7	4.3	6.1	2.9
2003	5.1	6.5	3.8	5.3	6.6	4.0	4.5	6.3	3.1
2002	5.0	6.4	3.8	5.2	6.6	3.9	4.5	6.1	3.2
2001	5.1	6.5	3.8	5.2	6.5	3.9	4.7	6.8	3.0
2000	5.1	6.5	3.8	5.1	6.6	3.8	4.7	6.4	3.3
1999	5.0	6.6	3.7	5.1	6.6	3.8	4.8	6.7	3.2
Other diseases of liver (K71-K72,K75-K76)									
2005	3.0	3.8	2.4	2.9	3.7	2.2	3.7	4.6	3.0
2004	3.0	3.7	2.4	2.9	3.6	2.3	3.9	4.9	3.1
2003	3.0	3.8	2.4	3.0	3.7	2.3	3.9	5.1	3.0
2002	2.9	3.8	2.1	2.8	3.7	2.1	3.8	5.0	2.9
2001	2.7	3.5	2.1	2.7	3.4	2.0	3.8	5.0	2.9
2000	2.6	3.3	2.0	2.5	3.2	1.9	3.8	4.9	2.9
1999	2.7	3.4	2.1	2.5	3.3	1.9	3.9	4.7	3.2
Cholelithiasis and other disorders of gallbladder (K80-K82)									
2005	1.0	1.1	0.9	1.0	1.1	0.9	1.2	1.3	1.2
2004	1.0	1.2	0.9	1.0	1.2	0.9	1.2	1.2	1.2
2003	1.0	1.2	0.9	1.0	1.1	0.9	1.1	1.3	1.0
2002	1.0	1.2	0.9	1.0	1.2	0.9	1.1	1.1	1.1
2001	1.0	1.2	0.9	1.0	1.2	0.9	1.3	1.4	1.2
2000	1.0	1.2	0.9	1.0	1.2	0.9	1.2	1.1	1.2
1999	1.0	1.2	0.9	1.0	1.2	0.9	1.1	1.1	1.0

Worktable 13. Age-adjusted death rates for 358 selected causes, by race and sex: United States, 1999-2005

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992) and year	All races 1/			White 2/			Black 2/		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
Disorders of biliary tract and pancreas (K83-K86)									
2005	1.6	2.0	1.3	1.5	1.9	1.3	2.1	2.7	1.7
2004	1.6	2.0	1.3	1.6	1.9	1.3	2.4	3.0	1.9
2003	1.7	2.0	1.4	1.6	1.9	1.3	2.4	3.2	1.8
2002	1.7	2.1	1.3	1.6	2.0	1.3	2.2	3.1	1.7
2001	1.8	2.1	1.5	1.7	2.0	1.4	2.5	3.1	2.0
2000	1.7	2.1	1.4	1.6	1.9	1.3	2.5	3.3	1.9
1999	1.6	2.0	1.3	1.5	1.9	1.3	2.4	3.2	1.8
Pancreatitis (K85,K86.0-K86.1)									
2005	1.1	1.4	0.8	1.1	1.4	0.8	1.5	2.0	1.1
2004	1.2	1.4	0.9	1.1	1.4	0.8	1.8	2.5	1.4
2003	1.2	1.5	0.9	1.1	1.4	0.9	1.8	2.6	1.3
2002	1.2	1.5	0.9	1.2	1.5	0.9	1.6	2.3	1.1
2001	1.3	1.5	1.1	1.2	1.5	1.0	1.9	2.4	1.5
2000	1.2	1.5	1.0	1.2	1.4	0.9	1.9	2.6	1.3
1999	1.2	1.5	1.0	1.1	1.4	0.9	1.9	2.6	1.3
Other disorders of biliary tract and pancreas (K83,K86.2-K86.9)									
2005	0.5	0.6	0.4	0.5	0.5	0.4	0.6	0.7	0.5
2004	0.5	0.5	0.4	0.5	0.5	0.4	0.5	0.6	0.5
2003	0.5	0.6	0.4	0.5	0.6	0.4	0.6	0.6	0.6
2002	0.5	0.6	0.4	0.5	0.6	0.4	0.6	0.7	0.6
2001	0.5	0.5	0.4	0.4	0.5	0.4	0.6	0.7	0.5
2000	0.5	0.5	0.4	0.4	0.5	0.4	0.6	0.7	0.5
1999	0.4	0.5	0.4	0.4	0.5	0.4	0.5	0.6	0.5

Worktable 13. Age-adjusted death rates for 358 selected causes, by race and sex: United States, 1999-2005

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992) and year	All races 1/			White 2/			Black 2/		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
All other diseases of digestive system (K90,K92)									
2005	2.6	3.0	2.3	2.6	2.9	2.3	3.3	4.0	2.9
2004	2.6	3.1	2.3	2.6	3.0	2.3	3.4	4.1	2.8
2003	2.8	3.2	2.4	2.8	3.2	2.4	3.6	4.2	3.1
2002	2.9	3.3	2.5	2.8	3.2	2.5	3.6	4.4	3.0
2001	2.9	3.4	2.5	2.9	3.3	2.5	3.8	4.7	3.1
2000	2.9	3.4	2.6	2.9	3.3	2.5	4.0	4.8	3.4
1999	2.9	3.3	2.5	2.8	3.3	2.5	3.6	4.4	3.0
XII. Diseases of the skin and subcutaneous tissue (L00-L98)									
2005	1.4	1.3	1.4	1.2	1.2	1.2	3.3	3.0	3.5
2004	1.4	1.3	1.5	1.2	1.1	1.2	3.6	3.1	3.8
2003	1.4	1.3	1.5	1.2	1.1	1.3	3.6	3.5	3.7
2002	1.4	1.3	1.4	1.1	1.1	1.1	3.9	3.3	4.2
2001	1.3	1.2	1.4	1.1	1.0	1.2	3.6	3.4	3.7
2000	1.4	1.2	1.4	1.1	1.0	1.2	3.9	3.7	4.0
1999	1.3	1.3	1.4	1.1	1.0	1.1	4.2	4.0	4.4
Infections of skin and subcutaneous tissue (L00-L08)									
2005	0.5	0.6	0.5	0.5	0.6	0.5	0.7	0.7	0.7
2004	0.5	0.6	0.5	0.5	0.6	0.5	0.7	0.8	0.7
2003	0.5	0.6	0.5	0.5	0.5	0.5	0.7	0.8	0.6
2002	0.5	0.5	0.4	0.4	0.5	0.4	0.6	0.6	0.6
2001	0.4	0.4	0.4	0.4	0.4	0.4	0.6	0.5	0.6
2000	0.4	0.4	0.4	0.4	0.4	0.4	0.5	0.5	0.5
1999	0.4	0.4	0.3	0.4	0.4	0.3	0.5	0.6	0.5

Worktable 13. Age-adjusted death rates for 358 selected causes, by race and sex: United States, 1999-2005

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992) and year	All races 1/			White 2/			Black 2/		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
Other and unspecified diseases of skin and subcutaneous tissue (L10-L98)									
2005	0.8	0.7	0.9	0.6	0.6	0.7	2.6	2.3	2.8
2004	0.8	0.7	0.9	0.6	0.5	0.7	2.8	2.4	3.0
2003	0.8	0.7	1.0	0.7	0.6	0.7	3.0	2.8	3.1
2002	0.9	0.8	1.0	0.7	0.6	0.8	3.3	2.7	3.6
2001	0.9	0.7	1.0	0.7	0.6	0.8	3.1	2.9	3.1
2000	0.9	0.8	1.1	0.7	0.6	0.8	3.4	3.2	3.5
1999	1.0	0.9	1.1	0.8	0.7	0.8	3.8	3.4	3.9
XIII. Diseases of the musculoskeletal system and connective tissue (M00-M99)									
2005	4.7	3.5	5.6	4.6	3.4	5.4	6.0	4.4	7.3
2004	4.8	3.5	5.7	4.6	3.3	5.4	6.5	5.2	7.5
2003	4.9	3.6	5.8	4.7	3.4	5.6	6.5	5.4	7.4
2002	5.0	3.6	5.9	4.8	3.5	5.7	6.4	5.1	7.4
2001	5.0	3.6	5.9	4.8	3.5	5.7	6.4	5.1	7.5
2000	5.0	3.7	5.9	4.8	3.5	5.7	6.6	5.7	7.4
1999	4.9	3.5	5.9	4.7	3.4	5.6	6.3	5.0	7.4
Rheumatoid arthritis and related inflammatory polyarthropathies (M05-M08)									
2005	0.8	0.5	1.1	0.9	0.5	1.1	0.6	0.3	0.8
2004	0.9	0.5	1.1	0.9	0.5	1.2	0.7	0.4	0.9
2003	0.9	0.5	1.2	1.0	0.5	1.2	0.7	0.5	0.8
2002	1.0	0.6	1.2	1.0	0.6	1.3	0.7	0.4	0.8
2001	0.9	0.5	1.2	1.0	0.5	1.3	0.7	0.4	0.8
2000	1.0	0.6	1.2	1.0	0.6	1.3	0.6	0.4	0.8
1999	1.0	0.6	1.3	1.0	0.6	1.3	0.6	0.4	0.7

Worktable 13. Age-adjusted death rates for 358 selected causes, by race and sex: United States, 1999-2005

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992) and year	All races 1/			White 2/			Black 2/		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
Systemic lupus erythematosus (M32)									
2005	0.4	0.1	0.8	0.3	0.1	0.5	1.3	0.4	2.1
2004	0.4	0.2	0.7	0.3	0.1	0.5	1.4	0.5	2.1
2003	0.5	0.1	0.7	0.3	0.1	0.5	1.2	0.3	2.0
2002	0.4	0.1	0.7	0.3	0.1	0.5	1.4	0.2	2.3
2001	0.5	0.2	0.8	0.3	0.1	0.6	1.3	0.3	2.1
2000	0.5	0.2	0.8	0.3	0.1	0.6	1.3	0.4	2.2
1999	0.5	0.2	0.8	0.4	0.1	0.6	1.3	0.3	2.2
Other arthropathies and related disorders (M00,M02,M10-M31,M33-M35)									
2005	1.6	1.1	1.9	1.5	1.1	1.8	2.1	1.4	2.5
2004	1.6	1.1	1.9	1.5	1.1	1.8	2.1	1.5	2.6
2003	1.6	1.2	1.9	1.5	1.1	1.8	2.3	1.9	2.7
2002	1.7	1.2	2.0	1.6	1.1	1.9	2.2	1.7	2.6
2001	1.7	1.3	2.0	1.6	1.2	1.9	2.3	1.8	2.6
2000	1.7	1.3	2.0	1.7	1.2	2.0	2.4	2.1	2.6
1999	1.7	1.3	2.0	1.6	1.2	1.9	2.4	1.9	2.9
Dorsopathies (M40-M54)									
2005	0.3	0.4	0.3	0.3	0.4	0.4	0.4	0.5	0.3
2004	0.4	0.3	0.4	0.4	0.3	0.4	0.3	0.5	0.2
2003	0.3	0.4	0.3	0.3	0.3	0.3	0.4	0.5	0.3
2002	0.3	0.3	0.3	0.3	0.3	0.3	0.3	0.5	0.2
2001	0.3	0.3	0.3	0.3	0.3	0.3	0.4	0.4	0.3
2000	0.3	0.3	0.3	0.3	0.3	0.3	0.3	0.4	0.2
1999	0.3	0.3	0.2	0.3	0.3	0.3	0.3	0.4	0.2

Worktable 13. Age-adjusted death rates for 358 selected causes, by race and sex: United States, 1999-2005

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992) and year	All races 1/			White 2/			Black 2/		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
Soft tissue disorders (M60-M79)									
2005	0.4	0.5	0.3	0.4	0.5	0.3	0.6	0.8	0.5
2004	0.4	0.5	0.4	0.4	0.5	0.3	0.8	1.0	0.6
2003	0.4	0.5	0.4	0.4	0.5	0.3	0.7	0.8	0.7
2002	0.4	0.5	0.3	0.4	0.5	0.3	0.7	0.9	0.5
2001	0.4	0.5	0.3	0.3	0.5	0.3	0.7	0.9	0.6
2000	0.4	0.5	0.3	0.3	0.4	0.2	0.8	0.9	0.7
1999	0.4	0.5	0.3	0.3	0.4	0.2	0.7	1.0	0.5
Osteopathies, chondropathies and other disorders of musculoskeletal system and connective tissue (M80-M99)									
2005	1.0	0.8	1.2	1.1	0.8	1.2	1.1	1.0	1.1
2004	1.1	0.8	1.2	1.1	0.8	1.3	1.1	1.3	1.0
2003	1.1	0.8	1.3	1.2	0.8	1.3	1.1	1.5	0.9
2002	1.2	0.8	1.3	1.2	0.8	1.3	1.1	1.3	1.0
2001	1.1	0.8	1.3	1.2	0.8	1.3	1.1	1.2	1.0
2000	1.1	0.8	1.2	1.1	0.8	1.3	1.1	1.4	0.9
1999	1.1	0.7	1.2	1.1	0.7	1.3	0.9	1.0	0.9
Osteoporosis (M80-M81)									
2005	0.5	0.2	0.7	0.6	0.2	0.8	0.1	*	0.1
2004	0.6	0.2	0.7	0.6	0.2	0.8	0.1	*	0.1
2003	0.6	0.2	0.8	0.6	0.2	0.9	*	*	*
2002	0.6	0.2	0.8	0.7	0.2	0.9	0.1	*	0.2
2001	0.6	0.2	0.8	0.6	0.2	0.8	0.1	*	0.1
2000	0.6	0.2	0.8	0.7	0.2	0.9	0.1	*	*
1999	0.6	0.2	0.8	0.7	0.2	0.9	0.1	*	*

Worktable 13. Age-adjusted death rates for 358 selected causes, by race and sex: United States, 1999-2005

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992) and year	All races 1/			White 2/			Black 2/		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
Other osteopathies, chondropathies and disorders of musculoskeletal system and connective tissue (M83-M99)									
2005	0.5	0.6	0.4	0.5	0.6	0.4	1.0	1.0	0.9
2004	0.5	0.6	0.5	0.5	0.6	0.5	1.0	1.2	0.9
2003	0.6	0.6	0.5	0.5	0.6	0.5	1.1	1.4	0.8
2002	0.5	0.6	0.4	0.5	0.6	0.4	1.0	1.3	0.8
2001	0.5	0.6	0.5	0.5	0.6	0.4	1.0	1.2	0.9
2000	0.5	0.6	0.4	0.4	0.5	0.4	1.0	1.3	0.8
1999	0.4	0.5	0.4	0.4	0.5	0.4	0.8	0.9	0.7
XIV. Diseases of the genitourinary system (N00-N98)									
2005	20.4	23.3	18.6	18.8	21.8	17.0	37.9	42.4	35.0
2004	20.1	23.0	18.3	18.5	21.4	16.7	37.9	43.0	34.8
2003	20.4	23.4	18.5	18.8	21.8	16.9	38.8	44.4	35.2
2002	20.2	23.2	18.3	18.6	21.6	16.7	38.2	44.4	34.6
2001	20.2	23.2	18.5	18.5	21.6	16.7	38.6	43.1	36.0
2000	19.8	23.0	18.0	18.1	21.4	16.3	37.7	42.6	34.8
1999	19.5	22.7	17.7	17.8	20.9	16.1	37.9	44.6	34.2
Diseases of urinary system (N00-N39)									
2005	20.0	22.7	18.3	18.5	21.2	16.7	37.4	41.7	34.6
2004	19.8	22.4	18.1	18.2	20.8	16.5	37.4	42.1	34.4
2003	20.1	22.9	18.3	18.4	21.3	16.7	38.4	43.7	34.9
2002	19.9	22.6	18.1	18.3	21.0	16.6	37.7	43.5	34.2
2001	19.9	22.6	18.2	18.2	21.0	16.5	38.1	42.4	35.6
2000	19.5	22.4	17.7	17.8	20.8	16.1	37.2	41.8	34.4
1999	19.2	22.0	17.5	17.5	20.2	15.9	37.3	43.7	33.7

Worktable 13. Age-adjusted death rates for 358 selected causes, by race and sex: United States, 1999-2005

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992) and year	All races 1/			White 2/			Black 2/		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
Glomerular and renal tubulo-interstitial diseases (N00-N15)									
2005	0.7	0.8	0.7	0.7	0.8	0.6	1.1	1.3	1.0
2004	0.6	0.7	0.6	0.6	0.7	0.6	0.9	1.0	0.8
2003	0.7	0.7	0.6	0.6	0.7	0.6	0.9	1.1	0.8
2002	0.7	0.8	0.6	0.6	0.7	0.6	0.8	1.0	0.7
2001	0.7	0.8	0.6	0.6	0.7	0.6	0.8	1.0	0.7
2000	0.7	0.8	0.7	0.6	0.7	0.6	1.1	1.2	1.0
1999	0.7	0.9	0.7	0.7	0.9	0.7	1.1	1.5	0.9
Glomerular diseases (N00-N07)									
2005	0.3	0.4	0.2	0.3	0.4	0.2	0.6	0.8	0.5
2004	0.2	0.3	0.2	0.2	0.2	0.2	0.4	0.4	0.4
2003	0.2	0.3	0.2	0.2	0.2	0.2	0.3	0.4	0.2
2002	0.2	0.3	0.2	0.2	0.3	0.2	0.4	0.4	0.4
2001	0.2	0.3	0.2	0.2	0.3	0.2	0.3	0.4	0.3
2000	0.2	0.3	0.2	0.2	0.3	0.2	0.4	0.5	0.4
1999	0.3	0.3	0.2	0.2	0.3	0.2	0.5	0.6	0.4
Acute and rapidly progressive nephritic syndrome (N00-N01)									
2005	*	*	*	*	*	*	*	*	*
2004	0.0	*	*	0.0	*	*	*	*	*
2003	*	*	*	*	*	*	*	*	*
2002	0.0	*	*	0.0	*	*	*	*	*
2001	0.0	*	*	0.0	*	*	*	*	*
2000	0.0	*	*	0.0	*	*	*	*	*
1999	0.0	*	*	0.0	*	*	*	*	*

Worktable 13. Age-adjusted death rates for 358 selected causes, by race and sex: United States, 1999-2005

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992) and year	All races 1/			White 2/			Black 2/		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
Nephrotic syndrome (N04)									
2005	0.0	0.0	0.0	0.0	0.0	0.0	*	*	*
2004	0.0	0.0	0.0	0.0	0.0	0.0	*	*	*
2003	0.0	0.0	0.0	0.0	0.0	0.0	*	*	*
2002	0.0	0.0	0.0	0.0	0.0	0.0	0.1	*	*
2001	0.0	0.0	0.0	0.0	0.0	0.0	*	*	*
2000	0.0	0.1	0.0	0.0	0.1	0.0	0.1	*	*
1999	0.1	0.1	0.0	0.0	0.0	0.0	0.1	*	*
Other and unspecified glomerular diseases (N02-N03,N05-N07)									
2005	0.3	0.3	0.2	0.2	0.3	0.2	0.6	0.7	0.5
2004	0.2	0.2	0.2	0.2	0.2	0.1	0.3	0.3	0.3
2003	0.2	0.2	0.1	0.1	0.2	0.1	0.3	0.3	0.2
2002	0.2	0.2	0.1	0.2	0.2	0.1	0.3	0.3	0.2
2001	0.2	0.2	0.1	0.1	0.2	0.1	0.2	0.3	0.2
2000	0.2	0.2	0.2	0.2	0.2	0.1	0.3	0.4	0.3
1999	0.2	0.2	0.1	0.2	0.2	0.1	0.4	0.4	0.3
Renal tubulo-interstitial diseases (N10-N15)									
2005	0.4	0.4	0.4	0.4	0.4	0.4	0.5	0.6	0.5
2004	0.4	0.5	0.4	0.4	0.4	0.4	0.5	0.6	0.4
2003	0.4	0.5	0.4	0.4	0.5	0.4	0.6	0.6	0.6
2002	0.4	0.5	0.4	0.4	0.5	0.4	0.4	0.5	0.4
2001	0.4	0.5	0.4	0.4	0.4	0.4	0.5	0.6	0.5
2000	0.5	0.5	0.5	0.4	0.4	0.4	0.6	0.7	0.6
1999	0.5	0.6	0.5	0.5	0.6	0.5	0.6	0.9	0.5

Worktable 13. Age-adjusted death rates for 358 selected causes, by race and sex: United States, 1999-2005

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992) and year	All races 1/			White 2/			Black 2/		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
Renal failure (N17-N19)									
2005	14.0	17.1	12.0	12.6	15.8	10.5	29.0	33.3	26.3
2004	13.9	17.0	12.0	12.5	15.6	10.6	29.1	33.8	26.3
2003	14.2	17.5	12.1	12.7	16.1	10.7	30.0	35.6	26.5
2002	13.9	17.3	11.9	12.5	15.8	10.5	29.3	34.8	25.9
2001	13.7	17.1	11.8	12.3	15.8	10.2	29.2	33.4	26.7
2000	13.2	16.6	11.3	11.8	15.2	9.8	28.2	32.6	25.5
1999	12.8	16.1	10.8	11.3	14.6	9.4	27.9	33.0	24.9
Urolithiasis (N20-N23)									
2005	0.1	0.1	0.1	0.1	0.1	0.1	0.1	*	*
2004	0.1	0.1	0.1	0.1	0.1	0.1	0.1	*	*
2003	0.1	0.1	0.1	0.1	0.1	0.1	0.1	*	*
2002	0.1	0.1	0.1	0.1	0.1	0.1	0.1	*	*
2001	0.1	0.1	0.1	0.1	0.1	0.1	0.1	*	*
2000	0.1	0.1	0.1	0.1	0.1	0.1	*	*	*
1999	0.1	0.1	0.1	0.1	0.1	0.1	*	*	*
Other and unspecified disorders of kidney (N25-N27)									
2005	0.0	0.0	0.0	0.0	0.0	0.0	*	*	*
2004	0.0	0.0	0.0	0.0	0.0	0.0	*	*	*
2003	0.0	0.0	0.0	0.0	0.0	0.0	*	*	*
2002	0.0	0.0	0.0	0.0	0.0	0.0	*	*	*
2001	0.0	*	0.0	0.0	*	0.0	*	*	*
2000	0.0	0.0	0.0	0.0	0.0	0.0	*	*	*
1999	0.0	0.0	0.0	0.0	0.0	0.0	0.1	*	*

Worktable 13. Age-adjusted death rates for 358 selected causes, by race and sex: United States, 1999-2005

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992) and year	All races 1/			White 2/			Black 2/		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
Urinary tract infection, site not specified (N39.0)									
2005	4.8	4.1	5.2	4.7	3.9	5.2	6.5	6.0	6.6
2004	4.6	4.0	5.0	4.5	3.8	4.9	6.5	6.4	6.5
2003	4.7	3.9	5.1	4.6	3.9	4.9	6.7	6.1	7.0
2002	4.7	4.0	5.2	4.6	3.8	5.0	6.8	6.7	6.9
2001	5.0	4.1	5.4	4.8	4.0	5.2	7.3	7.0	7.4
2000	5.0	4.4	5.3	4.9	4.2	5.2	7.2	7.0	7.1
1999	5.2	4.4	5.5	5.0	4.2	5.4	7.5	8.0	7.2
Other diseases of urinary system (N28-N36,N39.1-N39.9)									
2005	0.4	0.5	0.3	0.4	0.5	0.3	0.6	0.8	0.5
2004	0.4	0.5	0.3	0.4	0.5	0.3	0.7	0.8	0.7
2003	0.4	0.5	0.3	0.4	0.4	0.3	0.6	0.8	0.5
2002	0.4	0.5	0.3	0.4	0.4	0.3	0.7	0.8	0.6
2001	0.4	0.4	0.3	0.4	0.4	0.3	0.7	0.8	0.6
2000	0.4	0.5	0.3	0.4	0.5	0.3	0.7	0.9	0.6
1999	0.4	0.5	0.3	0.3	0.5	0.3	0.7	0.9	0.5
Diseases of male genital organs (N40-N50)									
2005	0.2	0.6	...	0.2	0.6	...	0.2	0.7	...
2004	0.2	0.6	...	0.2	0.6	...	0.3	0.8	...
2003	0.2	0.6	...	0.2	0.6	...	0.2	0.6	...
2002	0.2	0.6	...	0.2	0.5	...	0.3	0.8	...
2001	0.2	0.6	...	0.2	0.6	...	0.3	0.8	...
2000	0.2	0.6	...	0.2	0.6	...	0.3	0.8	...
1999	0.2	0.6	...	0.2	0.6	...	0.3	0.9	...

Worktable 13. Age-adjusted death rates for 358 selected causes, by race and sex: United States, 1999-2005

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992) and year	All races 1/			White 2/			Black 2/		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
Hyperplasia of prostate (N40)									
2005	0.2	0.5	...	0.2	0.5	...	0.1	0.4	...
2004	0.2	0.5	...	0.2	0.5	...	0.1	0.4	...
2003	0.2	0.5	...	0.2	0.5	...	0.1	0.4	...
2002	0.1	0.4	...	0.1	0.4	...	0.2	0.5	...
2001	0.1	0.5	...	0.2	0.5	...	0.1	0.4	...
2000	0.2	0.5	...	0.2	0.5	...	0.2	0.5	...
1999	0.2	0.5	...	0.2	0.5	...	0.2	0.5	...
Other diseases of male genital organs (N41-N50)									
2005	0.1	0.2	...	0.1	0.2	...	0.1	0.3	...
2004	0.1	0.2	...	0.1	0.1	...	0.1	0.4	...
2003	0.0	0.1	...	0.0	0.1	...	0.1	0.2	...
2002	0.1	0.1	...	0.0	0.1	...	0.1	0.3	...
2001	0.1	0.1	...	0.0	0.1	...	0.1	0.3	...
2000	0.0	0.1	...	0.0	0.1	...	0.1	0.3	...
1999	0.1	0.2	...	0.1	0.1	...	0.1	0.4	...
Disorders of breast (N60-N64)									
2005	0.0	*	0.0	0.0	*	0.0	*	*	*
2004	0.0	*	0.0	0.0	*	0.0	*	*	*
2003	0.0	*	0.0	0.0	*	0.0	*	*	*
2002	0.0	*	0.0	0.0	*	0.0	*	*	*
2001	0.0	*	0.0	0.0	*	0.0	*	*	*
2000	0.0	*	0.0	0.0	*	0.0	*	*	*
1999	0.0	*	0.0	0.0	*	0.0	*	*	*

Worktable 13. Age-adjusted death rates for 358 selected causes, by race and sex: United States, 1999-2005

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992) and year	All races 1/			White 2/			Black 2/		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
Disorders of female genital organs (N70-N98)									
2005	0.1	...	0.2	0.1	...	0.2	0.2	...	0.3
2004	0.1	...	0.2	0.1	...	0.2	0.2	...	0.3
2003	0.1	...	0.2	0.1	...	0.2	0.2	...	0.3
2002	0.1	...	0.2	0.1	...	0.2	0.2	...	0.3
2001	0.1	...	0.2	0.1	...	0.2	0.2	...	0.4
2000	0.1	...	0.2	0.1	...	0.2	0.2	...	0.3
1999	0.1	...	0.2	0.1	...	0.2	0.3	...	0.4
Inflammatory diseases of female pelvic organs (N70-N76)									
2005	0.0	...	0.1	0.0	...	0.1	*	...	*
2004	0.0	...	0.1	0.0	...	0.0	0.1	...	0.1
2003	0.0	...	0.1	0.0	...	0.1	0.1	...	0.1
2002	0.0	...	0.1	0.0	...	0.0	0.1	...	0.1
2001	0.0	...	0.1	0.0	...	0.1	0.1	...	0.1
2000	0.0	...	0.1	0.0	...	0.1	0.1	...	0.1
1999	0.0	...	0.1	0.0	...	0.0	0.1	...	0.2
Noninflammatory disorders of female genital tract (N80-N98)									
2005	0.1	...	0.1	0.1	...	0.1	0.1	...	0.2
2004	0.1	...	0.1	0.1	...	0.1	0.1	...	0.2
2003	0.1	...	0.1	0.1	...	0.1	0.1	...	0.2
2002	0.1	...	0.1	0.0	...	0.1	0.1	...	0.2
2001	0.1	...	0.1	0.1	...	0.1	0.1	...	0.2
2000	0.1	...	0.1	0.1	...	0.1	0.1	...	0.2
1999	0.1	...	0.1	0.1	...	0.1	0.1	...	0.3

Worktable 13. Age-adjusted death rates for 358 selected causes, by race and sex: United States, 1999-2005

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992) and year	All races 1/			White 2/			Black 2/		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
XV. Pregnancy, childbirth and the puerperium (O00-O99)									
2005	0.3	...	0.5	0.2	...	0.4	0.7	...	1.3
2004	0.2	...	0.5	0.2	...	0.4	0.6	...	1.2
2003	0.2	...	0.4	0.1	...	0.3	0.5	...	0.9
2002	0.1	...	0.3	0.1	...	0.2	0.4	...	0.8
2001	0.1	...	0.3	0.1	...	0.2	0.4	...	0.8
2000	0.1	...	0.3	0.1	...	0.2	0.4	...	0.7
1999	0.2	...	0.3	0.1	...	0.2	0.4	...	0.8
Pregnancy with abortive outcome (O00-O07)									
2005	0.0	...	0.0	*	...	*	*	...	*
2004	0.0	...	0.0	*	...	*	*	...	*
2003	0.0	...	0.0	*	...	*	*	...	*
2002	0.0	...	0.0	*	...	*	*	...	*
2001	0.0	...	0.0	0.0	...	0.0	*	...	*
2000	0.0	...	0.0	*	...	*	*	...	*
1999	0.0	...	0.0	*	...	*	*	...	*
Ectopic pregnancy (O00)									
2005	*	...	*	*	...	*	*	...	*
2004	*	...	*	*	...	*	*	...	*
2003	0.0	...	0.0	*	...	*	*	...	*
2002	*	...	*	*	...	*	*	...	*
2001	0.0	...	0.0	*	...	*	*	...	*
2000	0.0	...	0.0	*	...	*	*	...	*
1999	*	...	*	*	...	*	*	...	*

Worktable 13. Age-adjusted death rates for 358 selected causes, by race and sex: United States, 1999-2005

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992) and year	All races 1/			White 2/			Black 2/		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
Spontaneous abortion (O03)									
2005	*	...	*	*	...	*	*	...	*
2004	*	...	*	*	...	*	*	...	*
2003	*	...	*	*	...	*	*	...	*
2002	*	...	*	*	...	*	*	...	*
2001	*	...	*	*	...	*	*	...	*
2000	*	...	*	*	...	*	*	...	*
1999	*	...	*	*	...	*	*	...	*
Medical abortion (O04)									
2005	*	...	*	*	...	*	*	...	*
2004	*	...	*	*	...	*	*	...	*
2003	*	...	*	*	...	*	*	...	*
2002	*	...	*	*	...	*	*	...	*
2001	*	...	*	*	...	*	*	...	*
2000	*	...	*	*	...	*	*	...	*
1999	*	...	*	*	...	*	*	...	*
Other abortion (O05)									
2005	*	...	*	*	...	*	*	...	*
2004	*	...	*	*	...	*	*	...	*
2003	*	...	*	*	...	*	*	...	*
2002	*	...	*	*	...	*	*	...	*
2001	*	...	*	*	...	*	*	...	*
2000	*	...	*	*	...	*	*	...	*
1999	*	...	*	*	...	*	*	...	*

Worktable 13. Age-adjusted death rates for 358 selected causes, by race and sex: United States, 1999-2005

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992) and year	All races 1/			White 2/			Black 2/		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
Other and unspecified pregnancy with abortive outcome (O01-O02,O06-O07)									
2005	*	...	*	*	...	*	*	...	*
2004	*	...	*	*	...	*	*	...	*
2003	*	...	*	*	...	*	*	...	*
2002	*	...	*	*	...	*	*	...	*
2001	*	...	*	*	...	*	*	...	*
2000	*	...	*	*	...	*	*	...	*
1999	*	...	*	*	...	*	*	...	*
Other direct obstetric deaths (O10-O92)									
2005	0.1	...	0.3	0.1	...	0.2	0.4	...	0.7
2004	0.1	...	0.2	0.1	...	0.2	0.3	...	0.7
2003	0.1	...	0.2	0.1	...	0.2	0.3	...	0.7
2002	0.1	...	0.2	0.1	...	0.1	0.3	...	0.6
2001	0.1	...	0.2	0.1	...	0.2	0.3	...	0.6
2000	0.1	...	0.2	0.1	...	0.2	0.2	...	0.5
1999	0.1	...	0.2	0.1	...	0.1	0.3	...	0.6
Eclampsia and pre-eclampsia (O11,O13-O16)									
2005	0.0	...	0.0	0.0	...	0.0	0.0	...	0.1
2004	0.0	...	0.0	0.0	...	0.0	0.1	...	0.1
2003	0.0	...	0.0	0.0	...	0.0	0.1	...	0.1
2002	0.0	...	0.0	0.0	...	0.0	0.1	...	0.1
2001	0.0	...	0.1	0.0	...	0.0	0.1	...	0.1
2000	0.0	...	0.1	0.0	...	0.1	0.1	...	0.1
1999	0.0	...	0.1	0.0	...	0.0	0.1	...	0.2

Worktable 13. Age-adjusted death rates for 358 selected causes, by race and sex: United States, 1999-2005

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992) and year	All races 1/			White 2/			Black 2/		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
Hemorrhage of pregnancy and childbirth and placenta previa (O20,O44-O46,O67,O72)									
2005	0.0	...	0.0	0.0	...	0.0	*	...	*
2004	0.0	...	0.0	0.0	...	0.0	*	...	*
2003	0.0	...	0.0	0.0	...	0.0	*	...	*
2002	0.0	...	0.0	*	...	*	*	...	*
2001	0.0	...	0.0	0.0	...	0.0	*	...	*
2000	0.0	...	0.0	0.0	...	0.0	*	...	*
1999	0.0	...	0.0	0.0	...	0.0	*	...	*
Complications predominately related to the puerperium (O85-O92)									
2005	0.0	...	0.1	0.0	...	0.0	0.1	...	0.2
2004	0.0	...	0.1	0.0	...	0.1	0.1	...	0.2
2003	0.0	...	0.1	0.0	...	0.0	0.1	...	0.2
2002	0.0	...	0.1	0.0	...	0.0	0.1	...	0.2
2001	0.0	...	0.1	0.0	...	0.1	0.1	...	0.2
2000	0.0	...	0.1	0.0	...	0.0	0.1	...	0.1
1999	0.0	...	0.1	0.0	...	0.0	0.1	...	0.2
Obstetric embolism (O88)									
2005	0.0	...	0.0	0.0	...	0.0	*	...	*
2004	0.0	...	0.0	0.0	...	0.0	*	...	*
2003	0.0	...	0.0	0.0	...	0.0	*	...	*
2002	0.0	...	0.0	0.0	...	0.0	*	...	*
2001	0.0	...	0.0	0.0	...	0.0	0.1	...	0.1
2000	0.0	...	0.0	0.0	...	0.0	*	...	*
1999	0.0	...	0.0	0.0	...	0.0	*	...	*

Worktable 13. Age-adjusted death rates for 358 selected causes, by race and sex: United States, 1999-2005

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992) and year	All races 1/			White 2/			Black 2/		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
Other complications predominately related to the puerperium (O85-O87,O89-O92)									
2005	0.0	...	0.0	0.0	...	0.0	0.1	...	0.1
2004	0.0	...	0.0	0.0	...	0.0	0.1	...	0.1
2003	0.0	...	0.0	0.0	...	0.0	*	...	*
2002	0.0	...	0.0	*	...	*	0.1	...	0.1
2001	0.0	...	0.0	*	...	*	*	...	*
2000	0.0	...	0.0	*	...	*	*	...	*
1999	0.0	...	0.0	*	...	*	*	...	*
All other direct obstetric causes (O10,O12,O21-O43,O47-O66,O68-O71,O73-O75)									
2005	0.1	...	0.1	0.0	...	0.1	0.2	...	0.4
2004	0.1	...	0.1	0.0	...	0.1	0.1	...	0.3
2003	0.0	...	0.1	0.0	...	0.1	0.2	...	0.3
2002	0.0	...	0.1	0.0	...	0.1	0.1	...	0.2
2001	0.0	...	0.1	0.0	...	0.0	0.1	...	0.2
2000	0.0	...	0.1	0.0	...	0.1	0.1	...	0.2
1999	0.0	...	0.1	0.0	...	0.0	0.1	...	0.2
Obstetric death of unspecified cause (O95)									
2005	0.0	...	0.0	0.0	...	0.0	*	...	*
2004	0.0	...	0.0	*	...	*	*	...	*
2003	0.0	...	0.0	*	...	*	*	...	*
2002	*	...	*	*	...	*	*	...	*
2001	*	...	*	*	...	*	*	...	*
2000	*	...	*	*	...	*	*	...	*
1999	*	...	*	*	...	*	*	...	*

Worktable 13. Age-adjusted death rates for 358 selected causes, by race and sex: United States, 1999-2005

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992) and year	All races 1/			White 2/			Black 2/		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
Other deaths related to pregnancy, childbirth and the puerperium (O96-O97)									
2005	0.0	...	0.1	0.0	...	0.1	0.1	...	0.2
2004	0.1	...	0.1	0.1	...	0.1	0.1	...	0.2
2003	0.0	...	0.0	0.0	...	0.0	*	...	*
2002	0.0	...	0.0	*	...	*	*	...	*
2001	*	...	*	*	...	*	*	...	*
2000	*	...	*	*	...	*	*	...	*
1999	*	...	*	*	...	*	*	...	*
Indirect obstetric deaths (O98-O99)									
2005	0.1	...	0.1	0.0	...	0.1	0.1	...	0.3
2004	0.0	...	0.1	0.0	...	0.1	0.1	...	0.2
2003	0.0	...	0.1	0.0	...	0.1	0.1	...	0.1
2002	0.0	...	0.0	0.0	...	0.0	*	...	*
2001	0.0	...	0.0	0.0	...	0.0	0.0	...	0.1
2000	0.0	...	0.0	0.0	...	0.0	*	...	*
1999	0.0	...	0.0	0.0	...	0.0	*	...	*
XVI. Certain conditions originating in the perinatal period (P00-P96)									
2005	4.9	5.4	4.4	3.8	4.2	3.4	10.5	11.5	9.5
2004	4.8	5.4	4.2	3.8	4.3	3.3	10.3	11.3	9.2
2003	4.9	5.5	4.3	4.0	4.4	3.5	10.2	11.4	9.0
2002	4.9	5.4	4.3	3.9	4.3	3.4	10.1	10.9	9.2
2001	4.7	5.3	4.2	3.7	4.1	3.3	10.3	11.3	9.1
2000	5.1	5.6	4.5	4.0	4.4	3.5	11.3	12.6	9.9
1999	5.2	5.7	4.6	4.0	4.4	3.6	11.8	13.1	10.4

Worktable 13. Age-adjusted death rates for 358 selected causes, by race and sex: United States, 1999-2005

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992) and year	All races 1/			White 2/			Black 2/		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
Newborn affected by maternal factors and by complications of pregnancy, labor and delivery (P00-P04)									
2005	1.1	1.1	1.0	0.9	0.9	0.8	2.3	2.4	2.3
2004	1.1	1.1	1.0	0.8	0.9	0.7	2.3	2.4	2.1
2003	1.1	1.2	1.0	0.9	1.0	0.8	2.1	2.3	2.0
2002	1.0	1.2	0.9	0.8	0.9	0.7	2.1	2.3	1.9
2001	1.0	1.1	0.9	0.8	0.9	0.7	2.0	2.2	1.8
2000	1.0	1.1	0.9	0.8	0.9	0.7	2.1	2.2	1.9
1999	1.0	1.1	0.9	0.8	0.8	0.7	2.2	2.4	2.0
Disorders related to short gestation and low birth weight, not elsewhere classified (P07)									
2005	1.6	1.7	1.4	1.1	1.2	1.0	4.0	4.4	3.5
2004	1.6	1.8	1.4	1.2	1.3	1.0	3.8	4.2	3.3
2003	1.7	1.8	1.5	1.2	1.4	1.1	4.0	4.5	3.5
2002	1.6	1.7	1.4	1.1	1.2	1.0	3.9	4.2	3.5
2001	1.5	1.6	1.4	1.1	1.2	1.0	3.9	4.2	3.6
2000	1.6	1.7	1.5	1.1	1.2	1.1	4.2	4.6	3.8
1999	1.6	1.7	1.4	1.1	1.2	1.0	4.3	4.8	3.9
Birth trauma (P10-P15)									
2005	0.0	0.0	*	0.0	*	*	*	*	*
2004	0.0	*	*	0.0	*	*	*	*	*
2003	0.0	*	*	0.0	*	*	*	*	*
2002	0.1	0.1	0.1	0.1	0.1	0.1	0.2	0.2	0.1
2001	0.0	0.1	0.0	0.0	0.0	0.0	0.1	0.1	*
2000	0.0	0.0	0.0	0.0	*	*	*	*	*
1999	0.0	*	*	*	*	*	*	*	*

Worktable 13. Age-adjusted death rates for 358 selected causes, by race and sex: United States, 1999-2005

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992) and year	All races 1/			White 2/			Black 2/		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
Intrauterine hypoxia and birth asphyxia (P20-P21)									
2005	0.2	0.2	0.2	0.2	0.2	0.1	0.3	0.3	0.2
2004	0.2	0.2	0.2	0.2	0.2	0.1	0.3	0.3	0.3
2003	0.2	0.2	0.2	0.2	0.2	0.2	0.3	0.3	0.3
2002	0.2	0.2	0.2	0.2	0.2	0.2	0.3	0.3	0.3
2001	0.2	0.2	0.2	0.2	0.2	0.2	0.3	0.3	0.3
2000	0.2	0.3	0.2	0.2	0.2	0.2	0.4	0.4	0.3
1999	0.2	0.2	0.2	0.2	0.2	0.2	0.4	0.4	0.3
Respiratory distress of newborn (P22)									
2005	0.3	0.3	0.2	0.2	0.3	0.2	0.6	0.7	0.5
2004	0.3	0.3	0.3	0.2	0.3	0.2	0.6	0.7	0.5
2003	0.3	0.3	0.2	0.2	0.3	0.2	0.5	0.6	0.5
2002	0.3	0.4	0.3	0.3	0.3	0.2	0.7	0.7	0.7
2001	0.3	0.4	0.3	0.3	0.3	0.3	0.7	0.9	0.6
2000	0.4	0.5	0.3	0.3	0.4	0.2	0.8	0.9	0.6
1999	0.4	0.5	0.3	0.3	0.4	0.3	0.9	1.0	0.7
Other respiratory conditions originating in the perinatal period (P23-P28)									
2005	0.4	0.5	0.3	0.3	0.4	0.3	0.8	0.9	0.8
2004	0.4	0.5	0.4	0.3	0.4	0.3	0.9	0.9	0.8
2003	0.4	0.5	0.3	0.4	0.4	0.3	0.8	1.0	0.7
2002	0.4	0.5	0.4	0.4	0.4	0.3	0.9	1.0	0.8
2001	0.5	0.6	0.5	0.4	0.5	0.4	1.1	1.2	1.0
2000	0.6	0.6	0.5	0.5	0.5	0.4	1.2	1.4	1.0
1999	0.6	0.7	0.5	0.5	0.6	0.4	1.2	1.4	1.1

Worktable 13. Age-adjusted death rates for 358 selected causes, by race and sex: United States, 1999-2005

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992) and year	All races 1/			White 2/			Black 2/		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
Infections specific to the perinatal period (P35-P39)									
2005	0.3	0.4	0.3	0.3	0.3	0.2	0.7	0.8	0.6
2004	0.4	0.4	0.3	0.3	0.3	0.2	0.7	0.8	0.6
2003	0.3	0.4	0.3	0.3	0.3	0.2	0.7	0.8	0.6
2002	0.3	0.4	0.3	0.3	0.3	0.2	0.6	0.6	0.6
2001	0.3	0.3	0.3	0.2	0.3	0.2	0.6	0.6	0.6
2000	0.3	0.4	0.3	0.3	0.3	0.2	0.7	0.8	0.6
1999	0.3	0.4	0.3	0.2	0.3	0.2	0.7	0.8	0.6
Other and unspecified conditions originating in the perinatal period (P05,P08,P29,P50-P96)									
2005	1.0	1.1	0.8	0.8	0.9	0.7	1.8	2.1	1.5
2004	0.9	1.0	0.8	0.7	0.9	0.6	1.7	1.9	1.5
2003	0.9	1.0	0.8	0.8	0.9	0.7	1.7	1.9	1.4
2002	0.8	0.9	0.7	0.7	0.8	0.6	1.4	1.6	1.3
2001	0.8	1.0	0.7	0.7	0.8	0.6	1.5	1.8	1.3
2000	1.0	1.1	0.8	0.8	0.9	0.7	1.9	2.3	1.6
1999	1.0	1.1	0.8	0.8	0.9	0.7	2.0	2.2	1.9
XVII. Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)									
2005	3.5	3.7	3.3	3.5	3.6	3.3	3.9	4.1	3.7
2004	3.6	3.7	3.4	3.5	3.7	3.4	4.0	4.4	3.7
2003	3.6	3.8	3.4	3.6	3.7	3.4	4.0	4.4	3.6
2002	3.7	3.8	3.5	3.7	3.8	3.5	4.1	4.4	3.9
2001	3.6	3.8	3.4	3.6	3.8	3.4	4.1	4.3	3.9
2000	3.8	4.1	3.6	3.7	4.0	3.5	4.4	4.8	4.1
1999	3.8	3.9	3.6	3.7	3.9	3.6	4.3	4.7	4.1

Worktable 13. Age-adjusted death rates for 358 selected causes, by race and sex: United States, 1999-2005

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992) and year	All races 1/			White 2/			Black 2/		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
Anencephaly and similar malformations (Q00)									
2005	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1
2004	0.1	0.1	0.1	0.1	0.1	0.1	0.1	*	*
2003	0.1	0.1	0.1	0.1	0.1	0.1	0.1	*	0.1
2002	0.1	0.1	0.1	0.1	0.1	0.1	0.1	*	0.1
2001	0.1	0.1	0.1	0.1	0.1	0.1	0.1	*	0.1
2000	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1
1999	0.1	0.1	0.1	0.1	0.1	0.1	0.1	*	0.1
Spina bifida (Q05)									
2005	0.1	0.1	0.1	0.1	0.1	0.1	0.1	*	*
2004	0.0	0.0	0.0	0.1	0.0	0.1	*	*	*
2003	0.0	0.0	0.0	0.0	0.0	0.0	*	*	*
2002	0.1	0.1	0.1	0.1	0.1	0.1	*	*	*
2001	0.0	0.0	0.1	0.0	0.0	0.1	*	*	*
2000	0.0	0.0	0.0	0.0	0.0	0.0	*	*	*
1999	0.0	0.0	0.0	0.1	0.0	0.1	*	*	*
All other congenital malformations of nervous system (Q01-Q04,Q06-Q07)									
2005	0.3	0.3	0.3	0.3	0.3	0.3	0.4	0.3	0.3
2004	0.3	0.3	0.3	0.3	0.3	0.3	0.4	0.4	0.4
2003	0.3	0.3	0.3	0.3	0.3	0.3	0.3	0.4	0.3
2002	0.3	0.3	0.3	0.3	0.3	0.3	0.3	0.3	0.3
2001	0.3	0.3	0.3	0.3	0.3	0.2	0.3	0.3	0.4
2000	0.3	0.3	0.3	0.3	0.3	0.3	0.4	0.4	0.3
1999	0.3	0.3	0.3	0.3	0.3	0.3	0.4	0.3	0.4

Worktable 13. Age-adjusted death rates for 358 selected causes, by race and sex: United States, 1999-2005

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992) and year	All races 1/			White 2/			Black 2/		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
Congenital malformations of heart (Q20-Q24)									
2005	1.0	1.0	0.9	1.0	1.1	0.9	1.1	1.1	1.1
2004	1.0	1.1	0.9	1.0	1.1	0.9	1.2	1.3	1.0
2003	1.1	1.2	1.0	1.1	1.2	1.0	1.2	1.3	1.1
2002	1.1	1.2	1.0	1.1	1.2	1.0	1.2	1.3	1.2
2001	1.1	1.2	1.0	1.1	1.2	1.0	1.4	1.4	1.3
2000	1.2	1.3	1.2	1.2	1.3	1.1	1.4	1.6	1.3
1999	1.3	1.4	1.2	1.3	1.4	1.2	1.5	1.7	1.3
Other congenital malformations of circulatory system (Q25-Q28)									
2005	0.2	0.3	0.2	0.2	0.2	0.2	0.3	0.3	0.3
2004	0.3	0.3	0.3	0.3	0.3	0.3	0.4	0.5	0.3
2003	0.3	0.3	0.3	0.3	0.3	0.3	0.4	0.4	0.3
2002	0.3	0.4	0.3	0.3	0.3	0.3	0.5	0.6	0.4
2001	0.3	0.3	0.3	0.3	0.3	0.2	0.3	0.4	0.3
2000	0.3	0.4	0.3	0.3	0.4	0.3	0.4	0.5	0.4
1999	0.3	0.4	0.3	0.3	0.3	0.3	0.4	0.5	0.4
Congenital malformations of respiratory system (Q30-Q34)									
2005	0.2	0.2	0.2	0.2	0.2	0.2	0.3	0.3	0.2
2004	0.2	0.3	0.2	0.2	0.3	0.2	0.3	0.4	0.3
2003	0.2	0.2	0.2	0.2	0.2	0.2	0.3	0.3	0.2
2002	0.2	0.2	0.2	0.2	0.2	0.2	0.3	0.3	0.3
2001	0.2	0.2	0.2	0.2	0.2	0.2	0.3	0.4	0.2
2000	0.3	0.3	0.2	0.2	0.3	0.2	0.4	0.4	0.3
1999	0.2	0.2	0.2	0.2	0.2	0.2	0.3	0.4	0.2

Worktable 13. Age-adjusted death rates for 358 selected causes, by race and sex: United States, 1999-2005

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992) and year	All races 1/			White 2/			Black 2/		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
Other congenital malformations and deformations (Q10-Q18,Q35-Q89)									
2005	0.9	1.1	0.8	0.9	1.0	0.8	1.1	1.3	0.9
2004	0.9	1.0	0.8	0.9	1.0	0.8	1.0	1.2	0.8
2003	0.9	1.0	0.8	0.9	1.0	0.8	1.1	1.3	0.9
2002	0.9	1.0	0.8	0.9	1.0	0.8	1.0	1.2	0.9
2001	0.9	1.0	0.8	0.9	1.0	0.8	1.0	1.2	0.9
2000	0.9	1.1	0.8	0.9	1.0	0.8	1.1	1.2	0.9
1999	0.9	1.1	0.9	0.9	1.0	0.8	1.1	1.2	0.9
Down's syndrome (Q90)									
2005	0.3	0.3	0.3	0.3	0.3	0.3	0.2	0.2	0.3
2004	0.3	0.3	0.2	0.3	0.3	0.3	0.2	0.3	0.2
2003	0.3	0.3	0.3	0.3	0.3	0.3	0.2	0.3	0.2
2002	0.3	0.3	0.3	0.3	0.3	0.3	0.2	0.3	0.2
2001	0.3	0.3	0.3	0.3	0.3	0.3	0.2	0.2	0.2
2000	0.3	0.3	0.3	0.3	0.3	0.3	0.2	0.2	0.2
1999	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2
All other chromosomal abnormalities, not elsewhere classified (Q91-Q99)									
2005	0.3	0.3	0.4	0.3	0.3	0.4	0.4	0.3	0.5
2004	0.4	0.3	0.4	0.3	0.3	0.4	0.4	0.4	0.5
2003	0.4	0.3	0.4	0.4	0.3	0.4	0.4	0.3	0.5
2002	0.4	0.3	0.4	0.4	0.3	0.4	0.4	0.3	0.5
2001	0.3	0.3	0.4	0.3	0.3	0.4	0.4	0.3	0.4
2000	0.4	0.3	0.4	0.3	0.3	0.4	0.4	0.4	0.5
1999	0.3	0.3	0.4	0.3	0.3	0.4	0.4	0.3	0.5

Worktable 13. Age-adjusted death rates for 358 selected causes, by race and sex: United States, 1999-2005

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992) and year	All races 1/			White 2/			Black 2/		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
XVIII. Symptoms, signs and abnormal clinical and laboratory findings, not elsewhere classified (R00-R99)									
2005	10.4	11.1	9.4	10.0	10.6	9.1	14.6	17.1	12.5
2004	10.0	10.7	9.0	9.6	10.2	8.7	14.0	16.0	12.1
2003	10.6	11.4	9.5	10.2	11.0	9.2	14.8	17.4	12.5
2002	10.3	11.4	9.0	9.9	10.8	8.7	14.5	18.0	11.8
2001	11.3	12.7	9.8	11.0	12.2	9.5	15.1	17.7	12.8
2000	11.4	13.1	9.7	11.0	12.6	9.3	16.5	19.7	13.6
1999	9.7	11.0	8.4	9.1	10.1	7.9	15.4	19.1	12.4
Senility (R54)									
2005	1.5	1.2	1.6	1.5	1.2	1.6	1.4	1.2	1.5
2004	1.6	1.3	1.7	1.6	1.3	1.7	1.3	1.1	1.4
2003	1.6	1.2	1.7	1.6	1.3	1.8	1.1	0.9	1.2
2002	1.5	1.2	1.6	1.5	1.3	1.6	1.2	1.1	1.2
2001	1.5	1.2	1.6	1.5	1.3	1.6	1.3	1.3	1.3
2000	1.4	1.1	1.5	1.4	1.2	1.5	1.4	1.0	1.5
1999	1.2	1.0	1.2	1.2	1.0	1.3	1.0	1.1	1.0
Ill-defined and unknown causes of mortality (R95-R99)									
2005	5.6	6.7	4.4	5.3	6.4	4.2	8.5	10.6	6.8
2004	5.2	6.3	4.1	4.9	6.0	3.9	7.7	9.4	6.2
2003	5.9	7.2	4.7	5.6	6.8	4.4	8.7	11.2	6.8
2002	5.9	7.2	4.6	5.6	6.8	4.4	8.9	11.7	6.7
2001	7.1	8.7	5.6	6.8	8.4	5.3	9.5	11.6	7.7
2000	7.4	9.1	5.8	7.0	8.7	5.4	10.9	13.9	8.4
1999	6.0	7.3	4.8	5.5	6.6	4.4	10.1	13.3	7.6

Worktable 13. Age-adjusted death rates for 358 selected causes, by race and sex: United States, 1999-2005

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992) and year	All races 1/			White 2/			Black 2/		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
Sudden infant death syndrome (R95)									
2005	0.8	0.9	0.6	0.6	0.8	0.5	1.4	1.5	1.3
2004	0.8	0.9	0.6	0.6	0.7	0.5	1.5	1.8	1.2
2003	0.7	0.9	0.6	0.6	0.7	0.5	1.4	1.5	1.4
2002	0.8	0.9	0.7	0.7	0.8	0.5	1.4	1.6	1.3
2001	0.8	0.9	0.6	0.6	0.7	0.5	1.6	1.7	1.4
2000	0.9	1.1	0.8	0.7	0.9	0.6	1.9	2.1	1.6
1999	1.0	1.1	0.8	0.8	0.9	0.7	1.9	2.1	1.7
Other ill-defined and unknown causes of mortality (R96-R99)									
2005	4.9	5.8	3.9	4.7	5.6	3.7	7.1	9.1	5.5
2004	4.4	5.4	3.5	4.3	5.2	3.3	6.2	7.6	5.0
2003	5.2	6.3	4.0	5.0	6.1	4.0	7.3	9.6	5.5
2002	5.1	6.3	4.0	4.9	6.0	3.8	7.5	10.0	5.4
2001	6.3	7.8	4.9	6.2	7.6	4.8	7.9	9.8	6.3
2000	6.5	8.1	5.0	6.3	7.8	4.8	9.0	11.8	6.8
1999	5.1	6.2	4.0	4.7	5.7	3.7	8.2	11.1	6.0
All other symptoms, signs and abnormal clinical and laboratory findings, not elsewhere classified (R00-R53,R55-R94)									
2005	3.3	3.1	3.3	3.2	3.0	3.3	4.7	5.3	4.2
2004	3.2	3.1	3.2	3.1	2.9	3.2	5.0	5.4	4.5
2003	3.2	3.0	3.1	3.0	2.9	3.0	4.9	5.3	4.5
2002	2.9	3.0	2.8	2.8	2.8	2.7	4.5	5.3	3.9
2001	2.8	2.8	2.7	2.6	2.6	2.6	4.3	4.8	3.8
2000	2.7	2.8	2.5	2.6	2.7	2.4	4.2	4.8	3.7
1999	2.6	2.7	2.4	2.4	2.5	2.3	4.2	4.8	3.8

Worktable 13. Age-adjusted death rates for 358 selected causes, by race and sex: United States, 1999-2005

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992) and year	All races 1/			White 2/			Black 2/		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
XX. External causes of mortality (*U01-*U03,V01-Y89)									
2005	58.7	85.0	34.0	58.3	83.3	34.7	68.7	109.3	33.3
2004	57.3	82.7	33.7	57.2	81.5	34.3	65.5	103.8	32.5
2003	57.0	82.9	33.0	56.5	81.2	33.4	66.0	104.9	32.4
2002	56.7	82.7	32.6	55.9	80.7	32.9	67.2	107.7	32.6
2001 3/	56.1	82.6	31.8	55.2	80.3	31.8	68.2	108.4	33.7
2000	53.9	79.5	30.5	52.5	76.6	30.4	67.9	109.1	32.9
1999	54.3	80.1	31.0	52.7	76.9	30.7	69.8	111.7	34.6
Accidents (unintentional injuries) (V01-X59,Y85-Y86)									
2005	39.1	54.2	25.0	40.1	55.2	25.8	38.7	58.1	22.8
2004	37.7	52.1	24.5	38.8	53.2	25.3	36.3	54.4	21.7
2003	37.3	51.8	24.1	38.2	52.8	24.7	36.1	53.9	21.6
2002	36.9	51.5	23.5	37.5	52.0	24.0	36.9	56.2	21.3
2001	35.7	50.2	22.5	36.0	50.3	22.8	37.6	56.9	22.0
2000	34.9	49.3	22.0	35.1	49.2	22.2	37.7	58.1	21.3
1999	35.3	49.8	22.3	35.2	49.3	22.4	40.1	61.3	23.1
Transport accidents (V01-V99,Y85)									
2005	16.2	23.4	9.3	16.7	23.9	9.7	15.4	24.1	8.0
2004	16.2	23.1	9.7	16.7	23.6	10.0	15.7	24.4	8.4
2003	16.4	23.4	9.8	16.8	23.9	10.0	15.8	24.5	8.6
2002	16.7	23.9	9.9	17.1	24.2	10.2	16.0	25.0	8.5
2001	16.6	23.7	9.8	16.8	24.0	10.0	16.5	25.5	8.9
2000	16.6	23.7	10.0	16.8	23.8	10.2	16.7	26.1	8.7
1999	16.6	23.6	10.2	16.8	23.6	10.4	17.1	26.5	9.5

Worktable 13. Age-adjusted death rates for 358 selected causes, by race and sex: United States, 1999-2005

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992) and year	All races 1/			White 2/			Black 2/		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
Railway accidents (V05,V15,V80.6,V81.2-V81.9)									
2005	0.2	0.2	0.1	0.1	0.2	0.0	0.2	0.4	*
2004	0.1	0.3	0.1	0.1	0.3	0.0	0.2	0.4	*
2003	0.1	0.2	0.0	0.1	0.3	0.0	0.2	0.4	*
2002	0.2	0.3	0.0	0.2	0.3	0.0	0.3	0.5	*
2001	0.2	0.3	0.1	0.2	0.3	0.0	0.2	0.4	*
2000	0.2	0.3	0.1	0.2	0.3	0.0	0.3	0.5	*
1999	0.2	0.3	0.0	0.2	0.3	0.1	0.2	0.4	*
Motor vehicle accidents (V02-V04,V09.0,V09.2,V12-V14,V19.0-V19.2,V19.4-V19.6,V20-V79,V80.3-V80.5,V81.0-V81.1,V82.0-V82.1,V83-V86,V87.0-V87.8, V88.0-V88.8,V89.0,V89.2)									
2005	15.2	21.7	8.9	15.6	22.2	9.2	14.5	22.5	7.6
2004	15.2	21.4	9.3	15.6	21.8	9.5	14.8	22.7	8.1
2003	15.3	21.6	9.3	15.7	21.9	9.5	14.9	22.7	8.3
2002	15.7	22.1	9.6	16.0	22.4	9.8	15.0	23.2	8.2
2001	15.3	21.8	9.3	15.6	22.0	9.5	15.4	23.5	8.5
2000	15.4	21.7	9.5	15.6	21.8	9.8	15.7	24.4	8.4
1999	15.2	21.3	9.6	15.3	21.3	9.7	15.9	24.3	9.0
Pedestrian involved in collision with motor vehicle (V02-V04)									
2005	1.1	1.6	0.7	1.1	1.5	0.6	1.5	2.4	0.7
2004	1.1	1.6	0.7	1.1	1.6	0.6	1.5	2.4	0.8
2003	1.2	1.8	0.7	1.2	1.7	0.7	1.6	2.6	0.8
2002	1.3	1.8	0.8	1.2	1.7	0.7	1.7	2.8	0.9
2001	1.3	1.8	0.7	1.2	1.7	0.6	1.8	2.7	1.0
2000	1.2	1.8	0.7	1.1	1.6	0.7	1.8	3.0	0.9
1999	1.2	1.8	0.7	1.1	1.6	0.7	1.9	3.2	0.9

Worktable 13. Age-adjusted death rates for 358 selected causes, by race and sex: United States, 1999-2005

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992) and year	All races 1/			White 2/			Black 2/		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
Pedalcyclist involved in collision with motor vehicle (V12-V14,V19.0-V19.2,V19.4-V19.6)									
2005	0.2	0.4	0.1	0.2	0.4	0.1	0.3	0.5	*
2004	0.2	0.4	0.1	0.2	0.4	0.1	0.2	0.5	*
2003	0.2	0.4	0.0	0.2	0.3	0.1	0.2	0.5	*
2002	0.2	0.3	0.0	0.2	0.4	0.0	0.2	0.4	*
2001	0.2	0.4	0.0	0.2	0.4	0.0	0.3	0.6	*
2000	0.2	0.4	0.0	0.2	0.4	0.0	0.3	0.6	*
1999	0.2	0.4	0.1	0.2	0.4	0.1	0.3	0.5	*
Motorcyclist involved in any accident except collision with railway train (V20-V24,V26-V34,V36-V39)									
2005	1.5	2.6	0.3	1.6	2.8	0.3	1.2	2.3	0.1
2004	1.4	2.5	0.3	1.5	2.6	0.3	1.0	2.1	*
2003	1.3	2.3	0.3	1.4	2.4	0.3	1.0	2.1	*
2002	1.1	2.0	0.2	1.2	2.2	0.2	0.8	1.7	*
2001	1.1	1.9	0.2	1.2	2.0	0.2	0.8	1.7	*
2000	1.0	1.8	0.2	1.0	1.9	0.2	0.9	1.8	*
1999	0.8	1.5	0.2	0.9	1.6	0.2	0.7	1.5	*
Motor vehicle accident involving collision with railway train (V25,V35,V45,V55,V65,V75,V81.0-V81.1,V87.6,V88.6)									
2005	0.1	0.1	0.0	0.1	0.1	0.0	*	*	*
2004	0.1	0.1	0.0	0.1	0.1	0.0	0.1	0.2	*
2003	0.1	0.1	0.0	0.1	0.1	0.0	0.1	*	*
2002	0.1	0.1	0.1	0.1	0.1	0.1	0.1	*	*
2001	0.1	0.1	0.1	0.1	0.1	0.1	0.1	*	*
2000	0.1	0.1	0.0	0.1	0.2	0.0	0.1	*	*
1999	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	*

Worktable 13. Age-adjusted death rates for 358 selected causes, by race and sex: United States, 1999-2005

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992) and year	All races 1/			White 2/			Black 2/		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
Motorcyclist involved in collision with railway train (V25,V35)									
2005	*	*	*	*	*	*	*	*	*
2004	*	*	*	*	*	*	*	*	*
2003	*	*	*	*	*	*	*	*	*
2002	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*
Other motor vehicle accident involving collision with railway train (V45,V55,V65,V75,V81.0-V81.1,V87.6,V88.6)									
2005	0.1	0.1	0.0	0.1	0.1	0.0	*	*	*
2004	0.1	0.1	0.0	0.1	0.1	0.0	0.1	0.2	*
2003	0.1	0.1	0.0	0.1	0.1	0.0	0.1	*	*
2002	0.1	0.1	0.0	0.1	0.1	0.1	0.1	*	*
2001	0.1	0.1	0.1	0.1	0.1	0.1	0.1	*	*
2000	0.1	0.1	0.0	0.1	0.2	0.0	0.1	*	*
1999	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	*
Occupant of motor vehicle involved in collision with other (non-motorized) road vehicle, streetcar, animal or pedestrian (V40-V41,V46-V47, V50-V51,V56-V57,V60-V61,V66-V67,V70-V71,V76-V77)									
2005	1.4	2.2	0.7	1.5	2.2	0.8	1.5	2.4	0.6
2004	1.6	2.3	0.8	1.6	2.4	0.8	1.7	2.6	0.9
2003	1.7	2.5	0.9	1.8	2.6	0.9	1.6	2.7	0.7
2002	1.7	2.6	0.9	1.8	2.6	1.0	1.9	3.1	0.8
2001	1.5	2.2	0.8	1.5	2.2	0.8	1.7	2.8	0.8
2000	1.4	2.0	0.7	1.4	2.0	0.8	1.6	2.6	0.7
1999	1.3	2.0	0.7	1.4	2.0	0.7	1.7	2.7	0.8

Worktable 13. Age-adjusted death rates for 358 selected causes, by race and sex: United States, 1999-2005

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992) and year	All races 1/			White 2/			Black 2/		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
Occupant of car, pickup truck or van involved in collision with other motor vehicle (V42-V44,V49,V52-V54,V59)									
2005	3.4	4.3	2.6	3.5	4.4	2.7	3.1	4.2	2.2
2004	3.6	4.5	2.8	3.7	4.6	2.9	3.2	4.4	2.3
2003	3.7	4.6	2.8	3.8	4.7	3.0	3.4	4.6	2.5
2002	3.9	5.0	2.9	4.0	5.1	3.0	3.5	4.8	2.4
2001	3.7	4.7	2.8	3.8	4.8	2.9	3.6	4.9	2.4
2000	3.7	4.6	2.9	3.8	4.7	3.0	3.5	4.9	2.3
1999	3.7	4.6	2.9	3.7	4.6	3.0	3.6	5.1	2.4
Occupant of heavy transport vehicle or bus involved in collision with other motor vehicle (V62-V64,V69,V72-V74,V79)									
2005	0.1	0.1	0.0	0.1	0.1	0.0	0.1	0.1	*
2004	0.1	0.1	0.0	0.1	0.1	*	0.1	0.2	*
2003	0.1	0.1	0.0	0.1	0.1	0.0	0.1	0.1	*
2002	0.1	0.1	0.0	0.1	0.1	0.0	0.1	0.2	*
2001	0.1	0.1	0.0	0.1	0.1	0.0	*	*	*
2000	0.1	0.1	0.0	0.1	0.1	*	0.1	0.2	*
1999	0.1	0.1	0.0	0.1	0.1	0.0	0.1	*	*
Occupant of motor vehicle involved in noncollision accident (V48,V58,V68,V78)									
2005	1.3	1.8	0.8	1.4	2.0	0.8	1.1	1.6	0.7
2004	1.4	1.9	0.8	1.4	2.0	0.8	1.2	1.7	0.7
2003	1.5	2.0	0.9	1.5	2.1	1.0	1.2	1.8	0.7
2002	1.5	2.1	0.9	1.6	2.2	1.0	1.2	1.9	0.6
2001	1.4	1.9	0.8	1.4	2.0	0.8	1.1	1.7	0.5
2000	1.3	1.9	0.7	1.4	2.0	0.8	1.1	1.7	0.6
1999	1.3	1.9	0.8	1.3	1.9	0.8	1.2	1.7	0.7

Worktable 13. Age-adjusted death rates for 358 selected causes, by race and sex: United States, 1999-2005

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992) and year	All races 1/			White 2/			Black 2/		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
Occupant of special-use motor vehicle involved in any accident (V83-V86)									
2005	0.4	0.8	0.1	0.5	0.8	0.1	0.1	0.3	*
2004	0.4	0.7	0.1	0.4	0.8	0.1	0.1	0.3	*
2003	0.4	0.7	0.1	0.4	0.8	0.1	0.1	0.2	*
2002	0.3	0.6	0.1	0.4	0.7	0.1	0.1	0.1	*
2001	0.4	0.7	0.1	0.4	0.7	0.1	0.1	0.2	*
2000	0.4	0.7	0.1	0.4	0.8	0.1	0.1	0.3	*
1999	0.4	0.7	0.1	0.4	0.8	0.1	0.1	0.3	*
Other and unspecified motor vehicle accidents (V09.0,V09.2,V80.3-V80.5,V82.0-V82.1,V87.0-V87.5,V87.7-V87.8,V88.0-V88.5,V88.7-V88.8,V89.0,V89.2)									
2005	5.6	7.7	3.6	5.6	7.7	3.7	5.7	8.7	3.2
2004	5.4	7.3	3.7	5.5	7.3	3.7	5.6	8.3	3.3
2003	5.2	7.1	3.5	5.2	7.1	3.5	5.5	8.0	3.3
2002	5.4	7.4	3.6	5.5	7.5	3.7	5.4	8.1	3.2
2001	5.7	7.9	3.8	5.8	7.9	3.9	5.8	8.7	3.5
2000	6.0	8.2	4.1	6.1	8.2	4.1	6.2	9.2	3.7
1999	6.0	8.1	4.1	6.1	8.2	4.2	6.2	9.1	3.9
Streetcar accidents (V82.2-V82.9)									
2005	*	*	*	*	*	*	*	*	*
2004	*	*	*	*	*	*	*	*	*
2003	*	*	*	*	*	*	*	*	*
2002	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*

Worktable 13. Age-adjusted death rates for 358 selected causes, by race and sex: United States, 1999-2005

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992) and year	All races 1/			White 2/			Black 2/		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
Other and unspecified land transport accidents (V01,V06,V09.1,V09.3,V09.9,V10-V11,V16-V18,V19.3,V19.8-V19.9,V80.0-V80.2,V80.7-V80.9,V87.9, V88.9,V89.1,V89.3,V89.9)									
2005	0.3	0.4	0.1	0.3	0.4	0.2	0.3	0.5	0.1
2004	0.3	0.3	0.1	0.3	0.3	0.2	0.3	0.5	0.1
2003	0.3	0.5	0.2	0.3	0.5	0.2	0.3	0.6	0.1
2002	0.2	0.4	0.1	0.2	0.3	0.1	0.2	0.4	*
2001	0.3	0.5	0.2	0.3	0.5	0.1	0.4	0.6	0.2
2000	0.4	0.5	0.2	0.4	0.5	0.2	0.3	0.5	0.2
1999	0.6	0.9	0.3	0.6	0.9	0.3	0.5	0.8	0.2
Water transport accidents (V90-V94)									
2005	0.2	0.3	0.0	0.2	0.3	0.0	0.1	0.2	*
2004	0.2	0.3	0.1	0.2	0.4	0.1	0.1	0.3	*
2003	0.2	0.3	0.1	0.2	0.3	0.1	0.2	0.4	*
2002	0.2	0.4	0.0	0.2	0.4	0.0	0.2	0.4	*
2001	0.2	0.4	0.1	0.2	0.4	0.1	0.2	0.4	*
2000	0.2	0.4	0.0	0.2	0.4	0.0	0.2	0.4	*
1999	0.2	0.4	0.1	0.2	0.4	0.1	0.3	0.5	*
Air and space transport accidents (V95-V97)									
2005	0.2	0.3	0.1	0.2	0.4	0.1	*	*	*
2004	0.2	0.4	0.1	0.3	0.5	0.1	*	*	*
2003	0.3	0.4	0.1	0.3	0.5	0.1	*	*	*
2002	0.2	0.4	0.0	0.3	0.5	0.1	*	*	*
2001	0.3	0.5	0.2	0.4	0.6	0.2	0.1	*	*
2000	0.3	0.5	0.1	0.3	0.5	0.1	*	*	*
1999	0.2	0.4	0.1	0.3	0.5	0.1	*	*	*

Worktable 13. Age-adjusted death rates for 358 selected causes, by race and sex: United States, 1999-2005

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992) and year	All races 1/			White 2/			Black 2/		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
Other and unspecified transport accidents and their sequelae (V98-V99,Y85)									
2005	0.2	0.4	0.1	0.2	0.4	0.1	0.3	0.4	0.1
2004	0.2	0.4	0.1	0.2	0.3	0.1	0.2	0.4	*
2003	0.2	0.3	0.1	0.2	0.3	0.1	0.2	0.4	*
2002	0.2	0.3	0.1	0.2	0.3	0.1	0.2	0.5	*
2001	0.2	0.3	0.1	0.2	0.3	0.1	0.3	0.5	*
2000	0.2	0.3	0.1	0.2	0.3	0.1	0.2	0.4	*
1999	0.2	0.3	0.1	0.2	0.3	0.1	0.3	0.4	0.1
Falls (W00-W19)									
2005	6.4	8.3	4.9	6.7	8.6	5.2	3.7	5.2	2.6
2004	6.3	8.2	4.8	6.6	8.6	5.1	3.6	5.5	2.3
2003	5.9	7.6	4.5	6.1	7.9	4.8	3.4	4.7	2.4
2002	5.6	7.4	4.3	5.9	7.6	4.6	3.1	4.9	2.0
2001	5.3	7.2	3.9	5.6	7.5	4.1	3.3	5.1	2.0
2000	4.8	6.6	3.6	5.0	6.8	3.7	2.9	4.6	1.8
1999	4.8	6.6	3.5	5.0	6.8	3.7	3.4	5.3	2.1
Fall on same level (W00-W09,W18)									
2005	2.3	2.9	1.9	2.4	3.0	2.0	1.2	1.7	0.8
2004	2.1	2.6	1.7	2.2	2.7	1.8	1.1	1.5	0.8
2003	1.9	2.4	1.6	2.0	2.5	1.7	0.7	1.0	0.5
2002	1.8	2.3	1.5	1.9	2.4	1.6	0.8	1.3	0.5
2001	1.4	1.8	1.1	1.5	1.9	1.2	0.7	1.0	0.5
2000	1.2	1.5	1.0	1.2	1.5	1.0	0.5	0.7	0.4
1999	0.8	1.1	0.7	0.8	1.1	0.7	0.6	0.8	0.4

Worktable 13. Age-adjusted death rates for 358 selected causes, by race and sex: United States, 1999-2005

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992) and year	All races 1/			White 2/			Black 2/		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
Fall from one level to another (W10-W17)									
2005	1.1	1.8	0.5	1.2	2.0	0.6	0.7	1.1	0.3
2004	1.1	1.8	0.5	1.2	1.9	0.6	0.7	1.3	0.3
2003	1.1	1.8	0.5	1.2	1.9	0.6	0.8	1.4	0.4
2002	1.1	1.9	0.5	1.2	2.0	0.6	0.7	1.2	0.4
2001	1.2	2.0	0.5	1.2	2.0	0.6	0.9	1.5	0.3
2000	1.0	1.8	0.4	1.1	1.9	0.5	0.7	1.3	0.3
1999	1.1	1.9	0.5	1.2	1.9	0.6	0.8	1.3	0.3
Unspecified fall (W19)									
2005	2.9	3.5	2.5	3.1	3.6	2.7	1.8	2.4	1.4
2004	3.1	3.8	2.5	3.2	4.0	2.7	1.8	2.7	1.2
2003	2.8	3.4	2.4	2.9	3.5	2.5	1.8	2.4	1.5
2002	2.6	3.2	2.2	2.8	3.3	2.4	1.6	2.4	1.1
2001	2.7	3.5	2.2	2.8	3.6	2.3	1.7	2.6	1.2
2000	2.6	3.3	2.1	2.7	3.4	2.2	1.7	2.5	1.2
1999	2.9	3.7	2.3	3.0	3.8	2.4	2.1	3.1	1.4
Accidental discharge of firearms (W32-W34)									
2005	0.3	0.5	0.1	0.3	0.4	0.1	0.4	0.7	*
2004	0.2	0.4	0.0	0.2	0.4	0.1	0.3	0.6	*
2003	0.2	0.5	0.1	0.2	0.5	0.1	0.3	0.6	*
2002	0.3	0.4	0.1	0.2	0.4	0.1	0.4	0.7	*
2001	0.3	0.5	0.1	0.3	0.5	0.1	0.3	0.5	*
2000	0.3	0.5	0.1	0.3	0.5	0.1	0.3	0.6	0.1
1999	0.3	0.5	0.1	0.3	0.5	0.1	0.4	0.7	0.1

Worktable 13. Age-adjusted death rates for 358 selected causes, by race and sex: United States, 1999-2005

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992) and year	All races 1/			White 2/			Black 2/		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
Accidental explosion (W39-W40)									
2005	0.0	0.1	0.0	0.0	0.1	*	0.1	*	*
2004	0.1	0.1	0.0	0.1	0.1	0.0	*	*	*
2003	0.1	0.1	0.0	0.1	0.1	*	*	*	*
2002	0.0	0.1	*	0.1	0.1	*	*	*	*
2001	0.1	0.1	0.0	0.1	0.1	0.0	*	*	*
2000	0.1	0.1	0.0	0.1	0.1	0.0	*	*	*
1999	0.1	0.1	0.0	0.1	0.1	0.0	*	*	*
Accidental drowning and submersion (W65-W74)									
2005	1.2	1.9	0.5	1.2	1.8	0.5	1.4	2.5	0.4
2004	1.1	1.8	0.5	1.1	1.7	0.5	1.3	2.2	0.5
2003	1.1	1.8	0.5	1.1	1.7	0.5	1.2	2.2	0.4
2002	1.2	2.0	0.5	1.1	1.8	0.4	1.6	2.7	0.5
2001	1.1	1.8	0.5	1.1	1.7	0.5	1.5	2.6	0.5
2000	1.2	2.0	0.5	1.2	1.8	0.5	1.5	2.6	0.6
1999	1.3	2.0	0.5	1.2	1.9	0.5	1.7	3.0	0.5
Accidental inhalation and ingestion of food or other objects causing obstruction of respiratory tract (W78-W80)									
2005	1.4	1.8	1.2	1.4	1.7	1.2	2.0	2.4	1.6
2004	1.5	1.9	1.2	1.5	1.9	1.2	1.9	2.5	1.6
2003	1.5	1.9	1.2	1.4	1.9	1.2	1.8	2.3	1.4
2002	1.4	1.8	1.2	1.4	1.8	1.1	2.1	2.7	1.6
2001	1.5	1.9	1.2	1.4	1.8	1.2	2.2	3.1	1.6
2000	1.6	2.0	1.3	1.5	1.9	1.3	2.1	2.9	1.7
1999	1.4	1.8	1.2	1.4	1.7	1.2	2.2	2.9	1.7

Worktable 13. Age-adjusted death rates for 358 selected causes, by race and sex: United States, 1999-2005

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992) and year	All races 1/			White 2/			Black 2/		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
Other accidental and unspecified threats to breathing (W75-W77,W81-W84)									
2005	0.5	0.7	0.3	0.5	0.7	0.3	0.6	0.8	0.5
2004	0.5	0.6	0.3	0.5	0.7	0.3	0.6	0.7	0.5
2003	0.4	0.6	0.3	0.4	0.6	0.2	0.6	0.8	0.4
2002	0.5	0.7	0.3	0.5	0.6	0.3	0.7	0.8	0.5
2001	0.5	0.6	0.3	0.5	0.6	0.3	0.6	0.8	0.5
2000	0.5	0.7	0.3	0.5	0.7	0.3	0.6	0.8	0.4
1999	0.6	0.8	0.4	0.6	0.8	0.4	0.7	1.0	0.5
Accidental exposure to electric current (W85-W87)									
2005	0.1	0.2	0.0	0.1	0.3	*	0.1	0.2	*
2004	0.1	0.2	0.0	0.1	0.3	*	0.1	0.2	*
2003	0.1	0.2	*	0.1	0.3	*	0.1	0.2	*
2002	0.2	0.3	0.0	0.2	0.3	0.0	0.1	0.2	*
2001	0.1	0.3	*	0.2	0.3	*	0.1	0.2	*
2000	0.1	0.3	0.0	0.2	0.3	*	0.1	0.2	*
1999	0.2	0.3	0.0	0.2	0.3	0.0	0.1	0.2	*
Accidental exposure to radiation (W88-W91)									
2005	*	*	*	*	*	*	*	*	*
2004	*	*	*	*	*	*	*	*	*
2003	*	*	*	*	*	*	*	*	*
2002	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*

Worktable 13. Age-adjusted death rates for 358 selected causes, by race and sex: United States, 1999-2005

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992) and year	All races 1/			White 2/			Black 2/		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
Accidental exposure to smoke, fire and flames (X00-X09)									
2005	1.1	1.4	0.8	0.9	1.2	0.7	2.2	3.1	1.6
2004	1.1	1.4	0.8	0.9	1.2	0.7	2.4	3.3	1.8
2003	1.2	1.4	0.9	1.1	1.3	0.8	2.4	3.2	1.9
2002	1.1	1.4	0.8	0.9	1.3	0.7	2.4	3.3	1.7
2001	1.2	1.5	0.9	1.0	1.3	0.7	2.7	3.8	1.9
2000	1.2	1.6	0.9	1.0	1.3	0.7	3.0	4.2	2.2
1999	1.2	1.6	0.9	1.0	1.4	0.7	3.0	4.0	2.3
Accidental contact with heat and hot substances (X10-X19)									
2005	0.0	0.0	0.0	0.0	0.0	0.0	*	*	*
2004	0.0	0.0	0.0	0.0	0.0	0.0	0.1	*	*
2003	0.0	0.0	0.0	0.0	0.0	0.0	*	*	*
2002	0.0	0.0	0.0	0.0	0.0	0.0	0.1	*	*
2001	0.0	0.0	0.0	0.0	0.0	0.0	0.1	*	*
2000	0.0	0.0	0.0	0.0	0.0	0.0	0.1	*	*
1999	0.0	0.0	0.0	0.0	0.0	0.0	0.1	*	*
Victim of lightning (X33)									
2005	0.0	0.0	*	0.0	0.0	*	*	*	*
2004	0.0	0.0	*	0.0	0.0	*	*	*	*
2003	0.0	0.0	*	0.0	0.0	*	*	*	*
2002	0.0	0.0	*	0.0	0.1	*	*	*	*
2001	0.0	0.0	*	0.0	0.0	*	*	*	*
2000	0.0	0.0	*	0.0	0.0	*	*	*	*
1999	0.0	0.1	*	0.0	0.1	*	*	*	*

Worktable 13. Age-adjusted death rates for 358 selected causes, by race and sex: United States, 1999-2005

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992) and year	All races 1/			White 2/			Black 2/		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
Earthquake, volcanic eruption, avalanche, landslide and other earth movements (X34-X36)									
2005	0.0	0.0	*	0.0	0.0	*	*	*	*
2004	0.0	0.0	*	0.0	0.0	*	*	*	*
2003	0.0	0.0	*	0.0	0.0	*	*	*	*
2002	0.0	0.0	*	0.0	0.0	*	*	*	*
2001	0.0	0.0	*	0.0	0.0	*	*	*	*
2000	0.0	0.0	*	0.0	0.0	*	*	*	*
1999	0.0	0.0	*	0.0	0.0	*	*	*	*
Cataclysmic storm and flood (X37-X38)									
2005	0.3	0.3	0.2	0.2	0.2	0.2	1.3	2.0	0.9
2004	0.0	0.0	0.0	0.0	0.0	0.0	*	*	*
2003	0.0	0.0	0.0	0.0	0.0	0.0	*	*	*
2002	0.0	0.0	0.0	0.0	0.0	0.0	*	*	*
2001	0.0	0.0	0.0	0.0	0.0	0.0	*	*	*
2000	0.0	0.0	0.0	0.0	0.0	*	*	*	*
1999	0.1	0.1	0.1	0.1	0.0	0.1	0.1	*	*
Accidental poisoning and exposure to noxious substances (X40-X49)									
2005	7.9	10.7	5.1	8.4	11.3	5.5	8.2	11.9	4.9
2004	7.1	9.5	4.7	7.6	10.1	5.0	6.9	9.9	4.5
2003	6.7	9.1	4.3	7.1	9.6	4.5	6.8	9.9	4.1
2002	6.1	8.4	3.7	6.3	8.7	3.9	6.8	10.1	3.9
2001	4.9	7.0	2.9	5.0	7.1	2.9	6.3	9.5	3.7
2000	4.5	6.6	2.5	4.5	6.6	2.5	6.0	9.3	3.3
1999	4.4	6.4	2.3	4.3	6.3	2.3	6.2	9.8	3.2

Worktable 13. Age-adjusted death rates for 358 selected causes, by race and sex: United States, 1999-2005

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992) and year	All races 1/			White 2/			Black 2/		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
Accidental poisoning by and exposure to drugs and other biological substances (X40-X44)									
2005	7.6	10.1	5.0	8.0	10.7	5.3	7.7	11.2	4.7
2004	6.8	9.0	4.5	7.2	9.5	4.8	6.6	9.3	4.3
2003	6.3	8.5	4.1	6.7	9.0	4.3	6.4	9.2	4.0
2002	5.7	7.8	3.5	5.9	8.1	3.7	6.3	9.3	3.7
2001	4.6	6.5	2.7	4.7	6.5	2.7	5.9	8.7	3.3
2000	4.1	6.0	2.3	4.2	6.0	2.3	5.6	8.6	3.1
1999	4.0	5.9	2.1	3.9	5.8	2.1	5.8	9.0	2.9
Accidental poisoning by and exposure to other and unspecified solid or liquids substances (X45-X46,X48-X49)									
2005	0.2	0.3	0.1	0.2	0.3	0.1	0.1	0.2	*
2004	0.2	0.3	0.1	0.2	0.3	0.1	0.1	0.2	*
2003	0.2	0.3	0.1	0.2	0.3	0.1	0.1	0.3	*
2002	0.2	0.3	0.1	0.2	0.3	0.1	0.2	0.4	0.1
2001	0.1	0.2	0.1	0.1	0.2	0.1	0.2	0.2	0.1
2000	0.2	0.2	0.1	0.2	0.2	0.1	0.2	0.3	*
1999	0.2	0.2	0.1	0.2	0.2	0.1	0.2	0.3	*
Accidental poisoning by and exposure to other gases and vapors (X47)									
2005	0.2	0.3	0.1	0.2	0.3	0.1	0.3	0.5	0.2
2004	0.2	0.3	0.1	0.2	0.3	0.1	0.2	0.3	0.1
2003	0.2	0.3	0.1	0.2	0.3	0.1	0.3	0.4	0.1
2002	0.2	0.4	0.1	0.3	0.3	0.1	0.2	0.4	0.1
2001	0.2	0.3	0.1	0.2	0.3	0.1	0.3	0.5	0.2
2000	0.2	0.3	0.1	0.2	0.3	0.1	0.2	0.4	0.1
1999	0.2	0.3	0.1	0.2	0.3	0.1	0.3	0.4	0.1

Worktable 13. Age-adjusted death rates for 358 selected causes, by race and sex: United States, 1999-2005

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992) and year	All races 1/			White 2/			Black 2/		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
All other and unspecified accidents and their sequelae (W20-W31,W35-W38,W41-W64,W92-W99,X20-X32,X39,X50-X59,Y86)									
2005	3.6	4.9	2.4	3.7	5.0	2.4	3.4	5.0	2.1
2004	3.4	4.6	2.4	3.5	4.7	2.4	3.3	4.9	2.0
2003	3.6	4.9	2.5	3.7	4.9	2.6	3.6	5.3	2.3
2002	3.7	5.0	2.6	3.8	5.0	2.7	3.7	5.4	2.4
2001	4.0	5.3	2.9	4.1	5.4	2.9	3.9	5.5	2.7
2000	4.0	5.3	2.8	4.0	5.3	2.8	4.0	6.5	2.3
1999	4.4	5.9	3.1	4.4	5.8	3.1	4.9	7.7	3.0
Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)									
2005	10.9	18.0	4.4	12.0	19.6	4.9	5.2	9.2	1.9
2004	10.9	18.0	4.5	12.0	19.6	5.0	5.3	9.6	1.8
2003	10.8	18.0	4.2	11.8	19.6	4.6	5.2	9.2	1.9
2002	10.9	18.4	4.2	12.0	20.0	4.7	5.3	9.8	1.6
2001 3/	10.7	18.2	4.0	11.7	19.6	4.5	5.5	9.8	1.8
2000	10.4	17.7	4.0	11.3	19.1	4.3	5.5	10.0	1.8
1999	10.5	17.8	4.0	11.3	19.0	4.4	5.6	10.3	1.6
Intentional self-poisoning (suicide) by and exposure to drugs and other biological substances (X60-X64)									
2005	1.4	1.3	1.5	1.6	1.5	1.7	0.5	0.5	0.5
2004	1.4	1.3	1.5	1.6	1.5	1.7	0.4	0.4	0.5
2003	1.3	1.3	1.4	1.5	1.5	1.5	0.5	0.4	0.6
2002	1.3	1.3	1.4	1.5	1.4	1.6	0.4	0.5	0.4
2001	1.3	1.2	1.3	1.4	1.4	1.4	0.5	0.4	0.5
2000	1.2	1.1	1.2	1.3	1.3	1.4	0.5	0.5	0.4
1999	1.1	1.1	1.2	1.3	1.2	1.3	0.4	0.5	0.4

Worktable 13. Age-adjusted death rates for 358 selected causes, by race and sex: United States, 1999-2005

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992) and year	All races 1/			White 2/			Black 2/		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
Intentional self-poisoning (suicide) by and exposure to other and unspecified solid or liquid substances and their vapors (X65-X66,X68-X69)									
2005	0.1	0.1	0.0	0.1	0.1	0.0	*	*	*
2004	0.1	0.1	0.1	0.1	0.1	0.1	*	*	*
2003	0.1	0.1	0.0	0.1	0.1	0.0	*	*	*
2002	0.1	0.1	0.0	0.1	0.1	0.0	*	*	*
2001	0.1	0.1	0.0	0.1	0.1	0.0	*	*	*
2000	0.1	0.1	0.0	0.1	0.1	0.0	*	*	*
1999	0.1	0.1	0.0	0.1	0.1	0.0	*	*	*
Intentional self-poisoning (suicide) by and exposure to other gases and vapors (X67)									
2005	0.4	0.7	0.2	0.5	0.8	0.2	0.1	0.2	*
2004	0.5	0.7	0.2	0.5	0.9	0.2	0.1	0.2	*
2003	0.5	0.8	0.2	0.5	0.9	0.2	0.1	0.2	*
2002	0.5	0.8	0.2	0.6	0.9	0.2	0.1	0.2	*
2001	0.5	0.8	0.2	0.6	0.9	0.3	0.1	0.2	*
2000	0.5	0.8	0.2	0.6	0.9	0.3	0.1	0.2	*
1999	0.5	0.9	0.3	0.6	1.0	0.3	0.1	0.2	*
Intentional self-harm (suicide) by hanging, strangulation and suffocation (X70)									
2005	2.4	4.0	0.9	2.6	4.3	0.9	1.2	2.1	0.5
2004	2.5	4.1	0.9	2.6	4.4	0.9	1.4	2.4	0.4
2003	2.3	3.9	0.8	2.4	4.0	0.8	1.3	2.3	0.4
2002	2.2	3.7	0.8	2.4	4.0	0.8	1.1	2.1	0.3
2001	2.2	3.7	0.7	2.3	3.9	0.7	1.1	2.0	0.3
2000	2.0	3.4	0.7	2.1	3.6	0.7	1.2	2.1	0.3
1999	1.9	3.3	0.6	2.0	3.4	0.7	1.0	2.0	0.2

Worktable 13. Age-adjusted death rates for 358 selected causes, by race and sex: United States, 1999-2005

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992) and year	All races 1/			White 2/			Black 2/		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
Intentional self-harm (suicide) by discharge of firearms (X72-X74)									
2005	5.7	10.5	1.4	6.3	11.5	1.6	2.7	5.3	0.5
2004	5.6	10.3	1.5	6.3	11.4	1.7	2.7	5.4	0.5
2003	5.8	10.7	1.4	6.4	11.8	1.6	2.7	5.3	0.6
2002	5.9	11.1	1.4	6.5	12.1	1.6	3.0	5.9	0.5
2001	5.9	11.0	1.4	6.5	12.0	1.6	3.1	6.0	0.6
2000	5.9	11.0	1.5	6.5	11.9	1.6	3.1	6.0	0.7
1999	6.0	11.1	1.5	6.5	12.0	1.7	3.3	6.5	0.6
Intentional self-harm (suicide) by jumping from a high place (X80)									
2005	0.2	0.3	0.1	0.2	0.4	0.1	0.2	0.3	0.1
2004	0.2	0.3	0.1	0.2	0.3	0.1	0.2	0.3	*
2003	0.2	0.3	0.1	0.2	0.4	0.1	0.2	0.3	*
2002	0.2	0.4	0.1	0.3	0.4	0.1	0.2	0.3	*
2001	0.2	0.3	0.1	0.2	0.3	0.1	0.2	0.3	*
2000	0.2	0.3	0.1	0.2	0.3	0.1	0.2	0.3	*
1999	0.3	0.4	0.1	0.3	0.4	0.1	0.2	0.3	*
Intentional self-harm (suicide) by all other and unspecified means and their sequelae (*U03,X71,X75-X79,X81-X84,X87.0)									
2005	0.7	1.0	0.3	0.7	1.0	0.3	0.5	0.8	0.3
2004	0.6	1.0	0.3	0.7	1.0	0.3	0.5	0.8	0.2
2003	0.6	0.9	0.3	0.6	0.9	0.3	0.5	0.7	0.2
2002	0.6	1.0	0.3	0.7	1.0	0.3	0.5	0.8	0.2
2001 3/	0.6	0.9	0.3	0.6	1.0	0.3	0.5	0.7	0.3
2000	0.6	0.9	0.3	0.6	0.9	0.3	0.5	0.8	0.2
1999	0.6	0.9	0.3	0.6	0.9	0.3	0.5	0.8	0.3

Worktable 13. Age-adjusted death rates for 358 selected causes, by race and sex: United States, 1999-2005

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992) and year	All races 1/			White 2/			Black 2/		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)									
2005	6.1	9.6	2.5	3.7	5.3	1.9	21.1	37.3	6.1
2004	5.9	9.2	2.5	3.6	5.3	1.9	20.1	35.1	6.3
2003	6.0	9.4	2.6	3.7	5.3	2.0	21.0	36.7	6.4
2002	6.1	9.4	2.8	3.7	5.3	2.0	21.0	36.4	6.9
2001 3/	7.1	10.8	3.3	4.9	7.1	2.6	21.2	36.2	7.4
2000	5.9	9.0	2.8	3.6	5.2	2.1	20.5	35.4	7.1
1999	6.0	9.1	2.9	3.8	5.4	2.1	20.1	34.2	7.3
Assault (homicide) by drugs, biological substances and other and unspecified noxious substances (*U01.6-*U01.7,X85-X90)									
2005	0.0	0.0	0.0	0.0	0.0	0.0	0.1	*	*
2004	0.0	0.0	0.0	0.0	0.0	0.0	0.1	*	*
2003	0.0	0.0	0.0	0.0	0.0	0.0	0.1	*	*
2002	0.0	0.0	0.0	0.0	0.0	*	*	*	*
2001	0.0	0.0	0.0	0.0	0.0	0.0	*	*	*
2000	0.0	0.0	0.0	0.0	0.0	0.0	*	*	*
1999	0.0	0.0	0.0	0.0	0.0	0.0	*	*	*
Assault (homicide) by hanging, strangulation and suffocation (X91)									
2005	0.2	0.1	0.3	0.2	0.1	0.2	0.5	0.4	0.6
2004	0.2	0.1	0.3	0.2	0.1	0.2	0.5	0.4	0.7
2003	0.2	0.1	0.3	0.2	0.1	0.2	0.6	0.4	0.7
2002	0.2	0.2	0.3	0.2	0.1	0.2	0.6	0.4	0.7
2001	0.2	0.2	0.3	0.2	0.1	0.2	0.6	0.4	0.7
2000	0.2	0.2	0.3	0.2	0.1	0.2	0.6	0.3	0.8
1999	0.2	0.2	0.3	0.2	0.2	0.2	0.6	0.4	0.8

Worktable 13. Age-adjusted death rates for 358 selected causes, by race and sex: United States, 1999-2005

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992) and year	All races 1/			White 2/			Black 2/		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
Assault (homicide) by discharge of firearms (*U01.4,X93-X95)									
2005	4.2	7.0	1.2	2.2	3.5	0.9	16.0	29.8	3.0
2004	4.0	6.6	1.1	2.2	3.4	0.8	15.0	28.0	2.9
2003	4.1	6.8	1.2	2.2	3.4	0.9	15.7	29.0	3.2
2002	4.1	6.8	1.3	2.2	3.4	1.0	15.6	28.7	3.5
2001	3.9	6.6	1.3	2.2	3.5	1.0	14.8	27.4	3.0
2000	3.8	6.3	1.2	2.1	3.2	0.9	14.6	26.9	3.1
1999	3.8	6.3	1.3	2.1	3.3	1.0	14.3	26.1	3.5
Assault (homicide) by sharp object (X99)									
2005	0.7	1.0	0.4	0.5	0.7	0.3	2.2	3.4	1.1
2004	0.7	1.0	0.4	0.5	0.7	0.3	2.2	3.3	1.2
2003	0.7	1.0	0.4	0.5	0.7	0.3	2.2	3.5	1.1
2002	0.7	1.0	0.4	0.5	0.7	0.3	2.2	3.3	1.2
2001	0.7	1.0	0.4	0.5	0.6	0.3	2.1	3.1	1.2
2000	0.6	0.9	0.4	0.5	0.7	0.3	2.0	2.9	1.1
1999	0.7	1.0	0.4	0.4	0.6	0.3	2.0	3.3	1.0
Assault (homicide) by blunt object (Y00)									
2005	0.0	0.1	0.0	0.0	0.1	0.0	0.1	0.1	*
2004	0.0	0.0	0.0	0.0	0.0	0.0	0.1	*	*
2003	0.0	0.1	0.0	0.0	0.1	0.0	0.1	0.2	*
2002	0.1	0.1	0.0	0.1	0.1	0.0	0.1	0.3	*
2001	0.1	0.1	0.1	0.1	0.1	0.0	0.2	0.3	*
2000	0.1	0.1	0.1	0.1	0.1	0.0	0.2	0.3	0.1
1999	0.1	0.1	0.0	0.1	0.1	0.0	0.2	0.3	0.1

Worktable 13. Age-adjusted death rates for 358 selected causes, by race and sex: United States, 1999-2005

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992) and year	All races 1/			White 2/			Black 2/		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
Assault (homicide) by bodily force (Y04)									
2005	0.0	0.0	*	0.0	0.0	*	*	*	*
2004	0.0	0.0	*	0.0	0.0	*	0.1	*	*
2003	0.0	0.1	*	0.0	0.1	*	0.1	0.2	*
2002	0.0	0.1	*	0.0	0.0	*	0.1	0.2	*
2001	0.1	0.1	0.0	0.1	0.1	0.0	0.1	0.2	*
2000	0.1	0.1	*	0.0	0.1	*	0.1	0.2	*
1999	0.1	0.1	0.0	0.0	0.1	*	0.2	0.3	*
Sexual assault (homicide) by bodily force (Y05)									
2005	*	*	*	*	*	*	*	*	*
2004	*	*	*	*	*	*	*	*	*
2003	*	*	*	*	*	*	*	*	*
2002	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*
Neglect, abandonment and other maltreatment syndromes (Y06-Y07)									
2005	0.1	0.1	0.1	0.0	0.0	0.1	0.1	0.2	0.1
2004	0.1	0.1	0.1	0.0	0.1	0.0	0.1	0.1	0.1
2003	0.1	0.1	0.1	0.1	0.1	0.0	0.1	0.2	0.1
2002	0.1	0.1	0.1	0.1	0.1	0.1	0.2	0.2	0.2
2001	0.1	0.1	0.1	0.1	0.1	0.0	0.2	0.2	0.2
2000	0.1	0.1	0.1	0.1	0.1	0.0	0.2	0.2	0.2
1999	0.1	0.1	0.1	0.0	0.1	0.0	0.2	0.2	0.2

Worktable 13. Age-adjusted death rates for 358 selected causes, by race and sex: United States, 1999-2005

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992) and year	All races 1/			White 2/			Black 2/		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
Assault (homicide) by all other and unspecified means and their sequelae (*U01.0-*U01.3,*U01.5,*U01.8-*U01.9,*U02,X92,X96-X98,Y01-Y03,Y08-Y09,Y87.1)									
2005	0.9	1.2	0.5	0.7	0.9	0.4	2.1	3.2	1.2
2004	0.9	1.1	0.6	0.7	0.9	0.4	2.0	2.9	1.2
2003	0.9	1.2	0.6	0.7	0.9	0.5	2.1	3.2	1.2
2002	0.8	1.1	0.5	0.6	0.9	0.4	2.2	3.4	1.2
2001 3/	2.0	2.8	1.1	1.8	2.6	1.0	3.2	4.6	2.1
2000	1.0	1.4	0.7	0.7	0.9	0.5	2.9	4.4	1.7
1999	1.0	1.3	0.7	0.8	1.0	0.5	2.5	3.6	1.5
Event of undetermined intent (Y10-Y34,Y87.2,Y89.9)									
2005	1.6	1.9	1.2	1.6	1.9	1.3	1.8	2.5	1.2
2004	1.7	2.2	1.2	1.7	2.2	1.3	1.9	2.7	1.1
2003	1.7	2.3	1.2	1.8	2.3	1.3	1.9	2.9	1.0
2002	1.7	2.2	1.2	1.7	2.2	1.2	2.1	3.1	1.2
2001	1.5	2.0	0.9	1.5	2.0	1.0	1.8	2.9	0.9
2000	1.3	1.9	0.8	1.3	1.8	0.8	1.9	3.0	1.0
1999	1.4	2.0	0.9	1.4	1.8	0.9	2.0	3.2	0.9
Poisoning by and exposure to drugs and biological substances, undetermined intent (Y10-Y14)									
2005	1.0	1.2	0.9	1.1	1.2	1.0	1.0	1.2	0.8
2004	1.1	1.4	0.9	1.2	1.4	1.0	1.1	1.5	0.7
2003	1.2	1.5	0.9	1.3	1.6	1.0	1.2	1.8	0.7
2002	1.1	1.3	0.8	1.1	1.4	0.9	1.3	1.7	0.9
2001	1.0	1.3	0.7	1.0	1.3	0.7	1.1	1.8	0.5
2000	0.8	1.2	0.6	0.9	1.2	0.6	1.1	1.6	0.6
1999	0.9	1.2	0.6	0.9	1.1	0.6	1.2	1.9	0.6

Worktable 13. Age-adjusted death rates for 358 selected causes, by race and sex: United States, 1999-2005

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992) and year	All races 1/			White 2/			Black 2/		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
Poisoning by and exposure to other and unspecified solid or liquid substance, undetermined intent (Y15-Y16,Y18-Y19)									
2005	0.0	0.0	0.0	0.0	0.0	0.0	*	*	*
2004	0.0	0.0	*	0.0	0.0	*	*	*	*
2003	0.0	0.0	0.0	0.0	0.0	*	*	*	*
2002	0.0	0.0	*	0.0	0.0	*	*	*	*
2001	0.0	0.0	*	0.0	0.0	*	*	*	*
2000	0.0	0.0	*	0.0	0.0	*	*	*	*
1999	0.0	0.0	*	0.0	0.0	*	*	*	*
Poisoning by and exposure to other gases and vapors, undetermined intent (Y17)									
2005	0.0	0.0	0.0	0.0	0.0	0.0	*	*	*
2004	0.0	0.0	*	0.0	0.0	*	*	*	*
2003	0.0	0.1	0.0	0.0	0.1	0.0	*	*	*
2002	0.0	0.0	*	0.0	0.0	*	*	*	*
2001	0.0	0.1	*	0.0	0.1	*	*	*	*
2000	0.0	0.0	0.0	0.0	0.0	*	*	*	*
1999	0.0	0.0	0.0	0.0	0.0	0.0	*	*	*
Discharge of firearms, undetermined intent (Y22-Y24)									
2005	0.1	0.1	0.0	0.1	0.1	0.0	0.1	0.1	*
2004	0.1	0.1	0.0	0.1	0.1	0.0	0.1	0.1	*
2003	0.1	0.1	0.0	0.1	0.1	0.0	0.1	0.2	*
2002	0.1	0.1	0.0	0.1	0.1	0.0	0.1	0.2	*
2001	0.1	0.1	0.0	0.1	0.1	0.0	0.1	0.2	*
2000	0.1	0.1	0.0	0.1	0.1	0.0	0.1	0.3	*
1999	0.1	0.2	0.0	0.1	0.2	0.0	0.2	0.3	*

Worktable 13. Age-adjusted death rates for 358 selected causes, by race and sex: United States, 1999-2005

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992) and year	All races 1/			White 2/			Black 2/		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
Falling, jumping or pushed from a high place, undetermined intent (Y30)									
2005	0.0	0.0	*	0.0	0.0	*	*	*	*
2004	0.0	0.0	0.0	0.0	0.0	*	*	*	*
2003	0.0	0.0	0.0	0.0	0.0	*	*	*	*
2002	0.0	0.0	0.0	0.0	0.1	0.0	*	*	*
2001	0.0	0.0	0.0	0.0	0.0	*	*	*	*
2000	0.0	0.0	*	0.0	0.0	*	*	*	*
1999	0.0	0.0	*	0.0	0.0	*	*	*	*
All other and unspecified events of undetermined intent and their sequelae (Y20-Y21,Y25-Y29,Y31-Y34,Y87.2,Y89.9)									
2005	0.4	0.6	0.3	0.3	0.5	0.2	0.7	1.1	0.3
2004	0.4	0.6	0.3	0.4	0.5	0.2	0.7	1.0	0.4
2003	0.4	0.5	0.2	0.3	0.5	0.2	0.5	0.8	0.3
2002	0.4	0.5	0.2	0.4	0.5	0.2	0.6	1.1	0.3
2001	0.3	0.5	0.2	0.3	0.4	0.2	0.5	0.8	0.3
2000	0.3	0.5	0.2	0.3	0.4	0.2	0.6	1.0	0.3
1999	0.3	0.5	0.2	0.3	0.4	0.2	0.5	0.9	0.2
Legal intervention (Y35,Y89.0)									
2005	0.1	0.3	*	0.1	0.2	*	0.3	0.6	*
2004	0.1	0.3	*	0.1	0.2	*	0.2	0.5	*
2003	0.1	0.3	*	0.1	0.2	*	0.3	0.6	*
2002	0.1	0.3	*	0.1	0.2	*	0.3	0.6	*
2001	0.1	0.3	*	0.1	0.2	*	0.2	0.5	*
2000	0.1	0.2	*	0.1	0.2	*	0.3	0.6	*
1999	0.1	0.3	*	0.1	0.2	*	0.3	0.7	*

Worktable 13. Age-adjusted death rates for 358 selected causes, by race and sex: United States, 1999-2005

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992) and year	All races 1/			White 2/			Black 2/		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
Legal execution (Y35.5)									
2005	0.0	0.0	*	0.0	0.0	*	*	*	*
2004	0.0	0.0	*	0.0	0.0	*	*	*	*
2003	0.0	0.1	*	0.0	0.0	*	*	*	*
2002	0.0	0.1	*	0.0	0.0	*	*	*	*
2001	0.0	0.1	*	0.0	0.0	*	*	*	*
2000	0.0	0.0	*	0.0	0.0	*	0.1	0.2	*
1999	0.0	0.1	*	0.0	0.1	*	0.1	0.2	*
Other legal intervention and their sequelae (Y35.0-Y35.4,Y35.6-Y35.7,Y89.0)									
2005	0.1	0.2	*	0.1	0.2	*	0.2	0.5	*
2004	0.1	0.2	*	0.1	0.2	*	0.2	0.4	*
2003	0.1	0.2	*	0.1	0.2	*	0.3	0.5	*
2002	0.1	0.2	*	0.1	0.2	*	0.2	0.5	*
2001	0.1	0.2	*	0.1	0.2	*	0.2	0.4	*
2000	0.1	0.2	*	0.1	0.2	*	0.2	0.4	*
1999	0.1	0.2	*	0.1	0.2	*	0.2	0.5	*
Operations of war and their sequelae (Y36,Y89.1)									
2005	0.0	0.0	*	0.0	0.0	*	*	*	*
2004	0.0	0.0	*	0.0	0.0	*	*	*	*
2003	*	*	*	*	*	*	*	*	*
2002	0.0	0.0	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*
1999	0.0	0.0	*	0.0	0.0	*	*	*	*

Worktable 13. Age-adjusted death rates for 358 selected causes, by race and sex: United States, 1999-2005

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992) and year	All races 1/			White 2/			Black 2/		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
Complications of medical and surgical care (Y40-Y84,Y88)									
2005	0.9	0.9	0.8	0.8	0.9	0.8	1.5	1.5	1.5
2004	1.0	1.1	0.9	0.9	1.1	0.8	1.5	1.5	1.6
2003	1.0	1.1	0.9	0.9	1.0	0.8	1.5	1.5	1.6
2002	1.0	1.0	1.0	0.9	1.0	0.9	1.6	1.6	1.6
2001	1.1	1.1	1.0	1.0	1.1	0.9	2.0	2.3	1.7
2000	1.1	1.2	1.0	1.0	1.2	0.9	1.9	1.9	1.9
1999	1.0	1.1	0.9	1.0	1.1	0.9	1.8	2.0	1.7
Drugs, medicaments and biological substances causing adverse effects in therapeutic use, and their sequelae (Y40-Y59,Y88.0)									
2005	0.1	0.1	0.1	0.1	0.1	0.1	0.2	0.2	0.2
2004	0.1	0.1	0.1	0.1	0.1	0.1	0.2	0.2	0.2
2003	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.2	*
2002	0.1	0.1	0.1	0.1	0.1	0.1	0.1	*	0.1
2001	0.1	0.1	0.1	0.1	0.1	0.1	0.2	0.2	0.2
2000	0.1	0.1	0.1	0.1	0.1	0.1	0.1	*	0.2
1999	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	*
Misadventures to patients during medical and surgical care and their sequelae (Y60-Y69,Y88.1)									
2005	0.0	0.0	0.0	0.0	0.0	0.0	0.1	*	0.1
2004	0.1	0.0	0.1	0.1	0.0	0.1	0.1	*	*
2003	0.1	0.0	0.1	0.0	0.0	0.1	0.1	*	0.1
2002	0.0	0.0	0.1	0.0	0.0	0.0	0.1	*	0.1
2001	0.1	0.1	0.1	0.1	0.1	0.1	0.1	*	*
2000	0.1	0.1	0.1	0.1	0.1	0.1	0.1	*	0.2
1999	0.1	0.1	0.1	0.0	0.1	0.1	0.1	*	0.2

Worktable 13. Age-adjusted death rates for 358 selected causes, by race and sex: United States, 1999-2005

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992) and year	All races 1/			White 2/			Black 2/		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
Other complications of medical and surgical care and their sequelae (Y70-Y84,Y88.2-Y88.3)									
2005	0.7	0.8	0.7	0.7	0.7	0.6	1.2	1.3	1.2
2004	0.8	0.9	0.7	0.8	0.9	0.6	1.3	1.3	1.3
2003	0.8	0.9	0.8	0.8	0.9	0.7	1.3	1.2	1.3
2002	0.8	0.9	0.8	0.8	0.9	0.8	1.4	1.4	1.3
2001	0.9	1.0	0.9	0.8	0.9	0.8	1.6	2.0	1.4
2000	0.9	1.1	0.9	0.9	1.0	0.8	1.6	1.7	1.6
1999	0.9	1.0	0.8	0.8	0.9	0.8	1.5	1.7	1.4

* Figure does not meet standards of reliability or precision.
0.0 Quantity more than zero but less than 0.05.
... Category not applicable.
1/ Includes races other than white and black.
2/ Multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming in 2005, by California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and by California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005."
3/ Figures include September 11, 2001 related deaths for which death certificates were filed as of 10/24/02; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001."

Source: CDC/NCHS, National Vital Statistics System, Mortality.