Person with disabilities experiencing problems accessing NCHS birth and death tables should contact **nchsed@cdc.gov**, or call 301-458-4688

Person with disabilities experiencing problems accessing NCHS birth and death tables should contact **nchsed@cdc.gov**, or call 301-458-4688

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
All causes													
All origins, both sexes													
2005	2,448,017	28,440	4,756	6,602	34,234	41,925	84,785	183,530	275,301	398,355	686,665	703,169	255
2004	2,397,615	27,936	4,785	6,834	33,421	40,868	85,362	177,697	264,697	399,666	684,230	671,773	346
2003	2,448,288	28,025	4,965	6,954	33,568	41,300	89,461	176,781	262,519	413,497	703,024	687,852	342
2002	2,443,387	28,034	4,858	7,150	33,046	41,355	91,140	172,385	253,342	422,990	707,654	681,076	357
2001	2,416,425	27,568	5,107	7,095	32,252	41,683	91,674	168,065	244,139	430,960	701,929	665,531	422
2000	2,403,351	28,035	4,979	7,413	31,307	40,451	89,798	160,341	240,846	441,209	700,445	658,171	356
1999	2,391,399	27,937	5,249	7,595	30,656	41,066	89,256	152,974	238,979	452,600	698,590	646,141	356
All origins, male													
2005	1,207,675	16,018	2,765	3,853	25,509	29,283	53,309	114,472	165,429	222,807	335,235	238,796	199
2004	1,181,668	15,718	2,649	3,999	24,587	28,359	53,677	111,163	158,032	222,891	333,669	226,650	274
2003	1,201,964	15,902	2,826	4,150	24,670	28,602	56,435	110,682	156,461	231,421	342,332	228,212	271
2002	1,199,264	15,717	2,806	4,198	24,416	28,736	57,593	107,722	151,363	237,021	343,504	225,906	282
2001	1,183,421	15,477	2,899	4,168	23,963	28,757	58,164	104,848	144,958	241,581	340,742	217,533	331
2000	1,177,578	15,718	2,824	4,401	23,071	27,890	57,297	100,398	143,321	247,408	340,219	214,742	289
1999	1,175,460	15,646	2,975	4,492	22,414	28,276	57,118	95,659	142,724	254,920	340,970	209,989	277
All origins, female													
2005	1,240,342	12,422	1,991	2,749	8,725	12,642	31,476	69,058	109,872	175,548	351,430	464,373	56
2004	1,215,947	12,218	2,136	2,835	8,834	12,509	31,685	66,534	106,665	176,775	350,561	445,123	72
2003	1,246,324	12,123	2,139	2,804	8,898	12,698	33,026	66,099	106,058	182,076	360,692	459,640	71
2002	1,244,123	12,317	2,052	2,952	8,630	12,619	33,547	64,663	101,979	185,969	364,150	455,170	75
2001	1,233,004	12,091	2,208	2,927	8,289	12,926	33,510	63,217	99,181	189,379	361,187	447,998	91
2000	1,225,773	12,317	2,155	3,012	8,236	12,561	32,501	59,943	97,525	193,801	360,226	443,429	67
1999	1,215,939	12,291	2,274	3,103	8,242	12,790	32,138	57,315	96,255	197,680	357,620	436,152	79

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
All causes													
Hispanic, both sexes													
2005	131,161	5,724	1,041	1,068	5,837	6,411	8,876	13,569	16,089	21,422	29,010	22,058	56
2004	122,416	5,321	947	1,105	5,428	5,959	8,546	12,628	15,198	20,420	26,879	19,937	48
2003	122,026	5,281	993	1,173	5,394	6,189	8,576	12,516	14,913	20,596	26,219	20,129	47
2002	117,135	4,943	933	1,124	5,255	5,816	8,467	11,745	14,215	20,476	24,838	19,265	58
2001	113,413	4,680	917	1,038	5,058	5,797	8,441	11,362	13,337	20,172	23,637	18,907	67
2000	107,254	4,598	872	1,065	4,793	5,509	8,085	10,356	12,733	19,420	21,951	17,818	54
1999	103,740	4,412	882	1,014	4,505	5,397	7,798	9,876	12,319	19,201	20,912	17,358	66
Hispanic, male													
2005	73,788	3,192	610	612	4,604	4,960	6,145	8,994	9,954	11,984	14,539	8,143	51
2004	68,544	2,948	535	656	4,376	4,547	5,973	8,451	9,246	11,279	13,397	7,090	46
2003	68,119	2,940	569	706	4,244	4,742	5,887	8,336	9,122	11,465	12,904	7,165	39
2002	65,703	2,746	546	647	4,182	4,474	5,828	7,719	8,762	11,476	12,291	6,977	55
2001	63,317	2,601	518	600	4,019	4,366	5,895	7,477	8,026	11,231	11,759	6,767	58
2000	60,172	2,514	475	622	3,837	4,200	5,597	6,808	7,760	10,842	10,973	6,493	51
1999	57,991	2,411	481	592	3,547	4,084	5,476	6,497	7,490	10,783	10,317	6,253	60
Hispanic, female													
2005	57,373	2,532	431	456	1,233	1,451	2,731	4,575	6,135	9,438	14,471	13,915	5
2004	53,872	2,373	412	449	1,052	1,412	2,573	4,177	5,952	9,141	13,482	12,847	2
2003	53,907	2,341	424	467	1,150	1,447	2,689	4,180	5,791	9,131	13,315	12,964	8
2002	51,432	2,197	387	477	1,073	1,342	2,639	4,026	5,453	9,000	12,547	12,288	3
2001	50,096	2,079	399	438	1,039	1,431	2,546	3,885	5,311	8,941	11,878	12,140	9
2000	47,082	2,084	397	443	956	1,309	2,488	3,548	4,973	8,578	10,978	11,325	3
1999	45,749	2,001	401	422	958	1,313	2,322	3,379	4,829	8,418	10,595	11,105	6

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
All causes													
Non-Hispanic 1/, both sexes													
2005	2,312,028	22,488	3,701	5,515	28,309	35,380	75,626	169,355	258,493	376,169	656,613	680,266	113
2004	2,269,583	22,370	3,826	5,699	27,896	34,774	76,524	164,412	248,724	378,266	656,132	650,809	151
2003	2,319,476	22,475	3,956	5,761	28,044	34,955	80,503	163,533	246,665	391,778	675,309	666,350	147
2002	2,318,269	22,816	3,904	6,002	27,658	35,329	82,225	159,843	238,185	401,182	680,878	660,078	169
2001	2,295,244	22,629	4,171	6,033	27,059	35,656	82,763	155,916	229,812	409,500	676,512	645,020	173
2000	2,287,846	23,074	4,086	6,316	26,374	34,741	81,182	149,141	227,118	420,369	676,632	638,673	140
1999	2,279,325	23,126	4,343	6,560	26,043	35,471	80,949	142,302	225,659	431,952	675,730	627,035	155
Non-Hispanic 1/, male													
2005	1,131,013	12,701	2,147	3,231	20,835	24,225	46,957	105,044	154,967	210,333	320,125	230,371	77
2004	1,109,848	12,633	2,106	3,327	20,145	23,710	47,501	102,236	148,234	210,979	319,650	219,215	112
2003	1,129,927	12,799	2,247	3,432	20,328	23,740	50,284	101,828	146,696	219,240	328,640	220,575	118
2002	1,129,090	12,815	2,250	3,534	20,129	24,114	51,451	99,433	141,966	224,706	330,191	218,367	134
2001	1,115,683	12,739	2,371	3,554	19,836	24,225	51,918	96,815	136,214	229,539	328,079	210,256	137
2000	1,112,704	13,007	2,338	3,770	19,136	23,548	51,304	92,984	134,890	235,648	328,284	207,688	107
1999	1,112,718	12,998	2,479	3,885	18,797	24,047	51,272	88,567	134,540	243,210	329,682	203,128	113
Non-Hispanic 1/, female													
2005	1,181,015	9,787	1,554	2,284	7,474	11,155	28,669	64,311	103,526	165,836	336,488	449,895	36
2004	1,159,735	9,737	1,720	2,372	7,751	11,064	29,023	62,176	100,490	167,287	336,482	431,594	39
2003	1,189,549	9,676	1,709	2,329	7,716	11,215	30,219	61,705	99,969	172,538	346,669	445,775	29
2002	1,189,179	10,001	1,654	2,468	7,529	11,215	30,774	60,410	96,219	176,476	350,687	441,711	35
2001	1,179,561	9,890	1,800	2,479	7,223	11,431	30,845	59,101	93,598	179,961	348,433	434,764	36
2000	1,175,142	10,067	1,748	2,546	7,238	11,193	29,878	56,157	92,228	184,721	348,348	430,985	33
1999	1,166,607	10,128	1,864	2,675	7,246	11,424	29,677	53,735	91,119	188,742	346,048	423,907	42

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
All causes													
Non-Hispanic white, both sexes													
2005 2/	1,967,142	13,026	2,404	3,681	19,683	23,826	54,786	126,669	206,078	314,958	581,712	620,246	73
2004 2/	1,933,382	13,046	2,458	3,820	19,684	23,397	55,543	123,037	198,842	317,606	582,454	593,382	113
2003 2/	1,979,465	13,223	2,539	3,884	19,640	23,557	58,095	121,832	197,598	329,570	600,563	608,843	121
2002	1,981,973	13,463	2,502	4,031	19,458	23,758	59,529	119,812	190,843	338,961	606,431	603,063	122
2001	1,962,810	13,311	2,772	4,151	18,730	24,191	59,546	117,086	184,109	346,389	603,143	589,269	113
2000	1,959,919	13,513	2,641	4,397	18,261	23,191	58,231	111,964	182,348	356,869	604,658	583,757	89
1999	1,953,197	13,553	2,819	4,488	17,868	23,986	57,630	106,808	180,793	367,490	604,344	573,327	91
Non-Hispanic white, male													
2005 2/	954,402	7,418	1,408	2,163	14,267	16,366	34,719	79,860	124,229	177,517	285,608	210,795	52
2004 2/	938,143	7,399	1,365	2,264	14,017	16,098	35,253	77,741	119,171	178,642	285,752	200,357	84
2003 2/	956,194	7,591	1,423	2,319	14,022	16,097	37,123	77,106	118,383	186,112	293,774	202,148	96
2002	957,645	7,706	1,431	2,365	13,957	16,339	38,049	75,724	114,535	191,844	295,763	199,834	98
2001	945,967	7,508	1,592	2,428	13,542	16,613	38,213	73,731	109,993	196,158	294,013	192,084	92
2000	944,781	7,660	1,541	2,652	13,066	15,934	37,622	70,788	109,031	202,038	294,842	189,537	70
1999	944,913	7,718	1,605	2,643	12,677	16,449	37,291	67,151	108,692	209,155	296,528	184,938	66
Non-Hispanic white, female													
2005 2/	1,012,740	5,608	996	1,518	5,416	7,460	20,067	46,809	81,849	137,441	296,104	409,451	21
2004 2/	995,239	5,647	1,093	1,556	5,667	7,299	20,290	45,296	79,671	138,964	296,702	393,025	29
2003 2/	1,023,271	5,632	1,116	1,565	5,618	7,460	20,972	44,726	79,215	143,458	306,789	406,695	25
2002	1,024,328	5,757	1,071	1,666	5,501	7,419	21,480	44,088	76,308	147,117	310,668	403,229	24
2001	1,016,843	5,803	1,180	1,723	5,188	7,578	21,333	43,355	74,116	150,231	309,130	397,185	21
2000	1,015,138	5,853	1,100	1,745	5,195	7,257	20,609	41,176	73,317	154,831	309,816	394,220	19
1999	1,008,284	5,835	1,214	1,845	5,191	7,537	20,339	39,657	72,101	158,335	307,816	388,389	25

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
All causes													
Non-Hispanic black, both sexes													
2005 2/	289,163	8,335	1,065	1,500	7,244	9,850	17,930	37,298	44,937	51,166	61,319	48,481	38
2004 2/	283,859	8,221	1,128	1,551	6,924	9,768	18,237	36,247	42,998	51,126	60,748	46,877	34
2003 2/	287,968	8,159	1,169	1,530	7,021	9,795	19,561	36,584	42,350	52,583	61,770	47,426	20
2002	286,573	8,284	1,163	1,634	6,904	10,009	19,955	35,211	40,878	52,748	62,183	47,565	39
2001	284,343	8,277	1,156	1,557	7,024	9,859	20,436	34,098	39,523	53,803	61,592	46,968	50
2000	282,676	8,525	1,217	1,620	6,926	10,103	20,279	32,911	38,867	54,645	60,874	46,660	49
1999	281,979	8,589	1,283	1,771	6,980	9,992	20,770	31,377	39,063	55,567	60,634	45,895	58
Non-Hispanic black, male													
2005 2/	147,010	4,665	606	885	5,615	6,706	10,451	21,946	26,396	27,303	27,723	14,690	24
2004 2/	144,022	4,613	623	870	5,239	6,564	10,552	21,449	25,050	27,228	27,394	14,415	25
2003 2/	146,136	4,601	681	911	5,343	6,592	11,405	21,708	24,468	27,957	28,297	14,156	17
2002	144,802	4,520	681	977	5,248	6,723	11,661	20,839	23,706	27,859	28,137	14,422	29
2001	143,971	4,651	655	941	5,377	6,548	11,971	20,307	22,600	28,446	28,083	14,352	40
2000	143,297	4,757	672	959	5,224	6,649	12,061	19,647	22,402	28,745	27,715	14,431	35
1999	143,883	4,765	734	1,081	5,286	6,633	12,384	19,026	22,493	29,259	27,552	14,628	42
Non-Hispanic black, female													
2005 2/	142,153	3,670	459	615	1,629	3,144	7,479	15,352	18,541	23,863	33,596	33,791	14
2004 2/	139,837	3,608	505	681	1,685	3,204	7,685	14,798	17,948	23,898	33,354	32,462	9
2003 2/	141,832	3,558	488	619	1,678	3,203	8,156	14,876	17,882	24,626	33,473	33,270	3
2002	141,771	3,764	482	657	1,656	3,286	8,294	14,372	17,172	24,889	34,046	33,143	10
2001	140,372	3,626	501	616	1,647	3,311	8,465	13,791	16,923	25,357	33,509	32,616	10
2000	139,379	3,768	545	661	1,702	3,454	8,218	13,264	16,465	25,900	33,159	32,229	14
1999	138,096	3,824	549	690	1,694	3,359	8,386	12,351	16,570	26,308	33,082	31,267	16

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
All causes													
Origin not stated 3/, both sexes													
2005	4,828	228	14	19	88	134	283	606	719	764	1,042	845	86
2004	5,616	245	12	30	97	135	292	657	775	980	1,219	1,027	147
2003	6,786	269	16	20	130	156	382	732	941	1,123	1,496	1,373	148
2002	7,983	275	21	24	133	210	448	797	942	1,332	1,938	1,733	130
2001	7,768	259	19	24	135	230	470	787	990	1,288	1,780	1,604	182
2000	8,251	363	21	32	140	201	531	844	995	1,420	1,862	1,680	162
1999	8,334	399	24	21	108	198	509	796	1,001	1,447	1,948	1,748	135
Origin not stated 3/, male													
2005	2,874	125	8	10	70	98	207	434	508	490	571	282	71
2004	3,276	137	8	16	66	102	203	476	552	633	622	345	116
2003	3,918	163	10	12	98	120	264	518	643	716	788	472	114
2002	4,471	156	10	17	105	148	314	570	635	839	1,022	562	93
2001	4,421	137	10	14	108	166	351	556	718	811	904	510	136
2000	4,702	197	11	9	98	142	396	606	671	918	962	561	131
1999	4,751	237	15	15	70	145	370	595	694	927	971	608	104
Origin not stated 3/, female													
2005	1,954	103	6	9	18	36	76	172	211	274	471	563	15
2004	2,340	108	4	14	31	33	89	181	223	347	597	682	31
2003	2,868	106	6	8	32	36	118	214	298	407	708	901	34
2002	3,512	119	11	7	28	62	134	227	307	493	916	1,171	37
2001	3,347	122	9	10	27	64	119	231	272	477	876	1,094	46
2000	3,549	166	10	23	42	59	135	238	324	502	900	1,119	31
1999	3,583	162	9	6	38	53	139	201	307	520	977	1,140	31

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Salmonella infections (A01-A02)													
All origins, both sexes													
2005	30	-	-	-	1	2	2	2	6	3	9	5	-
2004	30	1	-	-	-	1	1	3	4	7	7	6	-
2003	43	3	-	1	-	1	3	2	6	8	10	9	-
2002	21	3	-	-	-	-	2	3	-	4	8	1	-
2001	41	2	-	-	-	2	1	3	3	7	14	9	-
2000	28	3	2	-	1	1	-	4	2	7	5	3	-
1999	38	3	-	-	2	-	2	2	3	7	12	7	-
All origins, male													
2005	11	-	-	-	-	-	1	1	5	1	2	1	-
2004	22	1	-	-	-	-	1	3	4	3	6	4	-
2003	23	1	-	-	-	-	2	2	4	4	4	6	-
2002	10	1	-	-	-	-	-	1	-	2	5	1	-
2001	21	-	-	-	-	1	1	3	2	6	7	1	-
2000	15	2	-	-	1	1	-	2	2	5	1	1	-
1999	21	2	-	-	2	-	1	2	2	5	5	2	-
All origins, female													
2005	19	-	-	-	1	2	1	1	1	2	7	4	-
2004	8	-	-	-	-	1	-	-	-	4	1	2	-
2003	20	2	-	1	-	1	1	-	2	4	6	3	-
2002	11	2	-	-	-	-	2	2	-	2	3	-	-
2001	20	2	-	-	-	1	-	-	1	1	7	8	-
2000	13	1	2	-	-	-	-	2	-	2	4	2	-
1999	17	1	-	-	-	-	1	-	1	2	7	5	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Salmonella infections (A01-A02)													
Hispanic, both sexes													
2005	4	-	-	-	-	-	1	1	1	-	1	-	-
2004	3	1	-	-	-	-	-	1	-	-	1	-	-
2003	6	-	-	1	-	-	1	1	-	2	1	-	-
2002	3	-	-	-	-	-	-	1	-	2	-	-	-
2001	5	-	-	-	-	-	-	1	1	1	2	-	-
2000	2	2	-	-	-	-	-	-	-	-	-	-	-
1999	4	-	-	-	-	-	-	-	1	1	2	-	-
Hispanic, male													
2005	3	-	-	-	-	-	1	1	1	-	-	-	-
2004	3	1	-	-	-	-	-	1	-	-	1	-	-
2003	3	-	-	-	-	-	1	1	-	1	-	-	-
2002	1	-	-	-	-	-	-	-	-	1	-	-	-
2001	5	-	-	-	-	-	-	1	1	1	2	-	-
2000	1	1	-	-	-	-	-	-	-	-	-	-	-
1999	3	-	-	-	-	-	-	-	1	1	1	-	-
Hispanic, female													
2005	1	-	-	-	-	-	-	-	-	-	1	-	-
2004	-	-	-	-	-	-	-	-	-	-	-	-	-
2003	3	-	-	1	-	-	-	-	-	1	1	-	-
2002	2	-	-	-	-	-	-	1	-	1	-	-	-
2001	-	-	-	-	-	-	-	-	-	-	-	-	-
2000	1	1	-	-	-	-	-	-	-	-	-	-	-
1999	1	-	-	-	-	-	-	-	-	-	1	-	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Salmonella infections (A01-A02)													
Non-Hispanic 1/, both sexes													
2005	26	-	-	-	1	2	1	1	5	3	8	5	-
2004	27	-	-	-	-	1	1	2	4	7	6	6	-
2003	37	3	-	-	-	1	2	1	6	6	9	9	-
2002	18	3	-	-	-	-	2	2	-	2	8	1	-
2001	36	2	-	-	-	2	1	2	2	6	12	9	-
2000	26	1	2	-	1	1	-	4	2	7	5	3	-
1999	34	3	-	-	2	-	2	2	2	6	10	7	-
Non-Hispanic 1/, male													
2005	8	-	-	-	-	-	-	-	4	1	2	1	-
2004	19	-	-	-	-	-	1	2	4	3	5	4	-
2003	20	1	-	-	-	-	1	1	4	3	4	6	-
2002	9	1	-	-	-	-	-	1	-	1	5	1	-
2001	16	-	-	-	-	1	1	2	1	5	5	1	-
2000	14	1	-	-	1	1	-	2	2	5	1	1	-
1999	18	2	-	-	2	-	1	2	1	4	4	2	-
Non-Hispanic 1/, female													
2005	18	-	-	-	1	2	1	1	1	2	6	4	-
2004	8	-	-	-	-	1	-	-	-	4	1	2	-
2003	17	2	-	-	-	1	1	-	2	3	5	3	-
2002	9	2	-	-	-	-	2	1	-	1	3	-	-
2001	20	2	-	-	-	1	-	-	1	1	7	8	-
2000	12	-	2	-	-	-	-	2	-	2	4	2	-
1999	16	1	-	-	-	-	1	-	1	2	6	5	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Salmonella infections (A01-A02)													
Non-Hispanic white, both sexes													
2005 2/	20	-	-	-	-	1	1	-	4	3	7	4	-
2004 2/	17	-	-	-	-	1	-	1	2	3	5	5	-
2003 2/	25	1	-	-	-	-	-	1	5	4	6	8	-
2002	15	2	-	-	-	-	2	2	-	1	7	1	-
2001	28	1	-	-	-	2	-	2	2	3	10	8	-
2000	23	-	2	-	-	1	-	4	2	6	5	3	-
1999	28	3	-	-	1	-	2	2	1	4	8	7	-
Non-Hispanic white, male													
2005 2/	6	-	-	-	-	-	-	-	3	1	1	1	-
2004 2/	11	-	-	-	-	-	-	1	2	1	4	3	-
2003 2/	14	-	-	-	-	-	-	1	3	3	2	5	-
2002	8	1	-	-	-	-	-	1	-	-	5	1	-
2001	13	-	-	-	-	1	-	2	1	3	5	1	-
2000	12	-	-	-	-	1	-	2	2	5	1	1	-
1999	13	2	-	-	1	-	1	2	-	2	3	2	-
Non-Hispanic white, female													
2005 2/	14	-	-	-	-	1	1	-	1	2	6	3	-
2004 2/	6	-	-	-	-	1	-	-	-	2	1	2	-
2003 2/	11	1	-	-	-	-	-	-	2	1	4	3	-
2002	7	1	-	-	-	-	2	1	-	1	2	-	-
2001	15	1	-	-	-	1	-	-	1	-	5	7	-
2000	11	-	2	-	-	-	-	2	-	1	4	2	-
1999	15	1	-	-	-	-	1	-	1	2	5	5	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Salmonella infections (A01-A02)													
Non-Hispanic black, both sexes													
2005 2/	5	-	-	-	-	1	-	1	1	-	1	1	-
2004 2/	4	-	-	-	-	-	-	1	1	2	-	-	-
2003 2/	8	2	-	-	-	1	1	-	-	1	3	-	-
2002	3	1	-	-	-	-	-	-	-	1	1	-	-
2001	5	1	-	-	-	-	-	-	-	1	2	1	-
2000	2	1	-	-	1	-	-	-	-	-	-	-	-
1999	4	-	-	-	1	-	-	-	-	1	2	-	-
Non-Hispanic black, male													
2005 2/	2	-	-	-	-	-	-	-	1	-	1	-	-
2004 2/	3	-	-	-	-	-	-	1	1	1	-	-	-
2003 2/	3	1	-	-	-	-	-	-	-	-	2	-	-
2002	1	-	-	-	-	-	-	-	-	1	-	-	-
2001	1	-	-	-	-	-	-	-	-	1	-	-	-
2000	2	1	-	-	1	-	-	-	-	-	-	-	-
1999	3	-	-	-	1	-	-	-	-	1	1	-	-
Non-Hispanic black, female													
2005 2/	3	-	-	-	-	1	-	1	-	-	-	1	-
2004 2/	1	-	-	-	-	-	-	-	-	1	-	-	-
2003 2/	5	1	-	-	-	1	1	-	-	1	1	-	-
2002	2	1	-	-	-	-	-	-	-	-	1	-	-
2001	4	1	-	-	-	-	-	-	-	-	2	1	-
2000	-	-	-	-	-	-	-	-	-	-	-	-	-
1999	1	-	-	-	-	-	-	-	-	-	1	-	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Salmonella infections (A01-A02)													
Origin not stated 3/, both sexes													
2005	-	-	-	-	-	-	-	-	-	-	-	-	-
2004	-	-	-	-	-	-	-	-	-	-	-	-	-
2003	-	-	-	-	-	-	-	-	-	-	-	-	-
2002	-	-	-	-	-	-	-	-	-	-	-	-	-
2001	-	-	-	-	-	-	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-	-	-	-
Origin not stated 3/, male													
2005	-	-	-	-	-	-	-	-	-	-	-	-	-
2004	-	-	-	-	-	-	-	-	-	-	-	-	-
2003	-	-	-	-	-	-	-	-	-	-	-	-	-
2002	-	-	-	-	-	-	-	-	-	-	-	-	-
2001	-	-	-	-	-	-	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-	-	-	-
Origin not stated 3/, female													
2005	-	-	-	-	-	-	-	-	-	-	-	-	-
2004	-	-	-	-	-	-	-	-	-	-	-	-	-
2003	-	-	-	-	-	-	-	-	-	-	-	-	-
2002	-	-	-	-	-	-	-	-	-	-	-	-	-
2001	-	-	-	-	-	-	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-	-	-	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Shigellosis and amebiasis (A03,A06)													
All origins, both sexes													
2005	10	-	1	2	-	-	-	1	2	2	2	-	-
2004	3	-	-	1	1	-	-	-	-	1	-	-	-
2003	7	-	-	-	-	-	3	-	2	1	1	-	-
2002	8	-	1	-	1	-	1	2	1	-	1	1	-
2001	7	1	1	1	-	-	-	3	-	-	1	-	-
2000	12	-	-	-	-	-	1	1	1	1	7	1	-
1999	10	-	1	1	-	-	1	-	1	1	1	4	-
All origins, male													
2005	3	-	-	1	-	-	-	1	1	-	-	-	-
2004	2	-	-	1	-	-	-	-	-	1	-	-	-
2003	5	-	-	-	-	-	2	-	1	1	1	-	-
2002	4	-	-	-	1	-	1	2	-	-	-	-	-
2001	6	1	1	1	-	-	-	3	-	-	-	-	-
2000	5	-	-	-	-	-	1	-	1	-	3	-	-
1999	6	-	1	1	-	-	1	-	1	1	-	1	-
All origins, female													
2005	7	-	1	1	-	-	-	-	1	2	2	-	-
2004	1	-	-	-	1	-	-	-	-	-	-	-	-
2003	2	-	-	-	-	-	1	-	1	-	-	-	-
2002	4	-	1	-	-	-	-	-	1	-	1	1	-
2001	1	-	-	-	-	-	-	-	-	-	1	-	-
2000	7	-	-	-	-	-	-	1	-	1	4	1	-
1999	4	-	-	-	-	-	-	-	-	-	1	3	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Shigellosis and amebiasis (A03,A06)													
Hispanic, both sexes													
2005	4	-	1	1	-	-	-	1	-	-	1	-	-
2004	2	-	-	1	1	-	-	-	-	-	-	-	-
2003	4	-	-	-	-	-	2	-	1	1	-	-	-
2002	1	-	-	-	-	-	-	1	-	-	-	-	-
2001	3	-	1	1	-	-	-	1	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-	-	-	-
1999	4	-	1	1	-	-	1	-	-	-	1	-	-
Hispanic, male													
2005	1	-	-	-	-	-	-	1	-	-	-	-	-
2004	1	-	-	1	-	-	-	-	-	-	-	-	-
2003	2	-	-	-	-	-	1	-	-	1	-	-	-
2002	1	-	-	-	-	-	-	1	-	-	-	-	-
2001	3	-	1	1	-	-	-	1	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-	-	-	-
1999	3	-	1	1	-	-	1	-	-	-	-	-	-
Hispanic, female													
2005	3	-	1	1	-	-	-	-	-	-	1	-	-
2004	1	-	-	-	1	-	-	-	-	-	-	-	-
2003	2	-	-	-	-	-	1	-	1	-	-	-	-
2002	-	-	-	-	-	-	-	-	-	-	-	-	-
2001	-	-	-	-	-	-	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-	-	-	-
1999	1	-	-	-	-	-	-	-	-	-	1	-	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Shigellosis and amebiasis (A03,A06)													
Non-Hispanic 1/, both sexes													
2005	6	-	-	1	-	-	-	-	2	2	1	-	-
2004	1	-	-	-	-	-	-	-	-	1	-	-	-
2003	3	-	-	-	-	-	1	-	1	-	1	-	-
2002	7	-	1	-	1	-	1	1	1	-	1	1	-
2001	4	1	-	-	-	-	-	2	-	-	1	-	-
2000	12	-	-	-	-	-	1	1	1	1	7	1	-
1999	6	-	-	-	-	-	-	-	1	1	-	4	-
Non-Hispanic 1/, male													
2005	2	-	-	1	-	-	-	-	1	-	-	-	-
2004	1	-	-	-	-	-	-	-	-	1	-	-	-
2003	3	-	-	-	-	-	1	-	1	-	1	-	-
2002	3	-	-	-	1	-	1	1	-	-	-	-	-
2001	3	1	-	-	-	-	-	2	-	-	-	-	-
2000	5	-	-	-	-	-	1	-	1	-	3	-	-
1999	3	-	-	-	-	-	-	-	1	1	-	1	-
Non-Hispanic 1/, female													
2005	4	-	-	-	-	-	-	-	1	2	1	-	-
2004	-	-	-	-	-	-	-	-	-	-	-	-	-
2003	-	-	-	-	-	-	-	-	-	-	-	-	-
2002	4	-	1	-	-	-	-	-	1	-	1	1	-
2001	1	-	-	-	-	-	-	-	-	-	1	-	-
2000	7	-	-	-	-	-	-	1	-	1	4	1	-
1999	3	-	-	-	-	-	-	-	-	-	-	3	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Shigellosis and amebiasis (A03,A06)													
Non-Hispanic white, both sexes													
2005 2/	6	-	-	1	-	-	-	-	2	2	1	-	-
2004 2/	1	-	-	-	-	-	-	-	-	1	-	-	-
2003 2/	2	-	-	-	-	-	-	-	1	-	1	-	-
2002	6	-	-	-	1	-	1	1	1	-	1	1	-
2001	4	1	-	-	-	-	-	2	-	-	1	-	-
2000	10	-	-	-	-	-	-	1	1	-	7	1	-
1999	6	-	-	-	-	-	-	-	1	1	-	4	-
Non-Hispanic white, male													
2005 2/	2	-	-	1	-	-	-	-	1	-	-	-	-
2004 2/	1	-	-	-	-	-	-	-	-	1	-	-	-
2003 2/	2	-	-	-	-	-	-	-	1	-	1	-	-
2002	3	-	-	-	1	-	1	1	-	-	-	-	-
2001	3	1	-	-	-	-	-	2	-	-	-	-	-
2000	4	-	-	-	-	-	-	-	1	-	3	-	-
1999	3	-	-	-	-	-	-	-	1	1	-	1	-
Non-Hispanic white, female													
2005 2/	4	-	-	-	-	-	-	-	1	2	1	-	-
2004 2/	-	-	-	-	-	-	-	-	-	-	-	-	-
2003 2/	-	-	-	-	-	-	-	-	-	-	-	-	-
2002	3	-	-	-	-	-	-	-	1	-	1	1	-
2001	1	-	-	-	-	-	-	-	-	-	1	-	-
2000	6	-	-	-	-	-	-	1	-	-	4	1	-
1999	3	-	-	-	-	-	-	-	-	-	-	3	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Shigellosis and amebiasis (A03,A06)													
Non-Hispanic black, both sexes													
2005 2/	-	-	-	-	-	-	-	-	-	-	-	-	-
2004 2/	-	-	-	-	-	-	-	-	-	-	-	-	-
2003 2/	1	-	-	-	-	-	1	-	-	-	-	-	-
2002	-	-	-	-	-	-	-	-	-	-	-	-	-
2001	-	-	-	-	-	-	-	-	-	-	-	-	-
2000	1	-	-	-	-	-	1	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-	-	-	-
Non-Hispanic black, male													
2005 2/	-	-	-	-	-	-	-	-	-	-	-	-	-
2004 2/	-	-	-	-	-	-	-	-	-	-	-	-	-
2003 2/	1	-	-	-	-	-	1	-	-	-	-	-	-
2002	-	-	-	-	-	-	-	-	-	-	-	-	-
2001	-	-	-	-	-	-	-	-	-	-	-	-	-
2000	1	-	-	-	-	-	1	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-	-	-	-
Non-Hispanic black, female													
2005 2/	-	-	-	-	-	-	-	-	-	-	-	-	-
2004 2/	-	-	-	-	-	-	-	-	-	-	-	-	-
2003 2/	-	-	-	-	-	-	-	-	-	-	-	-	-
2002	-	-	-	-	-	-	-	-	-	-	-	-	-
2001	-	-	-	-	-	-	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-	-	-	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Shigellosis and amebiasis (A03,A06)													
Origin not stated 3/, both sexes													
2005	-	-	-	-	-	-	-	-	-	-	-	-	-
2004	-	-	-	-	-	-	-	-	-	-	-	-	-
2003	-	-	-	-	-	-	-	-	-	-	-	-	-
2002	-	-	-	-	-	-	-	-	-	-	-	-	-
2001	-	-	-	-	-	-	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-	-	-	-
Origin not stated 3/, male													
2005	-	-	-	-	-	-	-	-	-	-	-	-	-
2004	-	-	-	-	-	-	-	-	-	-	-	-	-
2003	-	-	-	-	-	-	-	-	-	-	-	-	-
2002	-	-	-	-	-	-	-	-	-	-	-	-	-
2001	-	-	-	-	-	-	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-	-	-	-
Origin not stated 3/, female													
2005	-	-	-	-	-	-	-	-	-	-	-	-	-
2004	-	-	-	-	-	-	-	-	-	-	-	-	-
2003	-	-	-	-	-	-	-	-	-	-	-	-	-
2002	-	-	-	-	-	-	-	-	-	-	-	-	-
2001	-	-	-	-	-	-	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-	-	-	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Certain other intestinal infections (A04,A07-A09)													
All origins, both sexes													
2005	5,667	10	12	10	3	11	32	104	253	829	2,070	2,333	-
2004	4,357	13	15	1	2	12	31	94	252	608	1,662	1,667	-
2003	3,090	10	4	7	4	8	14	64	176	444	1,192	1,167	-
2002	2,465	13	8	3	6	4	18	56	142	362	889	964	-
2001	1,537	10	18	4	-	4	12	42	81	235	566	565	-
2000	1,328	13	10	-	1	3	19	29	70	222	462	499	-
1999	1,108	12	8	3	5	6	7	23	49	176	390	429	-
All origins, male													
2005	2,209	7	4	5	3	6	17	53	132	377	865	740	-
2004	1,622	8	5	-	1	5	19	48	125	264	665	482	-
2003	1,109	6	4	2	2	4	5	32	82	185	478	309	-
2002	947	3	5	1	3	1	8	29	71	185	358	283	-
2001	572	7	11	2	-	2	8	15	35	108	227	157	-
2000	524	10	6	-	-	3	10	16	39	105	195	140	-
1999	420	8	4	3	3	5	4	11	26	74	169	113	-
All origins, female													
2005	3,458	3	8	5	-	5	15	51	121	452	1,205	1,593	-
2004	2,735	5	10	1	1	7	12	46	127	344	997	1,185	-
2003	1,981	4	-	5	2	4	9	32	94	259	714	858	-
2002	1,518	10	3	2	3	3	10	27	71	177	531	681	-
2001	965	3	7	2	-	2	4	27	46	127	339	408	-
2000	804	3	4	-	1	-	9	13	31	117	267	359	-
1999	688	4	4	-	2	1	3	12	23	102	221	316	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Certain other intestinal infections (A04,A07-A09)													
Hispanic, both sexes													
2005	190	3	2	2	-	1	1	9	12	34	69	57	-
2004	149	2	3	-	1	1	1	9	17	26	50	39	-
2003	99	2	2	3	-	1	-	6	10	16	31	28	-
2002	78	5	5	-	1	-	1	3	6	10	31	16	-
2001	53	1	1	-	-	-	4	2	4	8	16	17	-
2000	46	3	2	-	-	-	2	-	8	9	11	11	-
1999	34	3	1	-	2	1	-	4	5	7	4	7	-
Hispanic, male													
2005	65	2	1	1	-	1	-	4	-	15	25	16	-
2004	57	1	1	-	-	1	-	6	5	12	16	15	-
2003	47	-	2	1	-	-	-	4	7	7	13	13	-
2002	31	1	3	-	-	-	-	2	4	5	12	4	-
2001	19	1	1	-	-	-	3	-	2	1	5	6	-
2000	23	2	1	-	-	-	1	-	5	6	4	4	-
1999	17	2	-	-	2	1	-	2	2	3	2	3	-
Hispanic, female													
2005	125	1	1	1	-	-	1	5	12	19	44	41	-
2004	92	1	2	-	1	-	1	3	12	14	34	24	-
2003	52	2	-	2	-	1	-	2	3	9	18	15	-
2002	47	4	2	-	1	-	1	1	2	5	19	12	-
2001	34	-	-	-	-	-	1	2	2	7	11	11	-
2000	23	1	1	-	-	-	1	-	3	3	7	7	-
1999	17	1	1	-	-	-	-	2	3	4	2	4	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Certain other intestinal infections (A04,A07-A09)													
Non-Hispanic 1/, both sexes													
2005	5,469	7	10	8	3	10	30	95	241	794	1,999	2,272	-
2004	4,200	9	12	1	1	11	30	85	234	582	1,610	1,625	-
2003	2,986	8	2	4	4	7	14	57	164	428	1,160	1,138	-
2002	2,381	8	3	3	5	4	17	52	135	351	857	946	-
2001	1,474	9	16	4	-	4	8	40	77	225	546	545	-
2000	1,277	10	8	-	1	3	17	29	62	211	449	487	-
1999	1,071	9	7	3	3	5	7	19	44	169	386	419	-
Non-Hispanic 1/, male													
2005	2,141	5	3	4	3	5	16	49	132	362	840	722	-
2004	1,560	5	4	-	1	4	19	42	119	252	648	466	-
2003	1,061	6	2	1	2	4	5	28	75	178	465	295	-
2002	915	2	2	1	3	1	8	26	67	180	346	279	-
2001	550	6	10	2	-	2	5	15	33	105	222	150	-
2000	497	8	5	-	-	3	9	16	34	98	189	135	-
1999	403	6	4	3	1	4	4	9	24	71	167	110	-
Non-Hispanic 1/, female													
2005	3,328	2	7	4	-	5	14	46	109	432	1,159	1,550	-
2004	2,640	4	8	1	-	7	11	43	115	330	962	1,159	-
2003	1,925	2	-	3	2	3	9	29	89	250	695	843	-
2002	1,466	6	1	2	2	3	9	26	68	171	511	667	-
2001	924	3	6	2	-	2	3	25	44	120	324	395	-
2000	780	2	3	-	1	-	8	13	28	113	260	352	-
1999	668	3	3	-	2	1	3	10	20	98	219	309	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Certain other intestinal infections (A04,A07-A09)													
Non-Hispanic white, both sexes													
2005 2/	5,082	3	7	6	2	8	22	77	196	719	1,874	2,168	-
2004 2/	3,919	6	9	1	1	8	21	67	206	528	1,511	1,561	-
2003 2/	2,776	3	1	3	3	5	11	44	140	397	1,088	1,081	-
2002	2,205	2	1	3	5	2	14	45	104	319	814	896	-
2001	1,357	5	13	3	-	2	6	35	62	202	509	520	-
2000	1,175	5	5	-	1	2	9	22	54	191	430	456	-
1999	990	5	6	1	2	5	4	15	36	155	360	401	-
Non-Hispanic white, male													
2005 2/	1,978	2	1	2	2	4	11	39	110	332	779	696	-
2004 2/	1,440	3	2	-	1	3	13	31	102	227	612	446	-
2003 2/	989	2	1	1	2	3	5	23	61	166	442	283	-
2002	840	-	1	1	3	1	6	24	52	159	328	265	-
2001	497	3	8	1	-	2	3	12	29	94	203	142	-
2000	453	4	4	-	-	2	6	12	29	89	180	127	-
1999	373	4	3	1	1	4	3	6	20	69	158	104	-
Non-Hispanic white, female													
2005 2/	3,104	1	6	4	-	4	11	38	86	387	1,095	1,472	-
2004 2/	2,479	3	7	1	-	5	8	36	104	301	899	1,115	-
2003 2/	1,787	1	-	2	1	2	6	21	79	231	646	798	-
2002	1,365	2	-	2	2	1	8	21	52	160	486	631	-
2001	860	2	5	2	-	-	3	23	33	108	306	378	-
2000	722	1	1	-	1	-	3	10	25	102	250	329	-
1999	617	1	3	-	1	1	1	9	16	86	202	297	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Certain other intestinal infections (A04,A07-A09)													
Non-Hispanic black, both sexes													
2005 2/	318	3	3	2	1	1	7	16	38	64	98	85	-
2004 2/	237	1	2	-	-	3	7	16	26	45	81	56	-
2003 2/	183	5	1	1	1	1	2	10	21	25	64	52	-
2002	155	5	1	-	-	2	3	5	29	31	35	44	-
2001	98	3	3	1	-	2	2	5	13	22	28	19	-
2000	83	4	2	-	-	1	6	7	6	15	16	26	-
1999	74	4	-	2	1	-	3	3	6	14	25	16	-
Non-Hispanic black, male													
2005 2/	129	2	2	2	1	1	4	9	17	26	47	18	-
2004 2/	96	1	1	-	-	1	4	10	16	21	26	16	-
2003 2/	65	4	1	-	-	1	-	4	12	9	23	11	-
2002	66	1	1	-	-	-	2	2	15	20	13	12	-
2001	42	2	2	1	-	-	2	3	4	10	12	6	-
2000	36	3	1	-	-	1	2	4	4	6	8	7	-
1999	25	2	-	2	-	-	1	2	2	2	9	5	-
Non-Hispanic black, female													
2005 2/	189	1	1	-	-	-	3	7	21	38	51	67	-
2004 2/	141	-	1	-	-	2	3	6	10	24	55	40	-
2003 2/	118	1	-	1	1	-	2	6	9	16	41	41	-
2002	89	4	-	-	-	2	1	3	14	11	22	32	-
2001	56	1	1	-	-	2	-	2	9	12	16	13	-
2000	47	1	1	-	-	-	4	3	2	9	8	19	-
1999	49	2	-	-	1	-	2	1	4	12	16	11	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Certain other intestinal infections (A04,A07-A09)													
Origin not stated 3/, both sexes													
2005	8	-	-	-	-	-	1	-	-	1	2	4	-
2004	8	2	-	-	-	-	-	-	1	-	2	3	-
2003	5	-	-	-	-	-	-	1	2	-	1	1	-
2002	6	-	-	-	-	-	-	1	1	1	1	2	-
2001	10	-	1	-	-	-	-	-	-	2	4	3	-
2000	5	-	-	-	-	-	-	-	-	2	2	1	-
1999	3	-	-	-	-	-	-	-	-	-	-	3	-
Origin not stated 3/, male													
2005	3	-	-	-	-	-	1	-	-	-	-	2	-
2004	5	2	-	-	-	-	-	-	1	-	1	1	-
2003	1	-	-	-	-	-	-	-	-	-	-	1	-
2002	1	-	-	-	-	-	-	1	-	-	-	-	-
2001	3	-	-	-	-	-	-	-	-	2	-	1	-
2000	4	-	-	-	-	-	-	-	-	1	2	1	-
1999	-	-	-	-	-	-	-	-	-	-	-	-	-
Origin not stated 3/, female													
2005	5	-	-	-	-	-	-	-	-	1	2	2	-
2004	3	-	-	-	-	-	-	-	-	-	1	2	-
2003	4	-	-	-	-	-	-	1	2	-	1	-	-
2002	5	-	-	-	-	-	-	-	1	1	1	2	-
2001	7	-	1	-	-	-	-	-	-	-	4	2	-
2000	1	-	-	-	-	-	-	-	-	1	-	-	-
1999	3	-	-	-	-	-	-	-	-	-	-	3	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Tuberculosis (A16-A19)													
All origins, both sexes													
2005	648	2	3	1	18	21	38	79	91	116	165	114	-
2004	657	1	1	2	7	16	53	80	79	137	165	116	-
2003	711	-	1	2	10	18	35	105	102	114	217	107	-
2002	784	1	-	-	9	18	52	81	107	161	234	121	-
2001	764	2	1	-	12	25	40	84	116	152	206	125	1
2000	776	1	6	1	11	20	42	91	81	155	252	115	1
1999	930	2	2	3	11	19	57	106	130	173	253	173	1
All origins, male													
2005	392	2	2	-	12	12	30	54	64	74	89	53	-
2004	406	-	1	1	4	12	40	53	60	93	84	58	-
2003	427	-	1	2	3	13	25	79	71	72	116	45	-
2002	462	-	-	-	5	11	38	55	75	102	121	55	-
2001	453	-	-	-	8	14	27	63	85	98	100	57	1
2000	468	-	1	-	7	14	28	76	59	87	139	56	1
1999	584	-	2	2	4	11	43	87	96	116	137	85	1
All origins, female													
2005	256	-	1	1	6	9	8	25	27	42	76	61	-
2004	251	1	-	1	3	4	13	27	19	44	81	58	-
2003	284	-	-	-	7	5	10	26	31	42	101	62	-
2002	322	1	-	-	4	7	14	26	32	59	113	66	-
2001	311	2	1	-	4	11	13	21	31	54	106	68	-
2000	308	1	5	1	4	6	14	15	22	68	113	59	-
1999	346	2	-	1	7	8	14	19	34	57	116	88	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Tuberculosis (A16-A19)													
Hispanic, both sexes													
2005	94	1	2	1	5	6	17	15	12	12	15	8	-
2004	110	1	-	1	3	6	12	11	14	23	26	13	-
2003	89	-	1	-	5	6	8	13	13	15	20	8	-
2002	95	-	-	-	2	4	11	10	17	21	19	11	-
2001	97	-	-	-	6	9	9	9	18	18	20	8	-
2000	83	-	2	1	3	7	3	9	12	13	23	10	-
1999	125	1	-	1	4	3	13	19	20	24	27	13	-
Hispanic, male													
2005	66	1	2	-	2	5	13	12	10	8	9	4	-
2004	73	-	-	1	2	4	8	7	11	16	17	7	-
2003	60	-	1	-	3	4	7	11	9	11	11	3	-
2002	52	-	-	-	1	3	9	6	6	11	11	5	-
2001	67	-	-	-	6	7	5	9	13	14	10	3	-
2000	46	-	-	-	1	4	2	5	7	8	14	5	-
1999	78	-	-	1	3	3	9	16	11	16	13	6	-
Hispanic, female													
2005	28	-	-	1	3	1	4	3	2	4	6	4	-
2004	37	1	-	-	1	2	4	4	3	7	9	6	-
2003	29	-	-	-	2	2	1	2	4	4	9	5	-
2002	43	-	-	-	1	1	2	4	11	10	8	6	-
2001	30	-	-	-	-	2	4	-	5	4	10	5	-
2000	37	-	2	1	2	3	1	4	5	5	9	5	-
1999	47	1	-	-	1	-	4	3	9	8	14	7	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Tuberculosis (A16-A19)													
Non-Hispanic 1/, both sexes													
2005	551	1	1	-	13	15	21	63	78	103	150	106	-
2004	542	-	1	1	4	10	41	69	65	113	138	100	-
2003	617	-	-	2	5	12	26	92	87	98	197	98	-
2002	682	1	-	-	6	14	39	70	88	139	215	110	-
2001	660	2	1	-	6	16	30	73	96	134	186	116	-
2000	683	1	4	-	8	13	39	80	68	140	226	104	-
1999	788	1	2	2	6	16	42	82	109	148	222	158	-
Non-Hispanic 1/, male													
2005	325	1	-	-	10	7	17	41	54	66	80	49	-
2004	330	-	1	-	2	8	32	46	49	76	66	50	-
2003	364	-	-	2	-	9	18	68	60	60	105	42	-
2002	404	-	-	-	4	8	27	48	67	90	110	50	-
2001	379	-	-	-	2	7	21	52	70	84	90	53	-
2000	414	-	1	-	6	10	26	69	51	78	122	51	-
1999	492	-	2	1	-	8	32	66	84	99	122	78	-
Non-Hispanic 1/, female													
2005	226	-	1	-	3	8	4	22	24	37	70	57	-
2004	212	-	-	1	2	2	9	23	16	37	72	50	-
2003	253	-	-	-	5	3	8	24	27	38	92	56	-
2002	278	1	-	-	2	6	12	22	21	49	105	60	-
2001	281	2	1	-	4	9	9	21	26	50	96	63	-
2000	269	1	3	-	2	3	13	11	17	62	104	53	-
1999	296	1	-	1	6	8	10	16	25	49	100	80	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Tuberculosis (A16-A19)													
Non-Hispanic white, both sexes													
2005 2/.....	283	1	-	-	4	4	8	29	34	51	82	70	-
2004 2/.....	282	-	1	-	-	2	17	28	28	55	88	63	-
2003 2/.....	347	-	-	2	1	2	3	32	47	60	130	70	-
2002.....	366	-	-	-	1	2	10	26	41	66	145	75	-
2001.....	368	1	1	-	2	2	12	32	48	72	120	78	-
2000.....	353	-	2	-	3	4	10	34	22	72	131	75	-
1999.....	430	-	-	-	-	2	11	30	53	84	142	108	-
Non-Hispanic white, male													
2005 2/.....	162	1	-	-	4	2	6	23	25	30	40	31	-
2004 2/.....	166	-	1	-	-	1	11	20	23	37	42	31	-
2003 2/.....	195	-	-	2	-	1	2	26	34	34	66	30	-
2002.....	218	-	-	-	1	-	7	20	31	52	72	35	-
2001.....	200	-	-	-	-	2	10	20	37	39	59	33	-
2000.....	205	-	-	-	3	3	9	28	17	44	63	38	-
1999.....	263	-	-	-	-	1	9	27	41	57	77	51	-
Non-Hispanic white, female													
2005 2/.....	121	-	-	-	-	2	2	6	9	21	42	39	-
2004 2/.....	116	-	-	-	-	1	6	8	5	18	46	32	-
2003 2/.....	152	-	-	-	1	1	1	6	13	26	64	40	-
2002.....	148	-	-	-	-	2	3	6	10	14	73	40	-
2001.....	168	1	1	-	2	-	2	12	11	33	61	45	-
2000.....	148	-	2	-	-	1	1	6	5	28	68	37	-
1999.....	167	-	-	-	-	1	2	3	12	27	65	57	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Tuberculosis (A16-A19)													
Non-Hispanic black, both sexes													
2005 2/.....	150	-	1	-	6	7	10	23	27	31	29	16	-
2004 2/.....	183	-	-	1	3	6	21	32	26	40	30	24	-
2003 2/.....	168	-	-	-	4	6	15	47	29	20	33	14	-
2002.....	198	-	-	-	3	8	23	34	31	43	37	19	-
2001.....	200	-	-	-	1	9	16	36	37	44	35	22	-
2000.....	238	-	2	-	2	7	25	41	33	52	58	18	-
1999.....	239	1	2	1	4	9	24	41	39	42	50	26	-
Non-Hispanic black, male													
2005 2/.....	92	-	-	-	4	2	9	15	18	25	13	6	-
2004 2/.....	114	-	-	-	2	5	18	18	20	25	15	11	-
2003 2/.....	111	-	-	-	-	5	11	35	19	15	20	6	-
2002.....	123	-	-	-	2	5	16	23	23	25	22	7	-
2001.....	124	-	-	-	-	3	10	29	26	30	14	12	-
2000.....	156	-	1	-	1	5	14	36	26	28	38	7	-
1999.....	154	-	2	-	-	4	18	33	31	28	25	13	-
Non-Hispanic black, female													
2005 2/.....	58	-	1	-	2	5	1	8	9	6	16	10	-
2004 2/.....	69	-	-	1	1	1	3	14	6	15	15	13	-
2003 2/.....	57	-	-	-	4	1	4	12	10	5	13	8	-
2002.....	75	-	-	-	1	3	7	11	8	18	15	12	-
2001.....	76	-	-	-	1	6	6	7	11	14	21	10	-
2000.....	82	-	1	-	1	2	11	5	7	24	20	11	-
1999.....	85	1	-	1	4	5	6	8	8	14	25	13	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Tuberculosis (A16-A19)													
Origin not stated 3/, both sexes													
2005	3	-	-	-	-	-	-	1	1	1	-	-	-
2004	5	-	-	-	-	-	-	-	-	1	1	3	-
2003	5	-	-	-	-	-	1	-	2	1	-	1	-
2002	7	-	-	-	1	-	2	1	2	1	-	-	-
2001	7	-	-	-	-	-	1	2	2	-	-	1	1
2000	10	-	-	-	-	-	-	2	1	2	3	1	1
1999	17	-	-	-	1	-	2	5	1	1	4	2	1
Origin not stated 3/, male													
2005	1	-	-	-	-	-	-	1	-	-	-	-	-
2004	3	-	-	-	-	-	-	-	-	1	1	1	-
2003	3	-	-	-	-	-	-	-	2	1	-	-	-
2002	6	-	-	-	-	-	2	1	2	1	-	-	-
2001	7	-	-	-	-	-	1	2	2	-	-	1	1
2000	8	-	-	-	-	-	-	2	1	1	3	-	1
1999	14	-	-	-	1	-	2	5	1	1	2	1	1
Origin not stated 3/, female													
2005	2	-	-	-	-	-	-	-	1	1	-	-	-
2004	2	-	-	-	-	-	-	-	-	-	-	2	-
2003	2	-	-	-	-	-	1	-	-	-	-	1	-
2002	1	-	-	-	1	-	-	-	-	-	-	-	-
2001	-	-	-	-	-	-	-	-	-	-	-	-	-
2000	2	-	-	-	-	-	-	-	-	1	-	1	-
1999	3	-	-	-	-	-	-	-	-	-	2	1	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Respiratory tuberculosis (A16)													
All origins, both sexes													
2005	480	2	1	-	9	11	28	62	62	81	129	95	-
2004	504	1	1	1	4	11	36	58	58	108	120	106	-
2003	562	-	1	2	9	6	26	86	81	88	177	86	-
2002	626	-	-	-	6	14	40	64	84	120	192	106	-
2001	600	2	1	-	7	17	28	61	96	115	166	106	1
2000	615	-	3	-	7	11	35	72	60	119	209	98	1
1999	764	1	1	1	8	14	45	76	109	139	216	153	1
All origins, male													
2005	291	2	1	-	4	3	23	43	46	57	68	44	-
2004	316	-	1	-	2	8	28	40	43	74	64	56	-
2003	344	-	1	2	3	6	17	64	57	61	97	36	-
2002	372	-	-	-	4	9	31	43	60	80	96	49	-
2001	371	-	-	-	5	10	18	49	78	76	84	50	1
2000	393	-	1	-	4	8	26	64	50	73	116	50	1
1999	486	-	1	1	2	8	33	64	83	97	120	76	1
All origins, female													
2005	189	-	-	-	5	8	5	19	16	24	61	51	-
2004	188	1	-	1	2	3	8	18	15	34	56	50	-
2003	218	-	-	-	6	-	9	22	24	27	80	50	-
2002	254	-	-	-	2	5	9	21	24	40	96	57	-
2001	229	2	1	-	2	7	10	12	18	39	82	56	-
2000	222	-	2	-	3	3	9	8	10	46	93	48	-
1999	278	1	-	-	6	6	12	12	26	42	96	77	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Respiratory tuberculosis (A16)													
Hispanic, both sexes													
2005	56	1	1	-	3	1	10	11	5	8	10	6	-
2004	83	1	-	-	1	2	9	8	11	17	22	12	-
2003	77	-	1	-	5	4	7	10	11	14	19	6	-
2002	78	-	-	-	1	4	10	8	12	16	16	11	-
2001	77	-	-	-	3	8	5	7	15	17	17	5	-
2000	60	-	1	-	2	3	2	5	9	12	18	8	-
1999	93	-	-	-	2	1	10	10	16	18	24	12	-
Hispanic, male													
2005	36	1	1	-	-	-	8	9	3	5	6	3	-
2004	54	-	-	-	-	1	7	5	8	11	15	7	-
2003	55	-	1	-	3	4	6	9	7	11	11	3	-
2002	44	-	-	-	1	3	8	4	5	9	9	5	-
2001	56	-	-	-	3	6	3	7	12	13	10	2	-
2000	40	-	-	-	1	3	2	3	6	8	12	5	-
1999	55	-	-	-	1	1	6	9	9	13	11	5	-
Hispanic, female													
2005	20	-	-	-	3	1	2	2	2	3	4	3	-
2004	29	1	-	-	1	1	2	3	3	6	7	5	-
2003	22	-	-	-	2	-	1	1	4	3	8	3	-
2002	34	-	-	-	-	1	2	4	7	7	7	6	-
2001	21	-	-	-	-	2	2	-	3	4	7	3	-
2000	20	-	1	-	1	-	-	2	3	4	6	3	-
1999	38	-	-	-	1	-	4	1	7	5	13	7	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Respiratory tuberculosis (A16)													
Non-Hispanic 1/, both sexes													
2005	423	1	-	-	6	10	18	51	56	73	119	89	-
2004	416	-	1	1	3	9	27	50	47	90	97	91	-
2003	481	-	-	2	4	2	18	76	68	74	158	79	-
2002	543	-	-	-	5	10	28	56	70	103	176	95	-
2001	517	2	1	-	4	9	22	53	79	98	149	100	-
2000	546	-	2	-	5	8	33	65	50	105	188	90	-
1999	657	1	1	1	5	13	33	63	92	120	189	139	-
Non-Hispanic 1/, male													
2005	255	1	-	-	4	3	15	34	43	52	62	41	-
2004	259	-	1	-	2	7	21	35	35	62	48	48	-
2003	287	-	-	2	-	2	11	55	48	50	86	33	-
2002	323	-	-	-	3	6	21	39	53	70	87	44	-
2001	309	-	-	-	2	4	14	41	64	63	74	47	-
2000	345	-	1	-	3	5	24	59	43	64	101	45	-
1999	419	-	1	1	-	7	25	52	73	83	107	70	-
Non-Hispanic 1/, female													
2005	168	-	-	-	2	7	3	17	13	21	57	48	-
2004	157	-	-	1	1	2	6	15	12	28	49	43	-
2003	194	-	-	-	4	-	7	21	20	24	72	46	-
2002	220	-	-	-	2	4	7	17	17	33	89	51	-
2001	208	2	1	-	2	5	8	12	15	35	75	53	-
2000	201	-	1	-	2	3	9	6	7	41	87	45	-
1999	238	1	-	-	5	6	8	11	19	37	82	69	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Respiratory tuberculosis (A16)													
Non-Hispanic white, both sexes													
2005 2/	219	1	-	-	2	3	8	23	26	36	65	55	-
2004 2/	219	-	1	-	-	2	11	20	21	46	61	57	-
2003 2/	270	-	-	2	1	-	2	25	37	43	106	54	-
2002	296	-	-	-	1	1	9	21	32	53	114	65	-
2001	288	1	1	-	1	-	8	23	41	52	97	64	-
2000	288	-	-	-	2	2	9	29	17	52	112	65	-
1999	371	-	-	-	-	2	9	27	45	69	119	100	-
Non-Hispanic white, male													
2005 2/	127	1	-	-	2	1	6	19	22	23	29	24	-
2004 2/	133	-	1	-	-	1	8	13	18	32	31	29	-
2003 2/	154	-	-	2	-	-	1	20	28	28	53	22	-
2002	177	-	-	-	1	-	7	16	24	44	53	32	-
2001	162	-	-	-	-	-	6	16	35	29	49	27	-
2000	174	-	-	-	2	1	9	26	14	34	54	34	-
1999	234	-	-	-	-	1	8	24	36	48	67	50	-
Non-Hispanic white, female													
2005 2/	92	-	-	-	-	2	2	4	4	13	36	31	-
2004 2/	86	-	-	-	-	1	3	7	3	14	30	28	-
2003 2/	116	-	-	-	1	-	1	5	9	15	53	32	-
2002	119	-	-	-	-	1	2	5	8	9	61	33	-
2001	126	1	1	-	1	-	2	7	6	23	48	37	-
2000	114	-	-	-	-	1	-	3	3	18	58	31	-
1999	137	-	-	-	-	1	1	3	9	21	52	50	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Respiratory tuberculosis (A16)													
Non-Hispanic black, both sexes													
2005 2/	113	-	-	-	3	5	10	19	18	23	20	15	-
2004 2/	135	-	-	1	2	5	15	22	17	30	21	22	-
2003 2/	134	-	-	-	3	2	10	41	23	17	26	12	-
2002	160	-	-	-	2	8	14	29	26	32	34	15	-
2001	157	-	-	-	1	6	12	28	30	32	28	20	-
2000	187	-	2	-	1	6	20	31	22	43	46	16	-
1999	194	1	1	-	4	8	19	29	35	32	43	22	-
Non-Hispanic black, male													
2005 2/	71	-	-	-	2	1	9	13	12	20	9	5	-
2004 2/	84	-	-	-	2	4	12	15	13	18	9	11	-
2003 2/	90	-	-	-	-	2	7	29	16	13	17	6	-
2002	101	-	-	-	1	5	10	20	20	19	21	5	-
2001	103	-	-	-	-	2	7	24	23	23	12	12	-
2000	129	-	1	-	-	4	12	28	22	26	30	6	-
1999	122	-	1	-	-	3	13	24	28	21	22	10	-
Non-Hispanic black, female													
2005 2/	42	-	-	-	1	4	1	6	6	3	11	10	-
2004 2/	51	-	-	1	-	1	3	7	4	12	12	11	-
2003 2/	44	-	-	-	3	-	3	12	7	4	9	6	-
2002	59	-	-	-	1	3	4	9	6	13	13	10	-
2001	54	-	-	-	1	4	5	4	7	9	16	8	-
2000	58	-	1	-	1	2	8	3	-	17	16	10	-
1999	72	1	-	-	4	5	6	5	7	11	21	12	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Respiratory tuberculosis (A16)													
Origin not stated 3/, both sexes													
2005	1	-	-	-	-	-	-	-	1	-	-	-	-
2004	5	-	-	-	-	-	-	-	-	1	1	3	-
2003	4	-	-	-	-	-	1	-	2	-	-	1	-
2002	5	-	-	-	-	-	2	-	2	1	-	-	-
2001	6	-	-	-	-	-	1	1	2	-	-	1	1
2000	9	-	-	-	-	-	-	2	1	2	3	-	1
1999	14	-	-	-	1	-	2	3	1	1	3	2	1
Origin not stated 3/, male													
2005	-	-	-	-	-	-	-	-	-	-	-	-	-
2004	3	-	-	-	-	-	-	-	-	1	1	1	-
2003	2	-	-	-	-	-	-	-	2	-	-	-	-
2002	5	-	-	-	-	-	2	-	2	1	-	-	-
2001	6	-	-	-	-	-	1	1	2	-	-	1	1
2000	8	-	-	-	-	-	-	2	1	1	3	-	1
1999	12	-	-	-	1	-	2	3	1	1	2	1	1
Origin not stated 3/, female													
2005	1	-	-	-	-	-	-	-	1	-	-	-	-
2004	2	-	-	-	-	-	-	-	-	-	-	2	-
2003	2	-	-	-	-	-	1	-	-	-	-	1	-
2002	-	-	-	-	-	-	-	-	-	-	-	-	-
2001	-	-	-	-	-	-	-	-	-	-	-	-	-
2000	1	-	-	-	-	-	-	-	-	1	-	-	-
1999	2	-	-	-	-	-	-	-	-	-	1	1	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Other tuberculosis (A17-A19)													
All origins, both sexes													
2005	168	-	2	1	9	10	10	17	29	35	36	19	-
2004	153	-	-	1	3	5	17	22	21	29	45	10	-
2003	149	-	-	-	1	12	9	19	21	26	40	21	-
2002	158	1	-	-	3	4	12	17	23	41	42	15	-
2001	164	-	-	-	5	8	12	23	20	37	40	19	-
2000	161	1	3	1	4	9	7	19	21	36	43	17	-
1999	166	1	1	2	3	5	12	30	21	34	37	20	-
All origins, male													
2005	101	-	1	-	8	9	7	11	18	17	21	9	-
2004	90	-	-	1	2	4	12	13	17	19	20	2	-
2003	83	-	-	-	-	7	8	15	14	11	19	9	-
2002	90	-	-	-	1	2	7	12	15	22	25	6	-
2001	82	-	-	-	3	4	9	14	7	22	16	7	-
2000	75	-	-	-	3	6	2	12	9	14	23	6	-
1999	98	-	1	1	2	3	10	23	13	19	17	9	-
All origins, female													
2005	67	-	1	1	1	1	3	6	11	18	15	10	-
2004	63	-	-	-	1	1	5	9	4	10	25	8	-
2003	66	-	-	-	1	5	1	4	7	15	21	12	-
2002	68	1	-	-	2	2	5	5	8	19	17	9	-
2001	82	-	-	-	2	4	3	9	13	15	24	12	-
2000	86	1	3	1	1	3	5	7	12	22	20	11	-
1999	68	1	-	1	1	2	2	7	8	15	20	11	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Other tuberculosis (A17-A19)													
Hispanic, both sexes													
2005	38	-	1	1	2	5	7	4	7	4	5	2	-
2004	27	-	-	1	2	4	3	3	3	6	4	1	-
2003	12	-	-	-	-	2	1	3	2	1	1	2	-
2002	17	-	-	-	1	-	1	2	5	5	3	-	-
2001	20	-	-	-	3	1	4	2	3	1	3	3	-
2000	23	-	1	1	1	4	1	4	3	1	5	2	-
1999	32	1	-	1	2	2	3	9	4	6	3	1	-
Hispanic, male													
2005	30	-	1	-	2	5	5	3	7	3	3	1	-
2004	19	-	-	1	2	3	1	2	3	5	2	-	-
2003	5	-	-	-	-	-	1	2	2	-	-	-	-
2002	8	-	-	-	-	-	1	2	1	2	2	-	-
2001	11	-	-	-	3	1	2	2	1	1	-	1	-
2000	6	-	-	-	-	1	-	2	1	-	2	-	-
1999	23	-	-	1	2	2	3	7	2	3	2	1	-
Hispanic, female													
2005	8	-	-	1	-	-	2	1	-	1	2	1	-
2004	8	-	-	-	-	1	2	1	-	1	2	1	-
2003	7	-	-	-	-	2	-	1	-	1	1	2	-
2002	9	-	-	-	1	-	-	-	4	3	1	-	-
2001	9	-	-	-	-	-	2	-	2	-	3	2	-
2000	17	-	1	1	1	3	1	2	2	1	3	2	-
1999	9	1	-	-	-	-	-	2	2	3	1	-	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Other tuberculosis (A17-A19)													
Non-Hispanic 1/, both sexes													
2005	128	-	1	-	7	5	3	12	22	30	31	17	-
2004	126	-	-	-	1	1	14	19	18	23	41	9	-
2003	136	-	-	-	1	10	8	16	19	24	39	19	-
2002	139	1	-	-	1	4	11	14	18	36	39	15	-
2001	143	-	-	-	2	7	8	20	17	36	37	16	-
2000	137	1	2	-	3	5	6	15	18	35	38	14	-
1999	131	-	1	1	1	3	9	19	17	28	33	19	-
Non-Hispanic 1/, male													
2005	70	-	-	-	6	4	2	7	11	14	18	8	-
2004	71	-	-	-	-	1	11	11	14	14	18	2	-
2003	77	-	-	-	-	7	7	13	12	10	19	9	-
2002	81	-	-	-	1	2	6	9	14	20	23	6	-
2001	70	-	-	-	-	3	7	11	6	21	16	6	-
2000	69	-	-	-	3	5	2	10	8	14	21	6	-
1999	73	-	1	-	-	1	7	14	11	16	15	8	-
Non-Hispanic 1/, female													
2005	58	-	1	-	1	1	1	5	11	16	13	9	-
2004	55	-	-	-	1	-	3	8	4	9	23	7	-
2003	59	-	-	-	1	3	1	3	7	14	20	10	-
2002	58	1	-	-	-	2	5	5	4	16	16	9	-
2001	73	-	-	-	2	4	1	9	11	15	21	10	-
2000	68	1	2	-	-	-	4	5	10	21	17	8	-
1999	58	-	-	1	1	2	2	5	6	12	18	11	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Other tuberculosis (A17-A19)													
Non-Hispanic white, both sexes													
2005 2/	64	-	-	-	2	1	-	6	8	15	17	15	-
2004 2/	63	-	-	-	-	-	6	8	7	9	27	6	-
2003 2/	77	-	-	-	-	2	1	7	10	17	24	16	-
2002	70	-	-	-	-	1	1	5	9	13	31	10	-
2001	80	-	-	-	1	2	4	9	7	20	23	14	-
2000	65	-	2	-	1	2	1	5	5	20	19	10	-
1999	59	-	-	-	-	-	2	3	8	15	23	8	-
Non-Hispanic white, male													
2005 2/	35	-	-	-	2	1	-	4	3	7	11	7	-
2004 2/	33	-	-	-	-	-	3	7	5	5	11	2	-
2003 2/	41	-	-	-	-	1	1	6	6	6	13	8	-
2002	41	-	-	-	-	-	-	4	7	8	19	3	-
2001	38	-	-	-	-	2	4	4	2	10	10	6	-
2000	31	-	-	-	1	2	-	2	3	10	9	4	-
1999	29	-	-	-	-	-	1	3	5	9	10	1	-
Non-Hispanic white, female													
2005 2/	29	-	-	-	-	-	-	2	5	8	6	8	-
2004 2/	30	-	-	-	-	-	3	1	2	4	16	4	-
2003 2/	36	-	-	-	-	1	-	1	4	11	11	8	-
2002	29	-	-	-	-	1	1	1	2	5	12	7	-
2001	42	-	-	-	1	-	-	5	5	10	13	8	-
2000	34	-	2	-	-	-	1	3	2	10	10	6	-
1999	30	-	-	-	-	-	1	-	3	6	13	7	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Other tuberculosis (A17-A19)													
Non-Hispanic black, both sexes													
2005 2/	37	-	1	-	3	2	-	4	9	8	9	1	-
2004 2/	48	-	-	-	1	1	6	10	9	10	9	2	-
2003 2/	34	-	-	-	1	4	5	6	6	3	7	2	-
2002	38	-	-	-	1	-	9	5	5	11	3	4	-
2001	43	-	-	-	-	3	4	8	7	12	7	2	-
2000	51	-	-	-	1	1	5	10	11	9	12	2	-
1999	45	-	1	1	-	1	5	12	4	10	7	4	-
Non-Hispanic black, male													
2005 2/	21	-	-	-	2	1	-	2	6	5	4	1	-
2004 2/	30	-	-	-	-	1	6	3	7	7	6	-	-
2003 2/	21	-	-	-	-	3	4	6	3	2	3	-	-
2002	22	-	-	-	1	-	6	3	3	6	1	2	-
2001	21	-	-	-	-	1	3	5	3	7	2	-	-
2000	27	-	-	-	1	1	2	8	4	2	8	1	-
1999	32	-	1	-	-	1	5	9	3	7	3	3	-
Non-Hispanic black, female													
2005 2/	16	-	1	-	1	1	-	2	3	3	5	-	-
2004 2/	18	-	-	-	1	-	-	7	2	3	3	2	-
2003 2/	13	-	-	-	1	1	1	-	3	1	4	2	-
2002	16	-	-	-	-	-	3	2	2	5	2	2	-
2001	22	-	-	-	-	2	1	3	4	5	5	2	-
2000	24	-	-	-	-	-	3	2	7	7	4	1	-
1999	13	-	-	1	-	-	-	3	1	3	4	1	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Other tuberculosis (A17-A19)													
Origin not stated 3/, both sexes													
2005	2	-	-	-	-	-	-	1	-	1	-	-	-
2004	-	-	-	-	-	-	-	-	-	-	-	-	-
2003	1	-	-	-	-	-	-	-	-	1	-	-	-
2002	2	-	-	-	1	-	-	1	-	-	-	-	-
2001	1	-	-	-	-	-	-	1	-	-	-	-	-
2000	1	-	-	-	-	-	-	-	-	-	-	1	-
1999	3	-	-	-	-	-	-	2	-	-	1	-	-
Origin not stated 3/, male													
2005	1	-	-	-	-	-	-	1	-	-	-	-	-
2004	-	-	-	-	-	-	-	-	-	-	-	-	-
2003	1	-	-	-	-	-	-	-	-	1	-	-	-
2002	1	-	-	-	-	-	-	1	-	-	-	-	-
2001	1	-	-	-	-	-	-	1	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-	-	-	-
1999	2	-	-	-	-	-	-	2	-	-	-	-	-
Origin not stated 3/, female													
2005	1	-	-	-	-	-	-	-	-	1	-	-	-
2004	-	-	-	-	-	-	-	-	-	-	-	-	-
2003	-	-	-	-	-	-	-	-	-	-	-	-	-
2002	1	-	-	-	1	-	-	-	-	-	-	-	-
2001	-	-	-	-	-	-	-	-	-	-	-	-	-
2000	1	-	-	-	-	-	-	-	-	-	-	1	-
1999	1	-	-	-	-	-	-	-	-	-	1	-	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Whooping cough (A37)													
All origins, both sexes													
2005	31	28	-	1	-	-	-	-	-	1	1	-	-
2004	16	15	-	1	-	-	-	-	-	-	-	-	-
2003	11	10	1	-	-	-	-	-	-	-	-	-	-
2002	18	18	-	-	-	-	-	-	-	-	-	-	-
2001	17	16	1	-	-	-	-	-	-	-	-	-	-
2000	12	12	-	-	-	-	-	-	-	-	-	-	-
1999	7	7	-	-	-	-	-	-	-	-	-	-	-
All origins, male													
2005	19	18	-	1	-	-	-	-	-	-	-	-	-
2004	10	9	-	1	-	-	-	-	-	-	-	-	-
2003	1	1	-	-	-	-	-	-	-	-	-	-	-
2002	6	6	-	-	-	-	-	-	-	-	-	-	-
2001	7	6	1	-	-	-	-	-	-	-	-	-	-
2000	5	5	-	-	-	-	-	-	-	-	-	-	-
1999	1	1	-	-	-	-	-	-	-	-	-	-	-
All origins, female													
2005	12	10	-	-	-	-	-	-	-	1	1	-	-
2004	6	6	-	-	-	-	-	-	-	-	-	-	-
2003	10	9	1	-	-	-	-	-	-	-	-	-	-
2002	12	12	-	-	-	-	-	-	-	-	-	-	-
2001	10	10	-	-	-	-	-	-	-	-	-	-	-
2000	7	7	-	-	-	-	-	-	-	-	-	-	-
1999	6	6	-	-	-	-	-	-	-	-	-	-	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Whooping cough (A37)													
Hispanic, both sexes													
2005	12	11	-	1	-	-	-	-	-	-	-	-	-
2004	6	5	-	1	-	-	-	-	-	-	-	-	-
2003	5	5	-	-	-	-	-	-	-	-	-	-	-
2002	3	3	-	-	-	-	-	-	-	-	-	-	-
2001	10	10	-	-	-	-	-	-	-	-	-	-	-
2000	7	7	-	-	-	-	-	-	-	-	-	-	-
1999	3	3	-	-	-	-	-	-	-	-	-	-	-
Hispanic, male													
2005	8	7	-	1	-	-	-	-	-	-	-	-	-
2004	4	3	-	1	-	-	-	-	-	-	-	-	-
2003	-	-	-	-	-	-	-	-	-	-	-	-	-
2002	2	2	-	-	-	-	-	-	-	-	-	-	-
2001	4	4	-	-	-	-	-	-	-	-	-	-	-
2000	2	2	-	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-	-	-	-
Hispanic, female													
2005	4	4	-	-	-	-	-	-	-	-	-	-	-
2004	2	2	-	-	-	-	-	-	-	-	-	-	-
2003	5	5	-	-	-	-	-	-	-	-	-	-	-
2002	1	1	-	-	-	-	-	-	-	-	-	-	-
2001	6	6	-	-	-	-	-	-	-	-	-	-	-
2000	5	5	-	-	-	-	-	-	-	-	-	-	-
1999	3	3	-	-	-	-	-	-	-	-	-	-	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Whooping cough (A37)													
Non-Hispanic 1/, both sexes													
2005	19	17	-	-	-	-	-	-	-	1	1	-	-
2004	9	9	-	-	-	-	-	-	-	-	-	-	-
2003	6	5	1	-	-	-	-	-	-	-	-	-	-
2002	15	15	-	-	-	-	-	-	-	-	-	-	-
2001	7	6	1	-	-	-	-	-	-	-	-	-	-
2000	5	5	-	-	-	-	-	-	-	-	-	-	-
1999	4	4	-	-	-	-	-	-	-	-	-	-	-
Non-Hispanic 1/, male													
2005	11	11	-	-	-	-	-	-	-	-	-	-	-
2004	5	5	-	-	-	-	-	-	-	-	-	-	-
2003	1	1	-	-	-	-	-	-	-	-	-	-	-
2002	4	4	-	-	-	-	-	-	-	-	-	-	-
2001	3	2	1	-	-	-	-	-	-	-	-	-	-
2000	3	3	-	-	-	-	-	-	-	-	-	-	-
1999	1	1	-	-	-	-	-	-	-	-	-	-	-
Non-Hispanic 1/, female													
2005	8	6	-	-	-	-	-	-	-	1	1	-	-
2004	4	4	-	-	-	-	-	-	-	-	-	-	-
2003	5	4	1	-	-	-	-	-	-	-	-	-	-
2002	11	11	-	-	-	-	-	-	-	-	-	-	-
2001	4	4	-	-	-	-	-	-	-	-	-	-	-
2000	2	2	-	-	-	-	-	-	-	-	-	-	-
1999	3	3	-	-	-	-	-	-	-	-	-	-	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Whooping cough (A37)													
Non-Hispanic white, both sexes													
2005 2/	11	10	-	-	-	-	-	-	-	-	1	-	-
2004 2/	8	8	-	-	-	-	-	-	-	-	-	-	-
2003 2/	5	4	1	-	-	-	-	-	-	-	-	-	-
2002	12	12	-	-	-	-	-	-	-	-	-	-	-
2001	6	5	1	-	-	-	-	-	-	-	-	-	-
2000	4	4	-	-	-	-	-	-	-	-	-	-	-
1999	3	3	-	-	-	-	-	-	-	-	-	-	-
Non-Hispanic white, male													
2005 2/	5	5	-	-	-	-	-	-	-	-	-	-	-
2004 2/	5	5	-	-	-	-	-	-	-	-	-	-	-
2003 2/	-	-	-	-	-	-	-	-	-	-	-	-	-
2002	3	3	-	-	-	-	-	-	-	-	-	-	-
2001	3	2	1	-	-	-	-	-	-	-	-	-	-
2000	2	2	-	-	-	-	-	-	-	-	-	-	-
1999	1	1	-	-	-	-	-	-	-	-	-	-	-
Non-Hispanic white, female													
2005 2/	6	5	-	-	-	-	-	-	-	-	1	-	-
2004 2/	3	3	-	-	-	-	-	-	-	-	-	-	-
2003 2/	5	4	1	-	-	-	-	-	-	-	-	-	-
2002	9	9	-	-	-	-	-	-	-	-	-	-	-
2001	3	3	-	-	-	-	-	-	-	-	-	-	-
2000	2	2	-	-	-	-	-	-	-	-	-	-	-
1999	2	2	-	-	-	-	-	-	-	-	-	-	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Whooping cough (A37)													
Non-Hispanic black, both sexes													
2005 2/	5	5	-	-	-	-	-	-	-	-	-	-	-
2004 2/	1	1	-	-	-	-	-	-	-	-	-	-	-
2003 2/	-	-	-	-	-	-	-	-	-	-	-	-	-
2002	2	2	-	-	-	-	-	-	-	-	-	-	-
2001	1	1	-	-	-	-	-	-	-	-	-	-	-
2000	1	1	-	-	-	-	-	-	-	-	-	-	-
1999	1	1	-	-	-	-	-	-	-	-	-	-	-
Non-Hispanic black, male													
2005 2/	4	4	-	-	-	-	-	-	-	-	-	-	-
2004 2/	-	-	-	-	-	-	-	-	-	-	-	-	-
2003 2/	-	-	-	-	-	-	-	-	-	-	-	-	-
2002	1	1	-	-	-	-	-	-	-	-	-	-	-
2001	-	-	-	-	-	-	-	-	-	-	-	-	-
2000	1	1	-	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-	-	-	-
Non-Hispanic black, female													
2005 2/	1	1	-	-	-	-	-	-	-	-	-	-	-
2004 2/	1	1	-	-	-	-	-	-	-	-	-	-	-
2003 2/	-	-	-	-	-	-	-	-	-	-	-	-	-
2002	1	1	-	-	-	-	-	-	-	-	-	-	-
2001	1	1	-	-	-	-	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-	-	-	-
1999	1	1	-	-	-	-	-	-	-	-	-	-	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Whooping cough (A37)													
Origin not stated 3/, both sexes													
2005	-	-	-	-	-	-	-	-	-	-	-	-	-
2004	1	1	-	-	-	-	-	-	-	-	-	-	-
2003	-	-	-	-	-	-	-	-	-	-	-	-	-
2002	-	-	-	-	-	-	-	-	-	-	-	-	-
2001	-	-	-	-	-	-	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-	-	-	-
Origin not stated 3/, male													
2005	-	-	-	-	-	-	-	-	-	-	-	-	-
2004	1	1	-	-	-	-	-	-	-	-	-	-	-
2003	-	-	-	-	-	-	-	-	-	-	-	-	-
2002	-	-	-	-	-	-	-	-	-	-	-	-	-
2001	-	-	-	-	-	-	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-	-	-	-
Origin not stated 3/, female													
2005	-	-	-	-	-	-	-	-	-	-	-	-	-
2004	-	-	-	-	-	-	-	-	-	-	-	-	-
2003	-	-	-	-	-	-	-	-	-	-	-	-	-
2002	-	-	-	-	-	-	-	-	-	-	-	-	-
2001	-	-	-	-	-	-	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-	-	-	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Scarlet fever and erysipelas (A38,A46)													
All origins, both sexes													
2005	3	-	1	-	-	-	-	-	-	-	-	2	-
2004	3	-	-	-	-	-	-	-	-	1	-	2	-
2003	1	-	-	-	-	-	-	-	-	-	-	1	-
2002	2	-	-	-	-	-	-	-	-	-	1	1	-
2001	3	-	-	1	-	-	-	-	-	1	-	1	-
2000	2	-	-	-	-	-	-	1	-	-	1	-	-
1999	5	-	-	-	-	-	-	-	-	2	1	2	-
All origins, male													
2005	2	-	1	-	-	-	-	-	-	-	-	1	-
2004	1	-	-	-	-	-	-	-	-	1	-	-	-
2003	-	-	-	-	-	-	-	-	-	-	-	-	-
2002	-	-	-	-	-	-	-	-	-	-	-	-	-
2001	2	-	-	1	-	-	-	-	-	-	-	1	-
2000	1	-	-	-	-	-	-	-	-	-	1	-	-
1999	-	-	-	-	-	-	-	-	-	-	-	-	-
All origins, female													
2005	1	-	-	-	-	-	-	-	-	-	-	1	-
2004	2	-	-	-	-	-	-	-	-	-	-	2	-
2003	1	-	-	-	-	-	-	-	-	-	-	1	-
2002	2	-	-	-	-	-	-	-	-	-	1	1	-
2001	1	-	-	-	-	-	-	-	-	1	-	-	-
2000	1	-	-	-	-	-	-	1	-	-	-	-	-
1999	5	-	-	-	-	-	-	-	-	2	1	2	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Scarlet fever and erysipelas (A38,A46)													
Hispanic, both sexes													
2005	1	-	1	-	-	-	-	-	-	-	-	-	-
2004	1	-	-	-	-	-	-	-	-	1	-	-	-
2003	-	-	-	-	-	-	-	-	-	-	-	-	-
2002	-	-	-	-	-	-	-	-	-	-	-	-	-
2001	-	-	-	-	-	-	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-	-	-	-
Hispanic, male													
2005	1	-	1	-	-	-	-	-	-	-	-	-	-
2004	1	-	-	-	-	-	-	-	-	1	-	-	-
2003	-	-	-	-	-	-	-	-	-	-	-	-	-
2002	-	-	-	-	-	-	-	-	-	-	-	-	-
2001	-	-	-	-	-	-	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-	-	-	-
Hispanic, female													
2005	-	-	-	-	-	-	-	-	-	-	-	-	-
2004	-	-	-	-	-	-	-	-	-	-	-	-	-
2003	-	-	-	-	-	-	-	-	-	-	-	-	-
2002	-	-	-	-	-	-	-	-	-	-	-	-	-
2001	-	-	-	-	-	-	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-	-	-	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Scarlet fever and erysipelas (A38,A46)													
Non-Hispanic 1/, both sexes													
2005	2	-	-	-	-	-	-	-	-	-	-	2	-
2004	2	-	-	-	-	-	-	-	-	-	-	2	-
2003	1	-	-	-	-	-	-	-	-	-	-	1	-
2002	2	-	-	-	-	-	-	-	-	-	1	1	-
2001	3	-	-	1	-	-	-	-	-	1	-	1	-
2000	2	-	-	-	-	-	-	1	-	-	1	-	-
1999	5	-	-	-	-	-	-	-	-	2	1	2	-
Non-Hispanic 1/, male													
2005	1	-	-	-	-	-	-	-	-	-	-	1	-
2004	-	-	-	-	-	-	-	-	-	-	-	-	-
2003	-	-	-	-	-	-	-	-	-	-	-	-	-
2002	-	-	-	-	-	-	-	-	-	-	-	-	-
2001	2	-	-	1	-	-	-	-	-	-	-	1	-
2000	1	-	-	-	-	-	-	-	-	-	1	-	-
1999	-	-	-	-	-	-	-	-	-	-	-	-	-
Non-Hispanic 1/, female													
2005	1	-	-	-	-	-	-	-	-	-	-	1	-
2004	2	-	-	-	-	-	-	-	-	-	-	2	-
2003	1	-	-	-	-	-	-	-	-	-	-	1	-
2002	2	-	-	-	-	-	-	-	-	-	1	1	-
2001	1	-	-	-	-	-	-	-	-	1	-	-	-
2000	1	-	-	-	-	-	-	1	-	-	-	-	-
1999	5	-	-	-	-	-	-	-	-	2	1	2	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Scarlet fever and erysipelas (A38,A46)													
Non-Hispanic white, both sexes													
2005 2/	2	-	-	-	-	-	-	-	-	-	-	2	-
2004 2/	2	-	-	-	-	-	-	-	-	-	-	2	-
2003 2/	1	-	-	-	-	-	-	-	-	-	-	1	-
2002	2	-	-	-	-	-	-	-	-	-	1	1	-
2001	3	-	-	1	-	-	-	-	-	1	-	1	-
2000	2	-	-	-	-	-	-	1	-	-	1	-	-
1999	4	-	-	-	-	-	-	-	-	1	1	2	-
Non-Hispanic white, male													
2005 2/	1	-	-	-	-	-	-	-	-	-	-	1	-
2004 2/	-	-	-	-	-	-	-	-	-	-	-	-	-
2003 2/	-	-	-	-	-	-	-	-	-	-	-	-	-
2002	-	-	-	-	-	-	-	-	-	-	-	-	-
2001	2	-	-	1	-	-	-	-	-	-	-	1	-
2000	1	-	-	-	-	-	-	-	-	-	1	-	-
1999	-	-	-	-	-	-	-	-	-	-	-	-	-
Non-Hispanic white, female													
2005 2/	1	-	-	-	-	-	-	-	-	-	-	1	-
2004 2/	2	-	-	-	-	-	-	-	-	-	-	2	-
2003 2/	1	-	-	-	-	-	-	-	-	-	-	1	-
2002	2	-	-	-	-	-	-	-	-	-	1	1	-
2001	1	-	-	-	-	-	-	-	-	1	-	-	-
2000	1	-	-	-	-	-	-	1	-	-	-	-	-
1999	4	-	-	-	-	-	-	-	-	1	1	2	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Scarlet fever and erysipelas (A38,A46)													
Non-Hispanic black, both sexes													
2005 2/	-	-	-	-	-	-	-	-	-	-	-	-	-
2004 2/	-	-	-	-	-	-	-	-	-	-	-	-	-
2003 2/	-	-	-	-	-	-	-	-	-	-	-	-	-
2002	-	-	-	-	-	-	-	-	-	-	-	-	-
2001	-	-	-	-	-	-	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-	-	-	-
1999	1	-	-	-	-	-	-	-	-	1	-	-	-
Non-Hispanic black, male													
2005 2/	-	-	-	-	-	-	-	-	-	-	-	-	-
2004 2/	-	-	-	-	-	-	-	-	-	-	-	-	-
2003 2/	-	-	-	-	-	-	-	-	-	-	-	-	-
2002	-	-	-	-	-	-	-	-	-	-	-	-	-
2001	-	-	-	-	-	-	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-	-	-	-
Non-Hispanic black, female													
2005 2/	-	-	-	-	-	-	-	-	-	-	-	-	-
2004 2/	-	-	-	-	-	-	-	-	-	-	-	-	-
2003 2/	-	-	-	-	-	-	-	-	-	-	-	-	-
2002	-	-	-	-	-	-	-	-	-	-	-	-	-
2001	-	-	-	-	-	-	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-	-	-	-
1999	1	-	-	-	-	-	-	-	-	1	-	-	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Scarlet fever and erysipelas (A38,A46)													
Origin not stated 3/, both sexes													
2005	-	-	-	-	-	-	-	-	-	-	-	-	-
2004	-	-	-	-	-	-	-	-	-	-	-	-	-
2003	-	-	-	-	-	-	-	-	-	-	-	-	-
2002	-	-	-	-	-	-	-	-	-	-	-	-	-
2001	-	-	-	-	-	-	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-	-	-	-
Origin not stated 3/, male													
2005	-	-	-	-	-	-	-	-	-	-	-	-	-
2004	-	-	-	-	-	-	-	-	-	-	-	-	-
2003	-	-	-	-	-	-	-	-	-	-	-	-	-
2002	-	-	-	-	-	-	-	-	-	-	-	-	-
2001	-	-	-	-	-	-	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-	-	-	-
Origin not stated 3/, female													
2005	-	-	-	-	-	-	-	-	-	-	-	-	-
2004	-	-	-	-	-	-	-	-	-	-	-	-	-
2003	-	-	-	-	-	-	-	-	-	-	-	-	-
2002	-	-	-	-	-	-	-	-	-	-	-	-	-
2001	-	-	-	-	-	-	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-	-	-	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Meningococcal infection (A39)													
All origins, both sexes													
2005	123	17	11	7	29	8	8	18	10	4	6	5	-
2004	138	8	18	13	31	14	15	15	6	7	3	8	-
2003	161	12	18	11	39	12	19	18	11	9	9	3	-
2002	161	17	12	20	33	16	9	19	13	7	10	5	-
2001	199	16	19	26	55	10	13	13	15	15	8	9	-
2000	211	22	19	18	56	15	13	29	9	8	15	7	-
1999	227	32	30	25	39	23	27	17	4	9	13	8	-
All origins, male													
2005	68	10	6	5	17	8	4	11	4	2	-	1	-
2004	79	2	11	9	19	10	8	12	4	3	1	-	-
2003	75	9	10	3	18	7	9	10	5	2	1	1	-
2002	83	8	9	13	17	10	5	10	4	2	3	2	-
2001	94	6	9	15	27	6	6	8	7	5	3	2	-
2000	106	14	12	7	35	6	9	11	2	2	7	1	-
1999	101	14	19	13	19	12	9	4	-	5	4	2	-
All origins, female													
2005	55	7	5	2	12	-	4	7	6	2	6	4	-
2004	59	6	7	4	12	4	7	3	2	4	2	8	-
2003	86	3	8	8	21	5	10	8	6	7	8	2	-
2002	78	9	3	7	16	6	4	9	9	5	7	3	-
2001	105	10	10	11	28	4	7	5	8	10	5	7	-
2000	105	8	7	11	21	9	4	18	7	6	8	6	-
1999	126	18	11	12	20	11	18	13	4	4	9	6	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Meningococcal infection (A39)													
Hispanic, both sexes													
2005	16	2	3	1	3	1	1	1	2	-	1	1	-
2004	20	2	5	1	3	3	-	4	-	1	-	1	-
2003	23	5	2	1	2	5	2	1	-	4	-	1	-
2002	19	7	-	1	5	3	1	1	1	-	-	-	-
2001	21	4	5	2	2	2	2	1	2	-	1	-	-
2000	34	9	3	2	9	5	-	3	1	-	2	-	-
1999	27	7	1	3	2	6	5	2	-	-	1	-	-
Hispanic, male													
2005	12	1	3	1	2	1	1	1	2	-	-	-	-
2004	12	1	2	1	-	3	-	4	-	1	-	-	-
2003	13	4	1	-	1	3	1	1	-	1	-	1	-
2002	9	3	-	1	2	2	-	-	1	-	-	-	-
2001	11	-	3	2	2	1	1	1	-	-	1	-	-
2000	16	4	2	-	6	2	-	1	-	-	1	-	-
1999	11	5	1	2	1	2	-	-	-	-	-	-	-
Hispanic, female													
2005	4	1	-	-	1	-	-	-	-	-	1	1	-
2004	8	1	3	-	3	-	-	-	-	-	-	1	-
2003	10	1	1	1	1	2	1	-	-	3	-	-	-
2002	10	4	-	-	3	1	1	1	-	-	-	-	-
2001	10	4	2	-	-	1	1	-	2	-	-	-	-
2000	18	5	1	2	3	3	-	2	1	-	1	-	-
1999	16	2	-	1	1	4	5	2	-	-	1	-	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Meningococcal infection (A39)													
Non-Hispanic 1/, both sexes													
2005	107	15	8	6	26	7	7	17	8	4	5	4	-
2004	118	6	13	12	28	11	15	11	6	6	3	7	-
2003	138	7	16	10	37	7	17	17	11	5	9	2	-
2002	142	10	12	19	28	13	8	18	12	7	10	5	-
2001	178	12	14	24	53	8	11	12	13	15	7	9	-
2000	177	13	16	16	47	10	13	26	8	8	13	7	-
1999	198	25	28	22	37	17	22	14	4	9	12	8	-
Non-Hispanic 1/, male													
2005	56	9	3	4	15	7	3	10	2	2	-	1	-
2004	67	1	9	8	19	7	8	8	4	2	1	-	-
2003	62	5	9	3	17	4	8	9	5	1	1	-	-
2002	74	5	9	12	15	8	5	10	3	2	3	2	-
2001	83	6	6	13	25	5	5	7	7	5	2	2	-
2000	90	10	10	7	29	4	9	10	2	2	6	1	-
1999	89	9	17	11	18	10	9	4	-	5	4	2	-
Non-Hispanic 1/, female													
2005	51	6	5	2	11	-	4	7	6	2	5	3	-
2004	51	5	4	4	9	4	7	3	2	4	2	7	-
2003	76	2	7	7	20	3	9	8	6	4	8	2	-
2002	68	5	3	7	13	5	3	8	9	5	7	3	-
2001	95	6	8	11	28	3	6	5	6	10	5	7	-
2000	87	3	6	9	18	6	4	16	6	6	7	6	-
1999	109	16	11	11	19	7	13	10	4	4	8	6	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Meningococcal infection (A39)													
Non-Hispanic white, both sexes													
2005 2/	86	12	8	5	20	5	4	14	7	3	4	4	-
2004 2/	90	5	10	11	21	7	10	7	5	5	3	6	-
2003 2/	99	7	10	6	28	3	12	10	7	5	9	2	-
2002	106	7	8	9	22	10	4	15	10	6	10	5	-
2001	138	10	10	18	38	5	9	11	9	14	6	8	-
2000	129	10	11	8	37	6	7	16	7	7	13	7	-
1999	155	23	24	15	27	8	14	12	4	9	11	8	-
Non-Hispanic white, male													
2005 2/	42	7	3	3	11	5	1	8	2	1	-	1	-
2004 2/	50	1	8	7	13	5	6	5	3	1	1	-	-
2003 2/	44	5	6	2	13	2	6	6	2	1	1	-	-
2002	58	5	6	8	12	7	3	8	3	1	3	2	-
2001	62	5	5	10	18	3	3	6	5	5	1	1	-
2000	60	7	6	4	22	1	5	4	2	2	6	1	-
1999	67	9	15	7	12	4	6	3	-	5	4	2	-
Non-Hispanic white, female													
2005 2/	44	5	5	2	9	-	3	6	5	2	4	3	-
2004 2/	40	4	2	4	8	2	4	2	2	4	2	6	-
2003 2/	55	2	4	4	15	1	6	4	5	4	8	2	-
2002	48	2	2	1	10	3	1	7	7	5	7	3	-
2001	76	5	5	8	20	2	6	5	4	9	5	7	-
2000	69	3	5	4	15	5	2	12	5	5	7	6	-
1999	88	14	9	8	15	4	8	9	4	4	7	6	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Meningococcal infection (A39)													
Non-Hispanic black, both sexes													
2005 2/	17	2	-	-	6	2	2	2	1	1	1	-	-
2004 2/	22	-	3	1	5	3	4	3	1	1	-	1	-
2003 2/	35	-	6	4	7	3	5	6	4	-	-	-	-
2002	29	3	3	8	5	2	4	3	1	-	-	-	-
2001	36	2	4	6	13	2	1	1	4	1	1	1	-
2000	44	3	5	8	7	4	6	9	1	1	-	-	-
1999	39	2	4	6	9	8	8	2	-	-	-	-	-
Non-Hispanic black, male													
2005 2/	11	1	-	-	4	2	1	2	-	1	-	-	-
2004 2/	13	-	1	1	4	2	1	2	1	1	-	-	-
2003 2/	16	-	3	1	3	1	2	3	3	-	-	-	-
2002	12	-	2	4	2	-	2	2	-	-	-	-	-
2001	18	1	1	3	6	1	1	1	2	-	1	1	-
2000	27	3	4	3	5	3	4	5	-	-	-	-	-
1999	20	-	2	3	6	5	3	1	-	-	-	-	-
Non-Hispanic black, female													
2005 2/	6	1	-	-	2	-	1	-	1	-	1	-	-
2004 2/	9	-	2	-	1	1	3	1	-	-	-	1	-
2003 2/	19	-	3	3	4	2	3	3	1	-	-	-	-
2002	17	3	1	4	3	2	2	1	1	-	-	-	-
2001	18	1	3	3	7	1	-	-	2	1	-	-	-
2000	17	-	1	5	2	1	2	4	1	1	-	-	-
1999	19	2	2	3	3	3	5	1	-	-	-	-	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Meningococcal infection (A39)													
Origin not stated 3/, both sexes													
2005	-	-	-	-	-	-	-	-	-	-	-	-	-
2004	-	-	-	-	-	-	-	-	-	-	-	-	-
2003	-	-	-	-	-	-	-	-	-	-	-	-	-
2002	-	-	-	-	-	-	-	-	-	-	-	-	-
2001	-	-	-	-	-	-	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-	-	-	-
1999	2	-	1	-	-	-	-	1	-	-	-	-	-
Origin not stated 3/, male													
2005	-	-	-	-	-	-	-	-	-	-	-	-	-
2004	-	-	-	-	-	-	-	-	-	-	-	-	-
2003	-	-	-	-	-	-	-	-	-	-	-	-	-
2002	-	-	-	-	-	-	-	-	-	-	-	-	-
2001	-	-	-	-	-	-	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-	-	-	-
1999	1	-	1	-	-	-	-	-	-	-	-	-	-
Origin not stated 3/, female													
2005	-	-	-	-	-	-	-	-	-	-	-	-	-
2004	-	-	-	-	-	-	-	-	-	-	-	-	-
2003	-	-	-	-	-	-	-	-	-	-	-	-	-
2002	-	-	-	-	-	-	-	-	-	-	-	-	-
2001	-	-	-	-	-	-	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-	-	-	-
1999	1	-	-	-	-	-	-	1	-	-	-	-	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Septicemia (A40-A41)													
All origins, both sexes													
2005	34,136	302	85	81	148	311	840	2,211	3,912	6,073	10,626	9,544	3
2004	33,373	271	84	66	135	328	846	2,251	3,745	5,983	10,586	9,075	3
2003	34,069	278	85	77	154	309	910	2,157	3,651	5,970	10,932	9,543	3
2002	33,865	296	79	95	118	314	856	2,074	3,360	6,336	11,010	9,324	3
2001	32,238	312	108	69	133	281	829	1,977	3,111	5,998	10,351	9,069	-
2000	31,224	274	99	63	100	280	877	1,845	2,899	5,704	9,938	9,144	1
1999	30,680	283	87	77	121	275	812	1,685	2,714	5,750	9,710	9,166	-
All origins, male													
2005	15,322	171	47	49	82	152	414	1,175	2,074	3,095	4,823	3,238	2
2004	15,011	167	41	31	71	165	433	1,207	1,934	3,017	4,870	3,074	1
2003	14,987	151	54	34	82	161	455	1,155	1,887	3,023	4,894	3,089	2
2002	14,947	158	41	52	59	166	441	1,124	1,748	3,154	5,028	2,973	3
2001	14,307	174	60	39	75	139	451	1,043	1,606	3,044	4,762	2,914	-
2000	13,537	139	53	38	60	158	451	1,039	1,513	2,867	4,482	2,736	1
1999	13,395	175	48	36	67	147	460	919	1,412	2,919	4,426	2,786	-
All origins, female													
2005	18,814	131	38	32	66	159	426	1,036	1,838	2,978	5,803	6,306	1
2004	18,362	104	43	35	64	163	413	1,044	1,811	2,966	5,716	6,001	2
2003	19,082	127	31	43	72	148	455	1,002	1,764	2,947	6,038	6,454	1
2002	18,918	138	38	43	59	148	415	950	1,612	3,182	5,982	6,351	-
2001	17,931	138	48	30	58	142	378	934	1,505	2,954	5,589	6,155	-
2000	17,687	135	46	25	40	122	426	806	1,386	2,837	5,456	6,408	-
1999	17,285	108	39	41	54	128	352	766	1,302	2,831	5,284	6,380	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Septicemia (A40-A41)													
Hispanic, both sexes													
2005	1,704	66	14	12	18	36	74	157	230	337	445	315	-
2004	1,648	55	24	9	13	35	88	162	216	319	437	290	-
2003	1,505	50	10	7	17	35	86	148	190	270	397	293	2
2002	1,521	65	16	14	15	38	67	136	165	341	382	282	-
2001	1,386	50	18	18	13	23	72	133	151	288	346	274	-
2000	1,226	62	15	7	12	31	79	122	145	235	300	218	-
1999	1,215	43	14	8	9	26	74	92	131	265	296	257	-
Hispanic, male													
2005	838	37	6	6	10	23	35	93	121	175	222	110	-
2004	801	36	14	6	7	16	53	106	109	154	195	105	-
2003	721	29	6	4	8	20	48	94	95	137	175	104	1
2002	744	34	9	8	8	25	47	94	84	161	176	98	-
2001	684	29	5	14	9	15	43	80	86	149	170	84	-
2000	625	33	9	5	6	21	46	78	80	126	143	78	-
1999	605	27	8	4	5	14	51	56	78	134	144	84	-
Hispanic, female													
2005	866	29	8	6	8	13	39	64	109	162	223	205	-
2004	847	19	10	3	6	19	35	56	107	165	242	185	-
2003	784	21	4	3	9	15	38	54	95	133	222	189	1
2002	777	31	7	6	7	13	20	42	81	180	206	184	-
2001	702	21	13	4	4	8	29	53	65	139	176	190	-
2000	601	29	6	2	6	10	33	44	65	109	157	140	-
1999	610	16	6	4	4	12	23	36	53	131	152	173	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Septicemia (A40-A41)													
Non-Hispanic 1/, both sexes													
2005	32,354	236	70	69	130	274	763	2,046	3,663	5,718	10,163	9,220	2
2004	31,656	212	60	57	122	292	758	2,080	3,517	5,652	10,132	8,772	2
2003	32,476	227	75	70	137	272	823	2,006	3,445	5,680	10,512	9,228	1
2002	32,225	227	63	81	102	275	787	1,931	3,182	5,969	10,593	9,012	3
2001	30,754	261	90	50	120	257	755	1,837	2,941	5,690	9,977	8,776	-
2000	29,864	208	83	56	88	248	793	1,713	2,739	5,442	9,601	8,892	1
1999	29,351	239	73	69	112	248	736	1,585	2,570	5,466	9,378	8,875	-
Non-Hispanic 1/, male													
2005	14,442	134	40	43	72	128	378	1,079	1,939	2,909	4,591	3,127	2
2004	14,174	128	27	25	64	148	380	1,094	1,817	2,857	4,670	2,964	-
2003	14,226	122	48	30	74	139	406	1,059	1,784	2,876	4,708	2,979	1
2002	14,136	120	32	44	50	141	392	1,025	1,657	2,975	4,830	2,867	3
2001	13,573	145	55	25	66	123	408	957	1,510	2,882	4,575	2,827	-
2000	12,842	104	44	33	54	137	401	957	1,424	2,724	4,316	2,647	1
1999	12,737	147	40	32	62	132	408	858	1,330	2,777	4,261	2,690	-
Non-Hispanic 1/, female													
2005	17,912	102	30	26	58	146	385	967	1,724	2,809	5,572	6,093	-
2004	17,482	84	33	32	58	144	378	986	1,700	2,795	5,462	5,808	2
2003	18,250	105	27	40	63	133	417	947	1,661	2,804	5,804	6,249	-
2002	18,089	107	31	37	52	134	395	906	1,525	2,994	5,763	6,145	-
2001	17,181	116	35	25	54	134	347	880	1,431	2,808	5,402	5,949	-
2000	17,022	104	39	23	34	111	392	756	1,315	2,718	5,285	6,245	-
1999	16,614	92	33	37	50	116	328	727	1,240	2,689	5,117	6,185	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Septicemia (A40-A41)													
Non-Hispanic white, both sexes													
2005 2/	25,496	100	44	47	77	157	480	1,342	2,641	4,413	8,381	7,814	-
2004 2/	25,041	112	36	41	70	172	474	1,359	2,513	4,328	8,426	7,509	1
2003 2/	25,664	96	43	47	86	155	513	1,295	2,471	4,315	8,803	7,840	-
2002	25,545	129	42	55	72	165	510	1,274	2,241	4,583	8,806	7,666	2
2001	24,389	122	63	35	69	145	458	1,181	2,081	4,385	8,352	7,498	-
2000	23,517	108	51	42	48	134	498	1,092	1,919	4,173	7,901	7,550	1
1999	23,023	121	47	47	71	155	439	1,015	1,792	4,140	7,717	7,479	-
Non-Hispanic white, male													
2005 2/	11,337	63	26	29	46	74	241	708	1,416	2,249	3,835	2,650	-
2004 2/	11,188	68	15	20	39	85	243	724	1,299	2,213	3,914	2,568	-
2003 2/	11,181	46	29	22	48	74	255	698	1,280	2,207	3,972	2,550	-
2002	11,148	67	25	34	37	86	249	680	1,204	2,310	4,006	2,448	2
2001	10,668	71	41	18	37	69	266	615	1,071	2,223	3,851	2,406	-
2000	10,056	55	26	25	33	74	262	613	1,015	2,131	3,579	2,242	1
1999	9,863	79	25	24	44	87	243	551	904	2,111	3,538	2,257	-
Non-Hispanic white, female													
2005 2/	14,159	37	18	18	31	83	239	634	1,225	2,164	4,546	5,164	-
2004 2/	13,853	44	21	21	31	87	231	635	1,214	2,115	4,512	4,941	1
2003 2/	14,483	50	14	25	38	81	258	597	1,191	2,108	4,831	5,290	-
2002	14,397	62	17	21	35	79	261	594	1,037	2,273	4,800	5,218	-
2001	13,721	51	22	17	32	76	192	566	1,010	2,162	4,501	5,092	-
2000	13,461	53	25	17	15	60	236	479	904	2,042	4,322	5,308	-
1999	13,160	42	22	23	27	68	196	464	888	2,029	4,179	5,222	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Septicemia (A40-A41)													
Non-Hispanic black, both sexes													
2005 2/	6,159	125	20	20	44	106	258	654	934	1,171	1,577	1,248	2
2004 2/	5,945	90	19	12	45	102	255	650	906	1,203	1,534	1,128	1
2003 2/	6,141	119	24	15	41	102	283	650	881	1,232	1,536	1,257	1
2002	6,074	85	17	20	24	102	254	593	868	1,251	1,633	1,226	1
2001	5,829	124	21	10	41	98	268	595	804	1,186	1,501	1,181	-
2000	5,863	96	25	11	33	107	269	572	756	1,173	1,576	1,245	-
1999	5,772	106	16	14	32	85	279	518	716	1,212	1,513	1,281	-
Non-Hispanic black, male													
2005 2/	2,755	65	13	13	24	49	121	344	485	586	654	399	2
2004 2/	2,660	51	9	4	22	52	125	334	465	582	675	341	-
2003 2/	2,748	70	14	6	19	56	134	338	462	601	663	384	1
2002	2,668	46	5	7	10	47	130	312	418	598	729	365	1
2001	2,611	67	12	3	25	46	128	304	411	586	656	373	-
2000	2,559	48	13	6	18	59	125	319	381	545	677	368	-
1999	2,594	60	10	7	14	41	154	279	392	609	648	380	-
Non-Hispanic black, female													
2005 2/	3,404	60	7	7	20	57	137	310	449	585	923	849	-
2004 2/	3,285	39	10	8	23	50	130	316	441	621	859	787	1
2003 2/	3,393	49	10	9	22	46	149	312	419	631	873	873	-
2002	3,406	39	12	13	14	55	124	281	450	653	904	861	-
2001	3,218	57	9	7	16	52	140	291	393	600	845	808	-
2000	3,304	48	12	5	15	48	144	253	375	628	899	877	-
1999	3,178	46	6	7	18	44	125	239	324	603	865	901	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Septicemia (A40-A41)													
Origin not stated 3/, both sexes													
2005	78	-	1	-	-	1	3	8	19	18	18	9	1
2004	69	4	-	-	-	1	-	9	12	12	17	13	1
2003	88	1	-	-	-	2	1	3	16	20	23	22	-
2002	119	4	-	-	1	1	2	7	13	26	35	30	-
2001	98	1	-	1	-	1	2	7	19	20	28	19	-
2000	134	4	1	-	-	1	5	10	15	27	37	34	-
1999	114	1	-	-	-	1	2	8	13	19	36	34	-
Origin not stated 3/, male													
2005	42	-	1	-	-	1	1	3	14	11	10	1	-
2004	36	3	-	-	-	1	-	7	8	6	5	5	1
2003	40	-	-	-	-	2	1	2	8	10	11	6	-
2002	67	4	-	-	1	-	2	5	7	18	22	8	-
2001	50	-	-	-	-	1	-	6	10	13	17	3	-
2000	70	2	-	-	-	-	4	4	9	17	23	11	-
1999	53	1	-	-	-	1	1	5	4	8	21	12	-
Origin not stated 3/, female													
2005	36	-	-	-	-	-	2	5	5	7	8	8	1
2004	33	1	-	-	-	-	-	2	4	6	12	8	-
2003	48	1	-	-	-	-	-	1	8	10	12	16	-
2002	52	-	-	-	-	1	-	2	6	8	13	22	-
2001	48	1	-	1	-	-	2	1	9	7	11	16	-
2000	64	2	1	-	-	1	1	6	6	10	14	23	-
1999	61	-	-	-	-	-	1	3	9	11	15	22	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Syphilis (A50-A53)													
All origins, both sexes													
2005	47	-	-	1	-	4	5	6	8	4	13	6	-
2004	43	-	-	-	-	3	3	3	4	6	12	12	-
2003	34	-	-	-	-	-	3	5	3	6	8	9	-
2002	41	1	1	-	-	-	2	5	5	6	12	9	-
2001	36	1	-	1	-	1	3	3	4	5	13	5	-
2000	41	1	-	-	-	1	-	6	3	7	13	10	-
1999	33	2	-	-	-	1	1	5	4	6	10	4	-
All origins, male													
2005	32	-	-	-	-	4	4	5	5	4	8	2	-
2004	21	-	-	-	-	2	2	1	2	4	6	4	-
2003	22	-	-	-	-	-	3	5	2	5	4	3	-
2002	21	-	1	-	-	-	2	4	2	5	4	3	-
2001	19	1	-	1	-	-	2	3	3	3	5	1	-
2000	24	1	-	-	-	1	-	5	3	2	7	5	-
1999	13	-	-	-	-	1	-	3	3	5	-	1	-
All origins, female													
2005	15	-	-	1	-	-	1	1	3	-	5	4	-
2004	22	-	-	-	-	1	1	2	2	2	6	8	-
2003	12	-	-	-	-	-	-	-	1	1	4	6	-
2002	20	1	-	-	-	-	-	1	3	1	8	6	-
2001	17	-	-	-	-	1	1	-	1	2	8	4	-
2000	17	-	-	-	-	-	-	1	-	5	6	5	-
1999	20	2	-	-	-	-	1	2	1	1	10	3	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Syphilis (A50-A53)													
Hispanic, both sexes													
2005	1	-	-	-	-	-	-	1	-	-	-	-	-
2004	1	-	-	-	-	-	-	-	-	-	1	-	-
2003	1	-	-	-	-	-	-	1	-	-	-	-	-
2002	1	-	-	-	-	-	-	-	-	-	1	-	-
2001	7	1	-	-	-	-	-	-	2	-	3	1	-
2000	1	-	-	-	-	-	-	-	-	-	1	-	-
1999	5	2	-	-	-	-	-	2	-	-	1	-	-
Hispanic, male													
2005	1	-	-	-	-	-	-	1	-	-	-	-	-
2004	1	-	-	-	-	-	-	-	-	-	1	-	-
2003	1	-	-	-	-	-	-	1	-	-	-	-	-
2002	1	-	-	-	-	-	-	-	-	-	1	-	-
2001	5	1	-	-	-	-	-	-	2	-	1	1	-
2000	-	-	-	-	-	-	-	-	-	-	-	-	-
1999	1	-	-	-	-	-	-	1	-	-	-	-	-
Hispanic, female													
2005	-	-	-	-	-	-	-	-	-	-	-	-	-
2004	-	-	-	-	-	-	-	-	-	-	-	-	-
2003	-	-	-	-	-	-	-	-	-	-	-	-	-
2002	-	-	-	-	-	-	-	-	-	-	-	-	-
2001	2	-	-	-	-	-	-	-	-	-	2	-	-
2000	1	-	-	-	-	-	-	-	-	-	1	-	-
1999	4	2	-	-	-	-	-	1	-	-	1	-	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Syphilis (A50-A53)													
Non-Hispanic 1/, both sexes													
2005	46	-	-	1	-	4	5	5	8	4	13	6	-
2004	42	-	-	-	-	3	3	3	4	6	11	12	-
2003	32	-	-	-	-	-	3	4	3	5	8	9	-
2002	39	1	1	-	-	-	2	5	5	6	11	8	-
2001	28	-	-	1	-	1	3	3	1	5	10	4	-
2000	39	1	-	-	-	1	-	6	3	7	11	10	-
1999	27	-	-	-	-	1	1	2	4	6	9	4	-
Non-Hispanic 1/, male													
2005	31	-	-	-	-	4	4	4	5	4	8	2	-
2004	20	-	-	-	-	2	2	1	2	4	5	4	-
2003	21	-	-	-	-	-	3	4	2	5	4	3	-
2002	20	-	1	-	-	-	2	4	2	5	3	3	-
2001	14	-	-	1	-	-	2	3	1	3	4	-	-
2000	24	1	-	-	-	1	-	5	3	2	7	5	-
1999	11	-	-	-	-	1	-	1	3	5	-	1	-
Non-Hispanic 1/, female													
2005	15	-	-	1	-	-	1	1	3	-	5	4	-
2004	22	-	-	-	-	1	1	2	2	2	6	8	-
2003	11	-	-	-	-	-	-	-	1	-	4	6	-
2002	19	1	-	-	-	-	-	1	3	1	8	5	-
2001	14	-	-	-	-	1	1	-	-	2	6	4	-
2000	15	-	-	-	-	-	-	1	-	5	4	5	-
1999	16	-	-	-	-	-	1	1	1	1	9	3	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Syphilis (A50-A53)													
Non-Hispanic white, both sexes													
2005 2/	15	-	-	-	-	-	1	-	3	1	8	2	-
2004 2/	16	-	-	-	-	-	3	1	1	3	5	3	-
2003 2/	16	-	-	-	-	-	1	-	2	2	5	6	-
2002	16	-	-	-	-	-	1	2	1	3	4	5	-
2001	14	-	-	-	-	-	1	1	1	2	7	2	-
2000	20	-	-	-	-	1	-	2	-	3	6	8	-
1999	14	-	-	-	-	1	1	2	3	3	3	1	-
Non-Hispanic white, male													
2005 2/	9	-	-	-	-	-	1	-	2	1	5	-	-
2004 2/	6	-	-	-	-	-	2	-	1	1	1	1	-
2003 2/	9	-	-	-	-	-	1	-	1	2	3	2	-
2002	8	-	-	-	-	-	1	2	-	3	-	2	-
2001	5	-	-	-	-	-	-	1	1	1	2	-	-
2000	7	-	-	-	-	1	-	1	-	-	2	3	-
1999	6	-	-	-	-	1	-	1	2	2	-	-	-
Non-Hispanic white, female													
2005 2/	6	-	-	-	-	-	-	-	1	-	3	2	-
2004 2/	10	-	-	-	-	-	1	1	-	2	4	2	-
2003 2/	7	-	-	-	-	-	-	-	1	-	2	4	-
2002	8	-	-	-	-	-	-	-	1	-	4	3	-
2001	9	-	-	-	-	-	1	-	-	1	5	2	-
2000	13	-	-	-	-	-	-	1	-	3	4	5	-
1999	8	-	-	-	-	-	1	1	1	1	3	1	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Syphilis (A50-A53)													
Non-Hispanic black, both sexes													
2005 2/	30	-	-	1	-	4	4	5	5	2	5	4	-
2004 2/	23	-	-	-	-	3	-	1	3	3	5	8	-
2003 2/	15	-	-	-	-	-	2	4	1	3	3	2	-
2002	23	1	1	-	-	-	1	3	4	3	7	3	-
2001	14	-	-	1	-	1	2	2	-	3	3	2	-
2000	19	1	-	-	-	-	-	4	3	4	5	2	-
1999	12	-	-	-	-	-	-	-	1	3	5	3	-
Non-Hispanic black, male													
2005 2/	21	-	-	-	-	4	3	4	3	2	3	2	-
2004 2/	13	-	-	-	-	2	-	1	1	3	3	3	-
2003 2/	11	-	-	-	-	-	2	4	1	3	1	-	-
2002	12	-	1	-	-	-	1	2	2	2	3	1	-
2001	9	-	-	1	-	-	2	2	-	2	2	-	-
2000	17	1	-	-	-	-	-	4	3	2	5	2	-
1999	5	-	-	-	-	-	-	-	1	3	-	1	-
Non-Hispanic black, female													
2005 2/	9	-	-	1	-	-	1	1	2	-	2	2	-
2004 2/	10	-	-	-	-	1	-	-	2	-	2	5	-
2003 2/	4	-	-	-	-	-	-	-	-	-	2	2	-
2002	11	1	-	-	-	-	-	1	2	1	4	2	-
2001	5	-	-	-	-	1	-	-	-	1	1	2	-
2000	2	-	-	-	-	-	-	-	-	2	-	-	-
1999	7	-	-	-	-	-	-	-	-	-	5	2	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Syphilis (A50-A53)													
Origin not stated 3/, both sexes													
2005	-	-	-	-	-	-	-	-	-	-	-	-	-
2004	-	-	-	-	-	-	-	-	-	-	-	-	-
2003	1	-	-	-	-	-	-	-	-	1	-	-	-
2002	1	-	-	-	-	-	-	-	-	-	-	1	-
2001	1	-	-	-	-	-	-	-	1	-	-	-	-
2000	1	-	-	-	-	-	-	-	-	-	1	-	-
1999	1	-	-	-	-	-	-	1	-	-	-	-	-
Origin not stated 3/, male													
2005	-	-	-	-	-	-	-	-	-	-	-	-	-
2004	-	-	-	-	-	-	-	-	-	-	-	-	-
2003	-	-	-	-	-	-	-	-	-	-	-	-	-
2002	-	-	-	-	-	-	-	-	-	-	-	-	-
2001	-	-	-	-	-	-	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-	-	-	-
1999	1	-	-	-	-	-	-	1	-	-	-	-	-
Origin not stated 3/, female													
2005	-	-	-	-	-	-	-	-	-	-	-	-	-
2004	-	-	-	-	-	-	-	-	-	-	-	-	-
2003	1	-	-	-	-	-	-	-	-	1	-	-	-
2002	1	-	-	-	-	-	-	-	-	-	-	1	-
2001	1	-	-	-	-	-	-	-	1	-	-	-	-
2000	1	-	-	-	-	-	-	-	-	-	1	-	-
1999	-	-	-	-	-	-	-	-	-	-	-	-	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Acute poliomyelitis (A80)													
All origins, both sexes													
2005	-	-	-	-	-	-	-	-	-	-	-	-	-
2004	-	-	-	-	-	-	-	-	-	-	-	-	-
2003	-	-	-	-	-	-	-	-	-	-	-	-	-
2002	-	-	-	-	-	-	-	-	-	-	-	-	-
2001	-	-	-	-	-	-	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-	-	-	-
All origins, male													
2005	-	-	-	-	-	-	-	-	-	-	-	-	-
2004	-	-	-	-	-	-	-	-	-	-	-	-	-
2003	-	-	-	-	-	-	-	-	-	-	-	-	-
2002	-	-	-	-	-	-	-	-	-	-	-	-	-
2001	-	-	-	-	-	-	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-	-	-	-
All origins, female													
2005	-	-	-	-	-	-	-	-	-	-	-	-	-
2004	-	-	-	-	-	-	-	-	-	-	-	-	-
2003	-	-	-	-	-	-	-	-	-	-	-	-	-
2002	-	-	-	-	-	-	-	-	-	-	-	-	-
2001	-	-	-	-	-	-	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-	-	-	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Acute poliomyelitis (A80)													
Hispanic, both sexes													
2005	-	-	-	-	-	-	-	-	-	-	-	-	-
2004	-	-	-	-	-	-	-	-	-	-	-	-	-
2003	-	-	-	-	-	-	-	-	-	-	-	-	-
2002	-	-	-	-	-	-	-	-	-	-	-	-	-
2001	-	-	-	-	-	-	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-	-	-	-
Hispanic, male													
2005	-	-	-	-	-	-	-	-	-	-	-	-	-
2004	-	-	-	-	-	-	-	-	-	-	-	-	-
2003	-	-	-	-	-	-	-	-	-	-	-	-	-
2002	-	-	-	-	-	-	-	-	-	-	-	-	-
2001	-	-	-	-	-	-	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-	-	-	-
Hispanic, female													
2005	-	-	-	-	-	-	-	-	-	-	-	-	-
2004	-	-	-	-	-	-	-	-	-	-	-	-	-
2003	-	-	-	-	-	-	-	-	-	-	-	-	-
2002	-	-	-	-	-	-	-	-	-	-	-	-	-
2001	-	-	-	-	-	-	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-	-	-	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Acute poliomyelitis (A80)													
Non-Hispanic 1/, both sexes													
2005	-	-	-	-	-	-	-	-	-	-	-	-	-
2004	-	-	-	-	-	-	-	-	-	-	-	-	-
2003	-	-	-	-	-	-	-	-	-	-	-	-	-
2002	-	-	-	-	-	-	-	-	-	-	-	-	-
2001	-	-	-	-	-	-	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-	-	-	-
Non-Hispanic 1/, male													
2005	-	-	-	-	-	-	-	-	-	-	-	-	-
2004	-	-	-	-	-	-	-	-	-	-	-	-	-
2003	-	-	-	-	-	-	-	-	-	-	-	-	-
2002	-	-	-	-	-	-	-	-	-	-	-	-	-
2001	-	-	-	-	-	-	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-	-	-	-
Non-Hispanic 1/, female													
2005	-	-	-	-	-	-	-	-	-	-	-	-	-
2004	-	-	-	-	-	-	-	-	-	-	-	-	-
2003	-	-	-	-	-	-	-	-	-	-	-	-	-
2002	-	-	-	-	-	-	-	-	-	-	-	-	-
2001	-	-	-	-	-	-	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-	-	-	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Acute poliomyelitis (A80)													
Non-Hispanic white, both sexes													
2005 2/	-	-	-	-	-	-	-	-	-	-	-	-	-
2004 2/	-	-	-	-	-	-	-	-	-	-	-	-	-
2003 2/	-	-	-	-	-	-	-	-	-	-	-	-	-
2002	-	-	-	-	-	-	-	-	-	-	-	-	-
2001	-	-	-	-	-	-	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-	-	-	-
Non-Hispanic white, male													
2005 2/	-	-	-	-	-	-	-	-	-	-	-	-	-
2004 2/	-	-	-	-	-	-	-	-	-	-	-	-	-
2003 2/	-	-	-	-	-	-	-	-	-	-	-	-	-
2002	-	-	-	-	-	-	-	-	-	-	-	-	-
2001	-	-	-	-	-	-	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-	-	-	-
Non-Hispanic white, female													
2005 2/	-	-	-	-	-	-	-	-	-	-	-	-	-
2004 2/	-	-	-	-	-	-	-	-	-	-	-	-	-
2003 2/	-	-	-	-	-	-	-	-	-	-	-	-	-
2002	-	-	-	-	-	-	-	-	-	-	-	-	-
2001	-	-	-	-	-	-	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-	-	-	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Acute poliomyelitis (A80)													
Non-Hispanic black, both sexes													
2005 2/	-	-	-	-	-	-	-	-	-	-	-	-	-
2004 2/	-	-	-	-	-	-	-	-	-	-	-	-	-
2003 2/	-	-	-	-	-	-	-	-	-	-	-	-	-
2002	-	-	-	-	-	-	-	-	-	-	-	-	-
2001	-	-	-	-	-	-	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-	-	-	-
Non-Hispanic black, male													
2005 2/	-	-	-	-	-	-	-	-	-	-	-	-	-
2004 2/	-	-	-	-	-	-	-	-	-	-	-	-	-
2003 2/	-	-	-	-	-	-	-	-	-	-	-	-	-
2002	-	-	-	-	-	-	-	-	-	-	-	-	-
2001	-	-	-	-	-	-	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-	-	-	-
Non-Hispanic black, female													
2005 2/	-	-	-	-	-	-	-	-	-	-	-	-	-
2004 2/	-	-	-	-	-	-	-	-	-	-	-	-	-
2003 2/	-	-	-	-	-	-	-	-	-	-	-	-	-
2002	-	-	-	-	-	-	-	-	-	-	-	-	-
2001	-	-	-	-	-	-	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-	-	-	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Acute poliomyelitis (A80)													
Origin not stated 3/, both sexes													
2005	-	-	-	-	-	-	-	-	-	-	-	-	-
2004	-	-	-	-	-	-	-	-	-	-	-	-	-
2003	-	-	-	-	-	-	-	-	-	-	-	-	-
2002	-	-	-	-	-	-	-	-	-	-	-	-	-
2001	-	-	-	-	-	-	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-	-	-	-
Origin not stated 3/, male													
2005	-	-	-	-	-	-	-	-	-	-	-	-	-
2004	-	-	-	-	-	-	-	-	-	-	-	-	-
2003	-	-	-	-	-	-	-	-	-	-	-	-	-
2002	-	-	-	-	-	-	-	-	-	-	-	-	-
2001	-	-	-	-	-	-	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-	-	-	-
Origin not stated 3/, female													
2005	-	-	-	-	-	-	-	-	-	-	-	-	-
2004	-	-	-	-	-	-	-	-	-	-	-	-	-
2003	-	-	-	-	-	-	-	-	-	-	-	-	-
2002	-	-	-	-	-	-	-	-	-	-	-	-	-
2001	-	-	-	-	-	-	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-	-	-	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Arthropod-borne viral encephalitis (A83-A84,A85.2)													
All origins, both sexes													
2005	6	-	1	1	-	1	-	-	2	-	1	-	-
2004	5	-	-	-	-	-	1	-	1	3	-	-	-
2003	13	1	-	1	-	-	-	1	3	2	5	-	-
2002	9	-	-	-	1	1	-	2	1	1	2	1	-
2001	9	1	-	-	-	-	-	1	-	5	2	-	-
2000	2	-	-	-	-	-	1	-	-	-	1	-	-
1999	3	-	-	-	-	-	-	-	-	2	-	1	-
All origins, male													
2005	4	-	-	1	-	1	-	-	1	-	1	-	-
2004	5	-	-	-	-	-	1	-	1	3	-	-	-
2003	9	1	-	1	-	-	-	1	2	2	2	-	-
2002	4	-	-	-	-	-	-	1	1	1	1	-	-
2001	6	-	-	-	-	-	-	-	-	4	2	-	-
2000	2	-	-	-	-	-	1	-	-	-	1	-	-
1999	3	-	-	-	-	-	-	-	-	2	-	1	-
All origins, female													
2005	2	-	1	-	-	-	-	-	1	-	-	-	-
2004	-	-	-	-	-	-	-	-	-	-	-	-	-
2003	4	-	-	-	-	-	-	-	1	-	3	-	-
2002	5	-	-	-	1	1	-	1	-	-	1	1	-
2001	3	1	-	-	-	-	-	1	-	1	-	-	-
2000	-	-	-	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-	-	-	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Arthropod-borne viral encephalitis (A83-A84,A85.2)													
Hispanic, both sexes													
2005	1	-	-	-	-	1	-	-	-	-	-	-	-
2004	-	-	-	-	-	-	-	-	-	-	-	-	-
2003	-	-	-	-	-	-	-	-	-	-	-	-	-
2002	-	-	-	-	-	-	-	-	-	-	-	-	-
2001	-	-	-	-	-	-	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-	-	-	-
Hispanic, male													
2005	1	-	-	-	-	1	-	-	-	-	-	-	-
2004	-	-	-	-	-	-	-	-	-	-	-	-	-
2003	-	-	-	-	-	-	-	-	-	-	-	-	-
2002	-	-	-	-	-	-	-	-	-	-	-	-	-
2001	-	-	-	-	-	-	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-	-	-	-
Hispanic, female													
2005	-	-	-	-	-	-	-	-	-	-	-	-	-
2004	-	-	-	-	-	-	-	-	-	-	-	-	-
2003	-	-	-	-	-	-	-	-	-	-	-	-	-
2002	-	-	-	-	-	-	-	-	-	-	-	-	-
2001	-	-	-	-	-	-	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-	-	-	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Arthropod-borne viral encephalitis (A83-A84,A85.2)													
Non-Hispanic 1/, both sexes													
2005	5	-	1	1	-	-	-	-	2	-	1	-	-
2004	5	-	-	-	-	-	1	-	1	3	-	-	-
2003	13	1	-	1	-	-	-	1	3	2	5	-	-
2002	9	-	-	-	1	1	-	2	1	1	2	1	-
2001	9	1	-	-	-	-	-	1	-	5	2	-	-
2000	2	-	-	-	-	-	1	-	-	-	1	-	-
1999	3	-	-	-	-	-	-	-	-	2	-	1	-
Non-Hispanic 1/, male													
2005	3	-	-	1	-	-	-	-	1	-	1	-	-
2004	5	-	-	-	-	-	1	-	1	3	-	-	-
2003	9	1	-	1	-	-	-	1	2	2	2	-	-
2002	4	-	-	-	-	-	-	1	1	1	1	-	-
2001	6	-	-	-	-	-	-	-	-	4	2	-	-
2000	2	-	-	-	-	-	1	-	-	-	1	-	-
1999	3	-	-	-	-	-	-	-	-	2	-	1	-
Non-Hispanic 1/, female													
2005	2	-	1	-	-	-	-	-	1	-	-	-	-
2004	-	-	-	-	-	-	-	-	-	-	-	-	-
2003	4	-	-	-	-	-	-	-	1	-	3	-	-
2002	5	-	-	-	1	1	-	1	-	-	1	1	-
2001	3	1	-	-	-	-	-	1	-	1	-	-	-
2000	-	-	-	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-	-	-	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Arthropod-borne viral encephalitis (A83-A84,A85.2)													
Non-Hispanic white, both sexes													
2005 2/	5	-	1	1	-	-	-	-	2	-	1	-	-
2004 2/	3	-	-	-	-	-	-	-	1	2	-	-	-
2003 2/	10	-	-	-	-	-	-	-	3	2	5	-	-
2002	6	-	-	-	1	1	-	1	-	-	2	1	-
2001	6	-	-	-	-	-	-	1	-	4	1	-	-
2000	2	-	-	-	-	-	1	-	-	-	1	-	-
1999	3	-	-	-	-	-	-	-	-	2	-	1	-
Non-Hispanic white, male													
2005 2/	3	-	-	1	-	-	-	-	1	-	1	-	-
2004 2/	3	-	-	-	-	-	-	-	1	2	-	-	-
2003 2/	6	-	-	-	-	-	-	-	2	2	2	-	-
2002	2	-	-	-	-	-	-	1	-	-	1	-	-
2001	5	-	-	-	-	-	-	-	-	4	1	-	-
2000	2	-	-	-	-	-	1	-	-	-	1	-	-
1999	3	-	-	-	-	-	-	-	-	2	-	1	-
Non-Hispanic white, female													
2005 2/	2	-	1	-	-	-	-	-	1	-	-	-	-
2004 2/	-	-	-	-	-	-	-	-	-	-	-	-	-
2003 2/	4	-	-	-	-	-	-	-	1	-	3	-	-
2002	4	-	-	-	1	1	-	-	-	-	1	1	-
2001	1	-	-	-	-	-	-	1	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-	-	-	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Arthropod-borne viral encephalitis (A83-A84,A85.2)													
Non-Hispanic black, both sexes													
2005 2/	-	-	-	-	-	-	-	-	-	-	-	-	-
2004 2/	2	-	-	-	-	-	1	-	-	1	-	-	-
2003 2/	1	-	-	-	-	-	-	1	-	-	-	-	-
2002	3	-	-	-	-	-	-	1	1	1	-	-	-
2001	2	1	-	-	-	-	-	-	-	1	-	-	-
2000	-	-	-	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-	-	-	-
Non-Hispanic black, male													
2005 2/	-	-	-	-	-	-	-	-	-	-	-	-	-
2004 2/	2	-	-	-	-	-	1	-	-	1	-	-	-
2003 2/	1	-	-	-	-	-	-	1	-	-	-	-	-
2002	2	-	-	-	-	-	-	-	1	1	-	-	-
2001	-	-	-	-	-	-	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-	-	-	-
Non-Hispanic black, female													
2005 2/	-	-	-	-	-	-	-	-	-	-	-	-	-
2004 2/	-	-	-	-	-	-	-	-	-	-	-	-	-
2003 2/	-	-	-	-	-	-	-	-	-	-	-	-	-
2002	1	-	-	-	-	-	-	1	-	-	-	-	-
2001	2	1	-	-	-	-	-	-	-	1	-	-	-
2000	-	-	-	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-	-	-	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Arthropod-borne viral encephalitis (A83-A84,A85.2)													
Origin not stated 3/, both sexes													
2005	-	-	-	-	-	-	-	-	-	-	-	-	-
2004	-	-	-	-	-	-	-	-	-	-	-	-	-
2003	-	-	-	-	-	-	-	-	-	-	-	-	-
2002	-	-	-	-	-	-	-	-	-	-	-	-	-
2001	-	-	-	-	-	-	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-	-	-	-
Origin not stated 3/, male													
2005	-	-	-	-	-	-	-	-	-	-	-	-	-
2004	-	-	-	-	-	-	-	-	-	-	-	-	-
2003	-	-	-	-	-	-	-	-	-	-	-	-	-
2002	-	-	-	-	-	-	-	-	-	-	-	-	-
2001	-	-	-	-	-	-	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-	-	-	-
Origin not stated 3/, female													
2005	-	-	-	-	-	-	-	-	-	-	-	-	-
2004	-	-	-	-	-	-	-	-	-	-	-	-	-
2003	-	-	-	-	-	-	-	-	-	-	-	-	-
2002	-	-	-	-	-	-	-	-	-	-	-	-	-
2001	-	-	-	-	-	-	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-	-	-	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Measles (B05)													
All origins, both sexes													
2005	1	-	1	-	-	-	-	-	-	-	-	-	-
2004	-	-	-	-	-	-	-	-	-	-	-	-	-
2003	1	-	1	-	-	-	-	-	-	-	-	-	-
2002	-	-	-	-	-	-	-	-	-	-	-	-	-
2001	1	-	-	-	-	-	1	-	-	-	-	-	-
2000	1	1	-	-	-	-	-	-	-	-	-	-	-
1999	2	1	-	-	-	-	-	-	1	-	-	-	-
All origins, male													
2005	1	-	1	-	-	-	-	-	-	-	-	-	-
2004	-	-	-	-	-	-	-	-	-	-	-	-	-
2003	1	-	1	-	-	-	-	-	-	-	-	-	-
2002	-	-	-	-	-	-	-	-	-	-	-	-	-
2001	1	-	-	-	-	-	1	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-	-	-	-
All origins, female													
2005	-	-	-	-	-	-	-	-	-	-	-	-	-
2004	-	-	-	-	-	-	-	-	-	-	-	-	-
2003	-	-	-	-	-	-	-	-	-	-	-	-	-
2002	-	-	-	-	-	-	-	-	-	-	-	-	-
2001	-	-	-	-	-	-	-	-	-	-	-	-	-
2000	1	1	-	-	-	-	-	-	-	-	-	-	-
1999	2	1	-	-	-	-	-	-	1	-	-	-	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Measles (B05)													
Hispanic, both sexes													
2005	-	-	-	-	-	-	-	-	-	-	-	-	-
2004	-	-	-	-	-	-	-	-	-	-	-	-	-
2003	-	-	-	-	-	-	-	-	-	-	-	-	-
2002	-	-	-	-	-	-	-	-	-	-	-	-	-
2001	-	-	-	-	-	-	-	-	-	-	-	-	-
2000	1	1	-	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-	-	-	-
Hispanic, male													
2005	-	-	-	-	-	-	-	-	-	-	-	-	-
2004	-	-	-	-	-	-	-	-	-	-	-	-	-
2003	-	-	-	-	-	-	-	-	-	-	-	-	-
2002	-	-	-	-	-	-	-	-	-	-	-	-	-
2001	-	-	-	-	-	-	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-	-	-	-
Hispanic, female													
2005	-	-	-	-	-	-	-	-	-	-	-	-	-
2004	-	-	-	-	-	-	-	-	-	-	-	-	-
2003	-	-	-	-	-	-	-	-	-	-	-	-	-
2002	-	-	-	-	-	-	-	-	-	-	-	-	-
2001	-	-	-	-	-	-	-	-	-	-	-	-	-
2000	1	1	-	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-	-	-	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Measles (B05)													
Non-Hispanic 1/, both sexes													
2005	1	-	1	-	-	-	-	-	-	-	-	-	-
2004	-	-	-	-	-	-	-	-	-	-	-	-	-
2003	1	-	1	-	-	-	-	-	-	-	-	-	-
2002	-	-	-	-	-	-	-	-	-	-	-	-	-
2001	1	-	-	-	-	-	1	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-	-	-	-
1999	2	1	-	-	-	-	-	-	1	-	-	-	-
Non-Hispanic 1/, male													
2005	1	-	1	-	-	-	-	-	-	-	-	-	-
2004	-	-	-	-	-	-	-	-	-	-	-	-	-
2003	1	-	1	-	-	-	-	-	-	-	-	-	-
2002	-	-	-	-	-	-	-	-	-	-	-	-	-
2001	1	-	-	-	-	-	1	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-	-	-	-
Non-Hispanic 1/, female													
2005	-	-	-	-	-	-	-	-	-	-	-	-	-
2004	-	-	-	-	-	-	-	-	-	-	-	-	-
2003	-	-	-	-	-	-	-	-	-	-	-	-	-
2002	-	-	-	-	-	-	-	-	-	-	-	-	-
2001	-	-	-	-	-	-	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-	-	-	-
1999	2	1	-	-	-	-	-	-	1	-	-	-	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Measles (B05)													
Non-Hispanic white, both sexes													
2005 2/	1	-	1	-	-	-	-	-	-	-	-	-	-
2004 2/	-	-	-	-	-	-	-	-	-	-	-	-	-
2003 2/	1	-	1	-	-	-	-	-	-	-	-	-	-
2002	-	-	-	-	-	-	-	-	-	-	-	-	-
2001	1	-	-	-	-	-	1	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-	-	-	-
1999	1	-	-	-	-	-	-	-	1	-	-	-	-
Non-Hispanic white, male													
2005 2/	1	-	1	-	-	-	-	-	-	-	-	-	-
2004 2/	-	-	-	-	-	-	-	-	-	-	-	-	-
2003 2/	1	-	1	-	-	-	-	-	-	-	-	-	-
2002	-	-	-	-	-	-	-	-	-	-	-	-	-
2001	1	-	-	-	-	-	1	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-	-	-	-
Non-Hispanic white, female													
2005 2/	-	-	-	-	-	-	-	-	-	-	-	-	-
2004 2/	-	-	-	-	-	-	-	-	-	-	-	-	-
2003 2/	-	-	-	-	-	-	-	-	-	-	-	-	-
2002	-	-	-	-	-	-	-	-	-	-	-	-	-
2001	-	-	-	-	-	-	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-	-	-	-
1999	1	-	-	-	-	-	-	-	1	-	-	-	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Measles (B05)													
Non-Hispanic black, both sexes													
2005 2/	-	-	-	-	-	-	-	-	-	-	-	-	-
2004 2/	-	-	-	-	-	-	-	-	-	-	-	-	-
2003 2/	-	-	-	-	-	-	-	-	-	-	-	-	-
2002	-	-	-	-	-	-	-	-	-	-	-	-	-
2001	-	-	-	-	-	-	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-	-	-	-
1999	1	1	-	-	-	-	-	-	-	-	-	-	-
Non-Hispanic black, male													
2005 2/	-	-	-	-	-	-	-	-	-	-	-	-	-
2004 2/	-	-	-	-	-	-	-	-	-	-	-	-	-
2003 2/	-	-	-	-	-	-	-	-	-	-	-	-	-
2002	-	-	-	-	-	-	-	-	-	-	-	-	-
2001	-	-	-	-	-	-	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-	-	-	-
Non-Hispanic black, female													
2005 2/	-	-	-	-	-	-	-	-	-	-	-	-	-
2004 2/	-	-	-	-	-	-	-	-	-	-	-	-	-
2003 2/	-	-	-	-	-	-	-	-	-	-	-	-	-
2002	-	-	-	-	-	-	-	-	-	-	-	-	-
2001	-	-	-	-	-	-	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-	-	-	-
1999	1	1	-	-	-	-	-	-	-	-	-	-	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Measles (B05)													
Origin not stated 3/, both sexes													
2005	-	-	-	-	-	-	-	-	-	-	-	-	-
2004	-	-	-	-	-	-	-	-	-	-	-	-	-
2003	-	-	-	-	-	-	-	-	-	-	-	-	-
2002	-	-	-	-	-	-	-	-	-	-	-	-	-
2001	-	-	-	-	-	-	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-	-	-	-
Origin not stated 3/, male													
2005	-	-	-	-	-	-	-	-	-	-	-	-	-
2004	-	-	-	-	-	-	-	-	-	-	-	-	-
2003	-	-	-	-	-	-	-	-	-	-	-	-	-
2002	-	-	-	-	-	-	-	-	-	-	-	-	-
2001	-	-	-	-	-	-	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-	-	-	-
Origin not stated 3/, female													
2005	-	-	-	-	-	-	-	-	-	-	-	-	-
2004	-	-	-	-	-	-	-	-	-	-	-	-	-
2003	-	-	-	-	-	-	-	-	-	-	-	-	-
2002	-	-	-	-	-	-	-	-	-	-	-	-	-
2001	-	-	-	-	-	-	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-	-	-	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Viral hepatitis (B15-B19)													
All origins, both sexes													
2005	5,529	-	-	1	14	61	509	2,314	1,377	648	482	123	-
2004	5,382	1	-	2	11	61	610	2,226	1,244	634	464	129	-
2003	5,431	1	-	-	14	64	652	2,259	1,085	698	543	115	-
2002	5,793	2	3	2	23	94	740	2,331	1,072	762	587	177	-
2001	5,585	1	-	3	20	74	797	2,060	1,014	836	613	166	1
2000	5,357	6	1	2	18	79	779	1,894	904	845	660	168	1
1999	4,853	2	3	3	14	88	722	1,506	786	881	651	197	-
All origins, male													
2005	3,612	-	-	1	3	34	349	1,697	977	317	193	41	-
2004	3,513	-	-	1	7	33	399	1,645	860	334	189	45	-
2003	3,546	-	-	-	7	42	461	1,666	729	352	249	40	-
2002	3,770	1	3	1	13	63	519	1,755	702	373	270	70	-
2001	3,539	1	-	2	11	42	553	1,570	643	396	263	57	1
2000	3,381	2	1	1	11	46	540	1,425	577	396	311	70	1
1999	3,043	2	2	2	10	53	520	1,138	491	426	317	82	-
All origins, female													
2005	1,917	-	-	-	11	27	160	617	400	331	289	82	-
2004	1,869	1	-	1	4	28	211	581	384	300	275	84	-
2003	1,885	1	-	-	7	22	191	593	356	346	294	75	-
2002	2,023	1	-	1	10	31	221	576	370	389	317	107	-
2001	2,046	-	-	1	9	32	244	490	371	440	350	109	-
2000	1,976	4	-	1	7	33	239	469	327	449	349	98	-
1999	1,810	-	1	1	4	35	202	368	295	455	334	115	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Viral hepatitis (B15-B19)													
Hispanic, both sexes													
2005	795	-	-	-	-	10	114	293	202	113	48	15	-
2004	716	-	-	1	2	9	110	275	171	89	51	8	-
2003	710	-	-	-	2	10	107	266	151	106	59	9	-
2002	758	-	-	-	5	17	101	291	164	102	63	15	-
2001	791	-	-	2	6	10	126	266	192	113	61	15	-
2000	708	3	-	-	3	14	126	218	147	120	66	11	-
1999	656	-	1	-	3	16	104	203	127	129	65	8	-
Hispanic, male													
2005	521	-	-	-	-	7	85	222	130	53	19	5	-
2004	470	-	-	1	1	6	85	206	110	45	14	2	-
2003	465	-	-	-	2	8	80	207	103	42	20	3	-
2002	491	-	-	-	5	14	75	218	108	40	25	6	-
2001	487	-	-	2	2	3	89	206	123	35	23	4	-
2000	456	1	-	-	2	11	100	165	94	52	25	6	-
1999	406	-	1	-	1	11	84	153	75	59	18	4	-
Hispanic, female													
2005	274	-	-	-	-	3	29	71	72	60	29	10	-
2004	246	-	-	-	1	3	25	69	61	44	37	6	-
2003	245	-	-	-	-	2	27	59	48	64	39	6	-
2002	267	-	-	-	-	3	26	73	56	62	38	9	-
2001	304	-	-	-	4	7	37	60	69	78	38	11	-
2000	252	2	-	-	1	3	26	53	53	68	41	5	-
1999	250	-	-	-	2	5	20	50	52	70	47	4	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Viral hepatitis (B15-B19)													
Non-Hispanic 1/, both sexes													
2005	4,714	-	-	1	14	51	391	2,011	1,171	534	433	108	-
2004	4,647	1	-	1	9	52	497	1,943	1,071	542	410	121	-
2003	4,704	1	-	-	12	54	545	1,984	931	587	484	106	-
2002	5,004	2	3	2	18	77	634	2,030	898	655	523	162	-
2001	4,776	1	-	1	14	64	669	1,785	817	721	552	151	1
2000	4,629	3	1	2	15	65	652	1,665	751	724	594	156	1
1999	4,176	2	2	3	11	71	613	1,294	656	750	585	189	-
Non-Hispanic 1/, male													
2005	3,076	-	-	1	3	27	261	1,467	843	264	174	36	-
2004	3,031	-	-	-	6	27	311	1,434	749	287	174	43	-
2003	3,068	-	-	-	5	34	381	1,451	625	306	229	37	-
2002	3,256	1	3	1	8	49	439	1,529	586	331	245	64	-
2001	3,038	1	-	-	9	39	462	1,357	515	361	240	53	1
2000	2,909	1	1	1	9	35	440	1,250	479	343	286	63	1
1999	2,620	2	1	2	9	41	432	978	414	365	298	78	-
Non-Hispanic 1/, female													
2005	1,638	-	-	-	11	24	130	544	328	270	259	72	-
2004	1,616	1	-	1	3	25	186	509	322	255	236	78	-
2003	1,636	1	-	-	7	20	164	533	306	281	255	69	-
2002	1,748	1	-	1	10	28	195	501	312	324	278	98	-
2001	1,738	-	-	1	5	25	207	428	302	360	312	98	-
2000	1,720	2	-	1	6	30	212	415	272	381	308	93	-
1999	1,556	-	1	1	2	30	181	316	242	385	287	111	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Viral hepatitis (B15-B19)													
Non-Hispanic white, both sexes													
2005 2/	3,603	-	-	-	9	31	304	1,602	858	360	351	88	-
2004 2/	3,533	1	-	-	3	31	390	1,517	766	401	322	102	-
2003 2/	3,585	-	-	-	7	39	425	1,566	642	427	390	89	-
2002	3,857	1	-	2	15	49	496	1,590	635	489	440	140	-
2001	3,663	1	-	1	7	43	522	1,364	586	544	467	128	-
2000	3,513	1	-	2	5	37	501	1,243	520	559	506	138	1
1999	3,176	1	2	2	6	51	464	958	465	567	497	163	-
Non-Hispanic white, male													
2005 2/	2,413	-	-	-	2	17	208	1,187	630	189	150	30	-
2004 2/	2,330	-	-	-	2	17	251	1,132	536	216	141	35	-
2003 2/	2,381	-	-	-	1	22	300	1,162	445	232	186	33	-
2002	2,551	1	-	1	6	33	351	1,212	415	263	214	55	-
2001	2,410	1	-	-	4	28	368	1,062	384	296	218	49	-
2000	2,234	-	-	1	4	19	342	958	339	265	248	57	1
1999	2,026	1	1	1	6	29	333	717	308	295	268	67	-
Non-Hispanic white, female													
2005 2/	1,190	-	-	-	7	14	96	415	228	171	201	58	-
2004 2/	1,203	1	-	-	1	14	139	385	230	185	181	67	-
2003 2/	1,204	-	-	-	6	17	125	404	197	195	204	56	-
2002	1,306	-	-	1	9	16	145	378	220	226	226	85	-
2001	1,253	-	-	1	3	15	154	302	202	248	249	79	-
2000	1,279	1	-	1	1	18	159	285	181	294	258	81	-
1999	1,150	-	1	1	-	22	131	241	157	272	229	96	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Viral hepatitis (B15-B19)													
Non-Hispanic black, both sexes													
2005 2/	856	-	-	1	3	17	64	349	253	106	50	13	-
2004 2/	846	-	-	1	4	14	79	362	247	78	49	12	-
2003 2/	842	-	-	-	4	11	97	352	221	89	57	11	-
2002	849	1	1	-	2	22	111	368	184	98	49	13	-
2001	819	-	-	-	6	15	121	349	167	91	53	16	1
2000	777	2	-	-	10	21	117	329	138	93	57	10	-
1999	718	1	-	1	4	17	119	277	118	109	53	19	-
Non-Hispanic black, male													
2005 2/	518	-	-	1	-	7	38	234	171	52	13	2	-
2004 2/	554	-	-	-	3	6	39	262	177	47	16	4	-
2003 2/	536	-	-	-	3	9	63	250	145	41	21	4	-
2002	532	-	1	-	1	11	67	262	122	48	17	3	-
2001	479	-	-	-	4	7	79	240	99	34	13	2	1
2000	490	1	-	-	5	11	77	234	86	48	25	3	-
1999	433	1	-	1	2	9	79	215	65	38	15	8	-
Non-Hispanic black, female													
2005 2/	338	-	-	-	3	10	26	115	82	54	37	11	-
2004 2/	292	-	-	1	1	8	40	100	70	31	33	8	-
2003 2/	306	-	-	-	1	2	34	102	76	48	36	7	-
2002	317	1	-	-	1	11	44	106	62	50	32	10	-
2001	340	-	-	-	2	8	42	109	68	57	40	14	-
2000	287	1	-	-	5	10	40	95	52	45	32	7	-
1999	285	-	-	-	2	8	40	62	53	71	38	11	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Viral hepatitis (B15-B19)													
Origin not stated 3/, both sexes													
2005	20	-	-	-	-	-	4	10	4	1	1	-	-
2004	19	-	-	-	-	-	3	8	2	3	3	-	-
2003	17	-	-	-	-	-	-	9	3	5	-	-	-
2002	31	-	-	-	-	-	5	10	10	5	1	-	-
2001	18	-	-	-	-	-	2	9	5	2	-	-	-
2000	20	-	-	-	-	-	1	11	6	1	-	1	-
1999	21	-	-	-	-	1	5	9	3	2	1	-	-
Origin not stated 3/, male													
2005	15	-	-	-	-	-	3	8	4	-	-	-	-
2004	12	-	-	-	-	-	3	5	1	2	1	-	-
2003	13	-	-	-	-	-	-	8	1	4	-	-	-
2002	23	-	-	-	-	-	5	8	8	2	-	-	-
2001	14	-	-	-	-	-	2	7	5	-	-	-	-
2000	16	-	-	-	-	-	-	10	4	1	-	1	-
1999	17	-	-	-	-	1	4	7	2	2	1	-	-
Origin not stated 3/, female													
2005	5	-	-	-	-	-	1	2	-	1	1	-	-
2004	7	-	-	-	-	-	-	3	1	1	2	-	-
2003	4	-	-	-	-	-	-	1	2	1	-	-	-
2002	8	-	-	-	-	-	-	2	2	3	1	-	-
2001	4	-	-	-	-	-	-	2	-	2	-	-	-
2000	4	-	-	-	-	-	1	1	2	-	-	-	-
1999	4	-	-	-	-	-	1	2	1	-	-	-	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Human immunodeficiency virus (HIV) disease (B20-B24)													
All origins, both sexes													
2005	12,543	2	2	13	159	1,318	4,363	4,516	1,612	438	104	15	1
2004	13,063	3	4	25	191	1,468	4,826	4,422	1,562	436	106	16	4
2003	13,658	5	2	36	178	1,588	5,340	4,442	1,517	438	95	14	3
2002	14,095	7	5	25	178	1,839	5,707	4,474	1,347	401	96	12	4
2001	14,175	10	13	38	225	2,101	5,867	4,120	1,317	384	88	10	2
2000	14,478	10	14	46	179	2,437	5,919	4,142	1,239	397	85	9	1
1999	14,802	14	24	66	198	2,729	6,232	3,907	1,149	398	70	14	1
All origins, male													
2005	9,189	-	1	7	89	809	3,146	3,417	1,292	348	73	6	1
2004	9,506	1	1	12	107	917	3,466	3,327	1,262	333	71	5	4
2003	10,119	2	2	19	95	1,038	3,884	3,438	1,215	336	75	12	3
2002	10,468	4	1	14	93	1,187	4,198	3,485	1,083	323	71	5	4
2001	10,590	7	8	23	105	1,421	4,360	3,241	1,051	292	73	7	2
2000	10,868	8	6	22	99	1,602	4,441	3,296	1,013	313	65	2	1
1999	11,234	9	16	36	95	1,914	4,706	3,134	948	317	50	8	1
All origins, female													
2005	3,354	2	1	6	70	509	1,217	1,099	320	90	31	9	-
2004	3,557	2	3	13	84	551	1,360	1,095	300	103	35	11	-
2003	3,539	3	-	17	83	550	1,456	1,004	302	102	20	2	-
2002	3,627	3	4	11	85	652	1,509	989	264	78	25	7	-
2001	3,585	3	5	15	120	680	1,507	879	266	92	15	3	-
2000	3,610	2	8	24	80	835	1,478	846	226	84	20	7	-
1999	3,568	5	8	30	103	815	1,526	773	201	81	20	6	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Human immunodeficiency virus (HIV) disease (B20-B24)													
Hispanic, both sexes													
2005	1,645	2	-	4	31	192	616	557	171	57	12	3	-
2004	1,758	1	1	4	25	211	676	569	186	64	18	-	3
2003	1,890	-	1	2	31	244	727	606	204	64	10	1	-
2002	1,855	4	-	3	29	262	779	537	188	40	9	1	3
2001	1,853	1	-	3	32	300	759	526	169	51	10	2	-
2000	1,908	-	-	9	23	365	762	531	163	41	14	-	-
1999	1,958	3	5	9	33	407	855	455	137	49	4	1	-
Hispanic, male													
2005	1,309	-	-	1	23	142	496	448	143	43	11	2	-
2004	1,371	1	1	1	19	169	522	445	152	45	13	-	3
2003	1,465	-	1	-	20	184	550	499	153	50	7	1	-
2002	1,440	3	-	3	19	201	607	421	140	36	7	-	3
2001	1,437	-	-	2	21	222	589	420	129	43	9	2	-
2000	1,493	-	-	5	18	274	603	426	123	34	10	-	-
1999	1,525	1	4	3	20	314	666	354	118	41	4	-	-
Hispanic, female													
2005	336	2	-	3	8	50	120	109	28	14	1	1	-
2004	387	-	-	3	6	42	154	124	34	19	5	-	-
2003	425	-	-	2	11	60	177	107	51	14	3	-	-
2002	415	1	-	-	10	61	172	116	48	4	2	1	-
2001	416	1	-	1	11	78	170	106	40	8	1	-	-
2000	415	-	-	4	5	91	159	105	40	7	4	-	-
1999	433	2	1	6	13	93	189	101	19	8	-	1	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Human immunodeficiency virus (HIV) disease (B20-B24)													
Non-Hispanic 1/, both sexes													
2005	10,813	-	2	9	128	1,118	3,715	3,927	1,433	377	91	12	1
2004	11,195	2	3	20	165	1,252	4,122	3,805	1,356	366	87	16	1
2003	11,629	5	1	34	147	1,333	4,561	3,785	1,298	366	84	13	2
2002	12,094	3	5	22	149	1,564	4,871	3,874	1,148	359	87	11	1
2001	12,141	9	13	35	193	1,779	5,044	3,533	1,123	325	77	8	2
2000	12,389	10	14	37	156	2,053	5,077	3,562	1,058	343	70	9	-
1999	12,638	11	19	56	165	2,301	5,305	3,377	987	339	65	13	-
Non-Hispanic 1/, male													
2005	7,811	-	1	6	66	663	2,622	2,944	1,142	301	61	4	1
2004	8,052	-	-	10	88	744	2,921	2,848	1,095	282	58	5	1
2003	8,550	2	1	19	75	846	3,299	2,900	1,050	278	67	11	2
2002	8,911	1	1	11	74	977	3,549	3,009	933	286	64	5	1
2001	9,011	7	8	21	84	1,187	3,725	2,766	902	241	63	5	2
2000	9,224	8	6	17	81	1,313	3,778	2,825	872	267	55	2	-
1999	9,538	8	12	33	75	1,584	3,984	2,715	806	268	45	8	-
Non-Hispanic 1/, female													
2005	3,002	-	1	3	62	455	1,093	983	291	76	30	8	-
2004	3,143	2	3	10	77	508	1,201	957	261	84	29	11	-
2003	3,079	3	-	15	72	487	1,262	885	248	88	17	2	-
2002	3,183	2	4	11	75	587	1,322	865	215	73	23	6	-
2001	3,130	2	5	14	109	592	1,319	767	221	84	14	3	-
2000	3,165	2	8	20	75	740	1,299	737	186	76	15	7	-
1999	3,100	3	7	23	90	717	1,321	662	181	71	20	5	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Human immunodeficiency virus (HIV) disease (B20-B24)													
Non-Hispanic white, both sexes													
2005 2/	3,755	-	2	-	20	277	1,352	1,408	514	131	42	9	-
2004 2/	3,862	1	1	2	23	296	1,529	1,329	500	128	42	11	-
2003 2/	4,126	1	-	3	28	375	1,703	1,360	485	125	38	7	1
2002	4,224	-	-	3	23	429	1,790	1,366	460	112	33	8	-
2001	4,286	2	3	8	30	519	1,911	1,248	423	110	27	5	-
2000	4,562	1	1	13	30	619	2,000	1,325	420	122	28	3	-
1999	4,744	2	2	15	34	794	2,082	1,286	369	121	32	7	-
Non-Hispanic white, male													
2005 2/	3,095	-	1	-	9	195	1,122	1,177	441	117	30	3	-
2004 2/	3,227	-	-	1	14	227	1,259	1,130	448	114	31	3	-
2003 2/	3,487	1	-	2	17	294	1,429	1,172	428	103	35	5	1
2002	3,582	-	-	1	12	335	1,517	1,183	409	99	22	4	-
2001	3,669	2	-	5	15	410	1,639	1,106	370	95	24	3	-
2000	3,837	1	-	5	20	475	1,678	1,161	369	103	25	-	-
1999	4,043	2	1	10	22	643	1,768	1,136	331	104	24	2	-
Non-Hispanic white, female													
2005 2/	660	-	1	-	11	82	230	231	73	14	12	6	-
2004 2/	635	1	1	1	9	69	270	199	52	14	11	8	-
2003 2/	639	-	-	1	11	81	274	188	57	22	3	2	-
2002	642	-	-	2	11	94	273	183	51	13	11	4	-
2001	617	-	3	3	15	109	272	142	53	15	3	2	-
2000	725	-	1	8	10	144	322	164	51	19	3	3	-
1999	701	-	1	5	12	151	314	150	38	17	8	5	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Human immunodeficiency virus (HIV) disease (B20-B24)													
Non-Hispanic black, both sexes													
2005 2/	6,908	-	-	8	108	824	2,298	2,475	902	240	49	3	1
2004 2/	7,163	1	2	17	141	936	2,522	2,423	841	229	45	5	1
2003 2/	7,351	3	1	31	117	938	2,786	2,382	802	239	45	6	1
2002	7,714	3	5	19	124	1,110	3,011	2,469	675	240	54	3	1
2001	7,710	7	10	26	161	1,234	3,072	2,249	690	208	49	2	2
2000	7,708	9	12	23	125	1,414	3,028	2,211	625	214	41	6	-
1999	7,751	9	17	41	130	1,479	3,154	2,057	613	213	33	5	-
Non-Hispanic black, male													
2005 2/	4,600	-	-	5	57	455	1,454	1,728	689	179	31	1	1
2004 2/	4,693	-	-	8	73	501	1,601	1,679	637	164	27	2	1
2003 2/	4,947	1	1	17	57	537	1,814	1,693	614	175	31	6	1
2002	5,203	1	1	10	60	624	1,978	1,791	512	182	42	1	1
2001	5,227	5	8	15	68	753	2,040	1,629	524	144	38	1	2
2000	5,291	7	6	12	60	822	2,060	1,642	492	159	29	2	-
1999	5,377	6	11	23	52	919	2,157	1,552	471	160	21	5	-
Non-Hispanic black, female													
2005 2/	2,308	-	-	3	51	369	844	747	213	61	18	2	-
2004 2/	2,470	1	2	9	68	435	921	744	204	65	18	3	-
2003 2/	2,404	2	-	14	60	401	972	689	188	64	14	-	-
2002	2,511	2	4	9	64	486	1,033	678	163	58	12	2	-
2001	2,483	2	2	11	93	481	1,032	620	166	64	11	1	-
2000	2,417	2	6	11	65	592	968	569	133	55	12	4	-
1999	2,374	3	6	18	78	560	997	505	142	53	12	-	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Human immunodeficiency virus (HIV) disease (B20-B24)													
Origin not stated 3/, both sexes													
2005	85	-	-	-	-	8	32	32	8	4	1	-	-
2004	110	-	-	1	1	5	28	48	20	6	1	-	-
2003	139	-	-	-	-	11	52	51	15	8	1	-	1
2002	146	-	-	-	-	13	57	63	11	2	-	-	-
2001	181	-	-	-	-	22	64	61	25	8	1	-	-
2000	181	-	-	-	-	19	80	49	18	13	1	-	1
1999	206	-	-	1	-	21	72	75	25	10	1	-	1
Origin not stated 3/, male													
2005	69	-	-	-	-	4	28	25	7	4	1	-	-
2004	83	-	-	1	-	4	23	34	15	6	-	-	-
2003	104	-	-	-	-	8	35	39	12	8	1	-	1
2002	117	-	-	-	-	9	42	55	10	1	-	-	-
2001	142	-	-	-	-	12	46	55	20	8	1	-	-
2000	151	-	-	-	-	15	60	45	18	12	-	-	1
1999	171	-	-	-	-	16	56	65	24	8	1	-	1
Origin not stated 3/, female													
2005	16	-	-	-	-	4	4	7	1	-	-	-	-
2004	27	-	-	-	1	1	5	14	5	-	1	-	-
2003	35	-	-	-	-	3	17	12	3	-	-	-	-
2002	29	-	-	-	-	4	15	8	1	1	-	-	-
2001	39	-	-	-	-	10	18	6	5	-	-	-	-
2000	30	-	-	-	-	4	20	4	-	1	1	-	-
1999	35	-	-	1	-	5	16	10	1	2	-	-	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Malaria (B50-B54)													
All origins, both sexes													
2005	6	-	-	-	2	1	1	1	-	1	-	-	-
2004	8	-	-	-	1	-	4	2	-	1	-	-	-
2003	4	-	-	-	-	-	1	-	2	1	-	-	-
2002	12	-	-	-	1	1	1	5	2	2	-	-	-
2001	9	-	-	1	-	-	2	2	2	1	1	-	-
2000	3	-	-	-	1	-	-	1	-	-	1	-	-
1999	7	1	-	-	-	2	1	-	1	2	-	-	-
All origins, male													
2005	4	-	-	-	-	1	1	1	-	1	-	-	-
2004	8	-	-	-	1	-	4	2	-	1	-	-	-
2003	4	-	-	-	-	-	1	-	2	1	-	-	-
2002	8	-	-	-	-	-	1	4	1	2	-	-	-
2001	5	-	-	1	-	-	-	1	1	1	1	-	-
2000	2	-	-	-	-	-	-	1	-	-	1	-	-
1999	4	1	-	-	-	1	-	-	1	1	-	-	-
All origins, female													
2005	2	-	-	-	2	-	-	-	-	-	-	-	-
2004	-	-	-	-	-	-	-	-	-	-	-	-	-
2003	-	-	-	-	-	-	-	-	-	-	-	-	-
2002	4	-	-	-	1	1	-	1	1	-	-	-	-
2001	4	-	-	-	-	-	2	1	1	-	-	-	-
2000	1	-	-	-	1	-	-	-	-	-	-	-	-
1999	3	-	-	-	-	1	1	-	-	1	-	-	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Malaria (B50-B54)													
Hispanic, both sexes													
2005	-	-	-	-	-	-	-	-	-	-	-	-	-
2004	1	-	-	-	1	-	-	-	-	-	-	-	-
2003	-	-	-	-	-	-	-	-	-	-	-	-	-
2002	-	-	-	-	-	-	-	-	-	-	-	-	-
2001	-	-	-	-	-	-	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-	-	-	-
Hispanic, male													
2005	-	-	-	-	-	-	-	-	-	-	-	-	-
2004	1	-	-	-	1	-	-	-	-	-	-	-	-
2003	-	-	-	-	-	-	-	-	-	-	-	-	-
2002	-	-	-	-	-	-	-	-	-	-	-	-	-
2001	-	-	-	-	-	-	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-	-	-	-
Hispanic, female													
2005	-	-	-	-	-	-	-	-	-	-	-	-	-
2004	-	-	-	-	-	-	-	-	-	-	-	-	-
2003	-	-	-	-	-	-	-	-	-	-	-	-	-
2002	-	-	-	-	-	-	-	-	-	-	-	-	-
2001	-	-	-	-	-	-	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-	-	-	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Malaria (B50-B54)													
Non-Hispanic 1/, both sexes													
2005	6	-	-	-	2	1	1	1	-	1	-	-	-
2004	7	-	-	-	-	-	4	2	-	1	-	-	-
2003	4	-	-	-	-	-	1	-	2	1	-	-	-
2002	12	-	-	-	1	1	1	5	2	2	-	-	-
2001	9	-	-	1	-	-	2	2	2	1	1	-	-
2000	3	-	-	-	1	-	-	1	-	-	1	-	-
1999	7	1	-	-	-	2	1	-	1	2	-	-	-
Non-Hispanic 1/, male													
2005	4	-	-	-	-	1	1	1	-	1	-	-	-
2004	7	-	-	-	-	-	4	2	-	1	-	-	-
2003	4	-	-	-	-	-	1	-	2	1	-	-	-
2002	8	-	-	-	-	-	1	4	1	2	-	-	-
2001	5	-	-	1	-	-	-	1	1	1	1	-	-
2000	2	-	-	-	-	-	-	1	-	-	1	-	-
1999	4	1	-	-	-	1	-	-	1	1	-	-	-
Non-Hispanic 1/, female													
2005	2	-	-	-	2	-	-	-	-	-	-	-	-
2004	-	-	-	-	-	-	-	-	-	-	-	-	-
2003	-	-	-	-	-	-	-	-	-	-	-	-	-
2002	4	-	-	-	1	1	-	1	1	-	-	-	-
2001	4	-	-	-	-	-	2	1	1	-	-	-	-
2000	1	-	-	-	1	-	-	-	-	-	-	-	-
1999	3	-	-	-	-	1	1	-	-	1	-	-	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Malaria (B50-B54)													
Non-Hispanic white, both sexes													
2005 2/	2	-	-	-	1	-	1	-	-	-	-	-	-
2004 2/	2	-	-	-	-	-	2	-	-	-	-	-	-
2003 2/	4	-	-	-	-	-	1	-	2	1	-	-	-
2002	7	-	-	-	-	-	1	3	1	2	-	-	-
2001	5	-	-	-	-	-	2	1	-	1	1	-	-
2000	1	-	-	-	1	-	-	-	-	-	-	-	-
1999	4	-	-	-	-	1	1	-	1	1	-	-	-
Non-Hispanic white, male													
2005 2/	1	-	-	-	-	-	1	-	-	-	-	-	-
2004 2/	2	-	-	-	-	-	2	-	-	-	-	-	-
2003 2/	4	-	-	-	-	-	1	-	2	1	-	-	-
2002	7	-	-	-	-	-	1	3	1	2	-	-	-
2001	2	-	-	-	-	-	-	-	-	1	1	-	-
2000	-	-	-	-	-	-	-	-	-	-	-	-	-
1999	3	-	-	-	-	1	-	-	1	1	-	-	-
Non-Hispanic white, female													
2005 2/	1	-	-	-	1	-	-	-	-	-	-	-	-
2004 2/	-	-	-	-	-	-	-	-	-	-	-	-	-
2003 2/	-	-	-	-	-	-	-	-	-	-	-	-	-
2002	-	-	-	-	-	-	-	-	-	-	-	-	-
2001	3	-	-	-	-	-	2	1	-	-	-	-	-
2000	1	-	-	-	1	-	-	-	-	-	-	-	-
1999	1	-	-	-	-	-	1	-	-	-	-	-	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Malaria (B50-B54)													
Non-Hispanic black, both sexes													
2005 2/	3	-	-	-	1	-	-	1	-	1	-	-	-
2004 2/	3	-	-	-	-	-	2	1	-	-	-	-	-
2003 2/	-	-	-	-	-	-	-	-	-	-	-	-	-
2002	4	-	-	-	1	1	-	2	-	-	-	-	-
2001	3	-	-	1	-	-	-	1	1	-	-	-	-
2000	1	-	-	-	-	-	-	1	-	-	-	-	-
1999	3	1	-	-	-	1	-	-	-	1	-	-	-
Non-Hispanic black, male													
2005 2/	2	-	-	-	-	-	-	1	-	1	-	-	-
2004 2/	3	-	-	-	-	-	2	1	-	-	-	-	-
2003 2/	-	-	-	-	-	-	-	-	-	-	-	-	-
2002	1	-	-	-	-	-	-	1	-	-	-	-	-
2001	2	-	-	1	-	-	-	1	-	-	-	-	-
2000	1	-	-	-	-	-	-	1	-	-	-	-	-
1999	1	1	-	-	-	-	-	-	-	-	-	-	-
Non-Hispanic black, female													
2005 2/	1	-	-	-	1	-	-	-	-	-	-	-	-
2004 2/	-	-	-	-	-	-	-	-	-	-	-	-	-
2003 2/	-	-	-	-	-	-	-	-	-	-	-	-	-
2002	3	-	-	-	1	1	-	1	-	-	-	-	-
2001	1	-	-	-	-	-	-	-	1	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-	-	-	-
1999	2	-	-	-	-	1	-	-	-	1	-	-	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Malaria (B50-B54)													
Origin not stated 3/, both sexes													
2005	-	-	-	-	-	-	-	-	-	-	-	-	-
2004	-	-	-	-	-	-	-	-	-	-	-	-	-
2003	-	-	-	-	-	-	-	-	-	-	-	-	-
2002	-	-	-	-	-	-	-	-	-	-	-	-	-
2001	-	-	-	-	-	-	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-	-	-	-
Origin not stated 3/, male													
2005	-	-	-	-	-	-	-	-	-	-	-	-	-
2004	-	-	-	-	-	-	-	-	-	-	-	-	-
2003	-	-	-	-	-	-	-	-	-	-	-	-	-
2002	-	-	-	-	-	-	-	-	-	-	-	-	-
2001	-	-	-	-	-	-	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-	-	-	-
Origin not stated 3/, female													
2005	-	-	-	-	-	-	-	-	-	-	-	-	-
2004	-	-	-	-	-	-	-	-	-	-	-	-	-
2003	-	-	-	-	-	-	-	-	-	-	-	-	-
2002	-	-	-	-	-	-	-	-	-	-	-	-	-
2001	-	-	-	-	-	-	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-	-	-	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Other and unspecified infectious and parasitic diseases and their sequelae (A00,A05,A20-A36,A42-A44,A48-A49,A54-A79,A81-A82,A85.0-A85.1,A85.8,A86-B04,B06-B09, B25-B49,B55-B99)													
All origins, both sexes													
2005	7,727	169	63	64	86	130	403	1,364	1,273	1,309	1,785	1,081	-
2004	7,540	167	65	47	95	144	434	1,236	1,237	1,309	1,714	1,092	-
2003	7,427	174	85	51	88	140	447	1,185	1,136	1,344	1,715	1,062	-
2002	6,707	224	61	84	97	169	453	916	889	1,223	1,594	997	-
2001	5,715	191	64	64	86	147	369	663	771	1,143	1,330	887	-
2000	5,532	167	71	67	96	141	350	585	707	1,074	1,375	899	-
1999	7,279	203	81	79	90	242	545	836	898	1,443	1,739	1,123	-
All origins, male													
2005	4,216	100	40	29	43	80	270	933	768	711	864	378	-
2004	4,097	100	35	25	59	81	257	834	760	710	790	446	-
2003	3,964	86	43	25	52	83	284	792	689	711	816	383	-
2002	3,622	125	39	40	52	96	291	626	532	663	785	373	-
2001	2,931	101	37	29	51	83	227	426	449	565	656	307	-
2000	2,861	94	41	35	54	75	220	358	383	597	665	339	-
1999	3,655	119	51	44	49	131	333	478	482	760	827	381	-
All origins, female													
2005	3,511	69	23	35	43	50	133	431	505	598	921	703	-
2004	3,443	67	30	22	36	63	177	402	477	599	924	646	-
2003	3,463	88	42	26	36	57	163	393	447	633	899	679	-
2002	3,085	99	22	44	45	73	162	290	357	560	809	624	-
2001	2,784	90	27	35	35	64	142	237	322	578	674	580	-
2000	2,671	73	30	32	42	66	130	227	324	477	710	560	-
1999	3,624	84	30	35	41	111	212	358	416	683	912	742	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Other and unspecified infectious and parasitic diseases and their sequelae (A00,A05,A20-A36,A42-A44,A48-A49,A54-A79,A81-A82,A85.0-A85.1,A85.8,A86-B04,B06-B09, B25-B49,B55-B99)													
Hispanic, both sexes													
2005	773	33	12	11	15	35	82	207	132	113	97	36	-
2004	726	31	20	10	17	22	82	174	134	97	100	39	-
2003	675	23	17	14	12	21	77	169	116	103	98	25	-
2002	593	41	11	17	19	51	67	127	85	75	68	32	-
2001	423	41	7	11	12	33	46	69	56	67	57	24	-
2000	412	23	18	13	19	26	50	61	54	62	57	29	-
1999	497	36	13	14	13	39	67	68	62	80	68	37	-
Hispanic, male													
2005	452	17	8	5	8	23	58	151	82	52	39	9	-
2004	439	15	11	6	15	15	56	122	88	50	49	12	-
2003	416	9	8	7	8	14	57	130	76	53	43	11	-
2002	363	27	7	11	7	31	48	95	52	40	30	15	-
2001	250	21	4	5	9	22	31	47	38	38	27	8	-
2000	237	14	12	8	12	20	37	33	35	28	27	11	-
1999	260	22	10	7	6	22	49	40	29	36	29	10	-
Hispanic, female													
2005	321	16	4	6	7	12	24	56	50	61	58	27	-
2004	287	16	9	4	2	7	26	52	46	47	51	27	-
2003	259	14	9	7	4	7	20	39	40	50	55	14	-
2002	230	14	4	6	12	20	19	32	33	35	38	17	-
2001	173	20	3	6	3	11	15	22	18	29	30	16	-
2000	175	9	6	5	7	6	13	28	19	34	30	18	-
1999	237	14	3	7	7	17	18	28	33	44	39	27	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Other and unspecified infectious and parasitic diseases and their sequelae (A00,A05,A20-A36,A42-A44,A48-A49,A54-A79,A81-A82,A85.0-A85.1,A85.8,A86-B04,B06-B09, B25-B49,B55-B99)													
Non-Hispanic 1/, both sexes													
2005	6,938	136	51	53	71	95	319	1,154	1,135	1,194	1,686	1,044	-
2004	6,797	136	45	37	77	121	352	1,059	1,102	1,208	1,610	1,050	-
2003	6,730	151	66	37	75	119	368	1,010	1,018	1,235	1,614	1,037	-
2002	6,097	181	49	67	77	117	384	787	801	1,148	1,522	964	-
2001	5,271	149	57	53	73	114	319	589	711	1,073	1,272	861	-
2000	5,100	144	53	54	75	115	297	521	649	1,012	1,312	868	-
1999	6,753	166	68	63	77	202	475	763	831	1,358	1,666	1,084	-
Non-Hispanic 1/, male													
2005	3,753	83	32	24	35	57	210	780	681	658	824	369	-
2004	3,651	85	24	19	43	66	201	711	671	658	739	434	-
2003	3,536	77	33	18	43	69	227	659	613	654	771	372	-
2002	3,250	97	31	29	44	65	242	530	479	623	753	357	-
2001	2,666	79	33	24	42	61	193	374	408	525	628	299	-
2000	2,613	80	29	27	42	55	182	322	347	569	633	327	-
1999	3,379	96	41	36	43	108	282	435	449	722	797	370	-
Non-Hispanic 1/, female													
2005	3,185	53	19	29	36	38	109	374	454	536	862	675	-
2004	3,146	51	21	18	34	55	151	348	431	550	871	616	-
2003	3,194	74	33	19	32	50	141	351	405	581	843	665	-
2002	2,847	84	18	38	33	52	142	257	322	525	769	607	-
2001	2,605	70	24	29	31	53	126	215	303	548	644	562	-
2000	2,487	64	24	27	33	60	115	199	302	443	679	541	-
1999	3,374	70	27	27	34	94	193	328	382	636	869	714	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Other and unspecified infectious and parasitic diseases and their sequelae (A00,A05,A20-A36,A42-A44,A48-A49,A54-A79,A81-A82,A85.0-A85.1,A85.8,A86-B04,B06-B09, B25-B49,B55-B99)													
Non-Hispanic white, both sexes													
2005 2/	5,662	85	34	41	40	50	217	896	908	996	1,459	936	-
2004 2/	5,545	86	28	23	53	67	243	848	868	986	1,399	944	-
2003 2/	5,509	87	36	25	51	76	257	773	794	1,056	1,399	955	-
2002	4,939	113	34	49	47	66	243	582	606	973	1,344	882	-
2001	4,257	80	37	38	45	63	211	435	551	899	1,123	775	-
2000	4,131	79	33	40	51	58	189	383	505	851	1,165	777	-
1999	5,288	101	43	42	51	104	287	509	628	1,126	1,438	959	-
Non-Hispanic white, male													
2005 2/	3,071	52	20	16	18	30	146	627	544	562	722	334	-
2004 2/	2,977	55	14	13	29	35	138	574	527	547	647	398	-
2003 2/	2,871	42	17	12	31	43	163	512	478	554	675	344	-
2002	2,608	57	21	24	27	33	161	398	358	537	665	327	-
2001	2,148	40	21	17	24	39	129	281	320	443	562	272	-
2000	2,092	45	16	18	29	28	115	239	275	483	557	287	-
1999	2,644	64	27	24	28	55	190	303	345	608	681	319	-
Non-Hispanic white, female													
2005 2/	2,591	33	14	25	22	20	71	269	364	434	737	602	-
2004 2/	2,568	31	14	10	24	32	105	274	341	439	752	546	-
2003 2/	2,638	45	19	13	20	33	94	261	316	502	724	611	-
2002	2,331	56	13	25	20	33	82	184	248	436	679	555	-
2001	2,109	40	16	21	21	24	82	154	231	456	561	503	-
2000	2,039	34	17	22	22	30	74	144	230	368	608	490	-
1999	2,644	37	16	18	23	49	97	206	283	518	757	640	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Other and unspecified infectious and parasitic diseases and their sequelae (A00,A05,A20-A36,A42-A44,A48-A49,A54-A79,A81-A82,A85.0-A85.1,A85.8,A86-B04,B06-B09, B25-B49,B55-B99)													
Non-Hispanic black, both sexes													
2005 2/	960	43	12	10	22	40	83	206	177	134	150	83	-
2004 2/	955	43	11	10	19	42	92	163	178	148	163	86	-
2003 2/	942	53	23	9	18	33	99	202	165	116	157	67	-
2002	927	58	13	10	25	46	121	179	151	122	130	72	-
2001	815	61	16	13	23	45	91	130	129	135	105	67	-
2000	815	60	14	10	21	50	95	124	116	129	120	76	-
1999	1,247	57	23	17	20	89	169	219	169	192	186	106	-
Non-Hispanic black, male													
2005 2/	514	27	7	6	11	22	54	117	108	67	70	25	-
2004 2/	520	24	8	5	11	23	49	110	114	75	71	30	-
2003 2/	515	30	12	5	9	20	57	124	97	73	66	22	-
2002	516	33	9	3	13	28	68	117	99	62	58	26	-
2001	419	36	10	6	15	19	56	80	70	66	42	19	-
2000	428	31	10	5	10	22	58	75	62	70	58	27	-
1999	617	29	14	10	12	48	81	114	82	94	94	39	-
Non-Hispanic black, female													
2005 2/	446	16	5	4	11	18	29	89	69	67	80	58	-
2004 2/	435	19	3	5	8	19	43	53	64	73	92	56	-
2003 2/	427	23	11	4	9	13	42	78	68	43	91	45	-
2002	411	25	4	7	12	18	53	62	52	60	72	46	-
2001	396	25	6	7	8	26	35	50	59	69	63	48	-
2000	387	29	4	5	11	28	37	49	54	59	62	49	-
1999	630	28	9	7	8	41	88	105	87	98	92	67	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Other and unspecified infectious and parasitic diseases and their sequelae (A00,A05,A20-A36,A42-A44,A48-A49,A54-A79,A81-A82,A85.0-A85.1,A85.8,A86-B04,B06-B09, B25-B49,B55-B99)													
Origin not stated 3/, both sexes													
2005	16	-	-	-	-	-	2	3	6	2	2	1	-
2004	17	-	-	-	1	1	-	3	1	4	4	3	-
2003	22	-	2	-	1	-	2	6	2	6	3	-	-
2002	17	2	1	-	1	1	2	2	3	-	4	1	-
2001	21	1	-	-	1	-	4	5	4	3	1	2	-
2000	20	-	-	-	2	-	3	3	4	-	6	2	-
1999	29	1	-	2	-	1	3	5	5	5	5	2	-
Origin not stated 3/, male													
2005	11	-	-	-	-	-	2	2	5	1	1	-	-
2004	7	-	-	-	1	-	-	1	1	2	2	-	-
2003	12	-	2	-	1	-	-	3	-	4	2	-	-
2002	9	1	1	-	1	-	1	1	1	-	2	1	-
2001	15	1	-	-	-	-	3	5	3	2	1	-	-
2000	11	-	-	-	-	-	1	3	1	-	5	1	-
1999	16	1	-	1	-	1	2	3	4	2	1	1	-
Origin not stated 3/, female													
2005	5	-	-	-	-	-	-	1	1	1	1	1	-
2004	10	-	-	-	-	1	-	2	-	2	2	3	-
2003	10	-	-	-	-	-	2	3	2	2	1	-	-
2002	8	1	-	-	-	1	1	1	2	-	2	-	-
2001	6	-	-	-	1	-	1	-	1	1	-	2	-
2000	9	-	-	-	2	-	2	-	3	-	1	1	-
1999	13	-	-	1	-	-	1	2	1	3	4	1	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Malignant neoplasms (C00-C97)													
All origins, both sexes													
2005	559,312	75	377	1,000	1,717	3,601	14,566	50,405	99,240	138,446	166,421	83,455	9
2004	553,888	74	399	1,019	1,709	3,633	14,723	49,520	96,956	139,417	166,085	80,345	8
2003	556,902	75	392	1,076	1,651	3,741	15,509	49,843	95,692	141,248	167,617	80,046	12
2002	557,271	74	402	1,072	1,730	3,872	16,085	49,637	93,391	144,757	167,062	79,182	7
2001	553,768	66	420	1,008	1,704	3,994	16,569	49,562	90,223	147,018	165,445	77,751	8
2000	553,091	92	420	1,014	1,713	3,916	16,520	48,034	89,005	150,131	165,099	77,136	11
1999	549,838	68	418	1,012	1,724	4,005	16,732	46,681	89,067	152,338	162,770	75,014	9
All origins, male													
2005	290,422	45	217	557	1,035	1,805	6,338	25,403	54,039	76,685	87,096	37,197	5
2004	286,830	37	212	572	1,027	1,750	6,408	25,432	52,738	76,487	86,734	35,427	6
2003	287,990	34	227	591	979	1,804	6,825	25,542	51,927	77,793	87,382	34,878	8
2002	288,768	42	216	600	1,016	1,868	7,056	25,192	51,115	80,085	86,952	34,621	5
2001	287,075	31	229	533	1,024	1,860	7,297	25,071	49,244	81,699	86,126	33,954	7
2000	286,082	50	233	569	1,022	1,854	7,344	24,215	48,427	83,191	85,907	33,260	10
1999	285,832	36	227	555	1,023	1,959	7,471	23,634	48,833	84,544	85,060	32,484	6
All origins, female													
2005	268,890	30	160	443	682	1,796	8,228	25,002	45,201	61,761	79,325	46,258	4
2004	267,058	37	187	447	682	1,883	8,315	24,088	44,218	62,930	79,351	44,918	2
2003	268,912	41	165	485	672	1,937	8,684	24,301	43,765	63,455	80,235	45,168	4
2002	268,503	32	186	472	714	2,004	9,029	24,445	42,276	64,672	80,110	44,561	2
2001	266,693	35	191	475	680	2,134	9,272	24,491	40,979	65,319	79,319	43,797	1
2000	267,009	42	187	445	691	2,062	9,176	23,819	40,578	66,940	79,192	43,876	1
1999	264,006	32	191	457	701	2,046	9,261	23,047	40,234	67,794	77,710	42,530	3

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Malignant neoplasms (C00-C97)													
Hispanic, both sexes													
2005	26,156	18	80	194	323	530	1,415	3,324	4,836	6,359	6,438	2,637	2
2004	24,522	19	70	211	305	526	1,345	3,040	4,659	5,996	5,970	2,378	3
2003	24,070	13	98	217	288	522	1,355	3,081	4,482	5,929	5,701	2,383	1
2002	23,141	18	72	210	304	502	1,346	2,859	4,247	5,857	5,365	2,361	-
2001	22,371	14	84	177	289	497	1,396	2,815	4,126	5,724	5,068	2,181	-
2000	21,160	17	75	198	275	476	1,295	2,575	3,780	5,557	4,827	2,085	-
1999	20,233	16	72	176	259	466	1,204	2,516	3,686	5,394	4,449	1,995	-
Hispanic, male													
2005	13,896	6	52	99	202	280	601	1,635	2,693	3,564	3,478	1,286	-
2004	13,002	7	29	118	207	267	604	1,566	2,528	3,313	3,283	1,077	3
2003	12,671	4	60	123	186	274	566	1,550	2,442	3,347	3,035	1,084	-
2002	12,235	10	35	108	191	252	557	1,430	2,377	3,256	2,902	1,117	-
2001	11,825	4	44	90	199	233	615	1,372	2,202	3,219	2,814	1,033	-
2000	11,138	10	43	122	162	241	534	1,232	2,034	3,086	2,657	1,017	-
1999	10,668	11	34	92	158	222	504	1,237	2,060	3,045	2,403	902	-
Hispanic, female													
2005	12,260	12	28	95	121	250	814	1,689	2,143	2,795	2,960	1,351	2
2004	11,520	12	41	93	98	259	741	1,474	2,131	2,683	2,687	1,301	-
2003	11,399	9	38	94	102	248	789	1,531	2,040	2,582	2,666	1,299	1
2002	10,906	8	37	102	113	250	789	1,429	1,870	2,601	2,463	1,244	-
2001	10,546	10	40	87	90	264	781	1,443	1,924	2,505	2,254	1,148	-
2000	10,022	7	32	76	113	235	761	1,343	1,746	2,471	2,170	1,068	-
1999	9,565	5	38	84	101	244	700	1,279	1,626	2,349	2,046	1,093	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Malignant neoplasms (C00-C97)													
Non-Hispanic 1/, both sexes													
2005	532,383	56	296	802	1,392	3,059	13,131	46,981	94,243	131,875	159,808	80,735	5
2004	528,395	54	328	804	1,397	3,098	13,352	46,391	92,093	133,165	159,862	77,847	4
2003	531,638	62	293	857	1,359	3,207	14,120	46,621	90,973	135,014	161,606	77,515	11
2002	532,712	55	329	861	1,418	3,350	14,693	46,628	88,886	138,545	161,303	76,639	5
2001	530,058	52	333	828	1,413	3,485	15,121	46,617	85,842	140,943	160,023	75,394	7
2000	530,531	75	341	812	1,432	3,431	15,170	45,305	84,977	144,200	159,906	74,871	11
1999	528,218	52	343	835	1,462	3,525	15,482	44,036	85,146	146,572	157,904	72,852	9
Non-Hispanic 1/, male													
2005	276,084	39	165	455	831	1,518	5,724	23,705	51,251	72,997	83,520	35,876	3
2004	273,282	29	182	452	816	1,477	5,794	23,812	50,078	73,022	83,325	34,293	2
2003	274,651	30	166	467	790	1,520	6,243	23,909	49,353	74,260	84,179	33,726	8
2002	275,770	31	181	492	821	1,610	6,475	23,682	48,587	76,621	83,839	33,427	4
2001	274,492	27	183	442	824	1,623	6,652	23,631	46,871	78,266	83,121	32,846	6
2000	274,185	40	186	446	858	1,607	6,787	22,902	46,253	79,878	83,047	32,171	10
1999	274,419	25	190	463	864	1,732	6,946	22,323	46,644	81,271	82,443	31,512	6
Non-Hispanic 1/, female													
2005	256,299	17	131	347	561	1,541	7,407	23,276	42,992	58,878	76,288	44,859	2
2004	255,113	25	146	352	581	1,621	7,558	22,579	42,015	60,143	76,537	43,554	2
2003	256,987	32	127	390	569	1,687	7,877	22,712	41,620	60,754	77,427	43,789	3
2002	256,942	24	148	369	597	1,740	8,218	22,946	40,299	61,924	77,464	43,212	1
2001	255,566	25	150	386	589	1,862	8,469	22,986	38,971	62,677	76,902	42,548	1
2000	256,346	35	155	366	574	1,824	8,383	22,403	38,724	64,322	76,859	42,700	1
1999	253,799	27	153	372	598	1,793	8,536	21,713	38,502	65,301	75,461	41,340	3

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Malignant neoplasms (C00-C97)													
Non-Hispanic white, both sexes													
2005 2/.....	456,133	46	215	597	1,048	2,175	9,976	36,433	77,617	112,940	142,314	72,770	2
2004 2/.....	453,534	44	244	587	1,016	2,270	10,062	36,016	76,400	114,148	142,461	70,283	3
2003 2/.....	457,127	51	221	637	1,039	2,354	10,723	36,168	75,733	116,198	144,012	69,981	10
2002.....	458,754	37	236	642	1,097	2,483	11,234	36,609	73,809	119,464	143,957	69,184	2
2001.....	456,709	42	268	607	1,076	2,594	11,551	36,836	71,389	121,529	142,760	68,053	4
2000.....	458,200	60	256	627	1,059	2,599	11,458	35,861	70,865	124,842	143,079	67,487	7
1999.....	456,315	37	281	608	1,070	2,682	11,848	34,944	70,788	127,119	141,295	65,636	7
Non-Hispanic white, male													
2005 2/.....	236,659	32	118	333	628	1,103	4,480	18,574	42,128	62,528	74,456	32,278	1
2004 2/.....	234,392	24	134	339	593	1,144	4,467	18,556	41,468	62,610	74,178	30,878	1
2003 2/.....	236,150	25	123	348	607	1,159	4,862	18,601	41,026	64,019	74,968	30,404	8
2002.....	237,323	24	134	355	639	1,218	5,041	18,699	40,210	66,250	74,639	30,112	2
2001.....	236,004	21	144	310	632	1,260	5,160	18,679	38,884	67,554	73,885	29,471	4
2000.....	235,932	34	146	338	636	1,259	5,201	18,092	38,368	69,085	73,952	28,814	7
1999.....	236,384	20	160	345	629	1,341	5,425	17,588	38,658	70,518	73,535	28,161	4
Non-Hispanic white, female													
2005 2/.....	219,474	14	97	264	420	1,072	5,496	17,859	35,489	50,412	67,858	40,492	1
2004 2/.....	219,142	20	110	248	423	1,126	5,595	17,460	34,932	51,538	68,283	39,405	2
2003 2/.....	220,977	26	98	289	432	1,195	5,861	17,567	34,707	52,179	69,044	39,577	2
2002.....	221,431	13	102	287	458	1,265	6,193	17,910	33,599	53,214	69,318	39,072	-
2001.....	220,705	21	124	297	444	1,334	6,391	18,157	32,505	53,975	68,875	38,582	-
2000.....	222,268	26	110	289	423	1,340	6,257	17,769	32,497	55,757	69,127	38,673	-
1999.....	219,931	17	121	263	441	1,341	6,423	17,356	32,130	56,601	67,760	37,475	3

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Malignant neoplasms (C00-C97)													
Non-Hispanic black, both sexes													
2005 2/	62,510	9	62	165	259	674	2,521	8,805	13,835	15,560	14,156	6,461	3
2004 2/	61,842	7	62	165	307	651	2,649	8,661	13,149	15,712	14,244	6,234	1
2003 2/	62,054	7	55	173	245	656	2,787	8,802	12,903	15,666	14,482	6,278	-
2002	61,996	12	71	168	242	663	2,870	8,376	12,693	16,084	14,518	6,296	3
2001	61,580	5	46	172	257	691	2,950	8,175	12,196	16,392	14,454	6,239	3
2000	61,390	11	66	147	297	641	3,072	8,068	12,019	16,485	14,244	6,337	3
1999	61,409	11	47	179	314	660	3,049	7,749	12,322	16,625	14,192	6,260	1
Non-Hispanic black, male													
2005 2/	32,363	6	36	98	151	313	960	4,309	7,695	8,630	7,304	2,859	2
2004 2/	32,089	2	35	82	174	254	1,025	4,439	7,276	8,642	7,422	2,737	1
2003 2/	32,131	4	30	96	141	263	1,112	4,533	7,135	8,593	7,540	2,684	-
2002	32,287	4	33	103	137	287	1,158	4,203	7,106	8,835	7,669	2,750	2
2001	32,335	2	29	100	148	267	1,201	4,196	6,752	9,134	7,700	2,804	2
2000	32,493	4	31	86	178	259	1,292	4,152	6,763	9,235	7,689	2,802	2
1999	32,542	3	24	94	181	294	1,255	4,098	6,933	9,228	7,603	2,828	1
Non-Hispanic black, female													
2005 2/	30,147	3	26	67	108	361	1,561	4,496	6,140	6,930	6,852	3,602	1
2004 2/	29,753	5	27	83	133	397	1,624	4,222	5,873	7,070	6,822	3,497	-
2003 2/	29,923	3	25	77	104	393	1,675	4,269	5,768	7,073	6,942	3,594	-
2002	29,709	8	38	65	105	376	1,712	4,173	5,587	7,249	6,849	3,546	1
2001	29,245	3	17	72	109	424	1,749	3,979	5,444	7,258	6,754	3,435	1
2000	28,897	7	35	61	119	382	1,780	3,916	5,256	7,250	6,555	3,535	1
1999	28,867	8	23	85	133	366	1,794	3,651	5,389	7,397	6,589	3,432	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Malignant neoplasms (C00-C97)													
Origin not stated 3/, both sexes													
2005	773	1	1	4	2	12	20	100	161	212	175	83	2
2004	971	1	1	4	7	9	26	89	204	256	253	120	1
2003	1,194	-	1	2	4	12	34	141	237	305	310	148	-
2002	1,418	1	1	1	8	20	46	150	258	355	394	182	2
2001	1,339	-	3	3	2	12	52	130	255	351	354	176	1
2000	1,400	-	4	4	6	9	55	154	248	374	366	180	-
1999	1,387	-	3	1	3	14	46	129	235	372	417	167	-
Origin not stated 3/, male													
2005	442	-	-	3	2	7	13	63	95	124	98	35	2
2004	546	1	1	2	4	6	10	54	132	152	126	57	1
2003	668	-	1	1	3	10	16	83	132	186	168	68	-
2002	763	1	-	-	4	6	24	80	151	208	211	77	1
2001	758	-	2	1	1	4	30	68	171	214	191	75	1
2000	759	-	4	1	2	6	23	81	140	227	203	72	-
1999	745	-	3	-	1	5	21	74	129	228	214	70	-
Origin not stated 3/, female													
2005	331	1	1	1	-	5	7	37	66	88	77	48	-
2004	425	-	-	2	3	3	16	35	72	104	127	63	-
2003	526	-	-	1	1	2	18	58	105	119	142	80	-
2002	655	-	1	1	4	14	22	70	107	147	183	105	1
2001	581	-	1	2	1	8	22	62	84	137	163	101	-
2000	641	-	-	3	4	3	32	73	108	147	163	108	-
1999	642	-	-	1	2	9	25	55	106	144	203	97	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Malignant neoplasms of lip, oral cavity and pharynx (C00-C14)													
All origins, both sexes													
2005	7,773	-	1	2	24	54	258	1,123	1,936	1,835	1,673	867	-
2004	7,826	-	2	5	23	43	256	1,193	1,781	1,874	1,774	875	-
2003	7,778	-	4	8	30	59	274	1,131	1,826	1,874	1,692	879	1
2002	7,737	-	3	8	23	57	258	1,101	1,811	1,898	1,721	857	-
2001	7,702	-	-	4	29	56	283	1,076	1,740	1,912	1,787	814	1
2000	7,492	-	-	4	26	50	293	1,068	1,642	1,959	1,626	824	-
1999	7,486	-	2	6	27	43	291	1,061	1,616	1,896	1,744	800	-
All origins, male													
2005	5,273	-	-	1	16	32	188	874	1,501	1,309	999	353	-
2004	5,312	-	-	2	18	35	176	937	1,400	1,312	1,057	375	-
2003	5,298	-	2	4	21	39	195	921	1,423	1,304	999	389	1
2002	5,184	-	2	3	11	39	179	870	1,392	1,317	997	374	-
2001	5,156	-	-	1	12	35	202	846	1,372	1,327	1,025	335	1
2000	4,952	-	-	4	13	28	215	825	1,270	1,326	939	332	-
1999	4,944	-	1	3	20	27	204	820	1,231	1,308	1,007	323	-
All origins, female													
2005	2,500	-	1	1	8	22	70	249	435	526	674	514	-
2004	2,514	-	2	3	5	8	80	256	381	562	717	500	-
2003	2,480	-	2	4	9	20	79	210	403	570	693	490	-
2002	2,553	-	1	5	12	18	79	231	419	581	724	483	-
2001	2,546	-	-	3	17	21	81	230	368	585	762	479	-
2000	2,540	-	-	-	13	22	78	243	372	633	687	492	-
1999	2,542	-	1	3	7	16	87	241	385	588	737	477	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Malignant neoplasms of lip, oral cavity and pharynx (C00-C14)													
Hispanic, both sexes													
2005	314	-	-	1	3	5	11	50	84	72	76	12	-
2004	332	-	1	-	2	5	12	62	80	82	59	29	-
2003	321	-	1	-	4	5	14	50	75	80	63	29	-
2002	284	-	-	2	3	4	17	47	66	62	54	29	-
2001	311	-	-	-	4	1	14	56	69	78	58	31	-
2000	287	-	-	1	1	6	13	42	75	73	48	28	-
1999	282	-	-	1	-	1	18	41	66	72	63	20	-
Hispanic, male													
2005	236	-	-	1	2	3	8	40	69	59	48	6	-
2004	241	-	-	-	2	5	6	52	63	62	36	15	-
2003	233	-	1	-	2	4	8	41	64	56	43	14	-
2002	215	-	-	-	1	3	14	39	53	51	36	18	-
2001	219	-	-	-	2	1	11	41	55	58	39	12	-
2000	218	-	-	1	-	4	11	34	62	58	28	20	-
1999	204	-	-	1	-	-	9	33	55	62	35	9	-
Hispanic, female													
2005	78	-	-	-	1	2	3	10	15	13	28	6	-
2004	91	-	1	-	-	-	6	10	17	20	23	14	-
2003	88	-	-	-	2	1	6	9	11	24	20	15	-
2002	69	-	-	2	2	1	3	8	13	11	18	11	-
2001	92	-	-	-	2	-	3	15	14	20	19	19	-
2000	69	-	-	-	1	2	2	8	13	15	20	8	-
1999	78	-	-	-	-	1	9	8	11	10	28	11	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Malignant neoplasms of lip, oral cavity and pharynx (C00-C14)													
Non-Hispanic 1/, both sexes													
2005	7,445	-	1	1	21	49	246	1,070	1,848	1,758	1,596	855	-
2004	7,473	-	1	5	21	38	243	1,125	1,698	1,786	1,712	844	-
2003	7,437	-	3	8	26	53	260	1,079	1,743	1,791	1,627	846	1
2002	7,431	-	3	6	20	53	238	1,049	1,739	1,830	1,666	827	-
2001	7,370	-	-	4	25	55	268	1,018	1,666	1,829	1,724	781	-
2000	7,181	-	-	3	24	44	280	1,022	1,560	1,877	1,575	796	-
1999	7,181	-	2	5	27	42	272	1,018	1,542	1,818	1,678	777	-
Non-Hispanic 1/, male													
2005	5,026	-	-	-	14	29	179	831	1,430	1,246	950	347	-
2004	5,055	-	-	2	16	30	169	881	1,334	1,245	1,020	358	-
2003	5,050	-	1	4	19	34	187	878	1,351	1,246	955	374	1
2002	4,950	-	2	3	10	36	162	826	1,335	1,261	960	355	-
2001	4,920	-	-	1	10	34	190	804	1,313	1,264	982	322	-
2000	4,719	-	-	3	13	24	204	788	1,204	1,263	908	312	-
1999	4,725	-	1	2	20	27	194	785	1,172	1,241	970	313	-
Non-Hispanic 1/, female													
2005	2,419	-	1	1	7	20	67	239	418	512	646	508	-
2004	2,418	-	1	3	5	8	74	244	364	541	692	486	-
2003	2,387	-	2	4	7	19	73	201	392	545	672	472	-
2002	2,481	-	1	3	10	17	76	223	404	569	706	472	-
2001	2,450	-	-	3	15	21	78	214	353	565	742	459	-
2000	2,462	-	-	-	11	20	76	234	356	614	667	484	-
1999	2,456	-	1	3	7	15	78	233	370	577	708	464	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Malignant neoplasms of lip, oral cavity and pharynx (C00-C14)													
Non-Hispanic white, both sexes													
2005 2/	6,096	-	1	1	9	26	177	779	1,453	1,458	1,412	780	-
2004 2/	6,106	-	1	3	12	24	151	789	1,322	1,500	1,522	782	-
2003 2/	6,079	-	2	5	15	36	182	774	1,385	1,483	1,419	777	1
2002	6,141	-	2	3	12	35	164	744	1,358	1,565	1,504	754	-
2001	6,052	-	-	2	18	36	165	736	1,288	1,524	1,560	723	-
2000	5,843	-	-	1	16	29	186	723	1,177	1,566	1,412	733	-
1999	5,826	-	2	3	14	29	176	677	1,180	1,527	1,497	721	-
Non-Hispanic white, male													
2005 2/	4,012	-	-	-	7	16	122	602	1,112	1,015	825	313	-
2004 2/	4,028	-	-	-	8	20	104	614	1,037	1,024	900	321	-
2003 2/	4,043	-	-	2	11	21	145	634	1,055	1,014	819	341	1
2002	4,035	-	2	2	7	25	114	587	1,042	1,067	867	322	-
2001	3,961	-	-	1	6	22	120	586	1,004	1,045	880	297	-
2000	3,726	-	-	1	7	16	128	546	904	1,036	797	291	-
1999	3,746	-	1	1	9	17	125	523	892	1,043	850	285	-
Non-Hispanic white, female													
2005 2/	2,084	-	1	1	2	10	55	177	341	443	587	467	-
2004 2/	2,078	-	1	3	4	4	47	175	285	476	622	461	-
2003 2/	2,036	-	2	3	4	15	37	140	330	469	600	436	-
2002	2,106	-	-	1	5	10	50	157	316	498	637	432	-
2001	2,091	-	-	1	12	14	45	150	284	479	680	426	-
2000	2,117	-	-	-	9	13	58	177	273	530	615	442	-
1999	2,080	-	1	2	5	12	51	154	288	484	647	436	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Malignant neoplasms of lip, oral cavity and pharynx (C00-C14)													
Non-Hispanic black, both sexes													
2005 2/	1,068	-	-	-	10	14	47	249	323	243	135	47	-
2004 2/	1,137	-	-	2	5	11	74	294	321	237	149	44	-
2003 2/	1,100	-	1	3	9	10	63	242	309	261	150	52	-
2002	1,034	-	-	2	6	11	63	254	318	213	114	53	-
2001	1,056	-	-	1	4	6	71	229	318	256	129	42	-
2000	1,089	-	-	2	7	7	60	249	327	254	136	47	-
1999	1,131	-	-	2	8	8	67	287	316	248	150	45	-
Non-Hispanic black, male													
2005 2/	832	-	-	-	5	9	40	200	265	191	99	23	-
2004 2/	863	-	-	2	5	9	49	235	254	190	92	27	-
2003 2/	831	-	1	2	6	9	32	197	260	201	99	24	-
2002	757	-	-	1	3	7	41	203	257	157	64	24	-
2001	773	-	-	-	2	5	45	179	263	184	78	17	-
2000	821	-	-	2	5	4	46	203	262	191	94	14	-
1999	819	-	-	1	6	6	47	224	243	169	102	21	-
Non-Hispanic black, female													
2005 2/	236	-	-	-	5	5	7	49	58	52	36	24	-
2004 2/	274	-	-	-	-	2	25	59	67	47	57	17	-
2003 2/	269	-	-	1	3	1	31	45	49	60	51	28	-
2002	277	-	-	1	3	4	22	51	61	56	50	29	-
2001	283	-	-	1	2	1	26	50	55	72	51	25	-
2000	268	-	-	-	2	3	14	46	65	63	42	33	-
1999	312	-	-	1	2	2	20	63	73	79	48	24	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Malignant neoplasms of lip, oral cavity and pharynx (C00-C14)													
Origin not stated 3/, both sexes													
2005	14	-	-	-	-	-	1	3	4	5	1	-	-
2004	21	-	-	-	-	-	1	6	3	6	3	2	-
2003	20	-	-	-	-	1	-	2	8	3	2	4	-
2002	22	-	-	-	-	-	3	5	6	6	1	1	-
2001	21	-	-	-	-	-	1	2	5	5	5	2	1
2000	24	-	-	-	1	-	-	4	7	9	3	-	-
1999	23	-	-	-	-	-	1	2	8	6	3	3	-
Origin not stated 3/, male													
2005	11	-	-	-	-	-	1	3	2	4	1	-	-
2004	16	-	-	-	-	-	1	4	3	5	1	2	-
2003	15	-	-	-	-	1	-	2	8	2	1	1	-
2002	19	-	-	-	-	-	3	5	4	5	1	1	-
2001	17	-	-	-	-	-	1	1	4	5	4	1	1
2000	15	-	-	-	-	-	-	3	4	5	3	-	-
1999	15	-	-	-	-	-	1	2	4	5	2	1	-
Origin not stated 3/, female													
2005	3	-	-	-	-	-	-	-	2	1	-	-	-
2004	5	-	-	-	-	-	-	2	-	1	2	-	-
2003	5	-	-	-	-	-	-	-	-	1	1	3	-
2002	3	-	-	-	-	-	-	-	2	1	-	-	-
2001	4	-	-	-	-	-	-	1	1	-	1	1	-
2000	9	-	-	-	1	-	-	1	3	4	-	-	-
1999	8	-	-	-	-	-	-	-	4	1	1	2	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Malignant neoplasm of esophagus (C15)													
All origins, both sexes													
2005	13,499	-	-	1	3	35	316	1,510	3,100	3,682	3,581	1,271	-
2004	13,023	-	-	-	5	36	291	1,423	2,991	3,554	3,467	1,256	-
2003	12,860	-	-	-	5	29	288	1,407	2,940	3,619	3,392	1,180	-
2002	12,701	-	-	1	6	19	311	1,390	2,763	3,691	3,389	1,130	1
2001	12,530	-	-	-	6	29	309	1,408	2,554	3,779	3,307	1,137	1
2000	12,232	1	-	-	3	39	297	1,348	2,610	3,671	3,163	1,100	-
1999	11,918	-	-	-	-	37	293	1,262	2,531	3,694	2,986	1,114	1
All origins, male													
2005	10,611	-	-	-	3	30	263	1,302	2,619	2,977	2,679	738	-
2004	10,081	-	-	-	4	28	242	1,211	2,502	2,817	2,546	731	-
2003	9,921	-	-	-	4	23	240	1,202	2,483	2,851	2,464	654	-
2002	9,789	-	-	-	3	14	257	1,185	2,321	2,952	2,454	603	-
2001	9,607	-	-	-	5	26	255	1,204	2,137	2,989	2,365	625	1
2000	9,279	1	-	-	2	31	265	1,161	2,169	2,831	2,233	586	-
1999	9,094	-	-	-	-	29	255	1,094	2,081	2,925	2,136	573	1
All origins, female													
2005	2,888	-	-	1	-	5	53	208	481	705	902	533	-
2004	2,942	-	-	-	1	8	49	212	489	737	921	525	-
2003	2,939	-	-	-	1	6	48	205	457	768	928	526	-
2002	2,912	-	-	1	3	5	54	205	442	739	935	527	1
2001	2,923	-	-	-	1	3	54	204	417	790	942	512	-
2000	2,953	-	-	-	1	8	32	187	441	840	930	514	-
1999	2,824	-	-	-	-	8	38	168	450	769	850	541	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Malignant neoplasm of esophagus (C15)													
Hispanic, both sexes													
2005	481	-	-	-	-	1	12	67	103	148	106	44	-
2004	440	-	-	-	-	4	19	39	105	131	103	39	-
2003	421	-	-	-	1	3	17	50	105	122	92	31	-
2002	411	-	-	-	1	1	23	53	80	120	91	42	-
2001	401	-	-	-	-	2	12	63	96	95	101	32	-
2000	384	-	-	-	-	2	10	38	86	120	98	30	-
1999	367	-	-	-	-	2	16	44	85	95	87	38	-
Hispanic, male													
2005	391	-	-	-	-	1	10	59	89	122	80	30	-
2004	346	-	-	-	-	3	13	39	91	103	76	21	-
2003	322	-	-	-	-	2	16	45	79	93	74	13	-
2002	329	-	-	-	1	1	19	49	75	96	66	22	-
2001	316	-	-	-	-	2	9	61	74	80	68	22	-
2000	292	-	-	-	-	2	8	34	64	92	72	20	-
1999	294	-	-	-	-	2	15	39	78	74	63	23	-
Hispanic, female													
2005	90	-	-	-	-	-	2	8	14	26	26	14	-
2004	94	-	-	-	-	1	6	-	14	28	27	18	-
2003	99	-	-	-	1	1	1	5	26	29	18	18	-
2002	82	-	-	-	-	-	4	4	5	24	25	20	-
2001	85	-	-	-	-	-	3	2	22	15	33	10	-
2000	92	-	-	-	-	-	2	4	22	28	26	10	-
1999	73	-	-	-	-	-	1	5	7	21	24	15	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Malignant neoplasm of esophagus (C15)													
Non-Hispanic 1/, both sexes													
2005	12,997	-	-	1	3	33	304	1,441	2,990	3,529	3,471	1,225	-
2004	12,557	-	-	-	5	32	272	1,377	2,877	3,418	3,360	1,216	-
2003	12,415	-	-	-	4	26	271	1,354	2,826	3,491	3,297	1,146	-
2002	12,254	-	-	1	5	17	287	1,334	2,671	3,560	3,296	1,082	1
2001	12,085	-	-	-	6	27	296	1,339	2,447	3,664	3,201	1,104	1
2000	11,819	1	-	-	3	37	287	1,308	2,521	3,538	3,058	1,066	-
1999	11,520	-	-	-	-	35	276	1,215	2,439	3,587	2,893	1,074	1
Non-Hispanic 1/, male													
2005	10,203	-	-	-	3	28	253	1,241	2,524	2,850	2,598	706	-
2004	9,715	-	-	-	4	25	229	1,166	2,406	2,710	2,466	709	-
2003	9,580	-	-	-	4	21	224	1,154	2,396	2,753	2,389	639	-
2002	9,431	-	-	-	2	12	237	1,133	2,235	2,848	2,386	578	-
2001	9,254	-	-	-	5	24	245	1,138	2,052	2,894	2,292	603	1
2000	8,964	1	-	-	2	29	257	1,126	2,102	2,728	2,156	563	-
1999	8,775	-	-	-	-	27	239	1,053	1,996	2,841	2,068	550	1
Non-Hispanic 1/, female													
2005	2,794	-	-	1	-	5	51	200	466	679	873	519	-
2004	2,842	-	-	-	1	7	43	211	471	708	894	507	-
2003	2,835	-	-	-	-	5	47	200	430	738	908	507	-
2002	2,823	-	-	1	3	5	50	201	436	712	910	504	1
2001	2,831	-	-	-	1	3	51	201	395	770	909	501	-
2000	2,855	-	-	-	1	8	30	182	419	810	902	503	-
1999	2,745	-	-	-	-	8	37	162	443	746	825	524	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Malignant neoplasm of esophagus (C15)													
Non-Hispanic white, both sexes													
2005 2/	11,295	-	-	1	3	29	242	1,170	2,504	3,064	3,140	1,142	-
2004 2/	10,809	-	-	-	3	24	227	1,095	2,381	2,934	3,036	1,109	-
2003 2/	10,631	-	-	-	4	22	222	1,042	2,314	2,992	2,971	1,064	-
2002	10,484	-	-	1	5	15	227	1,076	2,173	3,027	2,961	999	-
2001	10,193	-	-	-	6	22	238	1,039	1,925	3,060	2,888	1,014	1
2000	10,005	-	-	-	3	36	239	1,013	1,997	3,017	2,714	986	-
1999	9,703	-	-	-	-	30	213	936	1,923	3,038	2,576	987	-
Non-Hispanic white, male													
2005 2/	8,969	-	-	-	3	26	206	1,042	2,147	2,500	2,375	670	-
2004 2/	8,482	-	-	-	3	19	200	959	2,034	2,363	2,256	648	-
2003 2/	8,292	-	-	-	4	19	188	908	2,004	2,383	2,190	596	-
2002	8,169	-	-	-	2	11	196	938	1,849	2,465	2,171	537	-
2001	7,863	-	-	-	5	19	200	897	1,645	2,458	2,086	552	1
2000	7,680	-	-	-	2	29	221	895	1,702	2,366	1,941	524	-
1999	7,440	-	-	-	-	22	188	827	1,598	2,451	1,857	497	-
Non-Hispanic white, female													
2005 2/	2,326	-	-	1	-	3	36	128	357	564	765	472	-
2004 2/	2,327	-	-	-	-	5	27	136	347	571	780	461	-
2003 2/	2,339	-	-	-	-	3	34	134	310	609	781	468	-
2002	2,315	-	-	1	3	4	31	138	324	562	790	462	-
2001	2,330	-	-	-	1	3	38	142	280	602	802	462	-
2000	2,325	-	-	-	1	7	18	118	295	651	773	462	-
1999	2,263	-	-	-	-	8	25	109	325	587	719	490	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Malignant neoplasm of esophagus (C15)													
Non-Hispanic black, both sexes													
2005 2/	1,458	-	-	-	-	4	51	242	431	393	273	64	-
2004 2/	1,509	-	-	-	2	6	39	251	444	425	266	76	-
2003 2/	1,578	-	-	-	-	3	47	285	463	435	273	72	-
2002	1,572	-	-	-	-	2	56	234	454	475	282	68	1
2001	1,676	-	-	-	-	3	54	264	466	534	278	77	-
2000	1,645	1	-	-	-	1	44	278	481	462	307	71	-
1999	1,629	-	-	-	-	3	53	261	472	486	280	73	1
Non-Hispanic black, male													
2005 2/	1,065	-	-	-	-	2	39	182	334	294	186	28	-
2004 2/	1,047	-	-	-	1	5	26	187	326	296	167	39	-
2003 2/	1,122	-	-	-	-	2	36	222	351	313	162	36	-
2002	1,118	-	-	-	-	1	38	175	351	340	178	35	-
2001	1,222	-	-	-	-	3	41	210	363	379	185	41	-
2000	1,162	1	-	-	-	-	32	218	369	318	191	33	-
1999	1,187	-	-	-	-	3	44	209	365	340	183	42	1
Non-Hispanic black, female													
2005 2/	393	-	-	-	-	2	12	60	97	99	87	36	-
2004 2/	462	-	-	-	1	1	13	64	118	129	99	37	-
2003 2/	456	-	-	-	-	1	11	63	112	122	111	36	-
2002	454	-	-	-	-	1	18	59	103	135	104	33	1
2001	454	-	-	-	-	-	13	54	103	155	93	36	-
2000	483	-	-	-	-	1	12	60	112	144	116	38	-
1999	442	-	-	-	-	-	9	52	107	146	97	31	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Malignant neoplasm of esophagus (C15)													
Origin not stated 3/, both sexes													
2005	21	-	-	-	-	1	-	2	7	5	4	2	-
2004	26	-	-	-	-	-	-	7	9	5	4	1	-
2003	24	-	-	-	-	-	-	3	9	6	3	3	-
2002	36	-	-	-	-	1	1	3	12	11	2	6	-
2001	44	-	-	-	-	-	1	6	11	20	5	1	-
2000	29	-	-	-	-	-	-	2	3	13	7	4	-
1999	31	-	-	-	-	-	1	3	7	12	6	2	-
Origin not stated 3/, male													
2005	17	-	-	-	-	1	-	2	6	5	1	2	-
2004	20	-	-	-	-	-	-	6	5	4	4	1	-
2003	19	-	-	-	-	-	-	3	8	5	1	2	-
2002	29	-	-	-	-	1	1	3	11	8	2	3	-
2001	37	-	-	-	-	-	1	5	11	15	5	-	-
2000	23	-	-	-	-	-	-	1	3	11	5	3	-
1999	25	-	-	-	-	-	1	2	7	10	5	-	-
Origin not stated 3/, female													
2005	4	-	-	-	-	-	-	-	1	-	3	-	-
2004	6	-	-	-	-	-	-	1	4	1	-	-	-
2003	5	-	-	-	-	-	-	-	1	1	2	1	-
2002	7	-	-	-	-	-	-	-	1	3	-	3	-
2001	7	-	-	-	-	-	-	1	-	5	-	1	-
2000	6	-	-	-	-	-	-	1	-	2	2	1	-
1999	6	-	-	-	-	-	-	1	-	2	1	2	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Malignant neoplasm of stomach (C16)													
All origins, both sexes													
2005	11,514	-	-	-	20	115	448	1,103	1,783	2,615	3,470	1,960	-
2004	11,859	-	-	-	12	122	478	1,142	1,887	2,752	3,415	2,051	-
2003	12,110	-	-	1	19	127	480	1,210	1,874	2,793	3,549	2,057	-
2002	12,198	-	1	-	29	138	436	1,106	1,831	2,816	3,656	2,185	-
2001	12,319	-	-	-	20	122	464	1,127	1,771	3,021	3,793	2,001	-
2000	12,645	-	-	1	14	124	484	1,097	1,804	3,054	3,833	2,234	-
1999	12,711	-	-	1	13	136	505	1,101	1,860	3,152	3,860	2,083	-
All origins, male													
2005	6,752	-	-	-	9	58	266	693	1,204	1,681	1,972	869	-
2004	6,958	-	-	-	7	63	269	722	1,273	1,734	1,941	949	-
2003	7,055	-	-	1	7	63	262	767	1,282	1,826	1,980	867	-
2002	7,133	-	1	-	8	80	239	725	1,215	1,867	2,064	934	-
2001	7,191	-	-	-	9	71	253	731	1,238	1,928	2,148	813	-
2000	7,328	-	-	-	6	69	284	703	1,228	1,966	2,144	928	-
1999	7,459	-	-	1	6	76	283	746	1,252	2,005	2,239	851	-
All origins, female													
2005	4,762	-	-	-	11	57	182	410	579	934	1,498	1,091	-
2004	4,901	-	-	-	5	59	209	420	614	1,018	1,474	1,102	-
2003	5,055	-	-	-	12	64	218	443	592	967	1,569	1,190	-
2002	5,065	-	-	-	21	58	197	381	616	949	1,592	1,251	-
2001	5,128	-	-	-	11	51	211	396	533	1,093	1,645	1,188	-
2000	5,317	-	-	1	8	55	200	394	576	1,088	1,689	1,306	-
1999	5,252	-	-	-	7	60	222	355	608	1,147	1,621	1,232	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Malignant neoplasm of stomach (C16)													
Hispanic, both sexes													
2005	1,272	-	-	-	8	38	95	196	221	274	321	119	-
2004	1,355	-	-	-	4	37	125	187	259	310	301	132	-
2003	1,253	-	-	1	3	34	92	193	239	291	279	121	-
2002	1,185	-	-	-	6	38	97	154	218	259	283	130	-
2001	1,189	-	-	-	4	41	98	182	209	300	253	102	-
2000	1,111	-	-	1	4	24	105	151	167	306	254	99	-
1999	1,055	-	-	-	4	38	85	121	192	271	239	105	-
Hispanic, male													
2005	744	-	-	-	3	23	52	108	137	163	197	61	-
2004	771	-	-	-	4	22	64	96	157	181	187	60	-
2003	708	-	-	1	2	17	46	108	140	180	166	48	-
2002	671	-	-	-	2	26	44	85	128	172	150	64	-
2001	680	-	-	-	2	19	54	104	130	186	135	50	-
2000	655	-	-	-	1	16	52	78	110	205	144	49	-
1999	607	-	-	-	1	25	41	76	126	162	141	35	-
Hispanic, female													
2005	528	-	-	-	5	15	43	88	84	111	124	58	-
2004	584	-	-	-	-	15	61	91	102	129	114	72	-
2003	545	-	-	-	1	17	46	85	99	111	113	73	-
2002	514	-	-	-	4	12	53	69	90	87	133	66	-
2001	509	-	-	-	2	22	44	78	79	114	118	52	-
2000	456	-	-	1	3	8	53	73	57	101	110	50	-
1999	448	-	-	-	3	13	44	45	66	109	98	70	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Malignant neoplasm of stomach (C16)													
Non-Hispanic 1/, both sexes													
2005	10,223	-	-	-	12	77	352	903	1,559	2,335	3,146	1,839	-
2004	10,481	-	-	-	8	84	352	953	1,623	2,439	3,105	1,917	-
2003	10,827	-	-	-	16	91	388	1,014	1,630	2,496	3,260	1,932	-
2002	10,974	-	1	-	23	99	337	950	1,609	2,548	3,363	2,044	-
2001	11,099	-	-	-	16	81	365	943	1,558	2,712	3,529	1,895	-
2000	11,491	-	-	-	10	99	378	943	1,627	2,739	3,568	2,127	-
1999	11,615	-	-	1	9	98	420	970	1,662	2,867	3,615	1,973	-
Non-Hispanic 1/, male													
2005	5,997	-	-	-	6	35	213	582	1,065	1,516	1,773	807	-
2004	6,173	-	-	-	3	40	204	624	1,112	1,552	1,751	887	-
2003	6,326	-	-	-	5	45	216	656	1,138	1,641	1,808	817	-
2002	6,438	-	1	-	6	54	194	639	1,084	1,688	1,908	864	-
2001	6,492	-	-	-	7	52	199	625	1,105	1,737	2,007	760	-
2000	6,648	-	-	-	5	53	231	622	1,113	1,753	1,994	877	-
1999	6,827	-	-	1	5	51	242	662	1,122	1,834	2,096	814	-
Non-Hispanic 1/, female													
2005	4,226	-	-	-	6	42	139	321	494	819	1,373	1,032	-
2004	4,308	-	-	-	5	44	148	329	511	887	1,354	1,030	-
2003	4,501	-	-	-	11	46	172	358	492	855	1,452	1,115	-
2002	4,536	-	-	-	17	45	143	311	525	860	1,455	1,180	-
2001	4,607	-	-	-	9	29	166	318	453	975	1,522	1,135	-
2000	4,843	-	-	-	5	46	147	321	514	986	1,574	1,250	-
1999	4,788	-	-	-	4	47	178	308	540	1,033	1,519	1,159	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Malignant neoplasm of stomach (C16)													
Non-Hispanic white, both sexes													
2005 2/	7,455	-	-	-	6	33	203	584	1,084	1,674	2,397	1,474	-
2004 2/	7,732	-	-	-	3	46	209	601	1,121	1,745	2,436	1,571	-
2003 2/	8,035	-	-	-	12	52	234	637	1,172	1,861	2,540	1,527	-
2002	8,124	-	1	-	15	55	205	616	1,137	1,861	2,613	1,621	-
2001	8,308	-	-	-	6	43	211	615	1,110	2,010	2,779	1,534	-
2000	8,600	-	-	-	5	53	213	625	1,180	2,031	2,781	1,712	-
1999	8,712	-	-	-	4	50	237	646	1,156	2,181	2,859	1,579	-
Non-Hispanic white, male													
2005 2/	4,372	-	-	-	3	19	121	381	743	1,090	1,369	646	-
2004 2/	4,575	-	-	-	1	25	128	412	784	1,114	1,375	736	-
2003 2/	4,718	-	-	-	5	28	140	421	825	1,224	1,424	651	-
2002	4,864	-	1	-	3	31	123	431	791	1,265	1,523	696	-
2001	4,884	-	-	-	2	29	122	416	794	1,288	1,606	627	-
2000	5,033	-	-	-	2	29	129	424	825	1,330	1,580	714	-
1999	5,171	-	-	-	3	30	142	454	798	1,426	1,670	648	-
Non-Hispanic white, female													
2005 2/	3,083	-	-	-	3	14	82	203	341	584	1,028	828	-
2004 2/	3,157	-	-	-	2	21	81	189	337	631	1,061	835	-
2003 2/	3,317	-	-	-	7	24	94	216	347	637	1,116	876	-
2002	3,260	-	-	-	12	24	82	185	346	596	1,090	925	-
2001	3,424	-	-	-	4	14	89	199	316	722	1,173	907	-
2000	3,567	-	-	-	3	24	84	201	355	701	1,201	998	-
1999	3,541	-	-	-	1	20	95	192	358	755	1,189	931	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Malignant neoplasm of stomach (C16)													
Non-Hispanic black, both sexes													
2005 2/	1,949	-	-	-	3	23	99	226	340	463	519	276	-
2004 2/	2,004	-	-	-	4	26	93	255	364	519	495	248	-
2003 2/	2,017	-	-	-	1	22	101	271	347	458	528	289	-
2002	2,112	-	-	-	8	33	90	248	339	518	545	331	-
2001	2,051	-	-	-	5	25	121	237	324	505	548	286	-
2000	2,156	-	-	-	4	36	117	239	335	523	579	323	-
1999	2,183	-	-	1	3	36	127	237	373	507	580	319	-
Non-Hispanic black, male													
2005 2/	1,147	-	-	-	2	10	67	148	242	298	257	123	-
2004 2/	1,171	-	-	-	2	10	50	161	246	338	266	98	-
2003 2/	1,175	-	-	-	-	10	53	183	239	310	279	101	-
2002	1,184	-	-	-	3	20	49	165	223	323	276	125	-
2001	1,177	-	-	-	2	14	63	161	220	332	293	92	-
2000	1,202	-	-	-	2	20	77	154	222	314	295	118	-
1999	1,252	-	-	1	1	19	72	158	241	309	325	126	-
Non-Hispanic black, female													
2005 2/	802	-	-	-	1	13	32	78	98	165	262	153	-
2004 2/	833	-	-	-	2	16	43	94	118	181	229	150	-
2003 2/	842	-	-	-	1	12	48	88	108	148	249	188	-
2002	928	-	-	-	5	13	41	83	116	195	269	206	-
2001	874	-	-	-	3	11	58	76	104	173	255	194	-
2000	954	-	-	-	2	16	40	85	113	209	284	205	-
1999	931	-	-	-	2	17	55	79	132	198	255	193	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Malignant neoplasm of stomach (C16)													
Origin not stated 3/, both sexes													
2005	19	-	-	-	-	-	1	4	3	6	3	2	-
2004	23	-	-	-	-	1	1	2	5	3	9	2	-
2003	30	-	-	-	-	2	-	3	5	6	10	4	-
2002	39	-	-	-	-	1	2	2	4	9	10	11	-
2001	31	-	-	-	-	-	1	2	4	9	11	4	-
2000	43	-	-	-	-	1	1	3	10	9	11	8	-
1999	41	-	-	-	-	-	-	10	6	14	6	5	-
Origin not stated 3/, male													
2005	11	-	-	-	-	-	1	3	2	2	2	1	-
2004	14	-	-	-	-	1	1	2	4	1	3	2	-
2003	21	-	-	-	-	1	-	3	4	5	6	2	-
2002	24	-	-	-	-	-	1	1	3	7	6	6	-
2001	19	-	-	-	-	-	-	2	3	5	6	3	-
2000	25	-	-	-	-	-	1	3	5	8	6	2	-
1999	25	-	-	-	-	-	-	8	4	9	2	2	-
Origin not stated 3/, female													
2005	8	-	-	-	-	-	-	1	1	4	1	1	-
2004	9	-	-	-	-	-	-	-	1	2	6	-	-
2003	9	-	-	-	-	1	-	-	1	1	4	2	-
2002	15	-	-	-	-	1	1	1	1	2	4	5	-
2001	12	-	-	-	-	-	1	-	1	4	5	1	-
2000	18	-	-	-	-	1	-	-	5	1	5	6	-
1999	16	-	-	-	-	-	-	2	2	5	4	3	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Malignant neoplasms of colon, rectum and anus (C18-C21)													
All origins, both sexes													
2005	53,252	1	-	4	47	305	1,297	4,343	8,153	11,792	16,254	11,054	2
2004	53,772	-	-	1	54	279	1,326	4,273	8,208	11,954	16,714	10,963	-
2003	55,958	-	-	4	40	291	1,315	4,442	8,304	12,934	17,331	11,296	1
2002	56,741	1	-	1	50	290	1,425	4,573	8,421	13,392	17,498	11,089	1
2001	56,887	-	-	-	46	299	1,433	4,548	7,984	13,791	17,527	11,259	-
2000	57,477	-	-	3	40	272	1,369	4,237	7,940	14,101	18,053	11,462	-
1999	57,155	1	-	1	49	283	1,333	4,057	7,967	14,249	17,929	11,286	-
All origins, male													
2005	26,843	1	-	4	27	167	694	2,401	4,812	6,789	8,003	3,943	2
2004	26,930	-	-	1	34	152	695	2,442	4,785	6,792	8,189	3,840	-
2003	28,007	-	-	3	25	158	714	2,509	4,831	7,346	8,471	3,949	1
2002	28,501	1	-	1	34	161	797	2,564	5,035	7,580	8,504	3,823	1
2001	28,225	-	-	-	24	157	781	2,512	4,652	7,773	8,451	3,875	-
2000	28,462	-	-	2	24	153	759	2,359	4,608	7,924	8,760	3,873	-
1999	28,240	-	-	1	27	160	678	2,250	4,621	7,883	8,754	3,866	-
All origins, female													
2005	26,409	-	-	-	20	138	603	1,942	3,341	5,003	8,251	7,111	-
2004	26,842	-	-	-	20	127	631	1,831	3,423	5,162	8,525	7,123	-
2003	27,951	-	-	1	15	133	601	1,933	3,473	5,588	8,860	7,347	-
2002	28,240	-	-	-	16	129	628	2,009	3,386	5,812	8,994	7,266	-
2001	28,662	-	-	-	22	142	652	2,036	3,332	6,018	9,076	7,384	-
2000	29,015	-	-	1	16	119	610	1,878	3,332	6,177	9,293	7,589	-
1999	28,915	1	-	-	22	123	655	1,807	3,346	6,366	9,175	7,420	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Malignant neoplasms of colon, rectum and anus (C18-C21)													
Hispanic, both sexes													
2005	2,542	-	-	1	6	41	122	303	438	614	674	343	-
2004	2,438	-	-	-	9	42	114	289	471	604	599	310	-
2003	2,471	-	-	-	10	34	122	314	493	607	612	279	-
2002	2,364	-	-	1	2	27	121	282	454	605	573	299	-
2001	2,250	-	-	-	6	20	127	245	415	592	535	310	-
2000	2,117	-	-	-	8	33	118	263	395	530	497	273	-
1999	2,031	-	-	-	7	31	86	245	344	581	485	252	-
Hispanic, male													
2005	1,350	-	-	1	2	17	61	172	265	356	334	142	-
2004	1,327	-	-	-	3	21	72	172	281	336	319	123	-
2003	1,363	-	-	-	8	20	72	191	295	351	314	112	-
2002	1,287	-	-	1	-	13	62	170	291	351	295	104	-
2001	1,185	-	-	-	3	8	76	124	245	333	261	135	-
2000	1,124	-	-	-	3	17	52	140	227	301	267	117	-
1999	1,062	-	-	-	5	14	39	133	185	331	257	98	-
Hispanic, female													
2005	1,192	-	-	-	4	24	61	131	173	258	340	201	-
2004	1,111	-	-	-	6	21	42	117	190	268	280	187	-
2003	1,108	-	-	-	2	14	50	123	198	256	298	167	-
2002	1,077	-	-	-	2	14	59	112	163	254	278	195	-
2001	1,065	-	-	-	3	12	51	121	170	259	274	175	-
2000	993	-	-	-	5	16	66	123	168	229	230	156	-
1999	969	-	-	-	2	17	47	112	159	250	228	154	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Malignant neoplasms of colon, rectum and anus (C18-C21)													
Non-Hispanic 1/, both sexes													
2005	50,629	1	-	3	41	264	1,172	4,030	7,703	11,157	15,557	10,700	1
2004	51,252	-	-	1	45	236	1,210	3,978	7,728	11,326	16,088	10,640	-
2003	53,372	-	-	4	30	257	1,193	4,111	7,793	12,304	16,687	10,992	1
2002	54,244	1	-	-	48	259	1,299	4,281	7,948	12,758	16,882	10,767	1
2001	54,513	-	-	-	40	279	1,301	4,289	7,545	13,172	16,963	10,924	-
2000	55,244	-	-	3	32	238	1,247	3,963	7,523	13,544	17,527	11,167	-
1999	54,976	1	-	1	41	252	1,241	3,804	7,604	13,640	17,390	11,002	-
Non-Hispanic 1/, male													
2005	25,445	1	-	3	25	150	630	2,223	4,540	6,420	7,655	3,797	1
2004	25,561	-	-	1	31	130	622	2,268	4,499	6,442	7,855	3,713	-
2003	26,579	-	-	3	17	138	642	2,307	4,525	6,981	8,138	3,827	1
2002	27,140	1	-	-	34	146	730	2,388	4,731	7,208	8,188	3,713	1
2001	26,978	-	-	-	21	149	701	2,382	4,390	7,419	8,181	3,735	-
2000	27,287	-	-	2	21	135	705	2,215	4,373	7,609	8,476	3,751	-
1999	27,103	-	-	1	22	146	636	2,113	4,424	7,538	8,467	3,756	-
Non-Hispanic 1/, female													
2005	25,184	-	-	-	16	114	542	1,807	3,163	4,737	7,902	6,903	-
2004	25,691	-	-	-	14	106	588	1,710	3,229	4,884	8,233	6,927	-
2003	26,793	-	-	1	13	119	551	1,804	3,268	5,323	8,549	7,165	-
2002	27,104	-	-	-	14	113	569	1,893	3,217	5,550	8,694	7,054	-
2001	27,535	-	-	-	19	130	600	1,907	3,155	5,753	8,782	7,189	-
2000	27,957	-	-	1	11	103	542	1,748	3,150	5,935	9,051	7,416	-
1999	27,873	1	-	-	19	106	605	1,691	3,180	6,102	8,923	7,246	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Malignant neoplasms of colon, rectum and anus (C18-C21)													
Non-Hispanic white, both sexes													
2005 2/	42,479	1	-	3	30	188	875	2,989	6,068	9,253	13,529	9,543	-
2004 2/	43,390	-	-	1	26	166	885	3,011	6,213	9,414	14,120	9,554	-
2003 2/	45,205	-	-	4	19	184	845	3,052	6,241	10,297	14,700	9,862	1
2002	46,147	1	-	-	36	206	947	3,271	6,401	10,782	14,821	9,681	1
2001	46,450	-	-	-	28	203	966	3,284	6,092	11,124	14,903	9,850	-
2000	47,317	-	-	3	23	167	898	3,034	6,085	11,543	15,519	10,045	-
1999	47,275	-	-	1	26	182	933	2,956	6,184	11,643	15,427	9,923	-
Non-Hispanic white, male													
2005 2/	21,443	1	-	3	19	103	491	1,682	3,623	5,391	6,725	3,405	-
2004 2/	21,715	-	-	1	18	92	470	1,753	3,683	5,421	6,919	3,358	-
2003 2/	22,624	-	-	3	12	99	467	1,750	3,680	5,942	7,223	3,447	1
2002	23,119	1	-	-	24	114	534	1,853	3,859	6,189	7,211	3,333	1
2001	23,128	-	-	-	13	111	523	1,874	3,616	6,369	7,263	3,359	-
2000	23,369	-	-	2	13	101	515	1,732	3,561	6,553	7,539	3,353	-
1999	23,446	-	-	1	12	105	500	1,679	3,638	6,559	7,556	3,396	-
Non-Hispanic white, female													
2005 2/	21,036	-	-	-	11	85	384	1,307	2,445	3,862	6,804	6,138	-
2004 2/	21,675	-	-	-	8	74	415	1,258	2,530	3,993	7,201	6,196	-
2003 2/	22,581	-	-	1	7	85	378	1,302	2,561	4,355	7,477	6,415	-
2002	23,028	-	-	-	12	92	413	1,418	2,542	4,593	7,610	6,348	-
2001	23,322	-	-	-	15	92	443	1,410	2,476	4,755	7,640	6,491	-
2000	23,948	-	-	1	10	66	383	1,302	2,524	4,990	7,980	6,692	-
1999	23,829	-	-	-	14	77	433	1,277	2,546	5,084	7,871	6,527	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Malignant neoplasms of colon, rectum and anus (C18-C21)													
Non-Hispanic black, both sexes													
2005 2/	6,804	-	-	-	10	61	239	886	1,370	1,587	1,683	967	1
2004 2/	6,578	-	-	-	12	53	250	807	1,289	1,626	1,629	912	-
2003 2/	6,881	-	-	-	8	50	287	890	1,313	1,702	1,680	951	-
2002	6,817	-	-	-	10	38	283	857	1,297	1,656	1,756	920	-
2001	6,847	-	-	-	8	56	280	845	1,248	1,726	1,758	926	-
2000	6,820	-	-	-	6	54	290	819	1,215	1,720	1,732	984	-
1999	6,709	1	-	-	14	55	268	711	1,248	1,724	1,733	955	-
Non-Hispanic black, male													
2005 2/	3,301	-	-	-	6	39	115	457	775	833	762	313	1
2004 2/	3,173	-	-	-	10	27	121	428	685	862	767	273	-
2003 2/	3,307	-	-	-	4	25	141	480	716	887	752	302	-
2002	3,322	-	-	-	8	23	156	452	717	840	822	304	-
2001	3,221	-	-	-	7	30	152	428	652	881	766	305	-
2000	3,307	-	-	-	5	26	160	430	676	893	792	325	-
1999	3,160	-	-	-	9	30	123	374	692	850	786	296	-
Non-Hispanic black, female													
2005 2/	3,503	-	-	-	4	22	124	429	595	754	921	654	-
2004 2/	3,405	-	-	-	2	26	129	379	604	764	862	639	-
2003 2/	3,574	-	-	-	4	25	146	410	597	815	928	649	-
2002	3,495	-	-	-	2	15	127	405	580	816	934	616	-
2001	3,626	-	-	-	1	26	128	417	596	845	992	621	-
2000	3,513	-	-	-	1	28	130	389	539	827	940	659	-
1999	3,549	1	-	-	5	25	145	337	556	874	947	659	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Malignant neoplasms of colon, rectum and anus (C18-C21)													
Origin not stated 3/, both sexes													
2005	81	-	-	-	-	-	3	10	12	21	23	11	1
2004	82	-	-	-	-	1	2	6	9	24	27	13	-
2003	115	-	-	-	-	-	-	17	18	23	32	25	-
2002	133	-	-	-	-	4	5	10	19	29	43	23	-
2001	124	-	-	-	-	-	5	14	24	27	29	25	-
2000	116	-	-	-	-	1	4	11	22	27	29	22	-
1999	148	-	-	-	1	-	6	8	19	28	54	32	-
Origin not stated 3/, male													
2005	48	-	-	-	-	-	3	6	7	13	14	4	1
2004	42	-	-	-	-	1	1	2	5	14	15	4	-
2003	65	-	-	-	-	-	-	11	11	14	19	10	-
2002	74	-	-	-	-	2	5	6	13	21	21	6	-
2001	62	-	-	-	-	-	4	6	17	21	9	5	-
2000	51	-	-	-	-	1	2	4	8	14	17	5	-
1999	75	-	-	-	-	-	3	4	12	14	30	12	-
Origin not stated 3/, female													
2005	33	-	-	-	-	-	-	4	5	8	9	7	-
2004	40	-	-	-	-	-	1	4	4	10	12	9	-
2003	50	-	-	-	-	-	-	6	7	9	13	15	-
2002	59	-	-	-	-	2	-	4	6	8	22	17	-
2001	62	-	-	-	-	-	1	8	7	6	20	20	-
2000	65	-	-	-	-	-	2	7	14	13	12	17	-
1999	73	-	-	-	1	-	3	4	7	14	24	20	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Malignant neoplasms of liver and intrahepatic bile ducts (C22)													
All origins, both sexes													
2005	16,076	8	21	16	30	89	380	2,515	3,438	3,747	4,176	1,655	1
2004	15,321	1	22	16	43	84	380	2,348	3,123	3,670	4,059	1,575	-
2003	14,706	6	23	14	36	89	395	2,223	2,765	3,630	3,949	1,576	-
2002	14,047	3	28	18	27	105	414	1,976	2,558	3,581	3,728	1,609	-
2001	13,351	5	18	15	29	75	401	1,868	2,351	3,440	3,649	1,500	-
2000	12,916	2	23	20	30	86	429	1,686	2,256	3,439	3,558	1,387	-
1999	12,382	8	29	11	43	78	444	1,492	2,091	3,491	3,425	1,270	-
All origins, male													
2005	10,548	6	13	9	18	63	258	2,003	2,615	2,470	2,374	719	-
2004	9,832	1	12	8	25	55	289	1,889	2,318	2,382	2,227	626	-
2003	9,482	2	18	9	20	53	288	1,779	1,995	2,330	2,301	687	-
2002	9,011	2	19	12	18	62	308	1,611	1,858	2,367	2,112	642	-
2001	8,483	3	7	7	15	43	302	1,490	1,691	2,224	2,047	654	-
2000	8,176	1	14	8	20	59	297	1,324	1,607	2,246	2,052	548	-
1999	7,689	5	20	9	24	48	318	1,161	1,465	2,284	1,877	478	-
All origins, female													
2005	5,528	2	8	7	12	26	122	512	823	1,277	1,802	936	1
2004	5,489	-	10	8	18	29	91	459	805	1,288	1,832	949	-
2003	5,224	4	5	5	16	36	107	444	770	1,300	1,648	889	-
2002	5,036	1	9	6	9	43	106	365	700	1,214	1,616	967	-
2001	4,868	2	11	8	14	32	99	378	660	1,216	1,602	846	-
2000	4,740	1	9	12	10	27	132	362	649	1,193	1,506	839	-
1999	4,693	3	9	2	19	30	126	331	626	1,207	1,548	792	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Malignant neoplasms of liver and intrahepatic bile ducts (C22)													
Hispanic, both sexes													
2005	1,807	1	4	2	6	8	56	341	434	446	380	128	1
2004	1,565	1	2	-	1	11	61	289	369	391	333	107	-
2003	1,511	-	10	1	1	11	48	287	337	397	319	100	-
2002	1,363	-	3	-	1	8	48	217	310	403	265	108	-
2001	1,256	1	3	3	2	14	46	226	261	327	289	84	-
2000	1,176	1	3	7	4	7	59	176	234	335	262	88	-
1999	1,084	1	4	2	4	7	43	170	214	324	239	76	-
Hispanic, male													
2005	1,185	-	4	1	2	4	36	272	327	277	205	57	-
2004	1,046	1	1	-	-	9	52	226	291	239	176	51	-
2003	986	-	8	1	1	6	37	239	241	245	172	36	-
2002	891	-	2	-	1	4	33	179	225	256	146	45	-
2001	812	1	1	-	1	7	36	187	187	206	152	34	-
2000	762	1	2	4	3	5	45	141	165	208	152	36	-
1999	688	1	3	2	3	5	27	141	159	195	121	31	-
Hispanic, female													
2005	622	1	-	1	4	4	20	69	107	169	175	71	1
2004	519	-	1	-	1	2	9	63	78	152	157	56	-
2003	525	-	2	-	-	5	11	48	96	152	147	64	-
2002	472	-	1	-	-	4	15	38	85	147	119	63	-
2001	444	-	2	3	1	7	10	39	74	121	137	50	-
2000	414	-	1	3	1	2	14	35	69	127	110	52	-
1999	396	-	1	-	1	2	16	29	55	129	118	45	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Malignant neoplasms of liver and intrahepatic bile ducts (C22)													
Non-Hispanic 1/, both sexes													
2005	14,236	7	17	14	24	81	322	2,167	2,993	3,296	3,791	1,524	-
2004	13,718	-	20	16	42	73	318	2,052	2,742	3,272	3,718	1,465	-
2003	13,157	6	13	13	35	77	346	1,927	2,418	3,222	3,626	1,474	-
2002	12,638	3	25	18	26	96	365	1,751	2,234	3,171	3,452	1,497	-
2001	12,053	4	15	12	27	61	352	1,637	2,079	3,103	3,352	1,411	-
2000	11,702	1	20	13	26	78	368	1,502	2,018	3,091	3,287	1,298	-
1999	11,265	7	23	9	39	71	398	1,320	1,869	3,158	3,180	1,191	-
Non-Hispanic 1/, male													
2005	9,339	6	9	8	16	59	220	1,725	2,280	2,190	2,165	661	-
2004	8,761	-	11	8	25	46	237	1,657	2,017	2,139	2,047	574	-
2003	8,469	2	10	8	19	46	250	1,531	1,747	2,078	2,128	650	-
2002	8,085	2	17	12	17	57	274	1,426	1,621	2,106	1,957	596	-
2001	7,640	2	6	7	14	36	263	1,299	1,495	2,009	1,889	620	-
2000	7,384	-	12	4	17	53	251	1,175	1,440	2,028	1,893	511	-
1999	6,976	4	15	7	21	43	290	1,019	1,300	2,081	1,752	444	-
Non-Hispanic 1/, female													
2005	4,897	1	8	6	8	22	102	442	713	1,106	1,626	863	-
2004	4,957	-	9	8	17	27	81	395	725	1,133	1,671	891	-
2003	4,688	4	3	5	16	31	96	396	671	1,144	1,498	824	-
2002	4,553	1	8	6	9	39	91	325	613	1,065	1,495	901	-
2001	4,413	2	9	5	13	25	89	338	584	1,094	1,463	791	-
2000	4,318	1	8	9	9	25	117	327	578	1,063	1,394	787	-
1999	4,289	3	8	2	18	28	108	301	569	1,077	1,428	747	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Malignant neoplasms of liver and intrahepatic bile ducts (C22)													
Non-Hispanic white, both sexes													
2005 2/	10,950	6	15	9	13	39	190	1,485	2,132	2,547	3,188	1,326	-
2004 2/	10,554	-	17	10	26	36	188	1,390	1,962	2,527	3,140	1,258	-
2003 2/	10,149	4	9	7	22	34	216	1,271	1,741	2,506	3,071	1,268	-
2002	9,878	2	19	13	18	49	227	1,167	1,613	2,504	2,970	1,296	-
2001	9,433	4	11	5	20	34	189	1,089	1,505	2,477	2,857	1,242	-
2000	9,121	-	14	8	16	37	201	987	1,447	2,448	2,824	1,139	-
1999	8,890	5	20	4	25	29	238	880	1,340	2,546	2,755	1,048	-
Non-Hispanic white, male													
2005 2/	7,107	6	8	7	7	29	122	1,172	1,615	1,703	1,844	594	-
2004 2/	6,656	-	10	5	14	22	132	1,132	1,419	1,663	1,767	492	-
2003 2/	6,504	2	7	6	12	17	146	1,005	1,270	1,654	1,825	560	-
2002	6,215	2	12	7	9	23	168	951	1,167	1,672	1,681	523	-
2001	5,956	2	4	4	11	17	137	859	1,088	1,651	1,632	551	-
2000	5,679	-	9	4	10	22	134	759	1,020	1,624	1,655	442	-
1999	5,455	2	13	3	14	16	173	660	934	1,702	1,534	404	-
Non-Hispanic white, female													
2005 2/	3,843	-	7	2	6	10	68	313	517	844	1,344	732	-
2004 2/	3,898	-	7	5	12	14	56	258	543	864	1,373	766	-
2003 2/	3,645	2	2	1	10	17	70	266	471	852	1,246	708	-
2002	3,663	-	7	6	9	26	59	216	446	832	1,289	773	-
2001	3,477	2	7	1	9	17	52	230	417	826	1,225	691	-
2000	3,442	-	5	4	6	15	67	228	427	824	1,169	697	-
1999	3,435	3	7	1	11	13	65	220	406	844	1,221	644	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Malignant neoplasms of liver and intrahepatic bile ducts (C22)													
Non-Hispanic black, both sexes													
2005 2/	2,087	1	2	3	7	27	81	483	610	436	331	106	-
2004 2/	1,988	-	2	5	14	22	77	466	498	440	337	127	-
2003 2/	1,851	2	1	3	7	25	77	462	444	385	304	141	-
2002	1,708	1	4	3	4	26	83	400	385	384	292	126	-
2001	1,649	-	3	5	4	15	95	385	373	357	299	113	-
2000	1,581	1	2	2	6	20	101	360	344	383	262	100	-
1999	1,498	1	3	4	7	21	91	291	312	395	265	108	-
Non-Hispanic black, male													
2005 2/	1,460	-	1	1	6	19	56	392	482	288	182	33	-
2004 2/	1,330	-	-	2	9	14	59	372	382	290	155	47	-
2003 2/	1,215	-	-	2	4	15	58	371	316	234	155	60	-
2002	1,143	-	3	3	4	16	60	321	279	261	158	38	-
2001	1,036	-	2	2	1	10	70	302	266	204	142	37	-
2000	1,047	-	1	-	5	14	65	285	256	249	131	41	-
1999	939	1	2	3	2	14	65	238	210	245	133	26	-
Non-Hispanic black, female													
2005 2/	627	1	1	2	1	8	25	91	128	148	149	73	-
2004 2/	658	-	2	3	5	8	18	94	116	150	182	80	-
2003 2/	636	2	1	1	3	10	19	91	128	151	149	81	-
2002	565	1	1	-	-	10	23	79	106	123	134	88	-
2001	613	-	1	3	3	5	25	83	107	153	157	76	-
2000	534	1	1	2	1	6	36	75	88	134	131	59	-
1999	559	-	1	1	5	7	26	53	102	150	132	82	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Malignant neoplasms of liver and intrahepatic bile ducts (C22)													
Origin not stated 3/, both sexes													
2005	33	-	-	-	-	-	2	7	11	5	5	3	-
2004	38	-	-	-	-	-	1	7	12	7	8	3	-
2003	38	-	-	-	-	1	1	9	10	11	4	2	-
2002	46	-	-	-	-	1	1	8	14	7	11	4	-
2001	42	-	-	-	-	-	3	5	11	10	8	5	-
2000	38	-	-	-	-	1	2	8	4	13	9	1	-
1999	33	-	2	-	-	-	3	2	8	9	6	3	-
Origin not stated 3/, male													
2005	24	-	-	-	-	-	2	6	8	3	4	1	-
2004	25	-	-	-	-	-	-	6	10	4	4	1	-
2003	27	-	-	-	-	1	1	9	7	7	1	1	-
2002	35	-	-	-	-	1	1	6	12	5	9	1	-
2001	31	-	-	-	-	-	3	4	9	9	6	-	-
2000	30	-	-	-	-	1	1	8	2	10	7	1	-
1999	25	-	2	-	-	-	1	1	6	8	4	3	-
Origin not stated 3/, female													
2005	9	-	-	-	-	-	-	1	3	2	1	2	-
2004	13	-	-	-	-	-	1	1	2	3	4	2	-
2003	11	-	-	-	-	-	-	-	3	4	3	1	-
2002	11	-	-	-	-	-	-	2	2	2	2	3	-
2001	11	-	-	-	-	-	-	1	2	1	2	5	-
2000	8	-	-	-	-	-	1	-	2	3	2	-	-
1999	8	-	-	-	-	-	2	1	2	1	2	-	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Malignant neoplasm of pancreas (C25)													
All origins, both sexes													
2005	32,760	-	-	2	3	64	565	2,744	5,984	8,288	10,147	4,962	1
2004	31,772	-	-	-	11	62	521	2,641	5,756	8,178	9,853	4,749	1
2003	30,777	-	-	1	10	60	548	2,540	5,320	8,104	9,708	4,486	-
2002	30,264	1	-	-	7	39	596	2,433	5,105	8,187	9,555	4,340	1
2001	29,803	-	1	2	2	55	574	2,410	4,868	8,158	9,322	4,410	1
2000	29,332	2	1	1	5	55	548	2,362	4,708	8,103	9,459	4,087	1
1999	29,082	1	-	-	3	55	594	2,353	4,707	8,293	8,933	4,142	1
All origins, male													
2005	16,147	-	-	2	1	39	332	1,613	3,551	4,421	4,540	1,648	-
2004	15,776	-	-	-	6	39	323	1,647	3,377	4,430	4,416	1,538	-
2003	15,060	-	-	1	6	35	329	1,548	3,106	4,251	4,402	1,382	-
2002	14,877	-	-	-	5	26	364	1,526	3,012	4,270	4,270	1,403	1
2001	14,467	-	1	-	2	30	321	1,467	2,874	4,215	4,191	1,365	1
2000	14,238	1	1	-	3	33	324	1,486	2,749	4,189	4,189	1,262	1
1999	14,176	-	-	-	1	33	369	1,482	2,721	4,347	3,940	1,283	-
All origins, female													
2005	16,613	-	-	-	2	25	233	1,131	2,433	3,867	5,607	3,314	1
2004	15,996	-	-	-	5	23	198	994	2,379	3,748	5,437	3,211	1
2003	15,717	-	-	-	4	25	219	992	2,214	3,853	5,306	3,104	-
2002	15,387	1	-	-	2	13	232	907	2,093	3,917	5,285	2,937	-
2001	15,336	-	-	2	-	25	253	943	1,994	3,943	5,131	3,045	-
2000	15,094	1	-	1	2	22	224	876	1,959	3,914	5,270	2,825	-
1999	14,906	1	-	-	2	22	225	871	1,986	3,946	4,993	2,859	1

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Malignant neoplasm of pancreas (C25)													
Hispanic, both sexes													
2005	1,748	-	-	1	-	8	60	186	368	475	465	184	1
2004	1,570	-	-	-	3	10	45	166	321	448	430	147	-
2003	1,485	-	-	-	-	10	49	157	300	429	386	154	-
2002	1,363	-	-	-	1	4	51	148	273	410	333	143	-
2001	1,336	-	-	-	1	6	46	127	263	419	340	134	-
2000	1,243	1	-	-	2	6	37	128	234	397	313	125	-
1999	1,135	-	-	-	-	6	44	137	218	338	272	120	-
Hispanic, male													
2005	872	-	-	1	-	5	37	113	218	239	197	62	-
2004	789	-	-	-	1	5	31	96	184	240	191	41	-
2003	744	-	-	-	-	5	30	95	176	217	175	46	-
2002	667	-	-	-	1	2	35	96	147	214	136	36	-
2001	643	-	-	-	1	3	26	73	141	212	154	33	-
2000	620	1	-	-	1	2	21	80	133	214	128	40	-
1999	563	-	-	-	-	2	30	84	117	182	108	40	-
Hispanic, female													
2005	876	-	-	-	-	3	23	73	150	236	268	122	1
2004	781	-	-	-	2	5	14	70	137	208	239	106	-
2003	741	-	-	-	-	5	19	62	124	212	211	108	-
2002	696	-	-	-	-	2	16	52	126	196	197	107	-
2001	693	-	-	-	-	3	20	54	122	207	186	101	-
2000	623	-	-	-	1	4	16	48	101	183	185	85	-
1999	572	-	-	-	-	4	14	53	101	156	164	80	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Malignant neoplasm of pancreas (C25)													
Non-Hispanic 1/, both sexes													
2005	30,976	-	-	1	3	55	505	2,556	5,609	7,801	9,675	4,771	-
2004	30,148	-	-	-	8	52	476	2,473	5,420	7,716	9,405	4,597	1
2003	29,233	-	-	1	10	50	498	2,379	5,004	7,665	9,302	4,324	-
2002	28,839	1	-	-	6	35	545	2,278	4,821	7,763	9,200	4,189	1
2001	28,407	-	1	2	1	48	526	2,279	4,598	7,719	8,963	4,269	1
2000	28,026	1	1	1	3	49	510	2,228	4,465	7,692	9,124	3,951	1
1999	27,869	1	-	-	3	49	547	2,207	4,480	7,928	8,638	4,015	1
Non-Hispanic 1/, male													
2005	15,262	-	-	1	1	34	295	1,499	3,330	4,175	4,341	1,586	-
2004	14,953	-	-	-	5	34	292	1,550	3,180	4,180	4,217	1,495	-
2003	14,285	-	-	1	6	30	299	1,452	2,921	4,028	4,216	1,332	-
2002	14,181	-	-	-	4	24	329	1,428	2,856	4,051	4,126	1,362	1
2001	13,792	-	1	-	1	27	294	1,393	2,728	3,989	4,031	1,327	1
2000	13,590	-	1	-	2	31	302	1,401	2,609	3,970	4,054	1,219	1
1999	13,572	-	-	-	1	31	338	1,391	2,598	4,150	3,822	1,241	-
Non-Hispanic 1/, female													
2005	15,714	-	-	-	2	21	210	1,057	2,279	3,626	5,334	3,185	-
2004	15,195	-	-	-	3	18	184	923	2,240	3,536	5,188	3,102	1
2003	14,948	-	-	-	4	20	199	927	2,083	3,637	5,086	2,992	-
2002	14,658	1	-	-	2	11	216	850	1,965	3,712	5,074	2,827	-
2001	14,615	-	-	2	-	21	232	886	1,870	3,730	4,932	2,942	-
2000	14,436	1	-	1	1	18	208	827	1,856	3,722	5,070	2,732	-
1999	14,297	1	-	-	2	18	209	816	1,882	3,778	4,816	2,774	1

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Malignant neoplasm of pancreas (C25)													
Non-Hispanic white, both sexes													
2005 2/	26,366	-	-	1	1	40	393	1,981	4,617	6,597	8,481	4,255	-
2004 2/	25,662	-	-	-	3	37	358	1,931	4,427	6,496	8,305	4,104	1
2003 2/	24,919	-	-	-	7	39	372	1,830	4,097	6,503	8,206	3,865	-
2002	24,647	-	-	-	6	25	424	1,789	3,971	6,533	8,134	3,765	-
2001	24,424	-	1	2	1	36	415	1,816	3,847	6,564	7,931	3,810	1
2000	24,034	1	1	1	2	39	391	1,797	3,690	6,545	8,066	3,500	1
1999	23,903	-	-	-	2	34	421	1,771	3,675	6,784	7,625	3,590	1
Non-Hispanic white, male													
2005 2/	13,102	-	-	1	-	24	234	1,176	2,779	3,605	3,856	1,427	-
2004 2/	12,814	-	-	-	1	26	219	1,233	2,633	3,550	3,794	1,358	-
2003 2/	12,277	-	-	-	3	21	222	1,125	2,434	3,490	3,777	1,205	-
2002	12,242	-	-	-	4	17	257	1,141	2,392	3,480	3,715	1,236	-
2001	11,994	-	1	-	1	22	233	1,131	2,324	3,441	3,624	1,216	1
2000	11,764	-	1	-	2	25	236	1,141	2,183	3,444	3,638	1,093	1
1999	11,684	-	-	-	1	23	266	1,114	2,144	3,605	3,412	1,119	-
Non-Hispanic white, female													
2005 2/	13,264	-	-	-	1	16	159	805	1,838	2,992	4,625	2,828	-
2004 2/	12,848	-	-	-	2	11	139	698	1,794	2,946	4,511	2,746	1
2003 2/	12,642	-	-	-	4	18	150	705	1,663	3,013	4,429	2,660	-
2002	12,405	-	-	-	2	8	167	648	1,579	3,053	4,419	2,529	-
2001	12,430	-	-	2	-	14	182	685	1,523	3,123	4,307	2,594	-
2000	12,270	1	-	1	-	14	155	656	1,507	3,101	4,428	2,407	-
1999	12,219	-	-	-	1	11	155	657	1,531	3,179	4,213	2,471	1

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Malignant neoplasm of pancreas (C25)													
Non-Hispanic black, both sexes													
2005 2/	3,743	-	-	-	1	11	86	482	817	991	946	409	-
2004 2/	3,631	-	-	-	5	10	91	453	811	995	866	400	-
2003 2/	3,579	-	-	1	3	10	103	477	743	958	903	381	-
2002	3,466	1	-	-	-	6	101	399	726	1,016	858	358	1
2001	3,352	-	-	-	-	9	89	399	638	961	865	391	-
2000	3,391	-	-	-	1	9	97	379	664	970	890	381	-
1999	3,368	1	-	-	-	12	112	388	669	972	846	368	-
Non-Hispanic black, male													
2005 2/	1,740	-	-	-	1	7	44	273	460	470	378	107	-
2004 2/	1,717	-	-	-	4	5	55	256	458	518	320	101	-
2003 2/	1,662	-	-	1	3	8	64	292	403	448	343	100	-
2002	1,606	-	-	-	-	5	58	242	399	475	332	94	1
2001	1,501	-	-	-	-	4	47	229	339	460	335	87	-
2000	1,546	-	-	-	-	5	54	229	367	447	350	94	-
1999	1,601	-	-	-	-	8	63	246	386	457	344	97	-
Non-Hispanic black, female													
2005 2/	2,003	-	-	-	-	4	42	209	357	521	568	302	-
2004 2/	1,914	-	-	-	1	5	36	197	353	477	546	299	-
2003 2/	1,917	-	-	-	-	2	39	185	340	510	560	281	-
2002	1,860	1	-	-	-	1	43	157	327	541	526	264	-
2001	1,851	-	-	-	-	5	42	170	299	501	530	304	-
2000	1,845	-	-	-	1	4	43	150	297	523	540	287	-
1999	1,767	1	-	-	-	4	49	142	283	515	502	271	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Malignant neoplasm of pancreas (C25)													
Origin not stated 3/, both sexes													
2005	36	-	-	-	-	1	-	2	7	12	7	7	-
2004	54	-	-	-	-	-	-	2	15	14	18	5	-
2003	59	-	-	-	-	-	1	4	16	10	20	8	-
2002	62	-	-	-	-	-	-	7	11	14	22	8	-
2001	60	-	-	-	-	1	2	4	7	20	19	7	-
2000	63	-	-	-	-	-	1	6	9	14	22	11	-
1999	78	-	-	-	-	-	3	9	9	27	23	7	-
Origin not stated 3/, male													
2005	13	-	-	-	-	-	-	1	3	7	2	-	-
2004	34	-	-	-	-	-	-	1	13	10	8	2	-
2003	31	-	-	-	-	-	-	1	9	6	11	4	-
2002	29	-	-	-	-	-	-	2	9	5	8	5	-
2001	32	-	-	-	-	-	1	1	5	14	6	5	-
2000	28	-	-	-	-	-	1	5	7	5	7	3	-
1999	41	-	-	-	-	-	1	7	6	15	10	2	-
Origin not stated 3/, female													
2005	23	-	-	-	-	1	-	1	4	5	5	7	-
2004	20	-	-	-	-	-	-	1	2	4	10	3	-
2003	28	-	-	-	-	-	1	3	7	4	9	4	-
2002	33	-	-	-	-	-	-	5	2	9	14	3	-
2001	28	-	-	-	-	1	1	3	2	6	13	2	-
2000	35	-	-	-	-	-	-	1	2	9	15	8	-
1999	37	-	-	-	-	-	2	2	3	12	13	5	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Malignant neoplasm of larynx (C32)													
All origins, both sexes													
2005	3,797	-	-	-	-	3	64	423	916	1,156	914	320	1
2004	3,668	-	-	1	1	3	59	443	936	1,051	858	316	-
2003	3,792	-	-	-	-	1	79	420	956	1,136	891	308	1
2002	3,723	-	-	-	-	11	76	425	904	1,084	969	253	1
2001	3,797	-	-	-	-	3	86	419	913	1,170	910	296	-
2000	3,861	-	-	2	1	4	64	426	873	1,219	996	276	-
1999	3,815	-	-	-	-	6	78	405	904	1,239	892	291	-
All origins, male													
2005	2,983	-	-	-	-	2	47	320	752	916	693	252	1
2004	2,935	-	-	-	1	3	48	362	756	854	676	235	-
2003	3,020	-	-	-	-	1	60	354	777	899	708	221	-
2002	2,922	-	-	-	-	6	58	349	738	832	755	183	1
2001	3,018	-	-	-	-	2	66	346	737	919	710	238	-
2000	3,041	-	-	-	-	2	49	340	726	945	765	214	-
1999	2,992	-	-	-	-	5	64	346	735	948	681	213	-
All origins, female													
2005	814	-	-	-	-	1	17	103	164	240	221	68	-
2004	733	-	-	1	-	-	11	81	180	197	182	81	-
2003	772	-	-	-	-	-	19	66	179	237	183	87	1
2002	801	-	-	-	-	5	18	76	166	252	214	70	-
2001	779	-	-	-	-	1	20	73	176	251	200	58	-
2000	820	-	-	2	1	2	15	86	147	274	231	62	-
1999	823	-	-	-	-	1	14	59	169	291	211	78	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Malignant neoplasm of larynx (C32)													
Hispanic, both sexes													
2005	183	-	-	-	-	-	7	24	31	52	53	16	-
2004	190	-	-	-	-	-	2	21	45	62	47	13	-
2003	162	-	-	-	-	-	-	21	44	44	38	15	-
2002	165	-	-	-	-	1	8	19	43	48	33	13	-
2001	163	-	-	-	-	-	3	20	38	60	29	13	-
2000	143	-	-	1	-	1	4	17	29	45	35	11	-
1999	171	-	-	-	-	-	5	25	29	55	40	17	-
Hispanic, male													
2005	165	-	-	-	-	-	7	22	28	50	46	12	-
2004	178	-	-	-	-	-	2	21	42	60	42	11	-
2003	148	-	-	-	-	-	-	19	43	41	31	14	-
2002	139	-	-	-	-	-	7	16	40	42	27	7	-
2001	141	-	-	-	-	-	3	18	34	53	24	9	-
2000	124	-	-	-	-	-	3	16	28	37	30	10	-
1999	149	-	-	-	-	-	4	24	27	48	31	15	-
Hispanic, female													
2005	18	-	-	-	-	-	-	2	3	2	7	4	-
2004	12	-	-	-	-	-	-	-	3	2	5	2	-
2003	14	-	-	-	-	-	-	2	1	3	7	1	-
2002	26	-	-	-	-	1	1	3	3	6	6	6	-
2001	22	-	-	-	-	-	-	2	4	7	5	4	-
2000	19	-	-	1	-	1	1	1	1	8	5	1	-
1999	22	-	-	-	-	-	1	1	2	7	9	2	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Malignant neoplasm of larynx (C32)													
Non-Hispanic 1/, both sexes													
2005	3,605	-	-	-	-	3	57	395	884	1,102	861	303	-
2004	3,463	-	-	1	-	3	57	422	885	983	809	303	-
2003	3,610	-	-	-	-	1	78	396	908	1,086	847	293	1
2002	3,548	-	-	-	-	10	68	405	857	1,033	934	240	1
2001	3,614	-	-	-	-	3	82	398	865	1,106	877	283	-
2000	3,699	-	-	1	1	3	59	406	840	1,166	959	264	-
1999	3,627	-	-	-	-	6	73	377	871	1,180	847	273	-
Non-Hispanic 1/, male													
2005	2,811	-	-	-	-	2	40	295	723	865	647	239	-
2004	2,745	-	-	-	-	3	46	341	709	789	633	224	-
2003	2,853	-	-	-	-	1	59	332	730	853	671	207	-
2002	2,775	-	-	-	-	6	51	332	695	788	726	176	1
2001	2,858	-	-	-	-	2	62	327	694	862	682	229	-
2000	2,901	-	-	-	-	2	45	322	694	901	733	204	-
1999	2,832	-	-	-	-	5	60	320	706	897	646	198	-
Non-Hispanic 1/, female													
2005	794	-	-	-	-	1	17	100	161	237	214	64	-
2004	718	-	-	1	-	-	11	81	176	194	176	79	-
2003	757	-	-	-	-	-	19	64	178	233	176	86	1
2002	773	-	-	-	-	4	17	73	162	245	208	64	-
2001	756	-	-	-	-	1	20	71	171	244	195	54	-
2000	798	-	-	1	1	1	14	84	146	265	226	60	-
1999	795	-	-	-	-	1	13	57	165	283	201	75	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Malignant neoplasm of larynx (C32)													
Non-Hispanic white, both sexes													
2005 2/	2,877	-	-	-	-	2	43	280	654	883	757	258	-
2004 2/	2,793	-	-	1	-	3	40	304	689	799	691	266	-
2003 2/	2,879	-	-	-	-	1	61	269	697	878	722	250	1
2002	2,838	-	-	-	-	6	40	276	653	850	793	220	-
2001	2,876	-	-	-	-	3	58	268	657	893	747	250	-
2000	2,940	-	-	1	1	3	44	281	612	936	822	240	-
1999	2,924	-	-	-	-	4	40	256	660	978	751	235	-
Non-Hispanic white, male													
2005 2/	2,224	-	-	-	-	1	30	211	526	687	566	203	-
2004 2/	2,192	-	-	-	-	3	34	242	543	639	537	194	-
2003 2/	2,251	-	-	-	-	1	48	227	559	676	564	176	-
2002	2,205	-	-	-	-	3	30	230	518	645	619	160	-
2001	2,267	-	-	-	-	2	47	216	529	697	570	206	-
2000	2,291	-	-	-	-	2	31	220	500	721	632	185	-
1999	2,263	-	-	-	-	3	34	219	534	739	568	166	-
Non-Hispanic white, female													
2005 2/	653	-	-	-	-	1	13	69	128	196	191	55	-
2004 2/	601	-	-	1	-	-	6	62	146	160	154	72	-
2003 2/	628	-	-	-	-	-	13	42	138	202	158	74	1
2002	633	-	-	-	-	3	10	46	135	205	174	60	-
2001	609	-	-	-	-	1	11	52	128	196	177	44	-
2000	649	-	-	1	1	1	13	61	112	215	190	55	-
1999	661	-	-	-	-	1	6	37	126	239	183	69	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Malignant neoplasm of larynx (C32)													
Non-Hispanic black, both sexes													
2005 2/	673	-	-	-	-	1	13	111	213	200	97	38	-
2004 2/	625	-	-	-	-	-	17	111	185	173	111	28	-
2003 2/	687	-	-	-	-	-	16	120	206	193	114	38	-
2002	674	-	-	-	-	4	27	123	198	174	131	16	1
2001	668	-	-	-	-	-	23	124	195	188	111	27	-
2000	714	-	-	-	-	-	15	121	218	209	129	22	-
1999	657	-	-	-	-	2	32	117	203	188	89	26	-
Non-Hispanic black, male													
2005 2/	542	-	-	-	-	1	9	80	185	161	75	31	-
2004 2/	518	-	-	-	-	-	12	94	158	142	89	23	-
2003 2/	566	-	-	-	-	-	10	99	167	166	98	26	-
2002	541	-	-	-	-	3	20	96	172	135	101	13	1
2001	531	-	-	-	-	-	14	105	156	144	94	18	-
2000	574	-	-	-	-	-	14	99	186	164	94	17	-
1999	532	-	-	-	-	2	25	97	168	147	72	21	-
Non-Hispanic black, female													
2005 2/	131	-	-	-	-	-	4	31	28	39	22	7	-
2004 2/	107	-	-	-	-	-	5	17	27	31	22	5	-
2003 2/	121	-	-	-	-	-	6	21	39	27	16	12	-
2002	133	-	-	-	-	1	7	27	26	39	30	3	-
2001	137	-	-	-	-	-	9	19	39	44	17	9	-
2000	140	-	-	-	-	-	1	22	32	45	35	5	-
1999	125	-	-	-	-	-	7	20	35	41	17	5	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Malignant neoplasm of larynx (C32)													
Origin not stated 3/, both sexes													
2005	9	-	-	-	-	-	-	4	1	2	-	1	1
2004	15	-	-	-	1	-	-	-	6	6	2	-	-
2003	20	-	-	-	-	-	1	3	4	6	6	-	-
2002	10	-	-	-	-	-	-	1	4	3	2	-	-
2001	20	-	-	-	-	-	1	1	10	4	4	-	-
2000	19	-	-	-	-	-	1	3	4	8	2	1	-
1999	17	-	-	-	-	-	-	3	4	4	5	1	-
Origin not stated 3/, male													
2005	7	-	-	-	-	-	-	3	1	1	-	1	1
2004	12	-	-	-	1	-	-	-	5	5	1	-	-
2003	19	-	-	-	-	-	1	3	4	5	6	-	-
2002	8	-	-	-	-	-	-	1	3	2	2	-	-
2001	19	-	-	-	-	-	1	1	9	4	4	-	-
2000	16	-	-	-	-	-	1	2	4	7	2	-	-
1999	11	-	-	-	-	-	-	2	2	3	4	-	-
Origin not stated 3/, female													
2005	2	-	-	-	-	-	-	1	-	1	-	-	-
2004	3	-	-	-	-	-	-	-	1	1	1	-	-
2003	1	-	-	-	-	-	-	-	-	1	-	-	-
2002	2	-	-	-	-	-	-	-	1	1	-	-	-
2001	1	-	-	-	-	-	-	-	1	-	-	-	-
2000	3	-	-	-	-	-	-	1	-	1	-	1	-
1999	6	-	-	-	-	-	-	1	2	1	1	1	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Malignant neoplasms of trachea, bronchus and lung (C33-C34)													
All origins, both sexes													
2005	159,292	-	3	1	16	133	2,323	12,624	31,363	48,390	49,032	15,404	3
2004	158,091	2	4	4	23	147	2,426	12,448	31,086	49,029	48,464	14,455	3
2003	158,086	-	2	2	25	154	2,478	12,374	30,956	49,386	48,619	14,088	2
2002	157,713	-	1	4	21	165	2,698	12,150	30,684	50,256	48,082	13,652	-
2001	156,058	-	5	2	20	157	2,796	12,041	29,777	51,225	46,700	13,331	4
2000	155,521	2	2	1	22	197	2,756	11,915	29,721	52,258	45,836	12,806	5
1999	152,156	-	-	5	20	179	2,724	11,434	29,344	51,848	44,261	12,339	2
All origins, male													
2005	90,187	-	3	-	9	77	1,197	7,327	18,663	28,207	27,196	7,506	2
2004	89,630	1	1	3	14	87	1,280	7,317	18,440	28,453	26,992	7,039	3
2003	89,964	-	1	1	14	78	1,344	7,313	18,347	28,935	27,064	6,865	2
2002	90,171	-	1	3	16	90	1,386	7,194	18,415	29,543	26,821	6,702	-
2001	90,426	-	4	1	12	77	1,504	7,190	17,873	30,699	26,406	6,656	4
2000	90,469	2	1	-	14	102	1,542	7,123	17,937	31,382	25,966	6,395	5
1999	89,453	-	-	4	13	97	1,474	6,851	18,013	31,287	25,397	6,315	2
All origins, female													
2005	69,105	-	-	1	7	56	1,126	5,297	12,700	20,183	21,836	7,898	1
2004	68,461	1	3	1	9	60	1,146	5,131	12,646	20,576	21,472	7,416	-
2003	68,122	-	1	1	11	76	1,134	5,061	12,609	20,451	21,555	7,223	-
2002	67,542	-	-	1	5	75	1,312	4,956	12,269	20,713	21,261	6,950	-
2001	65,632	-	1	1	8	80	1,292	4,851	11,904	20,526	20,294	6,675	-
2000	65,052	-	1	1	8	95	1,214	4,792	11,784	20,876	19,870	6,411	-
1999	62,703	-	-	1	7	82	1,250	4,583	11,331	20,561	18,864	6,024	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Malignant neoplasms of trachea, bronchus and lung (C33-C34)													
Hispanic, both sexes													
2005	4,490	-	1	-	2	15	111	407	848	1,355	1,337	414	-
2004	4,215	-	1	2	2	25	86	379	786	1,303	1,223	406	2
2003	4,136	-	-	-	4	20	92	390	795	1,293	1,177	365	-
2002	4,038	-	-	2	4	27	90	362	808	1,303	1,101	341	-
2001	3,816	-	2	1	7	16	106	364	758	1,230	1,011	321	-
2000	3,643	-	-	-	3	17	90	342	672	1,191	1,006	322	-
1999	3,545	-	-	-	3	17	98	308	724	1,170	901	324	-
Hispanic, male													
2005	2,878	-	1	-	-	9	64	257	594	894	843	216	-
2004	2,704	-	-	1	1	15	56	255	529	858	782	205	2
2003	2,640	-	-	-	3	7	52	240	549	877	723	189	-
2002	2,617	-	-	1	2	14	45	229	567	846	729	184	-
2001	2,495	-	2	-	6	10	68	239	517	820	657	176	-
2000	2,411	-	-	-	3	10	56	229	471	793	666	183	-
1999	2,373	-	-	-	2	10	58	208	523	798	606	168	-
Hispanic, female													
2005	1,612	-	-	-	2	6	47	150	254	461	494	198	-
2004	1,511	-	1	1	1	10	30	124	257	445	441	201	-
2003	1,496	-	-	-	1	13	40	150	246	416	454	176	-
2002	1,421	-	-	1	2	13	45	133	241	457	372	157	-
2001	1,321	-	-	1	1	6	38	125	241	410	354	145	-
2000	1,232	-	-	-	-	7	34	113	201	398	340	139	-
1999	1,172	-	-	-	1	7	40	100	201	372	295	156	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Malignant neoplasms of trachea, bronchus and lung (C33-C34)													
Non-Hispanic 1/, both sexes													
2005	154,578	-	2	1	14	117	2,209	12,195	30,466	46,959	47,642	14,970	3
2004	153,581	2	3	2	21	121	2,335	12,045	30,234	47,629	47,164	14,024	1
2003	153,616	-	2	2	21	132	2,378	11,951	30,101	47,977	47,352	13,698	2
2002	153,292	-	1	2	17	135	2,601	11,752	29,802	48,836	46,862	13,284	-
2001	151,882	-	3	1	13	141	2,678	11,642	28,936	49,887	45,599	12,978	4
2000	151,483	2	2	1	19	180	2,653	11,525	28,965	50,940	44,742	12,449	5
1999	148,206	-	-	5	17	161	2,618	11,087	28,549	50,536	43,243	11,988	2
Non-Hispanic 1/, male													
2005	87,168	-	2	-	9	67	1,132	7,055	18,033	27,265	26,322	7,281	2
2004	86,740	1	1	2	13	71	1,222	7,049	17,867	27,527	26,166	6,820	1
2003	87,111	-	1	1	11	69	1,287	7,052	17,762	27,973	26,286	6,667	2
2002	87,335	-	1	2	14	76	1,337	6,942	17,806	28,628	26,026	6,503	-
2001	87,706	-	2	1	6	67	1,431	6,929	17,291	29,818	25,695	6,462	4
2000	87,816	2	1	-	11	92	1,479	6,866	17,409	30,511	25,245	6,195	5
1999	86,822	-	-	4	11	87	1,411	6,616	17,438	30,397	24,724	6,132	2
Non-Hispanic 1/, female													
2005	67,410	-	-	1	5	50	1,077	5,140	12,433	19,694	21,320	7,689	1
2004	66,841	1	2	-	8	50	1,113	4,996	12,367	20,102	20,998	7,204	-
2003	66,505	-	1	1	10	63	1,091	4,899	12,339	20,004	21,066	7,031	-
2002	65,957	-	-	-	3	59	1,264	4,810	11,996	20,208	20,836	6,781	-
2001	64,176	-	1	-	7	74	1,247	4,713	11,645	20,069	19,904	6,516	-
2000	63,667	-	1	1	8	88	1,174	4,659	11,556	20,429	19,497	6,254	-
1999	61,384	-	-	1	6	74	1,207	4,471	11,111	20,139	18,519	5,856	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Malignant neoplasms of trachea, bronchus and lung (C33-C34)													
Non-Hispanic white, both sexes													
2005 2/	134,924	-	2	1	12	82	1,743	9,647	25,619	41,162	43,076	13,578	2
2004 2/	134,118	1	2	2	16	83	1,818	9,507	25,573	41,792	42,515	12,809	-
2003 2/	134,437	-	2	1	18	107	1,852	9,391	25,613	42,204	42,752	12,495	2
2002	134,435	-	1	1	13	97	2,026	9,335	25,218	43,104	42,506	12,134	-
2001	133,266	-	2	1	9	114	2,100	9,301	24,713	43,972	41,223	11,829	2
2000	133,061	1	2	1	11	139	1,996	9,176	24,777	44,952	40,666	11,337	3
1999	129,967	-	-	3	9	131	2,025	8,832	24,329	44,577	39,153	10,907	1
Non-Hispanic white, male													
2005 2/	75,500	-	2	-	8	49	882	5,565	14,922	23,737	23,761	6,573	1
2004 2/	74,960	1	-	2	10	48	935	5,448	14,857	23,981	23,465	6,213	-
2003 2/	75,468	-	1	-	9	55	986	5,480	14,861	24,440	23,573	6,061	2
2002	75,754	-	1	1	11	58	1,016	5,415	14,818	25,102	23,408	5,924	-
2001	76,066	-	1	1	4	55	1,095	5,421	14,552	26,056	23,022	5,857	2
2000	76,218	1	1	-	7	73	1,076	5,355	14,682	26,704	22,718	5,598	3
1999	75,338	-	-	2	6	71	1,078	5,155	14,654	26,596	22,219	5,556	1
Non-Hispanic white, female													
2005 2/	59,424	-	-	1	4	33	861	4,082	10,697	17,425	19,315	7,005	1
2004 2/	59,158	-	2	-	6	35	883	4,059	10,716	17,811	19,050	6,596	-
2003 2/	58,969	-	1	1	9	52	866	3,911	10,752	17,764	19,179	6,434	-
2002	58,681	-	-	-	2	39	1,010	3,920	10,400	18,002	19,098	6,210	-
2001	57,200	-	1	-	5	59	1,005	3,880	10,161	17,916	18,201	5,972	-
2000	56,843	-	1	1	4	66	920	3,821	10,095	18,248	17,948	5,739	-
1999	54,629	-	-	1	3	60	947	3,677	9,675	17,981	16,934	5,351	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Malignant neoplasms of trachea, bronchus and lung (C33-C34)													
Non-Hispanic black, both sexes													
2005 2/	16,437	-	-	-	1	19	384	2,228	4,176	4,869	3,682	1,077	1
2004 2/	16,324	1	1	-	4	26	438	2,216	4,058	4,810	3,791	978	1
2003 2/	16,252	-	-	1	3	18	441	2,250	3,955	4,892	3,739	953	-
2002	16,170	-	-	1	4	23	503	2,165	3,994	4,930	3,641	909	-
2001	15,883	-	1	-	3	16	515	2,060	3,685	5,076	3,623	902	2
2000	15,924	1	-	-	5	31	578	2,113	3,688	5,178	3,414	915	1
1999	15,834	-	-	2	8	20	531	2,029	3,743	5,111	3,483	907	-
Non-Hispanic black, male													
2005 2/	9,794	-	-	-	1	11	202	1,316	2,694	2,987	2,053	529	1
2004 2/	9,906	-	1	-	3	14	233	1,414	2,646	2,940	2,173	481	1
2003 2/	9,914	-	-	1	2	12	251	1,399	2,570	3,009	2,192	478	-
2002	9,983	-	-	1	3	14	287	1,378	2,606	3,083	2,163	448	-
2001	10,030	-	1	-	1	7	298	1,352	2,400	3,266	2,226	477	2
2000	10,056	1	-	-	2	14	359	1,372	2,410	3,309	2,105	483	1
1999	10,039	-	-	2	5	11	301	1,338	2,500	3,267	2,127	488	-
Non-Hispanic black, female													
2005 2/	6,643	-	-	-	-	8	182	912	1,482	1,882	1,629	548	-
2004 2/	6,418	1	-	-	1	12	205	802	1,412	1,870	1,618	497	-
2003 2/	6,338	-	-	-	1	6	190	851	1,385	1,883	1,547	475	-
2002	6,187	-	-	-	1	9	216	787	1,388	1,847	1,478	461	-
2001	5,853	-	-	-	2	9	217	708	1,285	1,810	1,397	425	-
2000	5,868	-	-	-	3	17	219	741	1,278	1,869	1,309	432	-
1999	5,795	-	-	-	3	9	230	691	1,243	1,844	1,356	419	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Malignant neoplasms of trachea, bronchus and lung (C33-C34)													
Origin not stated 3/, both sexes													
2005	224	-	-	-	-	1	3	22	49	76	53	20	-
2004	295	-	-	-	-	1	5	24	66	97	77	25	-
2003	334	-	-	-	-	2	8	33	60	116	90	25	-
2002	383	-	-	-	-	3	7	36	74	117	119	27	-
2001	360	-	-	-	-	-	12	35	83	108	90	32	-
2000	395	-	-	-	-	-	13	48	84	127	88	35	-
1999	405	-	-	-	-	1	8	39	71	142	117	27	-
Origin not stated 3/, male													
2005	141	-	-	-	-	1	1	15	36	48	31	9	-
2004	186	-	-	-	-	1	2	13	44	68	44	14	-
2003	213	-	-	-	-	2	5	21	36	85	55	9	-
2002	219	-	-	-	-	-	4	23	42	69	66	15	-
2001	225	-	-	-	-	-	5	22	65	61	54	18	-
2000	242	-	-	-	-	-	7	28	57	78	55	17	-
1999	258	-	-	-	-	-	5	27	52	92	67	15	-
Origin not stated 3/, female													
2005	83	-	-	-	-	-	2	7	13	28	22	11	-
2004	109	-	-	-	-	-	3	11	22	29	33	11	-
2003	121	-	-	-	-	-	3	12	24	31	35	16	-
2002	164	-	-	-	-	3	3	13	32	48	53	12	-
2001	135	-	-	-	-	-	7	13	18	47	36	14	-
2000	153	-	-	-	-	-	6	20	27	49	33	18	-
1999	147	-	-	-	-	1	3	12	19	50	50	12	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Malignant melanoma of skin (C43)													
All origins, both sexes													
2005	8,345	-	1	3	40	216	524	1,197	1,575	1,748	2,031	1,010	-
2004	7,952	-	2	-	39	194	526	1,162	1,547	1,711	1,912	859	-
2003	7,818	-	1	2	38	191	561	1,179	1,547	1,593	1,875	831	-
2002	7,514	-	3	1	41	194	604	1,135	1,351	1,616	1,748	820	1
2001	7,542	-	-	-	41	186	622	1,208	1,333	1,689	1,678	785	-
2000	7,420	-	-	1	41	186	626	1,199	1,303	1,637	1,624	803	-
1999	7,215	1	-	1	39	194	625	1,120	1,263	1,652	1,639	681	-
All origins, male													
2005	5,283	-	1	-	23	138	313	748	1,030	1,177	1,294	559	-
2004	5,125	-	2	-	17	117	312	749	1,058	1,158	1,239	473	-
2003	4,963	-	-	-	24	119	336	753	1,042	1,050	1,203	436	-
2002	4,725	-	2	1	20	119	363	746	915	1,075	1,078	406	-
2001	4,763	-	-	-	24	103	380	756	893	1,156	1,063	388	-
2000	4,592	-	-	1	21	110	355	769	866	1,067	1,013	390	-
1999	4,529	1	-	-	23	110	397	717	851	1,096	1,006	328	-
All origins, female													
2005	3,062	-	-	3	17	78	211	449	545	571	737	451	-
2004	2,827	-	-	-	22	77	214	413	489	553	673	386	-
2003	2,855	-	1	2	14	72	225	426	505	543	672	395	-
2002	2,789	-	1	-	21	75	241	389	436	541	670	414	1
2001	2,779	-	-	-	17	83	242	452	440	533	615	397	-
2000	2,828	-	-	-	20	76	271	430	437	570	611	413	-
1999	2,686	-	-	1	16	84	228	403	412	556	633	353	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Malignant melanoma of skin (C43)													
Hispanic, both sexes													
2005	168	-	-	1	1	14	15	33	31	38	24	11	-
2004	138	-	-	-	4	6	16	29	30	23	24	6	-
2003	159	-	-	1	2	6	15	33	29	37	25	11	-
2002	147	-	-	-	2	7	21	27	24	19	29	18	-
2001	145	-	-	-	2	6	24	23	25	28	28	9	-
2000	144	-	-	-	2	5	26	28	27	26	19	11	-
1999	133	1	-	1	2	5	14	22	30	27	20	11	-
Hispanic, male													
2005	105	-	-	-	1	6	9	20	23	25	15	6	-
2004	79	-	-	-	1	4	9	19	18	13	12	3	-
2003	91	-	-	-	1	4	8	19	16	21	15	7	-
2002	77	-	-	-	1	6	13	13	16	11	13	4	-
2001	78	-	-	-	-	2	17	11	15	19	13	1	-
2000	92	-	-	-	1	4	17	19	19	14	13	5	-
1999	79	1	-	-	1	4	9	16	19	17	9	3	-
Hispanic, female													
2005	63	-	-	1	-	8	6	13	8	13	9	5	-
2004	59	-	-	-	3	2	7	10	12	10	12	3	-
2003	68	-	-	1	1	2	7	14	13	16	10	4	-
2002	70	-	-	-	1	1	8	14	8	8	16	14	-
2001	67	-	-	-	2	4	7	12	10	9	15	8	-
2000	52	-	-	-	1	1	9	9	8	12	6	6	-
1999	54	-	-	1	1	1	5	6	11	10	11	8	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Malignant melanoma of skin (C43)													
Non-Hispanic 1/, both sexes													
2005	8,169	-	1	2	39	202	509	1,163	1,542	1,709	2,004	998	-
2004	7,804	-	2	-	35	188	510	1,129	1,516	1,686	1,885	853	-
2003	7,640	-	1	1	36	183	544	1,144	1,513	1,555	1,846	817	-
2002	7,349	-	3	1	38	186	581	1,106	1,323	1,596	1,714	801	-
2001	7,387	-	-	-	39	179	597	1,184	1,307	1,660	1,646	775	-
2000	7,263	-	-	1	39	181	599	1,167	1,276	1,608	1,602	790	-
1999	7,062	-	-	-	37	189	611	1,094	1,231	1,620	1,612	668	-
Non-Hispanic 1/, male													
2005	5,171	-	1	-	22	132	304	727	1,005	1,152	1,276	552	-
2004	5,038	-	2	-	16	113	303	727	1,039	1,143	1,225	470	-
2003	4,859	-	-	-	23	113	326	732	1,025	1,028	1,185	427	-
2002	4,640	-	2	1	19	113	350	732	896	1,063	1,062	402	-
2001	4,678	-	-	-	24	100	363	744	877	1,136	1,047	387	-
2000	4,490	-	-	1	20	106	337	746	847	1,051	998	384	-
1999	4,438	-	-	-	22	106	388	698	831	1,076	994	323	-
Non-Hispanic 1/, female													
2005	2,998	-	-	2	17	70	205	436	537	557	728	446	-
2004	2,766	-	-	-	19	75	207	402	477	543	660	383	-
2003	2,781	-	1	1	13	70	218	412	488	527	661	390	-
2002	2,709	-	1	-	19	73	231	374	427	533	652	399	-
2001	2,709	-	-	-	15	79	234	440	430	524	599	388	-
2000	2,773	-	-	-	19	75	262	421	429	557	604	406	-
1999	2,624	-	-	-	15	83	223	396	400	544	618	345	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Malignant melanoma of skin (C43)													
Non-Hispanic white, both sexes													
2005 2/	7,973	-	1	1	34	197	491	1,137	1,498	1,677	1,967	970	-
2004 2/	7,654	-	1	-	32	185	500	1,112	1,494	1,651	1,845	834	-
2003 2/	7,471	-	1	1	34	178	533	1,116	1,479	1,526	1,805	798	-
2002	7,190	-	3	1	37	184	568	1,085	1,291	1,554	1,687	780	-
2001	7,250	-	-	-	38	177	590	1,168	1,286	1,616	1,614	761	-
2000	7,090	-	-	1	38	175	579	1,147	1,247	1,570	1,566	767	-
1999	6,936	-	-	-	34	183	604	1,081	1,213	1,593	1,577	651	-
Non-Hispanic white, male													
2005 2/	5,074	-	1	-	19	127	290	714	980	1,141	1,259	543	-
2004 2/	4,967	-	1	-	15	111	297	718	1,026	1,131	1,205	463	-
2003 2/	4,784	-	-	-	23	111	323	718	1,007	1,016	1,165	421	-
2002	4,555	-	2	1	18	111	344	719	876	1,039	1,050	395	-
2001	4,618	-	-	-	23	98	359	737	867	1,116	1,034	384	-
2000	4,414	-	-	1	20	104	327	737	833	1,032	981	379	-
1999	4,370	-	-	-	19	102	385	689	819	1,064	976	316	-
Non-Hispanic white, female													
2005 2/	2,899	-	-	1	15	70	201	423	518	536	708	427	-
2004 2/	2,687	-	-	-	17	74	203	394	468	520	640	371	-
2003 2/	2,687	-	1	1	11	67	210	398	472	510	640	377	-
2002	2,635	-	1	-	19	73	224	366	415	515	637	385	-
2001	2,632	-	-	-	15	79	231	431	419	500	580	377	-
2000	2,676	-	-	-	18	71	252	410	414	538	585	388	-
1999	2,566	-	-	-	15	81	219	392	394	529	601	335	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Malignant melanoma of skin (C43)													
Non-Hispanic black, both sexes													
2005 2/	123	-	-	1	2	3	10	13	26	19	29	20	-
2004 2/	118	-	1	-	3	2	8	12	16	27	33	16	-
2003 2/	120	-	-	-	1	3	10	16	23	26	28	13	-
2002	102	-	-	-	1	1	9	12	21	25	18	15	-
2001	102	-	-	-	1	1	7	9	13	36	24	11	-
2000	134	-	-	-	1	3	15	13	19	30	32	21	-
1999	94	-	-	-	1	4	4	9	14	23	25	14	-
Non-Hispanic black, male													
2005 2/	57	-	-	-	1	3	8	7	15	5	13	5	-
2004 2/	56	-	1	-	1	1	5	6	10	9	17	6	-
2003 2/	49	-	-	-	-	2	3	8	12	12	10	2	-
2002	52	-	-	-	1	1	4	7	13	12	8	6	-
2001	36	-	-	-	1	1	4	3	4	16	6	1	-
2000	59	-	-	-	-	2	9	6	10	14	15	3	-
1999	47	-	-	-	1	2	1	6	10	9	12	6	-
Non-Hispanic black, female													
2005 2/	66	-	-	1	1	-	2	6	11	14	16	15	-
2004 2/	62	-	-	-	2	1	3	6	6	18	16	10	-
2003 2/	71	-	-	-	1	1	7	8	11	14	18	11	-
2002	50	-	-	-	-	-	5	5	8	13	10	9	-
2001	66	-	-	-	-	-	3	6	9	20	18	10	-
2000	75	-	-	-	1	1	6	7	9	16	17	18	-
1999	47	-	-	-	-	2	3	3	4	14	13	8	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Malignant melanoma of skin (C43)													
Origin not stated 3/, both sexes													
2005	8	-	-	-	-	-	-	1	2	1	3	1	-
2004	10	-	-	-	-	-	-	4	1	2	3	-	-
2003	19	-	-	-	-	2	2	2	5	1	4	3	-
2002	18	-	-	-	1	1	2	2	4	1	5	1	1
2001	10	-	-	-	-	1	1	1	1	1	4	1	-
2000	13	-	-	-	-	-	1	4	-	3	3	2	-
1999	20	-	-	-	-	-	-	4	2	5	7	2	-
Origin not stated 3/, male													
2005	7	-	-	-	-	-	-	1	2	-	3	1	-
2004	8	-	-	-	-	-	-	3	1	2	2	-	-
2003	13	-	-	-	-	2	2	2	1	1	3	2	-
2002	8	-	-	-	-	-	-	1	3	1	3	-	-
2001	7	-	-	-	-	1	-	1	1	1	3	-	-
2000	10	-	-	-	-	-	1	4	-	2	2	1	-
1999	12	-	-	-	-	-	-	3	1	3	3	2	-
Origin not stated 3/, female													
2005	1	-	-	-	-	-	-	-	-	1	-	-	-
2004	2	-	-	-	-	-	-	1	-	-	1	-	-
2003	6	-	-	-	-	-	-	-	4	-	1	1	-
2002	10	-	-	-	1	1	2	1	1	-	2	1	1
2001	3	-	-	-	-	-	1	-	-	-	1	1	-
2000	3	-	-	-	-	-	-	-	-	1	1	1	-
1999	8	-	-	-	-	-	-	1	1	2	4	-	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Malignant neoplasm of breast (C50)													
All origins, both sexes													
2005	41,491	1	-	-	12	356	2,497	6,232	8,646	8,113	9,382	6,252	-
2004	41,316	-	-	-	11	396	2,511	6,232	8,474	8,285	9,371	6,036	-
2003	42,000	-	-	-	16	407	2,716	6,365	8,267	8,338	9,644	6,245	2
2002	41,883	-	-	-	10	425	2,722	6,454	7,838	8,505	9,748	6,181	-
2001	41,809	-	-	-	9	463	2,823	6,593	7,632	8,715	9,663	5,911	-
2000	42,300	-	-	-	11	467	2,841	6,385	7,573	9,024	9,750	6,249	-
1999	41,528	-	1	1	13	439	2,741	6,174	7,399	9,109	9,651	6,000	-
All origins, male													
2005	375	1	-	-	-	-	9	33	73	109	111	39	-
2004	362	-	-	-	1	1	10	33	67	91	109	50	-
2003	380	-	-	-	-	2	9	48	71	89	104	56	1
2002	369	-	-	-	-	3	12	44	67	84	112	47	-
2001	415	-	-	-	-	2	14	46	72	120	115	46	-
2000	428	-	-	-	-	3	19	46	85	115	109	51	-
1999	384	-	-	-	-	3	15	41	70	103	98	54	-
All origins, female													
2005	41,116	-	-	-	12	356	2,488	6,199	8,573	8,004	9,271	6,213	-
2004	40,954	-	-	-	10	395	2,501	6,199	8,407	8,194	9,262	5,986	-
2003	41,620	-	-	-	16	405	2,707	6,317	8,196	8,249	9,540	6,189	1
2002	41,514	-	-	-	10	422	2,710	6,410	7,771	8,421	9,636	6,134	-
2001	41,394	-	-	-	9	461	2,809	6,547	7,560	8,595	9,548	5,865	-
2000	41,872	-	-	-	11	464	2,822	6,339	7,488	8,909	9,641	6,198	-
1999	41,144	-	1	1	13	436	2,726	6,133	7,329	9,006	9,553	5,946	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Malignant neoplasm of breast (C50)													
Hispanic, both sexes													
2005	1,942	-	-	-	1	37	241	423	433	344	317	146	-
2004	1,926	-	-	-	1	48	225	397	448	373	282	152	-
2003	1,893	-	-	-	2	55	256	402	428	331	269	150	-
2002	1,736	-	-	-	2	54	218	401	348	323	262	128	-
2001	1,760	-	-	-	1	53	244	408	407	318	219	110	-
2000	1,689	-	-	-	1	54	218	381	359	337	231	108	-
1999	1,551	-	-	-	2	47	197	362	321	290	209	123	-
Hispanic, male													
2005	6	-	-	-	-	-	-	1	2	1	2	-	-
2004	17	-	-	-	-	-	-	5	3	6	2	1	-
2003	9	-	-	-	-	-	-	2	2	2	-	3	-
2002	12	-	-	-	-	1	-	5	-	3	2	1	-
2001	14	-	-	-	-	-	2	1	1	5	5	-	-
2000	19	-	-	-	-	-	2	2	4	6	4	1	-
1999	13	-	-	-	-	-	-	3	3	5	2	-	-
Hispanic, female													
2005	1,936	-	-	-	1	37	241	422	431	343	315	146	-
2004	1,909	-	-	-	1	48	225	392	445	367	280	151	-
2003	1,884	-	-	-	2	55	256	400	426	329	269	147	-
2002	1,724	-	-	-	2	53	218	396	348	320	260	127	-
2001	1,746	-	-	-	1	53	242	407	406	313	214	110	-
2000	1,670	-	-	-	1	54	216	379	355	331	227	107	-
1999	1,538	-	-	-	2	47	197	359	318	285	207	123	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Malignant neoplasm of breast (C50)													
Non-Hispanic 1/, both sexes													
2005	39,490	1	-	-	11	319	2,256	5,798	8,201	7,755	9,050	6,099	-
2004	39,329	-	-	-	10	347	2,282	5,828	8,017	7,897	9,077	5,871	-
2003	40,003	-	-	-	14	351	2,453	5,950	7,813	7,987	9,345	6,088	2
2002	40,037	-	-	-	8	368	2,495	6,033	7,470	8,161	9,464	6,038	-
2001	39,957	-	-	-	8	409	2,576	6,170	7,208	8,379	9,418	5,789	-
2000	40,506	-	-	-	10	413	2,615	5,989	7,192	8,670	9,494	6,123	-
1999	39,881	-	1	1	11	390	2,537	5,800	7,056	8,803	9,419	5,863	-
Non-Hispanic 1/, male													
2005	369	1	-	-	-	-	9	32	71	108	109	39	-
2004	343	-	-	-	1	1	10	28	64	85	107	47	-
2003	371	-	-	-	-	2	9	46	69	87	104	53	1
2002	355	-	-	-	-	2	12	38	66	81	110	46	-
2001	399	-	-	-	-	2	12	45	70	114	110	46	-
2000	409	-	-	-	-	3	17	44	81	109	105	50	-
1999	370	-	-	-	-	3	15	38	67	97	96	54	-
Non-Hispanic 1/, female													
2005	39,121	-	-	-	11	319	2,247	5,766	8,130	7,647	8,941	6,060	-
2004	38,986	-	-	-	9	346	2,272	5,800	7,953	7,812	8,970	5,824	-
2003	39,632	-	-	-	14	349	2,444	5,904	7,744	7,900	9,241	6,035	1
2002	39,682	-	-	-	8	366	2,483	5,995	7,404	8,080	9,354	5,992	-
2001	39,558	-	-	-	8	407	2,564	6,125	7,138	8,265	9,308	5,743	-
2000	40,097	-	-	-	10	410	2,598	5,945	7,111	8,561	9,389	6,073	-
1999	39,511	-	1	1	11	387	2,522	5,762	6,989	8,706	9,323	5,809	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Malignant neoplasm of breast (C50)													
Non-Hispanic white, both sexes													
2005 2/	32,736	1	-	-	6	200	1,553	4,227	6,573	6,594	8,073	5,509	-
2004 2/	32,833	-	-	-	6	215	1,560	4,305	6,523	6,763	8,157	5,304	-
2003 2/	33,451	-	-	-	9	216	1,728	4,475	6,354	6,824	8,356	5,488	1
2002	33,611	-	-	-	4	230	1,733	4,559	6,145	7,001	8,456	5,483	-
2001	33,629	-	-	-	3	255	1,828	4,737	5,895	7,185	8,472	5,254	-
2000	34,386	-	-	-	4	275	1,877	4,620	5,911	7,581	8,591	5,527	-
1999	33,790	-	-	1	6	257	1,802	4,470	5,779	7,623	8,488	5,364	-
Non-Hispanic white, male													
2005 2/	305	1	-	-	-	-	5	25	58	87	94	35	-
2004 2/	282	-	-	-	-	-	6	20	50	70	98	38	-
2003 2/	317	-	-	-	-	1	4	37	57	74	95	48	1
2002	288	-	-	-	-	1	9	27	51	65	94	41	-
2001	329	-	-	-	-	1	6	33	59	95	96	39	-
2000	341	-	-	-	-	1	9	35	63	97	94	42	-
1999	299	-	-	-	-	2	8	28	51	79	84	47	-
Non-Hispanic white, female													
2005 2/	32,431	-	-	-	6	200	1,548	4,202	6,515	6,507	7,979	5,474	-
2004 2/	32,551	-	-	-	6	215	1,554	4,285	6,473	6,693	8,059	5,266	-
2003 2/	33,134	-	-	-	9	215	1,724	4,438	6,297	6,750	8,261	5,440	-
2002	33,323	-	-	-	4	229	1,724	4,532	6,094	6,936	8,362	5,442	-
2001	33,300	-	-	-	3	254	1,822	4,704	5,836	7,090	8,376	5,215	-
2000	34,045	-	-	-	4	274	1,868	4,585	5,848	7,484	8,497	5,485	-
1999	33,491	-	-	1	6	255	1,794	4,442	5,728	7,544	8,404	5,317	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Malignant neoplasm of breast (C50)													
Non-Hispanic black, both sexes													
2005 2/	5,804	-	-	-	5	103	609	1,360	1,374	998	820	535	-
2004 2/	5,546	-	-	-	2	115	619	1,272	1,258	967	807	506	-
2003 2/	5,677	-	-	-	5	117	632	1,258	1,238	1,001	877	549	-
2002	5,579	-	-	-	3	123	670	1,247	1,122	1,005	900	509	-
2001	5,528	-	-	-	5	140	641	1,209	1,125	1,053	862	493	-
2000	5,384	-	-	-	5	120	641	1,167	1,103	961	825	562	-
1999	5,386	-	1	-	5	124	661	1,150	1,095	1,037	852	461	-
Non-Hispanic black, male													
2005 2/	56	-	-	-	-	-	2	6	12	21	13	2	-
2004 2/	56	-	-	-	-	1	3	8	12	14	9	9	-
2003 2/	51	-	-	-	-	1	5	8	12	13	7	5	-
2002	62	-	-	-	-	1	3	10	14	16	13	5	-
2001	64	-	-	-	-	1	5	12	10	17	13	6	-
2000	63	-	-	-	-	2	8	8	16	11	10	8	-
1999	67	-	-	-	-	1	7	9	16	16	11	7	-
Non-Hispanic black, female													
2005 2/	5,748	-	-	-	5	103	607	1,354	1,362	977	807	533	-
2004 2/	5,490	-	-	-	2	114	616	1,264	1,246	953	798	497	-
2003 2/	5,626	-	-	-	5	116	627	1,250	1,226	988	870	544	-
2002	5,517	-	-	-	3	122	667	1,237	1,108	989	887	504	-
2001	5,464	-	-	-	5	139	636	1,197	1,115	1,036	849	487	-
2000	5,321	-	-	-	5	118	633	1,159	1,087	950	815	554	-
1999	5,319	-	1	-	5	123	654	1,141	1,079	1,021	841	454	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Malignant neoplasm of breast (C50)													
Origin not stated 3/, both sexes													
2005	59	-	-	-	-	-	-	11	12	14	15	7	-
2004	61	-	-	-	-	1	4	7	9	15	12	13	-
2003	104	-	-	-	-	1	7	13	26	20	30	7	-
2002	110	-	-	-	-	3	9	20	20	21	22	15	-
2001	92	-	-	-	-	1	3	15	17	18	26	12	-
2000	105	-	-	-	-	-	8	15	22	17	25	18	-
1999	96	-	-	-	-	2	7	12	22	16	23	14	-
Origin not stated 3/, male													
2005	-	-	-	-	-	-	-	-	-	-	-	-	-
2004	2	-	-	-	-	-	-	-	-	-	-	2	-
2003	-	-	-	-	-	-	-	-	-	-	-	-	-
2002	2	-	-	-	-	-	-	1	1	-	-	-	-
2001	2	-	-	-	-	-	-	-	1	1	-	-	-
2000	-	-	-	-	-	-	-	-	-	-	-	-	-
1999	1	-	-	-	-	-	-	-	-	1	-	-	-
Origin not stated 3/, female													
2005	59	-	-	-	-	-	-	11	12	14	15	7	-
2004	59	-	-	-	-	1	4	7	9	15	12	11	-
2003	104	-	-	-	-	1	7	13	26	20	30	7	-
2002	108	-	-	-	-	3	9	19	19	21	22	15	-
2001	90	-	-	-	-	1	3	15	16	17	26	12	-
2000	105	-	-	-	-	-	8	15	22	17	25	18	-
1999	95	-	-	-	-	2	7	12	22	15	23	14	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Malignant neoplasm of cervix uteri (C53)													
All origins, both sexes													
2005	3,924	-	-	-	14	168	622	958	796	575	518	273	-
2004	3,850	-	-	-	19	213	622	850	815	601	484	246	-
2003	3,919	-	-	-	16	201	633	890	793	564	552	270	-
2002	3,952	-	-	-	17	208	666	894	725	630	528	284	-
2001	4,092	-	-	1	12	209	711	953	723	585	583	315	-
2000	4,200	-	-	1	17	230	694	915	748	686	599	310	-
1999	4,205	-	-	-	12	217	699	914	769	696	568	329	1
All origins, male													
2005	...	...	...	...	...	...	...	...	...	...	...	...	...
2004	...	...	...	...	...	...	...	...	...	...	...	...	...
2003	...	...	...	...	...	...	...	...	...	...	...	...	...
2002	...	...	...	...	...	...	...	...	...	...	...	...	...
2001	...	...	...	...	...	...	...	...	...	...	...	...	...
2000	...	...	...	...	...	...	...	...	...	...	...	...	...
1999	...	...	...	...	...	...	...	...	...	...	...	...	...
All origins, female													
2005	3,924	-	-	-	14	168	622	958	796	575	518	273	-
2004	3,850	-	-	-	19	213	622	850	815	601	484	246	-
2003	3,919	-	-	-	16	201	633	890	793	564	552	270	-
2002	3,952	-	-	-	17	208	666	894	725	630	528	284	-
2001	4,092	-	-	1	12	209	711	953	723	585	583	315	-
2000	4,200	-	-	1	17	230	694	915	748	686	599	310	-
1999	4,205	-	-	-	12	217	699	914	769	696	568	329	1

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Malignant neoplasm of cervix uteri (C53)													
Hispanic, both sexes													
2005	447	-	-	-	3	20	103	120	90	64	33	14	-
2004	430	-	-	-	3	32	97	96	95	54	39	14	-
2003	391	-	-	-	3	31	85	109	65	43	42	13	-
2002	388	-	-	-	2	30	86	87	71	55	40	17	-
2001	412	-	-	-	3	36	102	102	75	39	39	16	-
2000	434	-	-	-	4	33	108	87	78	60	43	21	-
1999	378	-	-	-	2	31	87	99	64	51	31	13	-
Hispanic, male													
2005	...	...	...	...	...	...	...	...	...	...	...	...	...
2004	...	...	...	...	...	...	...	...	...	...	...	...	...
2003	...	...	...	...	...	...	...	...	...	...	...	...	...
2002	...	...	...	...	...	...	...	...	...	...	...	...	...
2001	...	...	...	...	...	...	...	...	...	...	...	...	...
2000	...	...	...	...	...	...	...	...	...	...	...	...	...
1999	...	...	...	...	...	...	...	...	...	...	...	...	...
Hispanic, female													
2005	447	-	-	-	3	20	103	120	90	64	33	14	-
2004	430	-	-	-	3	32	97	96	95	54	39	14	-
2003	391	-	-	-	3	31	85	109	65	43	42	13	-
2002	388	-	-	-	2	30	86	87	71	55	40	17	-
2001	412	-	-	-	3	36	102	102	75	39	39	16	-
2000	434	-	-	-	4	33	108	87	78	60	43	21	-
1999	378	-	-	-	2	31	87	99	64	51	31	13	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Malignant neoplasm of cervix uteri (C53)													
Non-Hispanic 1/, both sexes													
2005	3,469	-	-	-	11	147	518	837	704	509	485	258	-
2004	3,415	-	-	-	16	181	523	753	720	547	443	232	-
2003	3,519	-	-	-	13	170	546	777	727	519	510	257	-
2002	3,551	-	-	-	15	177	577	805	650	573	487	267	-
2001	3,669	-	-	1	9	172	606	846	646	546	544	299	-
2000	3,755	-	-	1	13	197	583	825	668	625	554	289	-
1999	3,817	-	-	-	10	186	611	813	702	644	535	315	1
Non-Hispanic 1/, male													
2005	...	...	...	...	...	...	...	...	...	...	...	...	...
2004	...	...	...	...	...	...	...	...	...	...	...	...	...
2003	...	...	...	...	...	...	...	...	...	...	...	...	...
2002	...	...	...	...	...	...	...	...	...	...	...	...	...
2001	...	...	...	...	...	...	...	...	...	...	...	...	...
2000	...	...	...	...	...	...	...	...	...	...	...	...	...
1999	...	...	...	...	...	...	...	...	...	...	...	...	...
Non-Hispanic 1/, female													
2005	3,469	-	-	-	11	147	518	837	704	509	485	258	-
2004	3,415	-	-	-	16	181	523	753	720	547	443	232	-
2003	3,519	-	-	-	13	170	546	777	727	519	510	257	-
2002	3,551	-	-	-	15	177	577	805	650	573	487	267	-
2001	3,669	-	-	1	9	172	606	846	646	546	544	299	-
2000	3,755	-	-	1	13	197	583	825	668	625	554	289	-
1999	3,817	-	-	-	10	186	611	813	702	644	535	315	1

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Malignant neoplasm of cervix uteri (C53)													
Non-Hispanic white, both sexes													
2005 2/	2,538	-	-	-	7	113	369	595	506	378	367	203	-
2004 2/	2,474	-	-	-	11	128	372	539	540	385	335	164	-
2003 2/	2,564	-	-	-	10	110	392	565	549	371	370	197	-
2002	2,580	-	-	-	10	129	417	581	480	399	370	194	-
2001	2,725	-	-	-	8	125	448	632	478	392	409	233	-
2000	2,725	-	-	-	9	141	403	600	479	438	434	221	-
1999	2,854	-	-	-	7	147	451	630	508	454	414	242	1
Non-Hispanic white, male													
2005 2/	...	...	...	...	...	...	...	...	...	...	...	...	...
2004 2/	...	...	...	...	...	...	...	...	...	...	...	...	...
2003 2/	...	...	...	...	...	...	...	...	...	...	...	...	...
2002	...	...	...	...	...	...	...	...	...	...	...	...	...
2001	...	...	...	...	...	...	...	...	...	...	...	...	...
2000	...	...	...	...	...	...	...	...	...	...	...	...	...
1999	...	...	...	...	...	...	...	...	...	...	...	...	...
Non-Hispanic white, female													
2005 2/	2,538	-	-	-	7	113	369	595	506	378	367	203	-
2004 2/	2,474	-	-	-	11	128	372	539	540	385	335	164	-
2003 2/	2,564	-	-	-	10	110	392	565	549	371	370	197	-
2002	2,580	-	-	-	10	129	417	581	480	399	370	194	-
2001	2,725	-	-	-	8	125	448	632	478	392	409	233	-
2000	2,725	-	-	-	9	141	403	600	479	438	434	221	-
1999	2,854	-	-	-	7	147	451	630	508	454	414	242	1

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Malignant neoplasm of cervix uteri (C53)													
Non-Hispanic black, both sexes													
2005 2/	776	-	-	-	4	29	132	188	156	112	106	49	-
2004 2/	782	-	-	-	4	43	125	182	153	130	88	57	-
2003 2/	788	-	-	-	2	49	126	173	152	125	111	50	-
2002	820	-	-	-	3	43	144	188	131	152	101	58	-
2001	778	-	-	1	1	42	137	165	131	129	111	61	-
2000	860	-	-	1	4	51	147	177	152	158	108	62	-
1999	825	-	-	-	3	34	140	140	167	173	101	67	-
Non-Hispanic black, male													
2005 2/	...	...	...	...	...	...	...	...	...	...	...	...	...
2004 2/	...	...	...	...	...	...	...	...	...	...	...	...	...
2003 2/	...	...	...	...	...	...	...	...	...	...	...	...	...
2002	...	...	...	...	...	...	...	...	...	...	...	...	...
2001	...	...	...	...	...	...	...	...	...	...	...	...	...
2000	...	...	...	...	...	...	...	...	...	...	...	...	...
1999	...	...	...	...	...	...	...	...	...	...	...	...	...
Non-Hispanic black, female													
2005 2/	776	-	-	-	4	29	132	188	156	112	106	49	-
2004 2/	782	-	-	-	4	43	125	182	153	130	88	57	-
2003 2/	788	-	-	-	2	49	126	173	152	125	111	50	-
2002	820	-	-	-	3	43	144	188	131	152	101	58	-
2001	778	-	-	1	1	42	137	165	131	129	111	61	-
2000	860	-	-	1	4	51	147	177	152	158	108	62	-
1999	825	-	-	-	3	34	140	140	167	173	101	67	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Malignant neoplasm of cervix uteri (C53)													
Origin not stated 3/, both sexes													
2005	8	-	-	-	-	1	1	1	2	2	-	1	-
2004	5	-	-	-	-	-	2	1	-	-	2	-	-
2003	9	-	-	-	-	-	2	4	1	2	-	-	-
2002	13	-	-	-	-	1	3	2	4	2	1	-	-
2001	11	-	-	-	-	1	3	5	2	-	-	-	-
2000	11	-	-	-	-	-	3	3	2	1	2	-	-
1999	10	-	-	-	-	-	1	2	3	1	2	1	-
Origin not stated 3/, male													
2005	...	...	...	...	...	...	...	...	...	...	...	...	...
2004	...	...	...	...	...	...	...	...	...	...	...	...	...
2003	...	...	...	...	...	...	...	...	...	...	...	...	...
2002	...	...	...	...	...	...	...	...	...	...	...	...	...
2001	...	...	...	...	...	...	...	...	...	...	...	...	...
2000	...	...	...	...	...	...	...	...	...	...	...	...	...
1999	...	...	...	...	...	...	...	...	...	...	...	...	...
Origin not stated 3/, female													
2005	8	-	-	-	-	1	1	1	2	2	-	1	-
2004	5	-	-	-	-	-	2	1	-	-	2	-	-
2003	9	-	-	-	-	-	2	4	1	2	-	-	-
2002	13	-	-	-	-	1	3	2	4	2	1	-	-
2001	11	-	-	-	-	1	3	5	2	-	-	-	-
2000	11	-	-	-	-	-	3	3	2	1	2	-	-
1999	10	-	-	-	-	-	1	2	3	1	2	1	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Malignant neoplasms of corpus uteri and uterus, part unspecified (C54-C55)													
All origins, both sexes													
2005	7,096	-	-	-	4	22	146	568	1,468	1,808	1,955	1,125	-
2004	6,990	-	-	-	5	20	168	557	1,290	1,856	2,011	1,083	-
2003	6,899	-	-	-	4	27	154	534	1,301	1,851	1,951	1,077	-
2002	6,853	-	-	-	-	23	146	565	1,199	1,835	1,957	1,128	-
2001	6,784	-	-	-	-	38	139	566	1,105	1,784	2,063	1,088	1
2000	6,586	-	-	-	3	27	138	507	1,122	1,781	1,958	1,049	1
1999	6,468	-	-	-	2	23	142	468	1,019	1,807	2,008	999	-
All origins, male													
2005	...	...	...	...	...	...	...	...	...	...	...	...	...
2004	...	...	...	...	...	...	...	...	...	...	...	...	...
2003	...	...	...	...	...	...	...	...	...	...	...	...	...
2002	...	...	...	...	...	...	...	...	...	...	...	...	...
2001	...	...	...	...	...	...	...	...	...	...	...	...	...
2000	...	...	...	...	...	...	...	...	...	...	...	...	...
1999	...	...	...	...	...	...	...	...	...	...	...	...	...
All origins, female													
2005	7,096	-	-	-	4	22	146	568	1,468	1,808	1,955	1,125	-
2004	6,990	-	-	-	5	20	168	557	1,290	1,856	2,011	1,083	-
2003	6,899	-	-	-	4	27	154	534	1,301	1,851	1,951	1,077	-
2002	6,853	-	-	-	-	23	146	565	1,199	1,835	1,957	1,128	-
2001	6,784	-	-	-	-	38	139	566	1,105	1,784	2,063	1,088	1
2000	6,586	-	-	-	3	27	138	507	1,122	1,781	1,958	1,049	1
1999	6,468	-	-	-	2	23	142	468	1,019	1,807	2,008	999	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Malignant neoplasms of corpus uteri and uterus, part unspecified (C54-C55)													
Hispanic, both sexes													
2005	389	-	-	-	-	2	25	65	81	103	87	26	-
2004	325	-	-	-	-	3	17	35	78	100	72	20	-
2003	321	-	-	-	1	3	24	44	61	85	63	40	-
2002	337	-	-	-	-	5	20	39	64	106	66	37	-
2001	319	-	-	-	-	7	13	48	71	85	65	30	-
2000	277	-	-	-	-	2	19	35	62	75	54	30	-
1999	272	-	-	-	1	3	15	33	61	86	51	22	-
Hispanic, male													
2005	...	...	...	...	...	...	...	...	...	...	...	...	...
2004	...	...	...	...	...	...	...	...	...	...	...	...	...
2003	...	...	...	...	...	...	...	...	...	...	...	...	...
2002	...	...	...	...	...	...	...	...	...	...	...	...	...
2001	...	...	...	...	...	...	...	...	...	...	...	...	...
2000	...	...	...	...	...	...	...	...	...	...	...	...	...
1999	...	...	...	...	...	...	...	...	...	...	...	...	...
Hispanic, female													
2005	389	-	-	-	-	2	25	65	81	103	87	26	-
2004	325	-	-	-	-	3	17	35	78	100	72	20	-
2003	321	-	-	-	1	3	24	44	61	85	63	40	-
2002	337	-	-	-	-	5	20	39	64	106	66	37	-
2001	319	-	-	-	-	7	13	48	71	85	65	30	-
2000	277	-	-	-	-	2	19	35	62	75	54	30	-
1999	272	-	-	-	1	3	15	33	61	86	51	22	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Malignant neoplasms of corpus uteri and uterus, part unspecified (C54-C55)													
Non-Hispanic 1/, both sexes													
2005	6,693	-	-	-	4	20	121	502	1,380	1,703	1,866	1,097	-
2004	6,654	-	-	-	5	17	151	519	1,210	1,753	1,938	1,061	-
2003	6,561	-	-	-	3	24	130	488	1,237	1,761	1,881	1,037	-
2002	6,487	-	-	-	-	17	126	524	1,124	1,725	1,884	1,087	-
2001	6,449	-	-	-	-	31	126	517	1,029	1,697	1,996	1,052	1
2000	6,294	-	-	-	3	25	119	469	1,057	1,702	1,901	1,017	1
1999	6,188	-	-	-	1	20	127	435	956	1,719	1,954	976	-
Non-Hispanic 1/, male													
2005	...	...	...	...	...	...	...	...	...	...	...	...	...
2004	...	...	...	...	...	...	...	...	...	...	...	...	...
2003	...	...	...	...	...	...	...	...	...	...	...	...	...
2002	...	...	...	...	...	...	...	...	...	...	...	...	...
2001	...	...	...	...	...	...	...	...	...	...	...	...	...
2000	...	...	...	...	...	...	...	...	...	...	...	...	...
1999	...	...	...	...	...	...	...	...	...	...	...	...	...
Non-Hispanic 1/, female													
2005	6,693	-	-	-	4	20	121	502	1,380	1,703	1,866	1,097	-
2004	6,654	-	-	-	5	17	151	519	1,210	1,753	1,938	1,061	-
2003	6,561	-	-	-	3	24	130	488	1,237	1,761	1,881	1,037	-
2002	6,487	-	-	-	-	17	126	524	1,124	1,725	1,884	1,087	-
2001	6,449	-	-	-	-	31	126	517	1,029	1,697	1,996	1,052	1
2000	6,294	-	-	-	3	25	119	469	1,057	1,702	1,901	1,017	1
1999	6,188	-	-	-	1	20	127	435	956	1,719	1,954	976	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Malignant neoplasms of corpus uteri and uterus, part unspecified (C54-C55)													
Non-Hispanic white, both sexes													
2005 2/	5,360	-	-	-	2	11	86	387	1,035	1,313	1,557	969	-
2004 2/	5,392	-	-	-	3	13	115	397	934	1,314	1,654	962	-
2003 2/	5,321	-	-	-	2	13	86	386	955	1,366	1,584	929	-
2002	5,213	-	-	-	-	11	97	403	861	1,312	1,580	949	-
2001	5,283	-	-	-	-	19	94	398	802	1,319	1,712	939	-
2000	5,113	-	-	-	1	18	88	383	827	1,298	1,594	904	-
1999	5,114	-	-	-	1	15	96	353	753	1,351	1,687	858	-
Non-Hispanic white, male													
2005 2/	...	...	...	...	...	...	...	...	...	...	...	...	...
2004 2/	...	...	...	...	...	...	...	...	...	...	...	...	...
2003 2/	...	...	...	...	...	...	...	...	...	...	...	...	...
2002	...	...	...	...	...	...	...	...	...	...	...	...	...
2001	...	...	...	...	...	...	...	...	...	...	...	...	...
2000	...	...	...	...	...	...	...	...	...	...	...	...	...
1999	...	...	...	...	...	...	...	...	...	...	...	...	...
Non-Hispanic white, female													
2005 2/	5,360	-	-	-	2	11	86	387	1,035	1,313	1,557	969	-
2004 2/	5,392	-	-	-	3	13	115	397	934	1,314	1,654	962	-
2003 2/	5,321	-	-	-	2	13	86	386	955	1,366	1,584	929	-
2002	5,213	-	-	-	-	11	97	403	861	1,312	1,580	949	-
2001	5,283	-	-	-	-	19	94	398	802	1,319	1,712	939	-
2000	5,113	-	-	-	1	18	88	383	827	1,298	1,594	904	-
1999	5,114	-	-	-	1	15	96	353	753	1,351	1,687	858	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Malignant neoplasms of corpus uteri and uterus, part unspecified (C54-C55)													
Non-Hispanic black, both sexes													
2005 2/	1,165	-	-	-	2	7	24	93	298	353	271	117	-
2004 2/	1,102	-	-	-	2	4	29	93	229	395	258	92	-
2003 2/	1,092	-	-	-	1	8	30	80	245	357	269	102	-
2002	1,107	-	-	-	-	5	21	88	221	373	268	131	-
2001	1,037	-	-	-	-	9	22	93	190	352	265	105	1
2000	1,055	-	-	-	1	6	21	68	201	371	280	106	1
1999	980	-	-	-	-	5	27	57	189	346	245	111	-
Non-Hispanic black, male													
2005 2/	...	...	...	...	...	...	...	...	...	...	...	...	...
2004 2/	...	...	...	...	...	...	...	...	...	...	...	...	...
2003 2/	...	...	...	...	...	...	...	...	...	...	...	...	...
2002	...	...	...	...	...	...	...	...	...	...	...	...	...
2001	...	...	...	...	...	...	...	...	...	...	...	...	...
2000	...	...	...	...	...	...	...	...	...	...	...	...	...
1999	...	...	...	...	...	...	...	...	...	...	...	...	...
Non-Hispanic black, female													
2005 2/	1,165	-	-	-	2	7	24	93	298	353	271	117	-
2004 2/	1,102	-	-	-	2	4	29	93	229	395	258	92	-
2003 2/	1,092	-	-	-	1	8	30	80	245	357	269	102	-
2002	1,107	-	-	-	-	5	21	88	221	373	268	131	-
2001	1,037	-	-	-	-	9	22	93	190	352	265	105	1
2000	1,055	-	-	-	1	6	21	68	201	371	280	106	1
1999	980	-	-	-	-	5	27	57	189	346	245	111	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Malignant neoplasms of corpus uteri and uterus, part unspecified (C54-C55)													
Origin not stated 3/, both sexes													
2005	14	-	-	-	-	-	-	1	7	2	2	2	-
2004	11	-	-	-	-	-	-	3	2	3	1	2	-
2003	17	-	-	-	-	-	-	2	3	5	7	-	-
2002	29	-	-	-	-	1	-	2	11	4	7	4	-
2001	16	-	-	-	-	-	-	1	5	2	2	6	-
2000	15	-	-	-	-	-	-	3	3	4	3	2	-
1999	8	-	-	-	-	-	-	-	2	2	3	1	-
Origin not stated 3/, male													
2005	...	...	...	...	...	...	...	...	...	...	...	...	...
2004	...	...	...	...	...	...	...	...	...	...	...	...	...
2003	...	...	...	...	...	...	...	...	...	...	...	...	...
2002	...	...	...	...	...	...	...	...	...	...	...	...	...
2001	...	...	...	...	...	...	...	...	...	...	...	...	...
2000	...	...	...	...	...	...	...	...	...	...	...	...	...
1999	...	...	...	...	...	...	...	...	...	...	...	...	...
Origin not stated 3/, female													
2005	14	-	-	-	-	-	-	1	7	2	2	2	-
2004	11	-	-	-	-	-	-	3	2	3	1	2	-
2003	17	-	-	-	-	-	-	2	3	5	7	-	-
2002	29	-	-	-	-	1	-	2	11	4	7	4	-
2001	16	-	-	-	-	-	-	1	5	2	2	6	-
2000	15	-	-	-	-	-	-	3	3	4	3	2	-
1999	8	-	-	-	-	-	-	-	2	2	3	1	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Malignant neoplasm of ovary (C56)													
All origins, both sexes													
2005	14,787	-	-	3	18	83	413	1,682	3,048	3,569	4,031	1,940	-
2004	14,716	-	-	2	29	94	454	1,654	2,794	3,693	4,165	1,831	-
2003	14,657	-	-	-	28	84	452	1,629	2,891	3,569	4,194	1,810	-
2002	14,682	1	1	2	37	97	469	1,717	2,748	3,601	4,259	1,750	-
2001	14,414	-	-	2	21	102	472	1,626	2,612	3,666	4,170	1,743	-
2000	14,060	-	-	2	23	101	472	1,583	2,444	3,601	4,098	1,736	-
1999	13,627	-	1	2	27	77	496	1,549	2,426	3,612	3,875	1,562	-
All origins, male													
2005	...	...	...	...	...	...	...	...	...	...	...	...	...
2004	...	...	...	...	...	...	...	...	...	...	...	...	...
2003	...	...	...	...	...	...	...	...	...	...	...	...	...
2002	...	...	...	...	...	...	...	...	...	...	...	...	...
2001	...	...	...	...	...	...	...	...	...	...	...	...	...
2000	...	...	...	...	...	...	...	...	...	...	...	...	...
1999	...	...	...	...	...	...	...	...	...	...	...	...	...
All origins, female													
2005	14,787	-	-	3	18	83	413	1,682	3,048	3,569	4,031	1,940	-
2004	14,716	-	-	2	29	94	454	1,654	2,794	3,693	4,165	1,831	-
2003	14,657	-	-	-	28	84	452	1,629	2,891	3,569	4,194	1,810	-
2002	14,682	1	1	2	37	97	469	1,717	2,748	3,601	4,259	1,750	-
2001	14,414	-	-	2	21	102	472	1,626	2,612	3,666	4,170	1,743	-
2000	14,060	-	-	2	23	101	472	1,583	2,444	3,601	4,098	1,736	-
1999	13,627	-	1	2	27	77	496	1,549	2,426	3,612	3,875	1,562	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Malignant neoplasm of ovary (C56)													
Hispanic, both sexes													
2005	739	-	-	-	3	12	41	132	169	173	145	64	-
2004	689	-	-	1	4	11	51	111	145	164	143	59	-
2003	645	-	-	-	3	5	48	104	134	155	141	55	-
2002	622	-	-	2	3	10	47	116	111	145	145	43	-
2001	603	-	-	1	-	13	43	91	131	164	111	49	-
2000	578	-	-	1	4	9	30	94	111	154	126	49	-
1999	478	-	1	-	2	6	29	80	100	126	100	34	-
Hispanic, male													
2005	...	...	...	...	...	...	...	...	...	...	...	...	...
2004	...	...	...	...	...	...	...	...	...	...	...	...	...
2003	...	...	...	...	...	...	...	...	...	...	...	...	...
2002	...	...	...	...	...	...	...	...	...	...	...	...	...
2001	...	...	...	...	...	...	...	...	...	...	...	...	...
2000	...	...	...	...	...	...	...	...	...	...	...	...	...
1999	...	...	...	...	...	...	...	...	...	...	...	...	...
Hispanic, female													
2005	739	-	-	-	3	12	41	132	169	173	145	64	-
2004	689	-	-	1	4	11	51	111	145	164	143	59	-
2003	645	-	-	-	3	5	48	104	134	155	141	55	-
2002	622	-	-	2	3	10	47	116	111	145	145	43	-
2001	603	-	-	1	-	13	43	91	131	164	111	49	-
2000	578	-	-	1	4	9	30	94	111	154	126	49	-
1999	478	-	1	-	2	6	29	80	100	126	100	34	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Malignant neoplasm of ovary (C56)													
Non-Hispanic 1/, both sexes													
2005	14,041	-	-	3	15	71	372	1,549	2,877	3,394	3,884	1,876	-
2004	14,008	-	-	1	25	83	402	1,541	2,644	3,522	4,020	1,770	-
2003	13,985	-	-	-	25	79	403	1,519	2,749	3,406	4,050	1,754	-
2002	14,031	1	1	-	34	87	421	1,596	2,633	3,445	4,107	1,706	-
2001	13,783	-	-	1	21	89	428	1,533	2,475	3,495	4,052	1,689	-
2000	13,454	-	-	1	19	92	441	1,484	2,329	3,440	3,964	1,684	-
1999	13,103	-	-	2	25	71	466	1,466	2,316	3,474	3,756	1,527	-
Non-Hispanic 1/, male													
2005	...	...	...	...	...	...	...	...	...	...	...	...	...
2004	...	...	...	...	...	...	...	...	...	...	...	...	...
2003	...	...	...	...	...	...	...	...	...	...	...	...	...
2002	...	...	...	...	...	...	...	...	...	...	...	...	...
2001	...	...	...	...	...	...	...	...	...	...	...	...	...
2000	...	...	...	...	...	...	...	...	...	...	...	...	...
1999	...	...	...	...	...	...	...	...	...	...	...	...	...
Non-Hispanic 1/, female													
2005	14,041	-	-	3	15	71	372	1,549	2,877	3,394	3,884	1,876	-
2004	14,008	-	-	1	25	83	402	1,541	2,644	3,522	4,020	1,770	-
2003	13,985	-	-	-	25	79	403	1,519	2,749	3,406	4,050	1,754	-
2002	14,031	1	1	-	34	87	421	1,596	2,633	3,445	4,107	1,706	-
2001	13,783	-	-	1	21	89	428	1,533	2,475	3,495	4,052	1,689	-
2000	13,454	-	-	1	19	92	441	1,484	2,329	3,440	3,964	1,684	-
1999	13,103	-	-	2	25	71	466	1,466	2,316	3,474	3,756	1,527	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Malignant neoplasm of ovary (C56)													
Non-Hispanic white, both sexes													
2005 2/	12,472	-	-	2	15	56	293	1,316	2,529	2,980	3,538	1,743	-
2004 2/	12,432	-	-	1	16	60	326	1,280	2,312	3,103	3,701	1,633	-
2003 2/	12,477	-	-	-	18	64	329	1,282	2,432	3,024	3,720	1,608	-
2002	12,560	1	1	-	27	67	343	1,369	2,329	3,082	3,766	1,575	-
2001	12,338	-	-	-	14	68	347	1,335	2,151	3,125	3,741	1,557	-
2000	12,138	-	-	-	13	75	367	1,284	2,072	3,087	3,680	1,560	-
1999	11,795	-	-	2	20	51	394	1,268	2,061	3,119	3,456	1,424	-
Non-Hispanic white, male													
2005 2/	...	...	...	...	...	...	...	...	...	...	...	...	...
2004 2/	...	...	...	...	...	...	...	...	...	...	...	...	...
2003 2/	...	...	...	...	...	...	...	...	...	...	...	...	...
2002	...	...	...	...	...	...	...	...	...	...	...	...	...
2001	...	...	...	...	...	...	...	...	...	...	...	...	...
2000	...	...	...	...	...	...	...	...	...	...	...	...	...
1999	...	...	...	...	...	...	...	...	...	...	...	...	...
Non-Hispanic white, female													
2005 2/	12,472	-	-	2	15	56	293	1,316	2,529	2,980	3,538	1,743	-
2004 2/	12,432	-	-	1	16	60	326	1,280	2,312	3,103	3,701	1,633	-
2003 2/	12,477	-	-	-	18	64	329	1,282	2,432	3,024	3,720	1,608	-
2002	12,560	1	1	-	27	67	343	1,369	2,329	3,082	3,766	1,575	-
2001	12,338	-	-	-	14	68	347	1,335	2,151	3,125	3,741	1,557	-
2000	12,138	-	-	-	13	75	367	1,284	2,072	3,087	3,680	1,560	-
1999	11,795	-	-	2	20	51	394	1,268	2,061	3,119	3,456	1,424	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Malignant neoplasm of ovary (C56)													
Non-Hispanic black, both sexes													
2005 2/	1,204	-	-	1	-	14	54	158	261	324	278	114	-
2004 2/	1,247	-	-	-	7	17	52	192	248	346	272	113	-
2003 2/	1,180	-	-	-	6	12	56	181	236	297	267	125	-
2002	1,155	-	-	-	7	13	59	158	237	287	277	117	-
2001	1,140	-	-	1	4	15	65	134	251	301	252	117	-
2000	1,065	-	-	1	2	12	59	138	205	295	244	109	-
1999	1,065	-	-	-	4	13	47	143	211	301	256	90	-
Non-Hispanic black, male													
2005 2/	...	...	...	...	...	...	...	...	...	...	...	...	...
2004 2/	...	...	...	...	...	...	...	...	...	...	...	...	...
2003 2/	...	...	...	...	...	...	...	...	...	...	...	...	...
2002	...	...	...	...	...	...	...	...	...	...	...	...	...
2001	...	...	...	...	...	...	...	...	...	...	...	...	...
2000	...	...	...	...	...	...	...	...	...	...	...	...	...
1999	...	...	...	...	...	...	...	...	...	...	...	...	...
Non-Hispanic black, female													
2005 2/	1,204	-	-	1	-	14	54	158	261	324	278	114	-
2004 2/	1,247	-	-	-	7	17	52	192	248	346	272	113	-
2003 2/	1,180	-	-	-	6	12	56	181	236	297	267	125	-
2002	1,155	-	-	-	7	13	59	158	237	287	277	117	-
2001	1,140	-	-	1	4	15	65	134	251	301	252	117	-
2000	1,065	-	-	1	2	12	59	138	205	295	244	109	-
1999	1,065	-	-	-	4	13	47	143	211	301	256	90	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Malignant neoplasm of ovary (C56)													
Origin not stated 3/, both sexes													
2005	7	-	-	-	-	-	-	1	2	2	2	-	-
2004	19	-	-	-	-	-	1	2	5	7	2	2	-
2003	27	-	-	-	-	-	1	6	8	8	3	1	-
2002	29	-	-	-	-	-	1	5	4	11	7	1	-
2001	28	-	-	-	-	-	1	2	6	7	7	5	-
2000	28	-	-	-	-	-	1	5	4	7	8	3	-
1999	46	-	-	-	-	-	1	3	10	12	19	1	-
Origin not stated 3/, male													
2005	...	...	...	...	...	...	...	...	...	...	...	...	...
2004	...	...	...	...	...	...	...	...	...	...	...	...	...
2003	...	...	...	...	...	...	...	...	...	...	...	...	...
2002	...	...	...	...	...	...	...	...	...	...	...	...	...
2001	...	...	...	...	...	...	...	...	...	...	...	...	...
2000	...	...	...	...	...	...	...	...	...	...	...	...	...
1999	...	...	...	...	...	...	...	...	...	...	...	...	...
Origin not stated 3/, female													
2005	7	-	-	-	-	-	-	1	2	2	2	-	-
2004	19	-	-	-	-	-	1	2	5	7	2	2	-
2003	27	-	-	-	-	-	1	6	8	8	3	1	-
2002	29	-	-	-	-	-	1	5	4	11	7	1	-
2001	28	-	-	-	-	-	1	2	6	7	7	5	-
2000	28	-	-	-	-	-	1	5	4	7	8	3	-
1999	46	-	-	-	-	-	1	3	10	12	19	1	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Malignant neoplasm of prostate (C61)													
All origins, both sexes													
2005	28,905	-	1	-	1	3	24	395	2,154	5,764	11,666	8,897	-
2004	29,004	-	1	1	1	2	23	434	2,044	5,711	11,993	8,792	2
2003	29,554	-	-	1	1	1	25	418	2,074	6,033	12,284	8,717	-
2002	30,446	-	-	-	1	1	32	382	2,020	6,300	12,753	8,957	-
2001	30,719	-	-	-	1	1	30	412	1,907	6,543	12,976	8,849	-
2000	31,078	-	-	-	-	1	25	371	1,929	6,808	13,071	8,873	-
1999	31,729	1	-	-	1	2	33	374	1,946	7,316	13,435	8,620	1
All origins, male													
2005	28,905	-	1	-	1	3	24	395	2,154	5,764	11,666	8,897	-
2004	29,004	-	1	1	1	2	23	434	2,044	5,711	11,993	8,792	2
2003	29,554	-	-	1	1	1	25	418	2,074	6,033	12,284	8,717	-
2002	30,446	-	-	-	1	1	32	382	2,020	6,300	12,753	8,957	-
2001	30,719	-	-	-	1	1	30	412	1,907	6,543	12,976	8,849	-
2000	31,078	-	-	-	-	1	25	371	1,929	6,808	13,071	8,873	-
1999	31,729	1	-	-	1	2	33	374	1,946	7,316	13,435	8,620	1
All origins, female													
2005	...	...	...	...	...	...	...	...	...	...	...	...	...
2004	...	...	...	...	...	...	...	...	...	...	...	...	...
2003	...	...	...	...	...	...	...	...	...	...	...	...	...
2002	...	...	...	...	...	...	...	...	...	...	...	...	...
2001	...	...	...	...	...	...	...	...	...	...	...	...	...
2000	...	...	...	...	...	...	...	...	...	...	...	...	...
1999	...	...	...	...	...	...	...	...	...	...	...	...	...

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Malignant neoplasm of prostate (C61)													
Hispanic, both sexes													
2005	1,300	-	1	-	-	-	3	25	125	319	494	333	-
2004	1,239	-	-	-	-	-	-	25	109	284	546	274	1
2003	1,219	-	-	-	-	-	1	25	111	312	477	293	-
2002	1,209	-	-	-	-	1	1	29	106	292	472	308	-
2001	1,226	-	-	-	-	-	-	27	108	332	487	272	-
2000	1,033	-	-	-	-	-	1	26	78	277	401	250	-
1999	1,050	-	-	-	1	1	2	22	92	295	386	251	-
Hispanic, male													
2005	1,300	-	1	-	-	-	3	25	125	319	494	333	-
2004	1,239	-	-	-	-	-	-	25	109	284	546	274	1
2003	1,219	-	-	-	-	-	1	25	111	312	477	293	-
2002	1,209	-	-	-	-	1	1	29	106	292	472	308	-
2001	1,226	-	-	-	-	-	-	27	108	332	487	272	-
2000	1,033	-	-	-	-	-	1	26	78	277	401	250	-
1999	1,050	-	-	-	1	1	2	22	92	295	386	251	-
Hispanic, female													
2005	...	...	...	...	...	...	...	...	...	...	...	...	...
2004	...	...	...	...	...	...	...	...	...	...	...	...	...
2003	...	...	...	...	...	...	...	...	...	...	...	...	...
2002	...	...	...	...	...	...	...	...	...	...	...	...	...
2001	...	...	...	...	...	...	...	...	...	...	...	...	...
2000	...	...	...	...	...	...	...	...	...	...	...	...	...
1999	...	...	...	...	...	...	...	...	...	...	...	...	...

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Malignant neoplasm of prostate (C61)													
Non-Hispanic 1/, both sexes													
2005	27,566	-	-	-	1	3	21	370	2,025	5,430	11,158	8,558	-
2004	27,723	-	1	1	1	2	23	409	1,931	5,418	11,430	8,507	-
2003	28,279	-	-	1	1	1	24	391	1,959	5,705	11,787	8,410	-
2002	29,151	-	-	-	1	-	31	351	1,906	5,983	12,251	8,628	-
2001	29,407	-	-	-	1	1	29	385	1,790	6,197	12,449	8,555	-
2000	29,964	-	-	-	-	1	24	344	1,846	6,512	12,634	8,603	-
1999	30,614	1	-	-	-	1	31	351	1,848	7,004	13,025	8,352	1
Non-Hispanic 1/, male													
2005	27,566	-	-	-	1	3	21	370	2,025	5,430	11,158	8,558	-
2004	27,723	-	1	1	1	2	23	409	1,931	5,418	11,430	8,507	-
2003	28,279	-	-	1	1	1	24	391	1,959	5,705	11,787	8,410	-
2002	29,151	-	-	-	1	-	31	351	1,906	5,983	12,251	8,628	-
2001	29,407	-	-	-	1	1	29	385	1,790	6,197	12,449	8,555	-
2000	29,964	-	-	-	-	1	24	344	1,846	6,512	12,634	8,603	-
1999	30,614	1	-	-	-	1	31	351	1,848	7,004	13,025	8,352	1
Non-Hispanic 1/, female													
2005	...	...	...	...	...	...	...	...	...	...	...	...	...
2004	...	...	...	...	...	...	...	...	...	...	...	...	...
2003	...	...	...	...	...	...	...	...	...	...	...	...	...
2002	...	...	...	...	...	...	...	...	...	...	...	...	...
2001	...	...	...	...	...	...	...	...	...	...	...	...	...
2000	...	...	...	...	...	...	...	...	...	...	...	...	...
1999	...	...	...	...	...	...	...	...	...	...	...	...	...

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Malignant neoplasm of prostate (C61)													
Non-Hispanic white, both sexes													
2005 2/	22,319	-	-	-	1	2	10	241	1,442	4,086	9,217	7,320	-
2004 2/	22,509	-	1	1	1	2	17	264	1,411	4,147	9,388	7,277	-
2003 2/	23,015	-	-	1	1	1	16	264	1,390	4,383	9,700	7,259	-
2002	23,690	-	-	-	-	-	24	226	1,361	4,561	10,091	7,427	-
2001	23,811	-	-	-	1	-	18	258	1,285	4,755	10,230	7,264	-
2000	24,287	-	-	-	-	1	15	237	1,325	4,946	10,399	7,364	-
1999	24,936	1	-	-	-	1	23	244	1,310	5,413	10,850	7,093	1
Non-Hispanic white, male													
2005 2/	22,319	-	-	-	1	2	10	241	1,442	4,086	9,217	7,320	-
2004 2/	22,509	-	1	1	1	2	17	264	1,411	4,147	9,388	7,277	-
2003 2/	23,015	-	-	1	1	1	16	264	1,390	4,383	9,700	7,259	-
2002	23,690	-	-	-	-	-	24	226	1,361	4,561	10,091	7,427	-
2001	23,811	-	-	-	1	-	18	258	1,285	4,755	10,230	7,264	-
2000	24,287	-	-	-	-	1	15	237	1,325	4,946	10,399	7,364	-
1999	24,936	1	-	-	-	1	23	244	1,310	5,413	10,850	7,093	1
Non-Hispanic white, female													
2005 2/	...	...	...	...	...	...	...	...	...	...	...	...	...
2004 2/	...	...	...	...	...	...	...	...	...	...	...	...	...
2003 2/	...	...	...	...	...	...	...	...	...	...	...	...	...
2002	...	...	...	...	...	...	...	...	...	...	...	...	...
2001	...	...	...	...	...	...	...	...	...	...	...	...	...
2000	...	...	...	...	...	...	...	...	...	...	...	...	...
1999	...	...	...	...	...	...	...	...	...	...	...	...	...

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Malignant neoplasm of prostate (C61)													
Non-Hispanic black, both sexes													
2005 2/	4,771	-	-	-	-	-	10	119	542	1,249	1,754	1,097	-
2004 2/	4,768	-	-	-	-	-	4	140	493	1,187	1,854	1,090	-
2003 2/	4,845	-	-	-	-	-	8	123	539	1,242	1,909	1,024	-
2002	5,085	-	-	-	1	-	7	118	500	1,352	2,005	1,102	-
2001	5,215	-	-	-	-	1	11	124	481	1,372	2,054	1,172	-
2000	5,292	-	-	-	-	-	9	100	494	1,481	2,077	1,131	-
1999	5,315	-	-	-	-	-	7	105	512	1,524	2,027	1,140	-
Non-Hispanic black, male													
2005 2/	4,771	-	-	-	-	-	10	119	542	1,249	1,754	1,097	-
2004 2/	4,768	-	-	-	-	-	4	140	493	1,187	1,854	1,090	-
2003 2/	4,845	-	-	-	-	-	8	123	539	1,242	1,909	1,024	-
2002	5,085	-	-	-	1	-	7	118	500	1,352	2,005	1,102	-
2001	5,215	-	-	-	-	1	11	124	481	1,372	2,054	1,172	-
2000	5,292	-	-	-	-	-	9	100	494	1,481	2,077	1,131	-
1999	5,315	-	-	-	-	-	7	105	512	1,524	2,027	1,140	-
Non-Hispanic black, female													
2005 2/	...	...	...	...	...	...	...	...	...	...	...	...	...
2004 2/	...	...	...	...	...	...	...	...	...	...	...	...	...
2003 2/	...	...	...	...	...	...	...	...	...	...	...	...	...
2002	...	...	...	...	...	...	...	...	...	...	...	...	...
2001	...	...	...	...	...	...	...	...	...	...	...	...	...
2000	...	...	...	...	...	...	...	...	...	...	...	...	...
1999	...	...	...	...	...	...	...	...	...	...	...	...	...

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Malignant neoplasm of prostate (C61)													
Origin not stated 3/, both sexes													
2005	39	-	-	-	-	-	-	-	4	15	14	6	-
2004	42	-	-	-	-	-	-	-	4	9	17	11	1
2003	56	-	-	-	-	-	-	2	4	16	20	14	-
2002	86	-	-	-	-	-	-	2	8	25	30	21	-
2001	86	-	-	-	-	-	1	-	9	14	40	22	-
2000	81	-	-	-	-	-	-	1	5	19	36	20	-
1999	65	-	-	-	-	-	-	1	6	17	24	17	-
Origin not stated 3/, male													
2005	39	-	-	-	-	-	-	-	4	15	14	6	-
2004	42	-	-	-	-	-	-	-	4	9	17	11	1
2003	56	-	-	-	-	-	-	2	4	16	20	14	-
2002	86	-	-	-	-	-	-	2	8	25	30	21	-
2001	86	-	-	-	-	-	1	-	9	14	40	22	-
2000	81	-	-	-	-	-	-	1	5	19	36	20	-
1999	65	-	-	-	-	-	-	1	6	17	24	17	-
Origin not stated 3/, female													
2005	...	...	...	...	...	...	...	...	...	...	...	...	...
2004	...	...	...	...	...	...	...	...	...	...	...	...	...
2003	...	...	...	...	...	...	...	...	...	...	...	...	...
2002	...	...	...	...	...	...	...	...	...	...	...	...	...
2001	...	...	...	...	...	...	...	...	...	...	...	...	...
2000	...	...	...	...	...	...	...	...	...	...	...	...	...
1999	...	...	...	...	...	...	...	...	...	...	...	...	...

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Malignant neoplasms of kidney and renal pelvis (C64-C65)													
All origins, both sexes													
2005	12,517	8	16	30	17	49	294	1,277	2,514	3,143	3,510	1,659	-
2004	12,313	4	18	29	16	49	268	1,256	2,534	3,075	3,474	1,590	-
2003	12,286	1	14	39	24	56	321	1,201	2,498	3,090	3,475	1,567	-
2002	12,165	4	8	30	20	48	306	1,322	2,340	3,239	3,355	1,493	-
2001	12,078	1	22	35	20	52	358	1,311	2,328	3,221	3,306	1,424	-
2000	11,736	2	15	29	20	42	334	1,239	2,195	3,146	3,282	1,432	-
1999	11,116	2	19	33	17	54	342	1,147	2,111	3,076	3,010	1,305	-
All origins, male													
2005	7,783	4	9	13	11	35	181	890	1,752	2,078	1,999	811	-
2004	7,640	2	9	11	9	34	183	889	1,786	2,015	1,993	709	-
2003	7,667	1	4	19	16	36	231	859	1,787	1,975	2,036	703	-
2002	7,587	1	-	19	7	31	217	947	1,658	2,087	1,924	696	-
2001	7,524	1	10	9	16	27	240	941	1,634	2,093	1,866	687	-
2000	7,249	1	8	15	10	26	228	880	1,530	2,010	1,865	676	-
1999	6,867	1	11	16	11	37	236	816	1,474	1,958	1,717	590	-
All origins, female													
2005	4,734	4	7	17	6	14	113	387	762	1,065	1,511	848	-
2004	4,673	2	9	18	7	15	85	367	748	1,060	1,481	881	-
2003	4,619	-	10	20	8	20	90	342	711	1,115	1,439	864	-
2002	4,578	3	8	11	13	17	89	375	682	1,152	1,431	797	-
2001	4,554	-	12	26	4	25	118	370	694	1,128	1,440	737	-
2000	4,487	1	7	14	10	16	106	359	665	1,136	1,417	756	-
1999	4,249	1	8	17	6	17	106	331	637	1,118	1,293	715	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Malignant neoplasms of kidney and renal pelvis (C64-C65)													
Hispanic, both sexes													
2005	780	-	2	4	1	7	42	119	150	207	171	77	-
2004	763	2	3	1	3	7	43	125	182	178	160	59	-
2003	698	-	2	4	6	3	30	82	170	198	151	52	-
2002	659	2	3	2	1	6	37	103	160	152	146	47	-
2001	640	-	6	6	-	8	38	112	127	163	129	51	-
2000	567	-	3	3	3	7	32	88	135	132	125	39	-
1999	547	-	1	4	1	4	31	84	129	144	108	41	-
Hispanic, male													
2005	482	-	1	2	1	5	25	73	106	133	99	37	-
2004	507	1	2	1	2	6	27	90	118	132	97	31	-
2003	426	-	-	2	5	3	23	54	109	121	84	25	-
2002	426	-	-	2	-	4	26	72	106	97	95	24	-
2001	426	-	1	4	-	5	34	80	86	107	79	30	-
2000	372	-	1	3	2	4	27	60	92	85	78	20	-
1999	353	-	1	2	-	2	23	63	91	96	57	18	-
Hispanic, female													
2005	298	-	1	2	-	2	17	46	44	74	72	40	-
2004	256	1	1	-	1	1	16	35	64	46	63	28	-
2003	272	-	2	2	1	-	7	28	61	77	67	27	-
2002	233	2	3	-	1	2	11	31	54	55	51	23	-
2001	214	-	5	2	-	3	4	32	41	56	50	21	-
2000	195	-	2	-	1	3	5	28	43	47	47	19	-
1999	194	-	-	2	1	2	8	21	38	48	51	23	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Malignant neoplasms of kidney and renal pelvis (C64-C65)													
Non-Hispanic 1/, both sexes													
2005	11,719	8	14	25	16	42	252	1,153	2,362	2,931	3,336	1,580	-
2004	11,531	2	15	28	13	42	225	1,130	2,346	2,891	3,310	1,529	-
2003	11,566	1	12	35	17	53	291	1,117	2,325	2,884	3,320	1,511	-
2002	11,477	2	5	28	19	42	268	1,216	2,173	3,079	3,200	1,445	-
2001	11,415	1	16	29	20	44	319	1,196	2,198	3,051	3,172	1,369	-
2000	11,124	2	12	26	17	35	302	1,147	2,053	2,994	3,150	1,386	-
1999	10,542	2	18	29	16	49	309	1,060	1,979	2,927	2,892	1,261	-
Non-Hispanic 1/, male													
2005	7,286	4	8	10	10	30	156	813	1,644	1,940	1,898	773	-
2004	7,121	1	7	10	7	28	156	798	1,665	1,878	1,894	677	-
2003	7,224	1	4	17	10	33	208	804	1,675	1,848	1,948	676	-
2002	7,141	1	-	17	7	27	190	873	1,546	1,984	1,825	671	-
2001	7,084	1	9	5	16	22	205	859	1,546	1,981	1,784	656	-
2000	6,847	1	7	12	8	22	201	818	1,433	1,911	1,782	652	-
1999	6,501	1	10	14	11	35	212	751	1,381	1,858	1,657	571	-
Non-Hispanic 1/, female													
2005	4,433	4	6	15	6	12	96	340	718	991	1,438	807	-
2004	4,410	1	8	18	6	14	69	332	681	1,013	1,416	852	-
2003	4,342	-	8	18	7	20	83	313	650	1,036	1,372	835	-
2002	4,336	1	5	11	12	15	78	343	627	1,095	1,375	774	-
2001	4,331	-	7	24	4	22	114	337	652	1,070	1,388	713	-
2000	4,277	1	5	14	9	13	101	329	620	1,083	1,368	734	-
1999	4,041	1	8	15	5	14	97	309	598	1,069	1,235	690	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Malignant neoplasms of kidney and renal pelvis (C64-C65)													
Non-Hispanic white, both sexes													
2005 2/	10,312	7	10	18	6	17	193	956	2,036	2,598	3,032	1,439	-
2004 2/	10,218	2	10	23	8	22	170	952	2,067	2,551	2,999	1,414	-
2003 2/	10,225	1	12	26	10	32	219	936	2,004	2,570	3,031	1,384	-
2002	10,179	2	5	23	15	24	211	1,011	1,898	2,754	2,903	1,333	-
2001	10,073	-	11	17	8	23	235	998	1,904	2,741	2,880	1,256	-
2000	9,884	-	7	16	11	24	239	977	1,794	2,665	2,871	1,280	-
1999	9,319	2	11	21	9	28	238	877	1,740	2,613	2,636	1,144	-
Non-Hispanic white, male													
2005 2/	6,404	3	7	6	3	12	115	670	1,421	1,722	1,732	713	-
2004 2/	6,329	1	4	8	5	12	120	680	1,477	1,664	1,729	629	-
2003 2/	6,408	1	4	12	6	17	159	682	1,463	1,650	1,787	627	-
2002	6,363	1	-	14	5	15	150	741	1,355	1,785	1,668	629	-
2001	6,263	-	6	1	5	11	157	726	1,340	1,790	1,616	611	-
2000	6,082	-	4	6	6	13	165	701	1,258	1,697	1,628	604	-
1999	5,754	1	7	9	7	18	165	618	1,220	1,662	1,528	519	-
Non-Hispanic white, female													
2005 2/	3,908	4	3	12	3	5	78	286	615	876	1,300	726	-
2004 2/	3,889	1	6	15	3	10	50	272	590	887	1,270	785	-
2003 2/	3,817	-	8	14	4	15	60	254	541	920	1,244	757	-
2002	3,816	1	5	9	10	9	61	270	543	969	1,235	704	-
2001	3,810	-	5	16	3	12	78	272	564	951	1,264	645	-
2000	3,802	-	3	10	5	11	74	276	536	968	1,243	676	-
1999	3,565	1	4	12	2	10	73	259	520	951	1,108	625	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Malignant neoplasms of kidney and renal pelvis (C64-C65)													
Non-Hispanic black, both sexes													
2005 2/	1,141	-	2	7	10	21	48	163	270	266	241	113	-
2004 2/	1,084	-	4	4	5	19	44	151	231	270	256	100	-
2003 2/	1,105	-	-	9	7	19	63	153	261	246	232	115	-
2002	1,062	-	-	5	3	17	45	163	226	260	250	93	-
2001	1,105	-	3	12	10	19	72	163	240	249	241	96	-
2000	1,033	1	5	8	6	10	51	142	213	271	233	93	-
1999	1,028	-	4	6	7	20	61	154	202	263	213	98	-
Non-Hispanic black, male													
2005 2/	713	-	-	4	7	16	35	119	188	174	128	42	-
2004 2/	649	-	3	1	2	15	29	102	155	169	134	39	-
2003 2/	667	-	-	5	4	14	43	107	169	154	129	42	-
2002	626	-	-	3	2	11	31	103	164	151	125	36	-
2001	673	-	1	4	10	10	40	111	168	153	136	40	-
2000	625	-	3	4	2	8	27	94	142	179	124	42	-
1999	618	-	2	3	4	16	42	109	137	159	104	42	-
Non-Hispanic black, female													
2005 2/	428	-	2	3	3	5	13	44	82	92	113	71	-
2004 2/	435	-	1	3	3	4	15	49	76	101	122	61	-
2003 2/	438	-	-	4	3	5	20	46	92	92	103	73	-
2002	436	-	-	2	1	6	14	60	62	109	125	57	-
2001	432	-	2	8	-	9	32	52	72	96	105	56	-
2000	408	1	2	4	4	2	24	48	71	92	109	51	-
1999	410	-	2	3	3	4	19	45	65	104	109	56	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Malignant neoplasms of kidney and renal pelvis (C64-C65)													
Origin not stated 3/, both sexes													
2005	18	-	-	1	-	-	-	5	2	5	3	2	-
2004	19	-	-	-	-	-	-	1	6	6	4	2	-
2003	22	-	-	-	1	-	-	2	3	8	4	4	-
2002	29	-	-	-	-	-	1	3	7	8	9	1	-
2001	23	-	-	-	-	-	1	3	3	7	5	4	-
2000	45	-	-	-	-	-	-	4	7	20	7	7	-
1999	27	-	-	-	-	1	2	3	3	5	10	3	-
Origin not stated 3/, male													
2005	15	-	-	1	-	-	-	4	2	5	2	1	-
2004	12	-	-	-	-	-	-	1	3	5	2	1	-
2003	17	-	-	-	1	-	-	1	3	6	4	2	-
2002	20	-	-	-	-	-	1	2	6	6	4	1	-
2001	14	-	-	-	-	-	1	2	2	5	3	1	-
2000	30	-	-	-	-	-	-	2	5	14	5	4	-
1999	13	-	-	-	-	-	1	2	2	4	3	1	-
Origin not stated 3/, female													
2005	3	-	-	-	-	-	-	1	-	-	1	1	-
2004	7	-	-	-	-	-	-	-	3	1	2	1	-
2003	5	-	-	-	-	-	-	1	-	2	-	2	-
2002	9	-	-	-	-	-	-	1	1	2	5	-	-
2001	9	-	-	-	-	-	-	1	1	2	2	3	-
2000	15	-	-	-	-	-	-	2	2	6	2	3	-
1999	14	-	-	-	-	1	1	1	1	1	7	2	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Malignant neoplasm of bladder (C67)													
All origins, both sexes													
2005	13,253	-	-	-	-	15	122	523	1,482	2,772	4,859	3,480	-
2004	13,030	-	-	1	1	12	120	545	1,452	2,753	4,847	3,299	-
2003	12,483	-	-	-	1	4	98	522	1,391	2,736	4,679	3,052	-
2002	12,628	-	-	3	1	14	123	503	1,310	2,907	4,707	3,059	1
2001	12,225	-	-	1	3	15	128	475	1,265	2,814	4,488	3,036	-
2000	12,002	-	-	1	3	10	111	440	1,223	2,855	4,407	2,952	-
1999	11,911	-	-	1	1	16	100	463	1,244	2,923	4,317	2,845	1
All origins, male													
2005	9,190	-	-	-	-	9	81	358	1,115	2,072	3,436	2,119	-
2004	8,995	-	-	1	1	7	69	411	1,103	2,021	3,376	2,006	-
2003	8,576	-	-	-	1	2	64	401	1,049	1,998	3,254	1,807	-
2002	8,504	-	-	1	-	9	78	352	984	2,089	3,199	1,791	1
2001	8,343	-	-	-	3	8	70	351	956	2,088	3,116	1,751	-
2000	8,163	-	-	-	3	2	62	322	916	2,094	3,054	1,710	-
1999	8,105	-	-	1	-	9	63	350	933	2,132	3,008	1,608	1
All origins, female													
2005	4,063	-	-	-	-	6	41	165	367	700	1,423	1,361	-
2004	4,035	-	-	-	-	5	51	134	349	732	1,471	1,293	-
2003	3,907	-	-	-	-	2	34	121	342	738	1,425	1,245	-
2002	4,124	-	-	2	1	5	45	151	326	818	1,508	1,268	-
2001	3,882	-	-	1	-	7	58	124	309	726	1,372	1,285	-
2000	3,839	-	-	1	-	8	49	118	307	761	1,353	1,242	-
1999	3,806	-	-	-	1	7	37	113	311	791	1,309	1,237	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Malignant neoplasm of bladder (C67)													
Hispanic, both sexes													
2005	435	-	-	-	-	1	10	22	52	107	151	92	-
2004	397	-	-	-	-	1	6	26	61	93	142	68	-
2003	394	-	-	-	-	1	4	25	54	96	133	81	-
2002	387	-	-	1	-	-	5	24	57	91	110	99	-
2001	336	-	-	-	2	-	4	21	41	90	104	74	-
2000	307	-	-	-	-	1	8	17	50	71	89	71	-
1999	301	-	-	-	-	2	4	22	51	88	79	55	-
Hispanic, male													
2005	317	-	-	-	-	1	7	18	42	80	102	67	-
2004	253	-	-	-	-	-	2	22	45	62	80	42	-
2003	265	-	-	-	-	-	2	15	40	71	87	50	-
2002	252	-	-	-	-	-	2	16	39	67	68	60	-
2001	223	-	-	-	2	-	1	11	37	59	67	46	-
2000	216	-	-	-	-	-	6	12	37	50	69	42	-
1999	222	-	-	-	-	2	4	16	43	69	52	36	-
Hispanic, female													
2005	118	-	-	-	-	-	3	4	10	27	49	25	-
2004	144	-	-	-	-	1	4	4	16	31	62	26	-
2003	129	-	-	-	-	1	2	10	14	25	46	31	-
2002	135	-	-	1	-	-	3	8	18	24	42	39	-
2001	113	-	-	-	-	-	3	10	4	31	37	28	-
2000	91	-	-	-	-	1	2	5	13	21	20	29	-
1999	79	-	-	-	-	-	-	6	8	19	27	19	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Malignant neoplasm of bladder (C67)													
Non-Hispanic 1/, both sexes													
2005	12,803	-	-	-	-	14	112	499	1,428	2,662	4,703	3,385	-
2004	12,608	-	-	1	1	10	114	518	1,385	2,656	4,696	3,227	-
2003	12,067	-	-	-	1	3	94	495	1,336	2,637	4,534	2,967	-
2002	12,211	-	-	2	1	14	118	476	1,248	2,809	4,591	2,952	-
2001	11,858	-	-	1	1	15	124	452	1,220	2,718	4,372	2,955	-
2000	11,665	-	-	1	3	9	102	423	1,169	2,780	4,304	2,874	-
1999	11,582	-	-	1	1	14	96	439	1,190	2,832	4,228	2,780	1
Non-Hispanic 1/, male													
2005	8,861	-	-	-	-	8	74	338	1,071	1,989	3,330	2,051	-
2004	8,724	-	-	1	1	6	67	388	1,052	1,957	3,291	1,961	-
2003	8,294	-	-	-	1	2	62	384	1,008	1,925	3,157	1,755	-
2002	8,235	-	-	1	-	9	76	335	941	2,019	3,128	1,726	-
2001	8,100	-	-	-	1	8	69	339	917	2,025	3,041	1,700	-
2000	7,925	-	-	-	3	2	56	310	876	2,040	2,974	1,664	-
1999	7,865	-	-	1	-	7	59	332	888	2,060	2,950	1,567	1
Non-Hispanic 1/, female													
2005	3,942	-	-	-	-	6	38	161	357	673	1,373	1,334	-
2004	3,884	-	-	-	-	4	47	130	333	699	1,405	1,266	-
2003	3,773	-	-	-	-	1	32	111	328	712	1,377	1,212	-
2002	3,976	-	-	1	1	5	42	141	307	790	1,463	1,226	-
2001	3,758	-	-	1	-	7	55	113	303	693	1,331	1,255	-
2000	3,740	-	-	1	-	7	46	113	293	740	1,330	1,210	-
1999	3,717	-	-	-	1	7	37	107	302	772	1,278	1,213	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Malignant neoplasm of bladder (C67)													
Non-Hispanic white, both sexes													
2005 2/	11,669	-	-	-	-	12	100	417	1,269	2,405	4,347	3,119	-
2004 2/	11,544	-	-	1	-	8	79	433	1,253	2,420	4,375	2,975	-
2003 2/	11,003	-	-	-	1	3	72	407	1,198	2,399	4,168	2,755	-
2002	11,109	-	-	2	-	10	100	416	1,112	2,531	4,232	2,706	-
2001	10,823	-	-	1	1	14	100	386	1,076	2,479	4,026	2,740	-
2000	10,712	-	-	1	2	8	91	374	1,056	2,533	3,957	2,690	-
1999	10,548	-	-	1	-	8	77	375	1,042	2,556	3,921	2,567	1
Non-Hispanic white, male													
2005 2/	8,207	-	-	-	-	7	67	295	969	1,828	3,122	1,919	-
2004 2/	8,140	-	-	1	-	5	46	331	965	1,825	3,116	1,851	-
2003 2/	7,692	-	-	-	1	2	48	328	921	1,777	2,952	1,663	-
2002	7,649	-	-	1	-	6	66	302	866	1,864	2,921	1,623	-
2001	7,541	-	-	-	1	8	56	296	830	1,884	2,856	1,610	-
2000	7,390	-	-	-	2	2	52	279	803	1,897	2,784	1,571	-
1999	7,290	-	-	1	-	3	46	281	800	1,907	2,771	1,480	1
Non-Hispanic white, female													
2005 2/	3,462	-	-	-	-	5	33	122	300	577	1,225	1,200	-
2004 2/	3,404	-	-	-	-	3	33	102	288	595	1,259	1,124	-
2003 2/	3,311	-	-	-	-	1	24	79	277	622	1,216	1,092	-
2002	3,460	-	-	1	-	4	34	114	246	667	1,311	1,083	-
2001	3,282	-	-	1	-	6	44	90	246	595	1,170	1,130	-
2000	3,322	-	-	1	-	6	39	95	253	636	1,173	1,119	-
1999	3,258	-	-	-	-	5	31	94	242	649	1,150	1,087	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Malignant neoplasm of bladder (C67)													
Non-Hispanic black, both sexes													
2005 2/	947	-	-	-	-	2	12	74	139	215	294	211	-
2004 2/	904	-	-	-	1	2	27	76	117	200	268	213	-
2003 2/	907	-	-	-	-	-	20	78	109	204	315	181	-
2002	931	-	-	-	-	4	13	49	116	242	295	212	-
2001	870	-	-	-	-	1	23	55	125	197	291	178	-
2000	842	-	-	-	1	-	10	44	99	218	314	156	-
1999	906	-	-	-	1	4	17	60	130	244	262	188	-
Non-Hispanic black, male													
2005 2/	536	-	-	-	-	1	7	39	88	131	163	107	-
2004 2/	474	-	-	-	1	1	18	51	75	105	141	82	-
2003 2/	496	-	-	-	-	-	12	48	66	123	172	75	-
2002	475	-	-	-	-	3	7	26	63	128	164	84	-
2001	444	-	-	-	-	-	13	36	72	111	144	68	-
2000	460	-	-	-	1	-	4	27	63	119	171	75	-
1999	487	-	-	-	-	2	12	47	76	128	149	73	-
Non-Hispanic black, female													
2005 2/	411	-	-	-	-	1	5	35	51	84	131	104	-
2004 2/	430	-	-	-	-	1	9	25	42	95	127	131	-
2003 2/	411	-	-	-	-	-	8	30	43	81	143	106	-
2002	456	-	-	-	-	1	6	23	53	114	131	128	-
2001	426	-	-	-	-	1	10	19	53	86	147	110	-
2000	382	-	-	-	-	-	6	17	36	99	143	81	-
1999	419	-	-	-	1	2	5	13	54	116	113	115	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Malignant neoplasm of bladder (C67)													
Origin not stated 3/, both sexes													
2005	15	-	-	-	-	-	-	2	2	3	5	3	-
2004	25	-	-	-	-	1	-	1	6	4	9	4	-
2003	22	-	-	-	-	-	-	2	1	3	12	4	-
2002	30	-	-	-	-	-	-	3	5	7	6	8	1
2001	31	-	-	-	-	-	-	2	4	6	12	7	-
2000	30	-	-	-	-	-	1	-	4	4	14	7	-
1999	28	-	-	-	-	-	-	2	3	3	10	10	-
Origin not stated 3/, male													
2005	12	-	-	-	-	-	-	2	2	3	4	1	-
2004	18	-	-	-	-	1	-	1	6	2	5	3	-
2003	17	-	-	-	-	-	-	2	1	2	10	2	-
2002	17	-	-	-	-	-	-	1	4	3	3	5	1
2001	20	-	-	-	-	-	-	1	2	4	8	5	-
2000	22	-	-	-	-	-	-	-	3	4	11	4	-
1999	18	-	-	-	-	-	-	2	2	3	6	5	-
Origin not stated 3/, female													
2005	3	-	-	-	-	-	-	-	-	-	1	2	-
2004	7	-	-	-	-	-	-	-	-	2	4	1	-
2003	5	-	-	-	-	-	-	-	-	1	2	2	-
2002	13	-	-	-	-	-	-	2	1	4	3	3	-
2001	11	-	-	-	-	-	-	1	2	2	4	2	-
2000	8	-	-	-	-	-	1	-	1	-	3	3	-
1999	10	-	-	-	-	-	-	-	1	-	4	5	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Malignant neoplasms of meninges, brain and other parts of central nervous system (C70-C72)													
All origins, both sexes													
2005	13,152	18	84	324	234	379	948	1,943	2,839	2,943	2,633	807	-
2004	12,829	19	102	325	223	359	904	1,902	2,750	2,895	2,543	807	-
2003	12,901	18	104	336	236	405	1,079	1,962	2,698	2,836	2,510	717	-
2002	12,830	12	114	337	210	388	1,009	2,058	2,560	2,954	2,435	753	-
2001	12,609	16	100	300	224	416	1,028	1,951	2,464	2,823	2,566	721	-
2000	12,655	20	104	316	225	406	1,096	1,924	2,337	3,032	2,478	717	-
1999	12,765	10	113	308	218	444	1,065	1,880	2,453	3,119	2,492	663	-
All origins, male													
2005	7,317	12	38	171	144	248	604	1,160	1,653	1,643	1,335	309	-
2004	7,074	10	56	173	133	218	560	1,191	1,653	1,551	1,237	292	-
2003	7,122	9	56	175	136	232	639	1,233	1,530	1,556	1,267	289	-
2002	7,125	8	63	191	136	223	618	1,225	1,494	1,678	1,210	279	-
2001	7,015	8	52	145	129	262	638	1,173	1,470	1,589	1,288	261	-
2000	7,011	10	58	176	132	264	704	1,148	1,379	1,678	1,187	275	-
1999	7,001	8	56	154	131	276	638	1,120	1,417	1,728	1,226	247	-
All origins, female													
2005	5,835	6	46	153	90	131	344	783	1,186	1,300	1,298	498	-
2004	5,755	9	46	152	90	141	344	711	1,097	1,344	1,306	515	-
2003	5,779	9	48	161	100	173	440	729	1,168	1,280	1,243	428	-
2002	5,705	4	51	146	74	165	391	833	1,066	1,276	1,225	474	-
2001	5,594	8	48	155	95	154	390	778	994	1,234	1,278	460	-
2000	5,644	10	46	140	93	142	392	776	958	1,354	1,291	442	-
1999	5,764	2	57	154	87	168	427	760	1,036	1,391	1,266	416	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Malignant neoplasms of meninges, brain and other parts of central nervous system (C70-C72)													
Hispanic, both sexes													
2005	771	6	21	54	37	36	83	124	141	135	98	36	-
2004	671	6	16	57	38	33	66	95	131	114	85	30	-
2003	669	5	20	62	23	44	63	97	123	116	98	18	-
2002	649	2	21	52	24	33	55	123	115	118	82	24	-
2001	617	4	20	37	28	38	77	87	109	117	82	18	-
2000	631	4	16	63	28	45	76	92	106	105	72	24	-
1999	568	4	27	48	18	31	65	88	99	104	67	17	-
Hispanic, male													
2005	431	2	9	21	23	26	54	77	84	68	54	13	-
2004	365	2	8	25	23	19	44	59	76	57	43	9	-
2003	368	2	10	36	15	26	38	61	62	64	43	11	-
2002	359	2	12	27	16	16	30	80	62	65	39	10	-
2001	335	1	10	16	19	21	41	50	59	72	40	6	-
2000	346	1	10	41	16	27	40	41	62	59	38	11	-
1999	293	4	11	22	10	17	45	42	60	48	31	3	-
Hispanic, female													
2005	340	4	12	33	14	10	29	47	57	67	44	23	-
2004	306	4	8	32	15	14	22	36	55	57	42	21	-
2003	301	3	10	26	8	18	25	36	61	52	55	7	-
2002	290	-	9	25	8	17	25	43	53	53	43	14	-
2001	282	3	10	21	9	17	36	37	50	45	42	12	-
2000	285	3	6	22	12	18	36	51	44	46	34	13	-
1999	275	-	16	26	8	14	20	46	39	56	36	14	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Malignant neoplasms of meninges, brain and other parts of central nervous system (C70-C72)													
Non-Hispanic 1/, both sexes													
2005	12,364	12	62	270	196	340	864	1,816	2,695	2,805	2,533	771	-
2004	12,137	13	86	267	183	326	835	1,806	2,612	2,777	2,455	777	-
2003	12,213	13	84	273	213	361	1,012	1,863	2,573	2,715	2,408	698	-
2002	12,152	10	93	284	184	355	951	1,929	2,438	2,831	2,349	728	-
2001	11,967	12	79	261	196	377	950	1,862	2,346	2,703	2,479	702	-
2000	11,995	16	87	253	197	361	1,019	1,828	2,222	2,923	2,398	691	-
1999	12,173	6	86	260	199	413	1,000	1,789	2,348	3,010	2,418	644	-
Non-Hispanic 1/, male													
2005	6,877	10	29	150	120	220	550	1,080	1,567	1,574	1,281	296	-
2004	6,699	8	48	148	109	199	513	1,131	1,573	1,493	1,194	283	-
2003	6,746	7	46	139	121	206	598	1,171	1,468	1,489	1,223	278	-
2002	6,752	6	51	164	120	207	586	1,142	1,427	1,612	1,168	269	-
2001	6,667	7	41	128	110	241	596	1,122	1,406	1,516	1,245	255	-
2000	6,646	9	47	135	116	237	663	1,103	1,313	1,615	1,145	263	-
1999	6,696	4	45	132	121	259	593	1,076	1,353	1,678	1,191	244	-
Non-Hispanic 1/, female													
2005	5,487	2	33	120	76	120	314	736	1,128	1,231	1,252	475	-
2004	5,438	5	38	119	74	127	322	675	1,039	1,284	1,261	494	-
2003	5,467	6	38	134	92	155	414	692	1,105	1,226	1,185	420	-
2002	5,400	4	42	120	64	148	365	787	1,011	1,219	1,181	459	-
2001	5,300	5	38	133	86	136	354	740	940	1,187	1,234	447	-
2000	5,349	7	40	118	81	124	356	725	909	1,308	1,253	428	-
1999	5,477	2	41	128	78	154	407	713	995	1,332	1,227	400	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Malignant neoplasms of meninges, brain and other parts of central nervous system (C70-C72)													
Non-Hispanic white, both sexes													
2005 2/	11,313	9	47	208	161	293	767	1,643	2,481	2,609	2,374	721	-
2004 2/	11,121	12	68	194	145	274	737	1,616	2,427	2,598	2,321	729	-
2003 2/	11,179	10	65	209	174	314	911	1,678	2,373	2,520	2,276	649	-
2002	11,139	5	59	214	150	305	865	1,756	2,243	2,646	2,209	687	-
2001	11,004	8	63	205	155	335	864	1,695	2,155	2,509	2,356	659	-
2000	11,026	12	65	194	156	315	896	1,678	2,047	2,756	2,260	647	-
1999	11,238	3	70	195	152	372	902	1,635	2,171	2,839	2,293	606	-
Non-Hispanic white, male													
2005 2/	6,289	8	20	114	98	189	496	977	1,441	1,471	1,196	279	-
2004 2/	6,139	7	37	110	89	168	447	1,009	1,462	1,416	1,122	272	-
2003 2/	6,178	6	38	103	97	181	533	1,059	1,362	1,385	1,153	261	-
2002	6,195	3	34	116	101	175	540	1,051	1,306	1,515	1,097	257	-
2001	6,154	5	35	96	88	217	547	1,030	1,293	1,427	1,181	235	-
2000	6,128	8	35	101	93	200	591	1,015	1,219	1,531	1,086	249	-
1999	6,193	2	39	104	91	233	533	980	1,252	1,591	1,136	232	-
Non-Hispanic white, female													
2005 2/	5,024	1	27	94	63	104	271	666	1,040	1,138	1,178	442	-
2004 2/	4,982	5	31	84	56	106	290	607	965	1,182	1,199	457	-
2003 2/	5,001	4	27	106	77	133	378	619	1,011	1,135	1,123	388	-
2002	4,944	2	25	98	49	130	325	705	937	1,131	1,112	430	-
2001	4,850	3	28	109	67	118	317	665	862	1,082	1,175	424	-
2000	4,898	4	30	93	63	115	305	663	828	1,225	1,174	398	-
1999	5,045	1	31	91	61	139	369	655	919	1,248	1,157	374	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Malignant neoplasms of meninges, brain and other parts of central nervous system (C70-C72)													
Non-Hispanic black, both sexes													
2005 2/	781	3	13	48	23	28	64	136	164	146	121	35	-
2004 2/	750	-	17	48	32	36	70	137	141	134	98	37	-
2003 2/	799	2	17	48	34	33	82	145	151	149	102	36	-
2002	766	3	27	54	26	33	68	125	144	142	112	32	-
2001	736	1	10	44	29	28	63	133	145	153	92	38	-
2000	762	2	18	51	35	36	88	119	138	133	109	33	-
1999	754	3	11	54	37	36	76	128	142	132	104	31	-
Non-Hispanic black, male													
2005 2/	434	2	8	26	14	21	35	84	95	76	64	9	-
2004 2/	406	-	10	24	14	19	48	88	87	57	51	8	-
2003 2/	440	1	6	27	21	18	55	88	81	82	54	7	-
2002	400	2	12	35	14	22	35	56	92	70	55	7	-
2001	378	-	3	25	16	14	33	75	87	70	40	15	-
2000	407	1	10	30	21	28	52	71	76	66	43	9	-
1999	399	2	4	24	24	23	47	81	76	68	40	10	-
Non-Hispanic black, female													
2005 2/	347	1	5	22	9	7	29	52	69	70	57	26	-
2004 2/	344	-	7	24	18	17	22	49	54	77	47	29	-
2003 2/	359	1	11	21	13	15	27	57	70	67	48	29	-
2002	366	1	15	19	12	11	33	69	52	72	57	25	-
2001	358	1	7	19	13	14	30	58	58	83	52	23	-
2000	355	1	8	21	14	8	36	48	62	67	66	24	-
1999	355	1	7	30	13	13	29	47	66	64	64	21	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Malignant neoplasms of meninges, brain and other parts of central nervous system (C70-C72)													
Origin not stated 3/, both sexes													
2005	17	-	1	-	1	3	1	3	3	3	2	-	-
2004	21	-	-	1	2	-	3	1	7	4	3	-	-
2003	19	-	-	1	-	-	4	2	2	5	4	1	-
2002	29	-	-	1	2	-	3	6	7	5	4	1	-
2001	25	-	1	2	-	1	1	2	9	3	5	1	-
2000	29	-	1	-	-	-	1	4	9	4	8	2	-
1999	24	-	-	-	1	-	-	3	6	5	7	2	-
Origin not stated 3/, male													
2005	9	-	-	-	1	2	-	3	2	1	-	-	-
2004	10	-	-	-	1	-	3	1	4	1	-	-	-
2003	8	-	-	-	-	-	3	1	-	3	1	-	-
2002	14	-	-	-	-	-	2	3	5	1	3	-	-
2001	13	-	1	1	-	-	1	1	5	1	3	-	-
2000	19	-	1	-	-	-	1	4	4	4	4	1	-
1999	12	-	-	-	-	-	-	2	4	2	4	-	-
Origin not stated 3/, female													
2005	8	-	1	-	-	1	1	-	1	2	2	-	-
2004	11	-	-	1	1	-	-	-	3	3	3	-	-
2003	11	-	-	1	-	-	1	1	2	2	3	1	-
2002	15	-	-	1	2	-	1	3	2	4	1	1	-
2001	12	-	-	1	-	1	-	1	4	2	2	1	-
2000	10	-	-	-	-	-	-	-	5	-	4	1	-
1999	12	-	-	-	1	-	-	1	2	3	3	2	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Malignant neoplasms of lymphoid, hematopoietic and related tissue (C81-C96)													
All origins, both sexes													
2005	55,028	24	128	335	655	811	1,498	3,687	7,623	12,487	18,109	9,671	-
2004	54,645	25	122	359	645	822	1,514	3,741	7,491	12,588	18,132	9,206	-
2003	55,679	31	138	394	606	865	1,728	3,785	7,471	12,885	18,442	9,334	-
2002	56,225	28	119	379	703	916	1,751	3,960	7,500	13,508	18,289	9,072	-
2001	56,341	27	157	367	705	988	1,857	4,064	7,455	13,659	18,257	8,805	-
2000	56,518	39	146	372	718	918	1,853	4,025	7,284	14,041	18,448	8,674	-
1999	56,318	30	137	351	720	960	1,913	3,950	7,554	14,332	18,089	8,282	-
All origins, male													
2005	30,031	12	84	206	411	464	880	2,175	4,543	7,385	9,662	4,209	-
2004	29,522	10	72	214	399	489	915	2,253	4,497	7,368	9,462	3,843	-
2003	30,076	10	80	225	375	535	1,043	2,294	4,557	7,550	9,524	3,883	-
2002	30,302	15	63	217	417	545	1,038	2,341	4,470	7,903	9,553	3,740	-
2001	30,060	12	82	217	426	574	1,133	2,442	4,412	7,859	9,327	3,576	-
2000	29,888	21	76	223	455	539	1,075	2,399	4,205	8,038	9,399	3,458	-
1999	29,721	12	75	206	437	557	1,144	2,368	4,426	8,067	9,100	3,329	-
All origins, female													
2005	24,997	12	44	129	244	347	618	1,512	3,080	5,102	8,447	5,462	-
2004	25,123	15	50	145	246	333	599	1,488	2,994	5,220	8,670	5,363	-
2003	25,603	21	58	169	231	330	685	1,491	2,914	5,335	8,918	5,451	-
2002	25,923	13	56	162	286	371	713	1,619	3,030	5,605	8,736	5,332	-
2001	26,281	15	75	150	279	414	724	1,622	3,043	5,800	8,930	5,229	-
2000	26,630	18	70	149	263	379	778	1,626	3,079	6,003	9,049	5,216	-
1999	26,597	18	62	145	283	403	769	1,582	3,128	6,265	8,989	4,953	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Malignant neoplasms of lymphoid, hematopoietic and related tissue (C81-C96)													
Hispanic, both sexes													
2005	3,110	7	28	87	155	150	182	344	497	696	712	252	-
2004	2,866	8	26	109	141	149	197	312	441	613	653	217	-
2003	2,836	6	44	97	131	147	202	323	425	576	635	250	-
2002	2,829	3	28	93	145	139	225	282	437	634	604	239	-
2001	2,747	7	38	87	144	154	211	279	442	597	573	215	-
2000	2,643	8	29	74	129	124	200	280	427	633	523	216	-
1999	2,538	8	28	75	128	135	190	299	396	606	485	188	-
Hispanic, male													
2005	1,719	2	20	51	105	94	111	195	269	382	374	116	-
2004	1,619	3	12	69	100	93	133	200	253	344	327	85	-
2003	1,528	1	26	56	89	96	130	182	241	311	299	97	-
2002	1,518	1	12	47	97	92	132	163	245	331	290	108	-
2001	1,577	2	21	48	99	99	142	164	251	338	318	95	-
2000	1,460	5	14	47	83	83	115	172	247	338	258	98	-
1999	1,319	3	16	42	73	75	108	175	226	312	226	63	-
Hispanic, female													
2005	1,391	5	8	36	50	56	71	149	228	314	338	136	-
2004	1,247	5	14	40	41	56	64	112	188	269	326	132	-
2003	1,308	5	18	41	42	51	72	141	184	265	336	153	-
2002	1,311	2	16	46	48	47	93	119	192	303	314	131	-
2001	1,170	5	17	39	45	55	69	115	191	259	255	120	-
2000	1,183	3	15	27	46	41	85	108	180	295	265	118	-
1999	1,219	5	12	33	55	60	82	124	170	294	259	125	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Malignant neoplasms of lymphoid, hematopoietic and related tissue (C81-C96)													
Non-Hispanic 1/, both sexes													
2005	51,861	17	100	246	499	659	1,312	3,334	7,115	11,782	17,382	9,415	-
2004	51,695	17	96	247	501	673	1,312	3,423	7,038	11,959	17,455	8,974	-
2003	52,736	25	94	296	474	717	1,525	3,450	7,029	12,284	17,777	9,065	-
2002	53,249	24	91	286	554	775	1,518	3,663	7,048	12,839	17,636	8,815	-
2001	53,458	20	117	279	559	828	1,640	3,772	6,994	13,027	17,645	8,577	-
2000	53,749	31	116	295	586	793	1,645	3,735	6,833	13,379	17,893	8,443	-
1999	53,669	22	108	276	592	823	1,719	3,642	7,143	13,704	17,561	8,079	-
Non-Hispanic 1/, male													
2005	28,283	10	64	154	305	369	767	1,976	4,269	7,000	9,279	4,090	-
2004	27,865	7	60	143	297	396	780	2,047	4,237	7,020	9,127	3,751	-
2003	28,485	9	54	168	285	438	912	2,103	4,304	7,229	9,208	3,775	-
2002	28,698	13	51	170	317	452	900	2,168	4,217	7,546	9,236	3,628	-
2001	28,410	10	60	169	326	473	987	2,269	4,151	7,505	8,985	3,475	-
2000	28,358	16	61	175	371	455	956	2,219	3,941	7,682	9,127	3,355	-
1999	28,347	9	58	164	364	481	1,032	2,190	4,195	7,740	8,853	3,261	-
Non-Hispanic 1/, female													
2005	23,578	7	36	92	194	290	545	1,358	2,846	4,782	8,103	5,325	-
2004	23,830	10	36	104	204	277	532	1,376	2,801	4,939	8,328	5,223	-
2003	24,251	16	40	128	189	279	613	1,347	2,725	5,055	8,569	5,290	-
2002	24,551	11	40	116	237	323	618	1,495	2,831	5,293	8,400	5,187	-
2001	25,048	10	57	110	233	355	653	1,503	2,843	5,522	8,660	5,102	-
2000	25,391	15	55	120	215	338	689	1,516	2,892	5,697	8,766	5,088	-
1999	25,322	13	50	112	228	342	687	1,452	2,948	5,964	8,708	4,818	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Malignant neoplasms of lymphoid, hematopoietic and related tissue (C81-C96)													
Non-Hispanic white, both sexes													
2005 2/	45,619	14	66	186	370	434	994	2,570	5,967	10,276	15,954	8,788	-
2004 2/	45,383	13	69	183	373	479	1,002	2,705	5,933	10,394	15,920	8,312	-
2003 2/	46,553	23	72	218	355	524	1,166	2,727	5,996	10,779	16,253	8,440	-
2002	47,058	15	72	213	430	570	1,183	2,897	5,930	11,327	16,202	8,219	-
2001	47,190	17	97	198	422	595	1,255	3,057	5,878	11,492	16,199	7,980	-
2000	47,695	30	92	232	420	585	1,283	3,017	5,812	11,911	16,528	7,785	-
1999	47,608	16	91	199	437	618	1,352	2,980	6,042	12,208	16,172	7,493	-
Non-Hispanic white, male													
2005 2/	25,000	8	40	113	229	241	592	1,536	3,628	6,191	8,563	3,859	-
2004 2/	24,604	6	41	109	219	295	607	1,627	3,630	6,160	8,406	3,504	-
2003 2/	25,390	7	42	127	214	324	708	1,682	3,719	6,471	8,552	3,544	-
2002	25,523	9	42	125	254	333	701	1,755	3,597	6,746	8,557	3,404	-
2001	25,162	8	47	111	252	352	748	1,873	3,519	6,693	8,307	3,252	-
2000	25,272	16	50	134	268	338	775	1,803	3,379	6,899	8,491	3,119	-
1999	25,219	9	48	120	277	357	816	1,812	3,591	6,967	8,202	3,020	-
Non-Hispanic white, female													
2005 2/	20,619	6	26	73	141	193	402	1,034	2,339	4,085	7,391	4,929	-
2004 2/	20,779	7	28	74	154	184	395	1,078	2,303	4,234	7,514	4,808	-
2003 2/	21,163	16	30	91	141	200	458	1,045	2,277	4,308	7,701	4,896	-
2002	21,535	6	30	88	176	237	482	1,142	2,333	4,581	7,645	4,815	-
2001	22,028	9	50	87	170	243	507	1,184	2,359	4,799	7,892	4,728	-
2000	22,423	14	42	98	152	247	508	1,214	2,433	5,012	8,037	4,666	-
1999	22,389	7	43	79	160	261	536	1,168	2,451	5,241	7,970	4,473	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Malignant neoplasms of lymphoid, hematopoietic and related tissue (C81-C96)													
Non-Hispanic black, both sexes													
2005 2/	4,988	3	26	44	101	180	255	618	921	1,199	1,131	510	-
2004 2/	5,163	3	18	46	104	163	253	609	926	1,281	1,217	543	-
2003 2/	5,059	1	14	57	88	155	299	620	853	1,237	1,223	512	-
2002	5,080	5	14	51	87	156	266	624	935	1,268	1,166	508	-
2001	5,162	3	13	62	107	181	322	584	915	1,288	1,183	504	-
2000	5,026	1	19	43	130	158	302	615	868	1,232	1,101	557	-
1999	5,038	4	14	61	133	157	307	554	917	1,247	1,142	502	-
Non-Hispanic black, male													
2005 2/	2,604	2	18	31	61	99	140	361	520	644	551	177	-
2004 2/	2,608	-	13	24	61	86	139	352	505	698	536	194	-
2003 2/	2,517	1	8	31	52	87	170	366	491	613	521	177	-
2002	2,593	2	6	32	44	93	162	342	517	679	530	186	-
2001	2,642	2	9	43	57	98	199	326	512	679	538	179	-
2000	2,535	-	8	26	79	90	149	346	481	653	510	193	-
1999	2,566	-	8	38	73	97	180	318	500	645	521	186	-
Non-Hispanic black, female													
2005 2/	2,384	1	8	13	40	81	115	257	401	555	580	333	-
2004 2/	2,555	3	5	22	43	77	114	257	421	583	681	349	-
2003 2/	2,542	-	6	26	36	68	129	254	362	624	702	335	-
2002	2,487	3	8	19	43	63	104	282	418	589	636	322	-
2001	2,520	1	4	19	50	83	123	258	403	609	645	325	-
2000	2,491	1	11	17	51	68	153	269	387	579	591	364	-
1999	2,472	4	6	23	60	60	127	236	417	602	621	316	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Malignant neoplasms of lymphoid, hematopoietic and related tissue (C81-C96)													
Origin not stated 3/, both sexes													
2005	57	-	-	2	1	2	4	9	11	9	15	4	-
2004	84	-	-	3	3	-	5	6	12	16	24	15	-
2003	107	-	-	1	1	1	1	12	17	25	30	19	-
2002	147	1	-	-	4	2	8	15	15	35	49	18	-
2001	136	-	2	1	2	6	6	13	19	35	39	13	-
2000	126	-	1	3	3	1	8	10	24	29	32	15	-
1999	111	-	1	-	-	2	4	9	15	22	43	15	-
Origin not stated 3/, male													
2005	29	-	-	1	1	1	2	4	5	3	9	3	-
2004	38	-	-	2	2	-	2	6	7	4	8	7	-
2003	63	-	-	1	1	1	1	9	12	10	17	11	-
2002	86	1	-	-	3	1	6	10	8	26	27	4	-
2001	73	-	1	-	1	2	4	9	10	16	24	6	-
2000	70	-	1	1	1	1	4	8	17	18	14	5	-
1999	55	-	1	-	-	1	4	3	5	15	21	5	-
Origin not stated 3/, female													
2005	28	-	-	1	-	1	2	5	6	6	6	1	-
2004	46	-	-	1	1	-	3	-	5	12	16	8	-
2003	44	-	-	-	-	-	-	3	5	15	13	8	-
2002	61	-	-	-	1	1	2	5	7	9	22	14	-
2001	63	-	1	1	1	4	2	4	9	19	15	7	-
2000	56	-	-	2	2	-	4	2	7	11	18	10	-
1999	56	-	-	-	-	1	-	6	10	7	22	10	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Hodgkin's disease (C81)													
All origins, both sexes													
2005	1,272	-	-	11	61	116	118	168	184	205	304	105	-
2004	1,276	-	-	6	65	134	120	168	205	222	249	107	-
2003	1,347	-	-	5	64	167	154	154	198	216	277	112	-
2002	1,352	-	-	6	86	152	159	177	168	232	247	125	-
2001	1,323	-	-	6	78	167	161	182	143	211	278	97	-
2000	1,287	-	-	8	73	145	151	161	158	232	255	104	-
1999	1,403	-	1	9	94	148	165	165	178	283	262	98	-
All origins, male													
2005	684	-	-	7	27	65	71	94	121	118	131	50	-
2004	730	-	-	4	37	70	70	103	132	136	129	49	-
2003	725	-	-	4	30	88	95	91	119	128	132	38	-
2002	751	-	-	5	41	81	92	105	106	146	125	50	-
2001	725	-	-	3	43	90	101	110	85	129	130	34	-
2000	686	-	-	4	35	79	82	99	87	134	121	45	-
1999	785	-	1	3	55	86	103	98	115	160	119	45	-
All origins, female													
2005	588	-	-	4	34	51	47	74	63	87	173	55	-
2004	546	-	-	2	28	64	50	65	73	86	120	58	-
2003	622	-	-	1	34	79	59	63	79	88	145	74	-
2002	601	-	-	1	45	71	67	72	62	86	122	75	-
2001	598	-	-	3	35	77	60	72	58	82	148	63	-
2000	601	-	-	4	38	66	69	62	71	98	134	59	-
1999	618	-	-	6	39	62	62	67	63	123	143	53	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Hodgkin's disease (C81)													
Hispanic, both sexes													
2005	126	-	-	4	6	11	9	18	19	29	24	6	-
2004	95	-	-	2	8	12	8	11	12	17	15	10	-
2003	91	-	-	1	8	16	6	15	8	19	12	6	-
2002	99	-	-	4	9	15	12	15	10	13	18	3	-
2001	99	-	-	1	7	13	13	14	14	21	11	5	-
2000	95	-	-	-	9	14	8	13	13	11	23	4	-
1999	113	-	-	3	8	7	12	17	17	26	18	5	-
Hispanic, male													
2005	78	-	-	3	2	6	7	13	15	16	12	4	-
2004	57	-	-	1	6	7	6	8	6	13	5	5	-
2003	52	-	-	1	5	8	5	8	5	11	8	1	-
2002	53	-	-	3	4	7	8	9	7	6	8	1	-
2001	66	-	-	1	4	8	8	11	11	15	5	3	-
2000	63	-	-	-	7	7	6	11	8	10	12	2	-
1999	65	-	-	2	6	3	8	11	9	13	11	2	-
Hispanic, female													
2005	48	-	-	1	4	5	2	5	4	13	12	2	-
2004	38	-	-	1	2	5	2	3	6	4	10	5	-
2003	39	-	-	-	3	8	1	7	3	8	4	5	-
2002	46	-	-	1	5	8	4	6	3	7	10	2	-
2001	33	-	-	-	3	5	5	3	3	6	6	2	-
2000	32	-	-	-	2	7	2	2	5	1	11	2	-
1999	48	-	-	1	2	4	4	6	8	13	7	3	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Hodgkin's disease (C81)													
Non-Hispanic 1/, both sexes													
2005	1,141	-	-	6	55	104	108	149	165	176	280	98	-
2004	1,176	-	-	4	56	122	110	157	193	205	233	96	-
2003	1,253	-	-	4	56	151	147	138	190	197	264	106	-
2002	1,248	-	-	2	76	136	146	161	158	219	228	122	-
2001	1,221	-	-	5	71	154	147	167	129	189	267	92	-
2000	1,189	-	-	8	64	131	143	148	143	221	231	100	-
1999	1,286	-	1	6	86	141	153	147	159	257	243	93	-
Non-Hispanic 1/, male													
2005	602	-	-	4	25	58	63	80	106	102	119	45	-
2004	669	-	-	3	30	63	62	95	126	123	123	44	-
2003	672	-	-	3	25	80	89	83	114	117	124	37	-
2002	693	-	-	2	36	73	83	95	99	140	116	49	-
2001	657	-	-	2	39	82	92	98	74	114	125	31	-
2000	621	-	-	4	28	72	76	88	78	124	108	43	-
1999	719	-	1	1	49	83	95	87	105	147	108	43	-
Non-Hispanic 1/, female													
2005	539	-	-	2	30	46	45	69	59	74	161	53	-
2004	507	-	-	1	26	59	48	62	67	82	110	52	-
2003	581	-	-	1	31	71	58	55	76	80	140	69	-
2002	555	-	-	-	40	63	63	66	59	79	112	73	-
2001	564	-	-	3	32	72	55	69	55	75	142	61	-
2000	568	-	-	4	36	59	67	60	65	97	123	57	-
1999	567	-	-	5	37	58	58	60	54	110	135	50	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Hodgkin's disease (C81)													
Non-Hispanic white, both sexes													
2005 2/	968	-	-	6	42	61	79	119	145	160	266	90	-
2004 2/	1,020	-	-	3	43	86	91	135	169	184	220	89	-
2003 2/	1,099	-	-	4	43	118	125	123	162	177	246	101	-
2002	1,093	-	-	1	63	112	118	135	132	200	215	117	-
2001	1,069	-	-	4	55	113	124	146	110	178	249	90	-
2000	1,059	-	-	5	48	117	119	128	123	200	224	95	-
1999	1,149	-	1	5	67	116	134	129	144	234	235	84	-
Non-Hispanic white, male													
2005 2/	504	-	-	4	20	31	48	63	89	92	113	44	-
2004 2/	587	-	-	3	23	49	51	82	110	109	117	43	-
2003 2/	585	-	-	3	19	66	78	73	96	101	115	34	-
2002	599	-	-	1	32	57	65	75	80	130	111	48	-
2001	577	-	-	2	34	62	76	85	63	109	117	29	-
2000	551	-	-	2	23	63	64	76	65	112	105	41	-
1999	643	-	1	1	39	69	84	73	97	135	105	39	-
Non-Hispanic white, female													
2005 2/	464	-	-	2	22	30	31	56	56	68	153	46	-
2004 2/	433	-	-	-	20	37	40	53	59	75	103	46	-
2003 2/	514	-	-	1	24	52	47	50	66	76	131	67	-
2002	494	-	-	-	31	55	53	60	52	70	104	69	-
2001	492	-	-	2	21	51	48	61	47	69	132	61	-
2000	508	-	-	3	25	54	55	52	58	88	119	54	-
1999	506	-	-	4	28	47	50	56	47	99	130	45	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Hodgkin's disease (C81)													
Non-Hispanic black, both sexes													
2005 2/	141	-	-	-	9	36	25	27	16	13	10	5	-
2004 2/	133	-	-	1	13	32	17	20	18	18	11	3	-
2003 2/	119	-	-	-	10	31	18	14	21	17	8	-	-
2002	126	-	-	1	10	23	25	20	20	15	9	3	-
2001	136	-	-	1	15	39	21	17	18	11	12	2	-
2000	109	-	-	2	14	12	24	17	17	15	5	3	-
1999	125	-	-	-	19	25	18	18	14	18	6	7	-
Non-Hispanic black, male													
2005 2/	81	-	-	-	4	22	13	14	14	8	5	1	-
2004 2/	70	-	-	-	7	13	9	13	12	11	5	-	-
2003 2/	68	-	-	-	5	12	10	9	15	13	4	-	-
2002	80	-	-	1	4	15	17	15	15	8	4	1	-
2001	72	-	-	-	4	20	15	11	10	5	5	2	-
2000	57	-	-	1	5	8	12	9	11	7	3	1	-
1999	74	-	-	-	10	14	11	14	8	12	3	2	-
Non-Hispanic black, female													
2005 2/	60	-	-	-	5	14	12	13	2	5	5	4	-
2004 2/	63	-	-	1	6	19	8	7	6	7	6	3	-
2003 2/	51	-	-	-	5	19	8	5	6	4	4	-	-
2002	46	-	-	-	6	8	8	5	5	7	5	2	-
2001	64	-	-	1	11	19	6	6	8	6	7	-	-
2000	52	-	-	1	9	4	12	8	6	8	2	2	-
1999	51	-	-	-	9	11	7	4	6	6	3	5	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Hodgkin's disease (C81)													
Origin not stated 3/, both sexes													
2005	5	-	-	1	-	1	1	1	-	-	-	1	-
2004	5	-	-	-	1	-	2	-	-	-	1	1	-
2003	3	-	-	-	-	-	1	1	-	-	1	-	-
2002	5	-	-	-	1	1	1	1	-	-	1	-	-
2001	3	-	-	-	-	-	1	1	-	1	-	-	-
2000	3	-	-	-	-	-	-	-	2	-	1	-	-
1999	4	-	-	-	-	-	-	1	2	-	1	-	-
Origin not stated 3/, male													
2005	4	-	-	-	-	1	1	1	-	-	-	1	-
2004	4	-	-	-	1	-	2	-	-	-	1	-	-
2003	1	-	-	-	-	-	1	-	-	-	-	-	-
2002	5	-	-	-	1	1	1	1	-	-	1	-	-
2001	2	-	-	-	-	-	1	1	-	-	-	-	-
2000	2	-	-	-	-	-	-	-	1	-	1	-	-
1999	1	-	-	-	-	-	-	-	1	-	-	-	-
Origin not stated 3/, female													
2005	1	-	-	1	-	-	-	-	-	-	-	-	-
2004	1	-	-	-	-	-	-	-	-	-	-	1	-
2003	2	-	-	-	-	-	-	1	-	-	1	-	-
2002	-	-	-	-	-	-	-	-	-	-	-	-	-
2001	1	-	-	-	-	-	-	-	-	1	-	-	-
2000	1	-	-	-	-	-	-	-	1	-	-	-	-
1999	3	-	-	-	-	-	-	1	1	-	1	-	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Non-Hodgkin's lymphoma (C82-C85)													
All origins, both sexes													
2005	20,873	1	6	48	127	222	566	1,410	2,925	4,626	7,111	3,831	-
2004	20,938	1	2	41	144	256	543	1,538	2,866	4,820	7,080	3,647	-
2003	21,475	-	5	45	140	252	630	1,548	3,007	4,919	7,217	3,712	-
2002	21,910	2	15	57	142	281	687	1,581	3,049	5,264	7,283	3,549	-
2001	22,305	1	1	38	148	298	756	1,732	3,098	5,358	7,369	3,506	-
2000	22,729	1	7	47	180	293	751	1,745	3,170	5,598	7,535	3,402	-
1999	22,802	2	16	37	144	289	789	1,741	3,154	5,848	7,554	3,228	-
All origins, male													
2005	11,131	-	2	40	90	134	365	909	1,745	2,677	3,557	1,612	-
2004	11,056	-	2	28	93	164	367	983	1,725	2,738	3,511	1,445	-
2003	11,399	-	5	26	88	165	406	972	1,848	2,829	3,563	1,497	-
2002	11,538	1	6	40	93	178	439	978	1,838	2,986	3,594	1,385	-
2001	11,710	1	1	29	94	179	500	1,113	1,825	3,025	3,558	1,385	-
2000	11,812	-	5	28	126	172	482	1,112	1,849	3,105	3,634	1,299	-
1999	11,794	-	9	30	91	178	510	1,111	1,847	3,176	3,596	1,246	-
All origins, female													
2005	9,742	1	4	8	37	88	201	501	1,180	1,949	3,554	2,219	-
2004	9,882	1	-	13	51	92	176	555	1,141	2,082	3,569	2,202	-
2003	10,076	-	-	19	52	87	224	576	1,159	2,090	3,654	2,215	-
2002	10,372	1	9	17	49	103	248	603	1,211	2,278	3,689	2,164	-
2001	10,595	-	-	9	54	119	256	619	1,273	2,333	3,811	2,121	-
2000	10,917	1	2	19	54	121	269	633	1,321	2,493	3,901	2,103	-
1999	11,008	2	7	7	53	111	279	630	1,307	2,672	3,958	1,982	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Non-Hodgkin's lymphoma (C82-C85)													
Hispanic, both sexes													
2005	1,118	-	1	4	23	34	65	122	182	257	326	104	-
2004	1,062	-	-	9	25	47	75	131	167	238	278	92	-
2003	1,057	-	3	10	27	46	82	116	176	233	259	105	-
2002	1,068	-	2	7	25	41	78	108	186	268	262	91	-
2001	1,008	-	-	6	22	37	77	122	201	208	240	95	-
2000	1,026	-	2	2	21	38	83	113	190	277	210	90	-
1999	958	-	1	3	19	37	68	131	153	261	205	80	-
Hispanic, male													
2005	593	-	-	3	17	21	40	70	101	141	158	42	-
2004	605	-	-	6	17	28	58	99	93	136	132	36	-
2003	569	-	3	5	19	33	56	65	104	129	118	37	-
2002	549	-	-	3	19	25	45	65	109	136	111	36	-
2001	579	-	-	6	15	24	56	74	115	122	131	36	-
2000	560	-	2	1	17	23	51	72	112	143	98	41	-
1999	493	-	-	2	10	18	43	84	91	125	92	28	-
Hispanic, female													
2005	525	-	1	1	6	13	25	52	81	116	168	62	-
2004	457	-	-	3	8	19	17	32	74	102	146	56	-
2003	488	-	-	5	8	13	26	51	72	104	141	68	-
2002	519	-	2	4	6	16	33	43	77	132	151	55	-
2001	429	-	-	-	7	13	21	48	86	86	109	59	-
2000	466	-	-	1	4	15	32	41	78	134	112	49	-
1999	465	-	1	1	9	19	25	47	62	136	113	52	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Non-Hodgkin's lymphoma (C82-C85)													
Non-Hispanic 1/, both sexes													
2005	19,734	1	5	44	103	188	501	1,285	2,739	4,365	6,778	3,725	-
2004	19,842	1	2	31	119	209	465	1,404	2,694	4,575	6,793	3,549	-
2003	20,379	-	2	35	112	206	548	1,427	2,823	4,679	6,947	3,600	-
2002	20,780	2	13	50	115	240	603	1,466	2,857	4,983	6,999	3,452	-
2001	21,246	1	1	32	125	258	678	1,605	2,892	5,134	7,113	3,407	-
2000	21,657	1	5	45	159	254	668	1,626	2,970	5,311	7,312	3,306	-
1999	21,792	2	15	34	125	252	718	1,606	2,996	5,575	7,327	3,142	-
Non-Hispanic 1/, male													
2005	10,529	-	2	37	72	113	325	838	1,642	2,535	3,396	1,569	-
2004	10,437	-	2	22	76	136	309	881	1,628	2,601	3,377	1,405	-
2003	10,810	-	2	21	68	132	350	904	1,739	2,697	3,440	1,457	-
2002	10,953	1	6	37	73	153	389	910	1,725	2,838	3,472	1,349	-
2001	11,105	1	1	23	79	153	443	1,036	1,708	2,897	3,416	1,348	-
2000	11,225	-	3	27	109	148	431	1,036	1,728	2,954	3,533	1,256	-
1999	11,271	-	9	28	81	160	464	1,025	1,754	3,044	3,490	1,216	-
Non-Hispanic 1/, female													
2005	9,205	1	3	7	31	75	176	447	1,097	1,830	3,382	2,156	-
2004	9,405	1	-	9	43	73	156	523	1,066	1,974	3,416	2,144	-
2003	9,569	-	-	14	44	74	198	523	1,084	1,982	3,507	2,143	-
2002	9,827	1	7	13	42	87	214	556	1,132	2,145	3,527	2,103	-
2001	10,141	-	-	9	46	105	235	569	1,184	2,237	3,697	2,059	-
2000	10,432	1	2	18	50	106	237	590	1,242	2,357	3,779	2,050	-
1999	10,521	2	6	6	44	92	254	581	1,242	2,531	3,837	1,926	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Non-Hodgkin's lymphoma (C82-C85)													
Non-Hispanic white, both sexes													
2005 2/	17,852	1	4	34	70	118	369	1,016	2,369	3,948	6,377	3,546	-
2004 2/	17,970	1	2	20	85	139	362	1,125	2,360	4,142	6,379	3,355	-
2003 2/	18,526	-	2	22	85	132	417	1,158	2,500	4,242	6,533	3,435	-
2002	18,901	1	12	38	80	162	463	1,198	2,517	4,523	6,597	3,310	-
2001	19,300	1	1	26	87	182	523	1,337	2,535	4,683	6,702	3,223	-
2000	19,843	-	2	38	110	181	531	1,367	2,655	4,878	6,938	3,143	-
1999	19,956	-	12	23	91	177	559	1,372	2,667	5,145	6,935	2,975	-
Non-Hispanic white, male													
2005 2/	9,527	-	1	27	54	68	252	662	1,430	2,310	3,220	1,503	-
2004 2/	9,440	-	2	16	52	95	248	710	1,431	2,361	3,186	1,339	-
2003 2/	9,871	-	2	12	53	88	271	738	1,556	2,490	3,267	1,394	-
2002	9,961	-	6	29	55	106	297	763	1,526	2,600	3,287	1,292	-
2001	10,052	1	1	17	54	115	339	873	1,497	2,655	3,226	1,274	-
2000	10,264	-	2	23	79	109	353	870	1,538	2,725	3,365	1,200	-
1999	10,259	-	8	21	61	112	362	879	1,554	2,822	3,302	1,138	-
Non-Hispanic white, female													
2005 2/	8,325	1	3	7	16	50	117	354	939	1,638	3,157	2,043	-
2004 2/	8,530	1	-	4	33	44	114	415	929	1,781	3,193	2,016	-
2003 2/	8,655	-	-	10	32	44	146	420	944	1,752	3,266	2,041	-
2002	8,940	1	6	9	25	56	166	435	991	1,923	3,310	2,018	-
2001	9,248	-	-	9	33	67	184	464	1,038	2,028	3,476	1,949	-
2000	9,579	-	-	15	31	72	178	497	1,117	2,153	3,573	1,943	-
1999	9,697	-	4	2	30	65	197	493	1,113	2,323	3,633	1,837	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Non-Hodgkin's lymphoma (C82-C85)													
Non-Hispanic black, both sexes													
2005 2/	1,379	-	1	9	28	57	109	214	274	295	266	126	-
2004 2/	1,391	-	-	8	30	59	84	231	257	311	274	137	-
2003 2/	1,359	-	-	10	23	58	108	229	244	307	275	105	-
2002	1,446	1	1	10	24	63	112	215	277	356	284	103	-
2001	1,478	-	-	5	32	62	129	221	269	348	284	128	-
2000	1,355	1	2	5	35	55	118	206	256	318	244	115	-
1999	1,427	2	2	9	32	58	135	189	259	330	286	125	-
Non-Hispanic black, male													
2005 2/	744	-	1	9	15	34	60	143	163	165	111	43	-
2004 2/	728	-	-	3	20	36	50	145	150	165	117	42	-
2003 2/	682	-	-	8	12	34	63	143	138	143	108	33	-
2002	767	1	-	7	10	40	76	119	159	193	122	40	-
2001	795	-	-	5	22	32	88	135	151	190	124	48	-
2000	719	-	-	3	19	30	68	133	157	169	103	37	-
1999	778	-	1	6	18	39	86	120	153	174	131	50	-
Non-Hispanic black, female													
2005 2/	635	-	-	-	13	23	49	71	111	130	155	83	-
2004 2/	663	-	-	5	10	23	34	86	107	146	157	95	-
2003 2/	677	-	-	2	11	24	45	86	106	164	167	72	-
2002	679	-	1	3	14	23	36	96	118	163	162	63	-
2001	683	-	-	-	10	30	41	86	118	158	160	80	-
2000	636	1	2	2	16	25	50	73	99	149	141	78	-
1999	649	2	1	3	14	19	49	69	106	156	155	75	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Non-Hodgkin's lymphoma (C82-C85)													
Origin not stated 3/, both sexes													
2005	21	-	-	-	1	-	-	3	4	4	7	2	-
2004	34	-	-	1	-	-	3	3	5	7	9	6	-
2003	39	-	-	-	1	-	-	5	8	7	11	7	-
2002	62	-	-	-	2	-	6	7	6	13	22	6	-
2001	51	-	-	-	1	3	1	5	5	16	16	4	-
2000	46	-	-	-	-	1	-	6	10	10	13	6	-
1999	52	-	-	-	-	-	3	4	5	12	22	6	-
Origin not stated 3/, male													
2005	9	-	-	-	1	-	-	1	2	1	3	1	-
2004	14	-	-	-	-	-	-	3	4	1	2	4	-
2003	20	-	-	-	1	-	-	3	5	3	5	3	-
2002	36	-	-	-	1	-	5	3	4	12	11	-	-
2001	26	-	-	-	-	2	1	3	2	6	11	1	-
2000	27	-	-	-	-	1	-	4	9	8	3	2	-
1999	30	-	-	-	-	-	3	2	2	7	14	2	-
Origin not stated 3/, female													
2005	12	-	-	-	-	-	-	2	2	3	4	1	-
2004	20	-	-	1	-	-	3	-	1	6	7	2	-
2003	19	-	-	-	-	-	-	2	3	4	6	4	-
2002	26	-	-	-	1	-	1	4	2	1	11	6	-
2001	25	-	-	-	1	1	-	2	3	10	5	3	-
2000	19	-	-	-	-	-	-	2	1	2	10	4	-
1999	22	-	-	-	-	-	-	2	3	5	8	4	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Leukemia (C91-C95)													
All origins, both sexes													
2005	21,623	22	120	275	466	461	680	1,419	2,662	4,673	6,832	4,013	-
2004	21,395	23	120	312	435	416	713	1,367	2,688	4,694	6,797	3,830	-
2003	21,535	30	133	342	402	433	796	1,365	2,605	4,706	6,937	3,786	-
2002	21,498	21	104	316	472	471	767	1,459	2,611	4,818	6,763	3,696	-
2001	21,451	25	155	323	479	513	779	1,475	2,535	4,886	6,696	3,585	-
2000	21,339	38	137	315	464	473	801	1,411	2,414	5,010	6,733	3,543	-
1999	21,014	27	119	305	482	500	799	1,370	2,516	5,012	6,460	3,424	-
All origins, male													
2005	12,231	11	81	159	293	257	358	762	1,649	2,902	3,957	1,802	-
2004	12,007	9	70	182	268	242	398	782	1,663	2,903	3,821	1,669	-
2003	12,073	9	75	195	257	274	444	801	1,593	2,930	3,880	1,615	-
2002	12,016	11	57	172	282	280	429	821	1,558	3,010	3,804	1,592	-
2001	11,850	10	81	185	289	300	432	813	1,552	2,962	3,699	1,527	-
2000	11,777	21	69	190	293	283	423	791	1,452	3,090	3,695	1,470	-
1999	11,519	11	65	173	291	276	435	777	1,489	3,054	3,525	1,423	-
All origins, female													
2005	9,392	11	39	116	173	204	322	657	1,013	1,771	2,875	2,211	-
2004	9,388	14	50	130	167	174	315	585	1,025	1,791	2,976	2,161	-
2003	9,462	21	58	147	145	159	352	564	1,012	1,776	3,057	2,171	-
2002	9,482	10	47	144	190	191	338	638	1,053	1,808	2,959	2,104	-
2001	9,601	15	74	138	190	213	347	662	983	1,924	2,997	2,058	-
2000	9,562	17	68	125	171	190	378	620	962	1,920	3,038	2,073	-
1999	9,495	16	54	132	191	224	364	593	1,027	1,958	2,935	2,001	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Leukemia (C91-C95)													
Hispanic, both sexes													
2005	1,272	7	25	79	125	102	97	137	166	233	206	95	-
2004	1,150	8	26	98	108	86	95	119	139	199	196	76	-
2003	1,156	6	41	86	96	84	100	139	132	170	211	91	-
2002	1,088	2	26	82	109	80	121	93	143	183	164	85	-
2001	1,114	7	38	80	115	101	103	100	120	197	185	68	-
2000	1,038	8	27	72	99	71	97	110	133	182	165	74	-
1999	994	8	26	69	101	86	98	100	115	179	149	63	-
Hispanic, male													
2005	747	2	19	45	85	64	58	73	95	135	125	46	-
2004	669	3	12	62	77	54	57	67	86	114	106	31	-
2003	649	1	23	50	65	54	58	78	75	99	106	40	-
2002	601	1	12	41	73	57	69	51	67	97	94	39	-
2001	640	2	21	41	80	66	65	56	73	98	102	36	-
2000	595	5	12	46	59	52	52	64	82	98	87	38	-
1999	534	3	16	38	57	49	48	52	66	107	75	23	-
Hispanic, female													
2005	525	5	6	34	40	38	39	64	71	98	81	49	-
2004	481	5	14	36	31	32	38	52	53	85	90	45	-
2003	507	5	18	36	31	30	42	61	57	71	105	51	-
2002	487	1	14	41	36	23	52	42	76	86	70	46	-
2001	474	5	17	39	35	35	38	44	47	99	83	32	-
2000	443	3	15	26	40	19	45	46	51	84	78	36	-
1999	460	5	10	31	44	37	50	48	49	72	74	40	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Leukemia (C91-C95)													
Non-Hispanic 1/, both sexes													
2005	20,329	15	95	195	341	358	580	1,277	2,493	4,436	6,621	3,918	-
2004	20,214	15	94	212	325	330	618	1,246	2,543	4,487	6,596	3,748	-
2003	20,340	24	92	255	306	348	696	1,221	2,467	4,526	6,718	3,687	-
2002	20,363	18	78	234	362	390	645	1,360	2,464	4,622	6,586	3,604	-
2001	20,285	18	115	242	363	409	673	1,371	2,406	4,677	6,501	3,510	-
2000	20,248	30	109	240	362	402	697	1,299	2,274	4,817	6,555	3,463	-
1999	19,981	19	92	236	381	412	700	1,266	2,395	4,826	6,298	3,356	-
Non-Hispanic 1/, male													
2005	11,471	9	62	113	208	193	299	687	1,551	2,765	3,828	1,756	-
2004	11,322	6	58	118	190	188	341	713	1,574	2,787	3,711	1,636	-
2003	11,394	8	52	144	192	219	386	718	1,513	2,825	3,768	1,569	-
2002	11,386	9	45	131	208	223	360	765	1,489	2,904	3,702	1,550	-
2001	11,180	8	59	144	208	234	365	754	1,474	2,857	3,590	1,487	-
2000	11,155	16	56	143	233	231	367	725	1,366	2,987	3,601	1,430	-
1999	10,968	8	48	135	234	226	386	724	1,421	2,940	3,446	1,400	-
Non-Hispanic 1/, female													
2005	8,858	6	33	82	133	165	281	590	942	1,671	2,793	2,162	-
2004	8,892	9	36	94	135	142	277	533	969	1,700	2,885	2,112	-
2003	8,946	16	40	111	114	129	310	503	954	1,701	2,950	2,118	-
2002	8,977	9	33	103	154	167	285	595	975	1,718	2,884	2,054	-
2001	9,105	10	56	98	155	175	308	617	932	1,820	2,911	2,023	-
2000	9,093	14	53	97	129	171	330	574	908	1,830	2,954	2,033	-
1999	9,013	11	44	101	147	186	314	542	974	1,886	2,852	1,956	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Leukemia (C91-C95)													
Non-Hispanic white, both sexes													
2005 2/	18,152	12	62	145	258	250	462	1,024	2,167	3,962	6,127	3,683	-
2004 2/	17,936	11	67	160	244	246	470	1,018	2,194	3,956	6,074	3,496	-
2003 2/	18,190	22	70	190	227	265	534	1,004	2,151	4,054	6,221	3,452	-
2002	18,162	13	60	174	286	289	524	1,102	2,099	4,169	6,087	3,359	-
2001	18,105	15	95	168	280	298	524	1,123	2,084	4,185	6,031	3,302	-
2000	18,128	30	88	187	262	284	546	1,059	1,952	4,372	6,121	3,227	-
1999	17,885	15	78	171	279	315	564	1,040	2,075	4,348	5,848	3,152	-
Non-Hispanic white, male													
2005 2/	10,289	7	39	82	155	140	232	558	1,366	2,495	3,546	1,669	-
2004 2/	10,098	5	39	90	143	145	260	580	1,383	2,466	3,450	1,537	-
2003 2/	10,278	6	40	112	142	164	296	601	1,335	2,560	3,540	1,482	-
2002	10,218	8	36	95	167	168	294	626	1,290	2,631	3,446	1,457	-
2001	10,018	6	46	92	164	173	277	632	1,289	2,579	3,349	1,411	-
2000	10,009	16	46	108	166	164	298	593	1,183	2,726	3,377	1,332	-
1999	9,853	8	39	98	177	170	313	593	1,246	2,673	3,220	1,316	-
Non-Hispanic white, female													
2005 2/	7,863	5	23	63	103	110	230	466	801	1,467	2,581	2,014	-
2004 2/	7,838	6	28	70	101	101	210	438	811	1,490	2,624	1,959	-
2003 2/	7,912	16	30	78	85	101	238	403	816	1,494	2,681	1,970	-
2002	7,944	5	24	79	119	121	230	476	809	1,538	2,641	1,902	-
2001	8,087	9	49	76	116	125	247	491	795	1,606	2,682	1,891	-
2000	8,119	14	42	79	96	120	248	466	769	1,646	2,744	1,895	-
1999	8,032	7	39	73	102	145	251	447	829	1,675	2,628	1,836	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Leukemia (C91-C95)													
Non-Hispanic black, both sexes													
2005 2/	1,697	3	25	35	64	83	87	198	254	369	385	194	-
2004 2/	1,831	3	18	37	61	70	113	190	294	429	405	211	-
2003 2/	1,745	1	14	47	55	64	130	176	262	389	404	203	-
2002	1,752	4	13	40	53	68	89	197	295	372	405	216	-
2001	1,741	3	13	56	60	75	118	187	248	408	390	183	-
2000	1,729	-	17	36	80	88	116	204	258	373	349	208	-
1999	1,694	2	12	52	82	67	104	174	252	405	371	173	-
Non-Hispanic black, male													
2005 2/	902	2	17	22	42	40	48	101	143	209	211	67	-
2004 2/	959	-	13	21	34	36	60	107	161	262	185	80	-
2003 2/	908	1	8	23	35	40	73	99	150	218	187	74	-
2002	914	1	6	24	30	37	48	109	161	220	198	80	-
2001	915	2	9	38	31	44	67	93	147	232	187	65	-
2000	923	-	8	22	54	50	52	107	149	216	181	84	-
1999	887	-	7	32	45	38	54	103	142	222	178	66	-
Non-Hispanic black, female													
2005 2/	795	1	8	13	22	43	39	97	111	160	174	127	-
2004 2/	872	3	5	16	27	34	53	83	133	167	220	131	-
2003 2/	837	-	6	24	20	24	57	77	112	171	217	129	-
2002	838	3	7	16	23	31	41	88	134	152	207	136	-
2001	826	1	4	18	29	31	51	94	101	176	203	118	-
2000	806	-	9	14	26	38	64	97	109	157	168	124	-
1999	807	2	5	20	37	29	50	71	110	183	193	107	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Leukemia (C91-C95)													
Origin not stated 3/, both sexes													
2005	22	-	-	1	-	1	3	5	3	4	5	-	-
2004	31	-	-	2	2	-	-	2	6	8	5	6	-
2003	39	-	-	1	-	1	-	5	6	10	8	8	-
2002	47	1	-	-	1	1	1	6	4	13	13	7	-
2001	52	-	2	1	1	3	3	4	9	12	10	7	-
2000	53	-	1	3	3	-	7	2	7	11	13	6	-
1999	39	-	1	-	-	2	1	4	6	7	13	5	-
Origin not stated 3/, male													
2005	13	-	-	1	-	-	1	2	3	2	4	-	-
2004	16	-	-	2	1	-	-	2	3	2	4	2	-
2003	30	-	-	1	-	1	-	5	5	6	6	6	-
2002	29	1	-	-	1	-	-	5	2	9	8	3	-
2001	30	-	1	-	1	-	2	3	5	7	7	4	-
2000	27	-	1	1	1	-	4	2	4	5	7	2	-
1999	17	-	1	-	-	1	1	1	2	7	4	-	-
Origin not stated 3/, female													
2005	9	-	-	-	-	1	2	3	-	2	1	-	-
2004	15	-	-	-	1	-	-	-	3	6	1	4	-
2003	9	-	-	-	-	-	-	-	1	4	2	2	-
2002	18	-	-	-	-	1	1	1	2	4	5	4	-
2001	22	-	1	1	-	3	1	1	4	5	3	3	-
2000	26	-	-	2	2	-	3	-	3	6	6	4	-
1999	22	-	-	-	-	1	-	3	4	-	9	5	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Multiple myeloma and immunoproliferative neoplasms (C88,C90)													
All origins, both sexes													
2005	11,200	-	-	-	1	11	131	688	1,842	2,969	3,846	1,712	-
2004	10,965	1	-	-	1	14	136	663	1,720	2,840	3,980	1,610	-
2003	11,267	-	-	1	-	12	147	711	1,659	3,029	3,997	1,711	-
2002	11,392	2	-	-	2	12	138	736	1,669	3,176	3,971	1,686	-
2001	11,172	-	-	-	-	9	157	671	1,671	3,181	3,886	1,597	-
2000	11,063	-	-	1	-	7	146	700	1,530	3,179	3,896	1,604	-
1999	11,016	-	-	-	-	22	156	665	1,694	3,172	3,788	1,519	-
All origins, male													
2005	5,945	-	-	-	1	8	83	409	1,021	1,678	2,006	739	-
2004	5,694	1	-	-	1	11	78	381	972	1,587	1,990	673	-
2003	5,848	-	-	-	-	7	97	426	995	1,654	1,941	728	-
2002	5,953	-	-	-	-	6	78	433	966	1,746	2,018	706	-
2001	5,724	-	-	-	-	4	97	403	945	1,725	1,926	624	-
2000	5,553	-	-	-	-	5	85	391	808	1,697	1,932	635	-
1999	5,584	-	-	-	-	16	95	378	968	1,669	1,846	612	-
All origins, female													
2005	5,255	-	-	-	-	3	48	279	821	1,291	1,840	973	-
2004	5,271	-	-	-	-	3	58	282	748	1,253	1,990	937	-
2003	5,419	-	-	1	-	5	50	285	664	1,375	2,056	983	-
2002	5,439	2	-	-	2	6	60	303	703	1,430	1,953	980	-
2001	5,448	-	-	-	-	5	60	268	726	1,456	1,960	973	-
2000	5,510	-	-	1	-	2	61	309	722	1,482	1,964	969	-
1999	5,432	-	-	-	-	6	61	287	726	1,503	1,942	907	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Multiple myeloma and immunoproliferative neoplasms (C88,C90)													
Hispanic, both sexes													
2005	588	-	-	-	1	3	10	66	128	177	156	47	-
2004	558	-	-	-	-	4	19	51	123	158	164	39	-
2003	531	-	-	-	-	-	14	53	109	154	153	48	-
2002	572	1	-	-	1	3	14	66	98	169	160	60	-
2001	524	-	-	-	-	3	18	43	107	169	137	47	-
2000	478	-	-	-	-	1	12	43	90	161	124	47	-
1999	468	-	-	-	-	4	12	51	111	139	111	40	-
Hispanic, male													
2005	296	-	-	-	1	3	5	38	56	90	79	24	-
2004	287	-	-	-	-	4	12	26	68	80	84	13	-
2003	257	-	-	-	-	-	11	31	57	72	67	19	-
2002	314	-	-	-	-	3	10	38	62	92	77	32	-
2001	291	-	-	-	-	1	13	23	52	102	80	20	-
2000	239	-	-	-	-	1	6	24	44	87	61	16	-
1999	225	-	-	-	-	4	9	28	60	67	47	10	-
Hispanic, female													
2005	292	-	-	-	-	-	5	28	72	87	77	23	-
2004	271	-	-	-	-	-	7	25	55	78	80	26	-
2003	274	-	-	-	-	-	3	22	52	82	86	29	-
2002	258	1	-	-	1	-	4	28	36	77	83	28	-
2001	233	-	-	-	-	2	5	20	55	67	57	27	-
2000	239	-	-	-	-	-	6	19	46	74	63	31	-
1999	243	-	-	-	-	-	3	23	51	72	64	30	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Multiple myeloma and immunoproliferative neoplasms (C88,C90)													
Non-Hispanic 1/, both sexes													
2005	10,603	-	-	-	-	8	121	622	1,710	2,791	3,687	1,664	-
2004	10,393	1	-	-	1	10	117	611	1,596	2,681	3,807	1,569	-
2003	10,710	-	-	1	-	12	133	657	1,547	2,867	3,834	1,659	-
2002	10,787	1	-	-	1	9	124	669	1,566	2,998	3,798	1,621	-
2001	10,618	-	-	-	-	6	138	625	1,559	3,006	3,736	1,548	-
2000	10,561	-	-	1	-	6	133	655	1,435	3,010	3,767	1,554	-
1999	10,532	-	-	-	-	18	144	614	1,581	3,030	3,670	1,475	-
Non-Hispanic 1/, male													
2005	5,646	-	-	-	-	5	78	371	965	1,588	1,925	714	-
2004	5,403	1	-	-	1	7	66	354	904	1,506	1,905	659	-
2003	5,579	-	-	-	-	7	86	394	936	1,581	1,868	707	-
2002	5,623	-	-	-	-	3	68	394	902	1,649	1,934	673	-
2001	5,418	-	-	-	-	3	84	378	890	1,620	1,840	603	-
2000	5,300	-	-	-	-	4	79	365	761	1,605	1,868	618	-
1999	5,352	-	-	-	-	12	86	350	908	1,601	1,796	599	-
Non-Hispanic 1/, female													
2005	4,957	-	-	-	-	3	43	251	745	1,203	1,762	950	-
2004	4,990	-	-	-	-	3	51	257	692	1,175	1,902	910	-
2003	5,131	-	-	1	-	5	47	263	611	1,286	1,966	952	-
2002	5,164	1	-	-	1	6	56	275	664	1,349	1,864	948	-
2001	5,200	-	-	-	-	3	54	247	669	1,386	1,896	945	-
2000	5,261	-	-	1	-	2	54	290	674	1,405	1,899	936	-
1999	5,180	-	-	-	-	6	58	264	673	1,429	1,874	876	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Multiple myeloma and immunoproliferative neoplasms (C88,C90)													
Non-Hispanic white, both sexes													
2005 2/	8,599	-	-	-	-	4	82	410	1,279	2,194	3,170	1,460	-
2004 2/	8,397	1	-	-	1	7	77	423	1,200	2,101	3,225	1,362	-
2003 2/	8,689	-	-	1	-	9	89	437	1,181	2,292	3,240	1,440	-
2002	8,843	-	-	-	1	7	78	456	1,179	2,419	3,283	1,420	-
2001	8,641	-	-	-	-	2	81	447	1,144	2,427	3,194	1,346	-
2000	8,582	-	-	1	-	3	84	458	1,072	2,442	3,220	1,302	-
1999	8,547	-	-	-	-	10	92	432	1,145	2,465	3,133	1,270	-
Non-Hispanic white, male													
2005 2/	4,649	-	-	-	-	2	58	253	739	1,286	1,674	637	-
2004 2/	4,448	1	-	-	1	5	46	251	701	1,221	1,643	579	-
2003 2/	4,629	-	-	-	-	6	62	268	730	1,311	1,623	629	-
2002	4,708	-	-	-	-	2	45	287	699	1,371	1,703	601	-
2001	4,472	-	-	-	-	2	54	280	666	1,334	1,603	533	-
2000	4,400	-	-	-	-	2	58	261	586	1,325	1,628	540	-
1999	4,430	-	-	-	-	6	56	264	688	1,329	1,563	524	-
Non-Hispanic white, female													
2005 2/	3,950	-	-	-	-	2	24	157	540	908	1,496	823	-
2004 2/	3,949	-	-	-	-	2	31	172	499	880	1,582	783	-
2003 2/	4,060	-	-	1	-	3	27	169	451	981	1,617	811	-
2002	4,135	-	-	-	1	5	33	169	480	1,048	1,580	819	-
2001	4,169	-	-	-	-	-	27	167	478	1,093	1,591	813	-
2000	4,182	-	-	1	-	1	26	197	486	1,117	1,592	762	-
1999	4,117	-	-	-	-	4	36	168	457	1,136	1,570	746	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Multiple myeloma and immunoproliferative neoplasms (C88,C90)													
Non-Hispanic black, both sexes													
2005 2/	1,766	-	-	-	-	4	34	179	376	520	469	184	-
2004 2/	1,800	-	-	-	-	2	39	167	355	523	524	190	-
2003 2/	1,833	-	-	-	-	2	43	200	326	523	536	203	-
2002	1,750	-	-	-	-	2	40	192	343	524	465	184	-
2001	1,795	-	-	-	-	4	53	159	377	519	493	190	-
2000	1,824	-	-	-	-	3	44	186	336	526	500	229	-
1999	1,786	-	-	-	-	7	49	171	391	494	478	196	-
Non-Hispanic black, male													
2005 2/	874	-	-	-	-	3	19	103	199	260	224	66	-
2004 2/	850	-	-	-	-	1	20	87	182	260	229	71	-
2003 2/	858	-	-	-	-	1	24	114	188	239	222	70	-
2002	830	-	-	-	-	1	21	99	182	257	205	65	-
2001	853	-	-	-	-	1	28	87	203	251	220	63	-
2000	829	-	-	-	-	2	17	95	163	261	222	69	-
1999	824	-	-	-	-	6	29	80	196	237	208	68	-
Non-Hispanic black, female													
2005 2/	892	-	-	-	-	1	15	76	177	260	245	118	-
2004 2/	950	-	-	-	-	1	19	80	173	263	295	119	-
2003 2/	975	-	-	-	-	1	19	86	138	284	314	133	-
2002	920	-	-	-	-	1	19	93	161	267	260	119	-
2001	942	-	-	-	-	3	25	72	174	268	273	127	-
2000	995	-	-	-	-	1	27	91	173	265	278	160	-
1999	962	-	-	-	-	1	20	91	195	257	270	128	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Multiple myeloma and immunoproliferative neoplasms (C88,C90)													
Origin not stated 3/, both sexes													
2005	9	-	-	-	-	-	-	-	4	1	3	1	-
2004	14	-	-	-	-	-	-	1	1	1	9	2	-
2003	26	-	-	-	-	-	-	1	3	8	10	4	-
2002	33	-	-	-	-	-	-	1	5	9	13	5	-
2001	30	-	-	-	-	-	1	3	5	6	13	2	-
2000	24	-	-	-	-	-	1	2	5	8	5	3	-
1999	16	-	-	-	-	-	-	-	2	3	7	4	-
Origin not stated 3/, male													
2005	3	-	-	-	-	-	-	-	-	-	2	1	-
2004	4	-	-	-	-	-	-	1	-	1	1	1	-
2003	12	-	-	-	-	-	-	1	2	1	6	2	-
2002	16	-	-	-	-	-	-	1	2	5	7	1	-
2001	15	-	-	-	-	-	-	2	3	3	6	1	-
2000	14	-	-	-	-	-	-	2	3	5	3	1	-
1999	7	-	-	-	-	-	-	-	-	1	3	3	-
Origin not stated 3/, female													
2005	6	-	-	-	-	-	-	-	4	1	1	-	-
2004	10	-	-	-	-	-	-	-	1	-	8	1	-
2003	14	-	-	-	-	-	-	-	1	7	4	2	-
2002	17	-	-	-	-	-	-	-	3	4	6	4	-
2001	15	-	-	-	-	-	1	1	2	3	7	1	-
2000	10	-	-	-	-	-	1	-	2	3	2	2	-
1999	9	-	-	-	-	-	-	-	2	2	4	1	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Other and unspecified malignant neoplasms of lymphoid, hematopoietic and related tissue (C96)													
All origins, both sexes													
2005	60	1	2	1	-	1	3	2	10	14	16	10	-
2004	71	-	-	-	-	2	2	5	12	12	26	12	-
2003	55	1	-	1	-	1	1	7	2	15	14	13	-
2002	73	3	-	-	1	-	-	7	3	18	25	16	-
2001	90	1	1	-	-	1	4	4	8	23	28	20	-
2000	100	-	2	1	1	-	4	8	12	22	29	21	-
1999	83	1	1	-	-	1	4	9	12	17	25	13	-
All origins, male													
2005	40	1	1	-	-	-	3	1	7	10	11	6	-
2004	35	-	-	-	-	2	2	4	5	4	11	7	-
2003	31	1	-	-	-	1	1	4	2	9	8	5	-
2002	44	3	-	-	1	-	-	4	2	15	12	7	-
2001	51	1	-	-	-	1	3	3	5	18	14	6	-
2000	60	-	2	1	1	-	3	6	9	12	17	9	-
1999	39	1	-	-	-	1	1	4	7	8	14	3	-
All origins, female													
2005	20	-	1	1	-	1	-	1	3	4	5	4	-
2004	36	-	-	-	-	-	-	1	7	8	15	5	-
2003	24	-	-	1	-	-	-	3	-	6	6	8	-
2002	29	-	-	-	-	-	-	3	1	3	13	9	-
2001	39	-	1	-	-	-	1	1	3	5	14	14	-
2000	40	-	-	-	-	-	1	2	3	10	12	12	-
1999	44	-	1	-	-	-	3	5	5	9	11	10	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Other and unspecified malignant neoplasms of lymphoid, hematopoietic and related tissue (C96)													
Hispanic, both sexes													
2005	6	-	2	-	-	-	1	1	2	-	-	-	-
2004	1	-	-	-	-	-	-	-	-	1	-	-	-
2003	1	-	-	-	-	1	-	-	-	-	-	-	-
2002	2	-	-	-	1	-	-	-	-	1	-	-	-
2001	2	-	-	-	-	-	-	-	-	2	-	-	-
2000	6	-	-	-	-	-	-	1	1	2	1	1	-
1999	5	-	1	-	-	1	-	-	-	1	2	-	-
Hispanic, male													
2005	5	-	1	-	-	-	1	1	2	-	-	-	-
2004	1	-	-	-	-	-	-	-	-	1	-	-	-
2003	1	-	-	-	-	1	-	-	-	-	-	-	-
2002	1	-	-	-	1	-	-	-	-	-	-	-	-
2001	1	-	-	-	-	-	-	-	-	1	-	-	-
2000	3	-	-	-	-	-	-	1	1	-	-	1	-
1999	2	-	-	-	-	1	-	-	-	-	1	-	-
Hispanic, female													
2005	1	-	1	-	-	-	-	-	-	-	-	-	-
2004	-	-	-	-	-	-	-	-	-	-	-	-	-
2003	-	-	-	-	-	-	-	-	-	-	-	-	-
2002	1	-	-	-	-	-	-	-	-	1	-	-	-
2001	1	-	-	-	-	-	-	-	-	1	-	-	-
2000	3	-	-	-	-	-	-	-	-	2	1	-	-
1999	3	-	1	-	-	-	-	-	-	1	1	-	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Other and unspecified malignant neoplasms of lymphoid, hematopoietic and related tissue (C96)													
Non-Hispanic 1/, both sexes													
2005	54	1	-	1	-	1	2	1	8	14	16	10	-
2004	70	-	-	-	-	2	2	5	12	11	26	12	-
2003	54	1	-	1	-	-	1	7	2	15	14	13	-
2002	71	3	-	-	-	-	-	7	3	17	25	16	-
2001	88	1	1	-	-	1	4	4	8	21	28	20	-
2000	94	-	2	1	1	-	4	7	11	20	28	20	-
1999	78	1	-	-	-	-	4	9	12	16	23	13	-
Non-Hispanic 1/, male													
2005	35	1	-	-	-	-	2	-	5	10	11	6	-
2004	34	-	-	-	-	2	2	4	5	3	11	7	-
2003	30	1	-	-	-	-	1	4	2	9	8	5	-
2002	43	3	-	-	-	-	-	4	2	15	12	7	-
2001	50	1	-	-	-	1	3	3	5	17	14	6	-
2000	57	-	2	1	1	-	3	5	8	12	17	8	-
1999	37	1	-	-	-	-	1	4	7	8	13	3	-
Non-Hispanic 1/, female													
2005	19	-	-	1	-	1	-	1	3	4	5	4	-
2004	36	-	-	-	-	-	-	1	7	8	15	5	-
2003	24	-	-	1	-	-	-	3	-	6	6	8	-
2002	28	-	-	-	-	-	-	3	1	2	13	9	-
2001	38	-	1	-	-	-	1	1	3	4	14	14	-
2000	37	-	-	-	-	-	1	2	3	8	11	12	-
1999	41	-	-	-	-	-	3	5	5	8	10	10	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Other and unspecified malignant neoplasms of lymphoid, hematopoietic and related tissue (C96)													
Non-Hispanic white, both sexes													
2005 2/	48	1	-	1	-	1	2	1	7	12	14	9	-
2004 2/	60	-	-	-	-	1	2	4	10	11	22	10	-
2003 2/	49	1	-	1	-	-	1	5	2	14	13	12	-
2002	59	1	-	-	-	-	-	6	3	16	20	13	-
2001	75	1	1	-	-	-	3	4	5	19	23	19	-
2000	83	-	2	1	-	-	3	5	10	19	25	18	-
1999	71	1	-	-	-	-	3	7	11	16	21	12	-
Non-Hispanic white, male													
2005 2/	31	1	-	-	-	-	2	-	4	8	10	6	-
2004 2/	31	-	-	-	-	1	2	4	5	3	10	6	-
2003 2/	27	1	-	-	-	-	1	2	2	9	7	5	-
2002	37	1	-	-	-	-	-	4	2	14	10	6	-
2001	43	1	-	-	-	-	2	3	4	16	12	5	-
2000	48	-	2	1	-	-	2	3	7	11	16	6	-
1999	34	1	-	-	-	-	1	3	6	8	12	3	-
Non-Hispanic white, female													
2005 2/	17	-	-	1	-	1	-	1	3	4	4	3	-
2004 2/	29	-	-	-	-	-	-	-	5	8	12	4	-
2003 2/	22	-	-	1	-	-	-	3	-	5	6	7	-
2002	22	-	-	-	-	-	-	2	1	2	10	7	-
2001	32	-	1	-	-	-	1	1	1	3	11	14	-
2000	35	-	-	-	-	-	1	2	3	8	9	12	-
1999	37	-	-	-	-	-	2	4	5	8	9	9	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Other and unspecified malignant neoplasms of lymphoid, hematopoietic and related tissue (C96)													
Non-Hispanic black, both sexes													
2005 2/	5	-	-	-	-	-	-	-	1	2	1	1	-
2004 2/	8	-	-	-	-	-	-	1	2	-	3	2	-
2003 2/	3	-	-	-	-	-	-	1	-	1	-	1	-
2002	6	-	-	-	-	-	-	-	-	1	3	2	-
2001	12	-	-	-	-	1	1	-	3	2	4	1	-
2000	9	-	-	-	1	-	-	2	1	-	3	2	-
1999	6	-	-	-	-	-	1	2	1	-	1	1	-
Non-Hispanic black, male													
2005 2/	3	-	-	-	-	-	-	-	1	2	-	-	-
2004 2/	1	-	-	-	-	-	-	-	-	-	-	1	-
2003 2/	1	-	-	-	-	-	-	1	-	-	-	-	-
2002	2	-	-	-	-	-	-	-	-	1	1	-	-
2001	7	-	-	-	-	1	1	-	1	1	2	1	-
2000	7	-	-	-	1	-	-	2	1	-	1	2	-
1999	3	-	-	-	-	-	-	1	1	-	1	-	-
Non-Hispanic black, female													
2005 2/	2	-	-	-	-	-	-	-	-	-	1	1	-
2004 2/	7	-	-	-	-	-	-	1	2	-	3	1	-
2003 2/	2	-	-	-	-	-	-	-	-	1	-	1	-
2002	4	-	-	-	-	-	-	-	-	-	2	2	-
2001	5	-	-	-	-	-	-	-	2	1	2	-	-
2000	2	-	-	-	-	-	-	-	-	-	2	-	-
1999	3	-	-	-	-	-	1	1	-	-	-	1	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Other and unspecified malignant neoplasms of lymphoid, hematopoietic and related tissue (C96)													
Origin not stated 3/, both sexes													
2005	-	-	-	-	-	-	-	-	-	-	-	-	-
2004	-	-	-	-	-	-	-	-	-	-	-	-	-
2003	-	-	-	-	-	-	-	-	-	-	-	-	-
2002	-	-	-	-	-	-	-	-	-	-	-	-	-
2001	-	-	-	-	-	-	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-	-	-	-
Origin not stated 3/, male													
2005	-	-	-	-	-	-	-	-	-	-	-	-	-
2004	-	-	-	-	-	-	-	-	-	-	-	-	-
2003	-	-	-	-	-	-	-	-	-	-	-	-	-
2002	-	-	-	-	-	-	-	-	-	-	-	-	-
2001	-	-	-	-	-	-	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-	-	-	-
Origin not stated 3/, female													
2005	-	-	-	-	-	-	-	-	-	-	-	-	-
2004	-	-	-	-	-	-	-	-	-	-	-	-	-
2003	-	-	-	-	-	-	-	-	-	-	-	-	-
2002	-	-	-	-	-	-	-	-	-	-	-	-	-
2001	-	-	-	-	-	-	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-	-	-	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
All other and unspecified malignant neoplasms (C17,C23-C24,C26-C31,C37-C41,C44-C49,C51-C52,C57-C60,C62-C63,C66,C68-C69,C73-C80,C97)													
All origins, both sexes													
2005	62,851	15	122	279	579	701	1,827	5,558	10,422	14,019	18,480	10,848	1
2004	61,911	23	126	275	548	696	1,876	5,276	9,997	14,187	18,549	10,356	2
2003	62,639	19	106	274	516	690	1,885	5,611	9,820	14,277	18,880	10,556	5
2002	62,969	24	124	288	527	734	2,043	5,493	9,723	14,757	18,685	10,570	1
2001	62,708	17	117	279	516	728	2,055	5,506	9,441	15,023	18,700	10,326	-
2000	63,060	24	129	260	511	701	2,090	5,307	9,293	15,716	18,860	10,165	4
1999	66,251	14	116	291	519	762	2,314	5,477	9,863	16,834	19,656	10,403	2
All origins, male													
2005	32,194	9	68	151	362	440	1,001	3,111	6,002	7,687	9,137	4,226	-
2004	31,654	13	59	158	357	420	1,014	2,945	5,679	7,798	9,281	3,929	1
2003	31,845	12	66	152	329	427	1,046	3,143	5,573	7,800	9,321	3,973	3
2002	32,122	15	65	152	340	459	1,110	3,131	5,521	8,141	9,146	4,041	1
2001	31,663	7	73	153	346	442	1,108	3,164	5,326	8,177	9,032	3,835	-
2000	31,728	13	75	140	319	432	1,141	2,959	5,223	8,572	9,161	3,689	4
1999	33,449	8	64	160	329	490	1,300	3,098	5,597	9,157	9,439	3,806	1
All origins, female													
2005	30,657	6	54	128	217	261	826	2,447	4,420	6,332	9,343	6,622	1
2004	30,257	10	67	117	191	276	862	2,331	4,318	6,389	9,268	6,427	1
2003	30,794	7	40	122	187	263	839	2,468	4,247	6,477	9,559	6,583	2
2002	30,847	9	59	136	187	275	933	2,362	4,202	6,616	9,539	6,529	-
2001	31,045	10	44	126	170	286	947	2,342	4,115	6,846	9,668	6,491	-
2000	31,332	11	54	120	192	269	949	2,348	4,070	7,144	9,699	6,476	-
1999	32,802	6	52	131	190	272	1,014	2,379	4,266	7,677	10,217	6,597	1

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
All other and unspecified malignant neoplasms (C17,C23-C24,C26-C31,C37-C41,C44-C49,C51-C52,C57-C60,C62-C63,C66,C68-C69,C73-C80,C97)													
Hispanic, both sexes													
2005	3,238	4	23	43	97	135	196	343	540	737	794	326	-
2004	2,973	2	21	41	90	102	163	357	503	669	729	296	-
2003	3,085	2	21	51	94	110	193	375	494	717	701	326	1
2002	3,005	11	17	55	107	107	176	346	502	712	676	296	-
2001	2,844	2	15	42	85	82	188	334	481	690	615	310	-
2000	2,753	3	24	47	82	100	141	290	455	690	631	290	-
1999	2,747	2	11	45	84	99	175	314	471	671	587	288	-
Hispanic, male													
2005	1,715	2	16	21	63	86	117	183	315	396	388	128	-
2004	1,521	-	6	22	70	65	93	189	268	336	367	105	-
2003	1,621	1	15	27	60	84	103	214	274	385	332	126	-
2002	1,566	7	9	30	69	69	94	189	277	362	338	122	-
2001	1,455	-	9	22	64	56	95	181	262	339	315	112	-
2000	1,394	2	16	26	49	67	78	148	235	349	309	115	-
1999	1,399	2	3	23	62	63	90	162	256	351	278	109	-
Hispanic, female													
2005	1,523	2	7	22	34	49	79	160	225	341	406	198	-
2004	1,452	2	15	19	20	37	70	168	235	333	362	191	-
2003	1,464	1	6	24	34	26	90	161	220	332	369	200	1
2002	1,439	4	8	25	38	38	82	157	225	350	338	174	-
2001	1,389	2	6	20	21	26	93	153	219	351	300	198	-
2000	1,359	1	8	21	33	33	63	142	220	341	322	175	-
1999	1,348	-	8	22	22	36	85	152	215	320	309	179	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
All other and unspecified malignant neoplasms (C17,C23-C24,C26-C31,C37-C41,C44-C49,C51-C52,C57-C60,C62-C63,C66,C68-C69,C73-C80,C97)													
Non-Hispanic 1/, both sexes													
2005	59,519	10	99	235	482	563	1,627	5,203	9,862	13,258	17,668	10,511	1
2004	58,818	20	104	234	457	590	1,712	4,910	9,467	13,490	17,792	10,040	2
2003	59,402	17	84	223	420	578	1,686	5,216	9,289	13,529	18,150	10,206	4
2002	59,797	13	106	233	419	625	1,867	5,129	9,192	14,005	17,965	10,242	1
2001	59,685	15	102	237	431	645	1,858	5,155	8,935	14,278	18,042	9,987	-
2000	60,117	21	103	212	427	596	1,939	4,997	8,813	14,980	18,172	9,853	4
1999	63,328	12	105	245	434	655	2,130	5,149	9,361	16,121	19,020	10,094	2
Non-Hispanic 1/, male													
2005	30,420	7	52	129	299	352	881	2,918	5,674	7,277	8,738	4,093	-
2004	30,066	12	52	136	287	353	921	2,748	5,393	7,444	8,902	3,817	1
2003	30,140	11	50	125	268	341	940	2,916	5,275	7,396	8,976	3,839	3
2002	30,463	8	56	122	270	389	1,016	2,929	5,225	7,755	8,782	3,910	1
2001	30,107	7	64	131	282	385	1,006	2,971	5,046	7,800	8,701	3,714	-
2000	30,237	11	57	114	269	362	1,059	2,803	4,972	8,195	8,823	3,568	4
1999	31,956	6	61	137	266	423	1,206	2,928	5,325	8,779	9,132	3,692	1
Non-Hispanic 1/, female													
2005	29,099	3	47	106	183	211	746	2,285	4,188	5,981	8,930	6,418	1
2004	28,752	8	52	98	170	237	791	2,162	4,074	6,046	8,890	6,223	1
2003	29,262	6	34	98	152	237	746	2,300	4,014	6,133	9,174	6,367	1
2002	29,334	5	50	111	149	236	851	2,200	3,967	6,250	9,183	6,332	-
2001	29,578	8	38	106	149	260	852	2,184	3,889	6,478	9,341	6,273	-
2000	29,880	10	46	98	158	234	880	2,194	3,841	6,785	9,349	6,285	-
1999	31,372	6	44	108	168	232	924	2,221	4,036	7,342	9,888	6,402	1

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
All other and unspecified malignant neoplasms (C17,C23-C24,C26-C31,C37-C41,C44-C49,C51-C52,C57-C60,C62-C63,C66,C68-C69,C73-C80,C97)													
Non-Hispanic white, both sexes													
2005 2/	51,380	8	73	166	372	401	1,254	4,029	8,150	11,386	15,908	9,633	-
2004 2/	50,810	16	75	167	332	465	1,308	3,785	7,818	11,615	16,001	9,226	2
2003 2/	51,534	13	58	165	328	424	1,287	4,066	7,743	11,712	16,368	9,366	4
2002	51,731	11	73	171	319	465	1,433	4,032	7,635	12,071	16,159	9,361	1
2001	51,581	13	83	176	338	492	1,430	4,024	7,342	12,292	16,233	9,158	-
2000	52,223	16	75	168	328	479	1,452	3,908	7,330	13,019	16,395	9,050	3
1999	54,977	10	87	178	324	513	1,626	4,077	7,722	14,076	17,158	9,204	2
Non-Hispanic white, male													
2005 2/	26,332	5	40	89	231	258	697	2,285	4,722	6,274	7,952	3,779	-
2004 2/	26,000	9	40	102	209	296	705	2,114	4,457	6,442	8,101	3,524	1
2003 2/	26,189	9	31	94	209	261	729	2,281	4,419	6,440	8,169	3,544	3
2002	26,457	8	40	88	201	295	769	2,332	4,362	6,790	7,966	3,605	1
2001	26,007	6	50	96	220	296	792	2,326	4,139	6,789	7,882	3,411	-
2000	26,258	9	46	89	204	303	797	2,213	4,111	7,208	7,989	3,286	3
1999	27,780	5	52	104	190	338	943	2,305	4,423	7,714	8,322	3,383	1
Non-Hispanic white, female													
2005 2/	25,048	3	33	77	141	143	557	1,744	3,428	5,112	7,956	5,854	-
2004 2/	24,810	7	35	65	123	169	603	1,671	3,361	5,173	7,900	5,702	1
2003 2/	25,345	4	27	71	119	163	558	1,785	3,324	5,272	8,199	5,822	1
2002	25,274	3	33	83	118	170	664	1,700	3,273	5,281	8,193	5,756	-
2001	25,574	7	33	80	118	196	638	1,698	3,203	5,503	8,351	5,747	-
2000	25,965	7	29	79	124	176	655	1,695	3,219	5,811	8,406	5,764	-
1999	27,197	5	35	74	134	175	683	1,772	3,299	6,362	8,836	5,821	1

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
All other and unspecified malignant neoplasms (C17,C23-C24,C26-C31,C37-C41,C44-C49,C51-C52,C57-C60,C62-C63,C66,C68-C69,C73-C80,C97)													
Non-Hispanic black, both sexes													
2005 2/	6,591	2	19	61	80	127	303	976	1,404	1,497	1,445	676	1
2004 2/	6,582	3	19	60	101	96	339	944	1,367	1,550	1,449	654	-
2003 2/	6,537	2	22	51	70	122	326	978	1,316	1,498	1,458	694	-
2002	6,756	2	26	52	79	125	362	924	1,329	1,612	1,507	738	-
2001	6,725	1	16	46	76	124	339	963	1,333	1,659	1,468	700	-
2000	6,617	4	22	39	83	87	427	927	1,255	1,636	1,472	664	1
1999	7,009	1	14	49	83	106	421	928	1,407	1,704	1,539	757	-
Non-Hispanic black, male													
2005 2/	3,311	2	9	36	47	75	151	526	798	808	626	233	-
2004 2/	3,347	2	7	29	61	47	174	545	784	827	651	220	-
2003 2/	3,274	2	15	27	45	60	171	542	743	786	658	225	-
2002	3,340	-	12	28	54	67	200	509	739	813	675	243	-
2001	3,392	-	13	26	51	69	166	543	759	866	650	249	-
2000	3,337	1	9	24	56	46	227	510	733	827	687	216	1
1999	3,514	-	8	22	56	60	219	539	801	895	667	247	-
Non-Hispanic black, female													
2005 2/	3,280	-	10	25	33	52	152	450	606	689	819	443	1
2004 2/	3,235	1	12	31	40	49	165	399	583	723	798	434	-
2003 2/	3,263	-	7	24	25	62	155	436	573	712	800	469	-
2002	3,416	2	14	24	25	58	162	415	590	799	832	495	-
2001	3,333	1	3	20	25	55	173	420	574	793	818	451	-
2000	3,280	3	13	15	27	41	200	417	522	809	785	448	-
1999	3,495	1	6	27	27	46	202	389	606	809	872	510	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
All other and unspecified malignant neoplasms (C17,C23-C24,C26-C31,C37-C41,C44-C49,C51-C52,C57-C60,C62-C63,C66,C68-C69,C73-C80,C97)													
Origin not stated 3/, both sexes													
2005	94	1	-	1	-	3	4	12	20	24	18	11	-
2004	120	1	1	-	1	4	1	9	27	28	28	20	-
2003	152	-	1	-	2	2	6	20	37	31	29	24	-
2002	167	-	1	-	1	2	-	18	29	40	44	32	-
2001	179	-	-	-	-	1	9	17	25	55	43	29	-
2000	190	-	2	1	2	5	10	20	25	46	57	22	-
1999	176	-	-	1	1	8	9	14	31	42	49	21	-
Origin not stated 3/, male													
2005	59	-	-	1	-	2	3	10	13	14	11	5	-
2004	67	1	1	-	-	2	-	8	18	18	12	7	-
2003	84	-	1	-	1	2	3	13	24	19	13	8	-
2002	93	-	-	-	1	1	-	13	19	24	26	9	-
2001	101	-	-	-	-	1	7	12	18	38	16	9	-
2000	97	-	2	-	1	3	4	8	16	28	29	6	-
1999	94	-	-	-	1	4	4	8	16	27	29	5	-
Origin not stated 3/, female													
2005	35	1	-	-	-	1	1	2	7	10	7	6	-
2004	53	-	-	-	1	2	1	1	9	10	16	13	-
2003	68	-	-	-	1	-	3	7	13	12	16	16	-
2002	74	-	1	-	-	1	-	5	10	16	18	23	-
2001	78	-	-	-	-	-	2	5	7	17	27	20	-
2000	93	-	-	1	1	2	6	12	9	18	28	16	-
1999	82	-	-	1	-	4	5	6	15	15	20	16	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)													
All origins, both sexes													
2005	13,710	59	52	76	93	162	368	729	1,318	2,350	4,594	3,909	-
2004	13,580	71	53	84	110	173	318	698	1,310	2,343	4,644	3,774	2
2003	13,563	62	51	79	96	178	339	682	1,238	2,354	4,689	3,795	-
2002	13,299	69	60	89	87	163	366	714	1,193	2,348	4,597	3,613	-
2001	13,676	51	58	105	102	151	366	719	1,253	2,574	4,654	3,643	-
2000	13,546	60	53	99	95	165	368	690	1,224	2,546	4,645	3,600	1
1999	13,227	58	63	101	98	171	342	662	1,133	2,684	4,472	3,443	-
All origins, male													
2005	6,820	22	31	36	54	86	186	404	754	1,342	2,419	1,486	-
2004	6,764	35	28	46	63	94	172	382	750	1,322	2,378	1,492	2
2003	6,687	32	31	43	54	98	195	366	685	1,334	2,379	1,470	-
2002	6,536	30	30	49	50	84	196	365	680	1,315	2,332	1,405	-
2001	6,735	26	30	52	53	77	193	391	673	1,439	2,403	1,398	-
2000	6,559	32	33	49	62	82	188	385	657	1,415	2,319	1,336	1
1999	6,520	30	33	49	54	96	197	362	656	1,522	2,218	1,303	-
All origins, female													
2005	6,890	37	21	40	39	76	182	325	564	1,008	2,175	2,423	-
2004	6,816	36	25	38	47	79	146	316	560	1,021	2,266	2,282	-
2003	6,876	30	20	36	42	80	144	316	553	1,020	2,310	2,325	-
2002	6,763	39	30	40	37	79	170	349	513	1,033	2,265	2,208	-
2001	6,941	25	28	53	49	74	173	328	580	1,135	2,251	2,245	-
2000	6,987	28	20	50	33	83	180	305	567	1,131	2,326	2,264	-
1999	6,707	28	30	52	44	75	145	300	477	1,162	2,254	2,140	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)													
Hispanic, both sexes													
2005	572	16	12	10	13	24	28	55	75	107	152	80	-
2004	617	18	5	8	15	29	32	54	84	110	160	102	-
2003	603	15	7	13	21	32	25	42	64	106	185	93	-
2002	563	17	7	26	8	16	34	54	60	105	148	88	-
2001	543	5	13	21	8	20	39	50	67	100	128	92	-
2000	521	6	7	13	10	25	37	43	53	98	123	106	-
1999	497	8	12	17	12	25	30	39	60	109	111	74	-
Hispanic, male													
2005	306	7	7	7	8	14	11	33	44	63	82	30	-
2004	311	11	2	4	11	15	18	33	43	55	78	41	-
2003	301	6	3	7	13	21	14	22	39	54	88	34	-
2002	282	9	4	15	5	8	15	26	35	54	77	34	-
2001	276	2	7	12	6	6	24	30	35	50	67	37	-
2000	237	4	2	8	8	15	14	18	25	46	54	43	-
1999	233	5	4	8	8	14	21	26	30	47	47	23	-
Hispanic, female													
2005	266	9	5	3	5	10	17	22	31	44	70	50	-
2004	306	7	3	4	4	14	14	21	41	55	82	61	-
2003	302	9	4	6	8	11	11	20	25	52	97	59	-
2002	281	8	3	11	3	8	19	28	25	51	71	54	-
2001	267	3	6	9	2	14	15	20	32	50	61	55	-
2000	284	2	5	5	2	10	23	25	28	52	69	63	-
1999	264	3	8	9	4	11	9	13	30	62	64	51	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)													
Non-Hispanic 1/, both sexes													
2005	13,118	42	40	66	80	138	340	673	1,241	2,240	4,434	3,824	-
2004	12,939	53	48	76	95	144	286	643	1,222	2,225	4,477	3,669	1
2003	12,938	46	44	66	75	146	313	640	1,170	2,241	4,499	3,698	-
2002	12,708	52	53	63	78	147	328	659	1,131	2,241	4,436	3,520	-
2001	13,098	46	45	84	94	131	327	667	1,182	2,468	4,510	3,544	-
2000	12,990	53	46	85	85	140	330	645	1,167	2,445	4,511	3,482	1
1999	12,699	50	51	83	85	146	311	620	1,070	2,572	4,349	3,362	-
Non-Hispanic 1/, male													
2005	6,501	14	24	29	46	72	175	370	708	1,278	2,332	1,453	-
2004	6,440	24	26	42	52	79	154	349	704	1,263	2,295	1,451	1
2003	6,372	26	28	36	41	77	180	344	645	1,275	2,286	1,434	-
2002	6,237	21	26	34	44	76	179	339	645	1,260	2,245	1,368	-
2001	6,442	24	23	40	47	71	169	360	636	1,385	2,328	1,359	-
2000	6,302	28	31	41	54	67	173	365	629	1,367	2,258	1,288	1
1999	6,276	25	29	41	46	82	176	334	623	1,474	2,168	1,278	-
Non-Hispanic 1/, female													
2005	6,617	28	16	37	34	66	165	303	533	962	2,102	2,371	-
2004	6,499	29	22	34	43	65	132	294	518	962	2,182	2,218	-
2003	6,566	20	16	30	34	69	133	296	525	966	2,213	2,264	-
2002	6,471	31	27	29	34	71	149	320	486	981	2,191	2,152	-
2001	6,656	22	22	44	47	60	158	307	546	1,083	2,182	2,185	-
2000	6,688	25	15	44	31	73	157	280	538	1,078	2,253	2,194	-
1999	6,423	25	22	42	39	64	135	286	447	1,098	2,181	2,084	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)													
Non-Hispanic white, both sexes													
2005 2/	11,670	28	31	47	63	105	254	524	1,034	1,958	4,062	3,564	-
2004 2/	11,469	41	39	52	72	100	210	503	1,008	1,949	4,081	3,413	1
2003 2/	11,572	33	32	43	58	113	236	508	995	1,970	4,147	3,437	-
2002	11,332	42	36	41	59	105	251	530	962	1,985	4,063	3,258	-
2001	11,701	32	36	56	66	106	252	561	1,006	2,185	4,150	3,251	-
2000	11,636	41	37	63	63	105	255	516	1,004	2,144	4,171	3,237	-
1999	11,335	34	36	66	58	99	239	494	899	2,276	4,001	3,133	-
Non-Hispanic white, male													
2005 2/	5,850	10	20	22	39	55	135	294	599	1,143	2,165	1,368	-
2004 2/	5,754	19	21	29	41	59	115	279	585	1,128	2,111	1,366	1
2003 2/	5,732	20	19	24	34	57	143	281	544	1,142	2,125	1,343	-
2002	5,607	18	17	22	32	62	149	283	547	1,135	2,069	1,273	-
2001	5,801	18	18	25	31	58	135	312	547	1,250	2,152	1,255	-
2000	5,653	21	23	31	40	55	132	301	539	1,202	2,108	1,201	-
1999	5,653	17	18	32	30	53	143	276	536	1,326	2,013	1,209	-
Non-Hispanic white, female													
2005 2/	5,820	18	11	25	24	50	119	230	435	815	1,897	2,196	-
2004 2/	5,715	22	18	23	31	41	95	224	423	821	1,970	2,047	-
2003 2/	5,840	13	13	19	24	56	93	227	451	828	2,022	2,094	-
2002	5,725	24	19	19	27	43	102	247	415	850	1,994	1,985	-
2001	5,900	14	18	31	35	48	117	249	459	935	1,998	1,996	-
2000	5,983	20	14	32	23	50	123	215	465	942	2,063	2,036	-
1999	5,682	17	18	34	28	46	96	218	363	950	1,988	1,924	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)													
Non-Hispanic black, both sexes													
2005 2/	1,142	12	5	14	11	27	77	123	164	225	280	204	-
2004 2/	1,181	10	7	21	19	38	66	111	177	220	311	201	-
2003 2/	1,111	11	9	17	15	28	67	117	143	222	284	198	-
2002	1,134	9	14	18	15	31	67	112	144	211	300	213	-
2001	1,143	11	5	26	21	18	61	89	147	236	296	233	-
2000	1,122	9	9	18	16	33	62	112	133	245	282	202	1
1999	1,161	14	13	17	25	30	65	104	148	256	295	194	-
Non-Hispanic black, male													
2005 2/	493	4	2	4	5	13	35	62	84	102	123	59	-
2004 2/	547	4	4	12	10	16	33	57	100	110	140	61	-
2003 2/	516	5	8	8	6	17	31	54	82	114	127	64	-
2002	496	2	8	9	9	8	25	46	83	98	134	74	-
2001	522	5	3	13	13	11	30	38	73	119	141	76	-
2000	514	4	8	8	9	11	32	54	74	130	118	65	1
1999	516	7	10	9	15	20	30	44	75	128	124	54	-
Non-Hispanic black, female													
2005 2/	649	8	3	10	6	14	42	61	80	123	157	145	-
2004 2/	634	6	3	9	9	22	33	54	77	110	171	140	-
2003 2/	595	6	1	9	9	11	36	63	61	108	157	134	-
2002	638	7	6	9	6	23	42	66	61	113	166	139	-
2001	621	6	2	13	8	7	31	51	74	117	155	157	-
2000	608	5	1	10	7	22	30	58	59	115	164	137	-
1999	645	7	3	8	10	10	35	60	73	128	171	140	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)													
Origin not stated 3/, both sexes													
2005	20	1	-	-	-	-	-	1	2	3	8	5	-
2004	24	-	-	-	-	-	-	1	4	8	7	3	1
2003	22	1	-	-	-	-	1	-	4	7	5	4	-
2002	28	-	-	-	1	-	4	1	2	2	13	5	-
2001	35	-	-	-	-	-	-	2	4	6	16	7	-
2000	35	1	-	1	-	-	1	2	4	3	11	12	-
1999	31	-	-	1	1	-	1	3	3	3	12	7	-
Origin not stated 3/, male													
2005	13	1	-	-	-	-	-	1	2	1	5	3	-
2004	13	-	-	-	-	-	-	-	3	4	5	-	1
2003	14	-	-	-	-	-	1	-	1	5	5	2	-
2002	17	-	-	-	1	-	2	-	-	1	10	3	-
2001	17	-	-	-	-	-	-	1	2	4	8	2	-
2000	20	-	-	-	-	-	1	2	3	2	7	5	-
1999	11	-	-	-	-	-	-	2	3	1	3	2	-
Origin not stated 3/, female													
2005	7	-	-	-	-	-	-	-	-	2	3	2	-
2004	11	-	-	-	-	-	-	1	1	4	2	3	-
2003	8	1	-	-	-	-	-	-	3	2	-	2	-
2002	11	-	-	-	-	-	2	1	2	1	3	2	-
2001	18	-	-	-	-	-	-	1	2	2	8	5	-
2000	15	1	-	1	-	-	-	-	1	1	4	7	-
1999	20	-	-	1	1	-	1	1	-	2	9	5	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Anemias (D50-D64)													
All origins, both sexes													
2005	4,624	19	23	26	103	157	189	253	304	494	1,193	1,863	-
2004	4,575	15	33	46	128	164	201	236	309	473	1,119	1,850	1
2003	4,594	22	24	47	105	157	183	241	276	484	1,136	1,919	-
2002	4,614	18	21	53	111	180	196	250	264	508	1,161	1,852	-
2001	4,627	16	22	55	109	160	215	232	249	492	1,226	1,851	-
2000	4,494	10	35	52	85	153	190	227	260	515	1,171	1,796	-
1999	4,503	12	30	53	121	160	194	232	231	486	1,164	1,820	-
All origins, male													
2005	1,787	13	9	14	57	85	97	108	141	224	502	537	-
2004	1,833	9	18	21	69	90	107	112	164	226	466	550	1
2003	1,832	11	14	23	52	83	91	125	156	251	477	549	-
2002	1,781	12	9	26	67	101	96	109	131	244	454	532	-
2001	1,774	11	12	27	59	95	91	115	111	237	496	520	-
2000	1,669	6	16	29	39	70	78	108	130	236	465	492	-
1999	1,746	8	17	23	67	81	101	116	105	226	484	518	-
All origins, female													
2005	2,837	6	14	12	46	72	92	145	163	270	691	1,326	-
2004	2,742	6	15	25	59	74	94	124	145	247	653	1,300	-
2003	2,762	11	10	24	53	74	92	116	120	233	659	1,370	-
2002	2,833	6	12	27	44	79	100	141	133	264	707	1,320	-
2001	2,853	5	10	28	50	65	124	117	138	255	730	1,331	-
2000	2,825	4	19	23	46	83	112	119	130	279	706	1,304	-
1999	2,757	4	13	30	54	79	93	116	126	260	680	1,302	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Anemias (D50-D64)													
Hispanic, both sexes													
2005	184	6	4	4	7	8	7	10	9	32	47	50	-
2004	193	2	5	3	9	12	10	17	15	18	43	59	-
2003	191	2	5	4	5	7	9	12	18	25	48	56	-
2002	208	2	3	4	8	11	11	12	14	37	44	62	-
2001	185	-	3	5	11	12	12	11	17	25	43	46	-
2000	144	1	1	7	5	13	8	10	13	24	36	26	-
1999	164	-	6	3	6	8	10	9	10	27	35	50	-
Hispanic, male													
2005	81	4	1	4	5	7	3	3	6	10	21	17	-
2004	75	1	3	2	1	5	4	10	8	4	20	17	-
2003	84	2	2	2	2	5	3	6	11	12	19	20	-
2002	106	1	1	3	5	5	9	7	8	17	18	32	-
2001	82	-	2	5	6	8	6	4	7	12	21	11	-
2000	55	1	-	3	1	5	1	4	4	12	15	9	-
1999	79	-	4	1	3	5	6	4	3	17	17	19	-
Hispanic, female													
2005	103	2	3	-	2	1	4	7	3	22	26	33	-
2004	118	1	2	1	8	7	6	7	7	14	23	42	-
2003	107	-	3	2	3	2	6	6	7	13	29	36	-
2002	102	1	2	1	3	6	2	5	6	20	26	30	-
2001	103	-	1	-	5	4	6	7	10	13	22	35	-
2000	89	-	1	4	4	8	7	6	9	12	21	17	-
1999	85	-	2	2	3	3	4	5	7	10	18	31	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Anemias (D50-D64)													
Non-Hispanic 1/, both sexes													
2005	4,431	13	19	22	96	149	181	243	292	462	1,145	1,809	-
2004	4,371	13	28	43	119	152	190	219	292	454	1,071	1,790	-
2003	4,395	20	19	43	99	150	173	228	258	457	1,087	1,861	-
2002	4,391	14	17	49	103	167	182	237	250	470	1,114	1,788	-
2001	4,424	15	19	50	97	146	202	219	231	466	1,177	1,802	-
2000	4,336	8	34	44	79	140	181	215	246	489	1,132	1,768	-
1999	4,321	12	24	50	115	152	183	221	220	455	1,125	1,764	-
Non-Hispanic 1/, male													
2005	1,703	9	8	10	52	78	94	105	133	214	481	519	-
2004	1,751	8	15	19	68	85	102	102	154	222	443	533	-
2003	1,745	9	12	21	49	78	87	118	145	239	458	529	-
2002	1,671	9	8	23	62	96	86	102	123	227	435	500	-
2001	1,687	10	10	22	53	86	84	110	104	225	474	509	-
2000	1,608	4	16	26	37	65	77	104	126	222	448	483	-
1999	1,659	8	13	22	64	76	95	111	101	207	465	497	-
Non-Hispanic 1/, female													
2005	2,728	4	11	12	44	71	87	138	159	248	664	1,290	-
2004	2,620	5	13	24	51	67	88	117	138	232	628	1,257	-
2003	2,650	11	7	22	50	72	86	110	113	218	629	1,332	-
2002	2,720	5	9	26	41	71	96	135	127	243	679	1,288	-
2001	2,737	5	9	28	44	60	118	109	127	241	703	1,293	-
2000	2,728	4	18	18	42	75	104	111	120	267	684	1,285	-
1999	2,662	4	11	28	51	76	88	110	119	248	660	1,267	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Anemias (D50-D64)													
Non-Hispanic white, both sexes													
2005 2/	3,314	3	7	7	15	17	32	105	192	340	975	1,621	-
2004 2/	3,269	6	8	17	29	24	45	82	169	362	935	1,592	-
2003 2/	3,287	7	5	17	18	32	36	80	135	348	939	1,670	-
2002	3,253	5	10	19	23	33	34	90	147	361	944	1,587	-
2001	3,350	5	8	22	22	27	41	81	150	368	1,013	1,613	-
2000	3,299	3	14	14	16	16	36	82	155	384	994	1,585	-
1999	3,222	6	4	17	28	10	37	86	112	360	995	1,567	-
Non-Hispanic white, male													
2005 2/	1,208	2	3	1	5	9	14	47	89	161	408	469	-
2004 2/	1,247	3	3	6	18	14	26	45	87	183	380	482	-
2003 2/	1,236	2	3	9	9	17	19	45	74	178	405	475	-
2002	1,164	2	4	8	13	20	18	45	75	172	371	436	-
2001	1,220	4	3	11	12	12	15	37	68	175	419	464	-
2000	1,178	1	8	9	7	10	20	39	83	185	392	424	-
1999	1,162	4	2	6	11	2	18	46	45	163	416	449	-
Non-Hispanic white, female													
2005 2/	2,106	1	4	6	10	8	18	58	103	179	567	1,152	-
2004 2/	2,022	3	5	11	11	10	19	37	82	179	555	1,110	-
2003 2/	2,051	5	2	8	9	15	17	35	61	170	534	1,195	-
2002	2,089	3	6	11	10	13	16	45	72	189	573	1,151	-
2001	2,130	1	5	11	10	15	26	44	82	193	594	1,149	-
2000	2,121	2	6	5	9	6	16	43	72	199	602	1,161	-
1999	2,060	2	2	11	17	8	19	40	67	197	579	1,118	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Anemias (D50-D64)													
Non-Hispanic black, both sexes													
2005 2/	1,024	6	11	15	77	131	144	135	90	109	147	159	-
2004 2/	1,013	4	19	23	90	126	139	132	114	81	115	170	-
2003 2/	1,029	6	12	25	78	114	133	143	116	94	139	169	-
2002	1,048	5	7	26	78	127	142	141	94	100	153	175	-
2001	993	6	11	26	72	117	154	135	75	90	149	158	-
2000	961	4	19	30	61	124	139	128	83	89	121	163	-
1999	1,010	3	20	33	84	139	142	125	100	81	119	164	-
Non-Hispanic black, male													
2005 2/	452	6	5	9	45	69	77	58	34	47	62	40	-
2004 2/	465	3	12	12	50	70	74	55	62	35	49	43	-
2003 2/	483	4	7	12	39	58	68	73	67	55	53	47	-
2002	472	4	4	14	49	72	66	55	46	51	58	53	-
2001	443	4	7	10	41	73	66	73	35	46	51	37	-
2000	392	3	8	17	29	55	55	64	37	31	45	48	-
1999	453	2	11	16	50	72	73	61	53	37	41	37	-
Non-Hispanic black, female													
2005 2/	572	-	6	6	32	62	67	77	56	62	85	119	-
2004 2/	548	1	7	11	40	56	65	77	52	46	66	127	-
2003 2/	546	2	5	13	39	56	65	70	49	39	86	122	-
2002	576	1	3	12	29	55	76	86	48	49	95	122	-
2001	550	2	4	16	31	44	88	62	40	44	98	121	-
2000	569	1	11	13	32	69	84	64	46	58	76	115	-
1999	557	1	9	17	34	67	69	64	47	44	78	127	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Anemias (D50-D64)													
Origin not stated 3/, both sexes													
2005	9	-	-	-	-	-	1	-	3	-	1	4	-
2004	11	-	-	-	-	-	1	-	2	1	5	1	1
2003	8	-	-	-	1	-	1	1	-	2	1	2	-
2002	15	2	1	-	-	2	3	1	-	1	3	2	-
2001	18	1	-	-	1	2	1	2	1	1	6	3	-
2000	14	1	-	1	1	-	1	2	1	2	3	2	-
1999	18	-	-	-	-	-	1	2	1	4	4	6	-
Origin not stated 3/, male													
2005	3	-	-	-	-	-	-	-	2	-	-	1	-
2004	7	-	-	-	-	-	1	-	2	-	3	-	1
2003	3	-	-	-	1	-	1	1	-	-	-	-	-
2002	4	2	-	-	-	-	1	-	-	-	1	-	-
2001	5	1	-	-	-	1	1	1	-	-	1	-	-
2000	6	1	-	-	1	-	-	-	-	2	2	-	-
1999	8	-	-	-	-	-	-	1	1	2	2	2	-
Origin not stated 3/, female													
2005	6	-	-	-	-	-	1	-	1	-	1	3	-
2004	4	-	-	-	-	-	-	-	-	1	2	1	-
2003	5	-	-	-	-	-	-	-	-	2	1	2	-
2002	11	-	1	-	-	2	2	1	-	1	2	2	-
2001	13	-	-	-	1	1	-	1	1	1	5	3	-
2000	8	-	-	1	-	-	1	2	1	-	1	2	-
1999	10	-	-	-	-	-	1	1	-	2	2	4	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Diabetes mellitus (E10-E14)													
All origins, both sexes													
2005	75,119	1	2	38	202	617	2,045	5,691	11,301	16,183	23,136	15,903	-
2004	73,138	1	8	37	161	599	2,026	5,567	10,780	16,093	22,945	14,918	3
2003	74,219	3	8	31	160	657	2,049	5,658	10,731	16,656	23,299	14,964	3
2002	73,249	1	3	34	171	642	2,164	5,496	10,022	16,709	23,282	14,724	1
2001	71,372	2	6	39	151	595	1,958	5,343	9,570	16,731	22,805	14,171	1
2000	69,301	4	6	26	162	623	1,926	4,954	9,186	16,674	22,184	13,556	-
1999	68,399	2	8	32	155	582	1,942	4,735	9,097	16,908	21,757	13,178	3
All origins, male													
2005	36,538	-	-	15	115	372	1,270	3,467	6,570	8,505	10,701	5,523	-
2004	35,267	1	5	16	83	340	1,258	3,376	6,153	8,430	10,569	5,035	1
2003	35,438	-	6	16	96	379	1,264	3,341	6,044	8,687	10,679	4,923	3
2002	34,301	-	1	22	98	385	1,330	3,225	5,683	8,417	10,295	4,844	1
2001	32,841	1	2	23	85	341	1,181	3,154	5,241	8,437	9,918	4,457	1
2000	31,602	4	2	13	94	376	1,151	2,821	4,943	8,226	9,738	4,234	-
1999	31,150	-	4	16	80	319	1,139	2,791	4,856	8,279	9,539	4,125	2
All origins, female													
2005	38,581	1	2	23	87	245	775	2,224	4,731	7,678	12,435	10,380	-
2004	37,871	-	3	21	78	259	768	2,191	4,627	7,663	12,376	9,883	2
2003	38,781	3	2	15	64	278	785	2,317	4,687	7,969	12,620	10,041	-
2002	38,948	1	2	12	73	257	834	2,271	4,339	8,292	12,987	9,880	-
2001	38,531	1	4	16	66	254	777	2,189	4,329	8,294	12,887	9,714	-
2000	37,699	-	4	13	68	247	775	2,133	4,243	8,448	12,446	9,322	-
1999	37,249	2	4	16	75	263	803	1,944	4,241	8,629	12,218	9,053	1

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Diabetes mellitus (E10-E14)													
Hispanic, both sexes													
2005	6,665	-	-	3	14	69	216	632	1,168	1,670	1,947	946	-
2004	5,999	-	2	-	13	64	187	535	1,073	1,587	1,769	769	-
2003	6,179	-	1	3	19	64	197	593	1,121	1,611	1,743	826	1
2002	5,912	-	-	1	15	56	205	513	1,029	1,591	1,712	790	-
2001	5,663	1	1	2	19	49	167	507	982	1,604	1,548	783	-
2000	5,328	1	-	4	13	64	154	439	950	1,592	1,410	701	-
1999	5,182	-	-	4	14	43	157	413	927	1,551	1,369	704	-
Hispanic, male													
2005	3,296	-	-	2	9	40	153	378	662	850	877	325	-
2004	2,855	-	2	-	8	37	111	323	602	759	781	232	-
2003	2,976	-	1	1	11	33	121	345	618	793	775	277	1
2002	2,779	-	-	1	10	34	124	305	551	752	748	254	-
2001	2,590	-	1	2	11	28	109	311	510	758	638	222	-
2000	2,507	1	-	2	7	48	95	240	520	762	609	223	-
1999	2,335	-	-	2	8	21	84	252	480	709	562	217	-
Hispanic, female													
2005	3,369	-	-	1	5	29	63	254	506	820	1,070	621	-
2004	3,144	-	-	-	5	27	76	212	471	828	988	537	-
2003	3,203	-	-	2	8	31	76	248	503	818	968	549	-
2002	3,133	-	-	-	5	22	81	208	478	839	964	536	-
2001	3,073	1	-	-	8	21	58	196	472	846	910	561	-
2000	2,821	-	-	2	6	16	59	199	430	830	801	478	-
1999	2,847	-	-	2	6	22	73	161	447	842	807	487	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Diabetes mellitus (E10-E14)													
Non-Hispanic 1/, both sexes													
2005	68,325	1	2	35	188	548	1,823	5,045	10,104	14,491	21,149	14,939	-
2004	66,953	1	6	37	148	532	1,833	5,014	9,669	14,471	21,128	14,111	3
2003	67,862	3	7	27	138	592	1,847	5,046	9,583	15,014	21,498	14,105	2
2002	67,112	1	3	33	155	584	1,946	4,961	8,958	15,070	21,499	13,901	1
2001	65,498	1	5	37	131	542	1,787	4,810	8,562	15,066	21,198	13,358	1
2000	63,769	3	6	22	149	557	1,762	4,501	8,207	15,031	20,717	12,814	-
1999	63,017	2	8	28	140	534	1,781	4,305	8,138	15,318	20,325	12,436	2
Non-Hispanic 1/, male													
2005	33,178	-	-	13	106	332	1,113	3,079	5,892	7,645	9,805	5,193	-
2004	32,315	1	3	16	75	301	1,143	3,041	5,527	7,653	9,763	4,791	1
2003	32,368	-	5	15	82	345	1,139	2,986	5,408	7,876	9,875	4,635	2
2002	31,399	-	1	21	87	349	1,196	2,906	5,112	7,639	9,511	4,576	1
2001	30,144	1	1	21	74	311	1,069	2,828	4,715	7,643	9,255	4,225	1
2000	28,994	3	2	11	87	328	1,048	2,574	4,408	7,436	9,098	3,999	-
1999	28,719	-	4	14	72	295	1,053	2,525	4,358	7,548	8,955	3,894	1
Non-Hispanic 1/, female													
2005	35,147	1	2	22	82	216	710	1,966	4,212	6,846	11,344	9,746	-
2004	34,638	-	3	21	73	231	690	1,973	4,142	6,818	11,365	9,320	2
2003	35,494	3	2	12	56	247	708	2,060	4,175	7,138	11,623	9,470	-
2002	35,713	1	2	12	68	235	750	2,055	3,846	7,431	11,988	9,325	-
2001	35,354	-	4	16	57	231	718	1,982	3,847	7,423	11,943	9,133	-
2000	34,775	-	4	11	62	229	714	1,927	3,799	7,595	11,619	8,815	-
1999	34,298	2	4	14	68	239	728	1,780	3,780	7,770	11,370	8,542	1

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Diabetes mellitus (E10-E14)													
Non-Hispanic white, both sexes													
2005 2/	53,159	1	1	22	110	306	1,215	3,454	7,293	10,826	17,293	12,638	-
2004 2/	52,101	-	4	19	87	333	1,238	3,462	6,965	10,875	17,220	11,896	2
2003 2/	52,894	2	5	19	90	366	1,240	3,412	6,873	11,316	17,601	11,968	2
2002	52,463	1	3	19	95	382	1,345	3,385	6,403	11,337	17,692	11,801	-
2001	51,482	-	5	19	66	335	1,233	3,332	6,118	11,464	17,507	11,402	1
2000	50,163	2	6	13	92	357	1,260	3,078	5,923	11,475	17,040	10,917	-
1999	49,347	2	6	19	94	363	1,231	2,940	5,674	11,664	16,821	10,533	-
Non-Hispanic white, male													
2005 2/	26,365	-	-	9	64	189	760	2,179	4,384	5,941	8,302	4,537	-
2004 2/	25,773	-	2	7	45	191	795	2,184	4,128	5,984	8,276	4,161	-
2003 2/	25,940	-	4	12	54	212	794	2,122	4,041	6,206	8,417	4,076	2
2002	25,273	-	1	16	51	231	849	2,054	3,815	6,041	8,151	4,064	-
2001	24,302	-	1	13	42	184	767	2,014	3,524	6,142	7,874	3,740	1
2000	23,465	2	2	7	57	220	766	1,842	3,298	5,967	7,796	3,508	-
1999	23,177	-	3	9	52	203	747	1,801	3,200	6,066	7,722	3,374	-
Non-Hispanic white, female													
2005 2/	26,794	1	1	13	46	117	455	1,275	2,909	4,885	8,991	8,101	-
2004 2/	26,328	-	2	12	42	142	443	1,278	2,837	4,891	8,944	7,735	2
2003 2/	26,954	2	1	7	36	154	446	1,290	2,832	5,110	9,184	7,892	-
2002	27,190	1	2	3	44	151	496	1,331	2,588	5,296	9,541	7,737	-
2001	27,180	-	4	6	24	151	466	1,318	2,594	5,322	9,633	7,662	-
2000	26,698	-	4	6	35	137	494	1,236	2,625	5,508	9,244	7,409	-
1999	26,170	2	3	10	42	160	484	1,139	2,474	5,598	9,099	7,159	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Diabetes mellitus (E10-E14)													
Non-Hispanic black, both sexes													
2005 2/	12,835	-	1	12	73	218	534	1,396	2,415	3,101	3,205	1,880	-
2004 2/	12,685	1	1	16	60	180	540	1,371	2,304	3,078	3,269	1,864	1
2003 2/	12,787	1	2	7	45	208	547	1,431	2,347	3,145	3,265	1,789	-
2002	12,583	-	-	14	57	194	554	1,393	2,204	3,152	3,213	1,801	1
2001	12,167	1	-	15	56	198	506	1,306	2,133	3,110	3,132	1,710	-
2000	11,918	1	-	9	55	187	458	1,270	1,995	3,083	3,193	1,667	-
1999	11,830	-	2	9	43	160	502	1,219	2,136	3,120	2,995	1,642	2
Non-Hispanic black, male													
2005 2/	5,666	-	-	3	39	127	307	772	1,274	1,413	1,236	495	-
2004 2/	5,498	1	-	8	30	101	314	748	1,171	1,417	1,207	500	1
2003 2/	5,380	-	1	3	26	122	309	741	1,169	1,392	1,188	429	-
2002	5,158	-	-	5	34	112	316	744	1,104	1,320	1,135	387	1
2001	4,983	1	-	6	27	122	275	701	1,031	1,278	1,150	392	-
2000	4,720	1	-	4	28	98	252	648	952	1,248	1,100	389	-
1999	4,718	-	1	5	19	85	276	641	986	1,262	1,016	426	1
Non-Hispanic black, female													
2005 2/	7,169	-	1	9	34	91	227	624	1,141	1,688	1,969	1,385	-
2004 2/	7,187	-	1	8	30	79	226	623	1,133	1,661	2,062	1,364	-
2003 2/	7,407	1	1	4	19	86	238	690	1,178	1,753	2,077	1,360	-
2002	7,425	-	-	9	23	82	238	649	1,100	1,832	2,078	1,414	-
2001	7,184	-	-	9	29	76	231	605	1,102	1,832	1,982	1,318	-
2000	7,198	-	-	5	27	89	206	622	1,043	1,835	2,093	1,278	-
1999	7,112	-	1	4	24	75	226	578	1,150	1,858	1,979	1,216	1

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Diabetes mellitus (E10-E14)													
Origin not stated 3/, both sexes													
2005	129	-	-	-	-	-	6	14	29	22	40	18	-
2004	186	-	-	-	-	3	6	18	38	35	48	38	-
2003	178	-	-	1	3	1	5	19	27	31	58	33	-
2002	225	-	-	-	1	2	13	22	35	48	71	33	-
2001	211	-	-	-	1	4	4	26	26	61	59	30	-
2000	204	-	-	-	-	2	10	14	29	51	57	41	-
1999	200	-	-	-	1	5	4	17	32	39	63	38	1
Origin not stated 3/, male													
2005	64	-	-	-	-	-	4	10	16	10	19	5	-
2004	97	-	-	-	-	2	4	12	24	18	25	12	-
2003	94	-	-	-	3	1	4	10	18	18	29	11	-
2002	123	-	-	-	1	2	10	14	20	26	36	14	-
2001	107	-	-	-	-	2	3	15	16	36	25	10	-
2000	101	-	-	-	-	-	8	7	15	28	31	12	-
1999	96	-	-	-	-	3	2	14	18	22	22	14	1
Origin not stated 3/, female													
2005	65	-	-	-	-	-	2	4	13	12	21	13	-
2004	89	-	-	-	-	1	2	6	14	17	23	26	-
2003	84	-	-	1	-	-	1	9	9	13	29	22	-
2002	102	-	-	-	-	-	3	8	15	22	35	19	-
2001	104	-	-	-	1	2	1	11	10	25	34	20	-
2000	103	-	-	-	-	2	2	7	14	23	26	29	-
1999	104	-	-	-	1	2	2	3	14	17	41	24	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Nutritional deficiencies (E40-E64)													
All origins, both sexes													
2005	3,183	5	3	4	3	19	38	118	157	353	915	1,568	-
2004	3,019	7	2	3	6	13	39	117	162	298	897	1,474	1
2003	3,338	10	2	2	7	10	35	100	168	340	992	1,672	-
2002	3,779	4	1	1	7	16	60	102	168	382	1,132	1,906	-
2001	3,704	4	4	2	10	14	44	110	153	374	1,100	1,889	-
2000	4,242	14	2	5	10	18	56	99	173	460	1,285	2,120	-
1999	4,289	9	2	1	8	12	64	106	170	429	1,248	2,240	-
All origins, male													
2005	1,116	3	1	1	1	7	14	56	88	157	341	447	-
2004	1,087	5	1	2	5	6	15	50	87	145	357	413	1
2003	1,176	7	2	2	4	7	19	48	79	166	400	442	-
2002	1,321	3	1	-	4	7	31	48	93	186	434	514	-
2001	1,233	4	1	2	7	4	21	56	81	174	426	457	-
2000	1,412	10	2	2	2	9	16	46	97	221	471	536	-
1999	1,486	4	2	1	5	4	28	52	98	221	494	577	-
All origins, female													
2005	2,067	2	2	3	2	12	24	62	69	196	574	1,121	-
2004	1,932	2	1	1	1	7	24	67	75	153	540	1,061	-
2003	2,162	3	-	-	3	3	16	52	89	174	592	1,230	-
2002	2,458	1	-	1	3	9	29	54	75	196	698	1,392	-
2001	2,471	-	3	-	3	10	23	54	72	200	674	1,432	-
2000	2,830	4	-	3	8	9	40	53	76	239	814	1,584	-
1999	2,803	5	-	-	3	8	36	54	72	208	754	1,663	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Nutritional deficiencies (E40-E64)													
Hispanic, both sexes													
2005	139	-	-	-	-	1	2	7	5	20	50	54	-
2004	107	1	-	1	-	1	2	4	8	17	32	41	-
2003	126	2	2	-	-	1	1	3	7	19	36	55	-
2002	141	2	-	-	3	1	3	5	4	21	49	53	-
2001	136	-	1	1	-	-	7	3	5	12	45	62	-
2000	147	-	-	-	1	3	4	5	7	24	43	60	-
1999	138	-	-	-	-	-	4	6	7	17	31	73	-
Hispanic, male													
2005	59	-	-	-	-	1	1	3	2	12	21	19	-
2004	36	-	-	1	-	-	-	2	3	9	11	10	-
2003	54	2	2	-	-	1	-	2	6	10	13	18	-
2002	61	1	-	-	1	1	3	3	2	13	19	18	-
2001	53	-	-	1	-	-	2	1	4	2	20	23	-
2000	57	-	-	-	-	2	1	3	3	13	17	18	-
1999	54	-	-	-	-	-	2	5	2	8	13	24	-
Hispanic, female													
2005	80	-	-	-	-	-	1	4	3	8	29	35	-
2004	71	1	-	-	-	1	2	2	5	8	21	31	-
2003	72	-	-	-	-	-	1	1	1	9	23	37	-
2002	80	1	-	-	2	-	-	2	2	8	30	35	-
2001	83	-	1	-	-	-	5	2	1	10	25	39	-
2000	90	-	-	-	1	1	3	2	4	11	26	42	-
1999	84	-	-	-	-	-	2	1	5	9	18	49	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Nutritional deficiencies (E40-E64)													
Non-Hispanic 1/, both sexes													
2005	3,034	5	3	4	3	18	36	111	152	328	862	1,512	-
2004	2,906	6	2	2	6	12	37	113	153	281	862	1,432	-
2003	3,204	8	-	2	7	9	34	96	161	320	952	1,615	-
2002	3,621	2	1	1	4	15	56	97	164	358	1,077	1,846	-
2001	3,563	4	3	1	9	14	37	107	148	362	1,054	1,824	-
2000	4,080	14	2	5	9	15	52	94	166	435	1,233	2,055	-
1999	4,142	9	2	1	8	12	60	100	162	411	1,215	2,162	-
Non-Hispanic 1/, male													
2005	1,052	3	1	1	1	6	13	53	86	143	318	427	-
2004	1,047	5	1	1	5	6	15	48	83	136	345	402	-
2003	1,118	5	-	2	4	6	19	45	73	155	385	424	-
2002	1,255	2	1	-	3	6	28	45	91	173	412	494	-
2001	1,178	4	1	1	6	4	19	55	77	172	405	434	-
2000	1,348	10	2	2	2	7	15	43	94	208	448	517	-
1999	1,428	4	2	1	5	4	26	47	95	213	479	552	-
Non-Hispanic 1/, female													
2005	1,982	2	2	3	2	12	23	58	66	185	544	1,085	-
2004	1,859	1	1	1	1	6	22	65	70	145	517	1,030	-
2003	2,086	3	-	-	3	3	15	51	88	165	567	1,191	-
2002	2,366	-	-	1	1	9	28	52	73	185	665	1,352	-
2001	2,385	-	2	-	3	10	18	52	71	190	649	1,390	-
2000	2,732	4	-	3	7	8	37	51	72	227	785	1,538	-
1999	2,714	5	-	-	3	8	34	53	67	198	736	1,610	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Nutritional deficiencies (E40-E64)													
Non-Hispanic white, both sexes													
2005 2/	2,537	2	2	3	1	10	26	81	117	260	712	1,323	-
2004 2/	2,421	4	1	1	3	9	23	78	116	217	732	1,237	-
2003 2/	2,699	4	-	2	5	7	28	62	122	257	811	1,401	-
2002	3,064	1	1	-	3	12	37	76	117	280	922	1,615	-
2001	3,015	3	3	-	7	10	22	70	113	291	878	1,618	-
2000	3,469	8	1	2	7	6	36	68	123	345	1,062	1,811	-
1999	3,486	5	1	-	5	9	45	70	110	307	1,018	1,916	-
Non-Hispanic white, male													
2005 2/	865	1	-	-	-	3	10	39	62	115	260	375	-
2004 2/	847	3	-	-	3	5	7	31	60	101	289	348	-
2003 2/	915	3	-	2	3	5	16	29	50	121	317	369	-
2002	1,039	1	1	-	2	5	19	31	57	137	348	438	-
2001	948	3	1	-	6	3	10	27	52	131	340	375	-
2000	1,112	6	1	2	2	2	10	30	62	159	382	456	-
1999	1,142	2	1	-	3	3	18	34	61	150	392	478	-
Non-Hispanic white, female													
2005 2/	1,672	1	2	3	1	7	16	42	55	145	452	948	-
2004 2/	1,574	1	1	1	-	4	16	47	56	116	443	889	-
2003 2/	1,784	1	-	-	2	2	12	33	72	136	494	1,032	-
2002	2,025	-	-	-	1	7	18	45	60	143	574	1,177	-
2001	2,067	-	2	-	1	7	12	43	61	160	538	1,243	-
2000	2,357	2	-	-	5	4	26	38	61	186	680	1,355	-
1999	2,344	3	-	-	2	6	27	36	49	157	626	1,438	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Nutritional deficiencies (E40-E64)													
Non-Hispanic black, both sexes													
2005 2/	426	3	1	1	1	7	9	28	34	58	126	158	-
2004 2/	412	1	1	1	2	3	12	32	34	58	104	164	-
2003 2/	448	3	-	-	1	2	4	32	37	54	127	188	-
2002	495	-	-	1	1	3	16	20	46	70	139	199	-
2001	472	1	-	1	2	4	15	33	29	61	149	177	-
2000	537	5	1	3	2	8	16	24	41	78	147	212	-
1999	585	4	1	1	3	3	15	27	47	90	175	219	-
Non-Hispanic black, male													
2005 2/	153	2	1	1	1	2	3	14	23	22	45	39	-
2004 2/	170	1	1	1	2	1	7	16	21	32	43	45	-
2003 2/	181	1	-	-	-	1	2	16	22	31	62	46	-
2002	185	-	-	-	1	1	7	14	33	31	56	42	-
2001	202	1	-	1	-	1	9	25	21	38	55	51	-
2000	214	3	1	-	-	5	5	13	30	44	58	55	-
1999	250	2	1	1	2	1	8	12	30	55	74	64	-
Non-Hispanic black, female													
2005 2/	273	1	-	-	-	5	6	14	11	36	81	119	-
2004 2/	242	-	-	-	-	2	5	16	13	26	61	119	-
2003 2/	267	2	-	-	1	1	2	16	15	23	65	142	-
2002	310	-	-	1	-	2	9	6	13	39	83	157	-
2001	270	-	-	-	2	3	6	8	8	23	94	126	-
2000	323	2	-	3	2	3	11	11	11	34	89	157	-
1999	335	2	-	-	1	2	7	15	17	35	101	155	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Nutritional deficiencies (E40-E64)													
Origin not stated 3/, both sexes													
2005	10	-	-	-	-	-	-	-	-	5	3	2	-
2004	6	-	-	-	-	-	-	-	1	-	3	1	1
2003	8	-	-	-	-	-	-	1	-	1	4	2	-
2002	17	-	-	-	-	-	1	-	-	3	6	7	-
2001	5	-	-	-	1	-	-	-	-	-	1	3	-
2000	15	-	-	-	-	-	-	-	-	1	9	5	-
1999	9	-	-	-	-	-	-	-	1	1	2	5	-
Origin not stated 3/, male													
2005	5	-	-	-	-	-	-	-	-	2	2	1	-
2004	4	-	-	-	-	-	-	-	1	-	1	1	1
2003	4	-	-	-	-	-	-	1	-	1	2	-	-
2002	5	-	-	-	-	-	-	-	-	-	3	2	-
2001	2	-	-	-	1	-	-	-	-	-	1	-	-
2000	7	-	-	-	-	-	-	-	-	-	6	1	-
1999	4	-	-	-	-	-	-	-	1	-	2	1	-
Origin not stated 3/, female													
2005	5	-	-	-	-	-	-	-	-	3	1	1	-
2004	2	-	-	-	-	-	-	-	-	-	2	-	-
2003	4	-	-	-	-	-	-	-	-	-	2	2	-
2002	12	-	-	-	-	-	1	-	-	3	3	5	-
2001	3	-	-	-	-	-	-	-	-	-	-	3	-
2000	8	-	-	-	-	-	-	-	-	1	3	4	-
1999	5	-	-	-	-	-	-	-	-	1	-	4	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Malnutrition (E40-E46)													
All origins, both sexes													
2005	3,003	5	2	3	3	18	36	113	149	333	866	1,475	-
2004	2,847	6	1	2	5	11	35	109	152	288	864	1,373	1
2003	3,153	5	1	1	6	8	33	94	160	320	952	1,573	-
2002	3,510	2	1	1	6	13	55	91	156	354	1,067	1,764	-
2001	3,454	2	-	2	9	13	39	99	146	347	1,025	1,772	-
2000	3,926	11	1	3	8	16	53	90	158	423	1,199	1,964	-
1999	3,985	7	2	1	7	11	61	99	154	407	1,159	2,077	-
All origins, male													
2005	1,059	3	-	1	1	6	14	53	83	149	327	422	-
2004	1,026	5	1	2	5	6	13	48	81	141	342	381	1
2003	1,098	4	1	1	3	6	19	47	75	155	379	408	-
2002	1,233	1	1	-	4	5	28	42	84	174	406	488	-
2001	1,148	2	-	2	6	4	19	47	75	157	400	436	-
2000	1,302	8	1	1	2	8	15	41	86	207	434	499	-
1999	1,371	4	2	1	4	4	27	48	92	207	452	530	-
All origins, female													
2005	1,944	2	2	2	2	12	22	60	66	184	539	1,053	-
2004	1,821	1	-	-	-	5	22	61	71	147	522	992	-
2003	2,055	1	-	-	3	2	14	47	85	165	573	1,165	-
2002	2,277	1	-	1	2	8	27	49	72	180	661	1,276	-
2001	2,306	-	-	-	3	9	20	52	71	190	625	1,336	-
2000	2,624	3	-	2	6	8	38	49	72	216	765	1,465	-
1999	2,614	3	-	-	3	7	34	51	62	200	707	1,547	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Malnutrition (E40-E46)													
Hispanic, both sexes													
2005	132	-	-	-	-	1	2	7	5	18	48	51	-
2004	104	1	-	1	-	1	2	3	8	17	30	41	-
2003	112	1	1	-	-	1	-	3	7	19	30	50	-
2002	136	1	-	-	2	1	3	4	4	21	49	51	-
2001	132	-	-	1	-	-	6	3	5	12	44	61	-
2000	137	-	-	-	1	2	4	4	7	22	39	58	-
1999	132	-	-	-	-	-	4	6	7	17	30	68	-
Hispanic, male													
2005	57	-	-	-	-	1	1	3	2	12	20	18	-
2004	36	-	-	1	-	-	-	2	3	9	11	10	-
2003	47	1	1	-	-	1	-	2	6	10	10	16	-
2002	58	-	-	-	1	1	3	2	2	13	19	17	-
2001	52	-	-	1	-	-	2	1	4	2	20	22	-
2000	52	-	-	-	-	2	1	2	3	12	16	16	-
1999	50	-	-	-	-	-	2	5	2	8	12	21	-
Hispanic, female													
2005	75	-	-	-	-	-	1	4	3	6	28	33	-
2004	68	1	-	-	-	1	2	1	5	8	19	31	-
2003	65	-	-	-	-	-	-	1	1	9	20	34	-
2002	78	1	-	-	1	-	-	2	2	8	30	34	-
2001	80	-	-	-	-	-	4	2	1	10	24	39	-
2000	85	-	-	-	1	-	3	2	4	10	23	42	-
1999	82	-	-	-	-	-	2	1	5	9	18	47	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Malnutrition (E40-E46)													
Non-Hispanic 1/, both sexes													
2005	2,861	5	2	3	3	17	34	106	144	310	815	1,422	-
2004	2,737	5	1	1	5	10	33	106	143	271	831	1,331	-
2003	3,034	4	-	1	6	7	33	90	153	300	918	1,522	-
2002	3,358	1	1	1	4	12	51	87	152	330	1,013	1,706	-
2001	3,317	2	-	1	8	13	33	96	141	335	980	1,708	-
2000	3,774	11	1	3	7	14	49	86	151	400	1,151	1,901	-
1999	3,844	7	2	1	7	11	57	93	146	389	1,127	2,004	-
Non-Hispanic 1/, male													
2005	997	3	-	1	1	5	13	50	81	135	305	403	-
2004	986	5	1	1	5	6	13	46	77	132	330	370	-
2003	1,047	3	-	1	3	5	19	44	69	144	367	392	-
2002	1,171	1	1	-	3	4	25	40	82	161	385	469	-
2001	1,094	2	-	1	5	4	17	46	71	155	379	414	-
2000	1,243	8	1	1	2	6	14	39	83	195	412	482	-
1999	1,317	4	2	1	4	4	25	43	89	199	438	508	-
Non-Hispanic 1/, female													
2005	1,864	2	2	2	2	12	21	56	63	175	510	1,019	-
2004	1,751	-	-	-	-	4	20	60	66	139	501	961	-
2003	1,987	1	-	-	3	2	14	46	84	156	551	1,130	-
2002	2,187	-	-	1	1	8	26	47	70	169	628	1,237	-
2001	2,223	-	-	-	3	9	16	50	70	180	601	1,294	-
2000	2,531	3	-	2	5	8	35	47	68	205	739	1,419	-
1999	2,527	3	-	-	3	7	32	50	57	190	689	1,496	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Malnutrition (E40-E46)													
Non-Hispanic white, both sexes													
2005 2/	2,385	2	2	2	1	10	25	77	110	243	670	1,243	-
2004 2/	2,268	3	-	-	3	7	21	73	109	209	703	1,140	-
2003 2/	2,545	1	-	1	5	5	28	57	115	239	779	1,315	-
2002	2,836	1	1	-	3	9	34	67	108	262	864	1,487	-
2001	2,806	1	-	-	6	9	19	64	109	270	816	1,512	-
2000	3,204	7	-	1	6	6	33	62	112	312	993	1,672	-
1999	3,232	4	1	-	5	8	43	67	99	287	945	1,773	-
Non-Hispanic white, male													
2005 2/	818	1	-	-	-	3	10	36	58	107	249	354	-
2004 2/	792	3	-	-	3	5	6	29	55	98	275	318	-
2003 2/	852	1	-	1	3	4	16	28	47	111	301	340	-
2002	970	1	1	-	2	3	17	27	51	129	325	414	-
2001	881	1	-	-	5	3	8	23	49	119	316	357	-
2000	1,021	5	-	1	2	2	9	27	55	147	350	423	-
1999	1,047	2	1	-	3	3	17	31	58	138	359	435	-
Non-Hispanic white, female													
2005 2/	1,567	1	2	2	1	7	15	41	52	136	421	889	-
2004 2/	1,476	-	-	-	-	2	15	44	54	111	428	822	-
2003 2/	1,693	-	-	-	2	1	12	29	68	128	478	975	-
2002	1,866	-	-	-	1	6	17	40	57	133	539	1,073	-
2001	1,925	-	-	-	1	6	11	41	60	151	500	1,155	-
2000	2,183	2	-	-	4	4	24	35	57	165	643	1,249	-
1999	2,185	2	-	-	2	5	26	36	41	149	586	1,338	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Malnutrition (E40-E46)													
Non-Hispanic black, both sexes													
2005 2/	408	3	-	1	1	7	8	27	33	57	121	150	-
2004 2/	398	1	1	1	2	3	10	30	31	56	102	161	-
2003 2/	435	2	-	-	1	2	4	31	36	52	125	182	-
2002	465	-	-	1	1	3	14	19	44	60	134	189	-
2001	438	1	-	1	2	4	14	28	26	56	139	167	-
2000	505	3	1	2	1	7	16	23	37	77	136	202	-
1999	543	3	1	1	2	3	14	24	42	88	161	204	-
Non-Hispanic black, male													
2005 2/	146	2	-	1	1	2	3	14	22	22	43	36	-
2004 2/	164	1	1	1	2	1	6	16	20	31	42	43	-
2003 2/	175	1	-	-	-	1	2	16	21	30	60	44	-
2002	172	-	-	-	1	1	6	13	31	27	52	41	-
2001	186	1	-	1	-	1	9	20	18	34	53	49	-
2000	202	2	1	-	-	4	5	12	26	43	56	53	-
1999	234	2	1	1	1	1	8	11	27	53	66	63	-
Non-Hispanic black, female													
2005 2/	262	1	-	-	-	5	5	13	11	35	78	114	-
2004 2/	234	-	-	-	-	2	4	14	11	25	60	118	-
2003 2/	260	1	-	-	1	1	2	15	15	22	65	138	-
2002	293	-	-	1	-	2	8	6	13	33	82	148	-
2001	252	-	-	-	2	3	5	8	8	22	86	118	-
2000	303	1	-	2	1	3	11	11	11	34	80	149	-
1999	309	1	-	-	1	2	6	13	15	35	95	141	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Malnutrition (E40-E46)													
Origin not stated 3/, both sexes													
2005	10	-	-	-	-	-	-	-	-	5	3	2	-
2004	6	-	-	-	-	-	-	-	1	-	3	1	1
2003	7	-	-	-	-	-	-	1	-	1	4	1	-
2002	16	-	-	-	-	-	1	-	-	3	5	7	-
2001	5	-	-	-	1	-	-	-	-	-	1	3	-
2000	15	-	-	-	-	-	-	-	-	1	9	5	-
1999	9	-	-	-	-	-	-	-	1	1	2	5	-
Origin not stated 3/, male													
2005	5	-	-	-	-	-	-	-	-	2	2	1	-
2004	4	-	-	-	-	-	-	-	1	-	1	1	1
2003	4	-	-	-	-	-	-	1	-	1	2	-	-
2002	4	-	-	-	-	-	-	-	-	-	2	2	-
2001	2	-	-	-	1	-	-	-	-	-	1	-	-
2000	7	-	-	-	-	-	-	-	-	-	6	1	-
1999	4	-	-	-	-	-	-	-	1	-	2	1	-
Origin not stated 3/, female													
2005	5	-	-	-	-	-	-	-	-	3	1	1	-
2004	2	-	-	-	-	-	-	-	-	-	2	-	-
2003	3	-	-	-	-	-	-	-	-	-	2	1	-
2002	12	-	-	-	-	-	1	-	-	3	3	5	-
2001	3	-	-	-	-	-	-	-	-	-	-	3	-
2000	8	-	-	-	-	-	-	-	-	1	3	4	-
1999	5	-	-	-	-	-	-	-	-	1	-	4	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Other nutritional deficiencies (E50-E64)													
All origins, both sexes													
2005	180	-	1	1	-	1	2	5	8	20	49	93	-
2004	172	1	1	1	1	2	4	8	10	10	33	101	-
2003	185	5	1	1	1	2	2	6	8	20	40	99	-
2002	269	2	-	-	1	3	5	11	12	28	65	142	-
2001	250	2	4	-	1	1	5	11	7	27	75	117	-
2000	316	3	1	2	2	2	3	9	15	37	86	156	-
1999	304	2	-	-	1	1	3	7	16	22	89	163	-
All origins, male													
2005	57	-	1	-	-	1	-	3	5	8	14	25	-
2004	61	-	-	-	-	-	2	2	6	4	15	32	-
2003	78	3	1	1	1	1	-	1	4	11	21	34	-
2002	88	2	-	-	-	2	3	6	9	12	28	26	-
2001	85	2	1	-	1	-	2	9	6	17	26	21	-
2000	110	2	1	1	-	1	1	5	11	14	37	37	-
1999	115	-	-	-	1	-	1	4	6	14	42	47	-
All origins, female													
2005	123	-	-	1	-	-	2	2	3	12	35	68	-
2004	111	1	1	1	1	2	2	6	4	6	18	69	-
2003	107	2	-	-	-	1	2	5	4	9	19	65	-
2002	181	-	-	-	1	1	2	5	3	16	37	116	-
2001	165	-	3	-	-	1	3	2	1	10	49	96	-
2000	206	1	-	1	2	1	2	4	4	23	49	119	-
1999	189	2	-	-	-	1	2	3	10	8	47	116	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Other nutritional deficiencies (E50-E64)													
Hispanic, both sexes													
2005	7	-	-	-	-	-	-	-	-	2	2	3	-
2004	3	-	-	-	-	-	-	1	-	-	2	-	-
2003	14	1	1	-	-	-	1	-	-	-	6	5	-
2002	5	1	-	-	1	-	-	1	-	-	-	2	-
2001	4	-	1	-	-	-	1	-	-	-	1	1	-
2000	10	-	-	-	-	1	-	1	-	2	4	2	-
1999	6	-	-	-	-	-	-	-	-	-	1	5	-
Hispanic, male													
2005	2	-	-	-	-	-	-	-	-	-	1	1	-
2004	-	-	-	-	-	-	-	-	-	-	-	-	-
2003	7	1	1	-	-	-	-	-	-	-	3	2	-
2002	3	1	-	-	-	-	-	1	-	-	-	1	-
2001	1	-	-	-	-	-	-	-	-	-	-	1	-
2000	5	-	-	-	-	-	-	1	-	1	1	2	-
1999	4	-	-	-	-	-	-	-	-	-	1	3	-
Hispanic, female													
2005	5	-	-	-	-	-	-	-	-	2	1	2	-
2004	3	-	-	-	-	-	-	1	-	-	2	-	-
2003	7	-	-	-	-	-	1	-	-	-	3	3	-
2002	2	-	-	-	1	-	-	-	-	-	-	1	-
2001	3	-	1	-	-	-	1	-	-	-	1	-	-
2000	5	-	-	-	-	1	-	-	-	1	3	-	-
1999	2	-	-	-	-	-	-	-	-	-	-	2	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Other nutritional deficiencies (E50-E64)													
Non-Hispanic 1/, both sexes													
2005	173	-	1	1	-	1	2	5	8	18	47	90	-
2004	169	1	1	1	1	2	4	7	10	10	31	101	-
2003	170	4	-	1	1	2	1	6	8	20	34	93	-
2002	263	1	-	-	-	3	5	10	12	28	64	140	-
2001	246	2	3	-	1	1	4	11	7	27	74	116	-
2000	306	3	1	2	2	1	3	8	15	35	82	154	-
1999	298	2	-	-	1	1	3	7	16	22	88	158	-
Non-Hispanic 1/, male													
2005	55	-	1	-	-	1	-	3	5	8	13	24	-
2004	61	-	-	-	-	-	2	2	6	4	15	32	-
2003	71	2	-	1	1	1	-	1	4	11	18	32	-
2002	84	1	-	-	-	2	3	5	9	12	27	25	-
2001	84	2	1	-	1	-	2	9	6	17	26	20	-
2000	105	2	1	1	-	1	1	4	11	13	36	35	-
1999	111	-	-	-	1	-	1	4	6	14	41	44	-
Non-Hispanic 1/, female													
2005	118	-	-	1	-	-	2	2	3	10	34	66	-
2004	108	1	1	1	1	2	2	5	4	6	16	69	-
2003	99	2	-	-	-	1	1	5	4	9	16	61	-
2002	179	-	-	-	-	1	2	5	3	16	37	115	-
2001	162	-	2	-	-	1	2	2	1	10	48	96	-
2000	201	1	-	1	2	-	2	4	4	22	46	119	-
1999	187	2	-	-	-	1	2	3	10	8	47	114	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Other nutritional deficiencies (E50-E64)													
Non-Hispanic white, both sexes													
2005 2/	152	-	-	1	-	-	1	4	7	17	42	80	-
2004 2/	153	1	1	1	-	2	2	5	7	8	29	97	-
2003 2/	154	3	-	1	-	2	-	5	7	18	32	86	-
2002	228	-	-	-	-	3	3	9	9	18	58	128	-
2001	209	2	3	-	1	1	3	6	4	21	62	106	-
2000	265	1	1	1	1	-	3	6	11	33	69	139	-
1999	254	1	-	-	-	1	2	3	11	20	73	143	-
Non-Hispanic white, male													
2005 2/	47	-	-	-	-	-	-	3	4	8	11	21	-
2004 2/	55	-	-	-	-	-	1	2	5	3	14	30	-
2003 2/	63	2	-	1	-	1	-	1	3	10	16	29	-
2002	69	-	-	-	-	2	2	4	6	8	23	24	-
2001	67	2	1	-	1	-	2	4	3	12	24	18	-
2000	91	1	1	1	-	-	1	3	7	12	32	33	-
1999	95	-	-	-	-	-	1	3	3	12	33	43	-
Non-Hispanic white, female													
2005 2/	105	-	-	1	-	-	1	1	3	9	31	59	-
2004 2/	98	1	1	1	-	2	1	3	2	5	15	67	-
2003 2/	91	1	-	-	-	1	-	4	4	8	16	57	-
2002	159	-	-	-	-	1	1	5	3	10	35	104	-
2001	142	-	2	-	-	1	1	2	1	9	38	88	-
2000	174	-	-	-	1	-	2	3	4	21	37	106	-
1999	159	1	-	-	-	1	1	-	8	8	40	100	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Other nutritional deficiencies (E50-E64)													
Non-Hispanic black, both sexes													
2005 2/	18	-	1	-	-	-	1	1	1	1	5	8	-
2004 2/	14	-	-	-	-	-	2	2	3	2	2	3	-
2003 2/	13	1	-	-	-	-	-	1	1	2	2	6	-
2002	30	-	-	-	-	-	2	1	2	10	5	10	-
2001	34	-	-	-	-	-	1	5	3	5	10	10	-
2000	32	2	-	1	1	1	-	1	4	1	11	10	-
1999	42	1	-	-	1	-	1	3	5	2	14	15	-
Non-Hispanic black, male													
2005 2/	7	-	1	-	-	-	-	-	1	-	2	3	-
2004 2/	6	-	-	-	-	-	1	-	1	1	1	2	-
2003 2/	6	-	-	-	-	-	-	-	1	1	2	2	-
2002	13	-	-	-	-	-	1	1	2	4	4	1	-
2001	16	-	-	-	-	-	-	5	3	4	2	2	-
2000	12	1	-	-	-	1	-	1	4	1	2	2	-
1999	16	-	-	-	1	-	-	1	3	2	8	1	-
Non-Hispanic black, female													
2005 2/	11	-	-	-	-	-	1	1	-	1	3	5	-
2004 2/	8	-	-	-	-	-	1	2	2	1	1	1	-
2003 2/	7	1	-	-	-	-	-	1	-	1	-	4	-
2002	17	-	-	-	-	-	1	-	-	6	1	9	-
2001	18	-	-	-	-	-	1	-	-	1	8	8	-
2000	20	1	-	1	1	-	-	-	-	-	9	8	-
1999	26	1	-	-	-	-	1	2	2	-	6	14	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Other nutritional deficiencies (E50-E64)													
Origin not stated 3/, both sexes													
2005	-	-	-	-	-	-	-	-	-	-	-	-	-
2004	-	-	-	-	-	-	-	-	-	-	-	-	-
2003	1	-	-	-	-	-	-	-	-	-	-	1	-
2002	1	-	-	-	-	-	-	-	-	-	1	-	-
2001	-	-	-	-	-	-	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-	-	-	-
Origin not stated 3/, male													
2005	-	-	-	-	-	-	-	-	-	-	-	-	-
2004	-	-	-	-	-	-	-	-	-	-	-	-	-
2003	-	-	-	-	-	-	-	-	-	-	-	-	-
2002	1	-	-	-	-	-	-	-	-	-	1	-	-
2001	-	-	-	-	-	-	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-	-	-	-
Origin not stated 3/, female													
2005	-	-	-	-	-	-	-	-	-	-	-	-	-
2004	-	-	-	-	-	-	-	-	-	-	-	-	-
2003	1	-	-	-	-	-	-	-	-	-	-	1	-
2002	-	-	-	-	-	-	-	-	-	-	-	-	-
2001	-	-	-	-	-	-	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-	-	-	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Meningitis (G00,G03)													
All origins, both sexes													
2005	669	57	32	26	28	32	74	118	106	74	76	45	1
2004	674	66	25	25	26	37	67	117	92	92	85	42	-
2003	730	77	24	17	37	44	67	117	108	96	90	52	1
2002	700	74	19	31	26	36	61	116	88	109	84	55	1
2001	727	78	26	24	40	41	69	105	113	94	96	41	-
2000	758	91	36	30	15	43	72	100	90	108	123	50	-
1999	850	116	45	27	41	48	76	108	90	115	122	62	-
All origins, male													
2005	353	29	21	15	15	24	44	68	53	37	32	14	1
2004	349	32	15	13	10	22	46	73	48	33	38	19	-
2003	387	38	12	10	20	28	40	73	61	45	40	19	1
2002	389	41	13	16	15	19	43	71	48	62	41	19	1
2001	394	39	16	15	23	25	42	60	61	51	47	15	-
2000	405	48	20	16	4	20	46	56	54	60	62	19	-
1999	442	61	26	18	26	24	45	68	45	60	49	20	-
All origins, female													
2005	316	28	11	11	13	8	30	50	53	37	44	31	-
2004	325	34	10	12	16	15	21	44	44	59	47	23	-
2003	343	39	12	7	17	16	27	44	47	51	50	33	-
2002	311	33	6	15	11	17	18	45	40	47	43	36	-
2001	333	39	10	9	17	16	27	45	52	43	49	26	-
2000	353	43	16	14	11	23	26	44	36	48	61	31	-
1999	408	55	19	9	15	24	31	40	45	55	73	42	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Meningitis (G00,G03)													
Hispanic, both sexes													
2005	63	10	8	2	4	2	8	10	5	6	5	3	-
2004	74	8	3	2	5	5	8	15	11	11	4	2	-
2003	82	18	4	3	7	11	7	11	11	5	2	3	-
2002	70	8	5	-	6	5	9	8	8	11	9	1	-
2001	58	8	4	3	3	4	7	6	7	6	8	2	-
2000	52	9	4	3	3	5	8	6	5	5	4	-	-
1999	76	17	7	1	5	6	4	12	7	9	4	4	-
Hispanic, male													
2005	33	5	2	-	3	2	7	8	1	2	3	-	-
2004	40	3	2	2	2	5	6	8	6	4	2	-	-
2003	46	12	2	3	5	6	4	7	4	2	-	1	-
2002	42	6	4	-	5	3	5	5	4	6	3	1	-
2001	33	5	3	2	1	1	6	3	3	5	4	-	-
2000	32	6	2	1	2	3	5	5	3	1	4	-	-
1999	41	13	5	1	1	3	3	6	3	5	-	1	-
Hispanic, female													
2005	30	5	6	2	1	-	1	2	4	4	2	3	-
2004	34	5	1	-	3	-	2	7	5	7	2	2	-
2003	36	6	2	-	2	5	3	4	7	3	2	2	-
2002	28	2	1	-	1	2	4	3	4	5	6	-	-
2001	25	3	1	1	2	3	1	3	4	1	4	2	-
2000	20	3	2	2	1	2	3	1	2	4	-	-	-
1999	35	4	2	-	4	3	1	6	4	4	4	3	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Meningitis (G00,G03)													
Non-Hispanic 1/, both sexes													
2005	602	47	24	24	24	30	66	105	100	68	71	42	1
2004	600	58	22	23	21	32	59	102	81	81	81	40	-
2003	646	59	20	14	30	33	60	105	97	91	88	49	-
2002	625	66	14	31	20	31	51	107	79	98	73	54	1
2001	666	70	22	21	37	37	62	98	106	88	88	37	-
2000	705	82	32	27	12	37	64	94	85	103	119	50	-
1999	767	97	38	25	34	42	71	96	82	106	118	58	-
Non-Hispanic 1/, male													
2005	317	24	19	15	12	22	37	58	51	35	29	14	1
2004	309	29	13	11	8	17	40	65	42	29	36	19	-
2003	339	26	10	7	15	22	36	65	57	43	40	18	-
2002	344	35	9	16	10	16	37	66	43	56	37	18	1
2001	361	34	13	13	22	24	36	57	58	46	43	15	-
2000	372	42	18	15	2	16	41	51	51	59	58	19	-
1999	396	48	21	16	23	21	41	62	41	55	49	19	-
Non-Hispanic 1/, female													
2005	285	23	5	9	12	8	29	47	49	33	42	28	-
2004	291	29	9	12	13	15	19	37	39	52	45	21	-
2003	307	33	10	7	15	11	24	40	40	48	48	31	-
2002	281	31	5	15	10	15	14	41	36	42	36	36	-
2001	305	36	9	8	15	13	26	41	48	42	45	22	-
2000	333	40	14	12	10	21	23	43	34	44	61	31	-
1999	371	49	17	9	11	21	30	34	41	51	69	39	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Meningitis (G00,G03)													
Non-Hispanic white, both sexes													
2005 2/	433	26	12	16	17	18	41	76	77	54	58	37	1
2004 2/	417	38	16	17	14	14	41	62	57	58	65	35	-
2003 2/	444	35	14	10	18	15	36	68	61	71	77	39	-
2002	427	44	10	20	12	15	30	62	49	77	58	49	1
2001	469	39	15	15	24	25	33	66	77	73	74	28	-
2000	486	47	19	18	7	20	36	65	60	74	100	40	-
1999	524	51	26	18	15	28	45	60	63	80	92	46	-
Non-Hispanic white, male													
2005 2/	228	13	11	9	10	12	23	45	39	28	25	12	1
2004 2/	220	22	10	9	5	8	28	43	26	21	30	18	-
2003 2/	247	21	7	6	11	11	22	43	39	35	37	15	-
2002	227	23	6	11	5	6	20	37	27	45	32	14	1
2001	243	17	10	8	12	17	18	38	42	37	35	9	-
2000	264	27	12	9	1	7	22	38	39	47	48	14	-
1999	265	33	15	11	10	10	24	35	33	41	38	15	-
Non-Hispanic white, female													
2005 2/	205	13	1	7	7	6	18	31	38	26	33	25	-
2004 2/	197	16	6	8	9	6	13	19	31	37	35	17	-
2003 2/	197	14	7	4	7	4	14	25	22	36	40	24	-
2002	200	21	4	9	7	9	10	25	22	32	26	35	-
2001	226	22	5	7	12	8	15	28	35	36	39	19	-
2000	222	20	7	9	6	13	14	27	21	27	52	26	-
1999	259	18	11	7	5	18	21	25	30	39	54	31	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Meningitis (G00,G03)													
Non-Hispanic black, both sexes													
2005 2/	141	17	6	7	5	11	22	28	19	13	8	5	-
2004 2/	158	17	5	6	5	15	18	36	21	19	13	3	-
2003 2/	183	22	5	3	12	18	22	31	35	18	8	9	-
2002	175	20	4	10	6	15	19	40	23	21	13	4	-
2001	172	29	5	4	10	11	26	30	25	13	11	8	-
2000	202	34	13	9	5	17	27	26	22	27	13	9	-
1999	217	40	11	7	17	11	21	35	18	21	24	12	-
Non-Hispanic black, male													
2005 2/	75	8	3	6	1	9	13	13	10	7	3	2	-
2004 2/	76	7	2	2	2	7	12	20	13	6	5	-	-
2003 2/	82	5	2	1	4	11	12	17	18	8	2	2	-
2002	105	11	3	5	5	9	15	26	12	11	5	3	-
2001	104	16	2	3	8	6	17	18	14	8	6	6	-
2000	100	15	6	6	1	9	18	11	11	11	7	5	-
1999	119	13	6	5	11	9	14	27	8	12	10	4	-
Non-Hispanic black, female													
2005 2/	66	9	3	1	4	2	9	15	9	6	5	3	-
2004 2/	82	10	3	4	3	8	6	16	8	13	8	3	-
2003 2/	101	17	3	2	8	7	10	14	17	10	6	7	-
2002	70	9	1	5	1	6	4	14	11	10	8	1	-
2001	68	13	3	1	2	5	9	12	11	5	5	2	-
2000	102	19	7	3	4	8	9	15	11	16	6	4	-
1999	98	27	5	2	6	2	7	8	10	9	14	8	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Meningitis (G00,G03)													
Origin not stated 3/, both sexes													
2005	4	-	-	-	-	-	-	3	1	-	-	-	-
2004	-	-	-	-	-	-	-	-	-	-	-	-	-
2003	2	-	-	-	-	-	-	1	-	-	-	-	1
2002	5	-	-	-	-	-	1	1	1	-	2	-	-
2001	3	-	-	-	-	-	-	1	-	-	-	2	-
2000	1	-	-	-	-	1	-	-	-	-	-	-	-
1999	7	2	-	1	2	-	1	-	1	-	-	-	-
Origin not stated 3/, male													
2005	3	-	-	-	-	-	-	2	1	-	-	-	-
2004	-	-	-	-	-	-	-	-	-	-	-	-	-
2003	2	-	-	-	-	-	-	1	-	-	-	-	1
2002	3	-	-	-	-	-	1	-	1	-	1	-	-
2001	-	-	-	-	-	-	-	-	-	-	-	-	-
2000	1	-	-	-	-	1	-	-	-	-	-	-	-
1999	5	-	-	1	2	-	1	-	1	-	-	-	-
Origin not stated 3/, female													
2005	1	-	-	-	-	-	-	1	-	-	-	-	-
2004	-	-	-	-	-	-	-	-	-	-	-	-	-
2003	-	-	-	-	-	-	-	-	-	-	-	-	-
2002	2	-	-	-	-	-	-	1	-	-	1	-	-
2001	3	-	-	-	-	-	-	1	-	-	-	2	-
2000	-	-	-	-	-	-	-	-	-	-	-	-	-
1999	2	2	-	-	-	-	-	-	-	-	-	-	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Parkinson's disease (G20-G21)													
All origins, both sexes													
2005	19,544	-	-	-	3	4	7	66	434	2,414	9,294	7,322	-
2004	17,989	-	1	-	1	2	8	64	349	2,214	8,752	6,598	-
2003	17,997	1	-	1	-	2	11	64	352	2,330	8,723	6,513	-
2002	16,959	-	-	-	-	-	6	58	318	2,226	8,139	6,212	-
2001	16,544	-	-	-	-	2	15	45	301	2,147	8,125	5,909	-
2000	15,682	-	1	-	-	4	7	49	270	2,108	7,652	5,590	1
1999	14,593	1	-	-	2	2	5	43	242	2,020	7,110	5,168	-
All origins, male													
2005	11,247	-	-	-	3	2	3	44	275	1,596	5,658	3,666	-
2004	10,366	-	-	-	1	2	6	49	227	1,457	5,306	3,318	-
2003	10,180	1	-	-	-	-	6	33	219	1,517	5,191	3,213	-
2002	9,593	-	-	-	-	-	3	40	193	1,487	4,811	3,059	-
2001	9,405	-	-	-	-	1	11	33	214	1,394	4,837	2,915	-
2000	8,804	-	-	-	-	3	5	29	170	1,338	4,530	2,728	1
1999	8,262	-	-	-	2	2	3	30	161	1,303	4,307	2,454	-
All origins, female													
2005	8,297	-	-	-	-	2	4	22	159	818	3,636	3,656	-
2004	7,623	-	1	-	-	-	2	15	122	757	3,446	3,280	-
2003	7,817	-	-	1	-	2	5	31	133	813	3,532	3,300	-
2002	7,366	-	-	-	-	-	3	18	125	739	3,328	3,153	-
2001	7,139	-	-	-	-	1	4	12	87	753	3,288	2,994	-
2000	6,878	-	1	-	-	1	2	20	100	770	3,122	2,862	-
1999	6,331	1	-	-	-	-	2	13	81	717	2,803	2,714	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Parkinson's disease (G20-G21)													
Hispanic, both sexes													
2005	726	-	-	-	2	-	1	7	21	119	354	222	-
2004	594	-	-	-	-	1	-	2	23	87	302	179	-
2003	521	-	-	-	-	-	1	1	15	96	258	150	-
2002	491	-	-	-	-	-	1	6	21	102	223	138	-
2001	483	-	-	-	-	1	-	3	13	98	230	138	-
2000	425	-	-	-	-	-	-	5	15	84	191	130	-
1999	389	-	-	-	-	-	1	2	8	84	184	110	-
Hispanic, male													
2005	418	-	-	-	2	-	1	2	9	80	213	111	-
2004	348	-	-	-	-	1	-	2	15	54	186	90	-
2003	313	-	-	-	-	-	1	-	9	67	152	84	-
2002	281	-	-	-	-	-	1	3	16	71	122	68	-
2001	274	-	-	-	-	1	-	2	7	56	141	67	-
2000	240	-	-	-	-	-	-	2	9	46	112	71	-
1999	222	-	-	-	-	-	1	2	5	52	113	49	-
Hispanic, female													
2005	308	-	-	-	-	-	-	5	12	39	141	111	-
2004	246	-	-	-	-	-	-	-	8	33	116	89	-
2003	208	-	-	-	-	-	-	1	6	29	106	66	-
2002	210	-	-	-	-	-	-	3	5	31	101	70	-
2001	209	-	-	-	-	-	-	1	6	42	89	71	-
2000	185	-	-	-	-	-	-	3	6	38	79	59	-
1999	167	-	-	-	-	-	-	-	3	32	71	61	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Parkinson's disease (G20-G21)													
Non-Hispanic 1/, both sexes													
2005	18,804	-	-	-	1	4	6	59	411	2,294	8,934	7,095	-
2004	17,368	-	1	-	1	1	8	62	324	2,126	8,434	6,411	-
2003	17,443	-	-	1	-	2	10	63	336	2,232	8,444	6,355	-
2002	16,435	-	-	-	-	-	5	52	296	2,120	7,899	6,063	-
2001	16,019	-	-	-	-	1	15	41	288	2,046	7,873	5,755	-
2000	15,223	-	1	-	-	4	7	43	255	2,020	7,442	5,450	1
1999	14,177	1	-	-	2	2	4	41	234	1,930	6,917	5,046	-
Non-Hispanic 1/, male													
2005	10,823	-	-	-	1	2	2	42	266	1,516	5,442	3,552	-
2004	10,000	-	-	-	1	1	6	47	210	1,402	5,109	3,224	-
2003	9,846	-	-	-	-	-	5	33	209	1,448	5,025	3,126	-
2002	9,290	-	-	-	-	-	2	37	176	1,413	4,677	2,985	-
2001	9,104	-	-	-	-	-	11	30	207	1,335	4,681	2,840	-
2000	8,544	-	-	-	-	3	5	26	161	1,290	4,407	2,651	1
1999	8,029	-	-	-	2	2	2	28	156	1,248	4,189	2,402	-
Non-Hispanic 1/, female													
2005	7,981	-	-	-	-	2	4	17	145	778	3,492	3,543	-
2004	7,368	-	1	-	-	-	2	15	114	724	3,325	3,187	-
2003	7,597	-	-	1	-	2	5	30	127	784	3,419	3,229	-
2002	7,145	-	-	-	-	-	3	15	120	707	3,222	3,078	-
2001	6,915	-	-	-	-	1	4	11	81	711	3,192	2,915	-
2000	6,679	-	1	-	-	1	2	17	94	730	3,035	2,799	-
1999	6,148	1	-	-	-	-	2	13	78	682	2,728	2,644	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Parkinson's disease (G20-G21)													
Non-Hispanic white, both sexes													
2005 2/	17,767	-	-	-	1	4	6	53	372	2,105	8,445	6,781	-
2004 2/	16,428	-	1	-	-	1	7	53	296	1,988	7,999	6,083	-
2003 2/	16,499	-	-	-	-	2	9	54	292	2,061	8,020	6,061	-
2002	15,626	-	-	-	-	-	2	48	269	1,959	7,545	5,803	-
2001	15,239	-	-	-	-	1	11	36	257	1,911	7,521	5,502	-
2000	14,497	-	-	-	-	3	4	36	236	1,895	7,106	5,216	1
1999	13,475	1	-	-	2	1	4	37	207	1,807	6,594	4,822	-
Non-Hispanic white, male													
2005 2/	10,232	-	-	-	1	2	2	37	241	1,392	5,164	3,393	-
2004 2/	9,469	-	-	-	-	1	5	40	190	1,313	4,864	3,056	-
2003 2/	9,303	-	-	-	-	-	5	29	180	1,343	4,784	2,962	-
2002	8,838	-	-	-	-	-	1	33	163	1,312	4,473	2,856	-
2001	8,685	-	-	-	-	-	8	27	190	1,252	4,486	2,722	-
2000	8,116	-	-	-	-	3	3	22	149	1,214	4,191	2,533	1
1999	7,648	-	-	-	2	1	2	27	142	1,182	4,005	2,287	-
Non-Hispanic white, female													
2005 2/	7,535	-	-	-	-	2	4	16	131	713	3,281	3,388	-
2004 2/	6,959	-	1	-	-	-	2	13	106	675	3,135	3,027	-
2003 2/	7,196	-	-	-	-	2	4	25	112	718	3,236	3,099	-
2002	6,788	-	-	-	-	-	1	15	106	647	3,072	2,947	-
2001	6,554	-	-	-	-	1	3	9	67	659	3,035	2,780	-
2000	6,381	-	-	-	-	-	1	14	87	681	2,915	2,683	-
1999	5,827	1	-	-	-	-	2	10	65	625	2,589	2,535	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Parkinson's disease (G20-G21)													
Non-Hispanic black, both sexes													
2005 2/	705	-	-	-	-	-	-	5	25	126	339	210	-
2004 2/	623	-	-	-	-	-	1	5	21	92	290	214	-
2003 2/	624	-	-	1	-	-	1	7	33	120	282	180	-
2002	547	-	-	-	-	-	1	3	18	111	243	171	-
2001	547	-	-	-	-	-	4	5	20	91	238	189	-
2000	528	-	-	-	-	1	3	6	14	93	246	165	-
1999	502	-	-	-	-	-	-	4	22	95	221	160	-
Non-Hispanic black, male													
2005 2/	409	-	-	-	-	-	-	4	15	90	196	104	-
2004 2/	342	-	-	-	-	-	1	5	15	60	156	105	-
2003 2/	348	-	-	-	-	-	-	3	21	73	158	93	-
2002	306	-	-	-	-	-	-	3	9	76	138	80	-
2001	291	-	-	-	-	-	3	3	13	59	127	86	-
2000	304	-	-	-	-	-	2	4	9	60	154	75	-
1999	269	-	-	-	-	-	-	1	13	58	123	74	-
Non-Hispanic black, female													
2005 2/	296	-	-	-	-	-	-	1	10	36	143	106	-
2004 2/	281	-	-	-	-	-	-	-	6	32	134	109	-
2003 2/	276	-	-	1	-	-	1	4	12	47	124	87	-
2002	241	-	-	-	-	-	1	-	9	35	105	91	-
2001	256	-	-	-	-	-	1	2	7	32	111	103	-
2000	224	-	-	-	-	1	1	2	5	33	92	90	-
1999	233	-	-	-	-	-	-	3	9	37	98	86	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Parkinson's disease (G20-G21)													
Origin not stated 3/, both sexes													
2005	14	-	-	-	-	-	-	-	2	1	6	5	-
2004	27	-	-	-	-	-	-	-	2	1	16	8	-
2003	33	1	-	-	-	-	-	-	1	2	21	8	-
2002	33	-	-	-	-	-	-	-	1	4	17	11	-
2001	42	-	-	-	-	-	-	1	-	3	22	16	-
2000	34	-	-	-	-	-	-	1	-	4	19	10	-
1999	27	-	-	-	-	-	-	-	-	6	9	12	-
Origin not stated 3/, male													
2005	6	-	-	-	-	-	-	-	-	-	3	3	-
2004	18	-	-	-	-	-	-	-	2	1	11	4	-
2003	21	1	-	-	-	-	-	-	1	2	14	3	-
2002	22	-	-	-	-	-	-	-	1	3	12	6	-
2001	27	-	-	-	-	-	-	1	-	3	15	8	-
2000	20	-	-	-	-	-	-	1	-	2	11	6	-
1999	11	-	-	-	-	-	-	-	-	3	5	3	-
Origin not stated 3/, female													
2005	8	-	-	-	-	-	-	-	2	1	3	2	-
2004	9	-	-	-	-	-	-	-	-	-	5	4	-
2003	12	-	-	-	-	-	-	-	-	-	7	5	-
2002	11	-	-	-	-	-	-	-	-	1	5	5	-
2001	15	-	-	-	-	-	-	-	-	-	7	8	-
2000	14	-	-	-	-	-	-	-	-	2	8	4	-
1999	16	-	-	-	-	-	-	-	-	3	4	9	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Alzheimer's disease (G30)													
All origins, both sexes													
2005	71,599	-	-	-	-	2	10	80	648	3,813	23,139	43,906	1
2004	65,965	-	-	-	1	1	7	101	542	3,641	21,882	39,790	-
2003	63,457	-	-	-	-	-	11	78	554	3,836	21,157	37,821	-
2002	58,866	-	-	1	-	2	12	52	510	3,602	20,135	34,552	-
2001	53,852	-	-	-	-	2	3	73	529	3,422	18,542	31,281	-
2000	49,558	-	-	-	-	2	6	64	491	3,431	17,253	28,309	2
1999	44,536	-	-	-	1	-	7	60	448	3,204	15,836	24,980	-
All origins, male													
2005	20,559	-	-	-	-	-	6	30	287	1,648	8,326	10,262	-
2004	18,974	-	-	-	-	1	4	38	253	1,558	7,777	9,343	-
2003	18,335	-	-	-	-	-	4	34	243	1,684	7,606	8,764	-
2002	16,989	-	-	-	-	2	8	27	228	1,547	7,227	7,950	-
2001	15,762	-	-	-	-	2	3	38	228	1,472	6,795	7,224	-
2000	14,438	-	-	-	-	1	4	34	210	1,554	6,228	6,406	1
1999	13,391	-	-	-	1	-	3	28	222	1,437	5,904	5,796	-
All origins, female													
2005	51,040	-	-	-	-	2	4	50	361	2,165	14,813	33,644	1
2004	46,991	-	-	-	1	-	3	63	289	2,083	14,105	30,447	-
2003	45,122	-	-	-	-	-	7	44	311	2,152	13,551	29,057	-
2002	41,877	-	-	1	-	-	4	25	282	2,055	12,908	26,602	-
2001	38,090	-	-	-	-	-	-	35	301	1,950	11,747	24,057	-
2000	35,120	-	-	-	-	1	2	30	281	1,877	11,025	21,903	1
1999	31,145	-	-	-	-	-	4	32	226	1,767	9,932	19,184	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Alzheimer's disease (G30)													
Hispanic, both sexes													
2005	2,201	-	-	-	-	-	2	7	32	172	801	1,187	-
2004	1,897	-	-	-	1	-	-	8	22	153	681	1,032	-
2003	1,835	-	-	-	-	-	1	12	27	179	664	952	-
2002	1,474	-	-	-	-	-	1	-	20	127	535	791	-
2001	1,239	-	-	-	-	-	1	3	19	112	465	639	-
2000	1,127	-	-	-	-	-	-	6	19	128	401	573	-
1999	981	-	-	-	1	-	-	3	14	100	362	501	-
Hispanic, male													
2005	764	-	-	-	-	-	1	1	12	82	337	331	-
2004	606	-	-	-	-	-	-	4	10	65	260	267	-
2003	633	-	-	-	-	-	-	6	11	75	264	277	-
2002	464	-	-	-	-	-	1	-	8	55	196	204	-
2001	401	-	-	-	-	-	1	3	6	44	186	161	-
2000	383	-	-	-	-	-	-	4	9	63	166	141	-
1999	314	-	-	-	1	-	-	-	5	51	128	129	-
Hispanic, female													
2005	1,437	-	-	-	-	-	1	6	20	90	464	856	-
2004	1,291	-	-	-	1	-	-	4	12	88	421	765	-
2003	1,202	-	-	-	-	-	1	6	16	104	400	675	-
2002	1,010	-	-	-	-	-	-	-	12	72	339	587	-
2001	838	-	-	-	-	-	-	-	13	68	279	478	-
2000	744	-	-	-	-	-	-	2	10	65	235	432	-
1999	667	-	-	-	-	-	-	3	9	49	234	372	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Alzheimer's disease (G30)													
Non-Hispanic 1/, both sexes													
2005	69,315	-	-	-	-	2	8	73	613	3,637	22,314	42,667	1
2004	63,981	-	-	-	-	1	7	93	519	3,477	21,173	38,711	-
2003	61,506	-	-	-	-	-	10	66	525	3,647	20,459	36,799	-
2002	57,246	-	-	1	-	2	11	52	488	3,463	19,552	33,677	-
2001	52,516	-	-	-	-	2	2	70	509	3,305	18,051	30,577	-
2000	48,331	-	-	-	-	2	6	58	469	3,296	16,829	27,670	1
1999	43,471	-	-	-	-	-	7	57	434	3,099	15,447	24,427	-
Non-Hispanic 1/, male													
2005	19,764	-	-	-	-	-	5	29	274	1,562	7,977	9,917	-
2004	18,345	-	-	-	-	1	4	34	242	1,490	7,507	9,067	-
2003	17,666	-	-	-	-	-	4	28	231	1,603	7,332	8,468	-
2002	16,491	-	-	-	-	2	7	27	220	1,487	7,015	7,733	-
2001	15,334	-	-	-	-	2	2	35	222	1,427	6,600	7,046	-
2000	14,034	-	-	-	-	1	4	30	200	1,489	6,058	6,252	-
1999	13,052	-	-	-	-	-	3	28	217	1,384	5,768	5,652	-
Non-Hispanic 1/, female													
2005	49,551	-	-	-	-	2	3	44	339	2,075	14,337	32,750	1
2004	45,636	-	-	-	-	-	3	59	277	1,987	13,666	29,644	-
2003	43,840	-	-	-	-	-	6	38	294	2,044	13,127	28,331	-
2002	40,755	-	-	1	-	-	4	25	268	1,976	12,537	25,944	-
2001	37,182	-	-	-	-	-	-	35	287	1,878	11,451	23,531	-
2000	34,297	-	-	-	-	1	2	28	269	1,807	10,771	21,418	1
1999	30,419	-	-	-	-	-	4	29	217	1,715	9,679	18,775	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Alzheimer's disease (G30)													
Non-Hispanic white, both sexes													
2005 2/	63,958	-	-	-	-	2	8	66	559	3,208	20,469	39,646	-
2004 2/	59,153	-	-	-	-	1	6	89	472	3,107	19,438	36,040	-
2003 2/	57,268	-	-	-	-	-	10	62	489	3,269	18,937	34,501	-
2002	53,486	-	-	1	-	1	6	49	448	3,149	18,227	31,605	-
2001	49,030	-	-	-	-	2	2	64	480	3,020	16,742	28,720	-
2000	45,246	-	-	-	-	2	6	50	428	3,004	15,704	26,052	-
1999	40,835	-	-	-	-	-	5	51	399	2,821	14,453	23,106	-
Non-Hispanic white, male													
2005 2/	18,208	-	-	-	-	-	5	25	253	1,380	7,326	9,219	-
2004 2/	16,989	-	-	-	-	1	3	33	221	1,335	6,945	8,451	-
2003 2/	16,427	-	-	-	-	-	4	26	212	1,443	6,796	7,946	-
2002	15,399	-	-	-	-	1	4	24	203	1,353	6,544	7,270	-
2001	14,295	-	-	-	-	2	2	31	205	1,305	6,137	6,613	-
2000	13,123	-	-	-	-	1	4	25	183	1,367	5,686	5,857	-
1999	12,253	-	-	-	-	-	2	26	200	1,263	5,432	5,330	-
Non-Hispanic white, female													
2005 2/	45,750	-	-	-	-	2	3	41	306	1,828	13,143	30,427	-
2004 2/	42,164	-	-	-	-	-	3	56	251	1,772	12,493	27,589	-
2003 2/	40,841	-	-	-	-	-	6	36	277	1,826	12,141	26,555	-
2002	38,087	-	-	1	-	-	2	25	245	1,796	11,683	24,335	-
2001	34,735	-	-	-	-	-	-	33	275	1,715	10,605	22,107	-
2000	32,123	-	-	-	-	1	2	25	245	1,637	10,018	20,195	-
1999	28,582	-	-	-	-	-	3	25	199	1,558	9,021	17,776	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Alzheimer's disease (G30)													
Non-Hispanic black, both sexes													
2005 2/	4,579	-	-	-	-	-	-	5	45	390	1,579	2,559	1
2004 2/	4,196	-	-	-	-	-	1	4	37	327	1,510	2,317	-
2003 2/	3,693	-	-	-	-	-	-	4	29	326	1,347	1,987	-
2002	3,303	-	-	-	-	1	4	3	31	277	1,173	1,814	-
2001	3,104	-	-	-	-	-	-	5	26	253	1,176	1,644	-
2000	2,719	-	-	-	-	-	-	5	30	261	1,004	1,418	1
1999	2,325	-	-	-	-	-	1	4	29	249	883	1,159	-
Non-Hispanic black, male													
2005 2/	1,283	-	-	-	-	-	-	4	17	163	545	554	-
2004 2/	1,138	-	-	-	-	-	1	1	15	136	486	499	-
2003 2/	1,079	-	-	-	-	-	-	2	17	142	481	437	-
2002	940	-	-	-	-	1	3	3	14	122	418	379	-
2001	906	-	-	-	-	-	-	3	17	110	412	364	-
2000	774	-	-	-	-	-	-	3	16	113	320	322	-
1999	689	-	-	-	-	-	-	2	15	112	290	270	-
Non-Hispanic black, female													
2005 2/	3,296	-	-	-	-	-	-	1	28	227	1,034	2,005	1
2004 2/	3,058	-	-	-	-	-	-	3	22	191	1,024	1,818	-
2003 2/	2,614	-	-	-	-	-	-	2	12	184	866	1,550	-
2002	2,363	-	-	-	-	-	1	-	17	155	755	1,435	-
2001	2,198	-	-	-	-	-	-	2	9	143	764	1,280	-
2000	1,945	-	-	-	-	-	-	2	14	148	684	1,096	1
1999	1,636	-	-	-	-	-	1	2	14	137	593	889	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Alzheimer's disease (G30)													
Origin not stated 3/, both sexes													
2005	83	-	-	-	-	-	-	-	3	4	24	52	-
2004	87	-	-	-	-	-	-	-	1	11	28	47	-
2003	116	-	-	-	-	-	-	-	2	10	34	70	-
2002	146	-	-	-	-	-	-	-	2	12	48	84	-
2001	97	-	-	-	-	-	-	-	1	5	26	65	-
2000	100	-	-	-	-	-	-	-	3	7	23	66	1
1999	84	-	-	-	-	-	-	-	-	5	27	52	-
Origin not stated 3/, male													
2005	31	-	-	-	-	-	-	-	1	4	12	14	-
2004	23	-	-	-	-	-	-	-	1	3	10	9	-
2003	36	-	-	-	-	-	-	-	1	6	10	19	-
2002	34	-	-	-	-	-	-	-	-	5	16	13	-
2001	27	-	-	-	-	-	-	-	-	1	9	17	-
2000	21	-	-	-	-	-	-	-	1	2	4	13	1
1999	25	-	-	-	-	-	-	-	-	2	8	15	-
Origin not stated 3/, female													
2005	52	-	-	-	-	-	-	-	2	-	12	38	-
2004	64	-	-	-	-	-	-	-	-	8	18	38	-
2003	80	-	-	-	-	-	-	-	1	4	24	51	-
2002	112	-	-	-	-	-	-	-	2	7	32	71	-
2001	70	-	-	-	-	-	-	-	1	4	17	48	-
2000	79	-	-	-	-	-	-	-	2	5	19	53	-
1999	59	-	-	-	-	-	-	-	-	3	19	37	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Major cardiovascular diseases (I00-I78)													
All origins, both sexes													
2005	856,030	503	217	367	1,391	4,041	15,852	46,928	79,896	124,366	256,362	326,066	41
2004	861,190	564	234	336	1,333	4,006	16,138	46,153	77,984	128,816	264,037	321,544	45
2003	902,443	555	234	347	1,447	4,089	16,930	46,279	79,497	137,220	279,358	336,449	38
2002	918,628	640	223	358	1,283	3,986	16,962	45,800	78,430	144,071	287,415	339,409	51
2001	922,334	600	285	365	1,280	4,038	16,686	44,450	76,224	148,409	291,995	337,930	72
2000	936,923	636	234	362	1,309	3,792	16,624	43,583	77,482	156,187	296,202	340,450	62
1999	950,314	644	234	354	1,315	3,873	17,010	42,512	77,833	163,428	302,552	340,510	49
All origins, male													
2005	405,780	282	126	200	914	2,657	10,664	32,548	53,484	73,733	125,037	106,106	29
2004	406,458	307	120	170	867	2,602	10,806	32,057	52,014	76,179	128,490	102,808	38
2003	423,810	310	116	204	893	2,609	11,302	32,260	52,804	81,512	135,868	105,903	29
2002	429,682	343	122	194	845	2,568	11,356	31,928	52,037	85,319	138,770	106,164	36
2001	428,065	315	149	184	807	2,555	11,180	30,860	50,320	87,756	140,789	103,097	53
2000	435,912	341	128	206	783	2,378	11,110	30,535	51,423	92,510	142,797	103,658	43
1999	441,827	347	127	199	799	2,442	11,471	29,745	51,525	96,904	145,567	102,663	38
All origins, female													
2005	450,250	221	91	167	477	1,384	5,188	14,380	26,412	50,633	131,325	219,960	12
2004	454,732	257	114	166	466	1,404	5,332	14,096	25,970	52,637	135,547	218,736	7
2003	478,633	245	118	143	554	1,480	5,628	14,019	26,693	55,708	143,490	230,546	9
2002	488,946	297	101	164	438	1,418	5,606	13,872	26,393	58,752	148,645	233,245	15
2001	494,269	285	136	181	473	1,483	5,506	13,590	25,904	60,653	151,206	234,833	19
2000	501,011	295	106	156	526	1,414	5,514	13,048	26,059	63,677	153,405	236,792	19
1999	508,487	297	107	155	516	1,431	5,539	12,767	26,308	66,524	156,985	237,847	11

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Major cardiovascular diseases (I00-I78)													
Hispanic, both sexes													
2005	39,010	98	40	59	198	465	1,339	3,037	4,706	7,205	11,411	10,447	5
2004	36,900	104	43	46	179	424	1,304	2,874	4,494	7,058	10,733	9,636	5
2003	37,266	95	43	46	176	470	1,211	2,747	4,431	7,269	10,783	9,989	6
2002	36,543	109	36	51	176	445	1,184	2,690	4,460	7,342	10,362	9,687	1
2001	35,565	98	44	56	162	442	1,081	2,561	3,996	7,325	10,139	9,654	7
2000	34,059	106	37	47	169	419	1,094	2,351	4,041	7,114	9,380	9,296	5
1999	33,665	115	41	47	160	408	1,057	2,386	3,973	7,150	9,291	9,031	6
Hispanic, male													
2005	20,368	54	27	34	127	340	970	2,167	3,106	4,206	5,673	3,659	5
2004	19,348	54	28	21	132	323	948	2,044	2,986	4,132	5,391	3,284	5
2003	19,066	54	26	27	123	315	864	1,964	2,928	4,224	5,236	3,300	5
2002	18,824	52	16	21	122	322	829	1,876	2,958	4,304	5,039	3,284	1
2001	18,138	53	21	29	108	308	758	1,806	2,582	4,269	4,966	3,232	6
2000	17,442	54	21	30	110	283	739	1,671	2,657	4,123	4,622	3,128	4
1999	17,304	49	21	29	103	292	755	1,700	2,560	4,212	4,519	3,058	6
Hispanic, female													
2005	18,642	44	13	25	71	125	369	870	1,600	2,999	5,738	6,788	-
2004	17,552	50	15	25	47	101	356	830	1,508	2,926	5,342	6,352	-
2003	18,200	41	17	19	53	155	347	783	1,503	3,045	5,547	6,689	1
2002	17,719	57	20	30	54	123	355	814	1,502	3,038	5,323	6,403	-
2001	17,427	45	23	27	54	134	323	755	1,414	3,056	5,173	6,422	1
2000	16,617	52	16	17	59	136	355	680	1,384	2,991	4,758	6,168	1
1999	16,361	66	20	18	57	116	302	686	1,413	2,938	4,772	5,973	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Major cardiovascular diseases (I00-I78)													
Non-Hispanic 1/, both sexes													
2005	815,443	405	177	306	1,192	3,570	14,479	43,746	74,953	116,875	244,492	315,216	32
2004	822,392	457	191	288	1,149	3,575	14,778	43,114	73,224	121,373	252,787	311,423	33
2003	862,744	456	190	300	1,265	3,602	15,647	43,350	74,723	129,500	267,904	325,786	21
2002	879,170	528	187	307	1,105	3,522	15,707	42,913	73,650	136,175	276,232	328,806	38
2001	883,949	499	239	308	1,108	3,578	15,536	41,695	71,875	140,618	281,026	327,420	47
2000	899,801	526	196	313	1,137	3,357	15,433	41,021	73,064	148,517	285,960	330,234	43
1999	913,476	528	192	306	1,152	3,453	15,869	39,935	73,473	155,643	292,354	330,537	34
Non-Hispanic 1/, male													
2005	384,533	228	99	165	787	2,313	9,669	30,277	50,202	69,329	119,110	102,332	22
2004	386,042	252	92	147	732	2,273	9,822	29,889	48,830	71,787	122,820	99,372	26
2003	403,411	255	90	176	766	2,283	10,387	30,161	49,614	77,001	130,276	102,387	15
2002	409,343	289	106	173	721	2,234	10,477	29,901	48,851	80,666	133,303	102,597	25
2001	408,437	260	128	155	692	2,231	10,366	28,916	47,469	83,178	135,395	99,611	36
2000	416,837	284	107	176	672	2,088	10,294	28,702	48,495	88,005	137,754	100,232	28
1999	422,824	298	106	169	694	2,142	10,649	27,907	48,675	92,273	140,588	99,298	25
Non-Hispanic 1/, female													
2005	430,910	177	78	141	405	1,257	4,810	13,469	24,751	47,546	125,382	212,884	10
2004	436,350	205	99	141	417	1,302	4,956	13,225	24,394	49,586	129,967	212,051	7
2003	459,333	201	100	124	499	1,319	5,260	13,189	25,109	52,499	137,628	223,399	6
2002	469,827	239	81	134	384	1,288	5,230	13,012	24,799	55,509	142,929	226,209	13
2001	475,512	239	111	153	416	1,347	5,170	12,779	24,406	57,440	145,631	227,809	11
2000	482,964	242	89	137	465	1,269	5,139	12,319	24,569	60,512	148,206	230,002	15
1999	490,652	230	86	137	458	1,311	5,220	12,028	24,798	63,370	151,766	231,239	9

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Major cardiovascular diseases (I00-I78)													
Non-Hispanic white, both sexes													
2005 2/	698,139	227	101	183	733	2,124	9,763	30,953	57,249	94,862	214,270	287,655	19
2004 2/	705,435	277	109	182	721	2,157	10,113	30,593	56,141	98,940	222,269	283,914	19
2003 2/	741,886	250	107	191	780	2,207	10,600	30,708	57,448	105,736	236,254	297,588	17
2002	758,738	314	125	195	695	2,121	10,838	30,798	56,865	112,288	244,074	300,399	26
2001	764,021	289	140	204	671	2,240	10,524	29,967	55,480	116,098	248,922	299,458	28
2000	780,445	330	126	195	690	2,084	10,556	29,780	56,666	123,708	254,089	302,192	29
1999	793,945	324	121	203	730	2,184	10,807	29,134	57,094	130,111	260,395	302,824	18
Non-Hispanic white, male													
2005 2/	328,004	127	59	104	487	1,418	6,796	22,291	39,316	57,345	105,876	94,172	13
2004 2/	330,160	154	54	96	463	1,421	7,039	22,122	38,358	59,798	109,504	91,136	15
2003 2/	345,832	133	43	115	476	1,446	7,382	22,301	39,261	64,254	116,254	94,156	11
2002	352,520	177	77	111	462	1,387	7,559	22,435	38,736	68,016	119,331	94,211	18
2001	352,013	160	75	108	425	1,441	7,379	21,723	37,726	70,213	121,332	91,409	22
2000	360,757	180	75	118	418	1,361	7,422	21,785	38,681	75,026	123,834	91,837	20
1999	366,310	189	70	110	430	1,400	7,606	21,200	38,851	78,853	126,770	90,818	13
Non-Hispanic white, female													
2005 2/	370,135	100	42	79	246	706	2,967	8,662	17,933	37,517	108,394	193,483	6
2004 2/	375,275	123	55	86	258	736	3,074	8,471	17,783	39,142	112,765	192,778	4
2003 2/	396,054	117	64	76	304	761	3,218	8,407	18,187	41,482	120,000	203,432	6
2002	406,218	137	48	84	233	734	3,279	8,363	18,129	44,272	124,743	206,188	8
2001	412,008	129	65	96	246	799	3,145	8,244	17,754	45,885	127,590	208,049	6
2000	419,688	150	51	77	272	723	3,134	7,995	17,985	48,682	130,255	210,355	9
1999	427,635	135	51	93	300	784	3,201	7,934	18,243	51,258	133,625	212,006	5

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Major cardiovascular diseases (I00-I78)													
Non-Hispanic black, both sexes													
2005 2/	99,000	148	60	101	403	1,269	4,177	11,435	15,507	18,626	24,967	22,295	12
2004 2/	99,408	142	74	98	372	1,246	4,216	11,254	15,073	19,232	25,325	22,363	13
2003 2/	103,027	168	70	95	428	1,231	4,489	11,349	15,174	20,382	26,363	23,276	2
2002	103,054	183	50	93	356	1,252	4,363	10,923	14,834	20,461	26,932	23,596	11
2001	103,213	184	86	85	395	1,209	4,506	10,539	14,483	21,084	27,025	23,600	17
2000	103,239	166	60	95	396	1,160	4,396	10,109	14,502	21,561	26,954	23,826	14
1999	103,629	172	60	93	388	1,131	4,610	9,721	14,487	22,242	27,129	23,581	15
Non-Hispanic black, male													
2005 2/	47,004	86	30	51	265	770	2,500	7,046	9,471	10,022	10,682	6,073	8
2004 2/	46,787	80	37	47	242	725	2,467	6,873	9,157	10,200	10,768	6,181	10
2003 2/	48,237	102	41	53	262	721	2,628	6,959	9,006	10,809	11,435	6,219	2
2002	47,656	96	24	50	217	739	2,551	6,635	8,862	10,770	11,323	6,383	6
2001	47,646	93	45	39	240	706	2,625	6,386	8,489	11,055	11,536	6,419	13
2000	47,408	88	27	47	218	652	2,544	6,143	8,568	11,074	11,434	6,605	8
1999	47,982	91	32	53	240	659	2,716	5,984	8,609	11,567	11,329	6,691	11
Non-Hispanic black, female													
2005 2/	51,996	62	30	50	138	499	1,677	4,389	6,036	8,604	14,285	16,222	4
2004 2/	52,621	62	37	51	130	521	1,749	4,381	5,916	9,032	14,557	16,182	3
2003 2/	54,790	66	29	42	166	510	1,861	4,390	6,168	9,573	14,928	17,057	-
2002	55,398	87	26	43	139	513	1,812	4,288	5,972	9,691	15,609	17,213	5
2001	55,567	91	41	46	155	503	1,881	4,153	5,994	10,029	15,489	17,181	4
2000	55,831	78	33	48	178	508	1,852	3,966	5,934	10,487	15,520	17,221	6
1999	55,647	81	28	40	148	472	1,894	3,737	5,878	10,675	15,800	16,890	4

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Major cardiovascular diseases (I00-I78)													
Origin not stated 3/, both sexes													
2005	1,577	-	-	2	1	6	34	145	237	286	459	403	4
2004	1,898	3	-	2	5	7	56	165	266	385	517	485	7
2003	2,433	4	1	1	6	17	72	182	343	451	671	674	11
2002	2,915	3	-	-	2	19	71	197	320	554	821	916	12
2001	2,820	3	2	1	10	18	69	194	353	466	830	856	18
2000	3,063	4	1	2	3	16	97	211	377	556	862	920	14
1999	3,173	1	1	1	3	12	84	191	387	635	907	942	9
Origin not stated 3/, male													
2005	879	-	-	1	-	4	25	104	176	198	254	115	2
2004	1,068	1	-	2	3	6	36	124	198	260	279	152	7
2003	1,333	1	-	1	4	11	51	135	262	287	356	216	9
2002	1,515	2	-	-	2	12	50	151	228	349	428	283	10
2001	1,490	2	-	-	7	16	56	138	269	309	428	254	11
2000	1,633	3	-	-	1	7	77	162	271	382	421	298	11
1999	1,699	-	-	1	2	8	67	138	290	419	460	307	7
Origin not stated 3/, female													
2005	698	-	-	1	1	2	9	41	61	88	205	288	2
2004	830	2	-	-	2	1	20	41	68	125	238	333	-
2003	1,100	3	1	-	2	6	21	47	81	164	315	458	2
2002	1,400	1	-	-	-	7	21	46	92	205	393	633	2
2001	1,330	1	2	1	3	2	13	56	84	157	402	602	7
2000	1,430	1	1	2	2	9	20	49	106	174	441	622	3
1999	1,474	1	1	-	1	4	17	53	97	216	447	635	2

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Diseases of heart (I00-I09,I11,I13,I20-I51)													
All origins, both sexes													
2005	652,091	358	151	252	1,119	3,249	12,688	38,103	65,208	96,729	190,693	243,504	37
2004	652,486	421	187	245	1,038	3,163	12,925	37,556	63,613	99,999	195,379	237,924	36
2003	685,089	439	186	264	1,133	3,250	13,600	37,732	65,060	107,263	207,331	248,796	35
2002	696,947	500	165	255	1,022	3,165	13,688	37,570	64,234	112,547	213,581	250,173	47
2001	700,142	479	225	272	999	3,160	13,326	36,399	62,486	116,299	216,992	249,439	66
2000	710,760	493	181	271	1,031	2,958	13,181	35,480	63,399	122,405	220,060	251,242	59
1999	725,192	524	183	277	1,069	3,066	13,600	34,994	64,167	129,253	226,152	251,860	47
All origins, male													
2005	322,841	197	81	131	770	2,206	8,931	27,486	44,895	59,050	96,580	82,489	25
2004	321,973	228	94	122	691	2,143	9,011	27,050	43,788	61,000	98,812	79,002	32
2003	336,095	247	89	155	718	2,129	9,463	27,304	44,522	65,568	104,646	81,227	27
2002	340,933	267	87	141	686	2,131	9,638	27,239	43,900	68,658	107,217	80,935	34
2001	339,095	243	121	138	647	2,068	9,346	26,280	42,519	70,637	108,576	78,473	47
2000	344,807	260	107	163	638	1,922	9,286	25,927	43,285	74,589	109,693	78,896	41
1999	351,617	270	104	157	664	1,988	9,640	25,489	43,662	78,836	112,470	78,300	37
All origins, female													
2005	329,250	161	70	121	349	1,043	3,757	10,617	20,313	37,679	94,113	161,015	12
2004	330,513	193	93	123	347	1,020	3,914	10,506	19,825	38,999	96,567	158,922	4
2003	348,994	192	97	109	415	1,121	4,137	10,428	20,538	41,695	102,685	167,569	8
2002	356,014	233	78	114	336	1,034	4,050	10,331	20,334	43,889	106,364	169,238	13
2001	361,047	236	104	134	352	1,092	3,980	10,119	19,967	45,662	108,416	170,966	19
2000	365,953	233	74	108	393	1,036	3,895	9,553	20,114	47,816	110,367	172,346	18
1999	373,575	254	79	120	405	1,078	3,960	9,505	20,505	50,417	113,682	173,560	10

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Diseases of heart (I00-I09,I11,I13,I20-I51)													
Hispanic, both sexes													
2005	29,555	71	23	34	137	354	957	2,233	3,642	5,551	8,583	7,965	5
2004	27,788	81	39	34	129	302	934	2,110	3,460	5,389	8,061	7,246	3
2003	28,298	75	35	31	135	347	879	2,035	3,454	5,635	8,077	7,589	6
2002	27,887	85	24	30	128	321	852	2,020	3,492	5,719	7,779	7,436	1
2001	27,090	79	33	38	112	314	750	1,875	3,152	5,730	7,632	7,369	6
2000	25,819	86	29	38	125	280	760	1,754	3,136	5,453	7,067	7,086	5
1999	25,866	96	33	36	116	286	761	1,806	3,155	5,557	7,046	6,968	6
Hispanic, male													
2005	15,900	38	17	21	91	272	733	1,679	2,492	3,316	4,398	2,838	5
2004	14,983	41	24	15	101	244	715	1,569	2,376	3,236	4,163	2,496	3
2003	14,867	42	19	19	101	244	654	1,520	2,334	3,387	4,025	2,517	5
2002	14,798	41	11	14	94	242	625	1,467	2,393	3,441	3,898	2,571	1
2001	14,195	43	16	19	76	226	565	1,397	2,104	3,424	3,808	2,512	5
2000	13,566	44	21	26	87	199	550	1,307	2,128	3,245	3,525	2,430	4
1999	13,554	40	20	23	73	219	572	1,348	2,091	3,355	3,483	2,324	6
Hispanic, female													
2005	13,655	33	6	13	46	82	224	554	1,150	2,235	4,185	5,127	-
2004	12,805	40	15	19	28	58	219	541	1,084	2,153	3,898	4,750	-
2003	13,431	33	16	12	34	103	225	515	1,120	2,248	4,052	5,072	1
2002	13,089	44	13	16	34	79	227	553	1,099	2,278	3,881	4,865	-
2001	12,895	36	17	19	36	88	185	478	1,048	2,306	3,824	4,857	1
2000	12,253	42	8	12	38	81	210	447	1,008	2,208	3,542	4,656	1
1999	12,312	56	13	13	43	67	189	458	1,064	2,202	3,563	4,644	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Diseases of heart (I00-I09,I11,I13,I20-I51)													
Non-Hispanic 1/, both sexes													
2005	621,260	287	128	217	982	2,889	11,702	35,740	61,359	90,945	181,756	235,227	28
2004	623,165	338	148	209	904	2,855	11,942	35,311	59,930	94,288	186,908	230,305	27
2003	654,828	362	150	232	993	2,891	12,662	35,545	61,305	101,261	198,713	240,694	20
2002	666,690	413	141	225	892	2,827	12,780	35,376	60,465	106,351	205,156	242,030	34
2001	670,787	397	191	234	877	2,834	12,515	34,359	59,038	110,181	208,704	241,415	42
2000	682,470	403	151	232	904	2,664	12,341	33,551	59,943	116,476	212,328	243,436	41
1999	696,719	427	149	240	950	2,770	12,767	33,029	60,684	123,161	218,359	244,151	32
Non-Hispanic 1/, male													
2005	306,197	159	64	110	679	1,930	8,175	25,711	42,247	55,566	91,981	79,557	18
2004	306,083	186	70	105	587	1,893	8,263	25,378	41,242	57,539	94,418	76,379	23
2003	320,095	205	70	135	614	1,878	8,765	25,665	41,957	61,943	100,321	78,528	14
2002	324,848	225	76	127	590	1,878	8,972	25,636	41,305	64,912	102,972	78,132	23
2001	323,673	198	105	119	564	1,832	8,733	24,759	40,188	66,949	104,429	75,766	31
2000	329,888	213	86	137	550	1,716	8,673	24,487	40,924	71,010	105,834	76,231	27
1999	336,615	230	84	133	589	1,762	9,009	24,019	41,321	75,118	108,605	75,721	24
Non-Hispanic 1/, female													
2005	315,063	128	64	107	303	959	3,527	10,029	19,112	35,379	89,775	155,670	10
2004	317,082	152	78	104	317	962	3,679	9,933	18,688	36,749	92,490	153,926	4
2003	334,733	157	80	97	379	1,013	3,897	9,880	19,348	39,318	98,392	162,166	6
2002	341,842	188	65	98	302	949	3,808	9,740	19,160	41,439	102,184	163,898	11
2001	347,114	199	86	115	313	1,002	3,782	9,600	18,850	43,232	104,275	165,649	11
2000	352,582	190	65	95	354	948	3,668	9,064	19,019	45,466	106,494	167,205	14
1999	360,104	197	65	107	361	1,008	3,758	9,010	19,363	48,043	109,754	168,430	8

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Diseases of heart (I00-I09,I11,I13,I20-I51)													
Non-Hispanic white, both sexes													
2005 2/	535,101	162	70	127	604	1,742	8,153	26,135	47,744	74,716	160,084	215,545	19
2004 2/	537,512	210	80	125	553	1,745	8,411	25,858	46,934	77,652	165,259	210,669	16
2003 2/	565,808	200	83	146	605	1,788	8,894	25,982	48,091	83,479	176,035	220,489	16
2002	577,761	249	91	146	547	1,718	9,114	26,206	47,494	88,528	181,975	221,670	23
2001	582,349	234	108	152	522	1,775	8,734	25,544	46,433	91,855	185,536	221,430	26
2000	594,465	258	98	140	531	1,653	8,743	25,153	47,292	97,748	189,366	223,454	29
1999	607,704	259	92	156	585	1,755	8,936	24,823	47,950	103,750	195,187	224,193	18
Non-Hispanic white, male													
2005 2/	263,311	89	38	70	422	1,206	5,920	19,485	33,711	46,538	82,264	73,555	13
2004 2/	263,638	116	37	67	363	1,194	6,080	19,329	33,078	48,417	84,694	70,250	13
2003 2/	276,241	110	34	88	368	1,212	6,429	19,504	33,871	52,148	89,985	72,482	10
2002	281,442	139	54	85	370	1,177	6,657	19,774	33,313	55,265	92,616	71,976	16
2001	280,713	127	59	80	336	1,184	6,395	19,178	32,504	57,059	93,996	69,775	20
2000	287,210	139	59	87	326	1,115	6,446	19,101	33,179	61,024	95,576	70,138	20
1999	293,206	145	55	84	357	1,152	6,590	18,747	33,466	64,699	98,422	69,476	13
Non-Hispanic white, female													
2005 2/	271,790	73	32	57	182	536	2,233	6,650	14,033	28,178	77,820	141,990	6
2004 2/	273,874	94	43	58	190	551	2,331	6,529	13,856	29,235	80,565	140,419	3
2003 2/	289,567	90	49	58	237	576	2,465	6,478	14,220	31,331	86,050	148,007	6
2002	296,319	110	37	61	177	541	2,457	6,432	14,181	33,263	89,359	149,694	7
2001	301,636	107	49	72	186	591	2,339	6,366	13,929	34,796	91,540	151,655	6
2000	307,255	119	39	53	205	538	2,297	6,052	14,113	36,724	93,790	153,316	9
1999	314,498	114	37	72	228	603	2,346	6,076	14,484	39,051	96,765	154,717	5

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Diseases of heart (I00-I09,I11,I13,I20-I51)													
Non-Hispanic black, both sexes													
2005 2/	73,302	104	45	75	333	1,008	3,165	8,613	11,977	13,840	18,118	16,016	8
2004 2/	73,373	99	61	78	310	971	3,202	8,572	11,541	14,372	18,139	16,018	10
2003 2/	76,452	129	55	76	346	974	3,392	8,625	11,664	15,409	19,046	16,734	2
2002	76,694	142	40	65	299	993	3,312	8,318	11,554	15,411	19,520	17,030	10
2001	76,794	140	73	70	322	961	3,424	7,956	11,209	15,940	19,681	17,004	14
2000	76,706	118	47	75	331	929	3,259	7,606	11,257	16,413	19,600	17,059	12
1999	77,713	141	49	77	336	913	3,506	7,448	11,346	17,019	19,808	17,056	14
Non-Hispanic black, male													
2005 2/	35,874	61	19	35	228	624	1,969	5,497	7,404	7,586	7,952	4,495	4
2004 2/	35,749	55	32	35	202	588	1,936	5,393	7,149	7,798	7,931	4,621	9
2003 2/	36,962	79	31	42	225	572	2,066	5,443	7,040	8,377	8,488	4,597	2
2002	36,596	76	18	33	185	612	2,031	5,214	7,028	8,247	8,428	4,718	6
2001	36,541	65	40	34	205	578	2,071	4,955	6,713	8,491	8,621	4,758	10
2000	36,282	59	24	41	192	543	1,972	4,787	6,786	8,560	8,476	4,835	7
1999	37,077	71	27	45	212	546	2,165	4,716	6,907	9,016	8,416	4,945	11
Non-Hispanic black, female													
2005 2/	37,428	43	26	40	105	384	1,196	3,116	4,573	6,254	10,166	11,521	4
2004 2/	37,624	44	29	43	108	383	1,266	3,179	4,392	6,574	10,208	11,397	1
2003 2/	39,490	50	24	34	121	402	1,326	3,182	4,624	7,032	10,558	12,137	-
2002	40,098	66	22	32	114	381	1,281	3,104	4,526	7,164	11,092	12,312	4
2001	40,253	75	33	36	117	383	1,353	3,001	4,496	7,449	11,060	12,246	4
2000	40,424	59	23	34	139	386	1,287	2,819	4,471	7,853	11,124	12,224	5
1999	40,636	70	22	32	124	367	1,341	2,732	4,439	8,003	11,392	12,111	3

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Diseases of heart (I00-I09,I11,I13,I20-I51)													
Origin not stated 3/, both sexes													
2005	1,276	-	-	1	-	6	29	130	207	233	354	312	4
2004	1,533	2	-	2	5	6	49	135	223	322	410	373	6
2003	1,963	2	1	1	5	12	59	152	301	367	541	513	9
2002	2,370	2	-	-	2	17	56	174	277	477	646	707	12
2001	2,265	3	1	-	10	12	61	165	296	388	656	655	18
2000	2,471	4	1	1	2	14	80	175	320	476	665	720	13
1999	2,607	1	1	1	3	10	72	159	328	535	747	741	9
Origin not stated 3/, male													
2005	744	-	-	-	-	4	23	96	156	168	201	94	2
2004	907	1	-	2	3	6	33	103	170	225	231	127	6
2003	1,133	-	-	1	3	7	44	119	231	238	300	182	8
2002	1,287	1	-	-	2	11	41	136	202	305	347	232	10
2001	1,227	2	-	-	7	10	48	124	227	264	339	195	11
2000	1,353	3	-	-	1	7	63	133	233	334	334	235	10
1999	1,448	-	-	1	2	7	59	122	250	363	382	255	7
Origin not stated 3/, female													
2005	532	-	-	1	-	2	6	34	51	65	153	218	2
2004	626	1	-	-	2	-	16	32	53	97	179	246	-
2003	830	2	1	-	2	5	15	33	70	129	241	331	1
2002	1,083	1	-	-	-	6	15	38	75	172	299	475	2
2001	1,038	1	1	-	3	2	13	41	69	124	317	460	7
2000	1,118	1	1	1	1	7	17	42	87	142	331	485	3
1999	1,159	1	1	-	1	3	13	37	78	172	365	486	2

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Acute rheumatic fever and chronic rheumatic heart diseases (I00-I09)													
All origins, both sexes													
2005	3,365	1	1	4	9	27	82	191	354	613	1,114	969	-
2004	3,254	2	3	4	7	27	96	207	321	612	1,146	829	-
2003	3,624	1	1	6	11	25	99	224	367	693	1,233	964	-
2002	3,579	4	1	6	18	36	95	237	433	703	1,227	819	-
2001	3,489	3	1	5	11	34	130	232	361	744	1,188	780	-
2000	3,582	6	2	4	17	33	141	252	420	819	1,176	712	-
1999	3,676	6	1	5	13	42	124	245	411	854	1,227	748	-
All origins, male													
2005	1,044	1	-	2	7	14	38	81	141	210	326	224	-
2004	1,009	2	3	2	4	9	38	88	107	202	380	174	-
2003	1,153	1	-	4	7	14	44	78	146	241	384	234	-
2002	1,078	3	1	2	8	18	45	95	156	245	341	164	-
2001	1,031	3	1	2	7	15	54	87	135	231	330	166	-
2000	1,072	5	2	3	11	12	61	104	166	237	321	150	-
1999	1,042	2	1	2	10	17	49	90	133	290	325	123	-
All origins, female													
2005	2,321	-	1	2	2	13	44	110	213	403	788	745	-
2004	2,245	-	-	2	3	18	58	119	214	410	766	655	-
2003	2,471	-	1	2	4	11	55	146	221	452	849	730	-
2002	2,501	1	-	4	10	18	50	142	277	458	886	655	-
2001	2,458	-	-	3	4	19	76	145	226	513	858	614	-
2000	2,510	1	-	1	6	21	80	148	254	582	855	562	-
1999	2,634	4	-	3	3	25	75	155	278	564	902	625	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Acute rheumatic fever and chronic rheumatic heart diseases (I00-I09)													
Hispanic, both sexes													
2005	162	-	-	1	-	6	8	21	25	40	41	20	-
2004	158	-	1	1	1	2	14	15	29	44	43	8	-
2003	159	1	-	1	2	7	7	21	21	40	43	16	-
2002	198	-	-	-	2	11	11	24	35	54	42	19	-
2001	156	-	1	1	2	2	12	16	27	48	30	17	-
2000	154	1	-	-	1	6	13	19	30	47	27	10	-
1999	166	2	-	-	3	4	16	23	26	44	35	13	-
Hispanic, male													
2005	47	-	-	1	-	4	4	9	8	13	6	2	-
2004	56	-	1	1	1	1	3	6	5	18	17	3	-
2003	57	1	-	1	2	4	4	8	8	13	11	5	-
2002	55	-	-	-	1	5	4	9	13	15	4	4	-
2001	51	-	1	1	1	2	5	5	7	15	11	3	-
2000	49	1	-	-	-	1	4	9	12	13	6	3	-
1999	38	-	-	-	3	3	3	6	8	10	5	-	-
Hispanic, female													
2005	115	-	-	-	-	2	4	12	17	27	35	18	-
2004	102	-	-	-	-	1	11	9	24	26	26	5	-
2003	102	-	-	-	-	3	3	13	13	27	32	11	-
2002	143	-	-	-	1	6	7	15	22	39	38	15	-
2001	105	-	-	-	1	-	7	11	20	33	19	14	-
2000	105	-	-	-	1	5	9	10	18	34	21	7	-
1999	128	2	-	-	-	1	13	17	18	34	30	13	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Acute rheumatic fever and chronic rheumatic heart diseases (I00-I09)													
Non-Hispanic 1/, both sexes													
2005	3,200	1	1	3	9	21	74	168	329	572	1,073	949	-
2004	3,091	2	2	3	6	25	81	192	291	568	1,100	821	-
2003	3,460	-	1	5	9	18	92	203	344	653	1,190	945	-
2002	3,375	4	1	6	16	25	84	212	397	647	1,183	800	-
2001	3,327	3	-	4	8	32	118	214	333	696	1,157	762	-
2000	3,424	5	2	4	16	27	128	233	388	772	1,149	700	-
1999	3,498	4	1	5	10	38	106	221	383	806	1,190	734	-
Non-Hispanic 1/, male													
2005	995	1	-	1	7	10	34	71	133	196	320	222	-
2004	950	2	2	1	3	8	34	82	101	184	362	171	-
2003	1,095	-	-	3	5	10	40	70	137	228	373	229	-
2002	1,021	3	1	2	7	13	41	85	142	230	337	160	-
2001	978	3	-	1	5	13	49	81	128	216	319	163	-
2000	1,022	4	2	3	11	11	57	95	153	224	315	147	-
1999	999	2	1	2	7	14	45	83	123	279	320	123	-
Non-Hispanic 1/, female													
2005	2,205	-	1	2	2	11	40	97	196	376	753	727	-
2004	2,141	-	-	2	3	17	47	110	190	384	738	650	-
2003	2,365	-	1	2	4	8	52	133	207	425	817	716	-
2002	2,354	1	-	4	9	12	43	127	255	417	846	640	-
2001	2,349	-	-	3	3	19	69	133	205	480	838	599	-
2000	2,402	1	-	1	5	16	71	138	235	548	834	553	-
1999	2,499	2	-	3	3	24	61	138	260	527	870	611	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Acute rheumatic fever and chronic rheumatic heart diseases (I00-I09)													
Non-Hispanic white, both sexes													
2005 2/	2,866	1	1	3	4	11	49	112	259	506	1,003	917	-
2004 2/	2,754	2	1	1	3	14	43	139	228	499	1,034	790	-
2003 2/	3,066	-	1	3	3	12	55	128	254	571	1,129	910	-
2002	3,007	4	1	3	8	14	57	156	313	575	1,109	767	-
2001	2,924	1	-	2	7	13	73	139	265	601	1,094	729	-
2000	3,073	4	2	4	9	19	83	178	320	687	1,102	665	-
1999	3,130	2	1	3	5	21	68	163	314	730	1,125	698	-
Non-Hispanic white, male													
2005 2/	878	1	-	1	2	7	24	50	108	175	299	211	-
2004 2/	840	2	1	-	2	1	19	59	77	167	346	166	-
2003 2/	967	-	-	2	2	6	23	48	98	203	364	221	-
2002	897	3	1	2	3	7	26	63	113	205	319	155	-
2001	857	1	-	-	5	7	35	58	108	187	302	154	-
2000	917	3	2	3	6	9	42	78	136	197	303	138	-
1999	892	1	1	-	2	9	31	65	102	260	301	120	-
Non-Hispanic white, female													
2005 2/	1,988	-	1	2	2	4	25	62	151	331	704	706	-
2004 2/	1,914	-	-	1	1	13	24	80	151	332	688	624	-
2003 2/	2,099	-	1	1	1	6	32	80	156	368	765	689	-
2002	2,110	1	-	1	5	7	31	93	200	370	790	612	-
2001	2,067	-	-	2	2	6	38	81	157	414	792	575	-
2000	2,156	1	-	1	3	10	41	100	184	490	799	527	-
1999	2,238	1	-	3	3	12	37	98	212	470	824	578	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Acute rheumatic fever and chronic rheumatic heart diseases (I00-I09)													
Non-Hispanic black, both sexes													
2005 2/	225	-	-	-	2	8	16	47	44	41	44	23	-
2004 2/	245	-	1	2	3	6	33	48	46	45	40	21	-
2003 2/	305	-	-	2	4	5	29	63	72	66	40	24	-
2002	249	-	-	2	5	9	20	44	62	48	42	17	-
2001	302	2	-	1	1	13	41	60	53	69	40	22	-
2000	263	1	-	-	7	8	35	43	48	58	36	27	-
1999	269	1	-	2	4	13	30	49	51	54	39	26	-
Non-Hispanic black, male													
2005 2/	80	-	-	-	2	2	8	18	17	12	15	6	-
2004 2/	83	-	1	1	1	4	14	20	19	12	8	3	-
2003 2/	109	-	-	1	2	3	15	20	33	21	8	6	-
2002	91	-	-	-	3	4	12	17	21	20	11	3	-
2001	86	2	-	-	-	3	13	20	17	17	8	6	-
2000	77	1	-	-	5	2	12	13	12	16	8	8	-
1999	86	1	-	2	4	3	12	18	16	15	14	1	-
Non-Hispanic black, female													
2005 2/	145	-	-	-	-	6	8	29	27	29	29	17	-
2004 2/	162	-	-	1	2	2	19	28	27	33	32	18	-
2003 2/	196	-	-	1	2	2	14	43	39	45	32	18	-
2002	158	-	-	2	2	5	8	27	41	28	31	14	-
2001	216	-	-	1	1	10	28	40	36	52	32	16	-
2000	186	-	-	-	2	6	23	30	36	42	28	19	-
1999	183	-	-	-	-	10	18	31	35	39	25	25	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Acute rheumatic fever and chronic rheumatic heart diseases (I00-I09)													
Origin not stated 3/, both sexes													
2005	3	-	-	-	-	-	-	2	-	1	-	-	-
2004	5	-	-	-	-	-	1	-	1	-	3	-	-
2003	5	-	-	-	-	-	-	-	2	-	-	3	-
2002	6	-	-	-	-	-	-	1	1	2	2	-	-
2001	6	-	-	-	1	-	-	2	1	-	1	1	-
2000	4	-	-	-	-	-	-	-	2	-	-	2	-
1999	12	-	-	-	-	-	2	1	2	4	2	1	-
Origin not stated 3/, male													
2005	2	-	-	-	-	-	-	1	-	1	-	-	-
2004	3	-	-	-	-	-	1	-	1	-	1	-	-
2003	1	-	-	-	-	-	-	-	1	-	-	-	-
2002	2	-	-	-	-	-	-	1	1	-	-	-	-
2001	2	-	-	-	1	-	-	1	-	-	-	-	-
2000	1	-	-	-	-	-	-	-	1	-	-	-	-
1999	5	-	-	-	-	-	1	1	2	1	-	-	-
Origin not stated 3/, female													
2005	1	-	-	-	-	-	-	1	-	-	-	-	-
2004	2	-	-	-	-	-	-	-	-	-	2	-	-
2003	4	-	-	-	-	-	-	-	1	-	-	3	-
2002	4	-	-	-	-	-	-	-	-	2	2	-	-
2001	4	-	-	-	-	-	-	1	1	-	1	1	-
2000	3	-	-	-	-	-	-	-	1	-	-	2	-
1999	7	-	-	-	-	-	1	-	-	3	2	1	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Hypertensive heart disease (I11)													
All origins, both sexes													
2005	29,282	2	-	3	36	328	1,368	3,293	3,954	3,806	6,370	10,119	3
2004	28,585	-	-	2	44	302	1,317	3,265	3,885	4,001	6,398	9,368	3
2003	28,345	-	1	-	53	305	1,306	3,230	3,651	4,045	6,565	9,189	-
2002	26,551	-	-	2	30	297	1,213	2,882	3,320	3,895	6,321	8,587	4
2001	24,689	-	-	1	24	246	1,091	2,464	2,769	3,619	6,119	8,345	11
2000	23,761	-	1	2	32	213	981	2,104	2,672	3,605	5,962	8,186	3
1999	22,702	1	-	-	33	210	964	1,930	2,537	3,517	5,830	7,679	1
All origins, male													
2005	13,219	1	-	2	24	222	958	2,278	2,649	2,147	2,570	2,365	3
2004	12,929	-	-	-	32	223	896	2,192	2,584	2,303	2,504	2,192	3
2003	12,536	-	1	-	40	208	877	2,157	2,308	2,264	2,627	2,054	-
2002	11,660	-	-	1	26	215	849	1,938	2,159	2,088	2,509	1,871	4
2001	10,451	-	-	-	17	169	740	1,646	1,749	1,961	2,414	1,746	9
2000	9,876	-	-	1	24	139	643	1,436	1,684	1,971	2,269	1,707	2
1999	9,316	-	-	-	25	149	660	1,259	1,560	1,816	2,252	1,594	1
All origins, female													
2005	16,063	1	-	1	12	106	410	1,015	1,305	1,659	3,800	7,754	-
2004	15,656	-	-	2	12	79	421	1,073	1,301	1,698	3,894	7,176	-
2003	15,809	-	-	-	13	97	429	1,073	1,343	1,781	3,938	7,135	-
2002	14,891	-	-	1	4	82	364	944	1,161	1,807	3,812	6,716	-
2001	14,238	-	-	1	7	77	351	818	1,020	1,658	3,705	6,599	2
2000	13,885	-	1	1	8	74	338	668	988	1,634	3,693	6,479	1
1999	13,386	1	-	-	8	61	304	671	977	1,701	3,578	6,085	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Hypertensive heart disease (I11)													
Hispanic, both sexes													
2005	1,483	-	-	1	3	34	111	209	256	218	312	339	-
2004	1,491	-	-	1	4	24	105	193	244	282	322	316	-
2003	1,586	-	-	-	12	27	109	218	247	280	347	346	-
2002	1,424	-	-	-	4	31	97	180	241	264	308	299	-
2001	1,211	-	-	-	5	22	93	165	184	239	260	243	-
2000	991	-	-	-	1	20	64	118	141	198	233	216	-
1999	933	-	-	-	3	18	61	99	146	174	215	217	-
Hispanic, male													
2005	772	-	-	1	2	27	83	150	162	115	137	95	-
2004	806	-	-	-	2	19	86	124	169	157	155	94	-
2003	876	-	-	-	10	22	91	158	168	167	166	94	-
2002	781	-	-	-	4	25	78	135	159	149	149	82	-
2001	648	-	-	-	4	15	75	118	120	132	108	76	-
2000	495	-	-	-	1	16	43	84	86	106	103	56	-
1999	455	-	-	-	2	16	45	69	96	95	82	50	-
Hispanic, female													
2005	711	-	-	-	1	7	28	59	94	103	175	244	-
2004	685	-	-	1	2	5	19	69	75	125	167	222	-
2003	710	-	-	-	2	5	18	60	79	113	181	252	-
2002	643	-	-	-	-	6	19	45	82	115	159	217	-
2001	563	-	-	-	1	7	18	47	64	107	152	167	-
2000	496	-	-	-	-	4	21	34	55	92	130	160	-
1999	478	-	-	-	1	2	16	30	50	79	133	167	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Hypertensive heart disease (I11)													
Non-Hispanic 1/, both sexes													
2005	27,701	2	-	2	33	294	1,251	3,071	3,681	3,571	6,035	9,759	2
2004	26,972	-	-	1	39	278	1,204	3,049	3,618	3,690	6,059	9,032	2
2003	26,632	-	1	-	41	277	1,189	2,990	3,378	3,741	6,199	8,816	-
2002	24,976	-	-	2	26	265	1,109	2,679	3,060	3,591	5,985	8,257	2
2001	23,343	-	-	1	18	222	991	2,270	2,554	3,364	5,837	8,077	9
2000	22,646	-	1	2	31	193	905	1,968	2,518	3,374	5,706	7,946	2
1999	21,639	1	-	-	30	192	898	1,815	2,363	3,315	5,591	7,433	1
Non-Hispanic 1/, male													
2005	12,389	1	-	1	22	195	870	2,119	2,473	2,018	2,423	2,265	2
2004	12,051	-	-	-	29	204	806	2,050	2,396	2,129	2,343	2,092	2
2003	11,588	-	1	-	30	186	780	1,981	2,121	2,082	2,452	1,955	-
2002	10,790	-	-	1	22	190	765	1,784	1,988	1,916	2,344	1,778	2
2001	9,725	-	-	-	12	152	660	1,509	1,608	1,815	2,299	1,663	7
2000	9,298	-	-	1	23	123	591	1,338	1,586	1,839	2,152	1,644	1
1999	8,790	-	-	-	23	133	610	1,180	1,445	1,702	2,157	1,539	1
Non-Hispanic 1/, female													
2005	15,312	1	-	1	11	99	381	952	1,208	1,553	3,612	7,494	-
2004	14,921	-	-	1	10	74	398	999	1,222	1,561	3,716	6,940	-
2003	15,044	-	-	-	11	91	409	1,009	1,257	1,659	3,747	6,861	-
2002	14,186	-	-	1	4	75	344	895	1,072	1,675	3,641	6,479	-
2001	13,618	-	-	1	6	70	331	761	946	1,549	3,538	6,414	2
2000	13,348	-	1	1	8	70	314	630	932	1,535	3,554	6,302	1
1999	12,849	1	-	-	7	59	288	635	918	1,613	3,434	5,894	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Hypertensive heart disease (I11)													
Non-Hispanic white, both sexes													
2005 2/	20,219	1	-	1	9	140	666	1,735	2,355	2,381	4,616	8,313	2
2004 2/	19,630	-	-	-	19	121	652	1,718	2,229	2,508	4,661	7,720	2
2003 2/	19,286	-	-	-	15	133	623	1,606	2,087	2,465	4,823	7,534	-
2002	18,060	-	-	1	10	110	587	1,519	1,832	2,380	4,574	7,046	1
2001	16,991	-	-	-	5	102	493	1,213	1,520	2,198	4,540	6,915	5
2000	16,373	-	-	2	14	75	476	1,042	1,441	2,193	4,386	6,744	-
1999	15,654	1	-	-	9	91	428	954	1,371	2,147	4,294	6,359	-
Non-Hispanic white, male													
2005 2/	8,644	1	-	1	6	106	497	1,275	1,638	1,367	1,868	1,883	2
2004 2/	8,401	-	-	-	16	102	473	1,258	1,507	1,475	1,810	1,758	2
2003 2/	8,056	-	-	-	11	101	463	1,172	1,383	1,389	1,908	1,629	-
2002	7,427	-	-	1	9	83	436	1,088	1,239	1,294	1,793	1,483	1
2001	6,728	-	-	-	3	80	367	866	1,016	1,237	1,781	1,374	4
2000	6,390	-	-	1	11	53	341	771	956	1,242	1,640	1,375	-
1999	6,013	-	-	-	7	75	323	676	882	1,122	1,647	1,281	-
Non-Hispanic white, female													
2005 2/	11,575	-	-	-	3	34	169	460	717	1,014	2,748	6,430	-
2004 2/	11,229	-	-	-	3	19	179	460	722	1,033	2,851	5,962	-
2003 2/	11,230	-	-	-	4	32	160	434	704	1,076	2,915	5,905	-
2002	10,633	-	-	-	1	27	151	431	593	1,086	2,781	5,563	-
2001	10,263	-	-	-	2	22	126	347	504	961	2,759	5,541	1
2000	9,983	-	-	1	3	22	135	271	485	951	2,746	5,369	-
1999	9,641	1	-	-	2	16	105	278	489	1,025	2,647	5,078	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Hypertensive heart disease (I11)													
Non-Hispanic black, both sexes													
2005 2/	6,825	1	-	1	23	138	537	1,256	1,246	1,099	1,272	1,252	-
2004 2/	6,718	-	-	1	20	147	524	1,273	1,290	1,102	1,254	1,107	-
2003 2/	6,759	-	1	-	25	134	539	1,321	1,210	1,170	1,233	1,126	-
2002	6,385	-	-	1	15	149	491	1,115	1,158	1,124	1,267	1,065	-
2001	5,880	-	-	1	13	114	474	1,007	972	1,078	1,164	1,053	4
2000	5,828	-	1	-	17	114	414	886	1,025	1,093	1,194	1,082	2
1999	5,556	-	-	-	20	98	447	838	932	1,084	1,182	954	1
Non-Hispanic black, male													
2005 2/	3,402	-	-	-	15	76	338	785	782	591	495	320	-
2004 2/	3,336	-	-	-	13	92	312	749	820	614	469	267	-
2003 2/	3,248	-	1	-	19	78	299	762	687	646	482	274	-
2002	3,089	-	-	-	12	101	307	662	707	574	484	242	-
2001	2,760	-	-	-	9	67	275	609	551	538	458	250	3
2000	2,685	-	-	-	12	66	240	538	594	547	452	235	1
1999	2,571	-	-	-	15	55	267	487	528	536	458	224	1
Non-Hispanic black, female													
2005 2/	3,423	1	-	1	8	62	199	471	464	508	777	932	-
2004 2/	3,382	-	-	1	7	55	212	524	470	488	785	840	-
2003 2/	3,511	-	-	-	6	56	240	559	523	524	751	852	-
2002	3,296	-	-	1	3	48	184	453	451	550	783	823	-
2001	3,120	-	-	1	4	47	199	398	421	540	706	803	1
2000	3,143	-	1	-	5	48	174	348	431	546	742	847	1
1999	2,985	-	-	-	5	43	180	351	404	548	724	730	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Hypertensive heart disease (I11)													
Origin not stated 3/, both sexes													
2005	98	-	-	-	-	-	6	13	17	17	23	21	1
2004	122	-	-	-	1	-	8	23	23	29	17	20	1
2003	127	-	-	-	-	1	8	22	26	24	19	27	-
2002	151	-	-	-	-	1	7	23	19	40	28	31	2
2001	135	-	-	-	1	2	7	29	31	16	22	25	2
2000	124	-	-	-	-	-	12	18	13	33	23	24	1
1999	130	-	-	-	-	-	5	16	28	28	24	29	-
Origin not stated 3/, male													
2005	58	-	-	-	-	-	5	9	14	14	10	5	1
2004	72	-	-	-	1	-	4	18	19	17	6	6	1
2003	72	-	-	-	-	-	6	18	19	15	9	5	-
2002	89	-	-	-	-	-	6	19	12	23	16	11	2
2001	78	-	-	-	1	2	5	19	21	14	7	7	2
2000	83	-	-	-	-	-	9	14	12	26	14	7	1
1999	71	-	-	-	-	-	5	10	19	19	13	5	-
Origin not stated 3/, female													
2005	40	-	-	-	-	-	1	4	3	3	13	16	-
2004	50	-	-	-	-	-	4	5	4	12	11	14	-
2003	55	-	-	-	-	1	2	4	7	9	10	22	-
2002	62	-	-	-	-	1	1	4	7	17	12	20	-
2001	57	-	-	-	-	-	2	10	10	2	15	18	-
2000	41	-	-	-	-	-	3	4	1	7	9	17	-
1999	59	-	-	-	-	-	-	6	9	9	11	24	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Hypertensive heart and renal disease (I13)													
All origins, both sexes													
2005	3,172	-	-	-	5	33	83	189	240	426	908	1,288	-
2004	3,046	-	-	1	7	23	96	209	271	379	854	1,206	-
2003	3,108	-	1	-	5	28	93	201	267	401	903	1,209	-
2002	2,895	-	-	-	5	28	77	165	273	402	866	1,079	-
2001	2,826	1	-	-	7	28	99	146	247	417	855	1,026	-
2000	2,785	-	-	1	6	28	83	147	208	417	863	1,032	-
1999	3,327	-	-	2	6	21	114	156	262	470	1,087	1,209	-
All origins, male													
2005	1,369	-	-	-	4	20	52	125	142	209	378	439	-
2004	1,301	-	-	-	4	14	69	123	158	189	377	367	-
2003	1,326	-	1	-	4	13	62	130	159	189	382	386	-
2002	1,205	-	-	-	3	19	44	105	158	198	357	321	-
2001	1,169	1	-	-	4	17	63	99	126	206	339	314	-
2000	1,163	-	-	-	4	18	49	87	108	206	350	341	-
1999	1,390	-	-	1	3	16	67	100	146	223	460	374	-
All origins, female													
2005	1,803	-	-	-	1	13	31	64	98	217	530	849	-
2004	1,745	-	-	1	3	9	27	86	113	190	477	839	-
2003	1,782	-	-	-	1	15	31	71	108	212	521	823	-
2002	1,690	-	-	-	2	9	33	60	115	204	509	758	-
2001	1,657	-	-	-	3	11	36	47	121	211	516	712	-
2000	1,622	-	-	1	2	10	34	60	100	211	513	691	-
1999	1,937	-	-	1	3	5	47	56	116	247	627	835	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Hypertensive heart and renal disease (I13)													
Hispanic, both sexes													
2005	199	-	-	-	1	1	9	13	28	43	52	52	-
2004	190	-	-	-	1	4	12	18	29	23	61	42	-
2003	158	-	1	-	1	-	6	11	19	33	38	49	-
2002	172	-	-	-	2	3	4	11	26	34	51	41	-
2001	160	-	-	-	1	2	9	10	21	34	41	42	-
2000	138	-	-	-	1	3	8	14	15	27	32	38	-
1999	144	-	-	-	-	1	9	11	19	32	39	33	-
Hispanic, male													
2005	96	-	-	-	1	-	7	10	17	19	25	17	-
2004	96	-	-	-	1	4	8	10	16	11	29	17	-
2003	75	-	1	-	1	-	1	7	11	12	19	23	-
2002	80	-	-	-	1	1	2	5	18	18	20	15	-
2001	76	-	-	-	1	-	5	6	14	17	17	16	-
2000	67	-	-	-	1	2	4	7	5	19	18	11	-
1999	65	-	-	-	-	-	6	6	12	14	14	13	-
Hispanic, female													
2005	103	-	-	-	-	1	2	3	11	24	27	35	-
2004	94	-	-	-	-	-	4	8	13	12	32	25	-
2003	83	-	-	-	-	-	5	4	8	21	19	26	-
2002	92	-	-	-	1	2	2	6	8	16	31	26	-
2001	84	-	-	-	-	2	4	4	7	17	24	26	-
2000	71	-	-	-	-	1	4	7	10	8	14	27	-
1999	79	-	-	-	-	1	3	5	7	18	25	20	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Hypertensive heart and renal disease (I13)													
Non-Hispanic 1/, both sexes													
2005	2,963	-	-	-	4	32	74	174	210	380	854	1,235	-
2004	2,845	-	-	1	6	19	83	189	241	353	793	1,160	-
2003	2,944	-	-	-	4	28	86	190	247	368	863	1,158	-
2002	2,714	-	-	-	3	25	73	153	247	365	812	1,036	-
2001	2,650	1	-	-	6	26	87	135	225	380	810	980	-
2000	2,637	-	-	1	5	25	75	133	192	389	826	991	-
1999	3,171	-	-	2	6	20	102	142	242	436	1,047	1,174	-
Non-Hispanic 1/, male													
2005	1,266	-	-	-	3	20	45	114	123	188	351	422	-
2004	1,199	-	-	-	3	10	60	112	141	175	348	350	-
2003	1,248	-	-	-	3	13	60	123	147	177	362	363	-
2002	1,120	-	-	-	2	18	42	99	140	178	336	305	-
2001	1,086	1	-	-	3	17	56	92	111	188	321	297	-
2000	1,089	-	-	-	3	16	45	80	102	186	329	328	-
1999	1,318	-	-	1	3	16	59	91	134	208	446	360	-
Non-Hispanic 1/, female													
2005	1,697	-	-	-	1	12	29	60	87	192	503	813	-
2004	1,646	-	-	1	3	9	23	77	100	178	445	810	-
2003	1,696	-	-	-	1	15	26	67	100	191	501	795	-
2002	1,594	-	-	-	1	7	31	54	107	187	476	731	-
2001	1,564	-	-	-	3	9	31	43	114	192	489	683	-
2000	1,548	-	-	1	2	9	30	53	90	203	497	663	-
1999	1,853	-	-	1	3	4	43	51	108	228	601	814	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Hypertensive heart and renal disease (I13)													
Non-Hispanic white, both sexes													
2005 2/	1,989	-	-	-	1	6	19	50	67	195	605	1,046	-
2004 2/	1,878	-	-	1	1	4	16	41	90	187	576	962	-
2003 2/	1,942	-	-	-	1	7	14	42	92	196	646	944	-
2002	1,752	-	-	-	-	8	17	29	82	184	604	828	-
2001	1,759	-	-	-	2	5	16	43	95	204	601	793	-
2000	1,769	-	-	-	-	5	30	40	75	220	593	806	-
1999	2,091	-	-	1	-	6	19	43	91	215	768	948	-
Non-Hispanic white, male													
2005 2/	791	-	-	-	1	4	12	36	41	96	249	352	-
2004 2/	748	-	-	-	1	3	10	25	57	100	257	295	-
2003 2/	789	-	-	-	-	2	12	30	57	106	288	294	-
2002	679	-	-	-	-	5	13	19	47	96	256	243	-
2001	687	-	-	-	2	3	12	27	42	101	250	250	-
2000	710	-	-	-	-	4	18	24	41	109	256	258	-
1999	843	-	-	1	-	6	11	26	48	108	346	297	-
Non-Hispanic white, female													
2005 2/	1,198	-	-	-	-	2	7	14	26	99	356	694	-
2004 2/	1,130	-	-	1	-	1	6	16	33	87	319	667	-
2003 2/	1,153	-	-	-	1	5	2	12	35	90	358	650	-
2002	1,073	-	-	-	-	3	4	10	35	88	348	585	-
2001	1,072	-	-	-	-	2	4	16	53	103	351	543	-
2000	1,059	-	-	-	-	1	12	16	34	111	337	548	-
1999	1,248	-	-	-	-	-	8	17	43	107	422	651	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Hypertensive heart and renal disease (I13)													
Non-Hispanic black, both sexes													
2005 2/	879	-	-	-	3	25	54	121	136	168	215	157	-
2004 2/	877	-	-	-	5	15	65	139	142	146	194	171	-
2003 2/	910	-	-	-	3	20	71	142	148	154	191	181	-
2002	897	-	-	-	2	17	54	124	159	165	187	189	-
2001	821	1	-	-	4	21	65	92	125	164	187	162	-
2000	797	-	-	1	5	19	45	91	112	161	203	160	-
1999	989	-	-	1	6	13	82	92	143	205	243	204	-
Non-Hispanic black, male													
2005 2/	426	-	-	-	2	16	33	77	80	81	84	53	-
2004 2/	404	-	-	-	2	7	49	80	79	62	83	42	-
2003 2/	411	-	-	-	3	10	47	88	86	63	66	48	-
2002	402	-	-	-	1	13	27	80	88	70	66	57	-
2001	370	1	-	-	1	14	41	65	66	81	62	39	-
2000	348	-	-	-	3	12	27	55	57	74	65	55	-
1999	426	-	-	-	3	9	47	60	81	92	83	51	-
Non-Hispanic black, female													
2005 2/	453	-	-	-	1	9	21	44	56	87	131	104	-
2004 2/	473	-	-	-	3	8	16	59	63	84	111	129	-
2003 2/	499	-	-	-	-	10	24	54	62	91	125	133	-
2002	495	-	-	-	1	4	27	44	71	95	121	132	-
2001	451	-	-	-	3	7	24	27	59	83	125	123	-
2000	449	-	-	1	2	7	18	36	55	87	138	105	-
1999	563	-	-	1	3	4	35	32	62	113	160	153	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Hypertensive heart and renal disease (I13)													
Origin not stated 3/, both sexes													
2005	10	-	-	-	-	-	-	2	2	3	2	1	-
2004	11	-	-	-	-	-	1	2	1	3	-	4	-
2003	6	-	-	-	-	-	1	-	1	-	2	2	-
2002	9	-	-	-	-	-	-	1	-	3	3	2	-
2001	16	-	-	-	-	-	3	1	1	3	4	4	-
2000	10	-	-	-	-	-	-	-	1	1	5	3	-
1999	12	-	-	-	-	-	3	3	1	2	1	2	-
Origin not stated 3/, male													
2005	7	-	-	-	-	-	-	1	2	2	2	-	-
2004	6	-	-	-	-	-	1	1	1	3	-	-	-
2003	3	-	-	-	-	-	1	-	1	-	1	-	-
2002	5	-	-	-	-	-	-	1	-	2	1	1	-
2001	7	-	-	-	-	-	2	1	1	1	1	1	-
2000	7	-	-	-	-	-	-	-	1	1	3	2	-
1999	7	-	-	-	-	-	2	3	-	1	-	1	-
Origin not stated 3/, female													
2005	3	-	-	-	-	-	-	1	-	1	-	1	-
2004	5	-	-	-	-	-	-	1	-	-	-	4	-
2003	3	-	-	-	-	-	-	-	-	-	1	2	-
2002	4	-	-	-	-	-	-	-	-	1	2	1	-
2001	9	-	-	-	-	-	1	-	-	2	3	3	-
2000	3	-	-	-	-	-	-	-	-	-	2	1	-
1999	5	-	-	-	-	-	1	-	1	1	1	1	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Ischemic heart diseases (I20-I25)													
All origins, both sexes													
2005	445,687	13	7	14	151	1,014	6,860	25,310	46,799	70,121	134,435	160,935	28
2004	451,326	13	10	17	125	1,092	7,132	24,923	46,002	73,102	139,128	159,756	26
2003	480,028	22	12	24	160	1,082	7,547	25,361	47,377	78,880	149,370	170,164	29
2002	494,382	23	7	20	133	996	7,683	25,575	47,262	83,867	155,391	173,388	37
2001	502,189	26	9	20	163	1,116	7,534	25,073	46,574	87,463	159,854	174,309	48
2000	515,204	30	10	16	123	935	7,519	24,890	47,737	92,885	163,692	177,319	48
1999	529,659	27	5	22	137	1,014	7,751	24,775	48,377	98,840	169,296	179,376	39
All origins, male													
2005	232,115	7	4	9	121	774	5,188	19,233	33,548	44,497	71,352	57,364	18
2004	233,538	8	1	9	96	822	5,366	18,986	32,868	46,273	73,591	55,495	23
2003	246,342	13	6	10	107	813	5,679	19,481	33,767	49,999	78,573	57,872	22
2002	252,760	15	4	13	92	741	5,840	19,551	33,656	53,136	81,239	58,446	27
2001	254,005	17	3	13	119	830	5,712	19,141	33,006	55,072	82,953	57,106	33
2000	260,574	17	7	9	81	704	5,787	19,149	33,795	58,554	84,747	57,690	34
1999	267,268	13	3	17	93	739	5,970	19,061	34,123	62,187	87,305	57,726	31
All origins, female													
2005	213,572	6	3	5	30	240	1,672	6,077	13,251	25,624	63,083	103,571	10
2004	217,788	5	9	8	29	270	1,766	5,937	13,134	26,829	65,537	104,261	3
2003	233,686	9	6	14	53	269	1,868	5,880	13,610	28,881	70,797	112,292	7
2002	241,622	8	3	7	41	255	1,843	6,024	13,606	30,731	74,152	114,942	10
2001	248,184	9	6	7	44	286	1,822	5,932	13,568	32,391	76,901	117,203	15
2000	254,630	13	3	7	42	231	1,732	5,741	13,942	34,331	78,945	119,629	14
1999	262,391	14	2	5	44	275	1,781	5,714	14,254	36,653	81,991	121,650	8

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Ischemic heart diseases (I20-I25)													
Hispanic, both sexes													
2005	21,774	1	2	4	22	99	503	1,526	2,745	4,317	6,641	5,910	4
2004	20,482	4	2	4	15	95	477	1,424	2,609	4,147	6,208	5,494	3
2003	20,783	2	1	4	22	104	446	1,352	2,546	4,337	6,232	5,731	6
2002	20,941	4	1	3	18	88	442	1,389	2,683	4,510	6,060	5,742	1
2001	20,664	4	1	4	18	109	368	1,314	2,404	4,553	6,095	5,790	4
2000	19,744	6	1	1	12	69	392	1,240	2,457	4,364	5,626	5,572	4
1999	19,936	2	-	2	11	87	366	1,328	2,463	4,492	5,655	5,525	5
Hispanic, male													
2005	11,953	-	2	4	16	80	406	1,189	1,944	2,655	3,489	2,164	4
2004	11,245	2	-	3	14	81	387	1,116	1,852	2,577	3,286	1,924	3
2003	11,054	1	1	3	20	83	341	1,066	1,780	2,680	3,168	1,906	5
2002	11,295	2	-	2	12	76	346	1,058	1,907	2,775	3,099	2,017	1
2001	11,019	3	1	1	13	90	295	1,016	1,667	2,804	3,116	2,010	3
2000	10,562	4	1	1	9	56	322	973	1,726	2,661	2,888	1,918	3
1999	10,604	-	-	2	8	72	310	1,042	1,677	2,770	2,869	1,849	5
Hispanic, female													
2005	9,821	1	-	-	6	19	97	337	801	1,662	3,152	3,746	-
2004	9,237	2	2	1	1	14	90	308	757	1,570	2,922	3,570	-
2003	9,729	1	-	1	2	21	105	286	766	1,657	3,064	3,825	1
2002	9,646	2	1	1	6	12	96	331	776	1,735	2,961	3,725	-
2001	9,645	1	-	3	5	19	73	298	737	1,749	2,979	3,780	1
2000	9,182	2	-	-	3	13	70	267	731	1,703	2,738	3,654	1
1999	9,332	2	-	-	3	15	56	286	786	1,722	2,786	3,676	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Ischemic heart diseases (I20-I25)													
Non-Hispanic 1/, both sexes													
2005	423,007	12	5	10	129	914	6,346	23,703	43,905	65,631	127,530	154,801	21
2004	429,713	9	8	13	110	995	6,629	23,414	43,225	68,698	132,602	153,992	18
2003	457,794	20	11	20	137	974	7,070	23,913	44,605	74,256	142,711	164,061	16
2002	471,655	19	6	17	114	900	7,214	24,069	44,359	78,987	148,834	167,107	29
2001	479,842	21	8	16	144	1,004	7,135	23,663	43,946	82,604	153,239	168,032	30
2000	493,541	23	9	15	111	861	7,082	23,523	45,022	88,137	157,523	171,202	33
1999	507,700	25	5	20	125	926	7,344	23,331	45,666	93,919	163,035	173,278	26
Non-Hispanic 1/, male													
2005	219,626	7	2	5	105	693	4,773	17,980	31,494	41,719	67,707	55,128	13
2004	221,607	6	1	6	82	739	4,960	17,804	30,891	43,510	70,118	53,475	15
2003	234,433	12	5	7	86	727	5,316	18,341	31,813	47,124	75,165	55,826	11
2002	240,462	13	4	11	79	658	5,475	18,400	31,581	50,116	77,866	56,240	19
2001	242,054	13	2	12	105	737	5,392	18,047	31,160	52,059	79,549	54,956	22
2000	248,932	12	6	8	72	645	5,426	18,074	31,880	55,623	81,578	55,585	23
1999	255,524	13	3	15	85	666	5,623	17,928	32,249	59,117	84,127	55,678	20
Non-Hispanic 1/, female													
2005	203,381	5	3	5	24	221	1,573	5,723	12,411	23,912	59,823	99,673	8
2004	208,106	3	7	7	28	256	1,669	5,610	12,334	25,188	62,484	100,517	3
2003	223,361	8	6	13	51	247	1,754	5,572	12,792	27,132	67,546	108,235	5
2002	231,193	6	2	6	35	242	1,739	5,669	12,778	28,871	70,968	110,867	10
2001	237,788	8	6	4	39	267	1,743	5,616	12,786	30,545	73,690	113,076	8
2000	244,609	11	3	7	39	216	1,656	5,449	13,142	32,514	75,945	115,617	10
1999	252,176	12	2	5	40	260	1,721	5,403	13,417	34,802	78,908	117,600	6

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Ischemic heart diseases (I20-I25)													
Non-Hispanic white, both sexes													
2005 2/.....	368,505	6	3	8	85	641	4,826	18,307	35,153	54,742	112,817	141,904	13
2004 2/.....	374,900	6	5	7	57	690	5,128	18,140	34,877	57,266	117,816	140,898	10
2003 2/.....	400,101	14	5	11	91	695	5,434	18,508	36,009	62,110	126,817	150,394	13
2002.....	413,230	11	4	12	76	650	5,565	18,799	35,799	66,553	132,548	153,193	20
2001.....	420,959	14	5	10	86	718	5,456	18,518	35,421	69,681	136,815	154,217	18
2000.....	434,505	16	9	12	80	606	5,437	18,531	36,367	74,867	141,111	157,445	24
1999.....	447,960	16	4	12	76	665	5,607	18,425	37,066	80,129	146,594	159,351	15
Non-Hispanic white, male													
2005 2/.....	191,862	5	-	5	74	485	3,735	14,273	25,811	35,525	60,849	51,092	8
2004 2/.....	193,976	4	1	4	43	527	3,956	14,218	25,505	37,062	63,313	49,335	8
2003 2/.....	205,310	8	2	4	57	527	4,194	14,610	26,363	40,216	67,652	51,669	8
2002.....	211,391	8	2	8	54	493	4,346	14,856	26,108	43,205	70,370	51,928	13
2001.....	212,928	10	1	8	65	539	4,250	14,590	25,776	44,881	72,036	50,758	14
2000.....	219,732	9	6	6	52	473	4,319	14,723	26,398	48,374	74,042	51,313	17
1999.....	225,909	8	3	10	57	482	4,416	14,585	26,754	51,553	76,763	51,267	11
Non-Hispanic white, female													
2005 2/.....	176,643	1	3	3	11	156	1,091	4,034	9,342	19,217	51,968	90,812	5
2004 2/.....	180,924	2	4	3	14	163	1,172	3,922	9,372	20,204	54,503	91,563	2
2003 2/.....	194,791	6	3	7	34	168	1,240	3,898	9,646	21,894	59,165	98,725	5
2002.....	201,839	3	2	4	22	157	1,219	3,943	9,691	23,348	62,178	101,265	7
2001.....	208,031	4	4	2	21	179	1,206	3,928	9,645	24,800	64,779	103,459	4
2000.....	214,773	7	3	6	28	133	1,118	3,808	9,969	26,493	67,069	106,132	7
1999.....	222,051	8	1	2	19	183	1,191	3,840	10,312	28,576	69,831	108,084	4

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Ischemic heart diseases (I20-I25)													
Non-Hispanic black, both sexes													
2005 2/.....	45,435	5	1	2	35	218	1,310	4,718	7,576	9,103	12,096	10,364	7
2004 2/.....	46,064	2	1	6	46	257	1,315	4,664	7,292	9,717	12,200	10,557	7
2003 2/.....	48,617	4	5	8	41	229	1,407	4,726	7,439	10,360	13,207	11,190	1
2002.....	49,522	7	2	5	31	210	1,441	4,665	7,504	10,598	13,541	11,509	9
2001.....	50,295	7	3	5	52	246	1,474	4,533	7,451	11,107	13,782	11,625	10
2000.....	50,659	4	-	2	24	229	1,451	4,418	7,587	11,485	13,809	11,641	9
1999.....	51,368	8	1	8	44	213	1,552	4,362	7,542	11,929	13,890	11,808	11
Non-Hispanic black, male													
2005 2/.....	22,608	2	1	-	25	160	862	3,172	4,826	5,101	5,503	2,952	4
2004 2/.....	22,720	2	-	2	36	169	848	3,107	4,615	5,410	5,441	3,084	6
2003 2/.....	23,919	3	3	3	26	159	938	3,178	4,624	5,784	6,094	3,106	1
2002.....	23,976	4	2	3	18	131	947	3,053	4,722	5,811	6,013	3,266	6
2001.....	24,255	3	1	3	35	167	974	2,982	4,596	6,089	6,115	3,283	7
2000.....	24,289	2	-	2	14	150	946	2,886	4,713	6,121	6,118	3,331	6
1999.....	24,736	4	-	5	24	148	1,042	2,904	4,737	6,437	5,991	3,435	9
Non-Hispanic black, female													
2005 2/.....	22,827	3	-	2	10	58	448	1,546	2,750	4,002	6,593	7,412	3
2004 2/.....	23,344	-	1	4	10	88	467	1,557	2,677	4,307	6,759	7,473	1
2003 2/.....	24,698	1	2	5	15	70	469	1,548	2,815	4,576	7,113	8,084	-
2002.....	25,546	3	-	2	13	79	494	1,612	2,782	4,787	7,528	8,243	3
2001.....	26,040	4	2	2	17	79	500	1,551	2,855	5,018	7,667	8,342	3
2000.....	26,370	2	-	-	10	79	505	1,532	2,874	5,364	7,691	8,310	3
1999.....	26,632	4	1	3	20	65	510	1,458	2,805	5,492	7,899	8,373	2

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Ischemic heart diseases (I20-I25)													
Origin not stated 3/, both sexes													
2005	906	-	-	-	-	1	11	81	149	173	264	224	3
2004	1,131	-	-	-	-	2	26	85	168	257	318	270	5
2003	1,451	-	-	-	1	4	31	96	226	287	427	372	7
2002	1,786	-	-	-	1	8	27	117	220	370	497	539	7
2001	1,683	1	-	-	1	3	31	96	224	306	520	487	14
2000	1,919	1	-	-	-	5	45	127	258	384	543	545	11
1999	2,023	-	-	-	1	1	41	116	248	429	606	573	8
Origin not stated 3/, male													
2005	536	-	-	-	-	1	9	64	110	123	156	72	1
2004	686	-	-	-	-	2	19	66	125	186	187	96	5
2003	855	-	-	-	1	3	22	74	174	195	240	140	6
2002	1,003	-	-	-	1	7	19	93	168	245	274	189	7
2001	932	1	-	-	1	3	25	78	179	209	288	140	8
2000	1,080	1	-	-	-	3	39	102	189	270	281	187	8
1999	1,140	-	-	-	-	1	37	91	197	300	309	199	6
Origin not stated 3/, female													
2005	370	-	-	-	-	-	2	17	39	50	108	152	2
2004	445	-	-	-	-	-	7	19	43	71	131	174	-
2003	596	-	-	-	-	1	9	22	52	92	187	232	1
2002	783	-	-	-	-	1	8	24	52	125	223	350	-
2001	751	-	-	-	-	-	6	18	45	97	232	347	6
2000	839	-	-	-	-	2	6	25	69	114	262	358	3
1999	883	-	-	-	1	-	4	25	51	129	297	374	2

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Acute myocardial infarction (I21-I22)													
All origins, both sexes													
2005	151,004	11	2	7	68	392	2,734	10,070	18,553	26,674	45,449	47,041	3
2004	156,816	9	2	5	51	447	2,838	10,084	18,758	28,327	48,037	48,252	6
2003	170,564	9	6	12	73	413	3,094	10,420	19,767	30,923	53,108	52,734	5
2002	179,514	11	5	7	62	400	3,161	10,619	19,933	34,003	56,590	54,718	5
2001	184,757	15	2	7	64	475	3,176	10,817	19,830	35,380	59,254	55,727	10
2000	192,898	20	1	6	52	375	3,132	10,877	21,013	38,459	62,210	56,745	8
1999	199,454	13	2	11	60	434	3,249	10,823	21,406	41,441	64,711	57,297	7
All origins, male													
2005	80,079	6	2	3	54	297	2,025	7,563	13,151	16,676	23,540	16,759	3
2004	82,909	7	-	4	39	322	2,113	7,632	13,342	17,658	24,893	16,895	4
2003	89,515	4	2	6	46	300	2,301	7,984	13,972	19,290	27,350	18,256	4
2002	93,830	7	2	2	41	292	2,347	7,995	13,966	21,264	28,919	18,991	4
2001	95,875	10	2	4	48	333	2,384	8,158	13,858	21,963	30,125	18,984	6
2000	100,306	10	1	4	30	269	2,387	8,265	14,638	23,823	31,631	19,243	5
1999	104,169	6	1	7	40	306	2,470	8,300	15,029	25,708	32,943	19,357	2
All origins, female													
2005	70,925	5	-	4	14	95	709	2,507	5,402	9,998	21,909	30,282	-
2004	73,907	2	2	1	12	125	725	2,452	5,416	10,669	23,144	31,357	2
2003	81,049	5	4	6	27	113	793	2,436	5,795	11,633	25,758	34,478	1
2002	85,684	4	3	5	21	108	814	2,624	5,967	12,739	27,671	35,727	1
2001	88,882	5	-	3	16	142	792	2,659	5,972	13,417	29,129	36,743	4
2000	92,592	10	-	2	22	106	745	2,612	6,375	14,636	30,579	37,502	3
1999	95,285	7	1	4	20	128	779	2,523	6,377	15,733	31,768	37,940	5

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Acute myocardial infarction (I21-I22)													
Hispanic, both sexes													
2005	7,529	1	-	2	11	36	175	535	1,028	1,577	2,324	1,840	-
2004	7,132	2	1	2	5	35	178	518	982	1,530	2,161	1,717	1
2003	7,370	-	-	2	11	37	175	508	929	1,683	2,188	1,837	-
2002	7,631	3	1	1	7	34	177	518	1,089	1,726	2,184	1,891	-
2001	7,403	4	1	1	6	46	139	503	914	1,701	2,190	1,896	2
2000	7,106	5	-	-	5	31	154	486	941	1,661	2,049	1,774	-
1999	7,158	2	-	-	4	32	142	528	919	1,699	2,082	1,750	-
Hispanic, male													
2005	4,145	-	-	2	7	25	127	408	707	950	1,216	703	-
2004	3,931	2	-	2	4	27	141	406	692	928	1,128	600	1
2003	3,903	-	-	1	10	27	127	394	635	1,016	1,067	626	-
2002	4,125	2	-	-	5	29	126	382	755	1,043	1,121	662	-
2001	3,972	3	1	-	6	37	110	383	603	1,036	1,093	699	1
2000	3,825	3	-	-	4	24	125	363	643	984	1,047	632	-
1999	3,836	-	-	-	3	27	116	391	607	1,028	1,072	592	-
Hispanic, female													
2005	3,384	1	-	-	4	11	48	127	321	627	1,108	1,137	-
2004	3,201	-	1	-	1	8	37	112	290	602	1,033	1,117	-
2003	3,467	-	-	1	1	10	48	114	294	667	1,121	1,211	-
2002	3,506	1	1	1	2	5	51	136	334	683	1,063	1,229	-
2001	3,431	1	-	1	-	9	29	120	311	665	1,097	1,197	1
2000	3,281	2	-	-	1	7	29	123	298	677	1,002	1,142	-
1999	3,322	2	-	-	1	5	26	137	312	671	1,010	1,158	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Acute myocardial infarction (I21-I22)													
Non-Hispanic 1/, both sexes													
2005	143,235	10	2	5	57	356	2,558	9,500	17,481	25,052	43,068	45,143	3
2004	149,368	7	1	3	46	411	2,653	9,543	17,730	26,726	45,786	46,457	5
2003	162,813	9	6	10	62	375	2,909	9,886	18,788	29,159	50,806	50,800	3
2002	171,441	8	4	6	55	365	2,975	10,076	18,801	32,172	54,285	52,689	5
2001	176,895	11	1	6	58	428	3,023	10,294	18,844	33,604	56,923	53,696	7
2000	185,283	14	1	6	47	344	2,963	10,363	20,000	36,695	60,020	54,823	7
1999	191,729	11	2	11	56	402	3,090	10,264	20,409	39,614	62,466	55,400	4
Non-Hispanic 1/, male													
2005	75,793	6	2	1	47	272	1,897	7,129	12,408	15,700	22,288	16,040	3
2004	78,788	5	-	2	35	294	1,965	7,207	12,617	16,675	23,716	16,269	3
2003	85,386	4	2	5	36	272	2,168	7,571	13,297	18,224	26,216	17,588	3
2002	89,480	5	2	2	36	262	2,215	7,597	13,183	20,161	27,736	18,277	4
2001	91,655	7	1	4	42	295	2,261	7,758	13,202	20,875	28,956	18,249	5
2000	96,218	6	1	4	26	245	2,250	7,886	13,950	22,772	30,516	18,558	4
1999	100,014	6	1	7	37	279	2,338	7,887	14,363	24,598	31,785	18,712	1
Non-Hispanic 1/, female													
2005	67,442	4	-	4	10	84	661	2,371	5,073	9,352	20,780	29,103	-
2004	70,580	2	1	1	11	117	688	2,336	5,113	10,051	22,070	30,188	2
2003	77,427	5	4	5	26	103	741	2,315	5,491	10,935	24,590	33,212	-
2002	81,961	3	2	4	19	103	760	2,479	5,618	12,011	26,549	34,412	1
2001	85,240	4	-	2	16	133	762	2,536	5,642	12,729	27,967	35,447	2
2000	89,065	8	-	2	21	99	713	2,477	6,050	13,923	29,504	36,265	3
1999	91,715	5	1	4	19	123	752	2,377	6,046	15,016	30,681	36,688	3

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Acute myocardial infarction (I21-I22)													
Non-Hispanic white, both sexes													
2005 2/.....	124,831	5	-	3	39	244	1,989	7,626	14,459	21,282	38,105	41,077	2
2004 2/.....	130,419	4	1	1	22	303	2,089	7,605	14,748	22,566	40,840	42,237	3
2003 2/.....	142,402	7	3	7	43	273	2,277	7,840	15,624	24,791	45,199	46,335	3
2002.....	150,051	6	4	4	36	267	2,345	8,094	15,624	27,467	48,317	47,883	4
2001.....	155,312	8	1	1	36	313	2,367	8,302	15,677	28,749	50,908	48,946	4
2000.....	163,203	10	1	4	34	243	2,336	8,410	16,678	31,568	53,840	50,073	6
1999.....	169,545	7	2	6	35	293	2,446	8,404	17,116	34,339	56,289	50,606	2
Non-Hispanic white, male													
2005 2/.....	66,629	4	-	1	35	190	1,522	5,894	10,504	13,662	20,062	14,753	2
2004 2/.....	69,443	3	-	1	17	229	1,603	5,916	10,768	14,388	21,556	14,960	2
2003 2/.....	75,157	3	1	3	26	201	1,748	6,186	11,352	15,828	23,607	16,199	3
2002.....	79,037	4	2	2	25	200	1,793	6,306	11,266	17,673	25,036	16,727	3
2001.....	81,175	6	1	1	28	219	1,834	6,491	11,303	18,302	26,282	16,704	4
2000.....	85,412	5	1	2	20	183	1,839	6,635	11,919	20,062	27,749	16,994	3
1999.....	89,142	3	1	4	26	202	1,907	6,671	12,319	21,791	29,104	17,114	-
Non-Hispanic white, female													
2005 2/.....	58,202	1	-	2	4	54	467	1,732	3,955	7,620	18,043	26,324	-
2004 2/.....	60,976	1	1	-	5	74	486	1,689	3,980	8,178	19,284	27,277	1
2003 2/.....	67,245	4	2	4	17	72	529	1,654	4,272	8,963	21,592	30,136	-
2002.....	71,014	2	2	2	11	67	552	1,788	4,358	9,794	23,281	31,156	1
2001.....	74,137	2	-	-	8	94	533	1,811	4,374	10,447	24,626	32,242	-
2000.....	77,791	5	-	2	14	60	497	1,775	4,759	11,506	26,091	33,079	3
1999.....	80,403	4	1	2	9	91	539	1,733	4,797	12,548	27,185	33,492	2

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Acute myocardial infarction (I21-I22)													
Non-Hispanic black, both sexes													
2005 2/	15,359	4	1	2	16	94	497	1,647	2,581	3,139	4,077	3,300	1
2004 2/	16,035	2	-	2	21	93	497	1,725	2,616	3,511	4,123	3,444	1
2003 2/	17,268	1	2	3	18	87	573	1,790	2,699	3,714	4,671	3,710	-
2002	18,162	2	-	2	15	88	556	1,756	2,800	3,993	4,939	4,010	1
2001	18,399	3	-	4	21	97	572	1,740	2,752	4,173	5,039	3,995	3
2000	18,934	2	-	2	11	93	560	1,720	2,910	4,415	5,189	4,031	1
1999	19,120	4	-	5	19	88	580	1,653	2,862	4,542	5,256	4,109	2
Non-Hispanic black, male													
2005 2/	7,437	2	1	-	12	67	314	1,055	1,591	1,669	1,781	944	1
2004 2/	7,721	2	-	1	17	53	309	1,135	1,582	1,894	1,724	1,004	-
2003 2/	8,381	-	1	2	10	59	371	1,172	1,624	1,986	2,103	1,053	-
2002	8,607	1	-	-	7	54	359	1,109	1,651	2,086	2,154	1,185	1
2001	8,628	1	-	2	13	62	354	1,068	1,606	2,163	2,155	1,203	1
2000	8,959	1	-	2	5	55	355	1,068	1,731	2,279	2,247	1,215	1
1999	9,106	3	-	3	10	61	374	1,055	1,743	2,375	2,211	1,270	1
Non-Hispanic black, female													
2005 2/	7,922	2	-	2	4	27	183	592	990	1,470	2,296	2,356	-
2004 2/	8,314	-	-	1	4	40	188	590	1,034	1,617	2,399	2,440	1
2003 2/	8,887	1	1	1	8	28	202	618	1,075	1,728	2,568	2,657	-
2002	9,555	1	-	2	8	34	197	647	1,149	1,907	2,785	2,825	-
2001	9,771	2	-	2	8	35	218	672	1,146	2,010	2,884	2,792	2
2000	9,975	1	-	-	6	38	205	652	1,179	2,136	2,942	2,816	-
1999	10,014	1	-	2	9	27	206	598	1,119	2,167	3,045	2,839	1

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Acute myocardial infarction (I21-I22)													
Origin not stated 3/, both sexes													
2005	240	-	-	-	-	-	1	35	44	45	57	58	-
2004	316	-	-	-	-	1	7	23	46	71	90	78	-
2003	381	-	-	-	-	1	10	26	50	81	114	97	2
2002	442	-	-	-	-	1	9	25	43	105	121	138	-
2001	459	-	-	-	-	1	14	20	72	75	141	135	1
2000	509	1	-	-	-	-	15	28	72	103	141	148	1
1999	567	-	-	-	-	-	17	31	78	128	163	147	3
Origin not stated 3/, male													
2005	141	-	-	-	-	-	1	26	36	26	36	16	-
2004	190	-	-	-	-	1	7	19	33	55	49	26	-
2003	226	-	-	-	-	1	6	19	40	50	67	42	1
2002	225	-	-	-	-	1	6	16	28	60	62	52	-
2001	248	-	-	-	-	1	13	17	53	52	76	36	-
2000	263	1	-	-	-	-	12	16	45	67	68	53	1
1999	319	-	-	-	-	-	16	22	59	82	86	53	1
Origin not stated 3/, female													
2005	99	-	-	-	-	-	-	9	8	19	21	42	-
2004	126	-	-	-	-	-	-	4	13	16	41	52	-
2003	155	-	-	-	-	-	4	7	10	31	47	55	1
2002	217	-	-	-	-	-	3	9	15	45	59	86	-
2001	211	-	-	-	-	-	1	3	19	23	65	99	1
2000	246	-	-	-	-	-	3	12	27	36	73	95	-
1999	248	-	-	-	-	-	1	9	19	46	77	94	2

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Other acute ischemic heart diseases (I24)													
All origins, both sexes													
2005	3,565	-	1	1	3	11	82	351	515	606	935	1,060	-
2004	2,462	1	2	1	5	10	80	217	359	399	670	717	1
2003	3,158	4	1	1	5	11	98	277	403	579	837	941	1
2002	3,407	2	-	1	4	18	77	345	460	631	898	971	-
2001	3,351	3	-	2	7	14	94	307	463	645	871	945	-
2000	3,363	1	1	-	8	16	92	282	484	644	906	928	1
1999	3,232	3	1	1	3	6	81	252	448	674	855	908	-
All origins, male													
2005	1,823	-	-	1	2	7	59	245	371	364	450	324	-
2004	1,301	-	-	-	4	6	59	151	251	254	329	246	1
2003	1,621	2	1	-	5	7	73	197	278	361	414	282	1
2002	1,767	1	-	1	1	12	50	250	320	390	444	298	-
2001	1,720	2	-	1	6	10	65	230	320	393	424	269	-
2000	1,749	1	1	-	6	13	71	217	341	398	446	254	1
1999	1,665	1	1	1	3	4	63	178	315	410	423	266	-
All origins, female													
2005	1,742	-	1	-	1	4	23	106	144	242	485	736	-
2004	1,161	1	2	1	1	4	21	66	108	145	341	471	-
2003	1,537	2	-	1	-	4	25	80	125	218	423	659	-
2002	1,640	1	-	-	3	6	27	95	140	241	454	673	-
2001	1,631	1	-	1	1	4	29	77	143	252	447	676	-
2000	1,614	-	-	-	2	3	21	65	143	246	460	674	-
1999	1,567	2	-	-	-	2	18	74	133	264	432	642	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Other acute ischemic heart diseases (I24)													
Hispanic, both sexes													
2005	90	-	-	-	1	1	2	11	8	19	23	25	-
2004	35	1	1	1	-	1	2	3	6	6	11	2	1
2003	47	-	-	-	1	1	1	5	6	10	12	10	1
2002	64	-	-	1	1	1	1	10	12	14	15	9	-
2001	63	-	-	-	-	-	1	4	12	19	14	13	-
2000	69	-	1	-	1	1	1	6	10	15	17	17	-
1999	53	-	-	1	-	-	1	7	13	8	12	11	-
Hispanic, male													
2005	46	-	-	-	1	1	2	6	6	9	15	6	-
2004	21	-	-	-	-	1	2	3	4	3	6	1	1
2003	21	-	-	-	1	1	1	2	3	5	4	3	1
2002	40	-	-	1	-	1	1	8	9	10	7	3	-
2001	42	-	-	-	-	-	1	2	9	14	7	9	-
2000	39	-	1	-	1	1	1	3	5	11	9	7	-
1999	26	-	-	1	-	-	1	4	6	4	8	2	-
Hispanic, female													
2005	44	-	-	-	-	-	-	5	2	10	8	19	-
2004	14	1	1	1	-	-	-	-	2	3	5	1	-
2003	26	-	-	-	-	-	-	3	3	5	8	7	-
2002	24	-	-	-	1	-	-	2	3	4	8	6	-
2001	21	-	-	-	-	-	-	2	3	5	7	4	-
2000	30	-	-	-	-	-	-	3	5	4	8	10	-
1999	27	-	-	-	-	-	-	3	7	4	4	9	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Other acute ischemic heart diseases (I24)													
Non-Hispanic 1/, both sexes													
2005	3,461	-	1	1	2	10	80	335	506	583	910	1,033	-
2004	2,419	-	1	-	5	9	78	211	352	392	658	713	-
2003	3,104	4	1	1	4	10	97	272	395	566	824	930	-
2002	3,339	2	-	-	2	17	75	335	447	617	883	961	-
2001	3,284	3	-	2	7	14	93	303	450	625	855	932	-
2000	3,282	1	-	-	7	15	90	273	473	626	887	909	1
1999	3,167	3	1	-	3	6	80	245	433	664	841	891	-
Non-Hispanic 1/, male													
2005	1,769	-	-	1	1	6	57	236	364	351	435	318	-
2004	1,275	-	-	-	4	5	57	146	246	250	323	244	-
2003	1,595	2	1	-	4	6	72	195	273	354	409	279	-
2002	1,725	1	-	-	-	11	49	242	311	380	437	294	-
2001	1,675	2	-	1	6	10	64	228	310	378	416	260	-
2000	1,705	1	-	-	5	12	69	212	336	386	436	247	1
1999	1,635	1	1	-	3	4	62	174	307	406	414	263	-
Non-Hispanic 1/, female													
2005	1,692	-	1	-	1	4	23	99	142	232	475	715	-
2004	1,144	-	1	-	1	4	21	65	106	142	335	469	-
2003	1,509	2	-	1	-	4	25	77	122	212	415	651	-
2002	1,614	1	-	-	2	6	26	93	136	237	446	667	-
2001	1,609	1	-	1	1	4	29	75	140	247	439	672	-
2000	1,577	-	-	-	2	3	21	61	137	240	451	662	-
1999	1,532	2	-	-	-	2	18	71	126	258	427	628	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Other acute ischemic heart diseases (I24)													
Non-Hispanic white, both sexes													
2005 2/	2,829	-	1	1	2	6	58	225	375	471	779	911	-
2004 2/	1,988	-	1	-	3	2	58	140	262	311	559	652	-
2003 2/	2,613	2	-	1	3	5	68	200	304	463	716	851	-
2002	2,843	1	-	-	1	11	54	251	356	510	782	877	-
2001	2,756	3	-	2	4	11	62	222	342	511	757	842	-
2000	2,767	1	-	-	5	11	66	207	357	514	774	832	-
1999	2,689	1	1	-	3	5	52	169	344	553	736	825	-
Non-Hispanic white, male													
2005 2/	1,431	-	-	1	1	3	43	176	273	284	372	278	-
2004 2/	1,019	-	-	-	2	2	44	99	184	191	279	218	-
2003 2/	1,314	1	-	-	3	4	55	146	209	296	346	254	-
2002	1,454	1	-	-	-	7	41	179	251	313	390	272	-
2001	1,386	2	-	1	3	9	45	170	240	307	374	235	-
2000	1,412	1	-	-	3	8	52	165	261	314	382	226	-
1999	1,361	1	1	-	3	4	40	123	240	339	366	244	-
Non-Hispanic white, female													
2005 2/	1,398	-	1	-	1	3	15	49	102	187	407	633	-
2004 2/	969	-	1	-	1	-	14	41	78	120	280	434	-
2003 2/	1,299	1	-	1	-	1	13	54	95	167	370	597	-
2002	1,389	-	-	-	1	4	13	72	105	197	392	605	-
2001	1,370	1	-	1	1	2	17	52	102	204	383	607	-
2000	1,355	-	-	-	2	3	14	42	96	200	392	606	-
1999	1,328	-	-	-	-	1	12	46	104	214	370	581	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Other acute ischemic heart diseases (I24)													
Non-Hispanic black, both sexes													
2005 2/.....	551	-	-	-	-	4	19	97	115	98	116	102	-
2004 2/.....	389	-	-	-	1	7	18	66	85	70	88	54	-
2003 2/.....	432	1	1	-	1	3	27	62	81	91	97	68	-
2002.....	443	1	-	-	1	6	19	78	78	100	89	71	-
2001.....	453	-	-	-	3	2	29	69	98	95	79	78	-
2000.....	449	-	-	-	1	4	22	61	103	91	97	69	1
1999.....	429	2	-	-	-	1	26	73	83	93	97	54	-
Non-Hispanic black, male													
2005 2/.....	292	-	-	-	-	3	12	54	79	58	56	30	-
2004 2/.....	232	-	-	-	1	3	12	42	59	50	42	23	-
2003 2/.....	254	1	1	-	1	1	16	43	58	53	57	23	-
2002.....	242	-	-	-	-	4	6	58	50	64	40	20	-
2001.....	249	-	-	-	3	-	19	50	62	59	35	21	-
2000.....	255	-	-	-	1	4	16	44	67	58	46	18	1
1999.....	246	-	-	-	-	-	20	49	63	56	41	17	-
Non-Hispanic black, female													
2005 2/.....	259	-	-	-	-	1	7	43	36	40	60	72	-
2004 2/.....	157	-	-	-	-	4	6	24	26	20	46	31	-
2003 2/.....	178	-	-	-	-	2	11	19	23	38	40	45	-
2002.....	201	1	-	-	1	2	13	20	28	36	49	51	-
2001.....	204	-	-	-	-	2	10	19	36	36	44	57	-
2000.....	194	-	-	-	-	-	6	17	36	33	51	51	-
1999.....	183	2	-	-	-	1	6	24	20	37	56	37	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Other acute ischemic heart diseases (I24)													
Origin not stated 3/, both sexes													
2005	14	-	-	-	-	-	-	5	1	4	2	2	-
2004	8	-	-	-	-	-	-	3	1	1	1	2	-
2003	7	-	-	-	-	-	-	-	2	3	1	1	-
2002	4	-	-	-	1	-	1	-	1	-	-	1	-
2001	4	-	-	-	-	-	-	-	1	1	2	-	-
2000	12	-	-	-	-	-	1	3	1	3	2	2	-
1999	12	-	-	-	-	-	-	-	2	2	2	6	-
Origin not stated 3/, male													
2005	8	-	-	-	-	-	-	3	1	4	-	-	-
2004	5	-	-	-	-	-	-	2	1	1	-	1	-
2003	5	-	-	-	-	-	-	-	2	2	1	-	-
2002	2	-	-	-	1	-	-	-	-	-	-	1	-
2001	3	-	-	-	-	-	-	-	1	1	1	-	-
2000	5	-	-	-	-	-	1	2	-	1	1	-	-
1999	4	-	-	-	-	-	-	-	2	-	1	1	-
Origin not stated 3/, female													
2005	6	-	-	-	-	-	-	2	-	-	2	2	-
2004	3	-	-	-	-	-	-	1	-	-	1	1	-
2003	2	-	-	-	-	-	-	-	-	1	-	1	-
2002	2	-	-	-	-	-	1	-	1	-	-	-	-
2001	1	-	-	-	-	-	-	-	-	-	1	-	-
2000	7	-	-	-	-	-	-	1	1	2	1	2	-
1999	8	-	-	-	-	-	-	-	-	2	1	5	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Other forms of chronic ischemic heart disease (I20,I25)													
All origins, both sexes													
2005	291,118	2	4	6	80	611	4,044	14,889	27,731	42,841	88,051	112,834	25
2004	292,048	3	6	11	69	635	4,214	14,622	26,885	44,376	90,421	110,787	19
2003	306,306	9	5	11	82	658	4,355	14,664	27,207	47,378	95,425	116,489	23
2002	311,461	10	2	12	67	578	4,445	14,611	26,869	49,233	97,903	117,699	32
2001	314,081	8	7	11	92	627	4,264	13,949	26,281	51,438	99,729	117,637	38
2000	318,943	9	8	10	63	544	4,295	13,731	26,240	53,782	100,576	119,646	39
1999	326,973	11	2	10	74	574	4,421	13,700	26,523	56,725	103,730	121,171	32
All origins, male													
2005	150,213	1	2	5	65	470	3,104	11,425	20,026	27,457	47,362	40,281	15
2004	149,328	1	1	5	53	494	3,194	11,203	19,275	28,361	48,369	38,354	18
2003	155,206	7	3	4	56	506	3,305	11,300	19,517	30,348	50,809	39,334	17
2002	157,163	7	2	10	50	437	3,443	11,306	19,370	31,482	51,876	39,157	23
2001	156,410	5	1	8	65	487	3,263	10,753	18,828	32,716	52,404	37,853	27
2000	158,519	6	5	5	45	422	3,329	10,667	18,816	34,333	52,670	38,193	28
1999	161,434	6	1	9	50	429	3,437	10,583	18,779	36,069	53,939	38,103	29
All origins, female													
2005	140,905	1	2	1	15	141	940	3,464	7,705	15,384	40,689	72,553	10
2004	142,720	2	5	6	16	141	1,020	3,419	7,610	16,015	42,052	72,433	1
2003	151,100	2	2	7	26	152	1,050	3,364	7,690	17,030	44,616	77,155	6
2002	154,298	3	-	2	17	141	1,002	3,305	7,499	17,751	46,027	78,542	9
2001	157,671	3	6	3	27	140	1,001	3,196	7,453	18,722	47,325	79,784	11
2000	160,424	3	3	5	18	122	966	3,064	7,424	19,449	47,906	81,453	11
1999	165,539	5	1	1	24	145	984	3,117	7,744	20,656	49,791	83,068	3

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Other forms of chronic ischemic heart disease (I20,I25)													
Hispanic, both sexes													
2005	14,155	-	2	2	10	62	326	980	1,709	2,721	4,294	4,045	4
2004	13,315	1	-	1	10	59	297	903	1,621	2,611	4,036	3,775	1
2003	13,366	2	1	2	10	66	270	839	1,611	2,644	4,032	3,884	5
2002	13,246	1	-	1	10	53	264	861	1,582	2,770	3,861	3,842	1
2001	13,198	-	-	3	12	63	228	807	1,478	2,833	3,891	3,881	2
2000	12,569	1	-	1	6	37	237	748	1,506	2,688	3,560	3,781	4
1999	12,725	-	-	1	7	55	223	793	1,531	2,785	3,561	3,764	5
Hispanic, male													
2005	7,762	-	2	2	8	54	277	775	1,231	1,696	2,258	1,455	4
2004	7,293	-	-	1	10	53	244	707	1,156	1,646	2,152	1,323	1
2003	7,130	1	1	2	9	55	213	670	1,142	1,659	2,097	1,277	4
2002	7,130	-	-	1	7	46	219	668	1,143	1,722	1,971	1,352	1
2001	7,005	-	-	1	7	53	184	631	1,055	1,754	2,016	1,302	2
2000	6,698	1	-	1	4	31	196	607	1,078	1,666	1,832	1,279	3
1999	6,742	-	-	1	5	45	193	647	1,064	1,738	1,789	1,255	5
Hispanic, female													
2005	6,393	-	-	-	2	8	49	205	478	1,025	2,036	2,590	-
2004	6,022	1	-	-	-	6	53	196	465	965	1,884	2,452	-
2003	6,236	1	-	-	1	11	57	169	469	985	1,935	2,607	1
2002	6,116	1	-	-	3	7	45	193	439	1,048	1,890	2,490	-
2001	6,193	-	-	2	5	10	44	176	423	1,079	1,875	2,579	-
2000	5,871	-	-	-	2	6	41	141	428	1,022	1,728	2,502	1
1999	5,983	-	-	-	2	10	30	146	467	1,047	1,772	2,509	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Other forms of chronic ischemic heart disease (I20,I25)													
Non-Hispanic 1/, both sexes													
2005	276,311	2	2	4	70	548	3,708	13,868	25,918	39,996	83,552	108,625	18
2004	277,926	2	6	10	59	575	3,898	13,660	25,143	41,580	86,158	106,822	13
2003	291,877	7	4	9	71	589	4,064	13,755	25,422	44,531	91,081	112,331	13
2002	296,875	9	2	11	57	518	4,164	13,658	25,111	46,198	93,666	113,457	24
2001	299,663	7	7	8	79	562	4,019	13,066	24,652	48,375	95,461	113,404	23
2000	304,976	8	8	9	57	502	4,029	12,887	24,549	50,816	96,616	115,470	25
1999	312,804	11	2	9	66	518	4,174	12,822	24,824	53,641	99,728	116,987	22
Non-Hispanic 1/, male													
2005	142,064	1	-	3	57	415	2,819	10,615	18,722	25,668	44,984	38,770	10
2004	141,544	1	1	4	43	440	2,938	10,451	18,028	26,585	46,079	36,962	12
2003	147,452	6	2	2	46	449	3,076	10,575	18,243	28,546	48,540	37,959	8
2002	149,257	7	2	9	43	385	3,211	10,561	18,087	29,575	49,693	37,669	15
2001	148,724	4	1	7	57	432	3,067	10,061	17,648	30,806	50,177	36,447	17
2000	151,009	5	5	4	41	388	3,107	9,976	17,594	32,465	50,626	36,780	18
1999	153,875	6	1	8	45	383	3,223	9,867	17,579	34,113	51,928	36,703	19
Non-Hispanic 1/, female													
2005	134,247	1	2	1	13	133	889	3,253	7,196	14,328	38,568	69,855	8
2004	136,382	1	5	6	16	135	960	3,209	7,115	14,995	40,079	69,860	1
2003	144,425	1	2	7	25	140	988	3,180	7,179	15,985	42,541	74,372	5
2002	147,618	2	-	2	14	133	953	3,097	7,024	16,623	43,973	75,788	9
2001	150,939	3	6	1	22	130	952	3,005	7,004	17,569	45,284	76,957	6
2000	153,967	3	3	5	16	114	922	2,911	6,955	18,351	45,990	78,690	7
1999	158,929	5	1	1	21	135	951	2,955	7,245	19,528	47,800	80,284	3

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Other forms of chronic ischemic heart disease (I20,I25)													
Non-Hispanic white, both sexes													
2005 2/	240,845	1	2	4	44	391	2,779	10,456	20,319	32,989	73,933	99,916	11
2004 2/	242,493	2	3	6	32	385	2,981	10,395	19,867	34,389	76,417	98,009	7
2003 2/	255,086	5	2	3	45	417	3,089	10,468	20,081	36,856	80,902	103,208	10
2002	260,336	4	-	8	39	372	3,166	10,454	19,819	38,576	83,449	104,433	16
2001	262,891	3	4	7	46	394	3,027	9,994	19,402	40,421	85,150	104,429	14
2000	268,535	5	8	8	41	352	3,035	9,914	19,332	42,785	86,497	106,540	18
1999	275,726	8	1	6	38	367	3,109	9,852	19,606	45,237	89,569	107,920	13
Non-Hispanic white, male													
2005 2/	123,802	1	-	3	38	292	2,170	8,203	15,034	21,579	40,415	36,061	6
2004 2/	123,514	1	1	3	24	296	2,309	8,203	14,553	22,483	41,478	34,157	6
2003 2/	128,839	4	1	1	28	322	2,391	8,278	14,802	24,092	43,699	35,216	5
2002	130,900	3	-	6	29	286	2,512	8,371	14,591	25,219	44,944	34,929	10
2001	130,367	2	-	6	34	311	2,371	7,929	14,233	26,272	45,380	33,819	10
2000	132,908	3	5	4	29	282	2,428	7,923	14,218	27,998	45,911	34,093	14
1999	135,406	4	1	6	28	276	2,469	7,791	14,195	29,423	47,293	33,909	11
Non-Hispanic white, female													
2005 2/	117,043	-	2	1	6	99	609	2,253	5,285	11,410	33,518	63,855	5
2004 2/	118,979	1	2	3	8	89	672	2,192	5,314	11,906	34,939	63,852	1
2003 2/	126,247	1	1	2	17	95	698	2,190	5,279	12,764	37,203	67,992	5
2002	129,436	1	-	2	10	86	654	2,083	5,228	13,357	38,505	69,504	6
2001	132,524	1	4	1	12	83	656	2,065	5,169	14,149	39,770	70,610	4
2000	135,627	2	3	4	12	70	607	1,991	5,114	14,787	40,586	72,447	4
1999	140,320	4	-	-	10	91	640	2,061	5,411	15,814	42,276	74,011	2

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Other forms of chronic ischemic heart disease (I20,I25)													
Non-Hispanic black, both sexes													
2005 2/	29,525	1	-	-	19	120	794	2,974	4,880	5,866	7,903	6,962	6
2004 2/	29,640	-	1	4	24	157	800	2,873	4,591	6,136	7,989	7,059	6
2003 2/	30,917	2	2	5	22	139	807	2,874	4,659	6,555	8,439	7,412	1
2002	30,917	4	2	3	15	116	866	2,831	4,626	6,505	8,513	7,428	8
2001	31,443	4	3	1	28	147	873	2,724	4,601	6,839	8,664	7,552	7
2000	31,276	2	-	-	12	132	869	2,637	4,574	6,979	8,523	7,541	7
1999	31,819	2	1	3	25	124	946	2,636	4,597	7,294	8,537	7,645	9
Non-Hispanic black, male													
2005 2/	14,879	-	-	-	13	90	536	2,063	3,156	3,374	3,666	1,978	3
2004 2/	14,767	-	-	1	18	113	527	1,930	2,974	3,466	3,675	2,057	6
2003 2/	15,284	2	1	1	15	99	551	1,963	2,942	3,745	3,934	2,030	1
2002	15,127	3	2	3	11	73	582	1,886	3,021	3,661	3,819	2,061	5
2001	15,378	2	1	1	19	105	601	1,864	2,928	3,867	3,925	2,059	6
2000	15,075	1	-	-	8	91	575	1,774	2,915	3,784	3,825	2,098	4
1999	15,384	1	-	2	14	87	648	1,800	2,931	4,006	3,739	2,148	8
Non-Hispanic black, female													
2005 2/	14,646	1	-	-	6	30	258	911	1,724	2,492	4,237	4,984	3
2004 2/	14,873	-	1	3	6	44	273	943	1,617	2,670	4,314	5,002	-
2003 2/	15,633	-	1	4	7	40	256	911	1,717	2,810	4,505	5,382	-
2002	15,790	1	-	-	4	43	284	945	1,605	2,844	4,694	5,367	3
2001	16,065	2	2	-	9	42	272	860	1,673	2,972	4,739	5,493	1
2000	16,201	1	-	-	4	41	294	863	1,659	3,195	4,698	5,443	3
1999	16,435	1	1	1	11	37	298	836	1,666	3,288	4,798	5,497	1

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Other forms of chronic ischemic heart disease (I20,I25)													
Origin not stated 3/, both sexes													
2005	652	-	-	-	-	1	10	41	104	124	205	164	3
2004	807	-	-	-	-	1	19	59	121	185	227	190	5
2003	1,063	-	-	-	1	3	21	70	174	203	312	274	5
2002	1,340	-	-	-	-	7	17	92	176	265	376	400	7
2001	1,220	1	-	-	1	2	17	76	151	230	377	352	13
2000	1,398	-	-	-	-	5	29	96	185	278	400	395	10
1999	1,444	-	-	-	1	1	24	85	168	299	441	420	5
Origin not stated 3/, male													
2005	387	-	-	-	-	1	8	35	73	93	120	56	1
2004	491	-	-	-	-	1	12	45	91	130	138	69	5
2003	624	-	-	-	1	2	16	55	132	143	172	98	5
2002	776	-	-	-	-	6	13	77	140	185	212	136	7
2001	681	1	-	-	1	2	12	61	125	156	211	104	8
2000	812	-	-	-	-	3	26	84	144	202	212	134	7
1999	817	-	-	-	-	1	21	69	136	218	222	145	5
Origin not stated 3/, female													
2005	265	-	-	-	-	-	2	6	31	31	85	108	2
2004	316	-	-	-	-	-	7	14	30	55	89	121	-
2003	439	-	-	-	-	1	5	15	42	60	140	176	-
2002	564	-	-	-	-	1	4	15	36	80	164	264	-
2001	539	-	-	-	-	-	5	15	26	74	166	248	5
2000	586	-	-	-	-	2	3	12	41	76	188	261	3
1999	627	-	-	-	1	-	3	16	32	81	219	275	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Atherosclerotic cardiovascular disease, so described (I25.0)													
All origins, both sexes													
2005	62,799	1	-	-	16	182	1,420	5,884	9,565	10,246	16,286	19,182	17
2004	63,661	-	1	-	9	193	1,463	5,665	9,297	10,830	16,730	19,459	14
2003	67,232	2	-	-	16	211	1,488	5,684	9,110	11,494	18,339	20,874	14
2002	68,129	2	-	1	17	181	1,515	5,366	8,793	11,817	18,618	21,796	23
2001	68,103	-	-	-	20	179	1,457	5,047	8,438	12,005	19,196	21,736	25
2000	69,554	1	-	1	15	144	1,376	4,980	8,242	12,567	19,545	22,664	19
1999	71,926	-	1	1	13	160	1,488	4,737	8,383	13,395	20,321	23,403	24
All origins, male													
2005	34,453	-	-	-	15	146	1,120	4,643	7,160	6,709	8,384	6,266	10
2004	34,574	-	-	-	8	157	1,143	4,495	6,864	7,046	8,606	6,242	13
2003	36,042	2	-	-	12	181	1,157	4,552	6,694	7,422	9,333	6,677	12
2002	36,047	1	-	-	13	144	1,216	4,239	6,549	7,617	9,416	6,835	17
2001	35,440	-	-	-	16	133	1,130	4,037	6,245	7,668	9,671	6,521	19
2000	35,896	-	-	1	12	115	1,099	3,977	6,125	8,069	9,725	6,758	15
1999	36,633	-	-	1	10	119	1,192	3,752	6,123	8,546	9,963	6,905	22
All origins, female													
2005	28,346	1	-	-	1	36	300	1,241	2,405	3,537	7,902	12,916	7
2004	29,087	-	1	-	1	36	320	1,170	2,433	3,784	8,124	13,217	1
2003	31,190	-	-	-	4	30	331	1,132	2,416	4,072	9,006	14,197	2
2002	32,082	1	-	1	4	37	299	1,127	2,244	4,200	9,202	14,961	6
2001	32,663	-	-	-	4	46	327	1,010	2,193	4,337	9,525	15,215	6
2000	33,658	1	-	-	3	29	277	1,003	2,117	4,498	9,820	15,906	4
1999	35,293	-	1	-	3	41	296	985	2,260	4,849	10,358	16,498	2

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Atherosclerotic cardiovascular disease, so described (I25.0)													
Hispanic, both sexes													
2005	3,491	-	-	-	4	18	111	464	691	729	830	642	2
2004	3,277	-	-	-	3	17	102	392	652	709	814	588	-
2003	3,295	-	-	-	3	21	93	374	614	700	798	689	3
2002	3,166	1	-	-	4	26	108	327	586	722	744	647	1
2001	3,150	-	-	-	1	25	81	343	548	748	735	667	2
2000	2,953	-	-	-	2	7	83	299	522	707	649	682	2
1999	3,063	-	-	-	1	14	73	308	540	747	708	669	3
Hispanic, male													
2005	2,246	-	-	-	3	16	93	385	543	489	467	248	2
2004	2,098	-	-	-	3	16	88	336	498	471	474	212	-
2003	2,094	-	-	-	3	20	80	327	482	480	442	258	2
2002	1,946	-	-	-	3	24	89	270	461	475	375	248	1
2001	1,928	-	-	-	-	21	63	288	416	502	405	231	2
2000	1,794	-	-	-	2	6	71	257	406	438	368	245	1
1999	1,847	-	-	-	1	12	63	260	384	527	372	225	3
Hispanic, female													
2005	1,245	-	-	-	1	2	18	79	148	240	363	394	-
2004	1,179	-	-	-	-	1	14	56	154	238	340	376	-
2003	1,201	-	-	-	-	1	13	47	132	220	356	431	1
2002	1,220	1	-	-	1	2	19	57	125	247	369	399	-
2001	1,222	-	-	-	1	4	18	55	132	246	330	436	-
2000	1,159	-	-	-	-	1	12	42	116	269	281	437	1
1999	1,216	-	-	-	-	2	10	48	156	220	336	444	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Atherosclerotic cardiovascular disease, so described (I25.0)													
Non-Hispanic 1/, both sexes													
2005	59,053	1	-	-	12	163	1,306	5,401	8,815	9,450	15,386	18,507	12
2004	60,067	-	1	-	6	176	1,354	5,248	8,587	10,032	15,836	18,817	10
2003	63,550	2	-	-	13	189	1,385	5,269	8,403	10,715	17,436	20,132	6
2002	64,528	1	-	1	13	152	1,403	4,987	8,109	10,992	17,771	21,082	17
2001	64,534	-	-	-	19	154	1,366	4,664	7,816	11,159	18,346	20,999	11
2000	66,141	1	-	1	13	137	1,283	4,631	7,627	11,756	18,769	21,912	11
1999	68,373	-	1	1	12	146	1,404	4,387	7,754	12,524	19,465	22,661	18
Non-Hispanic 1/, male													
2005	32,029	-	-	-	12	129	1,024	4,241	6,570	6,167	7,876	6,003	7
2004	32,271	-	-	-	5	141	1,050	4,140	6,323	6,511	8,083	6,009	9
2003	33,677	2	-	-	9	160	1,067	4,193	6,136	6,883	8,828	6,394	5
2002	33,809	1	-	-	10	118	1,124	3,927	6,004	7,076	8,977	6,561	11
2001	33,247	-	-	-	16	112	1,059	3,717	5,770	7,093	9,200	6,270	10
2000	33,786	-	-	1	10	109	1,019	3,674	5,646	7,549	9,281	6,487	10
1999	34,451	-	-	1	9	107	1,118	3,457	5,663	7,922	9,504	6,654	16
Non-Hispanic 1/, female													
2005	27,024	1	-	-	-	34	282	1,160	2,245	3,283	7,510	12,504	5
2004	27,796	-	1	-	1	35	304	1,108	2,264	3,521	7,753	12,808	1
2003	29,873	-	-	-	4	29	318	1,076	2,267	3,832	8,608	13,738	1
2002	30,719	-	-	1	3	34	279	1,060	2,105	3,916	8,794	14,521	6
2001	31,287	-	-	-	3	42	307	947	2,046	4,066	9,146	14,729	1
2000	32,355	1	-	-	3	28	264	957	1,981	4,207	9,488	15,425	1
1999	33,922	-	1	-	3	39	286	930	2,091	4,602	9,961	16,007	2

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Atherosclerotic cardiovascular disease, so described (I25.0)													
Non-Hispanic white, both sexes													
2005 2/	48,146	-	-	-	10	106	961	3,948	6,498	7,273	12,868	16,475	7
2004 2/	49,197	-	-	-	4	115	1,036	3,856	6,443	7,739	13,280	16,720	4
2003 2/	52,268	1	-	-	9	136	1,047	3,870	6,273	8,237	14,706	17,985	4
2002	53,161	1	-	1	11	105	1,020	3,703	6,043	8,444	14,972	18,851	10
2001	53,031	-	-	-	11	108	999	3,421	5,649	8,504	15,521	18,811	7
2000	54,544	-	-	1	11	95	944	3,407	5,537	9,079	15,871	19,591	8
1999	56,375	-	-	1	5	96	1,017	3,155	5,626	9,665	16,517	20,283	10
Non-Hispanic white, male													
2005 2/	25,849	-	-	-	10	81	782	3,191	4,948	4,791	6,621	5,422	3
2004 2/	26,188	-	-	-	3	91	826	3,166	4,804	5,133	6,825	5,337	3
2003 2/	27,329	1	-	-	6	114	814	3,161	4,680	5,338	7,490	5,722	3
2002	27,493	1	-	-	8	84	844	3,008	4,559	5,526	7,583	5,874	6
2001	26,960	-	-	-	8	82	794	2,809	4,291	5,496	7,849	5,625	6
2000	27,461	-	-	1	8	77	775	2,786	4,210	5,949	7,864	5,783	8
1999	27,943	-	-	1	5	73	838	2,558	4,208	6,227	8,097	5,928	8
Non-Hispanic white, female													
2005 2/	22,297	-	-	-	-	25	179	757	1,550	2,482	6,247	11,053	4
2004 2/	23,009	-	-	-	1	24	210	690	1,639	2,606	6,455	11,383	1
2003 2/	24,939	-	-	-	3	22	233	709	1,593	2,899	7,216	12,263	1
2002	25,668	-	-	1	3	21	176	695	1,484	2,918	7,389	12,977	4
2001	26,071	-	-	-	3	26	205	612	1,358	3,008	7,672	13,186	1
2000	27,083	-	-	-	3	18	169	621	1,327	3,130	8,007	13,808	-
1999	28,432	-	-	-	-	23	179	597	1,418	3,438	8,420	14,355	2

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Atherosclerotic cardiovascular disease, so described (I25.0)													
Non-Hispanic black, both sexes													
2005 2/	9,442	1	-	-	1	41	296	1,278	2,036	1,893	2,152	1,740	4
2004 2/	9,497	-	1	-	2	49	271	1,244	1,893	2,036	2,226	1,769	6
2003 2/	9,764	1	-	-	3	46	281	1,213	1,880	2,157	2,344	1,838	1
2002	10,024	-	-	-	2	39	333	1,156	1,844	2,258	2,466	1,919	7
2001	10,156	-	-	-	7	38	324	1,115	1,927	2,361	2,472	1,909	3
2000	10,283	1	-	-	1	36	292	1,120	1,877	2,380	2,522	2,051	3
1999	10,643	-	1	-	7	41	358	1,105	1,904	2,553	2,593	2,073	8
Non-Hispanic black, male													
2005 2/	5,281	-	-	-	1	33	202	901	1,404	1,203	1,059	475	3
2004 2/	5,231	-	-	-	2	39	182	848	1,324	1,215	1,083	532	6
2003 2/	5,431	1	-	-	2	39	203	881	1,267	1,327	1,158	552	1
2002	5,483	-	-	-	2	27	233	806	1,278	1,357	1,223	552	5
2001	5,476	-	-	-	7	24	230	799	1,299	1,412	1,174	528	3
2000	5,528	-	-	-	1	26	207	802	1,282	1,398	1,204	606	2
1999	5,692	-	-	-	4	27	254	792	1,278	1,518	1,210	601	8
Non-Hispanic black, female													
2005 2/	4,161	1	-	-	-	8	94	377	632	690	1,093	1,265	1
2004 2/	4,266	-	1	-	-	10	89	396	569	821	1,143	1,237	-
2003 2/	4,333	-	-	-	1	7	78	332	613	830	1,186	1,286	-
2002	4,541	-	-	-	-	12	100	350	566	901	1,243	1,367	2
2001	4,680	-	-	-	-	14	94	316	628	949	1,298	1,381	-
2000	4,755	1	-	-	-	10	85	318	595	982	1,318	1,445	1
1999	4,951	-	1	-	3	14	104	313	626	1,035	1,383	1,472	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Atherosclerotic cardiovascular disease, so described (I25.0)													
Origin not stated 3/, both sexes													
2005	255	-	-	-	-	1	3	19	59	67	70	33	3
2004	317	-	-	-	-	-	7	25	58	89	80	54	4
2003	387	-	-	-	-	1	10	41	93	79	105	53	5
2002	435	-	-	-	-	3	4	52	98	103	103	67	5
2001	419	-	-	-	-	-	10	40	74	98	115	70	12
2000	460	-	-	-	-	-	10	50	93	104	127	70	6
1999	490	-	-	-	-	-	11	42	89	124	148	73	3
Origin not stated 3/, male													
2005	178	-	-	-	-	1	3	17	47	53	41	15	1
2004	205	-	-	-	-	-	5	19	43	64	49	21	4
2003	271	-	-	-	-	1	10	32	76	59	63	25	5
2002	292	-	-	-	-	2	3	42	84	66	64	26	5
2001	265	-	-	-	-	-	8	32	59	73	66	20	7
2000	316	-	-	-	-	-	9	46	73	82	76	26	4
1999	335	-	-	-	-	-	11	35	76	97	87	26	3
Origin not stated 3/, female													
2005	77	-	-	-	-	-	-	2	12	14	29	18	2
2004	112	-	-	-	-	-	2	6	15	25	31	33	-
2003	116	-	-	-	-	-	-	9	17	20	42	28	-
2002	143	-	-	-	-	1	1	10	14	37	39	41	-
2001	154	-	-	-	-	-	2	8	15	25	49	50	5
2000	144	-	-	-	-	-	1	4	20	22	51	44	2
1999	155	-	-	-	-	-	-	7	13	27	61	47	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
All other forms of chronic ischemic heart disease (I20,I25.1-I25.9)													
All origins, both sexes													
2005	228,319	1	4	6	64	429	2,624	9,005	18,166	32,595	71,765	93,652	8
2004	228,387	3	5	11	60	442	2,751	8,957	17,588	33,546	73,691	91,328	5
2003	239,074	7	5	11	66	447	2,867	8,980	18,097	35,884	77,086	95,615	9
2002	243,332	8	2	11	50	397	2,930	9,245	18,076	37,416	79,285	95,903	9
2001	245,978	8	7	11	72	448	2,807	8,902	17,843	39,433	80,533	95,901	13
2000	249,389	8	8	9	48	400	2,919	8,751	17,998	41,215	81,031	96,982	20
1999	255,047	11	1	9	61	414	2,933	8,963	18,140	43,330	83,409	97,768	8
All origins, male													
2005	115,760	1	2	5	50	324	1,984	6,782	12,866	20,748	38,978	34,015	5
2004	114,754	1	1	5	45	337	2,051	6,708	12,411	21,315	39,763	32,112	5
2003	119,164	5	3	4	44	325	2,148	6,748	12,823	22,926	41,476	32,657	5
2002	121,116	6	2	10	37	293	2,227	7,067	12,821	23,865	42,460	32,322	6
2001	120,970	5	1	8	49	354	2,133	6,716	12,583	25,048	42,733	31,332	8
2000	122,623	6	5	4	33	307	2,230	6,690	12,691	26,264	42,945	31,435	13
1999	124,801	6	1	8	40	310	2,245	6,831	12,656	27,523	43,976	31,198	7
All origins, female													
2005	112,559	-	2	1	14	105	640	2,223	5,300	11,847	32,787	59,637	3
2004	113,633	2	4	6	15	105	700	2,249	5,177	12,231	33,928	59,216	-
2003	119,910	2	2	7	22	122	719	2,232	5,274	12,958	35,610	62,958	4
2002	122,216	2	-	1	13	104	703	2,178	5,255	13,551	36,825	63,581	3
2001	125,008	3	6	3	23	94	674	2,186	5,260	14,385	37,800	64,569	5
2000	126,766	2	3	5	15	93	689	2,061	5,307	14,951	38,086	65,547	7
1999	130,246	5	-	1	21	104	688	2,132	5,484	15,807	39,433	66,570	1

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
All other forms of chronic ischemic heart disease (I20,I25.1-I25.9)													
Hispanic, both sexes													
2005	10,664	-	2	2	6	44	215	516	1,018	1,992	3,464	3,403	2
2004	10,038	1	-	1	7	42	195	511	969	1,902	3,222	3,187	1
2003	10,071	2	1	2	7	45	177	465	997	1,944	3,234	3,195	2
2002	10,080	-	-	1	6	27	156	534	996	2,048	3,117	3,195	-
2001	10,048	-	-	3	11	38	147	464	930	2,085	3,156	3,214	-
2000	9,616	1	-	1	4	30	154	449	984	1,981	2,911	3,099	2
1999	9,662	-	-	1	6	41	150	485	991	2,038	2,853	3,095	2
Hispanic, male													
2005	5,516	-	2	2	5	38	184	390	688	1,207	1,791	1,207	2
2004	5,195	-	-	1	7	37	156	371	658	1,175	1,678	1,111	1
2003	5,036	1	1	2	6	35	133	343	660	1,179	1,655	1,019	2
2002	5,184	-	-	1	4	22	130	398	682	1,247	1,596	1,104	-
2001	5,077	-	-	1	7	32	121	343	639	1,252	1,611	1,071	-
2000	4,904	1	-	1	2	25	125	350	672	1,228	1,464	1,034	2
1999	4,895	-	-	1	4	33	130	387	680	1,211	1,417	1,030	2
Hispanic, female													
2005	5,148	-	-	-	1	6	31	126	330	785	1,673	2,196	-
2004	4,843	1	-	-	-	5	39	140	311	727	1,544	2,076	-
2003	5,035	1	-	-	1	10	44	122	337	765	1,579	2,176	-
2002	4,896	-	-	-	2	5	26	136	314	801	1,521	2,091	-
2001	4,971	-	-	2	4	6	26	121	291	833	1,545	2,143	-
2000	4,712	-	-	-	2	5	29	99	312	753	1,447	2,065	-
1999	4,767	-	-	-	2	8	20	98	311	827	1,436	2,065	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
All other forms of chronic ischemic heart disease (I20,I25.1-I25.9)													
Non-Hispanic 1/, both sexes													
2005	217,258	1	2	4	58	385	2,402	8,467	17,103	30,546	68,166	90,118	6
2004	217,859	2	5	10	53	399	2,544	8,412	16,556	31,548	70,322	88,005	3
2003	228,327	5	4	9	58	400	2,679	8,486	17,019	33,816	73,645	92,199	7
2002	232,347	8	2	10	44	366	2,761	8,671	17,002	35,206	75,895	92,375	7
2001	235,129	7	7	8	60	408	2,653	8,402	16,836	37,216	77,115	92,405	12
2000	238,835	7	8	8	44	365	2,746	8,256	16,922	39,060	77,847	93,558	14
1999	244,431	11	1	8	54	372	2,770	8,435	17,070	41,117	80,263	94,326	4
Non-Hispanic 1/, male													
2005	110,035	1	-	3	45	286	1,795	6,374	12,152	19,501	37,108	32,767	3
2004	109,273	1	1	4	38	299	1,888	6,311	11,705	20,074	37,996	30,953	3
2003	113,775	4	2	2	37	289	2,009	6,382	12,107	21,663	39,712	31,565	3
2002	115,448	6	2	9	33	267	2,087	6,634	12,083	22,499	40,716	31,108	4
2001	115,477	4	1	7	41	320	2,008	6,344	11,878	23,713	40,977	30,177	7
2000	117,223	5	5	3	31	279	2,088	6,302	11,948	24,916	41,345	30,293	8
1999	119,424	6	1	7	36	276	2,105	6,410	11,916	26,191	42,424	30,049	3
Non-Hispanic 1/, female													
2005	107,223	-	2	1	13	99	607	2,093	4,951	11,045	31,058	57,351	3
2004	108,586	1	4	6	15	100	656	2,101	4,851	11,474	32,326	57,052	-
2003	114,552	1	2	7	21	111	670	2,104	4,912	12,153	33,933	60,634	4
2002	116,899	2	-	1	11	99	674	2,037	4,919	12,707	35,179	61,267	3
2001	119,652	3	6	1	19	88	645	2,058	4,958	13,503	36,138	62,228	5
2000	121,612	2	3	5	13	86	658	1,954	4,974	14,144	36,502	63,265	6
1999	125,007	5	-	1	18	96	665	2,025	5,154	14,926	37,839	64,277	1

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
All other forms of chronic ischemic heart disease (I20,I25.1-I25.9)													
Non-Hispanic white, both sexes													
2005 2/	192,699	1	2	4	34	285	1,818	6,508	13,821	25,716	61,065	83,441	4
2004 2/	193,296	2	3	6	28	270	1,945	6,539	13,424	26,650	63,137	81,289	3
2003 2/	202,818	4	2	3	36	281	2,042	6,598	13,808	28,619	66,196	85,223	6
2002	207,175	3	-	7	28	267	2,146	6,751	13,776	30,132	68,477	85,582	6
2001	209,860	3	4	7	35	286	2,028	6,573	13,753	31,917	69,629	85,618	7
2000	213,991	5	8	7	30	257	2,091	6,507	13,795	33,706	70,626	86,949	10
1999	219,351	8	1	5	33	271	2,092	6,697	13,980	35,572	73,052	87,637	3
Non-Hispanic white, male													
2005 2/	97,953	1	-	3	28	211	1,388	5,012	10,086	16,788	33,794	30,639	3
2004 2/	97,326	1	1	3	21	205	1,483	5,037	9,749	17,350	34,653	28,820	3
2003 2/	101,510	3	1	1	22	208	1,577	5,117	10,122	18,754	36,209	29,494	2
2002	103,407	2	-	6	21	202	1,668	5,363	10,032	19,693	37,361	29,055	4
2001	103,407	2	-	6	26	229	1,577	5,120	9,942	20,776	37,531	28,194	4
2000	105,447	3	5	3	21	205	1,653	5,137	10,008	22,049	38,047	28,310	6
1999	107,463	4	1	5	23	203	1,631	5,233	9,987	23,196	39,196	27,981	3
Non-Hispanic white, female													
2005 2/	94,746	-	2	1	6	74	430	1,496	3,735	8,928	27,271	52,802	1
2004 2/	95,970	1	2	3	7	65	462	1,502	3,675	9,300	28,484	52,469	-
2003 2/	101,308	1	1	2	14	73	465	1,481	3,686	9,865	29,987	55,729	4
2002	103,768	1	-	1	7	65	478	1,388	3,744	10,439	31,116	56,527	2
2001	106,453	1	4	1	9	57	451	1,453	3,811	11,141	32,098	57,424	3
2000	108,544	2	3	4	9	52	438	1,370	3,787	11,657	32,579	58,639	4
1999	111,888	4	-	-	10	68	461	1,464	3,993	12,376	33,856	59,656	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
All other forms of chronic ischemic heart disease (I20,I25.1-I25.9)													
Non-Hispanic black, both sexes													
2005 2/	20,083	-	-	-	18	79	498	1,696	2,844	3,973	5,751	5,222	2
2004 2/	20,143	-	-	4	22	108	529	1,629	2,698	4,100	5,763	5,290	-
2003 2/	21,153	1	2	5	19	93	526	1,661	2,779	4,398	6,095	5,574	-
2002	20,893	4	2	3	13	77	533	1,675	2,782	4,247	6,047	5,509	1
2001	21,287	4	3	1	21	109	549	1,609	2,674	4,478	6,192	5,643	4
2000	20,993	1	-	-	11	96	577	1,517	2,697	4,599	6,001	5,490	4
1999	21,176	2	-	3	18	83	588	1,531	2,693	4,741	5,944	5,572	1
Non-Hispanic black, male													
2005 2/	9,598	-	-	-	12	57	334	1,162	1,752	2,171	2,607	1,503	-
2004 2/	9,536	-	-	1	16	74	345	1,082	1,650	2,251	2,592	1,525	-
2003 2/	9,853	1	1	1	13	60	348	1,082	1,675	2,418	2,776	1,478	-
2002	9,644	3	2	3	9	46	349	1,080	1,743	2,304	2,596	1,509	-
2001	9,902	2	1	1	12	81	371	1,065	1,629	2,455	2,751	1,531	3
2000	9,547	1	-	-	7	65	368	972	1,633	2,386	2,621	1,492	2
1999	9,692	1	-	2	10	60	394	1,008	1,653	2,488	2,529	1,547	-
Non-Hispanic black, female													
2005 2/	10,485	-	-	-	6	22	164	534	1,092	1,802	3,144	3,719	2
2004 2/	10,607	-	-	3	6	34	184	547	1,048	1,849	3,171	3,765	-
2003 2/	11,300	-	1	4	6	33	178	579	1,104	1,980	3,319	4,096	-
2002	11,249	1	-	-	4	31	184	595	1,039	1,943	3,451	4,000	1
2001	11,385	2	2	-	9	28	178	544	1,045	2,023	3,441	4,112	1
2000	11,446	-	-	-	4	31	209	545	1,064	2,213	3,380	3,998	2
1999	11,484	1	-	1	8	23	194	523	1,040	2,253	3,415	4,025	1

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
All other forms of chronic ischemic heart disease (I20,I25.1-I25.9)													
Origin not stated 3/, both sexes													
2005	397	-	-	-	-	-	7	22	45	57	135	131	-
2004	490	-	-	-	-	1	12	34	63	96	147	136	1
2003	676	-	-	-	1	2	11	29	81	124	207	221	-
2002	905	-	-	-	-	4	13	40	78	162	273	333	2
2001	801	1	-	-	1	2	7	36	77	132	262	282	1
2000	938	-	-	-	-	5	19	46	92	174	273	325	4
1999	954	-	-	-	1	1	13	43	79	175	293	347	2
Origin not stated 3/, male													
2005	209	-	-	-	-	-	5	18	26	40	79	41	-
2004	286	-	-	-	-	1	7	26	48	66	89	48	1
2003	353	-	-	-	1	1	6	23	56	84	109	73	-
2002	484	-	-	-	-	4	10	35	56	119	148	110	2
2001	416	1	-	-	1	2	4	29	66	83	145	84	1
2000	496	-	-	-	-	3	17	38	71	120	136	108	3
1999	482	-	-	-	-	1	10	34	60	121	135	119	2
Origin not stated 3/, female													
2005	188	-	-	-	-	-	2	4	19	17	56	90	-
2004	204	-	-	-	-	-	5	8	15	30	58	88	-
2003	323	-	-	-	-	1	5	6	25	40	98	148	-
2002	421	-	-	-	-	-	3	5	22	43	125	223	-
2001	385	-	-	-	-	-	3	7	11	49	117	198	-
2000	442	-	-	-	-	2	2	8	21	54	137	217	1
1999	472	-	-	-	1	-	3	9	19	54	158	228	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Other heart diseases (I26-I51)													
All origins, both sexes													
2005	170,585	342	143	231	918	1,847	4,295	9,120	13,861	21,763	47,866	70,193	6
2004	166,275	406	174	221	855	1,719	4,284	8,952	13,134	21,905	47,853	66,765	7
2003	169,984	416	171	234	904	1,810	4,555	8,716	13,398	23,244	49,260	67,270	6
2002	169,540	473	157	227	836	1,808	4,620	8,711	12,946	23,680	49,776	66,300	6
2001	166,949	449	215	246	794	1,736	4,472	8,484	12,535	24,056	48,976	64,979	7
2000	165,428	457	168	248	853	1,749	4,457	8,087	12,362	24,679	48,367	63,993	8
1999	165,828	490	177	248	880	1,779	4,647	7,888	12,580	25,572	48,712	62,848	7
All origins, male													
2005	75,094	188	77	118	614	1,176	2,695	5,769	8,415	11,987	21,954	22,097	4
2004	73,196	218	90	111	555	1,075	2,642	5,661	8,071	12,033	21,960	20,774	6
2003	74,738	233	81	141	560	1,081	2,801	5,458	8,142	12,875	22,680	20,681	5
2002	74,230	249	82	125	557	1,138	2,860	5,550	7,771	12,991	22,771	20,133	3
2001	72,439	222	117	123	500	1,037	2,777	5,307	7,503	13,167	22,540	19,141	5
2000	72,122	238	98	150	518	1,049	2,746	5,151	7,532	13,621	22,006	19,008	5
1999	72,601	255	100	137	533	1,067	2,894	4,979	7,700	14,320	22,128	18,483	5
All origins, female													
2005	95,491	154	66	113	304	671	1,600	3,351	5,446	9,776	25,912	48,096	2
2004	93,079	188	84	110	300	644	1,642	3,291	5,063	9,872	25,893	45,991	1
2003	95,246	183	90	93	344	729	1,754	3,258	5,256	10,369	26,580	46,589	1
2002	95,310	224	75	102	279	670	1,760	3,161	5,175	10,689	27,005	46,167	3
2001	94,510	227	98	123	294	699	1,695	3,177	5,032	10,889	26,436	45,838	2
2000	93,306	219	70	98	335	700	1,711	2,936	4,830	11,058	26,361	44,985	3
1999	93,227	235	77	111	347	712	1,753	2,909	4,880	11,252	26,584	44,365	2

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Other heart diseases (I26-I51)													
Hispanic, both sexes													
2005	5,937	70	21	28	111	214	326	464	588	933	1,537	1,644	1
2004	5,467	77	36	28	108	177	326	460	549	893	1,427	1,386	-
2003	5,612	72	33	26	98	209	311	433	621	945	1,417	1,447	-
2002	5,152	81	23	27	102	188	298	416	507	857	1,318	1,335	-
2001	4,899	75	31	33	86	179	268	370	516	856	1,206	1,277	2
2000	4,792	79	28	37	110	182	283	363	493	817	1,149	1,250	1
1999	4,687	92	33	34	99	176	309	345	501	815	1,102	1,180	1
Hispanic, male													
2005	3,032	38	15	15	72	161	233	321	361	514	741	560	1
2004	2,780	39	23	11	83	139	231	313	334	473	676	458	-
2003	2,805	40	17	15	68	135	217	281	367	515	661	489	-
2002	2,587	39	11	12	76	135	195	260	296	484	626	453	-
2001	2,401	40	14	17	57	119	185	252	296	456	556	407	2
2000	2,393	39	20	25	76	124	177	234	299	446	510	442	1
1999	2,392	40	20	21	60	128	208	225	298	466	513	412	1
Hispanic, female													
2005	2,905	32	6	13	39	53	93	143	227	419	796	1,084	-
2004	2,687	38	13	17	25	38	95	147	215	420	751	928	-
2003	2,807	32	16	11	30	74	94	152	254	430	756	958	-
2002	2,565	42	12	15	26	53	103	156	211	373	692	882	-
2001	2,498	35	17	16	29	60	83	118	220	400	650	870	-
2000	2,399	40	8	12	34	58	106	129	194	371	639	808	-
1999	2,295	52	13	13	39	48	101	120	203	349	589	768	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Other heart diseases (I26-I51)													
Non-Hispanic 1/, both sexes													
2005	164,389	272	122	202	807	1,628	3,957	8,624	13,234	20,791	46,264	68,483	5
2004	160,544	327	138	191	743	1,538	3,945	8,467	12,555	20,979	46,354	65,300	7
2003	163,998	342	137	207	802	1,594	4,225	8,249	12,731	22,243	47,750	65,714	4
2002	163,970	390	134	200	733	1,612	4,300	8,263	12,402	22,761	48,342	64,830	3
2001	161,625	372	183	213	701	1,550	4,184	8,077	11,980	23,137	47,661	63,564	3
2000	160,222	375	139	210	741	1,558	4,151	7,694	11,823	23,804	47,124	62,597	6
1999	160,711	397	143	213	779	1,594	4,317	7,520	12,030	24,685	47,496	61,532	5
Non-Hispanic 1/, male													
2005	71,921	150	62	103	542	1,012	2,453	5,427	8,024	11,445	21,180	21,520	3
2004	70,276	178	67	98	470	932	2,403	5,330	7,713	11,541	21,247	20,291	6
2003	71,731	193	64	125	490	942	2,569	5,150	7,739	12,332	21,969	20,155	3
2002	71,455	209	71	113	480	999	2,649	5,268	7,454	12,472	22,089	19,649	2
2001	69,830	181	103	106	439	913	2,576	5,030	7,181	12,671	21,941	18,687	2
2000	69,547	197	78	125	441	921	2,554	4,900	7,203	13,138	21,460	18,527	3
1999	69,984	215	80	115	471	933	2,672	4,737	7,370	13,812	21,555	18,021	3
Non-Hispanic 1/, female													
2005	92,468	122	60	99	265	616	1,504	3,197	5,210	9,346	25,084	46,963	2
2004	90,268	149	71	93	273	606	1,542	3,137	4,842	9,438	25,107	45,009	1
2003	92,267	149	73	82	312	652	1,656	3,099	4,992	9,911	25,781	45,559	1
2002	92,515	181	63	87	253	613	1,651	2,995	4,948	10,289	26,253	45,181	1
2001	91,795	191	80	107	262	637	1,608	3,047	4,799	10,466	25,720	44,877	1
2000	90,675	178	61	85	300	637	1,597	2,794	4,620	10,666	25,664	44,070	3
1999	90,727	182	63	98	308	661	1,645	2,783	4,660	10,873	25,941	43,511	2

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Other heart diseases (I26-I51)													
Non-Hispanic white, both sexes													
2005 2/	141,522	154	66	115	505	944	2,593	5,931	9,910	16,892	41,043	63,365	4
2004 2/	138,350	202	74	116	473	916	2,572	5,820	9,510	17,192	41,172	60,299	4
2003 2/	141,413	186	77	132	495	941	2,768	5,698	9,649	18,137	42,620	60,707	3
2002	141,712	234	86	130	453	936	2,888	5,703	9,468	18,836	43,140	59,836	2
2001	139,716	219	103	140	422	937	2,696	5,631	9,132	19,171	42,486	58,776	3
2000	138,745	238	87	122	428	948	2,717	5,362	9,089	19,781	42,174	57,794	5
1999	138,869	240	87	140	495	972	2,814	5,238	9,108	20,529	42,406	56,837	3
Non-Hispanic white, male													
2005 2/	61,136	82	38	63	339	604	1,652	3,851	6,113	9,375	18,999	20,017	3
2004 2/	59,673	110	35	63	301	561	1,622	3,769	5,932	9,613	18,968	18,696	3
2003 2/	61,119	102	32	82	298	576	1,737	3,644	5,970	10,234	19,773	18,669	2
2002	61,048	128	51	74	304	589	1,836	3,748	5,806	10,465	19,878	18,167	2
2001	59,513	116	58	72	261	555	1,731	3,637	5,562	10,653	19,627	17,239	2
2000	59,461	127	51	77	257	576	1,726	3,505	5,648	11,102	19,335	17,054	3
1999	59,549	136	51	73	291	580	1,809	3,395	5,680	11,656	19,365	16,511	2
Non-Hispanic white, female													
2005 2/	80,386	72	28	52	166	340	941	2,080	3,797	7,517	22,044	43,348	1
2004 2/	78,677	92	39	53	172	355	950	2,051	3,578	7,579	22,204	41,603	1
2003 2/	80,294	84	45	50	197	365	1,031	2,054	3,679	7,903	22,847	42,038	1
2002	80,664	106	35	56	149	347	1,052	1,955	3,662	8,371	23,262	41,669	-
2001	80,203	103	45	68	161	382	965	1,994	3,570	8,518	22,859	41,537	1
2000	79,284	111	36	45	171	372	991	1,857	3,441	8,679	22,839	40,740	2
1999	79,320	104	36	67	204	392	1,005	1,843	3,428	8,873	23,041	40,326	1

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Other heart diseases (I26-I51)													
Non-Hispanic black, both sexes													
2005 2/	19,938	98	44	72	270	619	1,248	2,471	2,975	3,429	4,491	4,220	1
2004 2/	19,469	97	59	69	236	546	1,265	2,448	2,771	3,362	4,451	4,162	3
2003 2/	19,861	125	49	66	273	586	1,346	2,373	2,795	3,659	4,375	4,213	1
2002	19,641	135	38	57	246	608	1,306	2,370	2,671	3,476	4,483	4,250	1
2001	19,496	130	70	63	252	567	1,370	2,264	2,608	3,522	4,508	4,142	-
2000	19,159	113	46	72	278	559	1,314	2,168	2,485	3,616	4,358	4,149	1
1999	19,531	132	48	66	262	576	1,395	2,107	2,678	3,747	4,454	4,064	2
Non-Hispanic black, male													
2005 2/	9,358	59	18	35	184	370	728	1,445	1,699	1,801	1,855	1,164	-
2004 2/	9,206	53	31	32	150	316	713	1,437	1,616	1,700	1,930	1,225	3
2003 2/	9,275	76	27	38	175	322	767	1,395	1,610	1,863	1,838	1,163	1
2002	9,038	72	16	30	151	363	738	1,402	1,490	1,772	1,854	1,150	-
2001	9,070	59	39	31	160	327	768	1,279	1,483	1,766	1,978	1,180	-
2000	8,883	56	24	39	158	313	747	1,295	1,410	1,802	1,833	1,206	-
1999	9,258	66	27	38	166	331	797	1,247	1,545	1,936	1,870	1,234	1
Non-Hispanic black, female													
2005 2/	10,580	39	26	37	86	249	520	1,026	1,276	1,628	2,636	3,056	1
2004 2/	10,263	44	28	37	86	230	552	1,011	1,155	1,662	2,521	2,937	-
2003 2/	10,586	49	22	28	98	264	579	978	1,185	1,796	2,537	3,050	-
2002	10,603	63	22	27	95	245	568	968	1,181	1,704	2,629	3,100	1
2001	10,426	71	31	32	92	240	602	985	1,125	1,756	2,530	2,962	-
2000	10,276	57	22	33	120	246	567	873	1,075	1,814	2,525	2,943	1
1999	10,273	66	21	28	96	245	598	860	1,133	1,811	2,584	2,830	1

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Other heart diseases (I26-I51)													
Origin not stated 3/, both sexes													
2005	259	-	-	1	-	5	12	32	39	39	65	66	-
2004	264	2	-	2	4	4	13	25	30	33	72	79	-
2003	374	2	1	1	4	7	19	34	46	56	93	109	2
2002	418	2	-	-	1	8	22	32	37	62	116	135	3
2001	425	2	1	-	7	7	20	37	39	63	109	138	2
2000	414	3	1	1	2	9	23	30	46	58	94	146	1
1999	430	1	1	1	2	9	21	23	49	72	114	136	1
Origin not stated 3/, male													
2005	141	-	-	-	-	3	9	21	30	28	33	17	-
2004	140	1	-	2	2	4	8	18	24	19	37	25	-
2003	202	-	-	1	2	4	15	27	36	28	50	37	2
2002	188	1	-	-	1	4	16	22	21	35	56	31	1
2001	208	1	-	-	4	5	16	25	26	40	43	47	1
2000	182	2	-	-	1	4	15	17	30	37	36	39	1
1999	225	-	-	1	2	6	14	17	32	42	60	50	1
Origin not stated 3/, female													
2005	118	-	-	1	-	2	3	11	9	11	32	49	-
2004	124	1	-	-	2	-	5	7	6	14	35	54	-
2003	172	2	1	-	2	3	4	7	10	28	43	72	-
2002	230	1	-	-	-	4	6	10	16	27	60	104	2
2001	217	1	1	-	3	2	4	12	13	23	66	91	1
2000	232	1	1	1	1	5	8	13	16	21	58	107	-
1999	205	1	1	-	-	3	7	6	17	30	54	86	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Acute and subacute endocarditis (I33)													
All origins, both sexes													
2005	1,209	4	-	3	8	28	78	173	198	234	328	155	-
2004	1,196	1	1	-	12	35	84	190	165	234	318	155	1
2003	1,245	1	-	1	16	33	112	183	209	232	318	140	-
2002	1,154	2	2	2	7	44	90	174	168	238	306	121	-
2001	1,178	1	-	3	8	29	117	186	160	247	316	111	-
2000	1,172	3	1	3	13	40	116	186	165	224	302	119	-
1999	1,116	2	1	-	7	30	114	143	171	240	289	119	-
All origins, male													
2005	695	3	-	-	5	20	51	113	125	124	173	81	-
2004	659	-	-	-	8	22	55	122	98	119	162	72	1
2003	682	1	-	1	6	19	63	125	123	119	166	59	-
2002	643	1	1	1	6	31	60	117	91	133	153	49	-
2001	668	1	-	1	5	17	62	129	95	132	177	49	-
2000	665	2	-	1	7	24	73	120	96	122	169	51	-
1999	639	2	1	-	3	24	75	103	103	118	154	56	-
All origins, female													
2005	514	1	-	3	3	8	27	60	73	110	155	74	-
2004	537	1	1	-	4	13	29	68	67	115	156	83	-
2003	563	-	-	-	10	14	49	58	86	113	152	81	-
2002	511	1	1	1	1	13	30	57	77	105	153	72	-
2001	510	-	-	2	3	12	55	57	65	115	139	62	-
2000	507	1	1	2	6	16	43	66	69	102	133	68	-
1999	477	-	-	-	4	6	39	40	68	122	135	63	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Acute and subacute endocarditis (I33)													
Hispanic, both sexes													
2005	80	1	-	-	3	5	12	8	18	15	15	3	-
2004	98	-	-	-	2	6	19	18	18	15	16	4	-
2003	80	1	-	1	3	4	10	17	12	16	13	3	-
2002	69	1	1	-	-	7	5	13	12	14	13	3	-
2001	56	-	-	1	1	6	8	10	9	9	10	2	-
2000	70	-	1	1	-	6	11	16	8	18	7	2	-
1999	63	-	1	-	2	6	10	7	10	13	12	2	-
Hispanic, male													
2005	58	1	-	-	3	4	9	7	12	12	9	1	-
2004	68	-	-	-	1	6	15	15	13	8	7	3	-
2003	42	1	-	1	1	3	7	9	6	9	3	2	-
2002	45	-	-	-	-	6	5	11	7	9	6	1	-
2001	34	-	-	-	1	5	6	8	4	5	4	1	-
2000	39	-	-	-	-	4	7	10	4	11	2	1	-
1999	42	-	1	-	-	5	6	6	5	9	8	2	-
Hispanic, female													
2005	22	-	-	-	-	1	3	1	6	3	6	2	-
2004	30	-	-	-	1	-	4	3	5	7	9	1	-
2003	38	-	-	-	2	1	3	8	6	7	10	1	-
2002	24	1	1	-	-	1	-	2	5	5	7	2	-
2001	22	-	-	1	-	1	2	2	5	4	6	1	-
2000	31	-	1	1	-	2	4	6	4	7	5	1	-
1999	21	-	-	-	2	1	4	1	5	4	4	-	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Acute and subacute endocarditis (I33)													
Non-Hispanic 1/, both sexes													
2005	1,127	3	-	2	5	23	66	165	179	219	313	152	-
2004	1,097	1	1	-	10	29	65	172	146	219	302	151	1
2003	1,163	-	-	-	13	29	102	164	197	216	305	137	-
2002	1,082	1	1	2	7	37	85	159	155	224	293	118	-
2001	1,117	1	-	2	7	23	108	174	150	238	305	109	-
2000	1,096	3	-	2	13	34	102	170	157	205	293	117	-
1999	1,049	2	-	-	5	23	104	135	160	226	277	117	-
Non-Hispanic 1/, male													
2005	636	2	-	-	2	16	42	106	112	112	164	80	-
2004	591	-	-	-	7	16	40	107	85	111	155	69	1
2003	638	-	-	-	5	16	56	114	117	110	163	57	-
2002	596	1	1	1	6	25	55	105	83	124	147	48	-
2001	631	1	-	1	4	12	55	120	90	127	173	48	-
2000	625	2	-	1	7	20	65	110	92	111	167	50	-
1999	594	2	-	-	3	18	69	96	97	109	146	54	-
Non-Hispanic 1/, female													
2005	491	1	-	2	3	7	24	59	67	107	149	72	-
2004	506	1	1	-	3	13	25	65	61	108	147	82	-
2003	525	-	-	-	8	13	46	50	80	106	142	80	-
2002	486	-	-	1	1	12	30	54	72	100	146	70	-
2001	486	-	-	1	3	11	53	54	60	111	132	61	-
2000	471	1	-	1	6	14	37	60	65	94	126	67	-
1999	455	-	-	-	2	5	35	39	63	117	131	63	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Acute and subacute endocarditis (I33)													
Non-Hispanic white, both sexes													
2005 2/	870	2	-	1	3	15	35	105	116	182	271	140	-
2004 2/	859	1	-	-	8	21	47	96	107	180	263	136	-
2003 2/	912	-	-	-	11	22	68	100	137	172	271	131	-
2002	862	-	1	-	6	27	62	95	113	179	270	109	-
2001	859	1	-	2	5	15	65	114	107	182	265	103	-
2000	852	1	-	1	6	21	66	106	114	173	261	103	-
1999	814	1	-	-	4	17	59	82	107	182	252	110	-
Non-Hispanic white, male													
2005 2/	496	1	-	-	1	10	23	68	71	96	150	76	-
2004 2/	468	-	-	-	5	10	32	60	66	93	138	64	-
2003 2/	500	-	-	-	5	11	39	67	85	89	148	56	-
2002	481	-	1	-	5	20	41	68	61	102	139	44	-
2001	482	1	-	1	3	7	34	79	65	93	154	45	-
2000	496	1	-	-	3	15	42	73	72	93	151	46	-
1999	466	1	-	-	2	12	39	59	68	93	139	53	-
Non-Hispanic white, female													
2005 2/	374	1	-	1	2	5	12	37	45	86	121	64	-
2004 2/	391	1	-	-	3	11	15	36	41	87	125	72	-
2003 2/	412	-	-	-	6	11	29	33	52	83	123	75	-
2002	381	-	-	-	1	7	21	27	52	77	131	65	-
2001	377	-	-	1	2	8	31	35	42	89	111	58	-
2000	356	-	-	1	3	6	24	33	42	80	110	57	-
1999	348	-	-	-	2	5	20	23	39	89	113	57	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Acute and subacute endocarditis (I33)													
Non-Hispanic black, both sexes													
2005 2/	226	1	-	1	2	6	27	54	58	31	36	10	-
2004 2/	203	-	1	-	1	6	14	68	35	35	27	15	1
2003 2/	222	-	-	-	2	5	32	60	53	37	29	4	-
2002	197	-	-	1	1	10	22	57	41	38	19	8	-
2001	233	-	-	-	1	8	41	57	41	43	37	5	-
2000	213	2	-	-	6	11	31	57	39	30	26	11	-
1999	208	-	-	-	1	6	41	48	47	42	17	6	-
Non-Hispanic black, male													
2005 2/	122	1	-	-	1	5	18	33	37	13	11	3	-
2004 2/	104	-	-	-	1	4	7	42	18	16	10	5	1
2003 2/	119	-	-	-	-	4	15	43	27	17	13	-	-
2002	104	-	-	-	1	5	14	33	22	19	6	4	-
2001	134	-	-	-	-	5	20	39	24	26	18	2	-
2000	111	1	-	-	4	3	21	35	18	16	11	2	-
1999	115	-	-	-	1	6	27	34	26	15	5	1	-
Non-Hispanic black, female													
2005 2/	104	-	-	1	1	1	9	21	21	18	25	7	-
2004 2/	99	-	1	-	-	2	7	26	17	19	17	10	-
2003 2/	103	-	-	-	2	1	17	17	26	20	16	4	-
2002	93	-	-	1	-	5	8	24	19	19	13	4	-
2001	99	-	-	-	1	3	21	18	17	17	19	3	-
2000	102	1	-	-	2	8	10	22	21	14	15	9	-
1999	93	-	-	-	-	-	14	14	21	27	12	5	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Acute and subacute endocarditis (I33)													
Origin not stated 3/, both sexes													
2005	2	-	-	1	-	-	-	-	1	-	-	-	-
2004	1	-	-	-	-	-	-	-	1	-	-	-	-
2003	2	-	-	-	-	-	-	2	-	-	-	-	-
2002	3	-	-	-	-	-	-	2	1	-	-	-	-
2001	5	-	-	-	-	-	1	2	1	-	1	-	-
2000	6	-	-	-	-	-	3	-	-	1	2	-	-
1999	4	-	-	-	-	1	-	1	1	1	-	-	-
Origin not stated 3/, male													
2005	1	-	-	-	-	-	-	-	1	-	-	-	-
2004	-	-	-	-	-	-	-	-	-	-	-	-	-
2003	2	-	-	-	-	-	-	2	-	-	-	-	-
2002	2	-	-	-	-	-	-	1	1	-	-	-	-
2001	3	-	-	-	-	-	1	1	1	-	-	-	-
2000	1	-	-	-	-	-	1	-	-	-	-	-	-
1999	3	-	-	-	-	1	-	1	1	-	-	-	-
Origin not stated 3/, female													
2005	1	-	-	1	-	-	-	-	-	-	-	-	-
2004	1	-	-	-	-	-	-	-	1	-	-	-	-
2003	-	-	-	-	-	-	-	-	-	-	-	-	-
2002	1	-	-	-	-	-	-	1	-	-	-	-	-
2001	2	-	-	-	-	-	-	1	-	-	1	-	-
2000	5	-	-	-	-	-	2	-	-	1	2	-	-
1999	1	-	-	-	-	-	-	-	-	1	-	-	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Diseases of pericardium and acute myocarditis (I30-I31,I40)													
All origins, both sexes													
2005	864	21	17	30	34	53	84	132	110	132	155	96	-
2004	779	21	25	17	31	39	67	107	111	115	159	87	-
2003	868	33	22	28	55	48	86	111	104	151	153	77	-
2002	848	26	22	22	36	46	75	107	118	143	157	96	-
2001	839	32	26	21	40	50	93	111	106	120	154	85	1
2000	767	20	24	20	38	53	86	85	89	131	145	76	-
1999	890	27	25	21	32	50	114	105	125	146	160	85	-
All origins, male													
2005	439	9	10	9	21	31	46	77	57	71	78	30	-
2004	404	11	15	10	18	18	40	63	65	62	68	34	-
2003	449	15	8	12	34	29	52	71	57	84	65	22	-
2002	434	9	11	7	24	32	49	69	69	71	60	33	-
2001	401	18	15	7	22	33	41	59	61	59	61	24	1
2000	404	10	13	9	21	29	58	46	52	67	73	26	-
1999	442	13	14	7	20	34	64	56	70	69	65	30	-
All origins, female													
2005	425	12	7	21	13	22	38	55	53	61	77	66	-
2004	375	10	10	7	13	21	27	44	46	53	91	53	-
2003	419	18	14	16	21	19	34	40	47	67	88	55	-
2002	414	17	11	15	12	14	26	38	49	72	97	63	-
2001	438	14	11	14	18	17	52	52	45	61	93	61	-
2000	363	10	11	11	17	24	28	39	37	64	72	50	-
1999	448	14	11	14	12	16	50	49	55	77	95	55	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Diseases of pericardium and acute myocarditis (I30-I31,I40)													
Hispanic, both sexes													
2005	74	4	1	4	6	11	10	15	5	10	5	3	-
2004	64	7	8	3	4	3	6	7	11	4	9	2	-
2003	57	5	5	4	7	6	5	6	7	7	5	-	-
2002	64	4	2	3	5	3	6	10	14	7	5	5	-
2001	58	2	4	2	4	11	5	6	6	6	7	5	-
2000	61	5	4	1	5	9	10	4	9	5	7	2	-
1999	64	2	3	4	10	8	9	9	2	7	8	2	-
Hispanic, male													
2005	40	1	1	1	3	8	5	10	2	7	2	-	-
2004	38	3	7	1	3	1	4	6	8	1	4	-	-
2003	33	3	1	2	4	4	3	5	4	6	1	-	-
2002	35	3	-	-	3	3	3	7	9	3	1	3	-
2001	25	1	2	1	2	6	3	1	3	3	2	1	-
2000	29	2	4	-	1	4	6	2	3	-	6	1	-
1999	30	1	1	4	5	6	7	2	1	1	2	-	-
Hispanic, female													
2005	34	3	-	3	3	3	5	5	3	3	3	3	-
2004	26	4	1	2	1	2	2	1	3	3	5	2	-
2003	24	2	4	2	3	2	2	1	3	1	4	-	-
2002	29	1	2	3	2	-	3	3	5	4	4	2	-
2001	33	1	2	1	2	5	2	5	3	3	5	4	-
2000	32	3	-	1	4	5	4	2	6	5	1	1	-
1999	34	1	2	-	5	2	2	7	1	6	6	2	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Diseases of pericardium and acute myocarditis (I30-I31,I40)													
Non-Hispanic 1/, both sexes													
2005	789	17	16	26	28	42	74	117	105	122	149	93	-
2004	715	14	17	14	27	36	61	100	100	111	150	85	-
2003	806	28	17	24	48	41	81	102	97	144	148	76	-
2002	783	22	20	19	31	43	69	97	103	136	152	91	-
2001	779	30	22	19	36	38	87	105	100	114	147	80	1
2000	700	15	19	19	32	44	76	81	78	126	136	74	-
1999	824	25	22	17	22	41	104	96	123	139	152	83	-
Non-Hispanic 1/, male													
2005	399	8	9	8	18	23	41	67	55	64	76	30	-
2004	366	8	8	9	15	17	36	57	57	61	64	34	-
2003	414	12	7	10	30	25	49	64	53	78	64	22	-
2002	398	6	11	7	21	29	46	62	59	68	59	30	-
2001	375	17	13	6	20	26	38	58	58	56	59	23	1
2000	374	8	9	9	20	25	52	44	48	67	67	25	-
1999	410	12	13	3	15	27	56	54	69	68	63	30	-
Non-Hispanic 1/, female													
2005	390	9	7	18	10	19	33	50	50	58	73	63	-
2004	349	6	9	5	12	19	25	43	43	50	86	51	-
2003	392	16	10	14	18	16	32	38	44	66	84	54	-
2002	385	16	9	12	10	14	23	35	44	68	93	61	-
2001	404	13	9	13	16	12	49	47	42	58	88	57	-
2000	326	7	10	10	12	19	24	37	30	59	69	49	-
1999	414	13	9	14	7	14	48	42	54	71	89	53	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Diseases of pericardium and acute myocarditis (I30-I31,I40)													
Non-Hispanic white, both sexes													
2005 2/	591	5	3	12	20	23	51	88	77	104	125	83	-
2004 2/	533	4	5	6	14	20	34	73	83	92	126	76	-
2003 2/	616	10	8	13	31	18	50	75	83	126	131	71	-
2002	601	9	11	9	21	30	49	69	76	116	126	85	-
2001	600	13	10	14	23	26	55	75	79	98	134	72	1
2000	528	6	11	9	20	25	49	55	62	103	122	66	-
1999	639	10	10	9	14	26	70	73	104	115	134	74	-
Non-Hispanic white, male													
2005 2/	306	2	3	5	15	12	28	53	40	56	64	28	-
2004 2/	268	4	-	4	11	9	22	38	48	47	55	30	-
2003 2/	324	6	3	5	21	9	33	49	45	71	60	22	-
2002	308	3	7	3	14	21	33	43	45	60	51	28	-
2001	286	10	6	5	10	18	27	41	45	48	53	22	1
2000	280	4	7	5	12	14	34	30	41	52	59	22	-
1999	320	6	4	2	10	17	40	40	60	56	58	27	-
Non-Hispanic white, female													
2005 2/	285	3	-	7	5	11	23	35	37	48	61	55	-
2004 2/	265	-	5	2	3	11	12	35	35	45	71	46	-
2003 2/	292	4	5	8	10	9	17	26	38	55	71	49	-
2002	293	6	4	6	7	9	16	26	31	56	75	57	-
2001	314	3	4	9	13	8	28	34	34	50	81	50	-
2000	248	2	4	4	8	11	15	25	21	51	63	44	-
1999	319	4	6	7	4	9	30	33	44	59	76	47	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Diseases of pericardium and acute myocarditis (I30-I31,I40)													
Non-Hispanic black, both sexes													
2005 2/	161	10	11	9	5	14	17	27	25	14	21	8	-
2004 2/	162	8	11	8	10	12	26	26	16	17	22	6	-
2003 2/	168	14	9	9	15	22	30	23	12	16	13	5	-
2002	154	12	7	8	8	10	17	26	22	19	20	5	-
2001	157	14	12	4	13	10	24	30	19	13	11	7	-
2000	140	7	7	8	7	16	24	21	15	18	10	7	-
1999	158	10	11	8	6	14	30	20	17	20	16	6	-
Non-Hispanic black, male													
2005 2/	72	5	5	1	3	9	7	13	12	5	11	1	-
2004 2/	88	3	7	5	4	6	14	18	8	13	9	1	-
2003 2/	81	4	4	5	9	15	16	13	7	5	3	-	-
2002	75	3	3	3	5	6	10	17	13	7	7	1	-
2001	78	5	7	1	10	7	7	17	11	8	5	-	-
2000	72	4	2	3	3	10	16	10	6	11	5	2	-
1999	76	4	9	1	3	10	13	12	9	11	4	-	-
Non-Hispanic black, female													
2005 2/	89	5	6	8	2	5	10	14	13	9	10	7	-
2004 2/	74	5	4	3	6	6	12	8	8	4	13	5	-
2003 2/	87	10	5	4	6	7	14	10	5	11	10	5	-
2002	79	9	4	5	3	4	7	9	9	12	13	4	-
2001	79	9	5	3	3	3	17	13	8	5	6	7	-
2000	68	3	5	5	4	6	8	11	9	7	5	5	-
1999	82	6	2	7	3	4	17	8	8	9	12	6	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Diseases of pericardium and acute myocarditis (I30-I31,I40)													
Origin not stated 3/, both sexes													
2005	1	-	-	-	-	-	-	-	-	-	1	-	-
2004	-	-	-	-	-	-	-	-	-	-	-	-	-
2003	5	-	-	-	-	1	-	3	-	-	-	1	-
2002	1	-	-	-	-	-	-	-	1	-	-	-	-
2001	2	-	-	-	-	1	1	-	-	-	-	-	-
2000	6	-	1	-	1	-	-	-	2	-	2	-	-
1999	2	-	-	-	-	1	1	-	-	-	-	-	-
Origin not stated 3/, male													
2005	-	-	-	-	-	-	-	-	-	-	-	-	-
2004	-	-	-	-	-	-	-	-	-	-	-	-	-
2003	2	-	-	-	-	-	-	2	-	-	-	-	-
2002	1	-	-	-	-	-	-	-	1	-	-	-	-
2001	1	-	-	-	-	1	-	-	-	-	-	-	-
2000	1	-	-	-	-	-	-	-	1	-	-	-	-
1999	2	-	-	-	-	1	1	-	-	-	-	-	-
Origin not stated 3/, female													
2005	1	-	-	-	-	-	-	-	-	-	1	-	-
2004	-	-	-	-	-	-	-	-	-	-	-	-	-
2003	3	-	-	-	-	1	-	1	-	-	-	1	-
2002	-	-	-	-	-	-	-	-	-	-	-	-	-
2001	1	-	-	-	-	-	1	-	-	-	-	-	-
2000	5	-	1	-	1	-	-	-	1	-	2	-	-
1999	-	-	-	-	-	-	-	-	-	-	-	-	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Heart failure (I50)													
All origins, both sexes													
2005	58,933	27	6	11	42	94	316	1,087	2,610	5,888	17,331	31,521	-
2004	57,120	30	11	10	33	90	313	1,039	2,444	5,951	17,012	30,186	1
2003	57,448	38	5	12	38	97	315	981	2,469	6,138	17,136	30,219	-
2002	56,494	23	11	10	27	101	282	993	2,342	6,168	16,910	29,627	-
2001	56,934	24	17	15	34	92	307	954	2,374	6,274	17,134	29,708	1
2000	55,704	31	9	21	34	63	262	811	2,211	6,403	16,820	29,035	4
1999	54,913	11	10	12	31	65	272	782	2,107	6,524	16,771	28,327	1
All origins, male													
2005	23,026	14	4	5	27	61	207	653	1,509	3,069	7,804	9,673	-
2004	22,292	16	4	7	19	59	186	615	1,420	3,163	7,673	9,130	-
2003	22,427	28	3	7	24	60	193	587	1,454	3,313	7,658	9,100	-
2002	21,698	14	6	5	15	60	173	617	1,345	3,254	7,573	8,636	-
2001	21,632	16	5	8	16	56	182	588	1,347	3,344	7,672	8,398	-
2000	21,175	22	7	9	20	36	159	496	1,264	3,414	7,498	8,248	2
1999	20,755	5	7	6	23	32	164	486	1,228	3,526	7,318	7,959	1
All origins, female													
2005	35,907	13	2	6	15	33	109	434	1,101	2,819	9,527	21,848	-
2004	34,828	14	7	3	14	31	127	424	1,024	2,788	9,339	21,056	1
2003	35,021	10	2	5	14	37	122	394	1,015	2,825	9,478	21,119	-
2002	34,796	9	5	5	12	41	109	376	997	2,914	9,337	20,991	-
2001	35,302	8	12	7	18	36	125	366	1,027	2,930	9,462	21,310	1
2000	34,529	9	2	12	14	27	103	315	947	2,989	9,322	20,787	2
1999	34,158	6	3	6	8	33	108	296	879	2,998	9,453	20,368	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Heart failure (I50)													
Hispanic, both sexes													
2005	1,721	5	1	-	3	6	16	49	106	244	551	740	-
2004	1,545	5	5	-	3	5	22	54	104	244	499	604	-
2003	1,606	7	-	1	4	10	12	48	118	275	494	637	-
2002	1,412	4	-	2	-	3	13	43	91	235	451	570	-
2001	1,364	8	3	-	1	9	21	43	103	213	404	559	-
2000	1,270	7	1	2	-	3	10	22	100	203	365	557	-
1999	1,242	3	4	1	3	4	18	29	87	202	353	538	-
Hispanic, male													
2005	735	3	1	-	2	5	12	30	48	127	270	237	-
2004	668	4	3	-	2	3	10	34	64	125	235	188	-
2003	714	5	-	-	4	7	10	25	77	141	242	203	-
2002	593	2	-	2	-	2	6	20	54	121	209	177	-
2001	565	6	1	-	1	6	15	31	52	109	169	175	-
2000	535	5	1	1	-	2	6	9	62	109	151	189	-
1999	522	1	3	-	3	3	12	16	47	112	159	166	-
Hispanic, female													
2005	986	2	-	-	1	1	4	19	58	117	281	503	-
2004	877	1	2	-	1	2	12	20	40	119	264	416	-
2003	892	2	-	1	-	3	2	23	41	134	252	434	-
2002	819	2	-	-	-	1	7	23	37	114	242	393	-
2001	799	2	2	-	-	3	6	12	51	104	235	384	-
2000	735	2	-	1	-	1	4	13	38	94	214	368	-
1999	720	2	1	1	-	1	6	13	40	90	194	372	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Heart failure (I50)													
Non-Hispanic 1/, both sexes													
2005	57,140	22	5	11	39	88	299	1,033	2,499	5,635	16,756	30,753	-
2004	55,488	25	6	10	30	85	291	981	2,334	5,698	16,487	29,540	1
2003	55,732	31	5	11	34	87	302	933	2,345	5,843	16,614	29,527	-
2002	54,932	19	11	8	27	97	268	943	2,245	5,914	16,416	28,984	-
2001	55,437	16	14	15	33	82	286	906	2,258	6,044	16,688	29,094	1
2000	54,307	24	8	19	34	60	252	789	2,105	6,185	16,417	28,411	3
1999	53,528	8	6	11	28	61	253	750	2,007	6,308	16,377	27,718	1
Non-Hispanic 1/, male													
2005	22,255	11	3	5	25	56	194	618	1,457	2,936	7,522	9,428	-
2004	21,580	12	1	7	17	56	176	579	1,350	3,032	7,425	8,925	-
2003	21,658	23	3	7	20	53	182	562	1,371	3,161	7,400	8,876	-
2002	21,050	12	6	3	15	58	166	592	1,288	3,124	7,344	8,442	-
2001	21,012	10	4	8	15	49	167	555	1,287	3,225	7,489	8,203	-
2000	20,599	17	6	8	20	34	153	487	1,197	3,296	7,335	8,045	1
1999	20,170	4	4	6	20	29	152	469	1,173	3,406	7,140	7,766	1
Non-Hispanic 1/, female													
2005	34,885	11	2	6	14	32	105	415	1,042	2,699	9,234	21,325	-
2004	33,908	13	5	3	13	29	115	402	984	2,666	9,062	20,615	1
2003	34,074	8	2	4	14	34	120	371	974	2,682	9,214	20,651	-
2002	33,882	7	5	5	12	39	102	351	957	2,790	9,072	20,542	-
2001	34,425	6	10	7	18	33	119	351	971	2,819	9,199	20,891	1
2000	33,708	7	2	11	14	26	99	302	908	2,889	9,082	20,366	2
1999	33,358	4	2	5	8	32	101	281	834	2,902	9,237	19,952	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Heart failure (I50)													
Non-Hispanic white, both sexes													
2005 2/	50,835	11	2	4	22	42	161	640	1,835	4,601	14,914	28,603	-
2004 2/	49,628	15	5	5	18	46	171	601	1,719	4,767	14,804	27,476	1
2003 2/	49,788	13	4	6	17	48	165	588	1,743	4,832	14,932	27,440	-
2002	49,162	11	8	5	16	49	160	593	1,719	4,914	14,725	26,962	-
2001	49,788	11	12	11	17	49	167	608	1,730	5,052	15,033	27,097	1
2000	48,782	13	4	14	17	35	154	536	1,611	5,176	14,827	26,392	3
1999	47,977	5	4	8	15	30	156	480	1,537	5,302	14,702	25,738	-
Non-Hispanic white, male													
2005 2/	19,666	5	2	1	16	28	98	391	1,083	2,415	6,774	8,853	-
2004 2/	19,125	6	1	4	11	28	110	362	995	2,580	6,711	8,317	-
2003 2/	19,239	8	2	4	9	31	105	366	1,033	2,672	6,725	8,284	-
2002	18,705	8	5	2	7	28	105	387	1,000	2,620	6,663	7,880	-
2001	18,730	6	4	5	9	25	103	392	1,019	2,734	6,785	7,648	-
2000	18,386	10	4	6	10	20	103	342	942	2,828	6,671	7,449	1
1999	17,903	2	2	5	11	14	100	306	922	2,920	6,472	7,149	-
Non-Hispanic white, female													
2005 2/	31,169	6	-	3	6	14	63	249	752	2,186	8,140	19,750	-
2004 2/	30,503	9	4	1	7	18	61	239	724	2,187	8,093	19,159	1
2003 2/	30,549	5	2	2	8	17	60	222	710	2,160	8,207	19,156	-
2002	30,457	3	3	3	9	21	55	206	719	2,294	8,062	19,082	-
2001	31,058	5	8	6	8	24	64	216	711	2,318	8,248	19,449	1
2000	30,396	3	-	8	7	15	51	194	669	2,348	8,156	18,943	2
1999	30,074	3	2	3	4	16	56	174	615	2,382	8,230	18,589	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Heart failure (I50)													
Non-Hispanic black, both sexes													
2005 2/	5,570	7	3	6	16	45	131	366	616	941	1,616	1,823	-
2004 2/	5,198	8	1	4	11	36	111	367	561	829	1,487	1,783	-
2003 2/	5,294	15	-	5	17	37	131	328	563	921	1,466	1,811	-
2002	5,143	8	3	3	9	45	104	331	487	891	1,495	1,767	-
2001	5,094	5	2	3	16	32	112	284	494	905	1,475	1,766	-
2000	4,936	11	4	4	13	24	94	241	453	909	1,407	1,776	-
1999	4,961	3	2	3	12	30	93	257	446	922	1,468	1,724	1
Non-Hispanic black, male													
2005 2/	2,261	4	1	3	9	28	90	212	344	465	652	453	-
2004 2/	2,154	5	-	3	6	25	59	210	321	397	626	502	-
2003 2/	2,132	13	-	3	11	20	73	183	320	445	575	489	-
2002	2,055	4	1	1	6	27	59	192	266	450	602	447	-
2001	2,036	4	-	2	6	23	61	153	248	447	618	474	-
2000	1,942	7	2	2	8	13	47	138	231	420	572	502	-
1999	2,005	2	2	1	8	15	50	156	238	442	577	513	1
Non-Hispanic black, female													
2005 2/	3,309	3	2	3	7	17	41	154	272	476	964	1,370	-
2004 2/	3,044	3	1	1	5	11	52	157	240	432	861	1,281	-
2003 2/	3,162	2	-	2	6	17	58	145	243	476	891	1,322	-
2002	3,088	4	2	2	3	18	45	139	221	441	893	1,320	-
2001	3,058	1	2	1	10	9	51	131	246	458	857	1,292	-
2000	2,994	4	2	2	5	11	47	103	222	489	835	1,274	-
1999	2,956	1	-	2	4	15	43	101	208	480	891	1,211	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Heart failure (I50)													
Origin not stated 3/, both sexes													
2005	72	-	-	-	-	-	1	5	5	9	24	28	-
2004	87	-	-	-	-	-	-	4	6	9	26	42	-
2003	110	-	-	-	-	-	1	-	6	20	28	55	-
2002	150	-	-	-	-	1	1	7	6	19	43	73	-
2001	133	-	-	-	-	1	-	5	13	17	42	55	-
2000	127	-	-	-	-	-	-	-	6	15	38	67	1
1999	143	-	-	-	-	-	1	3	13	14	41	71	-
Origin not stated 3/, male													
2005	36	-	-	-	-	-	1	5	4	6	12	8	-
2004	44	-	-	-	-	-	-	2	6	6	13	17	-
2003	55	-	-	-	-	-	1	-	6	11	16	21	-
2002	55	-	-	-	-	-	1	5	3	9	20	17	-
2001	55	-	-	-	-	1	-	2	8	10	14	20	-
2000	41	-	-	-	-	-	-	-	5	9	12	14	1
1999	63	-	-	-	-	-	-	1	8	8	19	27	-
Origin not stated 3/, female													
2005	36	-	-	-	-	-	-	-	1	3	12	20	-
2004	43	-	-	-	-	-	-	2	-	3	13	25	-
2003	55	-	-	-	-	-	-	-	-	9	12	34	-
2002	95	-	-	-	-	1	-	2	3	10	23	56	-
2001	78	-	-	-	-	-	-	3	5	7	28	35	-
2000	86	-	-	-	-	-	-	-	1	6	26	53	-
1999	80	-	-	-	-	-	1	2	5	6	22	44	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
All other forms of heart disease (I26-I28,I34-I38,I42-I49,I51)													
All origins, both sexes													
2005	109,579	290	120	187	834	1,672	3,817	7,728	10,943	15,509	30,052	38,421	6
2004	107,180	354	137	194	779	1,555	3,820	7,616	10,414	15,605	30,364	36,337	5
2003	110,423	344	144	193	795	1,632	4,042	7,441	10,616	16,723	31,653	36,834	6
2002	111,044	422	122	193	766	1,617	4,173	7,437	10,318	17,131	32,403	36,456	6
2001	107,998	392	172	207	712	1,565	3,955	7,233	9,895	17,415	31,372	35,075	5
2000	107,785	403	134	204	768	1,593	3,993	7,005	9,897	17,921	31,100	34,763	4
1999	108,909	450	141	215	810	1,634	4,147	6,858	10,177	18,662	31,492	34,317	6
All origins, male													
2005	50,934	162	63	104	561	1,064	2,391	4,926	6,724	8,723	13,899	12,313	4
2004	49,841	191	71	94	510	976	2,361	4,861	6,488	8,689	14,057	11,538	5
2003	51,180	189	70	121	496	973	2,493	4,675	6,508	9,359	14,791	11,500	5
2002	51,455	225	64	112	512	1,015	2,578	4,747	6,266	9,533	14,985	11,415	3
2001	49,738	187	97	107	457	931	2,492	4,531	6,000	9,632	14,630	10,670	4
2000	49,878	204	78	131	470	960	2,456	4,489	6,120	10,018	14,266	10,683	3
1999	50,765	235	78	124	487	977	2,591	4,334	6,299	10,607	14,591	10,438	4
All origins, female													
2005	58,645	128	57	83	273	608	1,426	2,802	4,219	6,786	16,153	26,108	2
2004	57,339	163	66	100	269	579	1,459	2,755	3,926	6,916	16,307	24,799	-
2003	59,243	155	74	72	299	659	1,549	2,766	4,108	7,364	16,862	25,334	1
2002	59,589	197	58	81	254	602	1,595	2,690	4,052	7,598	17,418	25,041	3
2001	58,260	205	75	100	255	634	1,463	2,702	3,895	7,783	16,742	24,405	1
2000	57,907	199	56	73	298	633	1,537	2,516	3,777	7,903	16,834	24,080	1
1999	58,144	215	63	91	323	657	1,556	2,524	3,878	8,055	16,901	23,879	2

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
All other forms of heart disease (I26-I28,I34-I38,I42-I49,I51)													
Hispanic, both sexes													
2005	4,062	60	19	24	99	192	288	392	459	664	966	898	1
2004	3,760	65	23	25	99	163	279	381	416	630	903	776	-
2003	3,869	59	28	20	84	189	284	362	484	647	905	807	-
2002	3,607	72	20	22	97	175	274	350	390	601	849	757	-
2001	3,421	65	24	30	80	153	234	311	398	628	785	711	2
2000	3,391	67	22	33	105	164	252	321	376	591	770	689	1
1999	3,318	87	25	29	84	158	272	300	402	593	729	638	1
Hispanic, male													
2005	2,199	33	13	14	64	144	207	274	299	368	460	322	1
2004	2,006	32	13	10	77	129	202	258	249	339	430	267	-
2003	2,016	31	16	12	59	121	197	242	280	359	415	284	-
2002	1,914	34	11	10	73	124	181	222	226	351	410	272	-
2001	1,777	33	11	16	53	102	161	212	237	339	381	230	2
2000	1,790	32	15	24	75	114	158	213	230	326	351	251	1
1999	1,798	38	15	17	52	114	183	201	245	344	344	244	1
Hispanic, female													
2005	1,863	27	6	10	35	48	81	118	160	296	506	576	-
2004	1,754	33	10	15	22	34	77	123	167	291	473	509	-
2003	1,853	28	12	8	25	68	87	120	204	288	490	523	-
2002	1,693	38	9	12	24	51	93	128	164	250	439	485	-
2001	1,644	32	13	14	27	51	73	99	161	289	404	481	-
2000	1,601	35	7	9	30	50	94	108	146	265	419	438	-
1999	1,520	49	10	12	32	44	89	99	157	249	385	394	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
All other forms of heart disease (I26-I28,I34-I38,I42-I49,I51)													
Non-Hispanic 1/, both sexes													
2005	105,333	230	101	163	735	1,475	3,518	7,309	10,451	14,815	29,046	37,485	5
2004	103,244	287	114	167	676	1,388	3,528	7,214	9,975	14,951	29,415	35,524	5
2003	106,297	283	115	172	707	1,437	3,740	7,050	10,092	16,040	30,683	35,974	4
2002	107,173	348	102	171	668	1,435	3,878	7,064	9,899	16,487	31,481	35,637	3
2001	104,292	325	147	177	625	1,407	3,703	6,892	9,472	16,741	30,521	34,281	1
2000	104,119	333	112	170	662	1,420	3,721	6,654	9,483	17,288	30,278	33,995	3
1999	105,310	362	115	185	724	1,469	3,856	6,539	9,740	18,012	30,690	33,614	4
Non-Hispanic 1/, male													
2005	48,631	129	50	90	497	917	2,176	4,636	6,400	8,333	13,418	11,982	3
2004	47,739	158	58	82	431	843	2,151	4,587	6,221	8,337	13,603	11,263	5
2003	49,021	158	54	108	435	848	2,282	4,410	6,198	8,983	14,342	11,200	3
2002	49,411	190	53	102	438	887	2,382	4,509	6,024	9,156	14,539	11,129	2
2001	47,812	153	86	91	400	826	2,316	4,297	5,746	9,263	14,220	10,413	1
2000	47,949	170	63	107	394	842	2,284	4,259	5,866	9,664	13,891	10,407	2
1999	48,810	197	63	106	433	859	2,395	4,118	6,031	10,229	14,206	10,171	2
Non-Hispanic 1/, female													
2005	56,702	101	51	73	238	558	1,342	2,673	4,051	6,482	15,628	25,503	2
2004	55,505	129	56	85	245	545	1,377	2,627	3,754	6,614	15,812	24,261	-
2003	57,276	125	61	64	272	589	1,458	2,640	3,894	7,057	16,341	24,774	1
2002	57,762	158	49	69	230	548	1,496	2,555	3,875	7,331	16,942	24,508	1
2001	56,480	172	61	86	225	581	1,387	2,595	3,726	7,478	16,301	23,868	-
2000	56,170	163	49	63	268	578	1,437	2,395	3,617	7,624	16,387	23,588	1
1999	56,500	165	52	79	291	610	1,461	2,421	3,709	7,783	16,484	23,443	2

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
All other forms of heart disease (I26-I28,I34-I38,I42-I49,I51)													
Non-Hispanic white, both sexes													
2005 2/	89,226	136	61	98	460	864	2,346	5,098	7,882	12,005	25,733	34,539	4
2004 2/	87,330	182	64	105	433	829	2,320	5,050	7,601	12,153	25,979	32,611	3
2003 2/	90,097	163	65	113	436	853	2,485	4,935	7,686	13,007	27,286	33,065	3
2002	91,087	214	66	116	410	830	2,617	4,946	7,560	13,627	28,019	32,680	2
2001	88,469	194	81	113	377	847	2,409	4,834	7,216	13,839	27,054	31,504	1
2000	88,583	218	72	98	385	867	2,448	4,665	7,302	14,329	26,964	31,233	2
1999	89,439	224	73	123	462	899	2,529	4,603	7,360	14,930	27,318	30,915	3
Non-Hispanic white, male													
2005 2/	40,668	74	33	57	307	554	1,503	3,339	4,919	6,808	12,011	11,060	3
2004 2/	39,812	100	34	55	274	514	1,458	3,309	4,823	6,893	12,064	10,285	3
2003 2/	41,056	88	27	73	263	525	1,560	3,162	4,807	7,402	12,840	10,307	2
2002	41,554	117	38	69	278	520	1,657	3,250	4,700	7,683	13,025	10,215	2
2001	40,015	99	48	61	239	505	1,567	3,125	4,433	7,778	12,635	9,524	1
2000	40,299	112	40	66	232	527	1,547	3,060	4,593	8,129	12,454	9,537	2
1999	40,860	127	45	66	268	537	1,630	2,990	4,630	8,587	12,696	9,282	2
Non-Hispanic white, female													
2005 2/	48,558	62	28	41	153	310	843	1,759	2,963	5,197	13,722	23,479	1
2004 2/	47,518	82	30	50	159	315	862	1,741	2,778	5,260	13,915	22,326	-
2003 2/	49,041	75	38	40	173	328	925	1,773	2,879	5,605	14,446	22,758	1
2002	49,533	97	28	47	132	310	960	1,696	2,860	5,944	14,994	22,465	-
2001	48,454	95	33	52	138	342	842	1,709	2,783	6,061	14,419	21,980	-
2000	48,284	106	32	32	153	340	901	1,605	2,709	6,200	14,510	21,696	-
1999	48,579	97	28	57	194	362	899	1,613	2,730	6,343	14,622	21,633	1

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
All other forms of heart disease (I26-I28,I34-I38,I42-I49,I51)													
Non-Hispanic black, both sexes													
2005 2/	13,981	80	30	56	247	554	1,073	2,024	2,276	2,443	2,818	2,379	1
2004 2/	13,906	81	46	57	214	492	1,114	1,987	2,159	2,481	2,915	2,358	2
2003 2/	14,177	96	40	52	239	522	1,153	1,962	2,167	2,685	2,867	2,393	1
2002	14,147	115	28	45	228	543	1,163	1,956	2,121	2,528	2,949	2,470	1
2001	14,012	111	56	56	222	517	1,193	1,893	2,054	2,561	2,985	2,364	-
2000	13,870	93	35	60	252	508	1,165	1,849	1,978	2,659	2,915	2,355	1
1999	14,204	119	35	55	243	526	1,231	1,782	2,168	2,763	2,953	2,328	1
Non-Hispanic black, male													
2005 2/	6,903	49	12	31	171	328	613	1,187	1,306	1,318	1,181	707	-
2004 2/	6,860	45	24	24	139	281	633	1,167	1,269	1,274	1,285	717	2
2003 2/	6,943	59	23	30	155	283	663	1,156	1,256	1,396	1,247	674	1
2002	6,804	65	12	26	139	325	655	1,160	1,189	1,296	1,239	698	-
2001	6,822	50	32	28	144	292	680	1,070	1,200	1,285	1,337	704	-
2000	6,758	44	20	34	143	287	663	1,112	1,155	1,355	1,245	700	-
1999	7,062	60	16	36	154	300	707	1,045	1,272	1,468	1,284	720	-
Non-Hispanic black, female													
2005 2/	7,078	31	18	25	76	226	460	837	970	1,125	1,637	1,672	1
2004 2/	7,046	36	22	33	75	211	481	820	890	1,207	1,630	1,641	-
2003 2/	7,234	37	17	22	84	239	490	806	911	1,289	1,620	1,719	-
2002	7,343	50	16	19	89	218	508	796	932	1,232	1,710	1,772	1
2001	7,190	61	24	28	78	225	513	823	854	1,276	1,648	1,660	-
2000	7,112	49	15	26	109	221	502	737	823	1,304	1,670	1,655	1
1999	7,142	59	19	19	89	226	524	737	896	1,295	1,669	1,608	1

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
All other forms of heart disease (I26-I28,I34-I38,I42-I49,I51)													
Origin not stated 3/, both sexes													
2005	184	-	-	-	-	5	11	27	33	30	40	38	-
2004	176	2	-	2	4	4	13	21	23	24	46	37	-
2003	257	2	1	1	4	6	18	29	40	36	65	53	2
2002	264	2	-	-	1	7	21	23	29	43	73	62	3
2001	285	2	1	-	7	5	18	30	25	46	66	83	2
2000	275	3	-	1	1	9	20	30	38	42	52	79	-
1999	281	1	1	1	2	7	19	19	35	57	73	65	1
Origin not stated 3/, male													
2005	104	-	-	-	-	3	8	16	25	22	21	9	-
2004	96	1	-	2	2	4	8	16	18	13	24	8	-
2003	143	-	-	1	2	4	14	23	30	17	34	16	2
2002	130	1	-	-	1	4	15	16	16	26	36	14	1
2001	149	1	-	-	4	3	15	22	17	30	29	27	1
2000	139	2	-	-	1	4	14	17	24	28	24	25	-
1999	157	-	-	1	2	4	13	15	23	34	41	23	1
Origin not stated 3/, female													
2005	80	-	-	-	-	2	3	11	8	8	19	29	-
2004	80	1	-	-	2	-	5	5	5	11	22	29	-
2003	114	2	1	-	2	2	4	6	10	19	31	37	-
2002	134	1	-	-	-	3	6	7	13	17	37	48	2
2001	136	1	1	-	3	2	3	8	8	16	37	56	1
2000	136	1	-	1	-	5	6	13	14	14	28	54	-
1999	124	1	1	-	-	3	6	4	12	23	32	42	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Essential (primary) hypertension and hypertensive renal disease (I10,I12)													
All origins, both sexes													
2005	24,902	2	-	3	22	84	406	1,166	1,954	3,306	7,256	10,703	-
2004	23,076	2	1	1	21	103	348	1,138	1,839	3,152	6,819	9,648	4
2003	21,940	1	2	2	22	98	372	1,031	1,755	3,104	6,649	8,904	-
2002	20,261	5	1	1	21	85	354	923	1,526	2,922	6,138	8,285	-
2001	19,250	-	1	1	23	104	331	931	1,462	2,832	5,993	7,572	-
2000	18,073	1	1	4	17	70	348	878	1,440	2,784	5,623	6,907	-
1999	16,968	3	3	1	13	78	313	799	1,318	2,799	5,324	6,317	-
All origins, male													
2005	9,458	1	-	1	12	45	243	733	1,151	1,639	2,917	2,716	-
2004	8,869	1	1	-	13	59	222	714	1,036	1,535	2,782	2,503	3
2003	8,213	1	2	1	12	58	215	660	989	1,535	2,522	2,218	-
2002	7,647	2	1	1	11	47	219	560	850	1,419	2,405	2,132	-
2001	7,254	-	-	-	16	59	206	569	784	1,379	2,282	1,959	-
2000	6,895	1	-	3	4	39	216	555	818	1,308	2,204	1,747	-
1999	6,488	2	1	1	8	48	195	482	732	1,372	2,067	1,580	-
All origins, female													
2005	15,444	1	-	2	10	39	163	433	803	1,667	4,339	7,987	-
2004	14,207	1	-	1	8	44	126	424	803	1,617	4,037	7,145	1
2003	13,727	-	-	1	10	40	157	371	766	1,569	4,127	6,686	-
2002	12,614	3	-	-	10	38	135	363	676	1,503	3,733	6,153	-
2001	11,996	-	1	1	7	45	125	362	678	1,453	3,711	5,613	-
2000	11,178	-	1	1	13	31	132	323	622	1,476	3,419	5,160	-
1999	10,480	1	2	-	5	30	118	317	586	1,427	3,257	4,737	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Essential (primary) hypertension and hypertensive renal disease (I10,I12)													
Hispanic, both sexes													
2005	1,314	1	-	2	3	11	37	82	150	246	386	396	-
2004	1,116	-	1	-	4	9	22	89	121	205	303	362	-
2003	1,053	-	-	-	4	11	23	75	121	204	285	330	-
2002	942	2	-	-	5	9	30	39	123	183	283	268	-
2001	889	-	-	-	3	10	19	60	83	187	271	256	-
2000	821	1	1	-	4	7	27	55	109	162	232	223	-
1999	753	1	1	-	3	8	19	64	82	158	232	185	-
Hispanic, male													
2005	547	1	-	1	3	4	27	46	69	121	162	113	-
2004	492	-	1	-	3	6	11	58	68	104	136	105	-
2003	437	-	-	-	2	5	17	53	73	84	116	87	-
2002	385	1	-	-	1	5	18	22	66	80	113	79	-
2001	365	-	-	-	2	8	16	39	34	84	114	68	-
2000	386	1	-	-	2	5	20	44	65	72	107	70	-
1999	340	-	-	-	2	5	14	35	40	84	99	61	-
Hispanic, female													
2005	767	-	-	1	-	7	10	36	81	125	224	283	-
2004	624	-	-	-	1	3	11	31	53	101	167	257	-
2003	616	-	-	-	2	6	6	22	48	120	169	243	-
2002	557	1	-	-	4	4	12	17	57	103	170	189	-
2001	524	-	-	-	1	2	3	21	49	103	157	188	-
2000	435	-	1	-	2	2	7	11	44	90	125	153	-
1999	413	1	1	-	1	3	5	29	42	74	133	124	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Essential (primary) hypertension and hypertensive renal disease (I10,I12)													
Non-Hispanic 1/, both sexes													
2005	23,543	1	-	1	18	73	369	1,081	1,792	3,051	6,857	10,300	-
2004	21,905	2	-	1	17	94	325	1,045	1,709	2,933	6,505	9,270	4
2003	20,831	1	2	2	18	85	349	953	1,628	2,883	6,350	8,560	-
2002	19,254	3	1	1	16	75	322	881	1,400	2,725	5,840	7,990	-
2001	18,306	-	1	1	20	94	310	868	1,370	2,636	5,706	7,300	-
2000	17,182	-	-	4	13	63	318	815	1,320	2,616	5,373	6,660	-
1999	16,158	2	2	1	10	69	292	734	1,232	2,633	5,073	6,110	-
Non-Hispanic 1/, male													
2005	8,890	-	-	-	9	41	216	685	1,074	1,511	2,753	2,601	-
2004	8,353	1	-	-	10	53	210	653	963	1,422	2,643	2,395	3
2003	7,751	1	2	1	10	51	198	605	911	1,441	2,401	2,130	-
2002	7,232	1	1	1	10	42	200	536	781	1,328	2,285	2,047	-
2001	6,856	-	-	-	14	51	188	528	743	1,288	2,160	1,884	-
2000	6,475	-	-	3	2	34	194	504	746	1,232	2,088	1,672	-
1999	6,123	2	1	1	6	42	179	447	689	1,285	1,958	1,513	-
Non-Hispanic 1/, female													
2005	14,653	1	-	1	9	32	153	396	718	1,540	4,104	7,699	-
2004	13,552	1	-	1	7	41	115	392	746	1,511	3,862	6,875	1
2003	13,080	-	-	1	8	34	151	348	717	1,442	3,949	6,430	-
2002	12,022	2	-	-	6	33	122	345	619	1,397	3,555	5,943	-
2001	11,450	-	1	1	6	43	122	340	627	1,348	3,546	5,416	-
2000	10,707	-	-	1	11	29	124	311	574	1,384	3,285	4,988	-
1999	10,035	-	1	-	4	27	113	287	543	1,348	3,115	4,597	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Essential (primary) hypertension and hypertensive renal disease (I10,I12)													
Non-Hispanic white, both sexes													
2005 2/	17,949	-	-	1	6	19	160	523	1,059	1,978	5,415	8,788	-
2004 2/	16,569	-	-	-	9	35	139	510	960	1,940	5,098	7,875	3
2003 2/	15,878	-	1	1	8	32	165	493	918	1,862	5,026	7,372	-
2002	14,617	2	1	1	9	26	126	449	805	1,838	4,570	6,790	-
2001	13,837	-	1	1	11	35	121	404	776	1,747	4,499	6,242	-
2000	12,917	-	-	3	3	25	132	381	736	1,744	4,252	5,641	-
1999	12,048	1	-	-	5	28	112	343	686	1,788	3,940	5,145	-
Non-Hispanic white, male													
2005 2/	6,474	-	-	-	4	11	104	353	658	980	2,163	2,201	-
2004 2/	6,147	-	-	-	7	23	100	336	572	968	2,075	2,064	2
2003 2/	5,673	-	1	-	5	16	102	319	527	976	1,916	1,811	-
2002	5,325	1	1	1	7	16	82	290	477	920	1,784	1,746	-
2001	4,970	-	-	-	7	20	77	253	439	873	1,701	1,600	-
2000	4,721	-	-	3	1	17	89	255	434	852	1,671	1,399	-
1999	4,395	1	-	-	2	22	77	220	410	897	1,526	1,240	-
Non-Hispanic white, female													
2005 2/	11,475	-	-	1	2	8	56	170	401	998	3,252	6,587	-
2004 2/	10,422	-	-	-	2	12	39	174	388	972	3,023	5,811	1
2003 2/	10,205	-	-	1	3	16	63	174	391	886	3,110	5,561	-
2002	9,292	1	-	-	2	10	44	159	328	918	2,786	5,044	-
2001	8,867	-	1	1	4	15	44	151	337	874	2,798	4,642	-
2000	8,196	-	-	-	2	8	43	126	302	892	2,581	4,242	-
1999	7,653	-	-	-	3	6	35	123	276	891	2,414	3,905	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Essential (primary) hypertension and hypertensive renal disease (I10,I12)													
Non-Hispanic black, both sexes													
2005 2/	4,906	1	-	-	11	51	194	527	684	948	1,226	1,264	-
2004 2/	4,709	2	-	1	7	55	177	497	685	890	1,214	1,180	1
2003 2/	4,445	1	1	-	10	51	174	441	664	918	1,162	1,023	-
2002	4,165	1	-	-	7	44	189	413	550	795	1,133	1,033	-
2001	4,012	-	-	-	9	55	177	439	553	789	1,056	934	-
2000	3,862	-	-	1	10	35	176	405	555	801	980	899	-
1999	3,708	1	1	1	5	37	175	370	506	764	1,006	842	-
Non-Hispanic black, male													
2005 2/	2,108	-	-	-	5	28	103	307	395	465	496	309	-
2004 2/	1,945	1	-	-	3	29	103	291	361	404	488	264	1
2003 2/	1,875	1	1	-	5	33	92	278	359	424	419	263	-
2002	1,701	-	-	-	3	24	112	234	274	359	442	253	-
2001	1,698	-	-	-	7	29	102	263	284	374	399	240	-
2000	1,576	-	-	-	1	16	98	229	301	344	358	229	-
1999	1,550	1	1	1	4	17	97	218	256	348	381	226	-
Non-Hispanic black, female													
2005 2/	2,798	1	-	-	6	23	91	220	289	483	730	955	-
2004 2/	2,764	1	-	1	4	26	74	206	324	486	726	916	-
2003 2/	2,570	-	-	-	5	18	82	163	305	494	743	760	-
2002	2,464	1	-	-	4	20	77	179	276	436	691	780	-
2001	2,314	-	-	-	2	26	75	176	269	415	657	694	-
2000	2,286	-	-	1	9	19	78	176	254	457	622	670	-
1999	2,158	-	-	-	1	20	78	152	250	416	625	616	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Essential (primary) hypertension and hypertensive renal disease (I10,I12)													
Origin not stated 3/, both sexes													
2005	45	-	-	-	1	-	-	3	12	9	13	7	-
2004	55	-	-	-	-	-	1	4	9	14	11	16	-
2003	56	-	-	-	-	2	-	3	6	17	14	14	-
2002	65	-	-	-	-	1	2	3	3	14	15	27	-
2001	55	-	-	-	-	-	2	3	9	9	16	16	-
2000	70	-	-	-	-	-	3	8	11	6	18	24	-
1999	57	-	-	-	-	1	2	1	4	8	19	22	-
Origin not stated 3/, male													
2005	21	-	-	-	-	-	-	2	8	7	2	2	-
2004	24	-	-	-	-	-	1	3	5	9	3	3	-
2003	25	-	-	-	-	2	-	2	5	10	5	1	-
2002	30	-	-	-	-	-	1	2	3	11	7	6	-
2001	33	-	-	-	-	-	2	2	7	7	8	7	-
2000	34	-	-	-	-	-	2	7	7	4	9	5	-
1999	25	-	-	-	-	1	2	-	3	3	10	6	-
Origin not stated 3/, female													
2005	24	-	-	-	1	-	-	1	4	2	11	5	-
2004	31	-	-	-	-	-	-	1	4	5	8	13	-
2003	31	-	-	-	-	-	-	1	1	7	9	13	-
2002	35	-	-	-	-	1	1	1	-	3	8	21	-
2001	22	-	-	-	-	-	-	1	2	2	8	9	-
2000	36	-	-	-	-	-	1	1	4	2	9	19	-
1999	32	-	-	-	-	-	-	1	1	5	9	16	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Cerebrovascular diseases (I60-I69)													
All origins, both sexes													
2005	143,579	126	62	95	196	546	2,260	6,381	10,028	18,839	46,859	58,183	4
2004	150,074	127	41	77	211	567	2,361	6,181	9,966	19,901	50,092	60,545	5
2003	157,689	101	46	69	221	583	2,460	6,127	9,946	20,708	52,847	64,579	2
2002	162,672	117	53	91	171	567	2,425	6,055	9,897	21,992	54,889	66,412	3
2001	163,538	108	54	80	196	601	2,491	5,910	9,608	22,598	55,822	66,066	4
2000	167,661	124	47	76	199	602	2,599	6,011	9,956	23,649	57,020	67,376	2
1999	167,366	103	46	65	182	580	2,574	5,563	9,652	24,092	57,427	67,080	2
All origins, male													
2005	56,586	73	41	55	94	301	1,148	3,446	5,636	9,688	19,688	16,412	4
2004	58,800	71	23	40	116	284	1,240	3,409	5,523	10,207	21,025	16,859	3
2003	61,426	57	25	41	116	314	1,283	3,350	5,478	10,668	22,234	17,859	1
2002	62,622	67	30	47	103	284	1,178	3,278	5,456	11,207	22,655	18,316	1
2001	63,177	63	25	42	103	311	1,267	3,205	5,277	11,649	23,301	17,930	4
2000	64,769	74	19	33	106	308	1,297	3,235	5,496	12,042	23,946	18,212	1
1999	64,485	65	22	34	95	299	1,308	2,996	5,305	12,220	24,088	18,052	1
All origins, female													
2005	86,993	53	21	40	102	245	1,112	2,935	4,392	9,151	27,171	41,771	-
2004	91,274	56	18	37	95	283	1,121	2,772	4,443	9,694	29,067	43,686	2
2003	96,263	44	21	28	105	269	1,177	2,777	4,468	10,040	30,613	46,720	1
2002	100,050	50	23	44	68	283	1,247	2,777	4,441	10,785	32,234	48,096	2
2001	100,361	45	29	38	93	290	1,224	2,705	4,331	10,949	32,521	48,136	-
2000	102,892	50	28	43	93	294	1,302	2,776	4,460	11,607	33,074	49,164	1
1999	102,881	38	24	31	87	281	1,266	2,567	4,347	11,872	33,339	49,028	1

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Cerebrovascular diseases (I60-I69)													
Hispanic, both sexes													
2005	6,830	24	15	19	44	82	296	639	795	1,181	2,009	1,726	-
2004	6,781	20	3	10	36	93	304	590	776	1,232	2,006	1,709	2
2003	6,658	17	8	14	28	89	267	569	736	1,202	2,029	1,699	-
2002	6,451	20	12	20	29	93	256	560	708	1,215	1,918	1,620	-
2001	6,416	17	10	16	40	99	275	539	654	1,206	1,892	1,668	-
2000	6,187	16	6	8	31	108	267	477	681	1,219	1,742	1,632	-
1999	5,907	16	7	9	32	102	241	464	617	1,211	1,672	1,536	-
Hispanic, male													
2005	3,188	14	8	9	25	50	177	384	458	627	880	556	-
2004	3,207	11	3	5	20	61	190	354	454	647	904	556	2
2003	3,070	11	7	8	15	51	162	345	431	617	879	544	-
2002	3,003	9	5	7	17	61	154	340	421	652	825	512	-
2001	2,982	9	4	10	26	60	151	315	378	637	864	528	-
2000	2,865	8	-	3	17	65	145	280	392	645	799	511	-
1999	2,809	8	1	6	21	62	146	285	354	641	758	527	-
Hispanic, female													
2005	3,642	10	7	10	19	32	119	255	337	554	1,129	1,170	-
2004	3,574	9	-	5	16	32	114	236	322	585	1,102	1,153	-
2003	3,588	6	1	6	13	38	105	224	305	585	1,150	1,155	-
2002	3,448	11	7	13	12	32	102	220	287	563	1,093	1,108	-
2001	3,434	8	6	6	14	39	124	224	276	569	1,028	1,140	-
2000	3,322	8	6	5	14	43	122	197	289	574	943	1,121	-
1999	3,098	8	6	3	11	40	95	179	263	570	914	1,009	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Cerebrovascular diseases (I60-I69)													
Non-Hispanic 1/, both sexes													
2005	136,548	102	47	75	152	464	1,960	5,732	9,222	17,620	44,778	56,392	4
2004	143,044	106	38	67	175	473	2,051	5,571	9,163	18,631	48,009	58,758	2
2003	150,719	82	38	55	192	491	2,182	5,536	9,179	19,458	50,742	62,763	1
2002	155,852	96	41	71	142	474	2,161	5,477	9,160	20,731	52,850	64,646	3
2001	156,741	91	43	63	156	498	2,211	5,350	8,916	21,346	53,810	64,253	4
2000	161,057	108	41	67	167	493	2,323	5,509	9,238	22,373	55,129	65,608	1
1999	161,057	87	39	56	150	477	2,323	5,071	8,991	22,814	55,643	65,404	2
Non-Hispanic 1/, male													
2005	53,319	59	33	45	69	251	970	3,057	5,172	9,044	18,773	15,842	4
2004	55,489	60	20	35	96	223	1,048	3,041	5,053	9,541	20,086	16,286	-
2003	58,226	45	18	33	100	261	1,115	2,993	5,026	10,023	21,322	17,289	1
2002	59,470	57	25	40	86	223	1,019	2,926	5,018	10,532	21,773	17,770	1
2001	60,030	54	21	32	77	247	1,111	2,882	4,872	10,989	22,377	17,364	4
2000	61,715	66	19	30	89	243	1,145	2,936	5,079	11,364	23,084	17,660	-
1999	61,499	57	21	28	74	237	1,156	2,697	4,922	11,542	23,275	17,489	1
Non-Hispanic 1/, female													
2005	83,229	43	14	30	83	213	990	2,675	4,050	8,576	26,005	40,550	-
2004	87,555	46	18	32	79	250	1,003	2,530	4,110	9,090	27,923	42,472	2
2003	92,493	37	20	22	92	230	1,067	2,543	4,153	9,435	29,420	45,474	-
2002	96,382	39	16	31	56	251	1,142	2,551	4,142	10,199	31,077	46,876	2
2001	96,711	37	22	31	79	251	1,100	2,468	4,044	10,357	31,433	46,889	-
2000	99,342	42	22	37	78	250	1,178	2,573	4,159	11,009	32,045	47,948	1
1999	99,558	30	18	28	76	240	1,167	2,374	4,069	11,272	32,368	47,915	1

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Cerebrovascular diseases (I60-I69)													
Non-Hispanic white, both sexes													
2005 2/	115,074	56	30	45	92	274	1,158	3,461	6,351	13,674	38,830	51,103	-
2004 2/	121,065	59	25	48	117	280	1,256	3,355	6,321	14,563	41,670	53,371	-
2003 2/	127,969	46	23	37	124	298	1,239	3,294	6,293	15,259	44,277	57,078	1
2002	133,118	51	29	44	101	285	1,288	3,320	6,432	16,355	46,327	58,884	2
2001	133,879	46	28	42	98	320	1,333	3,200	6,208	16,838	47,304	58,461	1
2000	138,210	59	25	46	118	313	1,378	3,433	6,555	17,989	48,556	59,738	-
1999	138,716	55	27	40	105	301	1,406	3,143	6,311	18,332	49,178	59,818	-
Non-Hispanic white, male													
2005 2/	44,038	33	20	26	40	147	581	1,868	3,534	7,032	16,379	14,378	-
2004 2/	46,087	34	15	24	63	137	653	1,841	3,447	7,505	17,553	14,815	-
2003 2/	48,549	22	8	23	72	159	637	1,813	3,427	7,947	18,699	15,741	1
2002	49,908	32	18	23	62	136	617	1,797	3,511	8,368	19,181	16,162	1
2001	50,382	27	14	24	55	161	684	1,745	3,400	8,753	19,762	15,756	1
2000	52,000	38	14	23	67	161	694	1,856	3,605	9,176	20,379	15,987	-
1999	51,968	36	15	21	51	151	712	1,662	3,479	9,337	20,618	15,886	-
Non-Hispanic white, female													
2005 2/	71,036	23	10	19	52	127	577	1,593	2,817	6,642	22,451	36,725	-
2004 2/	74,978	25	10	24	54	143	603	1,514	2,874	7,058	24,117	38,556	-
2003 2/	79,420	24	15	14	52	139	602	1,481	2,866	7,312	25,578	41,337	-
2002	83,210	19	11	21	39	149	671	1,523	2,921	7,987	27,146	42,722	1
2001	83,497	19	14	18	43	159	649	1,455	2,808	8,085	27,542	42,705	-
2000	86,210	21	11	23	51	152	684	1,577	2,950	8,813	28,177	43,751	-
1999	86,748	19	12	19	54	150	694	1,481	2,832	8,995	28,560	43,932	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Cerebrovascular diseases (I60-I69)													
Non-Hispanic black, both sexes													
2005 2/	17,378	39	14	23	51	163	683	1,986	2,420	3,194	4,691	4,110	4
2004 2/	17,980	39	12	17	48	173	698	1,905	2,420	3,362	5,044	4,260	2
2003 2/	18,649	31	14	16	58	167	786	1,949	2,449	3,411	5,176	4,592	-
2002	18,691	36	10	23	38	166	741	1,874	2,303	3,600	5,277	4,622	1
2001	18,844	42	12	14	51	153	758	1,886	2,289	3,684	5,257	4,695	3
2000	19,053	46	13	15	42	158	829	1,819	2,265	3,660	5,344	4,861	1
1999	18,731	27	10	15	40	149	808	1,662	2,269	3,792	5,314	4,644	1
Non-Hispanic black, male													
2005 2/	7,448	22	10	14	24	89	329	1,039	1,411	1,636	1,821	1,049	4
2004 2/	7,586	23	5	10	30	77	344	1,012	1,381	1,685	1,959	1,060	-
2003 2/	7,798	19	9	9	25	86	385	1,031	1,363	1,673	2,071	1,127	-
2002	7,754	19	6	15	22	72	331	986	1,288	1,826	2,031	1,158	-
2001	7,828	26	5	5	20	75	355	998	1,252	1,844	2,068	1,177	3
2000	7,945	27	3	5	19	70	394	955	1,240	1,817	2,155	1,260	-
1999	7,823	17	4	7	19	72	387	900	1,235	1,857	2,100	1,225	-
Non-Hispanic black, female													
2005 2/	9,930	17	4	9	27	74	354	947	1,009	1,558	2,870	3,061	-
2004 2/	10,394	16	7	7	18	96	354	893	1,039	1,677	3,085	3,200	2
2003 2/	10,851	12	5	7	33	81	401	918	1,086	1,738	3,105	3,465	-
2002	10,937	17	4	8	16	94	410	888	1,015	1,774	3,246	3,464	1
2001	11,016	16	7	9	31	78	403	888	1,037	1,840	3,189	3,518	-
2000	11,108	19	10	10	23	88	435	864	1,025	1,843	3,189	3,601	1
1999	10,908	10	6	8	21	77	421	762	1,034	1,935	3,214	3,419	1

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Cerebrovascular diseases (I60-I69)													
Origin not stated 3/, both sexes													
2005	201	-	-	1	-	-	4	10	11	38	72	65	-
2004	249	1	-	-	-	1	6	20	27	38	77	78	1
2003	312	2	-	-	1	3	11	22	31	48	76	117	1
2002	369	1	-	-	-	-	8	18	29	46	121	146	-
2001	381	-	1	1	-	4	5	21	38	46	120	145	-
2000	417	-	-	1	1	1	9	25	37	57	149	136	1
1999	402	-	-	-	-	1	10	28	44	67	112	140	-
Origin not stated 3/, male													
2005	79	-	-	1	-	-	1	5	6	17	35	14	-
2004	104	-	-	-	-	-	2	14	16	19	35	17	1
2003	130	1	-	-	1	2	6	12	21	28	33	26	-
2002	149	1	-	-	-	-	5	12	17	23	57	34	-
2001	165	-	-	-	-	4	5	8	27	23	60	38	-
2000	189	-	-	-	-	-	7	19	25	33	63	41	1
1999	177	-	-	-	-	-	6	14	29	37	55	36	-
Origin not stated 3/, female													
2005	122	-	-	-	-	-	3	5	5	21	37	51	-
2004	145	1	-	-	-	1	4	6	11	19	42	61	-
2003	182	1	-	-	-	1	5	10	10	20	43	91	1
2002	220	-	-	-	-	-	3	6	12	23	64	112	-
2001	216	-	1	1	-	-	-	13	11	23	60	107	-
2000	228	-	-	1	1	1	2	6	12	24	86	95	-
1999	225	-	-	-	-	1	4	14	15	30	57	104	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Atherosclerosis (I70)													
All origins, both sexes													
2005	11,841	-	1	-	2	6	31	193	440	1,008	3,387	6,773	-
2004	11,861	-	1	-	2	6	43	153	459	1,132	3,406	6,659	-
2003	13,053	3	-	-	3	7	47	190	466	1,211	3,837	7,288	1
2002	13,821	4	-	-	3	9	50	182	487	1,300	4,062	7,723	1
2001	14,086	1	1	1	1	3	40	186	473	1,335	4,156	7,888	1
2000	14,393	2	-	-	3	8	48	176	451	1,423	4,277	8,005	-
1999	14,979	2	-	-	2	8	45	185	486	1,476	4,385	8,390	-
All origins, male													
2005	4,472	-	1	-	1	3	20	134	269	560	1,532	1,952	-
2004	4,510	-	-	-	2	3	25	104	275	619	1,549	1,933	-
2003	4,859	2	-	-	2	5	35	131	298	649	1,734	2,002	1
2002	5,171	1	-	-	2	5	30	119	305	728	1,803	2,177	1
2001	5,143	1	1	1	1	2	26	118	286	743	1,791	2,172	1
2000	5,357	-	-	-	2	6	35	119	294	800	1,935	2,166	-
1999	5,443	1	-	-	-	5	32	121	307	795	1,927	2,255	-
All origins, female													
2005	7,369	-	-	-	1	3	11	59	171	448	1,855	4,821	-
2004	7,351	-	1	-	-	3	18	49	184	513	1,857	4,726	-
2003	8,194	1	-	-	1	2	12	59	168	562	2,103	5,286	-
2002	8,650	3	-	-	1	4	20	63	182	572	2,259	5,546	-
2001	8,943	-	-	-	-	1	14	68	187	592	2,365	5,716	-
2000	9,036	2	-	-	1	2	13	57	157	623	2,342	5,839	-
1999	9,536	1	-	-	2	3	13	64	179	681	2,458	6,135	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Atherosclerosis (I70)													
Hispanic, both sexes													
2005	397	-	-	-	1	-	-	8	22	51	129	186	-
2004	369	-	-	-	-	-	3	5	22	55	127	157	-
2003	379	1	-	-	-	1	3	7	24	56	121	166	-
2002	416	1	-	-	2	-	2	9	20	50	136	196	-
2001	416	-	-	-	-	-	-	14	19	46	128	209	-
2000	414	-	-	-	-	-	2	9	16	63	124	200	-
1999	410	-	-	-	-	-	3	14	25	54	130	184	-
Hispanic, male													
2005	175	-	-	-	-	-	-	3	13	31	61	67	-
2004	172	-	-	-	-	-	2	3	11	29	64	63	-
2003	182	-	-	-	-	-	3	4	17	33	63	62	-
2002	175	1	-	-	2	-	1	6	12	27	65	61	-
2001	172	-	-	-	-	-	-	9	11	24	61	67	-
2000	173	-	-	-	-	-	2	7	8	39	65	52	-
1999	185	-	-	-	-	-	1	7	16	31	60	70	-
Hispanic, female													
2005	222	-	-	-	1	-	-	5	9	20	68	119	-
2004	197	-	-	-	-	-	1	2	11	26	63	94	-
2003	197	1	-	-	-	1	-	3	7	23	58	104	-
2002	241	-	-	-	-	-	1	3	8	23	71	135	-
2001	244	-	-	-	-	-	-	5	8	22	67	142	-
2000	241	-	-	-	-	-	-	2	8	24	59	148	-
1999	225	-	-	-	-	-	2	7	9	23	70	114	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Atherosclerosis (I70)													
Non-Hispanic 1/, both sexes													
2005	11,431	-	1	-	1	6	31	185	417	957	3,255	6,578	-
2004	11,478	-	1	-	2	6	40	148	435	1,075	3,275	6,496	-
2003	12,627	2	-	-	3	6	44	182	440	1,152	3,696	7,102	-
2002	13,369	3	-	-	1	9	48	173	466	1,248	3,911	7,509	1
2001	13,626	1	1	1	1	3	40	172	452	1,285	4,011	7,658	1
2000	13,945	2	-	-	3	8	46	167	434	1,357	4,145	7,783	-
1999	14,519	2	-	-	2	8	42	169	457	1,416	4,242	8,181	-
Non-Hispanic 1/, male													
2005	4,295	-	1	-	1	3	20	131	256	529	1,469	1,885	-
2004	4,331	-	-	-	2	3	23	101	262	589	1,482	1,869	-
2003	4,660	2	-	-	2	5	32	126	279	615	1,663	1,936	-
2002	4,984	-	-	-	-	5	29	113	292	700	1,733	2,111	1
2001	4,952	1	1	1	1	2	26	109	274	718	1,721	2,097	1
2000	5,166	-	-	-	2	6	33	112	285	759	1,865	2,104	-
1999	5,236	1	-	-	-	5	31	112	287	761	1,861	2,178	-
Non-Hispanic 1/, female													
2005	7,136	-	-	-	-	3	11	54	161	428	1,786	4,693	-
2004	7,147	-	1	-	-	3	17	47	173	486	1,793	4,627	-
2003	7,967	-	-	-	1	1	12	56	161	537	2,033	5,166	-
2002	8,385	3	-	-	1	4	19	60	174	548	2,178	5,398	-
2001	8,674	-	-	-	-	1	14	63	178	567	2,290	5,561	-
2000	8,779	2	-	-	1	2	13	55	149	598	2,280	5,679	-
1999	9,283	1	-	-	2	3	11	57	170	655	2,381	6,003	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Atherosclerosis (I70)													
Non-Hispanic white, both sexes													
2005 2/	10,443	-	-	-	-	5	26	140	345	806	2,959	6,162	-
2004 2/	10,514	-	1	-	2	5	28	120	366	921	3,007	6,064	-
2003 2/	11,475	-	-	-	2	4	36	141	359	978	3,349	6,606	-
2002	12,191	2	-	-	1	6	38	121	376	1,082	3,518	7,046	1
2001	12,517	1	1	1	-	2	33	137	361	1,119	3,675	7,186	1
2000	12,740	2	-	-	3	5	34	114	349	1,181	3,782	7,270	-
1999	13,286	1	-	-	2	6	30	135	374	1,241	3,880	7,617	-
Non-Hispanic white, male													
2005 2/	3,920	-	-	-	-	2	17	104	213	449	1,348	1,787	-
2004 2/	3,980	-	-	-	2	3	18	88	221	518	1,361	1,769	-
2003 2/	4,181	-	-	-	1	4	25	98	230	520	1,495	1,808	-
2002	4,508	-	-	-	-	4	23	79	237	613	1,561	1,990	1
2001	4,511	1	1	1	-	2	21	85	225	631	1,570	1,973	1
2000	4,688	-	-	-	2	5	26	79	238	673	1,716	1,949	-
1999	4,727	1	-	-	-	4	23	93	235	667	1,690	2,014	-
Non-Hispanic white, female													
2005 2/	6,523	-	-	-	-	3	9	36	132	357	1,611	4,375	-
2004 2/	6,534	-	1	-	-	2	10	32	145	403	1,646	4,295	-
2003 2/	7,294	-	-	-	1	-	11	43	129	458	1,854	4,798	-
2002	7,683	2	-	-	1	2	15	42	139	469	1,957	5,056	-
2001	8,006	-	-	-	-	-	12	52	136	488	2,105	5,213	-
2000	8,052	2	-	-	1	-	8	35	111	508	2,066	5,321	-
1999	8,559	-	-	-	2	2	7	42	139	574	2,190	5,603	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Atherosclerosis (I70)													
Non-Hispanic black, both sexes													
2005 2/	840	-	1	-	1	1	5	38	62	130	237	365	-
2004 2/	823	-	-	-	-	1	11	26	63	126	236	360	-
2003 2/	981	2	-	-	1	2	5	38	72	152	296	413	-
2002	1,009	1	-	-	-	3	9	46	77	134	349	390	-
2001	957	-	-	-	1	1	6	31	84	145	285	404	-
2000	1,049	-	-	-	-	3	12	47	79	147	320	441	-
1999	1,061	1	-	-	-	2	9	29	76	150	312	482	-
Non-Hispanic black, male													
2005 2/	310	-	1	-	1	1	3	21	37	70	95	81	-
2004 2/	296	-	-	-	-	-	4	12	37	60	106	77	-
2003 2/	400	2	-	-	1	1	4	26	44	83	140	99	-
2002	409	-	-	-	-	1	6	30	49	70	153	100	-
2001	365	-	-	-	1	-	4	21	43	74	124	98	-
2000	412	-	-	-	-	1	7	28	43	74	131	128	-
1999	430	-	-	-	-	1	6	15	46	80	142	140	-
Non-Hispanic black, female													
2005 2/	530	-	-	-	-	-	2	17	25	60	142	284	-
2004 2/	527	-	-	-	-	1	7	14	26	66	130	283	-
2003 2/	581	-	-	-	-	1	1	12	28	69	156	314	-
2002	600	1	-	-	-	2	3	16	28	64	196	290	-
2001	592	-	-	-	-	1	2	10	41	71	161	306	-
2000	637	-	-	-	-	2	5	19	36	73	189	313	-
1999	631	1	-	-	-	1	3	14	30	70	170	342	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Atherosclerosis (I70)													
Origin not stated 3/, both sexes													
2005	13	-	-	-	-	-	-	-	1	-	3	9	-
2004	14	-	-	-	-	-	-	-	2	2	4	6	-
2003	47	-	-	-	-	-	-	1	2	3	20	20	1
2002	36	-	-	-	-	-	-	-	1	2	15	18	-
2001	44	-	-	-	-	-	-	-	2	4	17	21	-
2000	34	-	-	-	-	-	-	-	1	3	8	22	-
1999	50	-	-	-	-	-	-	2	4	6	13	25	-
Origin not stated 3/, male													
2005	2	-	-	-	-	-	-	-	-	-	2	-	-
2004	7	-	-	-	-	-	-	-	2	1	3	1	-
2003	17	-	-	-	-	-	-	1	2	1	8	4	1
2002	12	-	-	-	-	-	-	-	1	1	5	5	-
2001	19	-	-	-	-	-	-	-	1	1	9	8	-
2000	18	-	-	-	-	-	-	-	1	2	5	10	-
1999	22	-	-	-	-	-	-	2	4	3	6	7	-
Origin not stated 3/, female													
2005	11	-	-	-	-	-	-	-	1	-	1	9	-
2004	7	-	-	-	-	-	-	-	-	1	1	5	-
2003	30	-	-	-	-	-	-	-	-	2	12	16	-
2002	24	-	-	-	-	-	-	-	-	1	10	13	-
2001	25	-	-	-	-	-	-	-	1	3	8	13	-
2000	16	-	-	-	-	-	-	-	-	1	3	12	-
1999	28	-	-	-	-	-	-	-	-	3	7	18	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Other diseases of circulatory system (I71-I78)													
All origins, both sexes													
2005	23,617	17	3	17	52	156	467	1,085	2,266	4,484	8,167	6,903	-
2004	23,693	14	4	13	61	167	461	1,125	2,107	4,632	8,341	6,768	-
2003	24,672	11	-	12	68	151	451	1,199	2,270	4,934	8,694	6,882	-
2002	24,927	14	4	11	66	160	445	1,070	2,286	5,310	8,745	6,816	-
2001	25,318	12	4	11	61	170	498	1,024	2,195	5,345	9,032	6,965	1
2000	26,036	16	5	11	59	154	448	1,038	2,236	5,926	9,222	6,920	1
1999	25,809	12	2	11	49	141	478	971	2,210	5,808	9,264	6,863	-
All origins, male													
2005	12,423	11	3	13	37	102	322	749	1,533	2,796	4,320	2,537	-
2004	12,306	7	2	8	45	113	308	780	1,392	2,818	4,322	2,511	-
2003	13,217	3	-	7	45	103	306	815	1,517	3,092	4,732	2,597	-
2002	13,309	6	4	5	43	101	291	732	1,526	3,307	4,690	2,604	-
2001	13,396	8	2	3	40	115	335	688	1,454	3,348	4,839	2,563	1
2000	14,084	6	2	7	33	103	276	699	1,530	3,771	5,019	2,637	1
1999	13,794	9	-	7	32	102	296	657	1,519	3,681	5,015	2,476	-
All origins, female													
2005	11,194	6	-	4	15	54	145	336	733	1,688	3,847	4,366	-
2004	11,387	7	2	5	16	54	153	345	715	1,814	4,019	4,257	-
2003	11,455	8	-	5	23	48	145	384	753	1,842	3,962	4,285	-
2002	11,618	8	-	6	23	59	154	338	760	2,003	4,055	4,212	-
2001	11,922	4	2	8	21	55	163	336	741	1,997	4,193	4,402	-
2000	11,952	10	3	4	26	51	172	339	706	2,155	4,203	4,283	-
1999	12,015	3	2	4	17	39	182	314	691	2,127	4,249	4,387	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Other diseases of circulatory system (I71-I78)													
Hispanic, both sexes													
2005	914	2	2	4	13	18	49	75	97	176	304	174	-
2004	846	3	-	2	10	20	41	80	115	177	236	162	-
2003	878	2	-	1	9	22	39	61	96	172	271	205	-
2002	847	1	-	1	12	22	44	62	117	175	246	167	-
2001	754	2	1	2	7	19	37	73	88	156	216	152	1
2000	818	3	1	1	9	24	38	56	99	217	215	155	-
1999	729	2	-	2	9	12	33	38	94	170	211	158	-
Hispanic, male													
2005	558	1	2	3	8	14	33	55	74	111	172	85	-
2004	494	2	-	1	8	12	30	60	77	116	124	64	-
2003	510	1	-	-	5	15	28	42	73	103	153	90	-
2002	463	-	-	-	8	14	31	41	66	104	138	61	-
2001	424	1	1	-	4	14	26	46	55	100	119	57	1
2000	452	1	-	1	4	14	22	33	64	122	126	65	-
1999	416	1	-	-	7	6	22	25	59	101	119	76	-
Hispanic, female													
2005	356	1	-	1	5	4	16	20	23	65	132	89	-
2004	352	1	-	1	2	8	11	20	38	61	112	98	-
2003	368	1	-	1	4	7	11	19	23	69	118	115	-
2002	384	1	-	1	4	8	13	21	51	71	108	106	-
2001	330	1	-	2	3	5	11	27	33	56	97	95	-
2000	366	2	1	-	5	10	16	23	35	95	89	90	-
1999	313	1	-	2	2	6	11	13	35	69	92	82	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Other diseases of circulatory system (I71-I78)													
Non-Hispanic 1/, both sexes													
2005	22,661	15	1	13	39	138	417	1,008	2,163	4,302	7,846	6,719	-
2004	22,800	11	4	11	51	147	420	1,039	1,987	4,446	8,090	6,594	-
2003	23,739	9	-	11	59	129	410	1,134	2,171	4,746	8,403	6,667	-
2002	24,005	13	4	10	54	137	396	1,006	2,159	5,120	8,475	6,631	-
2001	24,489	10	3	9	54	149	460	946	2,099	5,170	8,795	6,794	-
2000	25,147	13	4	10	50	129	405	979	2,129	5,695	8,985	6,747	1
1999	25,023	10	2	9	40	129	445	932	2,109	5,619	9,037	6,691	-
Non-Hispanic 1/, male													
2005	11,832	10	1	10	29	88	288	693	1,453	2,679	4,134	2,447	-
2004	11,786	5	2	7	37	101	278	716	1,310	2,696	4,191	2,443	-
2003	12,679	2	-	7	40	88	277	772	1,441	2,979	4,569	2,504	-
2002	12,809	6	4	5	35	86	257	690	1,455	3,194	4,540	2,537	-
2001	12,926	7	1	3	36	99	308	638	1,392	3,234	4,708	2,500	-
2000	13,593	5	2	6	29	89	249	663	1,461	3,640	4,883	2,565	1
1999	13,351	8	-	7	25	96	274	632	1,456	3,567	4,889	2,397	-
Non-Hispanic 1/, female													
2005	10,829	5	-	3	10	50	129	315	710	1,623	3,712	4,272	-
2004	11,014	6	2	4	14	46	142	323	677	1,750	3,899	4,151	-
2003	11,060	7	-	4	19	41	133	362	730	1,767	3,834	4,163	-
2002	11,196	7	-	5	19	51	139	316	704	1,926	3,935	4,094	-
2001	11,563	3	2	6	18	50	152	308	707	1,936	4,087	4,294	-
2000	11,554	8	2	4	21	40	156	316	668	2,055	4,102	4,182	-
1999	11,672	2	2	2	15	33	171	300	653	2,052	4,148	4,294	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Other diseases of circulatory system (I71-I78)													
Non-Hispanic white, both sexes													
2005 2/	19,572	9	1	10	31	84	266	694	1,750	3,688	6,982	6,057	-
2004 2/	19,775	8	3	9	40	92	279	750	1,560	3,864	7,235	5,935	-
2003 2/	20,756	4	-	7	41	85	266	798	1,787	4,158	7,567	6,043	-
2002	21,051	10	4	4	37	86	272	702	1,758	4,485	7,684	6,009	-
2001	21,439	8	2	8	40	108	303	682	1,702	4,539	7,908	6,139	-
2000	22,113	11	3	6	35	88	269	699	1,734	5,046	8,133	6,089	-
1999	22,191	8	2	7	33	94	323	690	1,773	5,000	8,210	6,051	-
Non-Hispanic white, male													
2005 2/	10,261	5	1	8	21	52	174	481	1,200	2,346	3,722	2,251	-
2004 2/	10,308	4	2	5	28	64	188	528	1,040	2,390	3,821	2,238	-
2003 2/	11,188	1	-	4	30	55	189	567	1,206	2,663	4,159	2,314	-
2002	11,337	5	4	2	23	54	180	495	1,198	2,850	4,189	2,337	-
2001	11,437	5	1	3	27	74	202	462	1,158	2,897	4,303	2,305	-
2000	12,138	3	2	5	22	63	167	494	1,225	3,301	4,492	2,364	-
1999	12,014	6	-	5	20	71	204	478	1,261	3,253	4,514	2,202	-
Non-Hispanic white, female													
2005 2/	9,311	4	-	2	10	32	92	213	550	1,342	3,260	3,806	-
2004 2/	9,467	4	1	4	12	28	91	222	520	1,474	3,414	3,697	-
2003 2/	9,568	3	-	3	11	30	77	231	581	1,495	3,408	3,729	-
2002	9,714	5	-	2	14	32	92	207	560	1,635	3,495	3,672	-
2001	10,002	3	1	5	13	34	101	220	544	1,642	3,605	3,834	-
2000	9,975	8	1	1	13	25	102	205	509	1,745	3,641	3,725	-
1999	10,177	2	2	2	13	23	119	212	512	1,747	3,696	3,849	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Other diseases of circulatory system (I71-I78)													
Non-Hispanic black, both sexes													
2005 2/	2,574	4	-	3	7	46	130	271	364	514	695	540	-
2004 2/	2,523	2	1	2	7	46	128	254	364	482	692	545	-
2003 2/	2,500	5	-	3	13	37	132	296	325	492	683	514	-
2002	2,495	3	-	5	12	46	112	272	350	521	653	521	-
2001	2,606	2	1	1	12	39	141	227	348	526	746	563	-
2000	2,569	2	-	4	13	35	120	232	346	540	710	566	1
1999	2,416	2	-	-	7	30	112	212	290	517	689	557	-
Non-Hispanic black, male													
2005 2/	1,264	3	-	2	7	28	96	182	224	265	318	139	-
2004 2/	1,211	1	-	2	7	31	80	165	229	253	284	159	-
2003 2/	1,202	1	-	2	6	29	81	181	200	252	317	133	-
2002	1,196	1	-	2	7	30	71	171	223	268	269	154	-
2001	1,214	2	-	-	7	24	93	149	197	272	324	146	-
2000	1,193	2	-	1	6	22	73	144	198	279	314	153	1
1999	1,102	2	-	-	5	23	61	135	165	266	290	155	-
Non-Hispanic black, female													
2005 2/	1,310	1	-	1	-	18	34	89	140	249	377	401	-
2004 2/	1,312	1	1	-	-	15	48	89	135	229	408	386	-
2003 2/	1,298	4	-	1	7	8	51	115	125	240	366	381	-
2002	1,299	2	-	3	5	16	41	101	127	253	384	367	-
2001	1,392	-	1	1	5	15	48	78	151	254	422	417	-
2000	1,376	-	-	3	7	13	47	88	148	261	396	413	-
1999	1,314	-	-	-	2	7	51	77	125	251	399	402	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Other diseases of circulatory system (I71-I78)													
Origin not stated 3/, both sexes													
2005	42	-	-	-	-	-	1	2	6	6	17	10	-
2004	47	-	-	-	-	-	-	6	5	9	15	12	-
2003	55	-	-	-	-	-	2	4	3	16	20	10	-
2002	75	-	-	-	-	1	5	2	10	15	24	18	-
2001	75	-	-	-	-	2	1	5	8	19	21	19	-
2000	71	-	-	-	-	1	5	3	8	14	22	18	-
1999	57	-	-	-	-	-	-	1	7	19	16	14	-
Origin not stated 3/, male													
2005	33	-	-	-	-	-	1	1	6	6	14	5	-
2004	26	-	-	-	-	-	-	4	5	6	7	4	-
2003	28	-	-	-	-	-	1	1	3	10	10	3	-
2002	37	-	-	-	-	1	3	1	5	9	12	6	-
2001	46	-	-	-	-	2	1	4	7	14	12	6	-
2000	39	-	-	-	-	-	5	3	5	9	10	7	-
1999	27	-	-	-	-	-	-	-	4	13	7	3	-
Origin not stated 3/, female													
2005	9	-	-	-	-	-	-	1	-	-	3	5	-
2004	21	-	-	-	-	-	-	2	-	3	8	8	-
2003	27	-	-	-	-	-	1	3	-	6	10	7	-
2002	38	-	-	-	-	-	2	1	5	6	12	12	-
2001	29	-	-	-	-	-	-	1	1	5	9	13	-
2000	32	-	-	-	-	1	-	-	3	5	12	11	-
1999	30	-	-	-	-	-	-	1	3	6	9	11	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Aortic aneurysm and dissection (I71)													
All origins, both sexes													
2005	13,843	1	-	4	34	114	329	758	1,456	2,870	4,971	3,306	-
2004	13,753	1	1	5	45	126	318	729	1,318	2,932	5,131	3,147	-
2003	14,810	1	-	3	41	112	316	820	1,477	3,271	5,471	3,298	-
2002	14,818	2	-	5	39	110	305	710	1,460	3,482	5,518	3,187	-
2001	15,234	-	2	4	43	117	332	661	1,394	3,517	5,751	3,412	1
2000	15,810	-	-	5	36	107	303	662	1,493	4,012	5,835	3,357	-
1999	15,807	1	1	3	31	93	329	638	1,437	4,001	5,936	3,337	-
All origins, male													
2005	8,168	1	-	4	26	83	265	577	1,063	1,911	2,778	1,460	-
2004	8,140	1	-	3	36	93	246	552	972	1,946	2,859	1,432	-
2003	8,976	-	-	2	31	83	246	616	1,107	2,174	3,192	1,525	-
2002	9,013	-	-	3	32	82	229	540	1,092	2,369	3,185	1,481	-
2001	9,114	-	2	3	31	90	264	506	1,031	2,382	3,313	1,491	1
2000	9,661	-	-	3	23	78	219	492	1,111	2,730	3,443	1,562	-
1999	9,615	1	-	2	20	78	232	486	1,088	2,712	3,501	1,495	-
All origins, female													
2005	5,675	-	-	-	8	31	64	181	393	959	2,193	1,846	-
2004	5,613	-	1	2	9	33	72	177	346	986	2,272	1,715	-
2003	5,834	1	-	1	10	29	70	204	370	1,097	2,279	1,773	-
2002	5,805	2	-	2	7	28	76	170	368	1,113	2,333	1,706	-
2001	6,120	-	-	1	12	27	68	155	363	1,135	2,438	1,921	-
2000	6,149	-	-	2	13	29	84	170	382	1,282	2,392	1,795	-
1999	6,192	-	1	1	11	15	97	152	349	1,289	2,435	1,842	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Aortic aneurysm and dissection (I71)													
Hispanic, both sexes													
2005	477	1	-	-	9	14	33	48	63	96	145	68	-
2004	465	1	-	1	8	16	26	52	74	100	123	64	-
2003	515	-	-	-	6	16	25	36	64	102	171	95	-
2002	431	-	-	-	8	16	24	36	75	89	127	56	-
2001	403	-	1	-	4	14	24	41	47	93	127	51	1
2000	465	-	-	1	8	15	21	41	58	121	137	63	-
1999	384	-	-	1	4	6	23	27	55	95	115	58	-
Hispanic, male													
2005	328	1	-	-	6	11	26	40	54	63	92	35	-
2004	315	1	-	-	7	10	24	44	58	71	71	29	-
2003	333	-	-	-	4	12	19	27	51	68	100	52	-
2002	284	-	-	-	5	12	20	29	51	63	78	26	-
2001	279	-	1	-	2	14	20	31	33	71	78	28	1
2000	294	-	-	1	4	11	14	28	42	76	86	32	-
1999	256	-	-	-	3	4	17	22	41	67	65	37	-
Hispanic, female													
2005	149	-	-	-	3	3	7	8	9	33	53	33	-
2004	150	-	-	1	1	6	2	8	16	29	52	35	-
2003	182	-	-	-	2	4	6	9	13	34	71	43	-
2002	147	-	-	-	3	4	4	7	24	26	49	30	-
2001	124	-	-	-	2	-	4	10	14	22	49	23	-
2000	171	-	-	-	4	4	7	13	16	45	51	31	-
1999	128	-	-	1	1	2	6	5	14	28	50	21	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Aortic aneurysm and dissection (I71)													
Non-Hispanic 1/, both sexes													
2005	13,344	-	-	4	25	100	295	708	1,389	2,770	4,820	3,233	-
2004	13,266	-	1	4	37	110	292	675	1,239	2,826	5,003	3,079	-
2003	14,259	1	-	3	35	96	289	781	1,411	3,159	5,286	3,198	-
2002	14,341	2	-	5	31	93	277	672	1,379	3,384	5,373	3,125	-
2001	14,782	-	1	4	39	102	307	617	1,341	3,408	5,610	3,353	-
2000	15,296	-	-	4	28	91	277	618	1,431	3,881	5,682	3,284	-
1999	15,393	1	1	2	27	87	306	611	1,378	3,893	5,811	3,276	-
Non-Hispanic 1/, male													
2005	7,822	-	-	4	20	72	238	536	1,005	1,844	2,681	1,422	-
2004	7,807	-	-	3	29	83	222	506	909	1,870	2,785	1,400	-
2003	8,622	-	-	2	27	71	226	588	1,054	2,098	3,085	1,471	-
2002	8,701	-	-	3	27	69	207	510	1,036	2,299	3,099	1,451	-
2001	8,798	-	1	3	29	75	243	472	992	2,299	3,225	1,459	-
2000	9,338	-	-	2	19	67	200	461	1,066	2,648	3,349	1,526	-
1999	9,342	1	-	2	17	74	215	464	1,045	2,635	3,432	1,457	-
Non-Hispanic 1/, female													
2005	5,522	-	-	-	5	28	57	172	384	926	2,139	1,811	-
2004	5,459	-	1	1	8	27	70	169	330	956	2,218	1,679	-
2003	5,637	1	-	1	8	25	63	193	357	1,061	2,201	1,727	-
2002	5,640	2	-	2	4	24	70	162	343	1,085	2,274	1,674	-
2001	5,984	-	-	1	10	27	64	145	349	1,109	2,385	1,894	-
2000	5,958	-	-	2	9	24	77	157	365	1,233	2,333	1,758	-
1999	6,051	-	1	-	10	13	91	147	333	1,258	2,379	1,819	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Aortic aneurysm and dissection (I71)													
Non-Hispanic white, both sexes													
2005 2/	11,684	-	-	3	22	63	179	473	1,148	2,443	4,390	2,963	-
2004 2/	11,719	-	-	4	28	69	189	479	985	2,533	4,573	2,859	-
2003 2/	12,712	-	-	1	27	61	188	548	1,178	2,844	4,883	2,982	-
2002	12,831	2	-	2	23	57	192	457	1,156	3,057	4,988	2,897	-
2001	13,217	-	1	4	28	77	203	435	1,117	3,079	5,155	3,118	-
2000	13,722	-	-	1	18	61	182	454	1,201	3,526	5,250	3,029	-
1999	13,934	1	1	1	23	64	217	440	1,195	3,548	5,407	3,037	-
Non-Hispanic white, male													
2005 2/	6,877	-	-	3	17	43	141	370	849	1,659	2,472	1,323	-
2004 2/	6,906	-	-	3	22	52	146	367	730	1,695	2,579	1,312	-
2003 2/	7,731	-	-	-	21	42	155	428	897	1,926	2,877	1,385	-
2002	7,834	-	-	1	19	43	142	355	875	2,105	2,917	1,377	-
2001	7,924	-	1	3	21	58	163	342	841	2,109	3,002	1,384	-
2000	8,488	-	-	1	14	45	134	355	914	2,453	3,136	1,436	-
1999	8,551	1	-	1	14	54	155	340	928	2,451	3,243	1,364	-
Non-Hispanic white, female													
2005 2/	4,807	-	-	-	5	20	38	103	299	784	1,918	1,640	-
2004 2/	4,813	-	-	1	6	17	43	112	255	838	1,994	1,547	-
2003 2/	4,981	-	-	1	6	19	33	120	281	918	2,006	1,597	-
2002	4,997	2	-	1	4	14	50	102	281	952	2,071	1,520	-
2001	5,293	-	-	1	7	19	40	93	276	970	2,153	1,734	-
2000	5,234	-	-	-	4	16	48	99	287	1,073	2,114	1,593	-
1999	5,383	-	1	-	9	10	62	100	267	1,097	2,164	1,673	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Aortic aneurysm and dissection (I71)													
Non-Hispanic black, both sexes													
2005 2/	1,288	-	-	1	3	29	98	201	207	254	307	188	-
2004 2/	1,215	-	1	-	6	33	93	173	215	224	315	155	-
2003 2/	1,220	1	-	1	4	29	94	202	196	242	292	159	-
2002	1,200	-	-	3	5	33	79	192	192	244	285	167	-
2001	1,238	-	-	-	10	23	91	154	189	240	352	179	-
2000	1,262	-	-	3	9	25	86	135	196	279	331	198	-
1999	1,172	-	-	-	4	21	80	152	152	271	303	189	-
Non-Hispanic black, male													
2005 2/	707	-	-	1	3	21	81	140	134	132	140	55	-
2004 2/	688	-	-	-	6	26	68	122	148	130	131	57	-
2003 2/	678	-	-	1	3	26	65	140	132	124	138	49	-
2002	664	-	-	2	5	24	61	138	138	132	119	45	-
2001	659	-	-	-	7	16	69	108	124	134	158	43	-
2000	661	-	-	1	5	19	60	89	123	151	153	60	-
1999	615	-	-	-	3	19	52	109	96	147	124	65	-
Non-Hispanic black, female													
2005 2/	581	-	-	-	-	8	17	61	73	122	167	133	-
2004 2/	527	-	1	-	-	7	25	51	67	94	184	98	-
2003 2/	542	1	-	-	1	3	29	62	64	118	154	110	-
2002	536	-	-	1	-	9	18	54	54	112	166	122	-
2001	579	-	-	-	3	7	22	46	65	106	194	136	-
2000	601	-	-	2	4	6	26	46	73	128	178	138	-
1999	557	-	-	-	1	2	28	43	56	124	179	124	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Aortic aneurysm and dissection (I71)													
Origin not stated 3/, both sexes													
2005	22	-	-	-	-	-	1	2	4	4	6	5	-
2004	22	-	-	-	-	-	-	2	5	6	5	4	-
2003	36	-	-	-	-	-	2	3	2	10	14	5	-
2002	46	-	-	-	-	1	4	2	6	9	18	6	-
2001	49	-	-	-	-	1	1	3	6	16	14	8	-
2000	49	-	-	-	-	1	5	3	4	10	16	10	-
1999	30	-	-	-	-	-	-	-	4	13	10	3	-
Origin not stated 3/, male													
2005	18	-	-	-	-	-	1	1	4	4	5	3	-
2004	18	-	-	-	-	-	-	2	5	5	3	3	-
2003	21	-	-	-	-	-	1	1	2	8	7	2	-
2002	28	-	-	-	-	1	2	1	5	7	8	4	-
2001	37	-	-	-	-	1	1	3	6	12	10	4	-
2000	29	-	-	-	-	-	5	3	3	6	8	4	-
1999	17	-	-	-	-	-	-	-	2	10	4	1	-
Origin not stated 3/, female													
2005	4	-	-	-	-	-	-	1	-	-	1	2	-
2004	4	-	-	-	-	-	-	-	-	1	2	1	-
2003	15	-	-	-	-	-	1	2	-	2	7	3	-
2002	18	-	-	-	-	-	2	1	1	2	10	2	-
2001	12	-	-	-	-	-	-	-	-	4	4	4	-
2000	20	-	-	-	-	1	-	-	1	4	8	6	-
1999	13	-	-	-	-	-	-	-	2	3	6	2	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Other diseases of arteries, arterioles and capillaries (I72-I78)													
All origins, both sexes													
2005	9,774	16	3	13	18	42	138	327	810	1,614	3,196	3,597	-
2004	9,940	13	3	8	16	41	143	396	789	1,700	3,210	3,621	-
2003	9,862	10	-	9	27	39	135	379	793	1,663	3,223	3,584	-
2002	10,109	12	4	6	27	50	140	360	826	1,828	3,227	3,629	-
2001	10,084	12	2	7	18	53	166	363	801	1,828	3,281	3,553	-
2000	10,226	16	5	6	23	47	145	376	743	1,914	3,387	3,563	1
1999	10,002	11	1	8	18	48	149	333	773	1,807	3,328	3,526	-
All origins, male													
2005	4,255	10	3	9	11	19	57	172	470	885	1,542	1,077	-
2004	4,166	6	2	5	9	20	62	228	420	872	1,463	1,079	-
2003	4,241	3	-	5	14	20	60	199	410	918	1,540	1,072	-
2002	4,296	6	4	2	11	19	62	192	434	938	1,505	1,123	-
2001	4,282	8	-	-	9	25	71	182	423	966	1,526	1,072	-
2000	4,423	6	2	4	10	25	57	207	419	1,041	1,576	1,075	1
1999	4,179	8	-	5	12	24	64	171	431	969	1,514	981	-
All origins, female													
2005	5,519	6	-	4	7	23	81	155	340	729	1,654	2,520	-
2004	5,774	7	1	3	7	21	81	168	369	828	1,747	2,542	-
2003	5,621	7	-	4	13	19	75	180	383	745	1,683	2,512	-
2002	5,813	6	-	4	16	31	78	168	392	890	1,722	2,506	-
2001	5,802	4	2	7	9	28	95	181	378	862	1,755	2,481	-
2000	5,803	10	3	2	13	22	88	169	324	873	1,811	2,488	-
1999	5,823	3	1	3	6	24	85	162	342	838	1,814	2,545	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Other diseases of arteries, arterioles and capillaries (I72-I78)													
Hispanic, both sexes													
2005	437	1	2	4	4	4	16	27	34	80	159	106	-
2004	381	2	-	1	2	4	15	28	41	77	113	98	-
2003	363	2	-	1	3	6	14	25	32	70	100	110	-
2002	416	1	-	1	4	6	20	26	42	86	119	111	-
2001	351	2	-	2	3	5	13	32	41	63	89	101	-
2000	353	3	1	-	1	9	17	15	41	96	78	92	-
1999	345	2	-	1	5	6	10	11	39	75	96	100	-
Hispanic, male													
2005	230	-	2	3	2	3	7	15	20	48	80	50	-
2004	179	1	-	1	1	2	6	16	19	45	53	35	-
2003	177	1	-	-	1	3	9	15	22	35	53	38	-
2002	179	-	-	-	3	2	11	12	15	41	60	35	-
2001	145	1	-	-	2	-	6	15	22	29	41	29	-
2000	158	1	-	-	-	3	8	5	22	46	40	33	-
1999	160	1	-	-	4	2	5	3	18	34	54	39	-
Hispanic, female													
2005	207	1	-	1	2	1	9	12	14	32	79	56	-
2004	202	1	-	-	1	2	9	12	22	32	60	63	-
2003	186	1	-	1	2	3	5	10	10	35	47	72	-
2002	237	1	-	1	1	4	9	14	27	45	59	76	-
2001	206	1	-	2	1	5	7	17	19	34	48	72	-
2000	195	2	1	-	1	6	9	10	19	50	38	59	-
1999	185	1	-	1	1	4	5	8	21	41	42	61	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Other diseases of arteries, arterioles and capillaries (I72-I78)													
Non-Hispanic 1/, both sexes													
2005	9,317	15	1	9	14	38	122	300	774	1,532	3,026	3,486	-
2004	9,534	11	3	7	14	37	128	364	748	1,620	3,087	3,515	-
2003	9,480	8	-	8	24	33	121	353	760	1,587	3,117	3,469	-
2002	9,664	11	4	5	23	44	119	334	780	1,736	3,102	3,506	-
2001	9,707	10	2	5	15	47	153	329	758	1,762	3,185	3,441	-
2000	9,851	13	4	6	22	38	128	361	698	1,814	3,303	3,463	1
1999	9,630	9	1	7	13	42	139	321	731	1,726	3,226	3,415	-
Non-Hispanic 1/, male													
2005	4,010	10	1	6	9	16	50	157	448	835	1,453	1,025	-
2004	3,979	5	2	4	8	18	56	210	401	826	1,406	1,043	-
2003	4,057	2	-	5	13	17	51	184	387	881	1,484	1,033	-
2002	4,108	6	4	2	8	17	50	180	419	895	1,441	1,086	-
2001	4,128	7	-	-	7	24	65	166	400	935	1,483	1,041	-
2000	4,255	5	2	4	10	22	49	202	395	992	1,534	1,039	1
1999	4,009	7	-	5	8	22	59	168	411	932	1,457	940	-
Non-Hispanic 1/, female													
2005	5,307	5	-	3	5	22	72	143	326	697	1,573	2,461	-
2004	5,555	6	1	3	6	19	72	154	347	794	1,681	2,472	-
2003	5,423	6	-	3	11	16	70	169	373	706	1,633	2,436	-
2002	5,556	5	-	3	15	27	69	154	361	841	1,661	2,420	-
2001	5,579	3	2	5	8	23	88	163	358	827	1,702	2,400	-
2000	5,596	8	2	2	12	16	79	159	303	822	1,769	2,424	-
1999	5,621	2	1	2	5	20	80	153	320	794	1,769	2,475	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Other diseases of arteries, arterioles and capillaries (I72-I78)													
Non-Hispanic white, both sexes													
2005 2/	7,888	9	1	7	9	21	87	221	602	1,245	2,592	3,094	-
2004 2/	8,056	8	3	5	12	23	90	271	575	1,331	2,662	3,076	-
2003 2/	8,044	4	-	6	14	24	78	250	609	1,314	2,684	3,061	-
2002	8,220	8	4	2	14	29	80	245	602	1,428	2,696	3,112	-
2001	8,222	8	1	4	12	31	100	247	585	1,460	2,753	3,021	-
2000	8,391	11	3	5	17	27	87	245	533	1,520	2,883	3,060	-
1999	8,257	7	1	6	10	30	106	250	578	1,452	2,803	3,014	-
Non-Hispanic white, male													
2005 2/	3,384	5	1	5	4	9	33	111	351	687	1,250	928	-
2004 2/	3,402	4	2	2	6	12	42	161	310	695	1,242	926	-
2003 2/	3,457	1	-	4	9	13	34	139	309	737	1,282	929	-
2002	3,503	5	4	1	4	11	38	140	323	745	1,272	960	-
2001	3,513	5	-	-	6	16	39	120	317	788	1,301	921	-
2000	3,650	3	2	4	8	18	33	139	311	848	1,356	928	-
1999	3,463	5	-	4	6	17	49	138	333	802	1,271	838	-
Non-Hispanic white, female													
2005 2/	4,504	4	-	2	5	12	54	110	251	558	1,342	2,166	-
2004 2/	4,654	4	1	3	6	11	48	110	265	636	1,420	2,150	-
2003 2/	4,587	3	-	2	5	11	44	111	300	577	1,402	2,132	-
2002	4,717	3	-	1	10	18	42	105	279	683	1,424	2,152	-
2001	4,709	3	1	4	6	15	61	127	268	672	1,452	2,100	-
2000	4,741	8	1	1	9	9	54	106	222	672	1,527	2,132	-
1999	4,794	2	1	2	4	13	57	112	245	650	1,532	2,176	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Other diseases of arteries, arterioles and capillaries (I72-I78)													
Non-Hispanic black, both sexes													
2005 2/	1,286	4	-	2	4	17	32	70	157	260	388	352	-
2004 2/	1,308	2	-	2	1	13	35	81	149	258	377	390	-
2003 2/	1,280	4	-	2	9	8	38	94	129	250	391	355	-
2002	1,295	3	-	2	7	13	33	80	158	277	368	354	-
2001	1,368	2	1	1	2	16	50	73	159	286	394	384	-
2000	1,307	2	-	1	4	10	34	97	150	261	379	368	1
1999	1,244	2	-	-	3	9	32	60	138	246	386	368	-
Non-Hispanic black, male													
2005 2/	557	3	-	1	4	7	15	42	90	133	178	84	-
2004 2/	523	1	-	2	1	5	12	43	81	123	153	102	-
2003 2/	524	1	-	1	3	3	16	41	68	128	179	84	-
2002	532	1	-	-	2	6	10	33	85	136	150	109	-
2001	555	2	-	-	-	8	24	41	73	138	166	103	-
2000	532	2	-	-	1	3	13	55	75	128	161	93	1
1999	487	2	-	-	2	4	9	26	69	119	166	90	-
Non-Hispanic black, female													
2005 2/	729	1	-	1	-	10	17	28	67	127	210	268	-
2004 2/	785	1	-	-	-	8	23	38	68	135	224	288	-
2003 2/	756	3	-	1	6	5	22	53	61	122	212	271	-
2002	763	2	-	2	5	7	23	47	73	141	218	245	-
2001	813	-	1	1	2	8	26	32	86	148	228	281	-
2000	775	-	-	1	3	7	21	42	75	133	218	275	-
1999	757	-	-	-	1	5	23	34	69	127	220	278	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Other diseases of arteries, arterioles and capillaries (I72-I78)													
Origin not stated 3/, both sexes													
2005	20	-	-	-	-	-	-	-	2	2	11	5	-
2004	25	-	-	-	-	-	-	4	-	3	10	8	-
2003	19	-	-	-	-	-	-	1	1	6	6	5	-
2002	29	-	-	-	-	-	1	-	4	6	6	12	-
2001	26	-	-	-	-	1	-	2	2	3	7	11	-
2000	22	-	-	-	-	-	-	-	4	4	6	8	-
1999	27	-	-	-	-	-	-	1	3	6	6	11	-
Origin not stated 3/, male													
2005	15	-	-	-	-	-	-	-	2	2	9	2	-
2004	8	-	-	-	-	-	-	2	-	1	4	1	-
2003	7	-	-	-	-	-	-	-	1	2	3	1	-
2002	9	-	-	-	-	-	1	-	-	2	4	2	-
2001	9	-	-	-	-	1	-	1	1	2	2	2	-
2000	10	-	-	-	-	-	-	-	2	3	2	3	-
1999	10	-	-	-	-	-	-	-	2	3	3	2	-
Origin not stated 3/, female													
2005	5	-	-	-	-	-	-	-	-	-	2	3	-
2004	17	-	-	-	-	-	-	2	-	2	6	7	-
2003	12	-	-	-	-	-	-	1	-	4	3	4	-
2002	20	-	-	-	-	-	-	-	4	4	2	10	-
2001	17	-	-	-	-	-	-	1	1	1	5	9	-
2000	12	-	-	-	-	-	-	-	2	1	4	5	-
1999	17	-	-	-	-	-	-	1	1	3	3	9	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Other disorders of circulatory system (I80-I99)													
All origins, both sexes													
2005	4,813	26	6	5	36	132	332	620	641	713	1,128	1,174	-
2004	4,673	29	2	5	35	137	313	559	623	672	1,165	1,133	-
2003	4,737	36	4	4	50	139	342	555	569	744	1,161	1,133	-
2002	4,711	27	4	2	35	139	305	531	595	697	1,253	1,123	-
2001	4,665	22	5	-	38	124	285	454	563	841	1,183	1,149	1
2000	4,603	27	4	12	37	133	319	515	516	795	1,161	1,084	-
1999	4,025	23	1	2	30	93	271	378	445	692	1,117	972	1
All origins, male													
2005	2,156	13	3	3	15	62	164	382	381	379	446	308	-
2004	2,083	18	-	3	19	72	162	329	351	328	505	296	-
2003	1,966	21	3	2	20	60	167	312	320	348	437	276	-
2002	1,944	15	1	-	21	69	154	284	309	323	483	285	-
2001	1,981	11	5	-	17	53	149	262	304	405	488	286	1
2000	1,981	16	2	6	16	57	190	295	264	390	472	273	-
1999	1,714	13	1	1	13	44	159	198	224	350	463	248	-
All origins, female													
2005	2,657	13	3	2	21	70	168	238	260	334	682	866	-
2004	2,590	11	2	2	16	65	151	230	272	344	660	837	-
2003	2,771	15	1	2	30	79	175	243	249	396	724	857	-
2002	2,767	12	3	2	14	70	151	247	286	374	770	838	-
2001	2,684	11	-	-	21	71	136	192	259	436	695	863	-
2000	2,622	11	2	6	21	76	129	220	252	405	689	811	-
1999	2,311	10	-	1	17	49	112	180	221	342	654	724	1

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Other disorders of circulatory system (I80-I99)													
Hispanic, both sexes													
2005	281	2	2	1	3	16	38	43	36	46	59	35	-
2004	224	5	1	-	4	15	20	47	36	26	33	37	-
2003	239	11	2	1	7	15	26	41	29	32	41	34	-
2002	219	7	1	-	5	15	21	35	27	37	38	33	-
2001	219	5	-	-	3	7	15	32	16	48	41	52	-
2000	229	6	-	4	3	11	32	34	23	35	44	37	-
1999	186	2	-	1	4	6	12	34	24	42	26	35	-
Hispanic, male													
2005	126	-	2	1	-	8	17	25	23	23	13	14	-
2004	106	3	-	-	1	9	11	29	23	13	13	4	-
2003	120	5	1	1	2	5	19	27	17	18	16	9	-
2002	95	5	-	-	3	8	9	17	16	16	16	5	-
2001	104	4	-	-	2	4	7	22	9	20	20	16	-
2000	116	3	-	2	1	7	20	19	13	16	22	13	-
1999	83	-	-	1	2	4	8	15	11	17	13	12	-
Hispanic, female													
2005	155	2	-	-	3	8	21	18	13	23	46	21	-
2004	118	2	1	-	3	6	9	18	13	13	20	33	-
2003	119	6	1	-	5	10	7	14	12	14	25	25	-
2002	124	2	1	-	2	7	12	18	11	21	22	28	-
2001	115	1	-	-	1	3	8	10	7	28	21	36	-
2000	113	3	-	2	2	4	12	15	10	19	22	24	-
1999	103	2	-	-	2	2	4	19	13	25	13	23	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Other disorders of circulatory system (I80-I99)													
Non-Hispanic 1/, both sexes													
2005	4,520	24	4	4	33	116	293	575	601	666	1,066	1,138	-
2004	4,439	24	1	5	31	122	293	512	584	644	1,130	1,093	-
2003	4,485	25	2	3	43	123	315	513	537	709	1,117	1,098	-
2002	4,479	20	3	2	30	124	281	494	566	659	1,212	1,088	-
2001	4,425	17	5	-	35	117	269	418	544	788	1,136	1,095	1
2000	4,359	20	4	8	34	120	283	479	491	758	1,117	1,045	-
1999	3,825	21	1	1	26	86	256	343	420	648	1,088	934	1
Non-Hispanic 1/, male													
2005	2,024	13	1	2	15	54	147	356	354	356	432	294	-
2004	1,971	15	-	3	18	63	151	300	326	314	491	290	-
2003	1,840	16	2	1	18	54	148	284	303	328	419	267	-
2002	1,842	10	1	-	18	61	143	265	292	306	466	280	-
2001	1,869	7	5	-	15	49	141	239	293	384	466	269	1
2000	1,856	12	2	4	15	49	167	274	250	373	450	260	-
1999	1,623	13	1	-	11	40	148	182	212	331	450	235	-
Non-Hispanic 1/, female													
2005	2,496	11	3	2	18	62	146	219	247	310	634	844	-
2004	2,468	9	1	2	13	59	142	212	258	330	639	803	-
2003	2,645	9	-	2	25	69	167	229	234	381	698	831	-
2002	2,637	10	2	2	12	63	138	229	274	353	746	808	-
2001	2,556	10	-	-	20	68	128	179	251	404	670	826	-
2000	2,503	8	2	4	19	71	116	205	241	385	667	785	-
1999	2,202	8	-	1	15	46	108	161	208	317	638	699	1

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Other disorders of circulatory system (I80-I99)													
Non-Hispanic white, both sexes													
2005 2/	3,659	19	1	4	18	71	185	415	455	539	910	1,042	-
2004 2/	3,571	17	1	3	19	68	181	354	456	517	978	977	-
2003 2/	3,607	18	2	2	23	72	198	381	413	575	952	971	-
2002	3,630	13	2	1	16	69	180	363	430	538	1,041	977	-
2001	3,617	12	3	-	18	73	175	306	414	642	987	987	-
2000	3,554	13	3	5	19	70	195	358	373	602	975	941	-
1999	3,149	17	-	1	12	57	168	243	342	524	949	835	1
Non-Hispanic white, male													
2005 2/	1,603	11	-	2	8	33	97	262	271	284	370	265	-
2004 2/	1,576	12	-	2	9	32	98	216	264	256	425	262	-
2003 2/	1,449	11	2	-	9	31	110	202	236	266	349	233	-
2002	1,450	8	-	-	9	33	92	194	224	248	396	246	-
2001	1,496	4	3	-	8	30	92	182	220	313	405	239	-
2000	1,496	9	1	3	8	26	118	207	188	307	398	231	-
1999	1,310	9	-	-	7	26	95	127	180	268	388	210	-
Non-Hispanic white, female													
2005 2/	2,056	8	1	2	10	38	88	153	184	255	540	777	-
2004 2/	1,995	5	1	1	10	36	83	138	192	261	553	715	-
2003 2/	2,158	7	-	2	14	41	88	179	177	309	603	738	-
2002	2,180	5	2	1	7	36	88	169	206	290	645	731	-
2001	2,121	8	-	-	10	43	83	124	194	329	582	748	-
2000	2,058	4	2	2	11	44	77	151	185	295	577	710	-
1999	1,839	8	-	1	5	31	73	116	162	256	561	625	1

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Other disorders of circulatory system (I80-I99)													
Non-Hispanic black, both sexes													
2005 2/	785	4	2	-	15	42	98	149	136	112	138	89	-
2004 2/	790	7	-	1	10	50	108	145	118	115	135	101	-
2003 2/	812	6	-	-	19	48	111	124	118	124	149	113	-
2002	789	7	1	1	13	54	94	120	128	113	159	99	-
2001	729	5	2	-	14	43	87	98	122	129	133	95	1
2000	751	5	1	3	14	48	86	116	109	145	130	94	-
1999	613	3	1	-	14	28	81	95	69	109	125	88	-
Non-Hispanic black, male													
2005 2/	380	2	-	-	7	18	44	87	76	64	57	25	-
2004 2/	355	3	-	-	7	28	52	78	57	51	54	25	-
2003 2/	363	5	-	-	9	22	36	76	64	58	62	31	-
2002	359	2	1	-	9	27	48	63	64	53	63	29	-
2001	335	3	2	-	5	18	46	48	72	62	53	25	1
2000	333	2	1	1	7	22	48	66	56	61	43	26	-
1999	281	3	1	-	4	14	49	52	28	56	55	19	-
Non-Hispanic black, female													
2005 2/	405	2	2	-	8	24	54	62	60	48	81	64	-
2004 2/	435	4	-	1	3	22	56	67	61	64	81	76	-
2003 2/	449	1	-	-	10	26	75	48	54	66	87	82	-
2002	430	5	-	1	4	27	46	57	64	60	96	70	-
2001	394	2	-	-	9	25	41	50	50	67	80	70	-
2000	418	3	-	2	7	26	38	50	53	84	87	68	-
1999	332	-	-	-	10	14	32	43	41	53	70	69	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Other disorders of circulatory system (I80-I99)													
Origin not stated 3/, both sexes													
2005	12	-	-	-	-	-	1	2	4	1	3	1	-
2004	10	-	-	-	-	-	-	-	3	2	2	3	-
2003	13	-	-	-	-	1	1	1	3	3	3	1	-
2002	13	-	-	-	-	-	3	2	2	1	3	2	-
2001	21	-	-	-	-	-	1	4	3	5	6	2	-
2000	15	1	-	-	-	2	4	2	2	2	-	2	-
1999	14	-	-	-	-	1	3	1	1	2	3	3	-
Origin not stated 3/, male													
2005	6	-	-	-	-	-	-	1	4	-	1	-	-
2004	6	-	-	-	-	-	-	-	2	1	1	2	-
2003	6	-	-	-	-	1	-	1	-	2	2	-	-
2002	7	-	-	-	-	-	2	2	1	1	1	-	-
2001	8	-	-	-	-	-	1	1	2	1	2	1	-
2000	9	1	-	-	-	1	3	2	1	1	-	-	-
1999	8	-	-	-	-	-	3	1	1	2	-	1	-
Origin not stated 3/, female													
2005	6	-	-	-	-	-	1	1	-	1	2	1	-
2004	4	-	-	-	-	-	-	-	1	1	1	1	-
2003	7	-	-	-	-	-	1	-	3	1	1	1	-
2002	6	-	-	-	-	-	1	-	1	-	2	2	-
2001	13	-	-	-	-	-	-	3	1	4	4	1	-
2000	6	-	-	-	-	1	1	-	1	1	-	2	-
1999	6	-	-	-	-	1	-	-	-	-	3	2	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Influenza and pneumonia (J10-J18)													
All origins, both sexes													
2005	63,001	265	110	106	172	354	934	2,183	3,422	6,623	18,563	30,267	2
2004	59,664	273	119	82	185	303	891	1,897	3,154	6,382	18,066	28,312	-
2003	65,163	322	163	147	224	373	992	2,140	3,130	6,831	19,442	31,397	2
2002	65,681	263	110	91	167	345	971	1,918	2,987	6,847	19,984	31,995	3
2001	62,034	299	112	92	181	339	983	1,801	2,704	6,650	18,677	30,191	5
2000	65,313	289	103	87	189	364	1,068	1,774	2,879	7,189	19,821	31,547	3
1999	63,730	320	130	93	179	339	1,063	1,697	2,625	6,861	19,192	31,229	2
All origins, male													
2005	28,052	148	56	53	117	196	540	1,299	1,988	3,648	9,203	10,802	2
2004	26,861	166	47	47	118	194	494	1,142	1,823	3,605	9,013	10,212	-
2003	28,778	192	91	70	124	229	560	1,279	1,771	3,882	9,629	10,950	1
2002	28,918	162	63	38	97	190	546	1,138	1,738	3,850	9,895	11,199	2
2001	27,342	167	57	47	106	191	556	1,099	1,587	3,732	9,294	10,502	4
2000	28,658	163	54	48	102	204	630	1,066	1,680	4,013	9,751	10,945	2
1999	27,718	171	66	53	90	197	592	1,059	1,462	3,893	9,458	10,676	1
All origins, female													
2005	34,949	117	54	53	55	158	394	884	1,434	2,975	9,360	19,465	-
2004	32,803	107	72	35	67	109	397	755	1,331	2,777	9,053	18,100	-
2003	36,385	130	72	77	100	144	432	861	1,359	2,949	9,813	20,447	1
2002	36,763	101	47	53	70	155	425	780	1,249	2,997	10,089	20,796	1
2001	34,692	132	55	45	75	148	427	702	1,117	2,918	9,383	19,689	1
2000	36,655	126	49	39	87	160	438	708	1,199	3,176	10,070	20,602	1
1999	36,012	149	64	40	89	142	471	638	1,163	2,968	9,734	20,553	1

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Influenza and pneumonia (J10-J18)													
Hispanic, both sexes													
2005	3,085	66	32	9	11	40	93	172	247	455	892	1,068	-
2004	2,912	53	24	13	32	48	82	153	214	422	857	1,014	-
2003	2,948	66	37	22	36	54	74	151	224	428	860	996	-
2002	2,824	48	17	14	19	43	77	124	207	399	855	1,021	-
2001	2,722	57	32	10	18	23	73	121	168	402	761	1,056	1
2000	2,625	60	17	21	21	48	82	120	191	405	690	970	-
1999	2,246	64	17	10	25	24	74	104	155	328	569	876	-
Hispanic, male													
2005	1,506	37	15	7	8	26	59	110	147	250	448	399	-
2004	1,403	37	9	10	24	31	48	100	122	220	425	377	-
2003	1,429	40	24	12	20	35	41	100	115	235	432	375	-
2002	1,398	33	11	6	11	26	52	86	130	229	422	392	-
2001	1,309	31	16	6	15	14	48	85	99	204	388	402	1
2000	1,303	30	5	10	10	27	56	82	119	218	367	379	-
1999	1,092	40	10	6	11	15	54	71	80	187	284	334	-
Hispanic, female													
2005	1,579	29	17	2	3	14	34	62	100	205	444	669	-
2004	1,509	16	15	3	8	17	34	53	92	202	432	637	-
2003	1,519	26	13	10	16	19	33	51	109	193	428	621	-
2002	1,426	15	6	8	8	17	25	38	77	170	433	629	-
2001	1,413	26	16	4	3	9	25	36	69	198	373	654	-
2000	1,322	30	12	11	11	21	26	38	72	187	323	591	-
1999	1,154	24	7	4	14	9	20	33	75	141	285	542	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Influenza and pneumonia (J10-J18)													
Non-Hispanic 1/, both sexes													
2005	59,804	197	78	96	160	313	838	2,003	3,167	6,151	17,633	29,167	1
2004	56,581	220	95	68	151	255	801	1,734	2,917	5,930	17,165	27,245	-
2003	62,042	254	126	124	188	319	917	1,975	2,883	6,373	18,544	30,337	2
2002	62,657	213	92	76	147	300	889	1,783	2,773	6,424	19,059	30,898	3
2001	59,103	240	79	82	163	312	906	1,669	2,516	6,209	17,863	29,062	2
2000	62,440	228	84	66	167	314	974	1,639	2,674	6,750	19,051	30,492	1
1999	61,218	255	110	83	153	315	979	1,577	2,446	6,499	18,553	30,247	1
Non-Hispanic 1/, male													
2005	26,482	109	41	46	108	169	479	1,184	1,834	3,385	8,733	10,393	1
2004	25,359	129	38	37	93	163	441	1,033	1,683	3,360	8,568	9,814	-
2003	27,255	150	67	58	104	194	518	1,167	1,641	3,626	9,178	10,551	1
2002	27,417	128	52	32	85	162	491	1,043	1,603	3,600	9,437	10,782	2
2001	25,926	135	41	41	91	175	504	1,007	1,475	3,500	8,881	10,075	1
2000	27,218	132	47	38	91	175	566	972	1,549	3,772	9,339	10,536	1
1999	26,489	131	55	47	79	182	531	974	1,367	3,687	9,138	10,298	-
Non-Hispanic 1/, female													
2005	33,322	88	37	50	52	144	359	819	1,333	2,766	8,900	18,774	-
2004	31,222	91	57	31	58	92	360	701	1,234	2,570	8,597	17,431	-
2003	34,787	104	59	66	84	125	399	808	1,242	2,747	9,366	19,786	1
2002	35,240	85	40	44	62	138	398	740	1,170	2,824	9,622	20,116	1
2001	33,177	105	38	41	72	137	402	662	1,041	2,709	8,982	18,987	1
2000	35,222	96	37	28	76	139	408	667	1,125	2,978	9,712	19,956	-
1999	34,729	124	55	36	74	133	448	603	1,079	2,812	9,415	19,949	1

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Influenza and pneumonia (J10-J18)													
Non-Hispanic white, both sexes													
2005 2/	52,431	94	46	67	123	224	593	1,449	2,457	5,022	15,638	26,717	1
2004 2/	49,456	112	55	43	108	183	552	1,242	2,295	4,894	15,143	24,829	-
2003 2/	54,617	143	80	86	137	206	637	1,424	2,210	5,262	16,545	27,886	1
2002	55,419	114	46	55	103	207	617	1,329	2,152	5,419	17,001	28,373	3
2001	51,952	117	48	57	108	207	605	1,193	1,963	5,214	15,837	26,602	1
2000	55,135	114	51	49	116	213	678	1,165	2,061	5,546	17,068	28,073	1
1999	54,280	121	58	62	95	199	637	1,109	1,894	5,481	16,785	27,838	1
Non-Hispanic white, male													
2005 2/	22,903	60	23	29	82	117	334	847	1,399	2,753	7,767	9,491	1
2004 2/	21,905	65	23	22	73	115	300	741	1,328	2,750	7,549	8,939	-
2003 2/	23,541	87	44	39	75	131	356	844	1,219	2,966	8,141	9,639	-
2002	23,934	73	24	20	62	111	340	768	1,228	3,050	8,402	9,854	2
2001	22,389	71	26	28	64	112	333	707	1,132	2,919	7,830	9,166	1
2000	23,609	68	24	29	68	111	394	667	1,176	3,084	8,346	9,641	1
1999	23,097	64	22	36	51	119	334	670	1,049	3,087	8,255	9,410	-
Non-Hispanic white, female													
2005 2/	29,528	34	23	38	41	107	259	602	1,058	2,269	7,871	17,226	-
2004 2/	27,551	47	32	21	35	68	252	501	967	2,144	7,594	15,890	-
2003 2/	31,076	56	36	47	62	75	281	580	991	2,296	8,404	18,247	1
2002	31,485	41	22	35	41	96	277	561	924	2,369	8,599	18,519	1
2001	29,563	46	22	29	44	95	272	486	831	2,295	8,007	17,436	-
2000	31,526	46	27	20	48	102	284	498	885	2,462	8,722	18,432	-
1999	31,183	57	36	26	44	80	303	439	845	2,394	8,530	18,428	1

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Influenza and pneumonia (J10-J18)													
Non-Hispanic black, both sexes													
2005 2/	5,716	95	28	22	31	77	214	494	603	914	1,497	1,741	-
2004 2/	5,620	93	34	20	40	63	223	443	537	850	1,539	1,778	-
2003 2/	5,798	91	27	28	42	101	251	487	565	915	1,514	1,776	1
2002	5,803	86	40	18	38	80	250	399	554	837	1,584	1,917	-
2001	5,686	103	24	17	45	87	274	430	471	796	1,571	1,868	-
2000	5,920	101	29	11	44	92	254	423	531	984	1,550	1,901	-
1999	5,794	114	40	15	49	107	305	425	486	880	1,410	1,963	-
Non-Hispanic black, male													
2005 2/	2,697	45	16	13	23	44	127	299	372	510	683	565	-
2004 2/	2,729	55	12	12	19	41	126	266	311	514	775	598	-
2003 2/	2,861	52	13	15	24	58	142	285	356	547	765	603	1
2002	2,734	47	22	10	20	43	136	239	338	454	789	636	-
2001	2,769	52	13	9	23	52	155	270	293	472	808	622	-
2000	2,880	55	20	7	21	58	151	271	322	562	763	650	-
1999	2,786	58	25	8	23	58	173	277	282	518	688	676	-
Non-Hispanic black, female													
2005 2/	3,019	50	12	9	8	33	87	195	231	404	814	1,176	-
2004 2/	2,891	38	22	8	21	22	97	177	226	336	764	1,180	-
2003 2/	2,937	39	14	13	18	43	109	202	209	368	749	1,173	-
2002	3,069	39	18	8	18	37	114	160	216	383	795	1,281	-
2001	2,917	51	11	8	22	35	119	160	178	324	763	1,246	-
2000	3,040	46	9	4	23	34	103	152	209	422	787	1,251	-
1999	3,008	56	15	7	26	49	132	148	204	362	722	1,287	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Influenza and pneumonia (J10-J18)													
Origin not stated 3/, both sexes													
2005	112	2	-	1	1	1	3	8	8	17	38	32	1
2004	171	-	-	1	2	-	8	10	23	30	44	53	-
2003	173	2	-	1	-	-	1	14	23	30	38	64	-
2002	200	2	1	1	1	2	5	11	7	24	70	76	-
2001	209	2	1	-	-	4	4	11	20	39	53	73	2
2000	248	1	2	-	1	2	12	15	14	34	80	85	2
1999	266	1	3	-	1	-	10	16	24	34	70	106	1
Origin not stated 3/, male													
2005	64	2	-	-	1	1	2	5	7	13	22	10	1
2004	99	-	-	-	1	-	5	9	18	25	20	21	-
2003	94	2	-	-	-	-	1	12	15	21	19	24	-
2002	103	1	-	-	1	2	3	9	5	21	36	25	-
2001	107	1	-	-	-	2	4	7	13	28	25	25	2
2000	137	1	2	-	1	2	8	12	12	23	45	30	1
1999	137	-	1	-	-	-	7	14	15	19	36	44	1
Origin not stated 3/, female													
2005	48	-	-	1	-	-	1	3	1	4	16	22	-
2004	72	-	-	1	1	-	3	1	5	5	24	32	-
2003	79	-	-	1	-	-	-	2	8	9	19	40	-
2002	97	1	1	1	-	-	2	2	2	3	34	51	-
2001	102	1	1	-	-	2	-	4	7	11	28	48	-
2000	111	-	-	-	-	-	4	3	2	11	35	55	1
1999	129	1	2	-	1	-	3	2	9	15	34	62	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Influenza (J10-J11)													
All origins, both sexes													
2005	1,812	19	19	20	12	11	19	45	66	137	482	982	-
2004	1,100	18	18	13	7	4	16	25	55	99	291	554	-
2003	1,792	31	59	46	30	13	24	61	85	183	457	803	-
2002	727	7	5	12	7	4	8	17	30	56	177	404	-
2001	257	7	6	12	7	10	6	15	21	21	56	96	-
2000	1,765	9	10	11	13	13	22	48	93	199	492	855	-
1999	1,665	13	12	11	11	9	16	26	71	164	451	881	-
All origins, male													
2005	657	9	10	10	6	3	11	26	37	61	192	292	-
2004	421	10	8	8	2	2	7	8	25	58	125	168	-
2003	699	18	33	19	12	9	10	33	42	89	189	245	-
2002	289	5	3	2	4	2	4	12	20	32	81	124	-
2001	101	6	2	4	6	6	2	9	9	8	20	29	-
2000	664	5	4	5	6	5	14	25	44	106	196	254	-
1999	602	6	9	5	2	5	9	15	37	74	202	238	-
All origins, female													
2005	1,155	10	9	10	6	8	8	19	29	76	290	690	-
2004	679	8	10	5	5	2	9	17	30	41	166	386	-
2003	1,093	13	26	27	18	4	14	28	43	94	268	558	-
2002	438	2	2	10	3	2	4	5	10	24	96	280	-
2001	156	1	4	8	1	4	4	6	12	13	36	67	-
2000	1,101	4	6	6	7	8	8	23	49	93	296	601	-
1999	1,063	7	3	6	9	4	7	11	34	90	249	643	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Influenza (J10-J11)													
Hispanic, both sexes													
2005	55	8	6	3	-	2	-	5	2	4	10	15	-
2004	26	6	2	2	1	-	-	3	-	4	2	6	-
2003	75	10	14	9	4	2	3	2	6	3	12	10	-
2002	19	2	-	4	-	1	1	1	-	2	3	5	-
2001	13	-	3	1	1	-	1	1	-	1	5	-	-
2000	47	2	1	3	1	4	2	2	2	4	10	16	-
1999	45	4	3	-	2	-	2	2	5	8	9	10	-
Hispanic, male													
2005	26	5	2	2	-	1	-	4	1	2	2	7	-
2004	11	3	-	2	1	-	-	-	-	2	2	1	-
2003	40	7	10	3	1	1	2	-	2	3	7	4	-
2002	10	2	-	2	-	1	1	1	-	1	1	1	-
2001	7	-	2	1	1	-	-	1	-	-	2	-	-
2000	21	-	1	1	1	1	1	1	2	2	5	6	-
1999	21	-	3	-	-	-	2	2	4	7	2	1	-
Hispanic, female													
2005	29	3	4	1	-	1	-	1	1	2	8	8	-
2004	15	3	2	-	-	-	-	3	-	2	-	5	-
2003	35	3	4	6	3	1	1	2	4	-	5	6	-
2002	9	-	-	2	-	-	-	-	-	1	2	4	-
2001	6	-	1	-	-	-	1	-	-	1	3	-	-
2000	26	2	-	2	-	3	1	1	-	2	5	10	-
1999	24	4	-	-	2	-	-	-	1	1	7	9	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Influenza (J10-J11)													
Non-Hispanic 1/, both sexes													
2005	1,752	11	13	17	12	9	19	40	63	133	472	963	-
2004	1,071	12	16	10	6	4	16	22	55	95	288	547	-
2003	1,714	21	45	37	26	11	21	59	78	180	443	793	-
2002	704	5	5	8	7	3	7	16	30	54	173	396	-
2001	244	7	3	11	6	10	5	14	21	20	51	96	-
2000	1,716	7	9	8	12	9	20	46	91	195	481	838	-
1999	1,612	9	9	11	9	9	14	24	66	154	439	868	-
Non-Hispanic 1/, male													
2005	631	4	8	8	6	2	11	22	36	59	190	285	-
2004	410	7	8	6	1	2	7	8	25	56	123	167	-
2003	657	11	23	16	11	8	8	33	39	86	181	241	-
2002	277	3	3	-	4	1	3	11	20	31	79	122	-
2001	94	6	-	3	5	6	2	8	9	8	18	29	-
2000	642	5	3	4	5	4	13	24	42	104	190	248	-
1999	580	6	6	5	2	5	7	13	33	66	200	237	-
Non-Hispanic 1/, female													
2005	1,121	7	5	9	6	7	8	18	27	74	282	678	-
2004	661	5	8	4	5	2	9	14	30	39	165	380	-
2003	1,057	10	22	21	15	3	13	26	39	94	262	552	-
2002	427	2	2	8	3	2	4	5	10	23	94	274	-
2001	150	1	3	8	1	4	3	6	12	12	33	67	-
2000	1,074	2	6	4	7	5	7	22	49	91	291	590	-
1999	1,032	3	3	6	7	4	7	11	33	88	239	631	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Influenza (J10-J11)													
Non-Hispanic white, both sexes													
2005 2/	1,654	6	10	10	10	8	13	30	57	124	450	936	-
2004 2/	1,004	6	12	8	5	3	14	17	50	88	268	533	-
2003 2/	1,578	13	29	22	16	8	17	47	69	167	426	764	-
2002	663	3	2	6	5	3	7	15	25	45	168	384	-
2001	216	7	2	6	4	7	3	14	17	18	47	91	-
2000	1,613	5	8	6	11	7	16	38	85	176	458	803	-
1999	1,522	3	5	10	8	8	8	20	60	139	417	844	-
Non-Hispanic white, male													
2005 2/	585	4	7	5	4	2	8	15	31	56	181	272	-
2004 2/	377	2	5	5	1	1	6	6	23	52	112	164	-
2003 2/	599	7	15	9	6	7	4	27	33	81	175	235	-
2002	258	1	1	-	2	1	3	11	16	27	76	120	-
2001	81	6	-	2	3	3	2	8	6	6	18	27	-
2000	595	5	3	3	4	2	9	18	39	97	181	234	-
1999	547	1	3	5	2	5	4	11	32	59	192	233	-
Non-Hispanic white, female													
2005 2/	1,069	2	3	5	6	6	5	15	26	68	269	664	-
2004 2/	627	4	7	3	4	2	8	11	27	36	156	369	-
2003 2/	979	6	14	13	10	1	13	20	36	86	251	529	-
2002	405	2	1	6	3	2	4	4	9	18	92	264	-
2001	135	1	2	4	1	4	1	6	11	12	29	64	-
2000	1,018	-	5	3	7	5	7	20	46	79	277	569	-
1999	975	2	2	5	6	3	4	9	28	80	225	611	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Influenza (J10-J11)													
Non-Hispanic black, both sexes													
2005 2/	67	5	2	5	1	1	6	7	6	5	14	15	-
2004 2/	50	4	3	1	1	1	1	3	4	7	16	9	-
2003 2/	95	7	6	10	7	3	1	12	7	10	12	20	-
2002	30	1	3	2	2	-	-	-	4	6	4	8	-
2001	20	-	-	2	1	2	1	-	3	2	4	5	-
2000	81	2	1	1	1	1	3	7	6	16	13	30	-
1999	65	4	1	-	1	1	5	3	5	12	15	18	-
Non-Hispanic black, male													
2005 2/	31	-	1	3	1	-	3	5	5	2	3	8	-
2004 2/	25	4	2	-	-	1	-	2	1	4	10	1	-
2003 2/	36	3	3	4	3	1	1	6	5	4	3	3	-
2002	15	1	2	-	2	-	-	-	3	3	2	2	-
2001	10	-	-	-	1	2	-	-	3	2	-	2	-
2000	35	-	-	1	1	1	3	5	3	5	5	11	-
1999	24	3	1	-	-	-	2	2	1	6	6	3	-
Non-Hispanic black, female													
2005 2/	36	5	1	2	-	1	3	2	1	3	11	7	-
2004 2/	25	-	1	1	1	-	1	1	3	3	6	8	-
2003 2/	59	4	3	6	4	2	-	6	2	6	9	17	-
2002	15	-	1	2	-	-	-	-	1	3	2	6	-
2001	10	-	-	2	-	-	1	-	-	-	4	3	-
2000	46	2	1	-	-	-	-	2	3	11	8	19	-
1999	41	1	-	-	1	1	3	1	4	6	9	15	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Influenza (J10-J11)													
Origin not stated 3/, both sexes													
2005	5	-	-	-	-	-	-	-	1	-	-	4	-
2004	3	-	-	1	-	-	-	-	-	-	1	1	-
2003	3	-	-	-	-	-	-	-	1	-	2	-	-
2002	4	-	-	-	-	-	-	-	-	-	1	3	-
2001	-	-	-	-	-	-	-	-	-	-	-	-	-
2000	2	-	-	-	-	-	-	-	-	-	1	1	-
1999	8	-	-	-	-	-	-	-	-	2	3	3	-
Origin not stated 3/, male													
2005	-	-	-	-	-	-	-	-	-	-	-	-	-
2004	-	-	-	-	-	-	-	-	-	-	-	-	-
2003	2	-	-	-	-	-	-	-	1	-	1	-	-
2002	2	-	-	-	-	-	-	-	-	-	1	1	-
2001	-	-	-	-	-	-	-	-	-	-	-	-	-
2000	1	-	-	-	-	-	-	-	-	-	1	-	-
1999	1	-	-	-	-	-	-	-	-	1	-	-	-
Origin not stated 3/, female													
2005	5	-	-	-	-	-	-	-	1	-	-	4	-
2004	3	-	-	1	-	-	-	-	-	-	1	1	-
2003	1	-	-	-	-	-	-	-	-	-	1	-	-
2002	2	-	-	-	-	-	-	-	-	-	-	2	-
2001	-	-	-	-	-	-	-	-	-	-	-	-	-
2000	1	-	-	-	-	-	-	-	-	-	-	1	-
1999	7	-	-	-	-	-	-	-	-	1	3	3	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Pneumonia (J12-J18)													
All origins, both sexes													
2005	61,189	246	91	86	160	343	915	2,138	3,356	6,486	18,081	29,285	2
2004	58,564	255	101	69	178	299	875	1,872	3,099	6,283	17,775	27,758	-
2003	63,371	291	104	101	194	360	968	2,079	3,045	6,648	18,985	30,594	2
2002	64,954	256	105	79	160	341	963	1,901	2,957	6,791	19,807	31,591	3
2001	61,777	292	106	80	174	329	977	1,786	2,683	6,629	18,621	30,095	5
2000	63,548	280	93	76	176	351	1,046	1,726	2,786	6,990	19,329	30,692	3
1999	62,065	307	118	82	168	330	1,047	1,671	2,554	6,697	18,741	30,348	2
All origins, male													
2005	27,395	139	46	43	111	193	529	1,273	1,951	3,587	9,011	10,510	2
2004	26,440	156	39	39	116	192	487	1,134	1,798	3,547	8,888	10,044	-
2003	28,079	174	58	51	112	220	550	1,246	1,729	3,793	9,440	10,705	1
2002	28,629	157	60	36	93	188	542	1,126	1,718	3,818	9,814	11,075	2
2001	27,241	161	55	43	100	185	554	1,090	1,578	3,724	9,274	10,473	4
2000	27,994	158	50	43	96	199	616	1,041	1,636	3,907	9,555	10,691	2
1999	27,116	165	57	48	88	192	583	1,044	1,425	3,819	9,256	10,438	1
All origins, female													
2005	33,794	107	45	43	49	150	386	865	1,405	2,899	9,070	18,775	-
2004	32,124	99	62	30	62	107	388	738	1,301	2,736	8,887	17,714	-
2003	35,292	117	46	50	82	140	418	833	1,316	2,855	9,545	19,889	1
2002	36,325	99	45	43	67	153	421	775	1,239	2,973	9,993	20,516	1
2001	34,536	131	51	37	74	144	423	696	1,105	2,905	9,347	19,622	1
2000	35,554	122	43	33	80	152	430	685	1,150	3,083	9,774	20,001	1
1999	34,949	142	61	34	80	138	464	627	1,129	2,878	9,485	19,910	1

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Pneumonia (J12-J18)													
Hispanic, both sexes													
2005	3,030	58	26	6	11	38	93	167	245	451	882	1,053	-
2004	2,886	47	22	11	31	48	82	150	214	418	855	1,008	-
2003	2,873	56	23	13	32	52	71	149	218	425	848	986	-
2002	2,805	46	17	10	19	42	76	123	207	397	852	1,016	-
2001	2,709	57	29	9	17	23	72	120	168	401	756	1,056	1
2000	2,578	58	16	18	20	44	80	118	189	401	680	954	-
1999	2,201	60	14	10	23	24	72	102	150	320	560	866	-
Hispanic, male													
2005	1,480	32	13	5	8	25	59	106	146	248	446	392	-
2004	1,392	34	9	8	23	31	48	100	122	218	423	376	-
2003	1,389	33	14	9	19	34	39	100	113	232	425	371	-
2002	1,388	31	11	4	11	25	51	85	130	228	421	391	-
2001	1,302	31	14	5	14	14	48	84	99	204	386	402	1
2000	1,282	30	4	9	9	26	55	81	117	216	362	373	-
1999	1,071	40	7	6	11	15	52	69	76	180	282	333	-
Hispanic, female													
2005	1,550	26	13	1	3	13	34	61	99	203	436	661	-
2004	1,494	13	13	3	8	17	34	50	92	200	432	632	-
2003	1,484	23	9	4	13	18	32	49	105	193	423	615	-
2002	1,417	15	6	6	8	17	25	38	77	169	431	625	-
2001	1,407	26	15	4	3	9	24	36	69	197	370	654	-
2000	1,296	28	12	9	11	18	25	37	72	185	318	581	-
1999	1,130	20	7	4	12	9	20	33	74	140	278	533	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Pneumonia (J12-J18)													
Non-Hispanic 1/, both sexes													
2005	58,052	186	65	79	148	304	819	1,963	3,104	6,018	17,161	28,204	1
2004	55,510	208	79	58	145	251	785	1,712	2,862	5,835	16,877	26,698	-
2003	60,328	233	81	87	162	308	896	1,916	2,805	6,193	18,101	29,544	2
2002	61,953	208	87	68	140	297	882	1,767	2,743	6,370	18,886	30,502	3
2001	58,859	233	76	71	157	302	901	1,655	2,495	6,189	17,812	28,966	2
2000	60,724	221	75	58	155	305	954	1,593	2,583	6,555	18,570	29,654	1
1999	59,606	246	101	72	144	306	965	1,553	2,380	6,345	18,114	29,379	1
Non-Hispanic 1/, male													
2005	25,851	105	33	38	102	167	468	1,162	1,798	3,326	8,543	10,108	1
2004	24,949	122	30	31	92	161	434	1,025	1,658	3,304	8,445	9,647	-
2003	26,598	139	44	42	93	186	510	1,134	1,602	3,540	8,997	10,310	1
2002	27,140	125	49	32	81	161	488	1,032	1,583	3,569	9,358	10,660	2
2001	25,832	129	41	38	86	169	502	999	1,466	3,492	8,863	10,046	1
2000	26,576	127	44	34	86	171	553	948	1,507	3,668	9,149	10,288	1
1999	25,909	125	49	42	77	177	524	961	1,334	3,621	8,938	10,061	-
Non-Hispanic 1/, female													
2005	32,201	81	32	41	46	137	351	801	1,306	2,692	8,618	18,096	-
2004	30,561	86	49	27	53	90	351	687	1,204	2,531	8,432	17,051	-
2003	33,730	94	37	45	69	122	386	782	1,203	2,653	9,104	19,234	1
2002	34,813	83	38	36	59	136	394	735	1,160	2,801	9,528	19,842	1
2001	33,027	104	35	33	71	133	399	656	1,029	2,697	8,949	18,920	1
2000	34,148	94	31	24	69	134	401	645	1,076	2,887	9,421	19,366	-
1999	33,697	121	52	30	67	129	441	592	1,046	2,724	9,176	19,318	1

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Pneumonia (J12-J18)													
Non-Hispanic white, both sexes													
2005 2/	50,777	88	36	57	113	216	580	1,419	2,400	4,898	15,188	25,781	1
2004 2/	48,452	106	43	35	103	180	538	1,225	2,245	4,806	14,875	24,296	-
2003 2/	53,039	130	51	64	121	198	620	1,377	2,141	5,095	16,119	27,122	1
2002	54,756	111	44	49	98	204	610	1,314	2,127	5,374	16,833	27,989	3
2001	51,736	110	46	51	104	200	602	1,179	1,946	5,196	15,790	26,511	1
2000	53,522	109	43	43	105	206	662	1,127	1,976	5,370	16,610	27,270	1
1999	52,758	118	53	52	87	191	629	1,089	1,834	5,342	16,368	26,994	1
Non-Hispanic white, male													
2005 2/	22,318	56	16	24	78	115	326	832	1,368	2,697	7,586	9,219	1
2004 2/	21,528	63	18	17	72	114	294	735	1,305	2,698	7,437	8,775	-
2003 2/	22,942	80	29	30	69	124	352	817	1,186	2,885	7,966	9,404	-
2002	23,676	72	23	20	60	110	337	757	1,212	3,023	8,326	9,734	2
2001	22,308	65	26	26	61	109	331	699	1,126	2,913	7,812	9,139	1
2000	23,014	63	21	26	64	109	385	649	1,137	2,987	8,165	9,407	1
1999	22,550	63	19	31	49	114	330	659	1,017	3,028	8,063	9,177	-
Non-Hispanic white, female													
2005 2/	28,459	32	20	33	35	101	254	587	1,032	2,201	7,602	16,562	-
2004 2/	26,924	43	25	18	31	66	244	490	940	2,108	7,438	15,521	-
2003 2/	30,097	50	22	34	52	74	268	560	955	2,210	8,153	17,718	1
2002	31,080	39	21	29	38	94	273	557	915	2,351	8,507	18,255	1
2001	29,428	45	20	25	43	91	271	480	820	2,283	7,978	17,372	-
2000	30,508	46	22	17	41	97	277	478	839	2,383	8,445	17,863	-
1999	30,208	55	34	21	38	77	299	430	817	2,314	8,305	17,817	1

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Pneumonia (J12-J18)													
Non-Hispanic black, both sexes													
2005 2/	5,649	90	26	17	30	76	208	487	597	909	1,483	1,726	-
2004 2/	5,570	89	31	19	39	62	222	440	533	843	1,523	1,769	-
2003 2/	5,703	84	21	18	35	98	250	475	558	905	1,502	1,756	1
2002	5,773	85	37	16	36	80	250	399	550	831	1,580	1,909	-
2001	5,666	103	24	15	44	85	273	430	468	794	1,567	1,863	-
2000	5,839	99	28	10	43	91	251	416	525	968	1,537	1,871	-
1999	5,729	110	39	15	48	106	300	422	481	868	1,395	1,945	-
Non-Hispanic black, male													
2005 2/	2,666	45	15	10	22	44	124	294	367	508	680	557	-
2004 2/	2,704	51	10	12	19	40	126	264	310	510	765	597	-
2003 2/	2,825	49	10	11	21	57	141	279	351	543	762	600	1
2002	2,719	46	20	10	18	43	136	239	335	451	787	634	-
2001	2,759	52	13	9	22	50	155	270	290	470	808	620	-
2000	2,845	55	20	6	20	57	148	266	319	557	758	639	-
1999	2,762	55	24	8	23	58	171	275	281	512	682	673	-
Non-Hispanic black, female													
2005 2/	2,983	45	11	7	8	32	84	193	230	401	803	1,169	-
2004 2/	2,866	38	21	7	20	22	96	176	223	333	758	1,172	-
2003 2/	2,878	35	11	7	14	41	109	196	207	362	740	1,156	-
2002	3,054	39	17	6	18	37	114	160	215	380	793	1,275	-
2001	2,907	51	11	6	22	35	118	160	178	324	759	1,243	-
2000	2,994	44	8	4	23	34	103	150	206	411	779	1,232	-
1999	2,967	55	15	7	25	48	129	147	200	356	713	1,272	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Pneumonia (J12-J18)													
Origin not stated 3/, both sexes													
2005	107	2	-	1	1	1	3	8	7	17	38	28	1
2004	168	-	-	-	2	-	8	10	23	30	43	52	-
2003	170	2	-	1	-	-	1	14	22	30	36	64	-
2002	196	2	1	1	1	2	5	11	7	24	69	73	-
2001	209	2	1	-	-	4	4	11	20	39	53	73	2
2000	246	1	2	-	1	2	12	15	14	34	79	84	2
1999	258	1	3	-	1	-	10	16	24	32	67	103	1
Origin not stated 3/, male													
2005	64	2	-	-	1	1	2	5	7	13	22	10	1
2004	99	-	-	-	1	-	5	9	18	25	20	21	-
2003	92	2	-	-	-	-	1	12	14	21	18	24	-
2002	101	1	-	-	1	2	3	9	5	21	35	24	-
2001	107	1	-	-	-	2	4	7	13	28	25	25	2
2000	136	1	2	-	1	2	8	12	12	23	44	30	1
1999	136	-	1	-	-	-	7	14	15	18	36	44	1
Origin not stated 3/, female													
2005	43	-	-	1	-	-	1	3	-	4	16	18	-
2004	69	-	-	-	1	-	3	1	5	5	23	31	-
2003	78	-	-	1	-	-	-	2	8	9	18	40	-
2002	95	1	1	1	-	-	2	2	2	3	34	49	-
2001	102	1	1	-	-	2	-	4	7	11	28	48	-
2000	110	-	-	-	-	-	4	3	2	11	35	54	1
1999	122	1	2	-	1	-	3	2	9	14	31	59	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Other acute lower respiratory infections (J20-J22)													
All origins, both sexes													
2005	404	51	16	4	4	6	15	15	19	26	78	170	-
2004	319	60	16	1	4	3	9	10	14	21	54	127	-
2003	421	54	28	6	6	2	11	19	21	30	80	164	-
2002	386	52	23	8	3	5	8	18	18	33	66	152	-
2001	347	50	18	3	1	2	12	14	16	28	58	145	-
2000	425	43	11	7	5	8	12	12	10	43	82	192	-
1999	540	63	25	5	5	4	11	15	21	35	107	249	-
All origins, male													
2005	172	30	10	4	3	2	6	12	5	10	41	49	-
2004	137	34	9	1	3	2	6	8	6	12	18	38	-
2003	170	35	17	2	2	1	6	8	12	17	35	35	-
2002	154	39	9	7	-	1	3	10	6	13	27	39	-
2001	117	26	8	2	1	-	7	7	6	10	20	30	-
2000	160	25	4	3	4	6	7	4	3	22	35	47	-
1999	196	36	15	2	1	3	5	8	10	17	51	48	-
All origins, female													
2005	232	21	6	-	1	4	9	3	14	16	37	121	-
2004	182	26	7	-	1	1	3	2	8	9	36	89	-
2003	251	19	11	4	4	1	5	11	9	13	45	129	-
2002	232	13	14	1	3	4	5	8	12	20	39	113	-
2001	230	24	10	1	-	2	5	7	10	18	38	115	-
2000	265	18	7	4	1	2	5	8	7	21	47	145	-
1999	344	27	10	3	4	1	6	7	11	18	56	201	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Other acute lower respiratory infections (J20-J22)													
Hispanic, both sexes													
2005	19	8	3	-	-	1	-	-	2	1	1	3	-
2004	32	15	5	1	-	1	1	-	2	2	1	4	-
2003	31	17	6	-	-	-	1	-	1	1	2	3	-
2002	21	10	2	2	-	1	-	-	2	-	-	4	-
2001	22	8	1	-	-	1	-	2	-	2	4	4	-
2000	24	9	2	-	-	1	-	1	-	1	2	8	-
1999	21	10	3	1	-	-	-	1	-	1	-	5	-
Hispanic, male													
2005	7	3	2	-	-	-	-	-	-	-	1	1	-
2004	15	8	2	1	-	-	-	-	2	-	-	2	-
2003	20	12	4	-	-	-	-	-	1	-	1	2	-
2002	9	5	1	1	-	-	-	-	1	-	-	1	-
2001	9	6	-	-	-	-	-	1	-	1	1	-	-
2000	15	6	1	-	-	1	-	1	-	1	2	3	-
1999	10	5	2	1	-	-	-	-	-	1	-	1	-
Hispanic, female													
2005	12	5	1	-	-	1	-	-	2	1	-	2	-
2004	17	7	3	-	-	1	1	-	-	2	1	2	-
2003	11	5	2	-	-	-	1	-	-	1	1	1	-
2002	12	5	1	1	-	1	-	-	1	-	-	3	-
2001	13	2	1	-	-	1	-	1	-	1	3	4	-
2000	9	3	1	-	-	-	-	-	-	-	-	5	-
1999	11	5	1	-	-	-	-	1	-	-	-	4	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Other acute lower respiratory infections (J20-J22)													
Non-Hispanic 1/, both sexes													
2005	385	43	13	4	4	5	15	15	17	25	77	167	-
2004	286	45	11	-	4	2	8	9	12	19	53	123	-
2003	388	37	22	6	6	2	10	19	19	29	77	161	-
2002	365	42	21	6	3	4	8	18	16	33	66	148	-
2001	325	42	17	3	1	1	12	12	16	26	54	141	-
2000	398	33	9	7	5	7	12	11	10	42	79	183	-
1999	518	53	22	4	5	4	10	14	21	34	107	244	-
Non-Hispanic 1/, male													
2005	165	27	8	4	3	2	6	12	5	10	40	48	-
2004	121	26	7	-	3	2	6	7	4	12	18	36	-
2003	148	23	13	2	2	1	6	8	10	17	33	33	-
2002	145	34	8	6	-	1	3	10	5	13	27	38	-
2001	108	20	8	2	1	-	7	6	6	9	19	30	-
2000	144	19	3	3	4	5	7	3	3	21	32	44	-
1999	185	31	13	1	1	3	4	8	10	16	51	47	-
Non-Hispanic 1/, female													
2005	220	16	5	-	1	3	9	3	12	15	37	119	-
2004	165	19	4	-	1	-	2	2	8	7	35	87	-
2003	240	14	9	4	4	1	4	11	9	12	44	128	-
2002	220	8	13	-	3	3	5	8	11	20	39	110	-
2001	217	22	9	1	-	1	5	6	10	17	35	111	-
2000	254	14	6	4	1	2	5	8	7	21	47	139	-
1999	333	22	9	3	4	1	6	6	11	18	56	197	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Other acute lower respiratory infections (J20-J22)													
Non-Hispanic white, both sexes													
2005 2/	334	23	11	3	4	4	12	11	16	21	71	158	-
2004 2/	246	28	4	-	3	2	8	7	11	17	48	118	-
2003 2/	342	25	11	5	3	1	10	17	16	25	73	156	-
2002	314	25	18	6	2	3	7	14	12	28	59	140	-
2001	278	24	12	3	1	-	12	8	14	20	50	134	-
2000	345	15	6	3	5	4	8	8	10	40	74	172	-
1999	440	26	12	3	4	3	8	10	17	26	99	232	-
Non-Hispanic white, male													
2005 2/	141	16	7	3	3	2	4	8	5	8	39	46	-
2004 2/	99	16	2	-	2	2	6	5	4	11	17	34	-
2003 2/	125	15	6	2	-	1	6	8	8	15	31	33	-
2002	122	22	6	6	-	-	3	9	3	12	25	36	-
2001	86	11	5	2	1	-	7	3	5	7	17	28	-
2000	118	6	3	2	4	3	6	3	3	19	30	39	-
1999	145	14	6	1	1	2	3	8	9	11	48	42	-
Non-Hispanic white, female													
2005 2/	193	7	4	-	1	2	8	3	11	13	32	112	-
2004 2/	147	12	2	-	1	-	2	2	7	6	31	84	-
2003 2/	217	10	5	3	3	-	4	9	8	10	42	123	-
2002	192	3	12	-	2	3	4	5	9	16	34	104	-
2001	192	13	7	1	-	-	5	5	9	13	33	106	-
2000	227	9	3	1	1	1	2	5	7	21	44	133	-
1999	295	12	6	2	3	1	5	2	8	15	51	190	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Other acute lower respiratory infections (J20-J22)													
Non-Hispanic black, both sexes													
2005 2/	40	18	1	1	-	1	3	2	1	3	4	6	-
2004 2/	31	13	6	-	1	-	-	1	-	2	5	3	-
2003 2/	40	12	7	1	3	1	-	2	2	4	3	5	-
2002	44	15	2	-	1	1	1	4	3	5	7	5	-
2001	33	14	2	-	-	1	-	4	1	4	-	7	-
2000	46	14	3	3	-	2	4	3	-	2	5	10	-
1999	64	24	6	-	1	1	2	4	2	8	6	10	-
Non-Hispanic black, male													
2005 2/	17	9	-	1	-	-	2	2	-	1	-	2	-
2004 2/	15	6	4	-	1	-	-	1	-	1	1	1	-
2003 2/	21	8	6	-	2	-	-	-	1	2	2	-	-
2002	22	12	1	-	-	1	-	1	2	1	2	2	-
2001	15	8	1	-	-	-	-	3	-	1	-	2	-
2000	22	10	-	1	-	2	1	-	-	2	2	4	-
1999	30	15	3	-	-	1	1	-	1	5	1	3	-
Non-Hispanic black, female													
2005 2/	23	9	1	-	-	1	1	-	1	2	4	4	-
2004 2/	16	7	2	-	-	-	-	-	-	1	4	2	-
2003 2/	19	4	1	1	1	1	-	2	1	2	1	5	-
2002	22	3	1	-	1	-	1	3	1	4	5	3	-
2001	18	6	1	-	-	1	-	1	1	3	-	5	-
2000	24	4	3	2	-	-	3	3	-	-	3	6	-
1999	34	9	3	-	1	-	1	4	1	3	5	7	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Other acute lower respiratory infections (J20-J22)													
Origin not stated 3/, both sexes													
2005	-	-	-	-	-	-	-	-	-	-	-	-	-
2004	1	-	-	-	-	-	-	1	-	-	-	-	-
2003	2	-	-	-	-	-	-	-	1	-	1	-	-
2002	-	-	-	-	-	-	-	-	-	-	-	-	-
2001	-	-	-	-	-	-	-	-	-	-	-	-	-
2000	3	1	-	-	-	-	-	-	-	-	1	1	-
1999	1	-	-	-	-	-	1	-	-	-	-	-	-
Origin not stated 3/, male													
2005	-	-	-	-	-	-	-	-	-	-	-	-	-
2004	1	-	-	-	-	-	-	1	-	-	-	-	-
2003	2	-	-	-	-	-	-	-	1	-	1	-	-
2002	-	-	-	-	-	-	-	-	-	-	-	-	-
2001	-	-	-	-	-	-	-	-	-	-	-	-	-
2000	1	-	-	-	-	-	-	-	-	-	1	-	-
1999	1	-	-	-	-	-	1	-	-	-	-	-	-
Origin not stated 3/, female													
2005	-	-	-	-	-	-	-	-	-	-	-	-	-
2004	-	-	-	-	-	-	-	-	-	-	-	-	-
2003	-	-	-	-	-	-	-	-	-	-	-	-	-
2002	-	-	-	-	-	-	-	-	-	-	-	-	-
2001	-	-	-	-	-	-	-	-	-	-	-	-	-
2000	2	1	-	-	-	-	-	-	-	-	-	1	-
1999	-	-	-	-	-	-	-	-	-	-	-	-	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Acute bronchitis and bronchiolitis (J20-J21)													
All origins, both sexes													
2005	283	50	14	4	3	3	13	15	13	17	53	98	-
2004	226	58	16	1	4	3	8	7	10	18	32	69	-
2003	304	53	27	6	6	2	9	16	17	22	45	101	-
2002	279	49	23	8	2	5	6	17	13	23	39	94	-
2001	263	49	18	3	1	2	11	12	14	23	39	91	-
2000	290	42	11	6	4	6	10	9	7	31	53	111	-
1999	388	62	24	4	5	3	10	14	18	29	69	150	-
All origins, male													
2005	123	29	8	4	2	-	5	12	3	7	26	27	-
2004	108	33	9	1	3	2	5	5	6	10	7	27	-
2003	126	34	16	2	2	1	5	7	11	12	17	19	-
2002	122	37	9	7	-	1	3	9	5	10	18	23	-
2001	89	25	8	2	1	-	6	6	6	8	10	17	-
2000	119	24	4	2	3	4	6	3	3	14	27	29	-
1999	154	36	14	2	1	2	4	7	8	15	34	31	-
All origins, female													
2005	160	21	6	-	1	3	8	3	10	10	27	71	-
2004	118	25	7	-	1	1	3	2	4	8	25	42	-
2003	178	19	11	4	4	1	4	9	6	10	28	82	-
2002	157	12	14	1	2	4	3	8	8	13	21	71	-
2001	174	24	10	1	-	2	5	6	8	15	29	74	-
2000	171	18	7	4	1	2	4	6	4	17	26	82	-
1999	234	26	10	2	4	1	6	7	10	14	35	119	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Acute bronchitis and bronchiolitis (J20-J21)													
Hispanic, both sexes													
2005	17	8	3	-	-	-	-	-	2	1	-	3	-
2004	30	14	5	1	-	1	1	-	2	2	1	3	-
2003	31	17	6	-	-	-	1	-	1	1	2	3	-
2002	19	9	2	2	-	1	-	-	2	-	-	3	-
2001	17	8	1	-	-	1	-	2	-	1	2	2	-
2000	22	9	2	-	-	1	-	1	-	1	2	6	-
1999	20	10	3	1	-	-	-	1	-	1	-	4	-
Hispanic, male													
2005	6	3	2	-	-	-	-	-	-	-	-	1	-
2004	14	7	2	1	-	-	-	-	2	-	-	2	-
2003	20	12	4	-	-	-	-	-	1	-	1	2	-
2002	9	5	1	1	-	-	-	-	1	-	-	1	-
2001	7	6	-	-	-	-	-	1	-	-	-	-	-
2000	15	6	1	-	-	1	-	1	-	1	2	3	-
1999	10	5	2	1	-	-	-	-	-	1	-	1	-
Hispanic, female													
2005	11	5	1	-	-	-	-	-	2	1	-	2	-
2004	16	7	3	-	-	1	1	-	-	2	1	1	-
2003	11	5	2	-	-	-	1	-	-	1	1	1	-
2002	10	4	1	1	-	1	-	-	1	-	-	2	-
2001	10	2	1	-	-	1	-	1	-	1	2	2	-
2000	7	3	1	-	-	-	-	-	-	-	-	3	-
1999	10	5	1	-	-	-	-	1	-	-	-	3	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Acute bronchitis and bronchiolitis (J20-J21)													
Non-Hispanic 1/, both sexes													
2005	266	42	11	4	3	3	13	15	11	16	53	95	-
2004	195	44	11	-	4	2	7	6	8	16	31	66	-
2003	272	36	21	6	6	2	8	16	15	21	43	98	-
2002	260	40	21	6	2	4	6	17	11	23	39	91	-
2001	246	41	17	3	1	1	11	10	14	22	37	89	-
2000	267	32	9	6	4	5	10	8	7	30	51	105	-
1999	367	52	21	3	5	3	9	13	18	28	69	146	-
Non-Hispanic 1/, male													
2005	117	26	6	4	2	-	5	12	3	7	26	26	-
2004	93	26	7	-	3	2	5	4	4	10	7	25	-
2003	105	22	12	2	2	1	5	7	9	12	16	17	-
2002	113	32	8	6	-	1	3	9	4	10	18	22	-
2001	82	19	8	2	1	-	6	5	6	8	10	17	-
2000	104	18	3	2	3	3	6	2	3	13	25	26	-
1999	143	31	12	1	1	2	3	7	8	14	34	30	-
Non-Hispanic 1/, female													
2005	149	16	5	-	1	3	8	3	8	9	27	69	-
2004	102	18	4	-	1	-	2	2	4	6	24	41	-
2003	167	14	9	4	4	1	3	9	6	9	27	81	-
2002	147	8	13	-	2	3	3	8	7	13	21	69	-
2001	164	22	9	1	-	1	5	5	8	14	27	72	-
2000	163	14	6	4	1	2	4	6	4	17	26	79	-
1999	224	21	9	2	4	1	6	6	10	14	35	116	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Acute bronchitis and bronchiolitis (J20-J21)													
Non-Hispanic white, both sexes													
2005 2/	225	22	10	3	3	2	11	11	11	13	50	89	-
2004 2/	159	28	4	-	3	2	7	5	7	14	27	62	-
2003 2/	231	24	10	5	3	1	8	15	12	17	41	95	-
2002	219	23	18	6	1	3	6	13	8	21	36	84	-
2001	202	23	12	3	1	-	11	6	12	17	34	83	-
2000	221	15	6	2	4	2	6	5	7	28	47	99	-
1999	299	25	12	2	4	3	7	9	14	20	63	140	-
Non-Hispanic white, male													
2005 2/	96	15	6	3	2	-	4	8	3	6	25	24	-
2004 2/	72	16	2	-	2	2	5	3	4	9	6	23	-
2003 2/	83	14	5	2	-	1	5	7	7	10	15	17	-
2002	92	20	6	6	-	-	3	8	3	9	17	20	-
2001	61	10	5	2	1	-	6	2	5	6	8	16	-
2000	83	6	3	1	3	1	5	2	3	11	24	24	-
1999	108	14	6	1	1	2	2	7	7	9	32	27	-
Non-Hispanic white, female													
2005 2/	129	7	4	-	1	2	7	3	8	7	25	65	-
2004 2/	87	12	2	-	1	-	2	2	3	5	21	39	-
2003 2/	148	10	5	3	3	-	3	8	5	7	26	78	-
2002	127	3	12	-	1	3	3	5	5	12	19	64	-
2001	141	13	7	1	-	-	5	4	7	11	26	67	-
2000	138	9	3	1	1	1	1	3	4	17	23	75	-
1999	191	11	6	1	3	1	5	2	7	11	31	113	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Acute bronchitis and bronchiolitis (J20-J21)													
Non-Hispanic black, both sexes													
2005 2/	31	18	1	1	-	1	2	2	-	2	1	3	-
2004 2/	28	12	6	-	1	-	-	1	-	2	4	2	-
2003 2/	35	12	7	1	3	1	-	1	2	4	1	3	-
2002	35	15	2	-	1	1	-	4	2	2	3	5	-
2001	31	14	2	-	-	1	-	4	1	3	-	6	-
2000	40	13	3	3	-	2	4	3	-	2	4	6	-
1999	55	24	5	-	1	-	2	4	2	8	4	5	-
Non-Hispanic black, male													
2005 2/	15	9	-	1	-	-	1	2	-	-	-	2	-
2004 2/	15	6	4	-	1	-	-	1	-	1	1	1	-
2003 2/	20	8	6	-	2	-	-	-	1	2	1	-	-
2002	20	12	1	-	-	1	-	1	1	1	1	2	-
2001	14	8	1	-	-	-	-	3	-	1	-	1	-
2000	18	9	-	1	-	2	1	-	-	2	1	2	-
1999	26	15	2	-	-	-	1	-	1	5	-	2	-
Non-Hispanic black, female													
2005 2/	16	9	1	-	-	1	1	-	-	2	1	1	-
2004 2/	13	6	2	-	-	-	-	-	-	1	3	1	-
2003 2/	15	4	1	1	1	1	-	1	1	2	-	3	-
2002	15	3	1	-	1	-	-	3	1	1	2	3	-
2001	17	6	1	-	-	1	-	1	1	2	-	5	-
2000	22	4	3	2	-	-	3	3	-	-	3	4	-
1999	29	9	3	-	1	-	1	4	1	3	4	3	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Acute bronchitis and bronchiolitis (J20-J21)													
Origin not stated 3/, both sexes													
2005	-	-	-	-	-	-	-	-	-	-	-	-	-
2004	1	-	-	-	-	-	-	1	-	-	-	-	-
2003	1	-	-	-	-	-	-	-	1	-	-	-	-
2002	-	-	-	-	-	-	-	-	-	-	-	-	-
2001	-	-	-	-	-	-	-	-	-	-	-	-	-
2000	1	1	-	-	-	-	-	-	-	-	-	-	-
1999	1	-	-	-	-	-	1	-	-	-	-	-	-
Origin not stated 3/, male													
2005	-	-	-	-	-	-	-	-	-	-	-	-	-
2004	1	-	-	-	-	-	-	1	-	-	-	-	-
2003	1	-	-	-	-	-	-	-	1	-	-	-	-
2002	-	-	-	-	-	-	-	-	-	-	-	-	-
2001	-	-	-	-	-	-	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-	-	-	-
1999	1	-	-	-	-	-	1	-	-	-	-	-	-
Origin not stated 3/, female													
2005	-	-	-	-	-	-	-	-	-	-	-	-	-
2004	-	-	-	-	-	-	-	-	-	-	-	-	-
2003	-	-	-	-	-	-	-	-	-	-	-	-	-
2002	-	-	-	-	-	-	-	-	-	-	-	-	-
2001	-	-	-	-	-	-	-	-	-	-	-	-	-
2000	1	1	-	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-	-	-	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Unspecified acute lower respiratory infection (J22)													
All origins, both sexes													
2005	121	1	2	-	1	3	2	-	6	9	25	72	-
2004	93	2	-	-	-	-	1	3	4	3	22	58	-
2003	117	1	1	-	-	-	2	3	4	8	35	63	-
2002	107	3	-	-	1	-	2	1	5	10	27	58	-
2001	84	1	-	-	-	-	1	2	2	5	19	54	-
2000	135	1	-	1	1	2	2	3	3	12	29	81	-
1999	152	1	1	1	-	1	1	1	3	6	38	99	-
All origins, male													
2005	49	1	2	-	1	2	1	-	2	3	15	22	-
2004	29	1	-	-	-	-	1	3	-	2	11	11	-
2003	44	1	1	-	-	-	1	1	1	5	18	16	-
2002	32	2	-	-	-	-	-	1	1	3	9	16	-
2001	28	1	-	-	-	-	1	1	-	2	10	13	-
2000	41	1	-	1	1	2	1	1	-	8	8	18	-
1999	42	-	1	-	-	1	1	1	2	2	17	17	-
All origins, female													
2005	72	-	-	-	-	1	1	-	4	6	10	50	-
2004	64	1	-	-	-	-	-	-	4	1	11	47	-
2003	73	-	-	-	-	-	1	2	3	3	17	47	-
2002	75	1	-	-	1	-	2	-	4	7	18	42	-
2001	56	-	-	-	-	-	-	1	2	3	9	41	-
2000	94	-	-	-	-	-	1	2	3	4	21	63	-
1999	110	1	-	1	-	-	-	-	1	4	21	82	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Unspecified acute lower respiratory infection (J22)													
Hispanic, both sexes													
2005	2	-	-	-	-	1	-	-	-	-	1	-	-
2004	2	1	-	-	-	-	-	-	-	-	-	1	-
2003	-	-	-	-	-	-	-	-	-	-	-	-	-
2002	2	1	-	-	-	-	-	-	-	-	-	1	-
2001	5	-	-	-	-	-	-	-	-	1	2	2	-
2000	2	-	-	-	-	-	-	-	-	-	-	2	-
1999	1	-	-	-	-	-	-	-	-	-	-	1	-
Hispanic, male													
2005	1	-	-	-	-	-	-	-	-	-	1	-	-
2004	1	1	-	-	-	-	-	-	-	-	-	-	-
2003	-	-	-	-	-	-	-	-	-	-	-	-	-
2002	-	-	-	-	-	-	-	-	-	-	-	-	-
2001	2	-	-	-	-	-	-	-	-	1	1	-	-
2000	-	-	-	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-	-	-	-
Hispanic, female													
2005	1	-	-	-	-	1	-	-	-	-	-	-	-
2004	1	-	-	-	-	-	-	-	-	-	-	1	-
2003	-	-	-	-	-	-	-	-	-	-	-	-	-
2002	2	1	-	-	-	-	-	-	-	-	-	1	-
2001	3	-	-	-	-	-	-	-	-	-	1	2	-
2000	2	-	-	-	-	-	-	-	-	-	-	2	-
1999	1	-	-	-	-	-	-	-	-	-	-	1	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Unspecified acute lower respiratory infection (J22)													
Non-Hispanic 1/, both sexes													
2005	119	1	2	-	1	2	2	-	6	9	24	72	-
2004	91	1	-	-	-	-	1	3	4	3	22	57	-
2003	116	1	1	-	-	-	2	3	4	8	34	63	-
2002	105	2	-	-	1	-	2	1	5	10	27	57	-
2001	79	1	-	-	-	-	1	2	2	4	17	52	-
2000	131	1	-	1	1	2	2	3	3	12	28	78	-
1999	151	1	1	1	-	1	1	1	3	6	38	98	-
Non-Hispanic 1/, male													
2005	48	1	2	-	1	2	1	-	2	3	14	22	-
2004	28	-	-	-	-	-	1	3	-	2	11	11	-
2003	43	1	1	-	-	-	1	1	1	5	17	16	-
2002	32	2	-	-	-	-	-	1	1	3	9	16	-
2001	26	1	-	-	-	-	1	1	-	1	9	13	-
2000	40	1	-	1	1	2	1	1	-	8	7	18	-
1999	42	-	1	-	-	1	1	1	2	2	17	17	-
Non-Hispanic 1/, female													
2005	71	-	-	-	-	-	1	-	4	6	10	50	-
2004	63	1	-	-	-	-	-	-	4	1	11	46	-
2003	73	-	-	-	-	-	1	2	3	3	17	47	-
2002	73	-	-	-	1	-	2	-	4	7	18	41	-
2001	53	-	-	-	-	-	-	1	2	3	8	39	-
2000	91	-	-	-	-	-	1	2	3	4	21	60	-
1999	109	1	-	1	-	-	-	-	1	4	21	81	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Unspecified acute lower respiratory infection (J22)													
Non-Hispanic white, both sexes													
2005 2/	109	1	1	-	1	2	1	-	5	8	21	69	-
2004 2/	87	-	-	-	-	-	1	2	4	3	21	56	-
2003 2/	111	1	1	-	-	-	2	2	4	8	32	61	-
2002	95	2	-	-	1	-	1	1	4	7	23	56	-
2001	76	1	-	-	-	-	1	2	2	3	16	51	-
2000	124	-	-	1	1	2	2	3	3	12	27	73	-
1999	141	1	-	1	-	-	1	1	3	6	36	92	-
Non-Hispanic white, male													
2005 2/	45	1	1	-	1	2	-	-	2	2	14	22	-
2004 2/	27	-	-	-	-	-	1	2	-	2	11	11	-
2003 2/	42	1	1	-	-	-	1	1	1	5	16	16	-
2002	30	2	-	-	-	-	-	1	-	3	8	16	-
2001	25	1	-	-	-	-	1	1	-	1	9	12	-
2000	35	-	-	1	1	2	1	1	-	8	6	15	-
1999	37	-	-	-	-	-	1	1	2	2	16	15	-
Non-Hispanic white, female													
2005 2/	64	-	-	-	-	-	1	-	3	6	7	47	-
2004 2/	60	-	-	-	-	-	-	-	4	1	10	45	-
2003 2/	69	-	-	-	-	-	1	1	3	3	16	45	-
2002	65	-	-	-	1	-	1	-	4	4	15	40	-
2001	51	-	-	-	-	-	-	1	2	2	7	39	-
2000	89	-	-	-	-	-	1	2	3	4	21	58	-
1999	104	1	-	1	-	-	-	-	1	4	20	77	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Unspecified acute lower respiratory infection (J22)													
Non-Hispanic black, both sexes													
2005 2/	9	-	-	-	-	-	1	-	1	1	3	3	-
2004 2/	3	1	-	-	-	-	-	-	-	-	1	1	-
2003 2/	5	-	-	-	-	-	-	1	-	-	2	2	-
2002	9	-	-	-	-	-	1	-	1	3	4	-	-
2001	2	-	-	-	-	-	-	-	-	1	-	1	-
2000	6	1	-	-	-	-	-	-	-	-	1	4	-
1999	9	-	1	-	-	1	-	-	-	-	2	5	-
Non-Hispanic black, male													
2005 2/	2	-	-	-	-	-	1	-	-	1	-	-	-
2004 2/	-	-	-	-	-	-	-	-	-	-	-	-	-
2003 2/	1	-	-	-	-	-	-	-	-	-	1	-	-
2002	2	-	-	-	-	-	-	-	1	-	1	-	-
2001	1	-	-	-	-	-	-	-	-	-	-	1	-
2000	4	1	-	-	-	-	-	-	-	-	1	2	-
1999	4	-	1	-	-	1	-	-	-	-	1	1	-
Non-Hispanic black, female													
2005 2/	7	-	-	-	-	-	-	-	1	-	3	3	-
2004 2/	3	1	-	-	-	-	-	-	-	-	1	1	-
2003 2/	4	-	-	-	-	-	-	1	-	-	1	2	-
2002	7	-	-	-	-	-	1	-	-	3	3	-	-
2001	1	-	-	-	-	-	-	-	-	1	-	-	-
2000	2	-	-	-	-	-	-	-	-	-	-	2	-
1999	5	-	-	-	-	-	-	-	-	-	1	4	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Unspecified acute lower respiratory infection (J22)													
Origin not stated 3/, both sexes													
2005	-	-	-	-	-	-	-	-	-	-	-	-	-
2004	-	-	-	-	-	-	-	-	-	-	-	-	-
2003	1	-	-	-	-	-	-	-	-	-	1	-	-
2002	-	-	-	-	-	-	-	-	-	-	-	-	-
2001	-	-	-	-	-	-	-	-	-	-	-	-	-
2000	2	-	-	-	-	-	-	-	-	-	1	1	-
1999	-	-	-	-	-	-	-	-	-	-	-	-	-
Origin not stated 3/, male													
2005	-	-	-	-	-	-	-	-	-	-	-	-	-
2004	-	-	-	-	-	-	-	-	-	-	-	-	-
2003	1	-	-	-	-	-	-	-	-	-	1	-	-
2002	-	-	-	-	-	-	-	-	-	-	-	-	-
2001	-	-	-	-	-	-	-	-	-	-	-	-	-
2000	1	-	-	-	-	-	-	-	-	-	1	-	-
1999	-	-	-	-	-	-	-	-	-	-	-	-	-
Origin not stated 3/, female													
2005	-	-	-	-	-	-	-	-	-	-	-	-	-
2004	-	-	-	-	-	-	-	-	-	-	-	-	-
2003	-	-	-	-	-	-	-	-	-	-	-	-	-
2002	-	-	-	-	-	-	-	-	-	-	-	-	-
2001	-	-	-	-	-	-	-	-	-	-	-	-	-
2000	1	-	-	-	-	-	-	-	-	-	-	1	-
1999	-	-	-	-	-	-	-	-	-	-	-	-	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Chronic lower respiratory diseases (J40-J47)													
All origins, both sexes													
2005	130,933	33	56	104	148	258	890	3,977	12,747	29,910	50,333	32,473	4
2004	121,987	35	48	120	179	255	887	3,511	11,754	28,390	47,568	29,239	1
2003	126,382	31	55	118	191	282	950	3,537	12,077	29,919	49,286	29,934	2
2002	124,816	39	65	136	192	301	1,008	3,475	11,280	29,788	49,241	29,284	7
2001	123,013	41	43	104	171	290	969	3,324	11,166	30,751	47,762	28,391	1
2000	122,009	36	51	139	190	296	930	3,251	10,739	31,157	47,722	27,496	2
1999	124,181	36	54	139	209	330	890	3,110	11,297	32,644	48,635	26,833	4
All origins, male													
2005	62,435	19	33	63	95	130	426	2,034	6,589	15,414	24,479	13,150	3
2004	58,646	24	30	75	110	151	431	1,806	6,029	14,503	23,255	12,231	1
2003	60,714	23	38	75	114	156	429	1,826	6,243	15,333	23,963	12,512	2
2002	60,713	23	44	74	120	165	497	1,795	5,783	15,339	24,432	12,436	5
2001	59,697	22	29	61	99	146	455	1,702	5,702	15,877	23,691	11,912	1
2000	60,004	23	28	80	120	165	437	1,661	5,564	16,210	23,838	11,876	2
1999	62,415	19	30	93	123	152	410	1,551	5,958	17,477	24,705	11,894	3
All origins, female													
2005	68,498	14	23	41	53	128	464	1,943	6,158	14,496	25,854	19,323	1
2004	63,341	11	18	45	69	104	456	1,705	5,725	13,887	24,313	17,008	-
2003	65,668	8	17	43	77	126	521	1,711	5,834	14,586	25,323	17,422	-
2002	64,103	16	21	62	72	136	511	1,680	5,497	14,449	24,809	16,848	2
2001	63,316	19	14	43	72	144	514	1,622	5,464	14,874	24,071	16,479	-
2000	62,005	13	23	59	70	131	493	1,590	5,175	14,947	23,884	15,620	-
1999	61,766	17	24	46	86	178	480	1,559	5,339	15,167	23,930	14,939	1

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Chronic lower respiratory diseases (J40-J47)													
Hispanic, both sexes													
2005	3,457	9	5	15	8	25	53	151	298	668	1,254	969	2
2004	3,107	9	11	12	17	30	66	124	272	657	1,086	823	-
2003	3,174	7	11	15	20	29	58	134	266	651	1,095	888	-
2002	3,058	4	9	13	15	31	70	127	274	635	1,055	825	-
2001	2,832	7	8	9	10	22	52	124	258	634	940	768	-
2000	2,689	7	9	15	16	20	55	105	244	572	921	725	-
1999	2,859	6	8	18	15	31	79	115	239	649	935	764	-
Hispanic, male													
2005	1,839	6	3	9	7	16	34	77	180	381	701	423	2
2004	1,631	6	7	7	11	20	45	56	145	387	577	370	-
2003	1,697	5	8	10	12	15	30	61	164	369	603	420	-
2002	1,625	2	4	8	12	19	31	70	164	362	593	360	-
2001	1,482	4	4	6	6	13	21	54	140	378	501	355	-
2000	1,451	4	5	9	13	12	24	53	146	343	509	333	-
1999	1,547	4	3	11	9	13	36	52	135	373	523	388	-
Hispanic, female													
2005	1,618	3	2	6	1	9	19	74	118	287	553	546	-
2004	1,476	3	4	5	6	10	21	68	127	270	509	453	-
2003	1,477	2	3	5	8	14	28	73	102	282	492	468	-
2002	1,433	2	5	5	3	12	39	57	110	273	462	465	-
2001	1,350	3	4	3	4	9	31	70	118	256	439	413	-
2000	1,238	3	4	6	3	8	31	52	98	229	412	392	-
1999	1,312	2	5	7	6	18	43	63	104	276	412	376	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Chronic lower respiratory diseases (J40-J47)													
Non-Hispanic 1/, both sexes													
2005	127,253	24	51	88	140	233	833	3,814	12,416	29,189	49,003	31,461	1
2004	118,650	26	37	108	162	224	818	3,375	11,455	27,656	46,413	28,375	1
2003	122,896	24	43	102	170	253	890	3,382	11,761	29,192	48,093	28,985	1
2002	121,424	34	56	121	176	267	930	3,335	10,975	29,069	48,058	28,397	6
2001	119,839	34	35	94	161	267	909	3,180	10,860	30,028	46,713	27,558	-
2000	118,963	29	42	123	173	272	872	3,122	10,447	30,486	46,687	26,709	1
1999	120,968	30	46	121	194	296	807	2,978	11,018	31,902	47,578	25,995	3
Non-Hispanic 1/, male													
2005	60,462	13	30	53	88	114	389	1,951	6,386	14,998	23,736	12,704	-
2004	56,884	18	23	68	99	131	386	1,742	5,867	14,066	22,638	11,845	1
2003	58,848	18	29	65	101	141	397	1,753	6,050	14,920	23,304	12,069	1
2002	58,896	20	40	64	107	143	460	1,719	5,604	14,924	23,761	12,050	4
2001	58,028	18	25	55	93	133	428	1,635	5,530	15,449	23,134	11,528	-
2000	58,349	19	23	71	107	150	411	1,593	5,388	15,805	23,266	11,515	1
1999	60,669	15	27	82	114	138	372	1,486	5,797	17,040	24,124	11,472	2
Non-Hispanic 1/, female													
2005	66,791	11	21	35	52	119	444	1,863	6,030	14,191	25,267	18,757	1
2004	61,766	8	14	40	63	93	432	1,633	5,588	13,590	23,775	16,530	-
2003	64,048	6	14	37	69	112	493	1,629	5,711	14,272	24,789	16,916	-
2002	62,528	14	16	57	69	124	470	1,616	5,371	14,145	24,297	16,347	2
2001	61,811	16	10	39	68	134	481	1,545	5,330	14,579	23,579	16,030	-
2000	60,614	10	19	52	66	122	461	1,529	5,059	14,681	23,421	15,194	-
1999	60,299	15	19	39	80	158	435	1,492	5,221	14,862	23,454	14,523	1

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Chronic lower respiratory diseases (J40-J47)													
Non-Hispanic white, both sexes													
2005 2/	117,337	11	20	31	64	129	593	3,054	11,058	26,824	45,941	29,611	1
2004 2/	109,670	14	12	41	75	120	583	2,695	10,190	25,565	43,613	26,761	1
2003 2/	113,541	17	15	39	82	140	617	2,725	10,465	26,902	45,177	27,361	1
2002	112,128	20	23	42	94	157	624	2,686	9,757	26,808	45,117	26,796	4
2001	110,753	16	13	39	75	134	613	2,570	9,678	27,693	43,972	25,950	-
2000	109,901	16	19	47	87	144	570	2,463	9,253	28,122	44,024	25,155	1
1999	111,620	10	23	55	110	146	533	2,372	9,786	29,400	44,662	24,521	2
Non-Hispanic white, male													
2005 2/	55,015	7	9	17	40	57	274	1,562	5,629	13,650	21,973	11,797	-
2004 2/	51,909	7	8	23	45	63	270	1,398	5,126	12,865	21,076	11,027	1
2003 2/	53,593	16	7	21	48	64	275	1,426	5,327	13,604	21,553	11,251	1
2002	53,662	13	17	23	48	81	304	1,383	4,909	13,615	22,034	11,232	3
2001	52,952	10	9	22	39	62	293	1,331	4,885	14,089	21,496	10,716	-
2000	53,231	12	8	24	52	71	267	1,259	4,728	14,398	21,703	10,708	1
1999	55,314	6	11	36	59	66	239	1,174	5,091	15,559	22,377	10,695	1
Non-Hispanic white, female													
2005 2/	62,322	4	11	14	24	72	319	1,492	5,429	13,174	23,968	17,814	1
2004 2/	57,761	7	4	18	30	57	313	1,297	5,064	12,700	22,537	15,734	-
2003 2/	59,948	1	8	18	34	76	342	1,299	5,138	13,298	23,624	16,110	-
2002	58,466	7	6	19	46	76	320	1,303	4,848	13,193	23,083	15,564	1
2001	57,801	6	4	17	36	72	320	1,239	4,793	13,604	22,476	15,234	-
2000	56,670	4	11	23	35	73	303	1,204	4,525	13,724	22,321	14,447	-
1999	56,306	4	12	19	51	80	294	1,198	4,695	13,841	22,285	13,826	1

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Chronic lower respiratory diseases (J40-J47)													
Non-Hispanic black, both sexes													
2005 2/	8,136	12	30	51	74	96	217	697	1,195	1,998	2,414	1,352	-
2004 2/	7,331	11	21	64	79	101	220	621	1,101	1,751	2,195	1,167	-
2003 2/	7,632	7	24	62	81	104	245	597	1,109	1,920	2,290	1,193	-
2002	7,730	14	29	74	78	102	292	589	1,067	1,892	2,366	1,226	1
2001	7,504	16	22	53	77	128	273	549	1,019	1,962	2,190	1,215	-
2000	7,528	11	21	70	80	114	272	593	1,063	1,993	2,128	1,183	-
1999	7,840	18	18	66	78	140	258	556	1,085	2,090	2,364	1,166	1
Non-Hispanic black, male													
2005 2/	4,406	6	21	33	47	52	103	351	660	1,146	1,360	627	-
2004 2/	4,010	10	14	44	48	66	107	317	644	1,000	1,200	560	-
2003 2/	4,202	2	19	43	48	72	106	293	615	1,093	1,340	571	-
2002	4,282	7	20	39	58	56	150	304	607	1,093	1,350	597	1
2001	4,143	6	16	32	48	69	124	271	554	1,147	1,285	591	-
2000	4,184	6	14	43	51	70	129	289	587	1,195	1,207	593	-
1999	4,452	9	13	46	52	67	121	284	624	1,238	1,402	595	1
Non-Hispanic black, female													
2005 2/	3,730	6	9	18	27	44	114	346	535	852	1,054	725	-
2004 2/	3,321	1	7	20	31	35	113	304	457	751	995	607	-
2003 2/	3,430	5	5	19	33	32	139	304	494	827	950	622	-
2002	3,448	7	9	35	20	46	142	285	460	799	1,016	629	-
2001	3,361	10	6	21	29	59	149	278	465	815	905	624	-
2000	3,344	5	7	27	29	44	143	304	476	798	921	590	-
1999	3,388	9	5	20	26	73	137	272	461	852	962	571	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Chronic lower respiratory diseases (J40-J47)													
Origin not stated 3/, both sexes													
2005	223	-	-	1	-	-	4	12	33	53	76	43	1
2004	230	-	-	-	-	1	3	12	27	77	69	41	-
2003	312	-	1	1	1	-	2	21	50	76	98	61	1
2002	334	1	-	2	1	3	8	13	31	84	128	62	1
2001	342	-	-	1	-	1	8	20	48	89	109	65	1
2000	357	-	-	1	1	4	3	24	48	99	114	62	1
1999	354	-	-	-	-	3	4	17	40	93	122	74	1
Origin not stated 3/, male													
2005	134	-	-	1	-	-	3	6	23	35	42	23	1
2004	131	-	-	-	-	-	-	8	17	50	40	16	-
2003	169	-	1	-	1	-	2	12	29	44	56	23	1
2002	192	1	-	2	1	3	6	6	15	53	78	26	1
2001	187	-	-	-	-	-	6	13	32	50	56	29	1
2000	204	-	-	-	-	3	2	15	30	62	63	28	1
1999	199	-	-	-	-	1	2	13	26	64	58	34	1
Origin not stated 3/, female													
2005	89	-	-	-	-	-	1	6	10	18	34	20	-
2004	99	-	-	-	-	1	3	4	10	27	29	25	-
2003	143	-	-	1	-	-	-	9	21	32	42	38	-
2002	142	-	-	-	-	-	2	7	16	31	50	36	-
2001	155	-	-	1	-	1	2	7	16	39	53	36	-
2000	153	-	-	1	1	1	1	9	18	37	51	34	-
1999	155	-	-	-	-	2	2	4	14	29	64	40	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Bronchitis, chronic and unspecified (J40-J42)													
All origins, both sexes													
2005	866	25	12	3	5	6	15	27	73	119	205	376	-
2004	773	17	12	4	7	5	12	35	67	97	199	318	-
2003	850	20	13	6	7	10	17	35	47	104	246	345	-
2002	955	24	13	5	6	6	18	36	54	135	286	372	-
2001	959	21	8	-	5	6	21	42	61	133	261	401	-
2000	1,167	18	14	5	5	8	20	38	67	158	333	501	-
1999	1,172	20	19	4	7	11	18	38	97	173	339	446	-
All origins, male													
2005	361	16	5	2	3	3	10	18	35	52	100	117	-
2004	338	12	9	4	4	4	5	22	35	54	83	106	-
2003	351	15	6	3	6	3	9	23	26	46	113	101	-
2002	420	15	8	2	3	2	12	27	28	74	127	122	-
2001	396	11	5	-	3	2	8	21	36	76	115	119	-
2000	503	9	7	3	4	7	14	16	32	79	174	158	-
1999	498	9	11	3	4	3	5	19	51	94	157	142	-
All origins, female													
2005	505	9	7	1	2	3	5	9	38	67	105	259	-
2004	435	5	3	-	3	1	7	13	32	43	116	212	-
2003	499	5	7	3	1	7	8	12	21	58	133	244	-
2002	535	9	5	3	3	4	6	9	26	61	159	250	-
2001	563	10	3	-	2	4	13	21	25	57	146	282	-
2000	664	9	7	2	1	1	6	22	35	79	159	343	-
1999	674	11	8	1	3	8	13	19	46	79	182	304	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Bronchitis, chronic and unspecified (J40-J42)													
Hispanic, both sexes													
2005	44	5	1	-	-	2	-	1	1	3	12	19	-
2004	38	4	2	1	-	-	-	2	3	3	13	10	-
2003	40	5	5	-	1	1	-	1	1	3	9	14	-
2002	51	3	1	-	2	1	3	3	2	2	14	20	-
2001	37	4	1	-	1	1	2	1	2	3	9	13	-
2000	52	4	3	-	-	2	3	1	1	9	13	16	-
1999	52	3	3	1	-	2	7	-	2	4	13	17	-
Hispanic, male													
2005	19	4	1	-	-	1	-	1	1	1	6	4	-
2004	19	3	2	1	-	-	-	1	2	-	6	4	-
2003	20	3	3	-	1	-	-	-	-	2	5	6	-
2002	20	1	1	-	1	1	2	3	1	1	4	5	-
2001	15	2	-	-	-	1	1	-	1	1	6	3	-
2000	28	2	2	-	-	2	2	-	1	7	6	6	-
1999	17	1	2	1	-	-	2	-	-	3	3	5	-
Hispanic, female													
2005	25	1	-	-	-	1	-	-	-	2	6	15	-
2004	19	1	-	-	-	-	-	1	1	3	7	6	-
2003	20	2	2	-	-	1	-	1	1	1	4	8	-
2002	31	2	-	-	1	-	1	-	1	1	10	15	-
2001	22	2	1	-	1	-	1	1	1	2	3	10	-
2000	24	2	1	-	-	-	1	1	-	2	7	10	-
1999	35	2	1	-	-	2	5	-	2	1	10	12	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Bronchitis, chronic and unspecified (J40-J42)													
Non-Hispanic 1/, both sexes													
2005	820	20	11	3	5	4	15	26	72	116	193	355	-
2004	733	13	10	3	7	5	12	33	64	94	186	306	-
2003	808	15	8	6	6	9	17	34	46	101	235	331	-
2002	900	20	12	5	4	5	15	33	52	132	271	351	-
2001	919	17	7	-	4	5	19	40	58	130	251	388	-
2000	1,112	14	11	5	5	6	17	37	66	149	320	482	-
1999	1,117	17	16	3	7	9	11	38	94	168	326	428	-
Non-Hispanic 1/, male													
2005	342	12	4	2	3	2	10	17	34	51	94	113	-
2004	318	9	7	3	4	4	5	21	33	54	77	101	-
2003	330	12	3	3	5	3	9	23	26	44	107	95	-
2002	398	13	7	2	2	1	10	24	27	73	122	117	-
2001	380	9	5	-	3	1	7	21	34	75	109	116	-
2000	474	7	5	3	4	5	12	16	31	72	168	151	-
1999	478	8	9	2	4	3	3	19	50	90	154	136	-
Non-Hispanic 1/, female													
2005	478	8	7	1	2	2	5	9	38	65	99	242	-
2004	415	4	3	-	3	1	7	12	31	40	109	205	-
2003	478	3	5	3	1	6	8	11	20	57	128	236	-
2002	502	7	5	3	2	4	5	9	25	59	149	234	-
2001	539	8	2	-	1	4	12	19	24	55	142	272	-
2000	638	7	6	2	1	1	5	21	35	77	152	331	-
1999	639	9	7	1	3	6	8	19	44	78	172	292	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Bronchitis, chronic and unspecified (J40-J42)													
Non-Hispanic white, both sexes													
2005 2/	738	10	9	3	4	4	11	22	57	103	176	339	-
2004 2/	653	8	4	2	5	5	8	24	53	88	174	282	-
2003 2/	712	11	4	3	5	5	13	23	37	92	214	305	-
2002	821	12	6	3	2	4	13	26	44	124	252	335	-
2001	839	7	5	-	3	4	16	35	54	113	232	370	-
2000	1,018	5	4	4	4	4	11	31	60	131	297	467	-
1999	1,002	8	12	3	6	6	9	24	86	150	293	405	-
Non-Hispanic white, male													
2005 2/	296	6	3	2	2	2	7	15	30	43	83	103	-
2004 2/	276	4	3	2	3	4	2	13	28	51	73	93	-
2003 2/	290	10	2	1	4	2	7	17	23	39	96	89	-
2002	354	9	2	2	1	1	8	17	22	67	113	112	-
2001	338	5	5	-	2	-	6	17	32	66	97	108	-
2000	429	3	1	2	3	3	6	15	27	64	159	146	-
1999	421	5	6	2	3	2	3	11	45	81	135	128	-
Non-Hispanic white, female													
2005 2/	442	4	6	1	2	2	4	7	27	60	93	236	-
2004 2/	377	4	1	-	2	1	6	11	25	37	101	189	-
2003 2/	422	1	2	2	1	3	6	6	14	53	118	216	-
2002	467	3	4	1	1	3	5	9	22	57	139	223	-
2001	501	2	-	-	1	4	10	18	22	47	135	262	-
2000	589	2	3	2	1	1	5	16	33	67	138	321	-
1999	581	3	6	1	3	4	6	13	41	69	158	277	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Bronchitis, chronic and unspecified (J40-J42)													
Non-Hispanic black, both sexes													
2005 2/	54	9	1	-	1	-	3	4	11	9	10	6	-
2004 2/	64	5	5	1	2	-	4	9	9	5	8	16	-
2003 2/	74	4	3	3	1	3	4	11	8	6	12	19	-
2002	66	8	6	2	2	1	2	7	5	6	13	14	-
2001	62	10	2	-	1	1	3	4	4	15	11	11	-
2000	80	7	6	1	1	1	6	6	4	16	20	12	-
1999	88	7	1	-	1	3	2	12	7	15	26	14	-
Non-Hispanic black, male													
2005 2/	33	6	1	-	1	-	2	2	4	6	7	4	-
2004 2/	34	5	4	1	1	-	3	8	3	3	2	4	-
2003 2/	29	2	-	2	1	1	2	6	3	2	7	3	-
2002	35	4	5	-	1	-	2	7	2	4	5	5	-
2001	32	4	-	-	1	1	1	4	2	8	6	5	-
2000	35	3	3	1	1	1	6	1	3	6	7	3	-
1999	43	3	1	-	1	1	-	7	4	7	16	3	-
Non-Hispanic black, female													
2005 2/	21	3	-	-	-	-	1	2	7	3	3	2	-
2004 2/	30	-	1	-	1	-	1	1	6	2	6	12	-
2003 2/	45	2	3	1	-	2	2	5	5	4	5	16	-
2002	31	4	1	2	1	1	-	-	3	2	8	9	-
2001	30	6	2	-	-	-	2	-	2	7	5	6	-
2000	45	4	3	-	-	-	-	5	1	10	13	9	-
1999	45	4	-	-	-	2	2	5	3	8	10	11	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Bronchitis, chronic and unspecified (J40-J42)													
Origin not stated 3/, both sexes													
2005	2	-	-	-	-	-	-	-	-	-	-	2	-
2004	2	-	-	-	-	-	-	-	-	-	-	2	-
2003	2	-	-	-	-	-	-	-	-	-	2	-	-
2002	4	1	-	-	-	-	-	-	-	1	1	1	-
2001	3	-	-	-	-	-	-	1	1	-	1	-	-
2000	3	-	-	-	-	-	-	-	-	-	-	3	-
1999	3	-	-	-	-	-	-	-	1	1	-	1	-
Origin not stated 3/, male													
2005	-	-	-	-	-	-	-	-	-	-	-	-	-
2004	1	-	-	-	-	-	-	-	-	-	-	1	-
2003	1	-	-	-	-	-	-	-	-	-	1	-	-
2002	2	1	-	-	-	-	-	-	-	-	1	-	-
2001	1	-	-	-	-	-	-	-	1	-	-	-	-
2000	1	-	-	-	-	-	-	-	-	-	-	1	-
1999	3	-	-	-	-	-	-	-	1	1	-	1	-
Origin not stated 3/, female													
2005	2	-	-	-	-	-	-	-	-	-	-	2	-
2004	1	-	-	-	-	-	-	-	-	-	-	1	-
2003	1	-	-	-	-	-	-	-	-	-	1	-	-
2002	2	-	-	-	-	-	-	-	-	1	-	1	-
2001	2	-	-	-	-	-	-	1	-	-	1	-	-
2000	2	-	-	-	-	-	-	-	-	-	-	2	-
1999	-	-	-	-	-	-	-	-	-	-	-	-	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Emphysema (J43)													
All origins, both sexes													
2005	14,002	2	1	-	-	8	77	444	1,677	3,749	5,400	2,643	1
2004	13,639	3	2	1	3	10	80	413	1,583	3,793	5,324	2,427	-
2003	14,861	-	1	-	4	5	87	450	1,755	4,063	5,920	2,574	2
2002	15,489	1	3	-	2	10	94	454	1,703	4,388	6,109	2,724	1
2001	16,242	3	-	1	2	7	75	448	1,829	4,804	6,337	2,736	-
2000	16,700	3	2	-	5	11	82	474	1,807	5,120	6,511	2,685	-
1999	17,787	1	-	1	1	9	97	468	2,019	5,656	6,955	2,580	-
All origins, male													
2005	7,264	-	1	-	-	5	54	277	973	2,039	2,767	1,147	1
2004	7,075	2	-	1	3	5	49	244	873	2,051	2,725	1,122	-
2003	7,549	-	1	-	2	3	49	279	932	2,141	2,995	1,145	2
2002	8,084	-	3	-	-	8	66	267	925	2,335	3,228	1,251	1
2001	8,415	1	-	1	1	4	52	266	966	2,554	3,255	1,315	-
2000	8,671	3	-	-	5	5	46	280	943	2,738	3,373	1,278	-
1999	9,480	-	-	-	1	6	49	268	1,085	3,136	3,677	1,258	-
All origins, female													
2005	6,738	2	-	-	-	3	23	167	704	1,710	2,633	1,496	-
2004	6,564	1	2	-	-	5	31	169	710	1,742	2,599	1,305	-
2003	7,312	-	-	-	2	2	38	171	823	1,922	2,925	1,429	-
2002	7,405	1	-	-	2	2	28	187	778	2,053	2,881	1,473	-
2001	7,827	2	-	-	1	3	23	182	863	2,250	3,082	1,421	-
2000	8,029	-	2	-	-	6	36	194	864	2,382	3,138	1,407	-
1999	8,307	1	-	1	-	3	48	200	934	2,520	3,278	1,322	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Emphysema (J43)													
Hispanic, both sexes													
2005	301	1	-	-	-	2	4	12	30	72	121	58	1
2004	269	1	1	-	-	1	4	6	20	63	110	63	-
2003	312	-	-	-	-	-	6	8	30	73	128	67	-
2002	295	-	-	-	-	2	1	5	29	79	113	66	-
2001	286	1	-	-	-	-	-	8	26	75	113	63	-
2000	266	-	-	-	1	-	3	11	26	65	113	47	-
1999	292	-	-	1	-	2	6	12	21	74	113	63	-
Hispanic, male													
2005	181	-	-	-	-	2	3	5	20	51	68	31	1
2004	152	-	-	-	-	1	4	4	15	37	57	34	-
2003	184	-	-	-	-	-	5	6	21	41	76	35	-
2002	168	-	-	-	-	1	1	4	17	48	65	32	-
2001	175	-	-	-	-	-	-	5	16	46	76	32	-
2000	167	-	-	-	1	-	3	8	17	43	66	29	-
1999	183	-	-	-	-	1	2	8	16	51	69	36	-
Hispanic, female													
2005	120	1	-	-	-	-	1	7	10	21	53	27	-
2004	117	1	1	-	-	-	-	2	5	26	53	29	-
2003	128	-	-	-	-	-	1	2	9	32	52	32	-
2002	127	-	-	-	-	1	-	1	12	31	48	34	-
2001	111	1	-	-	-	-	-	3	10	29	37	31	-
2000	99	-	-	-	-	-	-	3	9	22	47	18	-
1999	109	-	-	1	-	1	4	4	5	23	44	27	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Emphysema (J43)													
Non-Hispanic 1/, both sexes													
2005	13,678	1	1	-	-	6	72	430	1,645	3,671	5,271	2,581	-
2004	13,348	2	1	1	3	9	76	405	1,560	3,724	5,209	2,358	-
2003	14,514	-	1	-	4	5	81	438	1,716	3,982	5,781	2,505	1
2002	15,161	1	3	-	2	8	93	448	1,672	4,298	5,986	2,650	-
2001	15,914	2	-	1	2	7	74	439	1,794	4,718	6,210	2,667	-
2000	16,382	3	2	-	4	11	79	457	1,777	5,036	6,382	2,631	-
1999	17,441	1	-	-	1	7	88	450	1,992	5,571	6,825	2,506	-
Non-Hispanic 1/, male													
2005	7,067	-	1	-	-	3	50	271	951	1,983	2,695	1,113	-
2004	6,909	2	-	1	3	4	45	238	856	2,009	2,666	1,085	-
2003	7,342	-	1	-	2	3	44	270	908	2,094	2,910	1,109	1
2002	7,895	-	3	-	-	7	65	262	907	2,281	3,156	1,214	-
2001	8,213	1	-	1	1	4	51	260	943	2,500	3,172	1,280	-
2000	8,483	3	-	-	4	5	43	268	925	2,685	3,303	1,247	-
1999	9,260	-	-	-	1	5	45	255	1,063	3,076	3,598	1,217	-
Non-Hispanic 1/, female													
2005	6,611	1	-	-	-	3	22	159	694	1,688	2,576	1,468	-
2004	6,439	-	1	-	-	5	31	167	704	1,715	2,543	1,273	-
2003	7,172	-	-	-	2	2	37	168	808	1,888	2,871	1,396	-
2002	7,266	1	-	-	2	1	28	186	765	2,017	2,830	1,436	-
2001	7,701	1	-	-	1	3	23	179	851	2,218	3,038	1,387	-
2000	7,899	-	2	-	-	6	36	189	852	2,351	3,079	1,384	-
1999	8,181	1	-	-	-	2	43	195	929	2,495	3,227	1,289	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Emphysema (J43)													
Non-Hispanic white, both sexes													
2005 2/	12,675	-	1	-	-	4	53	353	1,492	3,404	4,930	2,438	-
2004 2/	12,471	-	-	1	2	6	58	334	1,408	3,491	4,938	2,233	-
2003 2/	13,518	-	1	-	4	3	64	376	1,557	3,717	5,438	2,357	1
2002	14,157	1	1	-	1	6	66	385	1,529	3,992	5,663	2,513	-
2001	14,930	1	-	-	1	4	64	374	1,633	4,414	5,913	2,526	-
2000	15,301	3	-	-	1	10	60	367	1,616	4,704	6,068	2,472	-
1999	16,338	-	-	-	-	5	64	372	1,825	5,238	6,469	2,365	-
Non-Hispanic white, male													
2005 2/	6,430	-	1	-	-	2	35	219	842	1,814	2,484	1,033	-
2004 2/	6,348	-	-	1	2	2	32	190	755	1,855	2,499	1,012	-
2003 2/	6,714	-	1	-	2	1	34	226	808	1,935	2,690	1,016	1
2002	7,241	-	1	-	-	5	47	219	810	2,081	2,946	1,132	-
2001	7,607	-	-	-	1	2	44	222	841	2,310	2,986	1,201	-
2000	7,805	3	-	-	1	4	31	206	830	2,470	3,121	1,139	-
1999	8,540	-	-	-	-	3	33	199	953	2,843	3,371	1,138	-
Non-Hispanic white, female													
2005 2/	6,245	-	-	-	-	2	18	134	650	1,590	2,446	1,405	-
2004 2/	6,123	-	-	-	-	4	26	144	653	1,636	2,439	1,221	-
2003 2/	6,804	-	-	-	2	2	30	150	749	1,782	2,748	1,341	-
2002	6,916	1	-	-	1	1	19	166	719	1,911	2,717	1,381	-
2001	7,323	1	-	-	-	2	20	152	792	2,104	2,927	1,325	-
2000	7,496	-	-	-	-	6	29	161	786	2,234	2,947	1,333	-
1999	7,798	-	-	-	-	2	31	173	872	2,395	3,098	1,227	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Emphysema (J43)													
Non-Hispanic black, both sexes													
2005 2/	810	1	-	-	-	2	17	69	137	221	263	100	-
2004 2/	716	2	-	-	1	2	15	64	131	185	219	97	-
2003 2/	805	-	-	-	-	2	15	55	142	220	266	105	-
2002	842	-	1	-	1	2	26	56	117	267	266	106	-
2001	800	1	-	1	1	3	9	59	138	254	219	115	-
2000	889	-	1	-	3	1	17	84	146	276	243	118	-
1999	923	1	-	-	-	-	22	75	146	279	283	117	-
Non-Hispanic black, male													
2005 2/	505	-	-	-	-	1	13	47	99	139	153	53	-
2004 2/	458	2	-	-	1	1	11	45	89	122	130	57	-
2003 2/	495	-	-	-	-	2	8	37	90	129	162	67	-
2002	538	-	1	-	-	2	18	38	77	173	169	60	-
2001	480	1	-	1	-	2	7	33	87	155	134	60	-
2000	551	-	-	-	3	1	11	56	87	183	134	76	-
1999	594	-	-	-	-	-	11	54	94	195	178	62	-
Non-Hispanic black, female													
2005 2/	305	1	-	-	-	1	4	22	38	82	110	47	-
2004 2/	258	-	-	-	-	1	4	19	42	63	89	40	-
2003 2/	310	-	-	-	-	-	7	18	52	91	104	38	-
2002	304	-	-	-	1	-	8	18	40	94	97	46	-
2001	320	-	-	-	1	1	2	26	51	99	85	55	-
2000	338	-	1	-	-	-	6	28	59	93	109	42	-
1999	329	1	-	-	-	-	11	21	52	84	105	55	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Emphysema (J43)													
Origin not stated 3/, both sexes													
2005	23	-	-	-	-	-	1	2	2	6	8	4	-
2004	22	-	-	-	-	-	-	2	3	6	5	6	-
2003	35	-	-	-	-	-	-	4	9	8	11	2	1
2002	33	-	-	-	-	-	-	1	2	11	10	8	1
2001	42	-	-	-	-	-	1	1	9	11	14	6	-
2000	52	-	-	-	-	-	-	6	4	19	16	7	-
1999	54	-	-	-	-	-	3	6	6	11	17	11	-
Origin not stated 3/, male													
2005	16	-	-	-	-	-	1	1	2	5	4	3	-
2004	14	-	-	-	-	-	-	2	2	5	2	3	-
2003	23	-	-	-	-	-	-	3	3	6	9	1	1
2002	21	-	-	-	-	-	-	1	1	6	7	5	1
2001	27	-	-	-	-	-	1	1	7	8	7	3	-
2000	21	-	-	-	-	-	-	4	1	10	4	2	-
1999	37	-	-	-	-	-	2	5	6	9	10	5	-
Origin not stated 3/, female													
2005	7	-	-	-	-	-	-	1	-	1	4	1	-
2004	8	-	-	-	-	-	-	-	1	1	3	3	-
2003	12	-	-	-	-	-	-	1	6	2	2	1	-
2002	12	-	-	-	-	-	-	-	1	5	3	3	-
2001	15	-	-	-	-	-	-	-	2	3	7	3	-
2000	31	-	-	-	-	-	-	2	3	9	12	5	-
1999	17	-	-	-	-	-	1	1	-	2	7	6	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Asthma (J45-J46)													
All origins, both sexes													
2005	3,884	4	37	97	131	207	369	595	520	475	709	740	-
2004	3,816	7	29	105	159	197	376	562	520	504	673	684	-
2003	4,099	7	37	110	158	227	411	632	562	532	752	671	-
2002	4,261	4	43	123	169	235	472	608	536	583	812	675	1
2001	4,269	10	31	99	140	233	455	603	553	634	802	709	-
2000	4,487	8	32	129	167	249	458	614	540	734	849	707	-
1999	4,657	5	27	126	182	262	447	607	583	773	925	720	-
All origins, male													
2005	1,315	1	25	61	83	101	151	215	170	150	218	140	-
2004	1,386	7	19	64	99	113	158	213	187	178	181	167	-
2003	1,493	4	29	71	91	127	160	236	214	173	234	154	-
2002	1,580	3	30	69	110	129	198	233	202	198	256	152	-
2001	1,479	6	22	57	80	116	195	219	182	220	225	157	-
2000	1,632	6	20	74	102	142	196	238	200	253	243	158	-
1999	1,620	3	16	85	108	128	184	197	207	267	273	152	-
All origins, female													
2005	2,569	3	12	36	48	106	218	380	350	325	491	600	-
2004	2,430	-	10	41	60	84	218	349	333	326	492	517	-
2003	2,606	3	8	39	67	100	251	396	348	359	518	517	-
2002	2,681	1	13	54	59	106	274	375	334	385	556	523	1
2001	2,790	4	9	42	60	117	260	384	371	414	577	552	-
2000	2,855	2	12	55	65	107	262	376	340	481	606	549	-
1999	3,037	2	11	41	74	134	263	410	376	506	652	568	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Asthma (J45-J46)													
Hispanic, both sexes													
2005	248	2	3	14	7	17	30	33	37	38	35	32	-
2004	281	3	7	11	14	23	33	46	42	42	36	24	-
2003	299	-	6	15	14	23	27	61	45	41	39	28	-
2002	287	1	6	13	11	23	42	44	38	45	42	22	-
2001	274	2	5	8	8	17	31	49	45	43	44	22	-
2000	292	1	5	13	14	16	29	41	48	60	40	25	-
1999	320	-	3	14	14	21	46	48	39	60	50	25	-
Hispanic, male													
2005	106	1	2	9	6	9	18	14	11	13	11	12	-
2004	111	3	4	6	9	15	19	16	9	15	8	7	-
2003	131	-	5	10	8	12	13	27	22	13	15	6	-
2002	114	1	3	8	9	14	14	22	15	10	15	3	-
2001	103	2	3	5	6	10	13	14	17	18	13	2	-
2000	129	1	2	8	11	9	14	20	19	25	12	8	-
1999	119	-	-	9	8	11	22	20	12	19	12	6	-
Hispanic, female													
2005	142	1	1	5	1	8	12	19	26	25	24	20	-
2004	170	-	3	5	5	8	14	30	33	27	28	17	-
2003	168	-	1	5	6	11	14	34	23	28	24	22	-
2002	173	-	3	5	2	9	28	22	23	35	27	19	-
2001	171	-	2	3	2	7	18	35	28	25	31	20	-
2000	163	-	3	5	3	7	15	21	29	35	28	17	-
1999	201	-	3	5	6	10	24	28	27	41	38	19	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Asthma (J45-J46)													
Non-Hispanic 1/, both sexes													
2005	3,626	2	34	82	124	190	338	561	481	435	672	707	-
2004	3,524	4	22	94	145	173	341	516	477	460	634	658	-
2003	3,786	7	30	94	143	204	382	568	516	491	713	638	-
2002	3,950	3	37	108	157	210	425	561	495	536	767	650	1
2001	3,976	8	26	90	132	215	421	550	507	587	756	684	-
2000	4,164	7	27	115	152	229	428	568	486	668	805	679	-
1999	4,324	5	24	112	168	238	400	557	543	711	874	692	-
Non-Hispanic 1/, male													
2005	1,205	-	23	51	77	92	133	200	159	136	206	128	-
2004	1,272	4	15	58	90	98	139	197	177	163	172	159	-
2003	1,356	4	23	61	82	115	145	208	192	160	219	147	-
2002	1,448	2	27	59	100	113	179	209	187	186	239	147	-
2001	1,365	4	19	52	74	106	180	202	165	200	211	152	-
2000	1,486	5	18	66	91	130	181	215	176	226	229	149	-
1999	1,495	3	16	76	100	116	162	176	195	246	260	145	-
Non-Hispanic 1/, female													
2005	2,421	2	11	31	47	98	205	361	322	299	466	579	-
2004	2,252	-	7	36	55	75	202	319	300	297	462	499	-
2003	2,430	3	7	33	61	89	237	360	324	331	494	491	-
2002	2,502	1	10	49	57	97	246	352	308	350	528	503	1
2001	2,611	4	7	38	58	109	241	348	342	387	545	532	-
2000	2,678	2	9	49	61	99	247	353	310	442	576	530	-
1999	2,829	2	8	36	68	122	238	381	348	465	614	547	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Asthma (J45-J46)													
Non-Hispanic white, both sexes													
2005 2/	2,475	-	7	26	52	101	196	329	307	286	539	632	-
2004 2/	2,377	2	7	34	63	80	195	274	296	342	507	577	-
2003 2/	2,589	4	8	34	64	108	215	318	334	348	581	575	-
2002	2,720	1	14	34	79	121	222	316	311	377	659	586	-
2001	2,717	4	7	37	59	97	219	326	306	432	628	602	-
2000	2,836	4	13	40	75	114	227	317	296	484	683	583	-
1999	3,011	1	8	47	92	107	222	327	351	511	747	598	-
Non-Hispanic white, male													
2005 2/	743	-	3	15	32	45	73	120	100	87	156	112	-
2004 2/	743	2	5	18	39	37	74	99	97	114	124	134	-
2003 2/	831	4	4	19	36	48	78	113	114	114	169	132	-
2002	895	-	12	20	43	59	80	117	113	125	196	130	-
2001	834	3	4	21	29	42	91	113	95	141	167	128	-
2000	920	2	7	20	44	60	90	130	97	159	193	118	-
1999	930	1	5	31	52	48	80	97	124	164	218	110	-
Non-Hispanic white, female													
2005 2/	1,732	-	4	11	20	56	123	209	207	199	383	520	-
2004 2/	1,634	-	2	16	24	43	121	175	199	228	383	443	-
2003 2/	1,758	-	4	15	28	60	137	205	220	234	412	443	-
2002	1,825	1	2	14	36	62	142	199	198	252	463	456	-
2001	1,883	1	3	16	30	55	128	213	211	291	461	474	-
2000	1,916	2	6	20	31	54	137	187	199	325	490	465	-
1999	2,081	-	3	16	40	59	142	230	227	347	529	488	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Asthma (J45-J46)													
Non-Hispanic black, both sexes													
2005 2/	1,002	2	27	50	70	84	131	210	162	124	98	44	-
2004 2/	1,001	2	13	57	74	91	139	221	162	98	91	53	-
2003 2/	1,019	3	20	59	74	90	155	229	140	120	89	40	-
2002	1,083	2	21	69	74	84	196	222	169	132	75	39	-
2001	1,092	3	19	51	67	115	186	203	170	126	99	53	-
2000	1,145	3	14	69	73	102	178	225	165	152	94	70	-
1999	1,134	4	14	65	71	126	166	211	162	157	94	64	-
Non-Hispanic black, male													
2005 2/	403	-	20	33	44	44	56	68	54	38	38	8	-
2004 2/	452	2	9	39	45	60	60	89	66	40	31	11	-
2003 2/	439	-	18	41	42	63	61	86	52	38	31	7	-
2002	488	2	14	37	56	50	96	83	67	51	23	9	-
2001	450	-	15	30	41	62	79	83	52	46	29	13	-
2000	472	3	11	42	44	62	79	71	65	53	21	21	-
1999	474	2	10	45	46	66	72	72	57	61	27	16	-
Non-Hispanic black, female													
2005 2/	599	2	7	17	26	40	75	142	108	86	60	36	-
2004 2/	549	-	4	18	29	31	79	132	96	58	60	42	-
2003 2/	580	3	2	18	32	27	94	143	88	82	58	33	-
2002	595	-	7	32	18	34	100	139	102	81	52	30	-
2001	642	3	4	21	26	53	107	120	118	80	70	40	-
2000	673	-	3	27	29	40	99	154	100	99	73	49	-
1999	660	2	4	20	25	60	94	139	105	96	67	48	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Asthma (J45-J46)													
Origin not stated 3/, both sexes													
2005	10	-	-	1	-	-	1	1	2	2	2	1	-
2004	11	-	-	-	-	1	2	-	1	2	3	2	-
2003	14	-	1	1	1	-	2	3	1	-	-	5	-
2002	24	-	-	2	1	2	5	3	3	2	3	3	-
2001	19	-	-	1	-	1	3	4	1	4	2	3	-
2000	31	-	-	1	1	4	1	5	6	6	4	3	-
1999	13	-	-	-	-	3	1	2	1	2	1	3	-
Origin not stated 3/, male													
2005	4	-	-	1	-	-	-	1	-	1	1	-	-
2004	3	-	-	-	-	-	-	-	1	-	1	1	-
2003	6	-	1	-	1	-	2	1	-	-	-	1	-
2002	18	-	-	2	1	2	5	2	-	2	2	2	-
2001	11	-	-	-	-	-	2	3	-	2	1	3	-
2000	17	-	-	-	-	3	1	3	5	2	2	1	-
1999	6	-	-	-	-	1	-	1	-	2	1	1	-
Origin not stated 3/, female													
2005	6	-	-	-	-	-	1	-	2	1	1	1	-
2004	8	-	-	-	-	1	2	-	-	2	2	1	-
2003	8	-	-	1	-	-	-	2	1	-	-	4	-
2002	6	-	-	-	-	-	-	1	3	-	1	1	-
2001	8	-	-	1	-	1	1	1	1	2	1	-	-
2000	14	-	-	1	1	1	-	2	1	4	2	2	-
1999	7	-	-	-	-	2	1	1	1	-	-	2	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Other chronic lower respiratory diseases (J44,J47)													
All origins, both sexes													
2005	112,181	2	6	4	12	37	429	2,911	10,477	25,567	44,019	28,714	3
2004	103,759	8	5	10	10	43	419	2,501	9,584	23,996	41,372	25,810	1
2003	106,572	4	4	2	22	40	435	2,420	9,713	25,220	42,368	26,344	-
2002	104,111	10	6	8	15	50	424	2,377	8,987	24,682	42,034	25,513	5
2001	101,543	7	4	4	24	44	418	2,231	8,723	25,180	40,362	24,545	1
2000	99,655	7	3	5	13	28	370	2,125	8,325	25,145	40,029	23,603	2
1999	100,565	10	8	8	19	48	328	1,997	8,598	26,042	40,416	23,087	4
All origins, male													
2005	53,495	2	2	-	9	21	211	1,524	5,411	13,173	21,394	11,746	2
2004	49,847	3	2	6	4	29	219	1,327	4,934	12,220	20,266	10,836	1
2003	51,321	4	2	1	15	23	211	1,288	5,071	12,973	20,621	11,112	-
2002	50,629	5	3	3	7	26	221	1,268	4,628	12,732	20,821	10,911	4
2001	49,407	4	2	3	15	24	200	1,196	4,518	13,027	20,096	10,321	1
2000	49,198	5	1	3	9	11	181	1,127	4,389	13,140	20,048	10,282	2
1999	50,817	7	3	5	10	15	172	1,067	4,615	13,980	20,598	10,342	3
All origins, female													
2005	58,686	-	4	4	3	16	218	1,387	5,066	12,394	22,625	16,968	1
2004	53,912	5	3	4	6	14	200	1,174	4,650	11,776	21,106	14,974	-
2003	55,251	-	2	1	7	17	224	1,132	4,642	12,247	21,747	15,232	-
2002	53,482	5	3	5	8	24	203	1,109	4,359	11,950	21,213	14,602	1
2001	52,136	3	2	1	9	20	218	1,035	4,205	12,153	20,266	14,224	-
2000	50,457	2	2	2	4	17	189	998	3,936	12,005	19,981	13,321	-
1999	49,748	3	5	3	9	33	156	930	3,983	12,062	19,818	12,745	1

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Other chronic lower respiratory diseases (J44,J47)													
Hispanic, both sexes													
2005	2,864	1	1	1	1	4	19	105	230	555	1,086	860	1
2004	2,519	1	1	-	3	6	29	70	207	549	927	726	-
2003	2,523	2	-	-	5	5	25	64	190	534	919	779	-
2002	2,425	-	2	-	2	5	24	75	205	509	886	717	-
2001	2,235	-	2	1	1	4	19	66	185	513	774	670	-
2000	2,079	2	1	2	1	2	20	52	169	438	755	637	-
1999	2,195	3	2	2	1	6	20	55	177	511	759	659	-
Hispanic, male													
2005	1,533	1	-	-	1	4	13	57	148	316	616	376	1
2004	1,349	-	1	-	2	4	22	35	119	335	506	325	-
2003	1,362	2	-	-	3	3	12	28	121	313	507	373	-
2002	1,323	-	-	-	2	3	14	41	131	303	509	320	-
2001	1,189	-	1	1	-	2	7	35	106	313	406	318	-
2000	1,127	1	1	1	1	1	5	25	109	268	425	290	-
1999	1,228	3	1	1	1	1	10	24	107	300	439	341	-
Hispanic, female													
2005	1,331	-	1	1	-	-	6	48	82	239	470	484	-
2004	1,170	1	-	-	1	2	7	35	88	214	421	401	-
2003	1,161	-	-	-	2	2	13	36	69	221	412	406	-
2002	1,102	-	2	-	-	2	10	34	74	206	377	397	-
2001	1,046	-	1	-	1	2	12	31	79	200	368	352	-
2000	952	1	-	1	-	1	15	27	60	170	330	347	-
1999	967	-	1	1	-	5	10	31	70	211	320	318	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Other chronic lower respiratory diseases (J44,J47)													
Non-Hispanic 1/, both sexes													
2005	109,129	1	5	3	11	33	408	2,797	10,218	24,967	42,867	27,818	1
2004	101,045	7	4	10	7	37	389	2,421	9,354	23,378	40,384	25,053	1
2003	103,788	2	4	2	17	35	410	2,342	9,483	24,618	41,364	25,511	-
2002	101,413	10	4	8	13	44	397	2,293	8,756	24,103	41,034	24,746	5
2001	99,030	7	2	3	23	40	395	2,151	8,501	24,593	39,496	23,819	-
2000	97,305	5	2	3	12	26	348	2,060	8,118	24,633	39,180	22,917	1
1999	98,086	7	6	6	18	42	308	1,933	8,389	25,452	39,553	22,369	3
Non-Hispanic 1/, male													
2005	51,848	1	2	-	8	17	196	1,463	5,242	12,828	20,741	11,350	-
2004	48,385	3	1	6	2	25	197	1,286	4,801	11,840	19,723	10,500	1
2003	49,820	2	2	1	12	20	199	1,252	4,924	12,622	20,068	10,718	-
2002	49,155	5	3	3	5	22	206	1,224	4,483	12,384	20,244	10,572	4
2001	48,070	4	1	2	15	22	190	1,152	4,388	12,674	19,642	9,980	-
2000	47,906	4	-	2	8	10	175	1,094	4,256	12,822	19,566	9,968	1
1999	49,436	4	2	4	9	14	162	1,036	4,489	13,628	20,112	9,974	2
Non-Hispanic 1/, female													
2005	57,281	-	3	3	3	16	212	1,334	4,976	12,139	22,126	16,468	1
2004	52,660	4	3	4	5	12	192	1,135	4,553	11,538	20,661	14,553	-
2003	53,968	-	2	1	5	15	211	1,090	4,559	11,996	21,296	14,793	-
2002	52,258	5	1	5	8	22	191	1,069	4,273	11,719	20,790	14,174	1
2001	50,960	3	1	1	8	18	205	999	4,113	11,919	19,854	13,839	-
2000	49,399	1	2	1	4	16	173	966	3,862	11,811	19,614	12,949	-
1999	48,650	3	4	2	9	28	146	897	3,900	11,824	19,441	12,395	1

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Other chronic lower respiratory diseases (J44,J47)													
Non-Hispanic white, both sexes													
2005 2/	101,449	1	3	2	8	20	333	2,350	9,202	23,031	40,296	26,202	1
2004 2/	94,169	4	1	4	5	29	322	2,063	8,433	21,644	37,994	23,669	1
2003 2/	96,722	2	2	2	9	24	325	2,008	8,537	22,745	38,944	24,124	-
2002	94,430	6	2	5	12	26	323	1,959	7,873	22,315	38,543	23,362	4
2001	92,267	4	1	2	12	29	314	1,835	7,685	22,734	37,199	22,452	-
2000	90,746	4	2	3	7	16	272	1,748	7,281	22,803	36,976	21,633	1
1999	91,269	1	3	5	12	28	238	1,649	7,524	23,501	37,153	21,153	2
Non-Hispanic white, male													
2005 2/	47,546	1	2	-	6	8	159	1,208	4,657	11,706	19,250	10,549	-
2004 2/	44,542	1	-	2	1	20	162	1,096	4,246	10,845	18,380	9,788	1
2003 2/	45,758	2	-	1	6	13	156	1,070	4,382	11,516	18,598	10,014	-
2002	45,172	4	2	1	4	16	169	1,030	3,964	11,342	18,779	9,858	3
2001	44,173	2	-	1	7	18	152	979	3,917	11,572	18,246	9,279	-
2000	44,077	4	-	2	4	4	140	908	3,774	11,705	18,230	9,305	1
1999	45,423	-	-	3	4	13	123	867	3,969	12,471	18,653	9,319	1
Non-Hispanic white, female													
2005 2/	53,903	-	1	2	2	12	174	1,142	4,545	11,325	21,046	15,653	1
2004 2/	49,627	3	1	2	4	9	160	967	4,187	10,799	19,614	13,881	-
2003 2/	50,964	-	2	1	3	11	169	938	4,155	11,229	20,346	14,110	-
2002	49,258	2	-	4	8	10	154	929	3,909	10,973	19,764	13,504	1
2001	48,094	2	1	1	5	11	162	856	3,768	11,162	18,953	13,173	-
2000	46,669	-	2	1	3	12	132	840	3,507	11,098	18,746	12,328	-
1999	45,846	1	3	2	8	15	115	782	3,555	11,030	18,500	11,834	1

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Other chronic lower respiratory diseases (J44,J47)													
Non-Hispanic black, both sexes													
2005 2/	6,270	-	2	1	3	10	66	414	885	1,644	2,043	1,202	-
2004 2/	5,550	2	3	6	2	8	62	327	799	1,463	1,877	1,001	-
2003 2/	5,734	-	1	-	6	9	71	302	819	1,574	1,923	1,029	-
2002	5,739	4	1	3	1	15	68	304	776	1,487	2,012	1,067	1
2001	5,550	2	1	1	8	9	75	283	707	1,567	1,861	1,036	-
2000	5,414	1	-	-	3	10	71	278	748	1,549	1,771	983	-
1999	5,695	6	3	1	6	11	68	258	770	1,639	1,961	971	1
Non-Hispanic black, male													
2005 2/	3,465	-	-	-	2	7	32	234	503	963	1,162	562	-
2004 2/	3,066	1	1	4	1	5	33	175	486	835	1,037	488	-
2003 2/	3,239	-	1	-	5	6	35	164	470	924	1,140	494	-
2002	3,221	1	-	2	1	4	34	176	461	865	1,153	523	1
2001	3,181	1	1	1	6	4	37	151	413	938	1,116	513	-
2000	3,126	-	-	-	3	6	33	161	432	953	1,045	493	-
1999	3,341	4	2	1	5	-	38	151	469	975	1,181	514	1
Non-Hispanic black, female													
2005 2/	2,805	-	2	1	1	3	34	180	382	681	881	640	-
2004 2/	2,484	1	2	2	1	3	29	152	313	628	840	513	-
2003 2/	2,495	-	-	-	1	3	36	138	349	650	783	535	-
2002	2,518	3	1	1	-	11	34	128	315	622	859	544	-
2001	2,369	1	-	-	2	5	38	132	294	629	745	523	-
2000	2,288	1	-	-	-	4	38	117	316	596	726	490	-
1999	2,354	2	1	-	1	11	30	107	301	664	780	457	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Other chronic lower respiratory diseases (J44,J47)													
Origin not stated 3/, both sexes													
2005	188	-	-	-	-	-	2	9	29	45	66	36	1
2004	195	-	-	-	-	-	1	10	23	69	61	31	-
2003	261	-	-	-	-	-	-	14	40	68	85	54	-
2002	273	-	-	-	-	1	3	9	26	70	114	50	-
2001	278	-	-	-	-	-	4	14	37	74	92	56	1
2000	271	-	-	-	-	-	2	13	38	74	94	49	1
1999	284	-	-	-	-	-	-	9	32	79	104	59	1
Origin not stated 3/, male													
2005	114	-	-	-	-	-	2	4	21	29	37	20	1
2004	113	-	-	-	-	-	-	6	14	45	37	11	-
2003	139	-	-	-	-	-	-	8	26	38	46	21	-
2002	151	-	-	-	-	1	1	3	14	45	68	19	-
2001	148	-	-	-	-	-	3	9	24	40	48	23	1
2000	165	-	-	-	-	-	1	8	24	50	57	24	1
1999	153	-	-	-	-	-	-	7	19	52	47	27	1
Origin not stated 3/, female													
2005	74	-	-	-	-	-	-	5	8	16	29	16	-
2004	82	-	-	-	-	-	1	4	9	24	24	20	-
2003	122	-	-	-	-	-	-	6	14	30	39	33	-
2002	122	-	-	-	-	-	2	6	12	25	46	31	-
2001	130	-	-	-	-	-	1	5	13	34	44	33	-
2000	106	-	-	-	-	-	1	5	14	24	37	25	-
1999	131	-	-	-	-	-	-	2	13	27	57	32	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Pneumoconioses and chemical effects (J60-J66,J68)													
All origins, both sexes													
2005	1,007	-	1	1	-	-	3	22	69	175	447	288	1
2004	1,033	-	-	-	1	-	4	17	70	209	442	290	-
2003	1,114	-	-	-	1	1	7	14	70	233	487	301	-
2002	1,114	2	-	-	-	-	5	17	75	223	480	312	-
2001	1,136	-	1	-	-	1	5	11	64	266	516	272	-
2000	1,161	-	-	-	-	3	7	12	57	281	484	317	-
1999	1,113	-	3	-	1	-	7	15	57	274	503	253	-
All origins, male													
2005	965	-	-	1	-	-	3	19	65	173	428	275	1
2004	993	-	-	-	1	-	4	14	68	204	425	277	-
2003	1,065	-	-	-	1	1	6	13	67	222	469	286	-
2002	1,057	-	-	-	-	-	4	13	69	214	457	300	-
2001	1,095	-	1	-	-	1	5	9	62	261	494	262	-
2000	1,113	-	-	-	-	3	6	12	54	265	467	306	-
1999	1,079	-	3	-	1	-	5	15	57	269	486	243	-
All origins, female													
2005	42	-	1	-	-	-	-	3	4	2	19	13	-
2004	40	-	-	-	-	-	-	3	2	5	17	13	-
2003	49	-	-	-	-	-	1	1	3	11	18	15	-
2002	57	2	-	-	-	-	1	4	6	9	23	12	-
2001	41	-	-	-	-	-	-	2	2	5	22	10	-
2000	48	-	-	-	-	-	1	-	3	16	17	11	-
1999	34	-	-	-	-	-	2	-	-	5	17	10	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Pneumoconioses and chemical effects (J60-J66,J68)													
Hispanic, both sexes													
2005	21	-	1	-	-	-	-	1	1	8	6	4	-
2004	25	-	-	-	-	-	2	1	2	4	10	6	-
2003	20	-	-	-	1	-	-	-	3	3	12	1	-
2002	33	-	-	-	-	-	1	2	4	12	5	9	-
2001	26	-	1	-	-	-	2	2	2	7	6	6	-
2000	27	-	-	-	-	1	1	1	1	5	13	5	-
1999	13	-	-	-	-	-	1	1	1	5	2	3	-
Hispanic, male													
2005	19	-	-	-	-	-	-	1	1	7	6	4	-
2004	25	-	-	-	-	-	2	1	2	4	10	6	-
2003	19	-	-	-	1	-	-	-	3	3	12	-	-
2002	31	-	-	-	-	-	1	1	4	11	5	9	-
2001	25	-	1	-	-	-	2	2	2	7	6	5	-
2000	26	-	-	-	-	1	1	1	1	5	13	4	-
1999	11	-	-	-	-	-	1	1	1	5	2	1	-
Hispanic, female													
2005	2	-	1	-	-	-	-	-	-	1	-	-	-
2004	-	-	-	-	-	-	-	-	-	-	-	-	-
2003	1	-	-	-	-	-	-	-	-	-	-	1	-
2002	2	-	-	-	-	-	-	1	-	1	-	-	-
2001	1	-	-	-	-	-	-	-	-	-	-	1	-
2000	1	-	-	-	-	-	-	-	-	-	-	1	-
1999	2	-	-	-	-	-	-	-	-	-	-	2	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Pneumoconioses and chemical effects (J60-J66,J68)													
Non-Hispanic 1/, both sexes													
2005	985	-	-	1	-	-	3	21	68	166	441	284	1
2004	1,007	-	-	-	1	-	2	16	68	205	431	284	-
2003	1,093	-	-	-	-	1	7	14	67	230	474	300	-
2002	1,078	2	-	-	-	-	4	15	70	211	473	303	-
2001	1,108	-	-	-	-	1	3	9	62	259	509	265	-
2000	1,133	-	-	-	-	2	6	11	56	276	470	312	-
1999	1,100	-	3	-	1	-	6	14	56	269	501	250	-
Non-Hispanic 1/, male													
2005	945	-	-	1	-	-	3	18	64	165	422	271	1
2004	967	-	-	-	1	-	2	13	66	200	414	271	-
2003	1,045	-	-	-	-	1	6	13	64	219	456	286	-
2002	1,023	-	-	-	-	-	3	12	64	203	450	291	-
2001	1,068	-	-	-	-	1	3	7	60	254	487	256	-
2000	1,086	-	-	-	-	2	5	11	53	260	453	302	-
1999	1,068	-	3	-	1	-	4	14	56	264	484	242	-
Non-Hispanic 1/, female													
2005	40	-	-	-	-	-	-	3	4	1	19	13	-
2004	40	-	-	-	-	-	-	3	2	5	17	13	-
2003	48	-	-	-	-	-	1	1	3	11	18	14	-
2002	55	2	-	-	-	-	1	3	6	8	23	12	-
2001	40	-	-	-	-	-	-	2	2	5	22	9	-
2000	47	-	-	-	-	-	1	-	3	16	17	10	-
1999	32	-	-	-	-	-	2	-	-	5	17	8	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Pneumoconioses and chemical effects (J60-J66,J68)													
Non-Hispanic white, both sexes													
2005 2/	931	-	-	1	-	-	2	19	61	160	418	269	1
2004 2/	951	-	-	-	1	-	2	15	63	190	415	265	-
2003 2/	1,037	-	-	-	-	-	5	12	64	218	447	291	-
2002	1,014	1	-	-	-	-	3	13	62	201	445	289	-
2001	1,046	-	-	-	-	1	3	5	53	245	487	252	-
2000	1,072	-	-	-	-	2	3	11	54	261	442	299	-
1999	1,024	-	-	-	-	-	3	10	46	250	482	233	-
Non-Hispanic white, male													
2005 2/	895	-	-	1	-	-	2	16	59	159	401	256	1
2004 2/	912	-	-	-	1	-	2	12	61	185	398	253	-
2003 2/	992	-	-	-	-	-	5	11	61	207	430	278	-
2002	966	-	-	-	-	-	2	10	59	193	424	278	-
2001	1,008	-	-	-	-	1	3	5	51	240	465	243	-
2000	1,027	-	-	-	-	2	2	11	51	245	427	289	-
1999	995	-	-	-	-	-	2	10	46	246	465	226	-
Non-Hispanic white, female													
2005 2/	36	-	-	-	-	-	-	3	2	1	17	13	-
2004 2/	39	-	-	-	-	-	-	3	2	5	17	12	-
2003 2/	45	-	-	-	-	-	-	1	3	11	17	13	-
2002	48	1	-	-	-	-	1	3	3	8	21	11	-
2001	38	-	-	-	-	-	-	-	2	5	22	9	-
2000	45	-	-	-	-	-	1	-	3	16	15	10	-
1999	29	-	-	-	-	-	1	-	-	4	17	7	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Pneumoconioses and chemical effects (J60-J66,J68)													
Non-Hispanic black, both sexes													
2005 2/	47	-	-	-	-	-	1	2	6	5	19	14	-
2004 2/	54	-	-	-	-	-	-	1	5	14	16	18	-
2003 2/	48	-	-	-	-	1	1	2	1	11	23	9	-
2002	56	1	-	-	-	-	1	2	6	10	22	14	-
2001	56	-	-	-	-	-	-	4	8	12	20	12	-
2000	54	-	-	-	-	-	3	-	2	15	23	11	-
1999	67	-	3	-	1	-	3	4	8	16	16	16	-
Non-Hispanic black, male													
2005 2/	43	-	-	-	-	-	1	2	4	5	17	14	-
2004 2/	54	-	-	-	-	-	-	1	5	14	16	18	-
2003 2/	45	-	-	-	-	1	-	2	1	11	22	8	-
2002	50	-	-	-	-	-	1	2	4	10	20	13	-
2001	54	-	-	-	-	-	-	2	8	12	20	12	-
2000	52	-	-	-	-	-	3	-	2	15	21	11	-
1999	64	-	3	-	1	-	2	4	8	15	16	15	-
Non-Hispanic black, female													
2005 2/	4	-	-	-	-	-	-	-	2	-	2	-	-
2004 2/	-	-	-	-	-	-	-	-	-	-	-	-	-
2003 2/	3	-	-	-	-	-	1	-	-	-	1	1	-
2002	6	1	-	-	-	-	-	-	2	-	2	1	-
2001	2	-	-	-	-	-	-	2	-	-	-	-	-
2000	2	-	-	-	-	-	-	-	-	-	2	-	-
1999	3	-	-	-	-	-	1	-	-	1	-	1	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Pneumoconioses and chemical effects (J60-J66,J68)													
Origin not stated 3/, both sexes													
2005	1	-	-	-	-	-	-	-	-	1	-	-	-
2004	1	-	-	-	-	-	-	-	-	-	1	-	-
2003	1	-	-	-	-	-	-	-	-	-	1	-	-
2002	3	-	-	-	-	-	-	-	1	-	2	-	-
2001	2	-	-	-	-	-	-	-	-	-	1	1	-
2000	1	-	-	-	-	-	-	-	-	-	1	-	-
1999	-	-	-	-	-	-	-	-	-	-	-	-	-
Origin not stated 3/, male													
2005	1	-	-	-	-	-	-	-	-	1	-	-	-
2004	1	-	-	-	-	-	-	-	-	-	1	-	-
2003	1	-	-	-	-	-	-	-	-	-	1	-	-
2002	3	-	-	-	-	-	-	-	1	-	2	-	-
2001	2	-	-	-	-	-	-	-	-	-	1	1	-
2000	1	-	-	-	-	-	-	-	-	-	1	-	-
1999	-	-	-	-	-	-	-	-	-	-	-	-	-
Origin not stated 3/, female													
2005	-	-	-	-	-	-	-	-	-	-	-	-	-
2004	-	-	-	-	-	-	-	-	-	-	-	-	-
2003	-	-	-	-	-	-	-	-	-	-	-	-	-
2002	-	-	-	-	-	-	-	-	-	-	-	-	-
2001	-	-	-	-	-	-	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-	-	-	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Pneumonitis due to solids and liquids (J69)													
All origins, both sexes													
2005	17,279	17	8	13	43	66	172	476	833	1,741	5,558	8,352	-
2004	16,780	12	9	11	39	68	159	394	743	1,767	5,564	8,014	-
2003	17,335	13	10	16	46	69	166	392	773	1,754	5,789	8,307	-
2002	17,593	2	7	16	44	74	192	351	671	1,793	5,900	8,543	-
2001	17,301	12	14	15	34	79	179	378	659	1,830	5,759	8,342	-
2000	16,636	10	11	13	37	62	181	366	598	1,899	5,505	7,954	-
1999	15,268	16	12	16	32	78	166	284	586	1,754	5,030	7,294	-
All origins, male													
2005	8,760	10	3	8	23	43	88	280	506	1,065	3,057	3,677	-
2004	8,593	4	5	6	20	40	97	250	451	1,085	3,059	3,576	-
2003	8,702	8	5	9	31	42	89	239	471	1,069	3,181	3,558	-
2002	8,876	-	3	9	27	42	119	226	393	1,083	3,243	3,731	-
2001	8,576	6	6	7	22	40	117	232	384	1,106	3,145	3,511	-
2000	8,390	4	5	9	23	35	115	225	354	1,195	3,038	3,387	-
1999	7,755	9	3	6	22	47	97	165	365	1,083	2,875	3,083	-
All origins, female													
2005	8,519	7	5	5	20	23	84	196	327	676	2,501	4,675	-
2004	8,187	8	4	5	19	28	62	144	292	682	2,505	4,438	-
2003	8,633	5	5	7	15	27	77	153	302	685	2,608	4,749	-
2002	8,717	2	4	7	17	32	73	125	278	710	2,657	4,812	-
2001	8,725	6	8	8	12	39	62	146	275	724	2,614	4,831	-
2000	8,246	6	6	4	14	27	66	141	244	704	2,467	4,567	-
1999	7,513	7	9	10	10	31	69	119	221	671	2,155	4,211	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Pneumonitis due to solids and liquids (J69)													
Hispanic, both sexes													
2005	591	3	2	3	1	8	8	28	34	70	177	257	-
2004	536	2	3	1	8	5	6	18	35	65	180	213	-
2003	524	2	3	2	6	5	9	18	35	80	146	218	-
2002	547	1	-	3	7	3	10	15	35	62	174	237	-
2001	555	-	3	5	3	4	10	18	33	69	152	258	-
2000	496	3	2	1	3	7	15	20	33	64	141	207	-
1999	413	2	2	5	2	4	3	8	26	57	118	186	-
Hispanic, male													
2005	317	2	1	2	1	7	4	19	25	36	96	124	-
2004	268	2	2	1	5	2	3	10	25	42	89	87	-
2003	263	2	2	1	5	3	6	13	20	47	84	80	-
2002	282	-	-	1	5	2	5	8	24	32	105	100	-
2001	283	-	1	2	1	3	5	10	22	42	81	116	-
2000	253	-	-	-	2	2	8	15	20	35	87	84	-
1999	195	1	1	1	-	3	1	4	14	31	65	74	-
Hispanic, female													
2005	274	1	1	1	-	1	4	9	9	34	81	133	-
2004	268	-	1	-	3	3	3	8	10	23	91	126	-
2003	261	-	1	1	1	2	3	5	15	33	62	138	-
2002	265	1	-	2	2	1	5	7	11	30	69	137	-
2001	272	-	2	3	2	1	5	8	11	27	71	142	-
2000	243	3	2	1	1	5	7	5	13	29	54	123	-
1999	218	1	1	4	2	1	2	4	12	26	53	112	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Pneumonitis due to solids and liquids (J69)													
Non-Hispanic 1/, both sexes													
2005	16,660	14	6	10	41	58	164	447	797	1,666	5,370	8,087	-
2004	16,213	10	6	10	31	63	153	376	706	1,698	5,375	7,785	-
2003	16,769	10	7	14	40	64	157	373	736	1,668	5,630	8,070	-
2002	17,006	1	7	13	36	71	182	336	634	1,723	5,717	8,286	-
2001	16,700	12	11	10	31	75	167	358	624	1,756	5,593	8,063	-
2000	16,094	7	9	12	34	55	166	343	562	1,833	5,346	7,727	-
1999	14,812	14	10	11	30	74	162	274	559	1,691	4,895	7,092	-
Non-Hispanic 1/, male													
2005	8,423	8	2	6	21	36	84	260	480	1,024	2,955	3,547	-
2004	8,312	2	3	5	15	38	94	240	424	1,039	2,966	3,486	-
2003	8,415	5	3	8	26	39	83	225	449	1,018	3,089	3,470	-
2002	8,565	-	3	8	21	40	114	218	367	1,045	3,130	3,619	-
2001	8,270	6	5	5	21	37	110	221	361	1,061	3,057	3,386	-
2000	8,107	4	5	9	21	33	107	207	333	1,158	2,937	3,293	-
1999	7,534	8	2	5	22	44	95	159	350	1,046	2,803	3,000	-
Non-Hispanic 1/, female													
2005	8,237	6	4	4	20	22	80	187	317	642	2,415	4,540	-
2004	7,901	8	3	5	16	25	59	136	282	659	2,409	4,299	-
2003	8,354	5	4	6	14	25	74	148	287	650	2,541	4,600	-
2002	8,441	1	4	5	15	31	68	118	267	678	2,587	4,667	-
2001	8,430	6	6	5	10	38	57	137	263	695	2,536	4,677	-
2000	7,987	3	4	3	13	22	59	136	229	675	2,409	4,434	-
1999	7,278	6	8	6	8	30	67	115	209	645	2,092	4,092	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Pneumonitis due to solids and liquids (J69)													
Non-Hispanic white, both sexes													
2005 2/	14,739	10	1	6	31	45	135	354	646	1,391	4,750	7,370	-
2004 2/	14,411	3	4	8	20	47	116	297	551	1,430	4,811	7,124	-
2003 2/	14,845	4	6	11	23	51	115	307	579	1,397	4,978	7,374	-
2002	15,203	1	5	11	27	56	138	262	519	1,437	5,131	7,616	-
2001	14,843	9	9	6	22	58	131	280	518	1,469	5,012	7,329	-
2000	14,343	6	7	11	24	40	128	259	449	1,515	4,793	7,111	-
1999	13,139	11	6	7	20	53	118	209	446	1,400	4,361	6,508	-
Non-Hispanic white, male													
2005 2/	7,430	6	-	5	16	25	71	201	387	853	2,611	3,255	-
2004 2/	7,367	-	2	4	10	30	74	194	321	868	2,664	3,200	-
2003 2/	7,454	1	3	6	16	30	65	187	354	857	2,741	3,194	-
2002	7,654	-	3	7	15	33	86	168	303	874	2,822	3,343	-
2001	7,411	6	3	3	17	32	87	179	298	902	2,765	3,119	-
2000	7,214	4	4	8	16	21	85	154	270	945	2,648	3,059	-
1999	6,642	6	1	4	14	30	70	122	276	860	2,492	2,767	-
Non-Hispanic white, female													
2005 2/	7,309	4	1	1	15	20	64	153	259	538	2,139	4,115	-
2004 2/	7,044	3	2	4	10	17	42	103	230	562	2,147	3,924	-
2003 2/	7,391	3	3	5	7	21	50	120	225	540	2,237	4,180	-
2002	7,549	1	2	4	12	23	52	94	216	563	2,309	4,273	-
2001	7,432	3	6	3	5	26	44	101	220	567	2,247	4,210	-
2000	7,129	2	3	3	8	19	43	105	179	570	2,145	4,052	-
1999	6,497	5	5	3	6	23	48	87	170	540	1,869	3,741	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Pneumonitis due to solids and liquids (J69)													
Non-Hispanic black, both sexes													
2005 2/	1,606	3	5	3	9	12	26	86	136	232	525	569	-
2004 2/	1,488	6	2	1	11	14	35	65	140	239	463	512	-
2003 2/	1,598	4	1	3	14	12	41	61	141	231	533	557	-
2002	1,528	-	2	2	8	14	38	64	98	250	499	553	-
2001	1,584	3	2	4	7	16	33	71	100	251	497	600	-
2000	1,507	1	2	1	9	14	37	79	102	289	468	505	-
1999	1,495	2	4	3	10	18	40	59	105	264	469	521	-
Non-Hispanic black, male													
2005 2/	819	2	2	1	4	11	12	56	83	143	289	216	-
2004 2/	762	1	1	-	5	8	20	39	92	151	241	204	-
2003 2/	776	3	-	2	8	9	17	36	86	137	276	202	-
2002	755	-	-	1	5	6	23	42	56	152	253	217	-
2001	731	-	2	2	2	4	20	38	60	140	252	211	-
2000	757	-	1	1	4	11	21	52	55	190	243	179	-
1999	768	1	1	1	8	12	22	33	68	168	261	193	-
Non-Hispanic black, female													
2005 2/	787	1	3	2	5	1	14	30	53	89	236	353	-
2004 2/	726	5	1	1	6	6	15	26	48	88	222	308	-
2003 2/	822	1	1	1	6	3	24	25	55	94	257	355	-
2002	773	-	2	1	3	8	15	22	42	98	246	336	-
2001	853	3	-	2	5	12	13	33	40	111	245	389	-
2000	750	1	1	-	5	3	16	27	47	99	225	326	-
1999	727	1	3	2	2	6	18	26	37	96	208	328	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Pneumonitis due to solids and liquids (J69)													
Origin not stated 3/, both sexes													
2005	28	-	-	-	1	-	-	1	2	5	11	8	-
2004	31	-	-	-	-	-	-	-	2	4	9	16	-
2003	42	1	-	-	-	-	-	1	2	6	13	19	-
2002	40	-	-	-	1	-	-	-	2	8	9	20	-
2001	46	-	-	-	-	-	2	2	2	5	14	21	-
2000	46	-	-	-	-	-	-	3	3	2	18	20	-
1999	43	-	-	-	-	-	1	2	1	6	17	16	-
Origin not stated 3/, male													
2005	20	-	-	-	1	-	-	1	1	5	6	6	-
2004	13	-	-	-	-	-	-	-	2	4	4	3	-
2003	24	1	-	-	-	-	-	1	2	4	8	8	-
2002	29	-	-	-	1	-	-	-	2	6	8	12	-
2001	23	-	-	-	-	-	2	1	1	3	7	9	-
2000	30	-	-	-	-	-	-	3	1	2	14	10	-
1999	26	-	-	-	-	-	1	2	1	6	7	9	-
Origin not stated 3/, female													
2005	8	-	-	-	-	-	-	-	1	-	5	2	-
2004	18	-	-	-	-	-	-	-	-	-	5	13	-
2003	18	-	-	-	-	-	-	-	-	2	5	11	-
2002	11	-	-	-	-	-	-	-	-	2	1	8	-
2001	23	-	-	-	-	-	-	1	1	2	7	12	-
2000	16	-	-	-	-	-	-	-	2	-	4	10	-
1999	17	-	-	-	-	-	-	-	-	-	10	7	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Other diseases of respiratory system (J00-J06,J30-J39,J67,J70-J98)													
All origins, both sexes													
2005	27,056	303	94	61	110	181	474	1,360	2,676	5,373	9,466	6,955	3
2004	27,310	266	105	62	123	174	537	1,349	2,769	5,523	9,629	6,773	-
2003	25,520	279	86	60	136	210	495	1,335	2,442	5,156	8,810	6,510	1
2002	25,039	278	85	84	126	185	528	1,288	2,440	5,237	8,658	6,130	-
2001	26,178	294	93	78	141	216	570	1,356	2,432	5,510	9,202	6,283	3
2000	25,535	273	87	74	131	253	615	1,332	2,411	5,448	8,900	6,008	3
1999	25,031	252	91	76	115	257	576	1,219	2,260	5,409	8,646	6,130	-
All origins, male													
2005	13,168	187	57	33	60	91	226	737	1,486	2,974	4,787	2,529	1
2004	13,434	138	59	35	61	90	269	723	1,509	3,106	4,898	2,546	-
2003	12,391	163	49	38	76	115	251	704	1,367	2,849	4,395	2,384	-
2002	12,057	164	56	39	68	88	281	656	1,339	2,929	4,261	2,176	-
2001	12,495	186	47	43	76	104	304	737	1,322	3,025	4,453	2,197	1
2000	12,391	154	45	38	73	127	335	712	1,324	2,961	4,415	2,205	2
1999	12,110	145	55	39	69	127	317	637	1,213	3,005	4,328	2,175	-
All origins, female													
2005	13,888	116	37	28	50	90	248	623	1,190	2,399	4,679	4,426	2
2004	13,876	128	46	27	62	84	268	626	1,260	2,417	4,731	4,227	-
2003	13,129	116	37	22	60	95	244	631	1,075	2,307	4,415	4,126	1
2002	12,982	114	29	45	58	97	247	632	1,101	2,308	4,397	3,954	-
2001	13,683	108	46	35	65	112	266	619	1,110	2,485	4,749	4,086	2
2000	13,144	119	42	36	58	126	280	620	1,087	2,487	4,485	3,803	1
1999	12,921	107	36	37	46	130	259	582	1,047	2,404	4,318	3,955	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Other diseases of respiratory system (J00-J06,J30-J39,J67,J70-J98)													
Hispanic, both sexes													
2005	1,591	54	16	9	20	32	47	111	209	356	494	243	-
2004	1,508	53	16	11	17	18	54	97	173	346	486	237	-
2003	1,374	57	14	16	16	27	52	94	185	311	373	229	-
2002	1,273	43	18	14	11	27	39	99	164	308	348	202	-
2001	1,275	41	17	7	18	17	39	106	165	287	363	215	-
2000	1,225	43	17	10	15	34	56	81	167	273	325	204	-
1999	1,157	38	10	8	15	23	45	94	157	269	307	191	-
Hispanic, male													
2005	779	30	5	5	11	20	23	64	112	172	245	92	-
2004	725	26	9	7	8	8	27	58	96	176	230	80	-
2003	686	29	8	9	9	12	33	59	112	167	165	83	-
2002	635	21	14	8	9	18	23	48	92	181	156	65	-
2001	643	21	8	3	9	12	26	58	92	151	175	88	-
2000	608	23	8	4	8	19	34	46	96	139	165	66	-
1999	583	20	6	3	7	12	29	45	89	139	158	75	-
Hispanic, female													
2005	812	24	11	4	9	12	24	47	97	184	249	151	-
2004	783	27	7	4	9	10	27	39	77	170	256	157	-
2003	688	28	6	7	7	15	19	35	73	144	208	146	-
2002	638	22	4	6	2	9	16	51	72	127	192	137	-
2001	632	20	9	4	9	5	13	48	73	136	188	127	-
2000	617	20	9	6	7	15	22	35	71	134	160	138	-
1999	574	18	4	5	8	11	16	49	68	130	149	116	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Other diseases of respiratory system (J00-J06,J30-J39,J67,J70-J98)													
Non-Hispanic 1/, both sexes													
2005	25,434	248	78	52	90	147	427	1,248	2,463	5,010	8,959	6,709	3
2004	25,769	212	89	51	106	156	482	1,252	2,590	5,168	9,131	6,532	-
2003	24,094	221	72	43	120	180	443	1,237	2,251	4,833	8,425	6,268	1
2002	23,706	235	67	70	115	158	487	1,177	2,268	4,916	8,296	5,917	-
2001	24,835	251	76	71	123	198	528	1,245	2,265	5,203	8,815	6,057	3
2000	24,236	228	69	64	115	218	556	1,246	2,238	5,158	8,552	5,790	2
1999	23,802	211	81	68	100	233	529	1,120	2,092	5,127	8,315	5,926	-
Non-Hispanic 1/, male													
2005	12,369	156	52	28	49	70	203	672	1,370	2,799	4,533	2,436	1
2004	12,687	112	50	28	53	82	241	665	1,408	2,923	4,660	2,465	-
2003	11,675	133	41	29	67	101	218	642	1,250	2,674	4,222	2,298	-
2002	11,397	143	42	31	59	70	258	602	1,243	2,743	4,098	2,108	-
2001	11,818	165	39	40	67	92	276	677	1,228	2,860	4,265	2,108	1
2000	11,752	130	37	34	65	108	298	663	1,227	2,815	4,239	2,135	1
1999	11,483	123	49	36	62	115	286	589	1,117	2,857	4,153	2,096	-
Non-Hispanic 1/, female													
2005	13,065	92	26	24	41	77	224	576	1,093	2,211	4,426	4,273	2
2004	13,082	100	39	23	53	74	241	587	1,182	2,245	4,471	4,067	-
2003	12,419	88	31	14	53	79	225	595	1,001	2,159	4,203	3,970	1
2002	12,309	92	25	39	56	88	229	575	1,025	2,173	4,198	3,809	-
2001	13,017	86	37	31	56	106	252	568	1,037	2,343	4,550	3,949	2
2000	12,484	98	32	30	50	110	258	583	1,011	2,343	4,313	3,655	1
1999	12,319	88	32	32	38	118	243	531	975	2,270	4,162	3,830	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Other diseases of respiratory system (J00-J06,J30-J39,J67,J70-J98)													
Non-Hispanic white, both sexes													
2005 2/	22,272	140	50	41	57	92	298	889	2,003	4,349	8,166	6,185	2
2004 2/	22,756	104	52	32	76	100	322	904	2,140	4,559	8,389	6,078	-
2003 2/	21,311	113	39	30	73	120	284	896	1,858	4,285	7,759	5,853	1
2002	20,879	109	39	43	70	105	339	859	1,850	4,329	7,619	5,517	-
2001	21,835	116	44	52	82	122	350	883	1,860	4,597	8,133	5,595	1
2000	21,397	122	46	43	71	148	385	928	1,847	4,557	7,866	5,384	-
1999	20,959	111	52	47	75	155	351	826	1,718	4,511	7,642	5,471	-
Non-Hispanic white, male													
2005 2/	10,794	86	33	21	32	46	149	487	1,111	2,477	4,131	2,220	1
2004 2/	11,218	54	30	17	40	53	155	492	1,178	2,603	4,302	2,294	-
2003 2/	10,314	64	20	20	37	70	149	486	1,038	2,391	3,897	2,142	-
2002	10,043	67	26	19	39	45	178	446	1,023	2,448	3,782	1,970	-
2001	10,433	76	20	31	46	52	202	498	1,018	2,574	3,963	1,953	-
2000	10,349	77	22	24	38	74	205	493	1,021	2,508	3,913	1,974	-
1999	10,154	66	28	26	51	86	200	443	919	2,561	3,848	1,926	-
Non-Hispanic white, female													
2005 2/	11,478	54	17	20	25	46	149	402	892	1,872	4,035	3,965	1
2004 2/	11,538	50	22	15	36	47	167	412	962	1,956	4,087	3,784	-
2003 2/	10,997	49	19	10	36	50	135	410	820	1,894	3,862	3,711	1
2002	10,836	42	13	24	31	60	161	413	827	1,881	3,837	3,547	-
2001	11,402	40	24	21	36	70	148	385	842	2,023	4,170	3,642	1
2000	11,048	45	24	19	33	74	180	435	826	2,049	3,953	3,410	-
1999	10,805	45	24	21	24	69	151	383	799	1,950	3,794	3,545	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Other diseases of respiratory system (J00-J06,J30-J39,J67,J70-J98)													
Non-Hispanic black, both sexes													
2005 2/	2,459	99	23	10	28	49	120	311	366	508	563	381	1
2004 2/	2,374	97	28	15	23	49	150	306	384	470	524	328	-
2003 2/	2,207	97	27	7	41	54	139	304	329	410	493	306	-
2002	2,257	112	28	25	41	47	133	278	353	446	494	300	-
2001	2,462	116	26	17	38	67	163	326	343	482	527	355	2
2000	2,336	102	20	18	39	58	152	284	325	495	527	314	2
1999	2,339	90	26	19	21	73	160	258	317	503	523	349	-
Non-Hispanic black, male													
2005 2/	1,195	66	15	6	15	21	52	156	204	246	273	141	-
2004 2/	1,147	53	15	9	11	27	82	151	194	244	245	116	-
2003 2/	1,083	60	18	6	27	27	59	138	183	217	239	109	-
2002	1,076	67	16	10	19	23	73	136	187	222	229	94	-
2001	1,116	76	14	8	19	34	68	157	173	222	230	114	1
2000	1,126	50	14	10	23	30	84	151	166	242	240	115	1
1999	1,081	52	19	9	9	27	74	132	164	237	235	123	-
Non-Hispanic black, female													
2005 2/	1,264	33	8	4	13	28	68	155	162	262	290	240	1
2004 2/	1,227	44	13	6	12	22	68	155	190	226	279	212	-
2003 2/	1,124	37	9	1	14	27	80	166	146	193	254	197	-
2002	1,181	45	12	15	22	24	60	142	166	224	265	206	-
2001	1,346	40	12	9	19	33	95	169	170	260	297	241	1
2000	1,210	52	6	8	16	28	68	133	159	253	287	199	1
1999	1,258	38	7	10	12	46	86	126	153	266	288	226	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Other diseases of respiratory system (J00-J06,J30-J39,J67,J70-J98)													
Origin not stated 3/, both sexes													
2005	31	1	-	-	-	2	-	1	4	7	13	3	-
2004	33	1	-	-	-	-	1	-	6	9	12	4	-
2003	52	1	-	1	-	3	-	4	6	12	12	13	-
2002	60	-	-	-	-	-	2	12	8	13	14	11	-
2001	68	2	-	-	-	1	3	5	2	20	24	11	-
2000	74	2	1	-	1	1	3	5	6	17	23	14	1
1999	72	3	-	-	-	1	2	5	11	13	24	13	-
Origin not stated 3/, male													
2005	20	1	-	-	-	1	-	1	4	3	9	1	-
2004	22	-	-	-	-	-	1	-	5	7	8	1	-
2003	30	1	-	-	-	2	-	3	5	8	8	3	-
2002	25	-	-	-	-	-	-	6	4	5	7	3	-
2001	34	-	-	-	-	-	2	2	2	14	13	1	-
2000	31	1	-	-	-	-	3	3	1	7	11	4	1
1999	44	2	-	-	-	-	2	3	7	9	17	4	-
Origin not stated 3/, female													
2005	11	-	-	-	-	1	-	-	-	4	4	2	-
2004	11	1	-	-	-	-	-	-	1	2	4	3	-
2003	22	-	-	1	-	1	-	1	1	4	4	10	-
2002	35	-	-	-	-	-	2	6	4	8	7	8	-
2001	34	2	-	-	-	1	1	3	-	6	11	10	-
2000	43	1	1	-	1	1	-	2	5	10	12	10	-
1999	28	1	-	-	-	1	-	2	4	4	7	9	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Peptic ulcer (K25-K28)													
All origins, both sexes													
2005	3,478	2	1	1	8	25	108	281	376	544	1,040	1,092	-
2004	3,682	2	3	2	10	17	90	269	374	582	1,148	1,185	-
2003	3,913	2	1	3	7	25	101	281	383	587	1,279	1,244	-
2002	4,079	1	1	1	-	22	113	272	411	665	1,291	1,302	-
2001	4,491	-	1	5	6	32	118	284	418	718	1,438	1,470	1
2000	4,558	-	-	-	9	14	113	307	443	748	1,415	1,509	-
1999	4,604	2	2	-	2	21	123	255	422	836	1,500	1,439	2
All origins, male													
2005	1,722	1	-	1	6	17	73	203	244	305	512	360	-
2004	1,690	1	2	1	6	14	57	176	228	304	530	371	-
2003	1,880	-	1	1	6	13	60	192	237	346	606	418	-
2002	1,951	-	1	1	-	14	72	177	247	378	622	439	-
2001	2,074	-	1	2	3	21	87	186	273	397	649	454	1
2000	2,201	-	-	-	7	6	79	217	284	442	667	499	-
1999	2,189	-	-	-	2	12	76	169	265	504	712	448	1
All origins, female													
2005	1,756	1	1	-	2	8	35	78	132	239	528	732	-
2004	1,992	1	1	1	4	3	33	93	146	278	618	814	-
2003	2,033	2	-	2	1	12	41	89	146	241	673	826	-
2002	2,128	1	-	-	-	8	41	95	164	287	669	863	-
2001	2,417	-	-	3	3	11	31	98	145	321	789	1,016	-
2000	2,357	-	-	-	2	8	34	90	159	306	748	1,010	-
1999	2,415	2	2	-	-	9	47	86	157	332	788	991	1

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Peptic ulcer (K25-K28)													
Hispanic, both sexes													
2005	175	-	-	-	2	3	4	26	21	29	51	39	-
2004	169	-	-	-	1	2	4	18	30	30	49	35	-
2003	200	1	-	-	1	5	6	30	26	42	56	33	-
2002	187	-	-	-	-	1	5	20	32	39	53	37	-
2001	166	-	-	-	1	-	1	16	22	33	54	38	1
2000	186	-	-	-	-	-	9	17	28	36	50	46	-
1999	172	-	1	-	-	8	4	15	25	37	44	38	-
Hispanic, male													
2005	92	-	-	-	2	2	4	19	14	13	28	10	-
2004	93	-	-	-	1	2	4	12	19	15	27	13	-
2003	114	-	-	-	1	3	6	21	16	30	30	7	-
2002	115	-	-	-	-	1	3	17	23	28	27	16	-
2001	99	-	-	-	1	-	1	11	19	18	37	11	1
2000	97	-	-	-	-	-	6	14	23	21	17	16	-
1999	101	-	-	-	-	5	3	10	18	22	27	16	-
Hispanic, female													
2005	83	-	-	-	-	1	-	7	7	16	23	29	-
2004	76	-	-	-	-	-	-	6	11	15	22	22	-
2003	86	1	-	-	-	2	-	9	10	12	26	26	-
2002	72	-	-	-	-	-	2	3	9	11	26	21	-
2001	67	-	-	-	-	-	-	5	3	15	17	27	-
2000	89	-	-	-	-	-	3	3	5	15	33	30	-
1999	71	-	1	-	-	3	1	5	7	15	17	22	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Peptic ulcer (K25-K28)													
Non-Hispanic 1/, both sexes													
2005	3,293	2	1	1	6	22	103	253	353	513	988	1,051	-
2004	3,506	2	3	2	9	15	85	250	344	551	1,099	1,146	-
2003	3,692	1	1	3	6	20	94	246	354	544	1,215	1,208	-
2002	3,879	1	1	1	-	21	108	251	374	625	1,235	1,262	-
2001	4,308	-	1	5	5	32	117	267	393	681	1,380	1,427	-
2000	4,357	-	-	-	9	14	102	290	414	709	1,361	1,458	-
1999	4,415	2	1	-	2	13	118	239	393	796	1,451	1,398	2
Non-Hispanic 1/, male													
2005	1,624	1	-	1	4	15	68	182	229	291	483	350	-
2004	1,592	1	2	1	5	12	52	163	209	288	503	356	-
2003	1,753	-	1	1	5	10	53	166	219	315	574	409	-
2002	1,829	-	1	1	-	13	69	159	222	349	594	421	-
2001	1,966	-	1	2	2	21	86	175	252	375	609	443	-
2000	2,099	-	-	-	7	6	72	203	261	419	648	483	-
1999	2,079	-	-	-	2	7	72	158	244	481	682	432	1
Non-Hispanic 1/, female													
2005	1,669	1	1	-	2	7	35	71	124	222	505	701	-
2004	1,914	1	1	1	4	3	33	87	135	263	596	790	-
2003	1,939	1	-	2	1	10	41	80	135	229	641	799	-
2002	2,050	1	-	-	-	8	39	92	152	276	641	841	-
2001	2,342	-	-	3	3	11	31	92	141	306	771	984	-
2000	2,258	-	-	-	2	8	30	87	153	290	713	975	-
1999	2,336	2	1	-	-	6	46	81	149	315	769	966	1

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Peptic ulcer (K25-K28)													
Non-Hispanic white, both sexes													
2005 2/	2,818	1	-	1	2	14	72	182	292	429	874	951	-
2004 2/	3,045	1	1	1	8	8	57	181	279	462	983	1,064	-
2003 2/	3,197	-	1	2	3	15	65	174	287	472	1,067	1,111	-
2002	3,394	1	1	-	-	11	72	192	288	537	1,123	1,169	-
2001	3,766	-	1	4	2	16	77	197	308	583	1,248	1,330	-
2000	3,825	-	-	-	9	9	70	210	343	609	1,233	1,342	-
1999	3,859	1	1	-	2	7	73	171	322	677	1,313	1,291	1
Non-Hispanic white, male													
2005 2/	1,361	1	-	1	1	10	45	126	189	240	432	316	-
2004 2/	1,356	-	1	1	4	6	35	120	168	246	442	333	-
2003 2/	1,471	-	1	1	2	7	35	114	175	269	496	371	-
2002	1,569	-	1	-	-	9	47	116	169	296	539	392	-
2001	1,649	-	1	2	1	11	55	122	194	311	543	409	-
2000	1,799	-	-	-	7	4	49	150	217	357	580	435	-
1999	1,783	-	-	-	2	3	41	111	195	409	622	399	1
Non-Hispanic white, female													
2005 2/	1,457	-	-	-	1	4	27	56	103	189	442	635	-
2004 2/	1,689	1	-	-	4	2	22	61	111	216	541	731	-
2003 2/	1,726	-	-	1	1	8	30	60	112	203	571	740	-
2002	1,825	1	-	-	-	2	25	76	119	241	584	777	-
2001	2,117	-	-	2	1	5	22	75	114	272	705	921	-
2000	2,026	-	-	-	2	5	21	60	126	252	653	907	-
1999	2,076	1	1	-	-	4	32	60	127	268	691	892	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Peptic ulcer (K25-K28)													
Non-Hispanic black, both sexes													
2005 2/	358	1	1	-	4	8	28	62	49	69	72	64	-
2004 2/	362	-	1	1	1	6	28	61	59	73	76	56	-
2003 2/	376	1	-	1	1	5	23	68	59	53	107	58	-
2002	376	-	-	1	-	10	35	56	68	69	75	62	-
2001	403	-	-	1	2	15	35	59	68	73	83	67	-
2000	420	-	-	-	-	4	29	70	63	79	91	84	-
1999	443	1	-	-	-	4	43	60	65	92	98	79	1
Non-Hispanic black, male													
2005 2/	207	-	-	-	3	5	20	50	36	42	30	21	-
2004 2/	187	-	1	-	1	5	17	39	38	33	39	14	-
2003 2/	219	-	-	-	1	3	13	49	41	36	57	19	-
2002	209	-	-	1	-	4	21	41	45	41	38	18	-
2001	238	-	-	-	1	10	27	46	47	47	39	21	-
2000	228	-	-	-	-	2	21	45	40	48	45	27	-
1999	240	-	-	-	-	3	30	43	46	55	39	24	-
Non-Hispanic black, female													
2005 2/	151	1	1	-	1	3	8	12	13	27	42	43	-
2004 2/	175	-	-	1	-	1	11	22	21	40	37	42	-
2003 2/	157	1	-	1	-	2	10	19	18	17	50	39	-
2002	167	-	-	-	-	6	14	15	23	28	37	44	-
2001	165	-	-	1	1	5	8	13	21	26	44	46	-
2000	192	-	-	-	-	2	8	25	23	31	46	57	-
1999	203	1	-	-	-	1	13	17	19	37	59	55	1

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Peptic ulcer (K25-K28)													
Origin not stated 3/, both sexes													
2005	10	-	-	-	-	-	1	2	2	2	1	2	-
2004	7	-	-	-	-	-	1	1	-	1	-	4	-
2003	21	-	-	-	-	-	1	5	3	1	8	3	-
2002	13	-	-	-	-	-	-	1	5	1	3	3	-
2001	17	-	-	-	-	-	-	1	3	4	4	5	-
2000	15	-	-	-	-	-	2	-	1	3	4	5	-
1999	17	-	-	-	-	-	1	1	4	3	5	3	-
Origin not stated 3/, male													
2005	6	-	-	-	-	-	1	2	1	1	1	-	-
2004	5	-	-	-	-	-	1	1	-	1	-	2	-
2003	13	-	-	-	-	-	1	5	2	1	2	2	-
2002	7	-	-	-	-	-	-	1	2	1	1	2	-
2001	9	-	-	-	-	-	-	-	2	4	3	-	-
2000	5	-	-	-	-	-	1	-	-	2	2	-	-
1999	9	-	-	-	-	-	1	1	3	1	3	-	-
Origin not stated 3/, female													
2005	4	-	-	-	-	-	-	-	1	1	-	2	-
2004	2	-	-	-	-	-	-	-	-	-	-	2	-
2003	8	-	-	-	-	-	-	-	1	-	6	1	-
2002	6	-	-	-	-	-	-	-	3	-	2	1	-
2001	8	-	-	-	-	-	-	1	1	-	1	5	-
2000	10	-	-	-	-	-	1	-	1	1	2	5	-
1999	8	-	-	-	-	-	-	-	1	2	2	3	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Diseases of appendix (K35-K38)													
All origins, both sexes													
2005	439	2	4	12	9	9	19	33	57	76	110	108	-
2004	452	3	4	13	8	9	14	45	52	81	138	85	-
2003	439	4	5	11	11	16	25	50	43	64	132	78	-
2002	480	3	-	7	12	10	27	54	64	89	120	94	-
2001	428	1	8	6	15	15	26	40	45	89	90	93	-
2000	435	3	5	15	11	11	27	43	60	71	121	68	-
1999	390	3	2	26	5	6	24	49	48	69	96	62	-
All origins, male													
2005	247	-	2	9	7	4	15	18	37	53	57	45	-
2004	253	3	4	11	4	7	13	32	29	43	70	37	-
2003	258	2	2	5	9	9	16	30	30	46	75	34	-
2002	275	1	-	3	7	9	17	39	45	56	67	31	-
2001	252	-	7	5	8	11	17	28	30	57	52	37	-
2000	237	2	2	4	7	10	16	29	43	40	60	24	-
1999	236	1	2	15	5	5	17	37	34	38	47	35	-
All origins, female													
2005	192	2	2	3	2	5	4	15	20	23	53	63	-
2004	199	-	-	2	4	2	1	13	23	38	68	48	-
2003	181	2	3	6	2	7	9	20	13	18	57	44	-
2002	205	2	-	4	5	1	10	15	19	33	53	63	-
2001	176	1	1	1	7	4	9	12	15	32	38	56	-
2000	198	1	3	11	4	1	11	14	17	31	61	44	-
1999	154	2	-	11	-	1	7	12	14	31	49	27	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Diseases of appendix (K35-K38)													
Hispanic, both sexes													
2005	25	1	1	2	1	2	2	2	5	3	4	2	-
2004	33	1	-	2	2	2	3	8	4	8	1	2	-
2003	31	1	-	5	1	2	1	8	1	4	4	4	-
2002	28	1	-	2	3	1	5	1	6	3	4	2	-
2001	30	1	4	1	2	1	1	5	2	5	7	1	-
2000	25	1	1	4	5	2	1	1	5	2	3	-	-
1999	29	1	-	8	2	-	4	4	3	3	3	1	-
Hispanic, male													
2005	12	-	-	2	1	2	1	2	1	1	2	-	-
2004	21	1	-	2	1	1	3	7	2	3	-	1	-
2003	20	1	-	1	1	2	1	4	1	3	3	3	-
2002	16	-	-	2	1	1	2	1	5	2	2	-	-
2001	16	-	3	1	-	-	-	3	1	5	3	-	-
2000	16	-	1	3	3	2	-	1	4	-	2	-	-
1999	20	1	-	4	2	-	4	3	2	2	1	1	-
Hispanic, female													
2005	13	1	1	-	-	-	1	-	4	2	2	2	-
2004	12	-	-	-	1	1	-	1	2	5	1	1	-
2003	11	-	-	4	-	-	-	4	-	1	1	1	-
2002	12	1	-	-	2	-	3	-	1	1	2	2	-
2001	14	1	1	-	2	1	1	2	1	-	4	1	-
2000	9	1	-	1	2	-	1	-	1	2	1	-	-
1999	9	-	-	4	-	-	-	1	1	1	2	-	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Diseases of appendix (K35-K38)													
Non-Hispanic 1/, both sexes													
2005	413	1	3	9	8	7	17	31	52	73	106	106	-
2004	418	2	4	11	6	7	11	37	47	73	137	83	-
2003	407	3	5	6	10	14	24	41	42	60	128	74	-
2002	451	2	-	5	9	9	22	53	57	86	116	92	-
2001	398	-	4	5	13	14	25	35	43	84	83	92	-
2000	410	2	4	11	6	9	26	42	55	69	118	68	-
1999	358	2	2	17	3	6	20	44	45	65	93	61	-
Non-Hispanic 1/, male													
2005	234	-	2	6	6	2	14	16	36	52	55	45	-
2004	231	2	4	9	3	6	10	25	26	40	70	36	-
2003	237	1	2	4	8	7	15	25	29	43	72	31	-
2002	258	1	-	1	6	8	15	38	39	54	65	31	-
2001	236	-	4	4	8	11	17	25	29	52	49	37	-
2000	221	2	1	1	4	8	16	28	39	40	58	24	-
1999	213	-	2	10	3	5	13	33	32	35	46	34	-
Non-Hispanic 1/, female													
2005	179	1	1	3	2	5	3	15	16	21	51	61	-
2004	187	-	-	2	3	1	1	12	21	33	67	47	-
2003	170	2	3	2	2	7	9	16	13	17	56	43	-
2002	193	1	-	4	3	1	7	15	18	32	51	61	-
2001	162	-	-	1	5	3	8	10	14	32	34	55	-
2000	189	-	3	10	2	1	10	14	16	29	60	44	-
1999	145	2	-	7	-	1	7	11	13	30	47	27	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Diseases of appendix (K35-K38)													
Non-Hispanic white, both sexes													
2005 2/	348	1	3	4	3	6	12	22	41	62	94	100	-
2004 2/	344	2	3	6	3	3	7	27	39	62	116	76	-
2003 2/	323	2	3	1	6	8	13	29	31	49	113	68	-
2002	377	1	-	2	4	7	13	39	50	69	108	84	-
2001	313	-	1	3	8	7	16	24	27	75	70	82	-
2000	323	2	3	9	3	2	17	28	45	52	97	65	-
1999	284	2	-	10	1	4	15	33	27	54	81	57	-
Non-Hispanic white, male													
2005 2/	196	-	2	3	2	2	9	11	30	45	50	42	-
2004 2/	189	2	3	6	2	2	6	18	23	34	63	30	-
2003 2/	192	1	2	-	5	3	11	19	22	35	64	30	-
2002	210	-	-	-	2	6	11	28	35	43	60	25	-
2001	182	-	1	2	5	5	12	17	21	47	43	29	-
2000	167	2	1	1	2	2	9	18	30	31	49	22	-
1999	163	-	-	6	1	3	9	24	22	28	38	32	-
Non-Hispanic white, female													
2005 2/	152	1	1	1	1	4	3	11	11	17	44	58	-
2004 2/	155	-	-	-	1	1	1	9	16	28	53	46	-
2003 2/	131	1	1	1	1	5	2	10	9	14	49	38	-
2002	167	1	-	2	2	1	2	11	15	26	48	59	-
2001	131	-	-	1	3	2	4	7	6	28	27	53	-
2000	156	-	2	8	1	-	8	10	15	21	48	43	-
1999	121	2	-	4	-	1	6	9	5	26	43	25	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Diseases of appendix (K35-K38)													
Non-Hispanic black, both sexes													
2005 2/	55	-	-	4	5	1	5	9	9	9	9	4	-
2004 2/	58	-	1	3	2	4	3	8	7	9	16	5	-
2003 2/	70	-	2	5	4	5	11	8	10	9	11	5	-
2002	61	1	-	3	5	2	5	13	6	13	5	8	-
2001	72	-	2	2	5	6	8	9	16	7	11	6	-
2000	77	-	-	2	2	7	9	12	10	16	16	3	-
1999	67	-	2	7	1	2	5	10	17	9	11	3	-
Non-Hispanic black, male													
2005 2/	33	-	-	3	4	-	5	5	5	6	4	1	-
2004 2/	38	-	1	2	1	4	3	7	3	6	6	5	-
2003 2/	37	-	-	4	3	3	4	4	6	7	5	1	-
2002	40	1	-	1	4	2	1	10	3	8	4	6	-
2001	47	-	2	2	3	6	5	7	8	4	5	5	-
2000	48	-	-	-	1	6	7	8	9	9	6	2	-
1999	47	-	2	4	1	2	4	9	9	6	8	2	-
Non-Hispanic black, female													
2005 2/	22	-	-	1	1	1	-	4	4	3	5	3	-
2004 2/	20	-	-	1	1	-	-	1	4	3	10	-	-
2003 2/	33	-	2	1	1	2	7	4	4	2	6	4	-
2002	21	-	-	2	1	-	4	3	3	5	1	2	-
2001	25	-	-	-	2	-	3	2	8	3	6	1	-
2000	29	-	-	2	1	1	2	4	1	7	10	1	-
1999	20	-	-	3	-	-	1	1	8	3	3	1	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Diseases of appendix (K35-K38)													
Origin not stated 3/, both sexes													
2005	1	-	-	1	-	-	-	-	-	-	-	-	-
2004	1	-	-	-	-	-	-	-	1	-	-	-	-
2003	1	-	-	-	-	-	-	1	-	-	-	-	-
2002	1	-	-	-	-	-	-	-	1	-	-	-	-
2001	-	-	-	-	-	-	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-	-	-	-
1999	3	-	-	1	-	-	-	1	-	1	-	-	-
Origin not stated 3/, male													
2005	1	-	-	1	-	-	-	-	-	-	-	-	-
2004	1	-	-	-	-	-	-	-	1	-	-	-	-
2003	1	-	-	-	-	-	-	1	-	-	-	-	-
2002	1	-	-	-	-	-	-	-	1	-	-	-	-
2001	-	-	-	-	-	-	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-	-	-	-
1999	3	-	-	1	-	-	-	1	-	1	-	-	-
Origin not stated 3/, female													
2005	-	-	-	-	-	-	-	-	-	-	-	-	-
2004	-	-	-	-	-	-	-	-	-	-	-	-	-
2003	-	-	-	-	-	-	-	-	-	-	-	-	-
2002	-	-	-	-	-	-	-	-	-	-	-	-	-
2001	-	-	-	-	-	-	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-	-	-	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Hernia (K40-K46)													
All origins, both sexes													
2005	1,639	43	4	2	2	5	39	83	181	210	454	616	-
2004	1,602	40	4	4	9	10	36	88	130	239	509	533	-
2003	1,613	37	6	6	3	5	28	105	161	237	462	563	-
2002	1,595	29	3	7	5	11	35	77	126	225	489	588	-
2001	1,522	43	2	3	3	15	34	82	131	213	443	553	-
2000	1,522	46	4	3	5	5	31	84	113	210	466	555	-
1999	1,464	44	3	2	3	16	39	55	107	215	451	529	-
All origins, male													
2005	646	28	1	1	-	1	15	45	91	86	175	203	-
2004	652	20	1	2	5	7	22	45	62	120	198	170	-
2003	668	20	4	4	1	4	16	59	72	111	199	178	-
2002	599	18	3	5	4	7	17	35	56	103	189	162	-
2001	587	28	1	-	1	10	16	44	64	94	173	156	-
2000	586	28	3	3	3	1	13	45	49	92	195	154	-
1999	597	26	2	1	3	8	23	32	60	93	192	157	-
All origins, female													
2005	993	15	3	1	2	4	24	38	90	124	279	413	-
2004	950	20	3	2	4	3	14	43	68	119	311	363	-
2003	945	17	2	2	2	1	12	46	89	126	263	385	-
2002	996	11	-	2	1	4	18	42	70	122	300	426	-
2001	935	15	1	3	2	5	18	38	67	119	270	397	-
2000	936	18	1	-	2	4	18	39	64	118	271	401	-
1999	867	18	1	1	-	8	16	23	47	122	259	372	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Hernia (K40-K46)													
Hispanic, both sexes													
2005	86	9	1	-	1	-	6	3	9	13	18	26	-
2004	88	11	-	1	-	2	2	7	8	19	24	14	-
2003	88	3	2	-	-	-	2	14	15	15	23	14	-
2002	81	3	-	2	2	1	5	6	14	14	19	15	-
2001	92	9	-	-	2	1	7	8	9	12	24	20	-
2000	85	9	-	1	1	1	1	7	9	14	23	19	-
1999	66	3	1	-	-	-	3	4	5	16	17	17	-
Hispanic, male													
2005	28	3	-	-	-	-	2	1	5	5	4	8	-
2004	31	4	-	-	-	2	2	4	1	8	6	4	-
2003	38	2	2	-	-	-	1	8	5	7	7	6	-
2002	35	2	-	2	2	1	2	4	7	4	7	4	-
2001	32	7	-	-	1	-	3	3	3	4	6	5	-
2000	33	6	-	1	-	1	1	5	3	3	10	3	-
1999	27	2	1	-	-	-	2	3	4	6	6	3	-
Hispanic, female													
2005	58	6	1	-	1	-	4	2	4	8	14	18	-
2004	57	7	-	1	-	-	-	3	7	11	18	10	-
2003	50	1	-	-	-	-	1	6	10	8	16	8	-
2002	46	1	-	-	-	-	3	2	7	10	12	11	-
2001	60	2	-	-	1	1	4	5	6	8	18	15	-
2000	52	3	-	-	1	-	-	2	6	11	13	16	-
1999	39	1	-	-	-	-	1	1	1	10	11	14	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Hernia (K40-K46)													
Non-Hispanic 1/, both sexes													
2005	1,553	34	3	2	1	5	33	80	172	197	436	590	-
2004	1,510	29	4	3	9	8	34	79	121	220	484	519	-
2003	1,518	34	4	6	3	5	26	90	146	221	435	548	-
2002	1,504	26	3	5	3	10	29	71	111	209	465	572	-
2001	1,425	33	2	3	1	14	27	74	121	201	417	532	-
2000	1,431	37	4	2	4	4	30	76	103	194	442	535	-
1999	1,390	41	2	2	3	16	36	50	102	198	431	509	-
Non-Hispanic 1/, male													
2005	618	25	1	1	-	1	13	44	86	81	171	195	-
2004	617	16	1	2	5	5	20	39	60	112	191	166	-
2003	626	18	2	4	1	4	15	50	67	103	190	172	-
2002	558	16	3	3	2	6	14	31	49	98	178	158	-
2001	552	20	1	-	-	10	13	41	60	90	166	151	-
2000	550	22	3	2	3	-	12	40	45	88	184	151	-
1999	566	24	1	1	3	8	21	28	56	86	185	153	-
Non-Hispanic 1/, female													
2005	935	9	2	1	1	4	20	36	86	116	265	395	-
2004	893	13	3	1	4	3	14	40	61	108	293	353	-
2003	892	16	2	2	2	1	11	40	79	118	245	376	-
2002	946	10	-	2	1	4	15	40	62	111	287	414	-
2001	873	13	1	3	1	4	14	33	61	111	251	381	-
2000	881	15	1	-	1	4	18	36	58	106	258	384	-
1999	824	17	1	1	-	8	15	22	46	112	246	356	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Hernia (K40-K46)													
Non-Hispanic white, both sexes													
2005 2/	1,372	26	3	1	-	4	23	62	140	169	398	546	-
2004 2/	1,335	22	2	3	7	7	23	61	99	189	444	478	-
2003 2/	1,327	26	1	3	2	3	18	70	116	192	389	507	-
2002	1,343	21	2	5	3	5	20	52	88	176	431	540	-
2001	1,248	22	2	1	-	12	18	59	97	173	374	490	-
2000	1,266	27	2	2	2	3	21	59	90	169	399	492	-
1999	1,227	31	2	2	2	10	27	36	89	167	390	471	-
Non-Hispanic white, male													
2005 2/	540	19	1	-	-	1	10	37	71	71	153	177	-
2004 2/	540	12	-	2	4	4	13	33	51	95	175	151	-
2003 2/	544	14	1	2	1	2	10	40	57	88	172	157	-
2002	493	12	2	3	2	2	11	26	41	81	167	146	-
2001	469	11	1	-	-	8	10	29	46	81	149	134	-
2000	486	17	2	2	1	-	9	29	40	78	170	138	-
1999	505	18	1	1	2	5	18	22	53	79	171	135	-
Non-Hispanic white, female													
2005 2/	832	7	2	1	-	3	13	25	69	98	245	369	-
2004 2/	795	10	2	1	3	3	10	28	48	94	269	327	-
2003 2/	783	12	-	1	1	1	8	30	59	104	217	350	-
2002	850	9	-	2	1	3	9	26	47	95	264	394	-
2001	779	11	1	1	-	4	8	30	51	92	225	356	-
2000	780	10	-	-	1	3	12	30	50	91	229	354	-
1999	722	13	1	1	-	5	9	14	36	88	219	336	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Hernia (K40-K46)													
Non-Hispanic black, both sexes													
2005 2/	151	5	-	1	1	1	8	16	28	25	32	34	-
2004 2/	148	6	2	-	2	1	9	16	20	28	31	33	-
2003 2/	163	8	3	3	-	2	8	18	28	24	38	31	-
2002	137	4	1	-	-	3	8	17	21	27	30	26	-
2001	155	10	-	1	1	2	9	15	23	25	33	36	-
2000	142	7	2	-	2	1	8	13	12	21	36	40	-
1999	139	9	-	-	1	4	9	12	12	24	37	31	-
Non-Hispanic black, male													
2005 2/	66	4	-	1	-	-	3	7	12	10	16	13	-
2004 2/	65	4	1	-	1	1	5	6	9	16	11	11	-
2003 2/	71	4	1	2	-	2	5	9	10	13	14	11	-
2002	56	3	1	-	-	2	3	5	7	15	11	9	-
2001	74	8	-	-	-	2	3	12	13	7	14	15	-
2000	52	3	1	-	2	-	2	9	4	9	11	11	-
1999	53	6	-	-	1	1	3	6	2	7	13	14	-
Non-Hispanic black, female													
2005 2/	85	1	-	-	1	1	5	9	16	15	16	21	-
2004 2/	83	2	1	-	1	-	4	10	11	12	20	22	-
2003 2/	92	4	2	1	-	-	3	9	18	11	24	20	-
2002	81	1	-	-	-	1	5	12	14	12	19	17	-
2001	81	2	-	1	1	-	6	3	10	18	19	21	-
2000	90	4	1	-	-	1	6	4	8	12	25	29	-
1999	86	3	-	-	-	3	6	6	10	17	24	17	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Hernia (K40-K46)													
Origin not stated 3/, both sexes													
2005	-	-	-	-	-	-	-	-	-	-	-	-	-
2004	4	-	-	-	-	-	-	2	1	-	1	-	-
2003	7	-	-	-	-	-	-	1	-	1	4	1	-
2002	10	-	-	-	-	-	1	-	1	2	5	1	-
2001	5	1	-	-	-	-	-	-	1	-	2	1	-
2000	6	-	-	-	-	-	-	1	1	2	1	1	-
1999	8	-	-	-	-	-	-	1	-	1	3	3	-
Origin not stated 3/, male													
2005	-	-	-	-	-	-	-	-	-	-	-	-	-
2004	4	-	-	-	-	-	-	2	1	-	1	-	-
2003	4	-	-	-	-	-	-	1	-	1	2	-	-
2002	6	-	-	-	-	-	1	-	-	1	4	-	-
2001	3	1	-	-	-	-	-	-	1	-	1	-	-
2000	3	-	-	-	-	-	-	-	1	1	1	-	-
1999	4	-	-	-	-	-	-	1	-	1	1	1	-
Origin not stated 3/, female													
2005	-	-	-	-	-	-	-	-	-	-	-	-	-
2004	-	-	-	-	-	-	-	-	-	-	-	-	-
2003	3	-	-	-	-	-	-	-	-	-	2	1	-
2002	4	-	-	-	-	-	-	-	1	1	1	1	-
2001	2	-	-	-	-	-	-	-	-	-	1	1	-
2000	3	-	-	-	-	-	-	1	-	1	-	1	-
1999	4	-	-	-	-	-	-	-	-	-	2	2	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Chronic liver disease and cirrhosis (K70,K73-K74)													
All origins, both sexes													
2005	27,530	10	1	2	23	311	2,688	7,517	7,126	5,066	3,781	1,002	3
2004	27,013	9	1	1	19	309	2,799	7,496	6,569	5,119	3,733	957	1
2003	27,503	2	-	-	16	358	3,020	7,466	6,428	5,406	3,858	946	3
2002	27,257	6	3	6	31	374	3,154	7,216	6,097	5,381	4,000	985	4
2001	27,035	3	-	2	26	387	3,336	7,259	5,750	5,486	3,801	979	6
2000	26,552	3	3	3	36	415	3,371	6,654	5,774	5,482	3,829	979	3
1999	26,259	5	2	2	26	407	3,302	6,368	5,637	5,642	3,901	965	2
All origins, male													
2005	17,937	6	1	1	13	196	1,786	5,423	5,094	3,052	1,920	442	3
2004	17,545	6	-	-	9	221	1,916	5,382	4,587	3,122	1,917	385	-
2003	17,912	2	-	-	8	252	2,033	5,514	4,537	3,265	1,901	397	3
2002	17,401	6	1	3	14	236	2,079	5,285	4,228	3,197	1,953	395	4
2001	17,393	1	-	2	12	259	2,206	5,295	4,028	3,335	1,833	416	6
2000	17,214	3	1	2	19	280	2,314	4,975	4,005	3,317	1,874	421	3
1999	17,115	4	-	1	11	257	2,314	4,829	3,943	3,454	1,905	395	2
All origins, female													
2005	9,593	4	-	1	10	115	902	2,094	2,032	2,014	1,861	560	-
2004	9,468	3	1	1	10	88	883	2,114	1,982	1,997	1,816	572	1
2003	9,591	-	-	-	8	106	987	1,952	1,891	2,141	1,957	549	-
2002	9,856	-	2	3	17	138	1,075	1,931	1,869	2,184	2,047	590	-
2001	9,642	2	-	-	14	128	1,130	1,964	1,722	2,151	1,968	563	-
2000	9,338	-	2	1	17	135	1,057	1,679	1,769	2,165	1,955	558	-
1999	9,144	1	2	1	15	150	988	1,539	1,694	2,188	1,996	570	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Chronic liver disease and cirrhosis (K70,K73-K74)													
Hispanic, both sexes													
2005	3,555	4	1	-	3	78	467	1,003	877	645	394	81	2
2004	3,383	1	-	1	5	73	468	998	769	601	412	55	-
2003	3,382	1	-	-	2	100	469	927	802	624	397	60	-
2002	3,409	1	1	2	9	81	507	983	758	663	321	82	1
2001	3,301	-	-	-	5	81	530	951	721	618	335	59	1
2000	3,187	-	1	-	7	92	501	854	708	628	323	73	-
1999	2,972	-	-	-	8	80	518	787	638	583	285	73	-
Hispanic, male													
2005	2,561	3	1	-	3	62	369	814	664	408	206	29	2
2004	2,376	1	-	-	3	61	377	825	554	336	198	21	-
2003	2,387	1	-	-	2	79	380	768	616	332	186	23	-
2002	2,437	1	-	1	5	62	410	805	561	407	157	27	1
2001	2,410	-	-	-	2	67	425	782	555	386	170	22	1
2000	2,312	-	-	-	5	82	399	707	531	395	163	30	-
1999	2,143	-	-	-	5	62	421	647	481	360	136	31	-
Hispanic, female													
2005	994	1	-	-	-	16	98	189	213	237	188	52	-
2004	1,007	-	-	1	2	12	91	173	215	265	214	34	-
2003	995	-	-	-	-	21	89	159	186	292	211	37	-
2002	972	-	1	1	4	19	97	178	197	256	164	55	-
2001	891	-	-	-	3	14	105	169	166	232	165	37	-
2000	875	-	1	-	2	10	102	147	177	233	160	43	-
1999	829	-	-	-	3	18	97	140	157	223	149	42	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Chronic liver disease and cirrhosis (K70,K73-K74)													
Non-Hispanic 1/, both sexes													
2005	23,882	6	-	2	20	232	2,207	6,480	6,226	4,408	3,379	921	1
2004	23,542	8	1	-	14	236	2,325	6,455	5,779	4,505	3,317	901	1
2003	24,030	1	-	-	14	257	2,541	6,508	5,596	4,771	3,454	886	2
2002	23,733	5	2	4	22	291	2,632	6,198	5,307	4,700	3,668	902	2
2001	23,629	3	-	1	21	305	2,796	6,280	5,002	4,846	3,455	917	3
2000	23,251	3	2	3	29	319	2,856	5,762	5,041	4,833	3,498	903	2
1999	23,185	5	2	2	18	326	2,767	5,547	4,978	5,044	3,606	888	2
Non-Hispanic 1/, male													
2005	15,302	3	-	1	10	134	1,406	4,579	4,411	2,633	1,711	413	1
2004	15,104	5	-	-	6	160	1,535	4,527	4,016	2,775	1,717	363	-
2003	15,452	1	-	-	6	172	1,646	4,719	3,894	2,925	1,713	374	2
2002	14,876	5	1	2	9	172	1,658	4,452	3,641	2,778	1,788	368	2
2001	14,914	1	-	1	10	192	1,777	4,493	3,450	2,934	1,661	392	3
2000	14,814	3	1	2	14	195	1,903	4,237	3,452	2,908	1,708	389	2
1999	14,896	4	-	1	6	194	1,881	4,155	3,444	3,081	1,765	363	2
Non-Hispanic 1/, female													
2005	8,580	3	-	1	10	98	801	1,901	1,815	1,775	1,668	508	-
2004	8,438	3	1	-	8	76	790	1,928	1,763	1,730	1,600	538	1
2003	8,578	-	-	-	8	85	895	1,789	1,702	1,846	1,741	512	-
2002	8,857	-	1	2	13	119	974	1,746	1,666	1,922	1,880	534	-
2001	8,715	2	-	-	11	113	1,019	1,787	1,552	1,912	1,794	525	-
2000	8,437	-	1	1	15	124	953	1,525	1,589	1,925	1,790	514	-
1999	8,289	1	2	1	12	132	886	1,392	1,534	1,963	1,841	525	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Chronic liver disease and cirrhosis (K70,K73-K74)													
Non-Hispanic white, both sexes													
2005 2/	20,428	4	-	2	12	156	1,814	5,323	5,258	3,929	3,079	851	-
2004 2/	20,144	6	1	-	7	178	1,863	5,330	4,838	4,003	3,084	833	1
2003 2/	20,602	-	-	-	7	186	2,068	5,310	4,706	4,256	3,239	828	2
2002	20,353	4	2	3	17	204	2,148	5,023	4,483	4,205	3,417	845	2
2001	20,065	3	-	1	5	227	2,212	5,035	4,189	4,307	3,222	862	2
2000	19,739	-	2	2	14	217	2,240	4,637	4,235	4,285	3,264	842	1
1999	19,602	4	-	1	7	243	2,148	4,430	4,117	4,482	3,347	821	2
Non-Hispanic white, male													
2005 2/	13,088	2	-	1	4	92	1,167	3,799	3,738	2,348	1,552	385	-
2004 2/	12,931	3	-	-	4	121	1,240	3,765	3,387	2,463	1,606	342	-
2003 2/	13,254	-	-	-	3	126	1,366	3,894	3,285	2,627	1,603	348	2
2002	12,800	4	1	2	8	130	1,371	3,670	3,097	2,489	1,681	345	2
2001	12,609	1	-	1	4	145	1,403	3,645	2,875	2,606	1,557	370	2
2000	12,640	-	1	1	6	140	1,518	3,477	2,935	2,602	1,597	362	1
1999	12,613	4	-	-	3	151	1,478	3,365	2,885	2,746	1,641	338	2
Non-Hispanic white, female													
2005 2/	7,340	2	-	1	8	64	647	1,524	1,520	1,581	1,527	466	-
2004 2/	7,213	3	1	-	3	57	623	1,565	1,451	1,540	1,478	491	1
2003 2/	7,348	-	-	-	4	60	702	1,416	1,421	1,629	1,636	480	-
2002	7,553	-	1	1	9	74	777	1,353	1,386	1,716	1,736	500	-
2001	7,456	2	-	-	1	82	809	1,390	1,314	1,701	1,665	492	-
2000	7,099	-	1	1	8	77	722	1,160	1,300	1,683	1,667	480	-
1999	6,989	-	-	1	4	92	670	1,065	1,232	1,736	1,706	483	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Chronic liver disease and cirrhosis (K70,K73-K74)													
Non-Hispanic black, both sexes													
2005 2/	2,495	2	-	-	4	33	234	884	751	352	190	44	1
2004 2/	2,506	2	-	-	4	22	307	890	728	371	140	42	-
2003 2/	2,582	1	-	-	3	41	323	936	700	401	135	42	-
2002	2,545	1	-	1	5	49	313	956	649	358	173	40	-
2001	2,733	-	-	-	10	45	420	993	649	412	169	34	1
2000	2,705	2	-	1	12	56	446	929	622	429	163	44	1
1999	2,793	-	2	1	10	54	453	905	691	432	193	52	-
Non-Hispanic black, male													
2005 2/	1,634	1	-	-	4	14	141	599	528	225	105	16	1
2004 2/	1,650	2	-	-	-	14	197	608	501	242	72	14	-
2003 2/	1,696	1	-	-	2	25	188	664	492	233	70	21	-
2002	1,592	1	-	-	1	26	180	637	436	221	75	15	-
2001	1,819	-	-	-	2	25	273	702	462	268	75	11	1
2000	1,709	2	-	1	6	26	279	633	408	255	81	17	1
1999	1,812	-	-	1	3	27	293	649	453	273	96	17	-
Non-Hispanic black, female													
2005 2/	861	1	-	-	-	19	93	285	223	127	85	28	-
2004 2/	856	-	-	-	4	8	110	282	227	129	68	28	-
2003 2/	886	-	-	-	1	16	135	272	208	168	65	21	-
2002	953	-	-	1	4	23	133	319	213	137	98	25	-
2001	914	-	-	-	8	20	147	291	187	144	94	23	-
2000	996	-	-	-	6	30	167	296	214	174	82	27	-
1999	981	-	2	-	7	27	160	256	238	159	97	35	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Chronic liver disease and cirrhosis (K70,K73-K74)													
Origin not stated 3/, both sexes													
2005	93	-	-	-	-	1	14	34	23	13	8	-	-
2004	88	-	-	-	-	-	6	43	21	13	4	1	-
2003	91	-	-	-	-	1	10	31	30	11	7	-	1
2002	115	-	-	-	-	2	15	35	32	18	11	1	1
2001	105	-	-	1	-	1	10	28	27	22	11	3	2
2000	114	-	-	-	-	4	14	38	25	21	8	3	1
1999	102	-	-	-	-	1	17	34	21	15	10	4	-
Origin not stated 3/, male													
2005	74	-	-	-	-	-	11	30	19	11	3	-	-
2004	65	-	-	-	-	-	4	30	17	11	2	1	-
2003	73	-	-	-	-	1	7	27	27	8	2	-	1
2002	88	-	-	-	-	2	11	28	26	12	8	-	1
2001	69	-	-	1	-	-	4	20	23	15	2	2	2
2000	88	-	-	-	-	3	12	31	22	14	3	2	1
1999	76	-	-	-	-	1	12	27	18	13	4	1	-
Origin not stated 3/, female													
2005	19	-	-	-	-	1	3	4	4	2	5	-	-
2004	23	-	-	-	-	-	2	13	4	2	2	-	-
2003	18	-	-	-	-	-	3	4	3	3	5	-	-
2002	27	-	-	-	-	-	4	7	6	6	3	1	-
2001	36	-	-	-	-	1	6	8	4	7	9	1	-
2000	26	-	-	-	-	1	2	7	3	7	5	1	-
1999	26	-	-	-	-	-	5	7	3	2	6	3	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Alcoholic liver disease (K70)													
All origins, both sexes													
2005	12,928	-	-	-	7	217	1,806	4,490	3,721	1,844	747	94	2
2004	12,548	-	-	-	11	228	1,871	4,392	3,393	1,841	709	102	1
2003	12,360	-	-	-	6	243	1,971	4,288	3,247	1,788	729	86	2
2002	12,121	-	1	-	8	242	1,971	4,137	3,083	1,849	730	98	2
2001	12,207	-	-	-	8	264	2,145	4,203	2,913	1,848	717	104	5
2000	12,109	-	-	-	8	279	2,226	3,888	2,969	1,891	751	95	2
1999	11,958	-	-	-	8	259	2,186	3,802	2,859	1,935	789	118	2
All origins, male													
2005	9,425	-	-	-	6	147	1,189	3,274	2,850	1,352	537	68	2
2004	9,155	-	-	-	8	167	1,267	3,176	2,549	1,392	527	69	-
2003	9,104	-	-	-	3	182	1,317	3,228	2,459	1,343	518	52	2
2002	8,814	-	1	-	6	162	1,305	3,074	2,337	1,367	495	65	2
2001	8,932	-	-	-	5	179	1,446	3,099	2,242	1,379	505	72	5
2000	8,985	-	-	-	5	202	1,533	2,968	2,216	1,433	561	65	2
1999	8,922	-	-	-	6	165	1,520	2,927	2,194	1,497	538	73	2
All origins, female													
2005	3,503	-	-	-	1	70	617	1,216	871	492	210	26	-
2004	3,393	-	-	-	3	61	604	1,216	844	449	182	33	1
2003	3,256	-	-	-	3	61	654	1,060	788	445	211	34	-
2002	3,307	-	-	-	2	80	666	1,063	746	482	235	33	-
2001	3,275	-	-	-	3	85	699	1,104	671	469	212	32	-
2000	3,124	-	-	-	3	77	693	920	753	458	190	30	-
1999	3,036	-	-	-	2	94	666	875	665	438	251	45	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Alcoholic liver disease (K70)													
Hispanic, both sexes													
2005	1,805	-	-	-	1	61	316	615	450	267	83	10	2
2004	1,685	-	-	-	2	55	325	611	383	205	100	4	-
2003	1,712	-	-	-	1	67	332	551	447	216	90	8	-
2002	1,679	-	-	-	4	53	321	587	392	246	63	12	1
2001	1,729	-	-	-	-	65	341	582	396	240	94	10	1
2000	1,680	-	-	-	4	66	353	526	400	247	78	6	-
1999	1,547	-	-	-	2	50	342	499	356	213	69	16	-
Hispanic, male													
2005	1,508	-	-	-	1	50	255	514	380	225	71	10	2
2004	1,414	-	-	-	2	48	266	539	314	162	79	4	-
2003	1,425	-	-	-	1	53	273	487	380	157	69	5	-
2002	1,418	-	-	-	4	47	271	493	339	211	44	8	1
2001	1,457	-	-	-	-	55	282	494	338	210	68	9	1
2000	1,442	-	-	-	4	62	282	455	344	221	69	5	-
1999	1,294	-	-	-	2	41	276	426	307	180	52	10	-
Hispanic, female													
2005	297	-	-	-	-	11	61	101	70	42	12	-	-
2004	271	-	-	-	-	7	59	72	69	43	21	-	-
2003	287	-	-	-	-	14	59	64	67	59	21	3	-
2002	261	-	-	-	-	6	50	94	53	35	19	4	-
2001	272	-	-	-	-	10	59	88	58	30	26	1	-
2000	238	-	-	-	-	4	71	71	56	26	9	1	-
1999	253	-	-	-	-	9	66	73	49	33	17	6	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Alcoholic liver disease (K70)													
Non-Hispanic 1/, both sexes													
2005	11,067	-	-	-	6	156	1,480	3,853	3,257	1,567	664	84	-
2004	10,814	-	-	-	9	173	1,545	3,753	2,997	1,630	608	98	1
2003	10,599	-	-	-	5	176	1,633	3,718	2,783	1,566	638	78	2
2002	10,377	-	1	-	4	188	1,639	3,527	2,671	1,597	663	86	1
2001	10,423	-	-	-	8	198	1,801	3,603	2,499	1,596	622	94	2
2000	10,370	-	-	-	4	211	1,865	3,337	2,554	1,639	671	88	1
1999	10,353	-	-	-	6	208	1,831	3,283	2,491	1,713	717	102	2
Non-Hispanic 1/, male													
2005	7,870	-	-	-	5	97	927	2,742	2,457	1,118	466	58	-
2004	7,700	-	-	-	6	119	1,000	2,615	2,223	1,225	447	65	-
2003	7,638	-	-	-	2	129	1,041	2,725	2,063	1,181	448	47	2
2002	7,342	-	1	-	2	114	1,027	2,562	1,981	1,150	447	57	1
2001	7,431	-	-	-	5	124	1,162	2,592	1,888	1,158	437	63	2
2000	7,492	-	-	-	1	139	1,243	2,492	1,859	1,208	490	59	1
1999	7,580	-	-	-	4	123	1,235	2,485	1,877	1,308	483	63	2
Non-Hispanic 1/, female													
2005	3,197	-	-	-	1	59	553	1,111	800	449	198	26	-
2004	3,114	-	-	-	3	54	545	1,138	774	405	161	33	1
2003	2,961	-	-	-	3	47	592	993	720	385	190	31	-
2002	3,035	-	-	-	2	74	612	965	690	447	216	29	-
2001	2,992	-	-	-	3	74	639	1,011	611	438	185	31	-
2000	2,878	-	-	-	3	72	622	845	695	431	181	29	-
1999	2,773	-	-	-	2	85	596	798	614	405	234	39	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Alcoholic liver disease (K70)													
Non-Hispanic white, both sexes													
2005 2/	9,372	-	-	-	3	105	1,204	3,229	2,785	1,368	600	78	-
2004 2/	9,153	-	-	-	5	134	1,261	3,153	2,524	1,429	555	91	1
2003 2/	8,902	-	-	-	3	131	1,325	3,068	2,342	1,368	592	71	2
2002	8,713	-	1	-	1	133	1,335	2,907	2,255	1,403	601	76	1
2001	8,635	-	-	-	2	146	1,431	2,935	2,092	1,381	566	80	2
2000	8,556	-	-	-	2	150	1,454	2,703	2,126	1,439	601	81	-
1999	8,470	-	-	-	3	155	1,414	2,649	2,045	1,480	636	86	2
Non-Hispanic white, male													
2005 2/	6,714	-	-	-	2	67	768	2,307	2,119	977	420	54	-
2004 2/	6,558	-	-	-	4	92	818	2,219	1,883	1,073	408	61	-
2003 2/	6,467	-	-	-	1	96	858	2,262	1,753	1,039	413	43	2
2002	6,242	-	1	-	1	87	838	2,143	1,701	1,012	409	49	1
2001	6,209	-	-	-	2	96	922	2,152	1,583	997	401	54	2
2000	6,296	-	-	-	-	105	979	2,068	1,579	1,065	446	54	-
1999	6,288	-	-	-	2	95	967	2,041	1,566	1,131	431	53	2
Non-Hispanic white, female													
2005 2/	2,658	-	-	-	1	38	436	922	666	391	180	24	-
2004 2/	2,595	-	-	-	1	42	443	934	641	356	147	30	1
2003 2/	2,435	-	-	-	2	35	467	806	589	329	179	28	-
2002	2,471	-	-	-	-	46	497	764	554	391	192	27	-
2001	2,426	-	-	-	-	50	509	783	509	384	165	26	-
2000	2,260	-	-	-	2	45	475	635	547	374	155	27	-
1999	2,182	-	-	-	1	60	447	608	479	349	205	33	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Alcoholic liver disease (K70)													
Non-Hispanic black, both sexes													
2005 2/	1,154	-	-	-	2	15	149	445	342	151	45	5	-
2004 2/	1,160	-	-	-	2	9	164	437	360	154	31	3	-
2003 2/	1,206	-	-	-	-	19	187	465	340	157	33	5	-
2002	1,173	-	-	-	3	26	169	461	315	143	48	8	-
2001	1,331	-	-	-	1	26	244	508	324	176	43	9	-
2000	1,370	-	-	-	-	34	285	505	321	169	50	5	1
1999	1,445	-	-	-	2	28	281	507	358	191	64	14	-
Non-Hispanic black, male													
2005 2/	790	-	-	-	2	6	82	309	246	110	32	3	-
2004 2/	805	-	-	-	-	7	106	290	260	121	19	2	-
2003 2/	850	-	-	-	-	14	110	344	242	112	24	4	-
2002	783	-	-	-	1	14	103	314	212	103	30	6	-
2001	926	-	-	-	-	12	160	349	239	135	26	5	-
2000	910	-	-	-	-	17	185	339	208	123	33	4	1
1999	1,001	-	-	-	2	14	178	358	254	146	41	8	-
Non-Hispanic black, female													
2005 2/	364	-	-	-	-	9	67	136	96	41	13	2	-
2004 2/	355	-	-	-	2	2	58	147	100	33	12	1	-
2003 2/	356	-	-	-	-	5	77	121	98	45	9	1	-
2002	390	-	-	-	2	12	66	147	103	40	18	2	-
2001	405	-	-	-	1	14	84	159	85	41	17	4	-
2000	460	-	-	-	-	17	100	166	113	46	17	1	-
1999	444	-	-	-	-	14	103	149	104	45	23	6	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Alcoholic liver disease (K70)													
Origin not stated 3/, both sexes													
2005	56	-	-	-	-	-	10	22	14	10	-	-	-
2004	49	-	-	-	-	-	1	28	13	6	1	-	-
2003	49	-	-	-	-	-	6	19	17	6	1	-	-
2002	65	-	-	-	-	1	11	23	20	6	4	-	-
2001	55	-	-	-	-	1	3	18	18	12	1	-	2
2000	59	-	-	-	-	2	8	25	15	5	2	1	1
1999	58	-	-	-	-	1	13	20	12	9	3	-	-
Origin not stated 3/, male													
2005	47	-	-	-	-	-	7	18	13	9	-	-	-
2004	41	-	-	-	-	-	1	22	12	5	1	-	-
2003	41	-	-	-	-	-	3	16	16	5	1	-	-
2002	54	-	-	-	-	1	7	19	17	6	4	-	-
2001	44	-	-	-	-	-	2	13	16	11	-	-	2
2000	51	-	-	-	-	1	8	21	13	4	2	1	1
1999	48	-	-	-	-	1	9	16	10	9	3	-	-
Origin not stated 3/, female													
2005	9	-	-	-	-	-	3	4	1	1	-	-	-
2004	8	-	-	-	-	-	-	6	1	1	-	-	-
2003	8	-	-	-	-	-	3	3	1	1	-	-	-
2002	11	-	-	-	-	-	4	4	3	-	-	-	-
2001	11	-	-	-	-	1	1	5	2	1	1	-	-
2000	8	-	-	-	-	1	-	4	2	1	-	-	-
1999	10	-	-	-	-	-	4	4	2	-	-	-	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Other chronic liver disease and cirrhosis (K73-K74)													
All origins, both sexes													
2005	14,602	10	1	2	16	94	882	3,027	3,405	3,222	3,034	908	1
2004	14,465	9	1	1	8	81	928	3,104	3,176	3,278	3,024	855	-
2003	15,143	2	-	-	10	115	1,049	3,178	3,181	3,618	3,129	860	1
2002	15,136	6	2	6	23	132	1,183	3,079	3,014	3,532	3,270	887	2
2001	14,828	3	-	2	18	123	1,191	3,056	2,837	3,638	3,084	875	1
2000	14,443	3	3	3	28	136	1,145	2,766	2,805	3,591	3,078	884	1
1999	14,301	5	2	2	18	148	1,116	2,566	2,778	3,707	3,112	847	-
All origins, male													
2005	8,512	6	1	1	7	49	597	2,149	2,244	1,700	1,383	374	1
2004	8,390	6	-	-	1	54	649	2,206	2,038	1,730	1,390	316	-
2003	8,808	2	-	-	5	70	716	2,286	2,078	1,922	1,383	345	1
2002	8,587	6	-	3	8	74	774	2,211	1,891	1,830	1,458	330	2
2001	8,461	1	-	2	7	80	760	2,196	1,786	1,956	1,328	344	1
2000	8,229	3	1	2	14	78	781	2,007	1,789	1,884	1,313	356	1
1999	8,193	4	-	1	5	92	794	1,902	1,749	1,957	1,367	322	-
All origins, female													
2005	6,090	4	-	1	9	45	285	878	1,161	1,522	1,651	534	-
2004	6,075	3	1	1	7	27	279	898	1,138	1,548	1,634	539	-
2003	6,335	-	-	-	5	45	333	892	1,103	1,696	1,746	515	-
2002	6,549	-	2	3	15	58	409	868	1,123	1,702	1,812	557	-
2001	6,367	2	-	-	11	43	431	860	1,051	1,682	1,756	531	-
2000	6,214	-	2	1	14	58	364	759	1,016	1,707	1,765	528	-
1999	6,108	1	2	1	13	56	322	664	1,029	1,750	1,745	525	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Other chronic liver disease and cirrhosis (K73-K74)													
Hispanic, both sexes													
2005	1,750	4	1	-	2	17	151	388	427	378	311	71	-
2004	1,698	1	-	1	3	18	143	387	386	396	312	51	-
2003	1,670	1	-	-	1	33	137	376	355	408	307	52	-
2002	1,730	1	1	2	5	28	186	396	366	417	258	70	-
2001	1,572	-	-	-	5	16	189	369	325	378	241	49	-
2000	1,507	-	1	-	3	26	148	328	308	381	245	67	-
1999	1,425	-	-	-	6	30	176	288	282	370	216	57	-
Hispanic, male													
2005	1,053	3	1	-	2	12	114	300	284	183	135	19	-
2004	962	1	-	-	1	13	111	286	240	174	119	17	-
2003	962	1	-	-	1	26	107	281	236	175	117	18	-
2002	1,019	1	-	1	1	15	139	312	222	196	113	19	-
2001	953	-	-	-	2	12	143	288	217	176	102	13	-
2000	870	-	-	-	1	20	117	252	187	174	94	25	-
1999	849	-	-	-	3	21	145	221	174	180	84	21	-
Hispanic, female													
2005	697	1	-	-	-	5	37	88	143	195	176	52	-
2004	736	-	-	1	2	5	32	101	146	222	193	34	-
2003	708	-	-	-	-	7	30	95	119	233	190	34	-
2002	711	-	1	1	4	13	47	84	144	221	145	51	-
2001	619	-	-	-	3	4	46	81	108	202	139	36	-
2000	637	-	1	-	2	6	31	76	121	207	151	42	-
1999	576	-	-	-	3	9	31	67	108	190	132	36	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Other chronic liver disease and cirrhosis (K73-K74)													
Non-Hispanic 1/, both sexes													
2005	12,815	6	-	2	14	76	727	2,627	2,969	2,841	2,715	837	1
2004	12,728	8	1	-	5	63	780	2,702	2,782	2,875	2,709	803	-
2003	13,431	1	-	-	9	81	908	2,790	2,813	3,205	2,816	808	-
2002	13,356	5	1	4	18	103	993	2,671	2,636	3,103	3,005	816	1
2001	13,206	3	-	1	13	107	995	2,677	2,503	3,250	2,833	823	1
2000	12,881	3	2	3	25	108	991	2,425	2,487	3,194	2,827	815	1
1999	12,832	5	2	2	12	118	936	2,264	2,487	3,331	2,889	786	-
Non-Hispanic 1/, male													
2005	7,432	3	-	1	5	37	479	1,837	1,954	1,515	1,245	355	1
2004	7,404	5	-	-	-	41	535	1,912	1,793	1,550	1,270	298	-
2003	7,814	1	-	-	4	43	605	1,994	1,831	1,744	1,265	327	-
2002	7,534	5	-	2	7	58	631	1,890	1,660	1,628	1,341	311	1
2001	7,483	1	-	1	5	68	615	1,901	1,562	1,776	1,224	329	1
2000	7,322	3	1	2	13	56	660	1,745	1,593	1,700	1,218	330	1
1999	7,316	4	-	1	2	71	646	1,670	1,567	1,773	1,282	300	-
Non-Hispanic 1/, female													
2005	5,383	3	-	1	9	39	248	790	1,015	1,326	1,470	482	-
2004	5,324	3	1	-	5	22	245	790	989	1,325	1,439	505	-
2003	5,617	-	-	-	5	38	303	796	982	1,461	1,551	481	-
2002	5,822	-	1	2	11	45	362	781	976	1,475	1,664	505	-
2001	5,723	2	-	-	8	39	380	776	941	1,474	1,609	494	-
2000	5,559	-	1	1	12	52	331	680	894	1,494	1,609	485	-
1999	5,516	1	2	1	10	47	290	594	920	1,558	1,607	486	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Other chronic liver disease and cirrhosis (K73-K74)													
Non-Hispanic white, both sexes													
2005 2/	11,056	4	-	2	9	51	610	2,094	2,473	2,561	2,479	773	-
2004 2/	10,991	6	1	-	2	44	602	2,177	2,314	2,574	2,529	742	-
2003 2/	11,700	-	-	-	4	55	743	2,242	2,364	2,888	2,647	757	-
2002	11,640	4	1	3	16	71	813	2,116	2,228	2,802	2,816	769	1
2001	11,430	3	-	1	3	81	781	2,100	2,097	2,926	2,656	782	-
2000	11,183	-	2	2	12	67	786	1,934	2,109	2,846	2,663	761	1
1999	11,132	4	-	1	4	88	734	1,781	2,072	3,002	2,711	735	-
Non-Hispanic white, male													
2005 2/	6,374	2	-	1	2	25	399	1,492	1,619	1,371	1,132	331	-
2004 2/	6,373	3	-	-	-	29	422	1,546	1,504	1,390	1,198	281	-
2003 2/	6,787	-	-	-	2	30	508	1,632	1,532	1,588	1,190	305	-
2002	6,558	4	-	2	7	43	533	1,527	1,396	1,477	1,272	296	1
2001	6,400	1	-	1	2	49	481	1,493	1,292	1,609	1,156	316	-
2000	6,344	-	1	1	6	35	539	1,409	1,356	1,537	1,151	308	1
1999	6,325	4	-	-	1	56	511	1,324	1,319	1,615	1,210	285	-
Non-Hispanic white, female													
2005 2/	4,682	2	-	1	7	26	211	602	854	1,190	1,347	442	-
2004 2/	4,618	3	1	-	2	15	180	631	810	1,184	1,331	461	-
2003 2/	4,913	-	-	-	2	25	235	610	832	1,300	1,457	452	-
2002	5,082	-	1	1	9	28	280	589	832	1,325	1,544	473	-
2001	5,030	2	-	-	1	32	300	607	805	1,317	1,500	466	-
2000	4,839	-	1	1	6	32	247	525	753	1,309	1,512	453	-
1999	4,807	-	-	1	3	32	223	457	753	1,387	1,501	450	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Other chronic liver disease and cirrhosis (K73-K74)													
Non-Hispanic black, both sexes													
2005 2/	1,341	2	-	-	2	18	85	439	409	201	145	39	1
2004 2/	1,346	2	-	-	2	13	143	453	368	217	109	39	-
2003 2/	1,376	1	-	-	3	22	136	471	360	244	102	37	-
2002	1,372	1	-	1	2	23	144	495	334	215	125	32	-
2001	1,402	-	-	-	9	19	176	485	325	236	126	25	1
2000	1,335	2	-	1	12	22	161	424	301	260	113	39	-
1999	1,348	-	2	1	8	26	172	398	333	241	129	38	-
Non-Hispanic black, male													
2005 2/	844	1	-	-	2	8	59	290	282	115	73	13	1
2004 2/	845	2	-	-	-	7	91	318	241	121	53	12	-
2003 2/	846	1	-	-	2	11	78	320	250	121	46	17	-
2002	809	1	-	-	-	12	77	323	224	118	45	9	-
2001	893	-	-	-	2	13	113	353	223	133	49	6	1
2000	799	2	-	1	6	9	94	294	200	132	48	13	-
1999	811	-	-	1	1	13	115	291	199	127	55	9	-
Non-Hispanic black, female													
2005 2/	497	1	-	-	-	10	26	149	127	86	72	26	-
2004 2/	501	-	-	-	2	6	52	135	127	96	56	27	-
2003 2/	530	-	-	-	1	11	58	151	110	123	56	20	-
2002	563	-	-	1	2	11	67	172	110	97	80	23	-
2001	509	-	-	-	7	6	63	132	102	103	77	19	-
2000	536	-	-	-	6	13	67	130	101	128	65	26	-
1999	537	-	2	-	7	13	57	107	134	114	74	29	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Other chronic liver disease and cirrhosis (K73-K74)													
Origin not stated 3/, both sexes													
2005	37	-	-	-	-	1	4	12	9	3	8	-	-
2004	39	-	-	-	-	-	5	15	8	7	3	1	-
2003	42	-	-	-	-	1	4	12	13	5	6	-	1
2002	50	-	-	-	-	1	4	12	12	12	7	1	1
2001	50	-	-	1	-	-	7	10	9	10	10	3	-
2000	55	-	-	-	-	2	6	13	10	16	6	2	-
1999	44	-	-	-	-	-	4	14	9	6	7	4	-
Origin not stated 3/, male													
2005	27	-	-	-	-	-	4	12	6	2	3	-	-
2004	24	-	-	-	-	-	3	8	5	6	1	1	-
2003	32	-	-	-	-	1	4	11	11	3	1	-	1
2002	34	-	-	-	-	1	4	9	9	6	4	-	1
2001	25	-	-	1	-	-	2	7	7	4	2	2	-
2000	37	-	-	-	-	2	4	10	9	10	1	1	-
1999	28	-	-	-	-	-	3	11	8	4	1	1	-
Origin not stated 3/, female													
2005	10	-	-	-	-	1	-	-	3	1	5	-	-
2004	15	-	-	-	-	-	2	7	3	1	2	-	-
2003	10	-	-	-	-	-	-	1	2	2	5	-	-
2002	16	-	-	-	-	-	-	3	3	6	3	1	-
2001	25	-	-	-	-	-	5	3	2	6	8	1	-
2000	18	-	-	-	-	-	2	3	1	6	5	1	-
1999	16	-	-	-	-	-	1	3	1	2	6	3	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Cholelithiasis and other disorders of gallbladder (K80-K82)													
All origins, both sexes													
2005	3,072	-	-	4	8	16	42	113	217	448	1,055	1,169	-
2004	3,086	-	-	-	8	17	55	138	247	458	1,008	1,155	-
2003	2,948	-	-	1	4	25	40	126	182	489	1,054	1,027	-
2002	2,979	-	-	-	9	21	54	129	216	504	992	1,054	-
2001	2,965	1	-	1	7	19	44	137	242	526	945	1,043	-
2000	2,810	-	-	2	9	19	53	109	189	450	926	1,053	-
1999	2,830	2	-	1	10	16	59	124	203	475	948	992	-
All origins, male													
2005	1,363	-	-	1	5	8	17	59	118	271	495	389	-
2004	1,370	-	-	-	3	10	30	73	138	248	483	385	-
2003	1,318	-	-	-	4	11	21	64	106	265	519	328	-
2002	1,341	-	-	-	3	11	29	83	122	258	500	335	-
2001	1,341	1	-	-	3	9	25	79	142	269	464	349	-
2000	1,244	-	-	1	4	7	27	53	110	242	461	339	-
1999	1,248	1	-	1	5	4	30	58	120	252	447	330	-
All origins, female													
2005	1,709	-	-	3	3	8	25	54	99	177	560	780	-
2004	1,716	-	-	-	5	7	25	65	109	210	525	770	-
2003	1,630	-	-	1	-	14	19	62	76	224	535	699	-
2002	1,638	-	-	-	6	10	25	46	94	246	492	719	-
2001	1,624	-	-	1	4	10	19	58	100	257	481	694	-
2000	1,566	-	-	1	5	12	26	56	79	208	465	714	-
1999	1,582	1	-	-	5	12	29	66	83	223	501	662	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Cholelithiasis and other disorders of gallbladder (K80-K82)													
Hispanic, both sexes													
2005	225	-	-	1	1	2	10	16	24	45	56	70	-
2004	215	-	-	-	4	1	10	18	23	36	66	57	-
2003	178	-	-	-	2	2	6	13	14	44	61	36	-
2002	196	-	-	-	2	6	11	11	25	42	55	44	-
2001	197	-	-	-	2	3	8	13	23	51	47	50	-
2000	183	-	-	-	2	3	5	18	19	41	51	44	-
1999	185	-	-	-	-	3	11	14	20	34	57	46	-
Hispanic, male													
2005	102	-	-	1	1	2	5	6	9	24	31	23	-
2004	97	-	-	-	2	1	5	9	16	16	34	14	-
2003	94	-	-	-	2	-	3	11	7	24	36	11	-
2002	93	-	-	-	-	2	7	10	14	21	23	16	-
2001	92	-	-	-	-	1	3	5	16	25	25	17	-
2000	99	-	-	-	-	1	5	9	15	25	32	12	-
1999	84	-	-	-	-	1	6	6	12	15	29	15	-
Hispanic, female													
2005	123	-	-	-	-	-	5	10	15	21	25	47	-
2004	118	-	-	-	2	-	5	9	7	20	32	43	-
2003	84	-	-	-	-	2	3	2	7	20	25	25	-
2002	103	-	-	-	2	4	4	1	11	21	32	28	-
2001	105	-	-	-	2	2	5	8	7	26	22	33	-
2000	84	-	-	-	2	2	-	9	4	16	19	32	-
1999	101	-	-	-	-	2	5	8	8	19	28	31	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Cholelithiasis and other disorders of gallbladder (K80-K82)													
Non-Hispanic 1/, both sexes													
2005	2,841	-	-	3	6	14	32	97	191	402	999	1,097	-
2004	2,867	-	-	-	4	16	45	120	224	421	941	1,096	-
2003	2,765	-	-	1	2	23	34	112	168	445	990	990	-
2002	2,776	-	-	-	7	15	43	118	190	460	935	1,008	-
2001	2,764	1	-	1	5	16	36	124	219	473	896	993	-
2000	2,619	-	-	2	7	16	48	91	170	405	873	1,007	-
1999	2,639	2	-	1	10	13	48	110	183	438	889	945	-
Non-Hispanic 1/, male													
2005	1,257	-	-	-	3	6	12	53	107	246	464	366	-
2004	1,272	-	-	-	1	9	25	64	122	231	449	371	-
2003	1,223	-	-	-	2	11	18	52	99	241	483	317	-
2002	1,244	-	-	-	3	9	22	73	107	236	476	318	-
2001	1,248	1	-	-	3	8	22	74	126	243	439	332	-
2000	1,141	-	-	1	4	6	22	44	95	215	428	326	-
1999	1,163	1	-	1	5	3	24	52	108	236	418	315	-
Non-Hispanic 1/, female													
2005	1,584	-	-	3	3	8	20	44	84	156	535	731	-
2004	1,595	-	-	-	3	7	20	56	102	190	492	725	-
2003	1,542	-	-	1	-	12	16	60	69	204	507	673	-
2002	1,532	-	-	-	4	6	21	45	83	224	459	690	-
2001	1,516	-	-	1	2	8	14	50	93	230	457	661	-
2000	1,478	-	-	1	3	10	26	47	75	190	445	681	-
1999	1,476	1	-	-	5	10	24	58	75	202	471	630	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Cholelithiasis and other disorders of gallbladder (K80-K82)													
Non-Hispanic white, both sexes													
2005 2/	2,427	-	-	3	4	11	19	59	149	313	881	988	-
2004 2/	2,460	-	-	-	2	9	35	78	170	338	823	1,005	-
2003 2/	2,391	-	-	1	1	16	22	72	128	377	878	896	-
2002	2,420	-	-	-	5	10	33	82	152	390	818	930	-
2001	2,368	1	-	1	3	12	25	86	168	381	787	904	-
2000	2,272	-	-	1	5	7	33	58	132	325	775	936	-
1999	2,299	-	-	1	6	10	26	85	134	361	805	871	-
Non-Hispanic white, male													
2005 2/	1,087	-	-	-	2	6	8	31	85	198	422	335	-
2004 2/	1,110	-	-	-	1	6	19	41	98	195	401	349	-
2003 2/	1,056	-	-	-	1	6	10	36	75	208	430	290	-
2002	1,087	-	-	-	3	6	17	49	87	207	424	294	-
2001	1,065	1	-	-	2	5	17	52	96	200	395	297	-
2000	1,012	-	-	1	3	4	17	31	77	181	391	307	-
1999	1,017	-	-	1	4	1	14	40	79	201	389	288	-
Non-Hispanic white, female													
2005 2/	1,340	-	-	3	2	5	11	28	64	115	459	653	-
2004 2/	1,350	-	-	-	1	3	16	37	72	143	422	656	-
2003 2/	1,335	-	-	1	-	10	12	36	53	169	448	606	-
2002	1,333	-	-	-	2	4	16	33	65	183	394	636	-
2001	1,303	-	-	1	1	7	8	34	72	181	392	607	-
2000	1,260	-	-	-	2	3	16	27	55	144	384	629	-
1999	1,282	-	-	-	2	9	12	45	55	160	416	583	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Cholelithiasis and other disorders of gallbladder (K80-K82)													
Non-Hispanic black, both sexes													
2005 2/	324	-	-	-	2	3	11	31	35	72	93	77	-
2004 2/	313	-	-	-	-	4	9	36	45	61	90	68	-
2003 2/	285	-	-	-	1	7	11	34	33	50	78	71	-
2002	266	-	-	-	1	5	8	29	28	54	91	50	-
2001	309	-	-	-	2	4	11	31	42	78	77	64	-
2000	274	-	-	1	2	8	14	27	29	61	75	57	-
1999	257	2	-	-	4	3	20	21	35	59	65	48	-
Non-Hispanic black, male													
2005 2/	135	-	-	-	1	-	4	17	17	38	37	21	-
2004 2/	120	-	-	-	-	2	5	21	20	26	32	14	-
2003 2/	126	-	-	-	1	5	8	14	18	23	37	20	-
2002	106	-	-	-	-	3	4	18	16	17	37	11	-
2001	130	-	-	-	1	3	5	18	24	34	21	24	-
2000	92	-	-	-	1	2	4	10	14	23	26	12	-
1999	102	1	-	-	1	2	9	9	19	25	23	13	-
Non-Hispanic black, female													
2005 2/	189	-	-	-	1	3	7	14	18	34	56	56	-
2004 2/	193	-	-	-	-	2	4	15	25	35	58	54	-
2003 2/	159	-	-	-	-	2	3	20	15	27	41	51	-
2002	160	-	-	-	1	2	4	11	12	37	54	39	-
2001	179	-	-	-	1	1	6	13	18	44	56	40	-
2000	182	-	-	1	1	6	10	17	15	38	49	45	-
1999	155	1	-	-	3	1	11	12	16	34	42	35	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Cholelithiasis and other disorders of gallbladder (K80-K82)													
Origin not stated 3/, both sexes													
2005	6	-	-	-	1	-	-	-	2	1	-	2	-
2004	4	-	-	-	-	-	-	-	-	1	1	2	-
2003	5	-	-	-	-	-	-	1	-	-	3	1	-
2002	7	-	-	-	-	-	-	-	1	2	2	2	-
2001	4	-	-	-	-	-	-	-	-	2	2	-	-
2000	8	-	-	-	-	-	-	-	-	4	2	2	-
1999	6	-	-	-	-	-	-	-	-	3	2	1	-
Origin not stated 3/, male													
2005	4	-	-	-	1	-	-	-	2	1	-	-	-
2004	1	-	-	-	-	-	-	-	-	1	-	-	-
2003	1	-	-	-	-	-	-	1	-	-	-	-	-
2002	4	-	-	-	-	-	-	-	1	1	1	1	-
2001	1	-	-	-	-	-	-	-	-	1	-	-	-
2000	4	-	-	-	-	-	-	-	-	2	1	1	-
1999	1	-	-	-	-	-	-	-	-	1	-	-	-
Origin not stated 3/, female													
2005	2	-	-	-	-	-	-	-	-	-	-	2	-
2004	3	-	-	-	-	-	-	-	-	-	1	2	-
2003	4	-	-	-	-	-	-	-	-	-	3	1	-
2002	3	-	-	-	-	-	-	-	-	1	1	1	-
2001	3	-	-	-	-	-	-	-	-	1	2	-	-
2000	4	-	-	-	-	-	-	-	-	2	1	1	-
1999	5	-	-	-	-	-	-	-	-	2	2	1	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)													
All origins, both sexes													
2005	43,901	159	16	23	91	285	742	2,028	4,141	7,320	14,403	14,693	-
2004	42,480	174	14	21	87	246	801	2,067	3,963	7,119	14,058	13,928	2
2003	42,453	181	14	22	95	282	796	2,001	3,806	7,345	14,093	13,816	2
2002	40,974	173	14	21	84	269	749	1,893	3,455	7,164	13,896	13,256	-
2001	39,480	133	16	18	92	225	768	1,822	3,284	7,356	13,097	12,668	1
2000	37,251	162	14	25	88	244	740	1,651	3,100	6,990	12,458	11,777	2
1999	35,525	166	15	31	72	252	707	1,480	2,864	6,841	11,927	11,170	-
All origins, male													
2005	21,268	88	12	10	45	153	423	1,237	2,352	3,733	7,177	6,038	-
2004	20,370	110	10	11	50	113	451	1,208	2,120	3,677	7,007	5,612	1
2003	20,481	106	7	9	59	147	459	1,193	2,065	3,785	7,049	5,600	2
2002	19,695	109	7	12	41	129	407	1,086	1,824	3,802	6,920	5,358	-
2001	18,852	76	7	6	45	112	414	1,053	1,714	3,826	6,485	5,114	-
2000	17,811	103	7	17	43	129	429	982	1,651	3,564	6,213	4,671	2
1999	17,016	90	9	18	33	125	402	846	1,530	3,504	6,031	4,428	-
All origins, female													
2005	22,633	71	4	13	46	132	319	791	1,789	3,587	7,226	8,655	-
2004	22,110	64	4	10	37	133	350	859	1,843	3,442	7,051	8,316	1
2003	21,972	75	7	13	36	135	337	808	1,741	3,560	7,044	8,216	-
2002	21,279	64	7	9	43	140	342	807	1,631	3,362	6,976	7,898	-
2001	20,628	57	9	12	47	113	354	769	1,570	3,530	6,612	7,554	1
2000	19,440	59	7	8	45	115	311	669	1,449	3,426	6,245	7,106	-
1999	18,509	76	6	13	39	127	305	634	1,334	3,337	5,896	6,742	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)													
Hispanic, both sexes													
2005	2,329	34	4	5	15	30	68	165	351	506	699	452	-
2004	2,196	37	3	11	12	22	72	182	319	523	629	386	-
2003	2,170	35	1	4	19	30	76	174	318	484	578	450	1
2002	1,898	35	5	5	9	33	65	157	267	427	532	363	-
2001	1,857	18	2	4	16	25	74	132	237	470	524	355	-
2000	1,646	25	4	3	14	26	53	109	232	404	429	346	1
1999	1,584	20	3	9	6	21	62	107	230	369	426	331	-
Hispanic, male													
2005	1,186	18	3	1	8	16	37	112	201	244	338	208	-
2004	1,101	25	2	4	8	8	45	107	180	251	306	165	-
2003	1,069	19	-	1	13	16	49	107	160	230	280	193	1
2002	937	21	1	3	5	12	38	88	141	214	273	141	-
2001	892	13	-	2	9	10	46	64	132	217	251	148	-
2000	805	16	3	2	7	11	36	69	115	194	206	145	1
1999	814	15	2	7	2	10	31	64	137	178	212	156	-
Hispanic, female													
2005	1,143	16	1	4	7	14	31	53	150	262	361	244	-
2004	1,095	12	1	7	4	14	27	75	139	272	323	221	-
2003	1,101	16	1	3	6	14	27	67	158	254	298	257	-
2002	961	14	4	2	4	21	27	69	126	213	259	222	-
2001	965	5	2	2	7	15	28	68	105	253	273	207	-
2000	841	9	1	1	7	15	17	40	117	210	223	201	-
1999	770	5	1	2	4	11	31	43	93	191	214	175	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)													
Non-Hispanic 1/, both sexes													
2005	41,510	124	12	17	76	254	672	1,856	3,784	6,804	13,690	14,221	-
2004	40,192	137	10	10	75	223	729	1,876	3,637	6,579	13,392	13,523	1
2003	40,172	143	13	18	76	252	717	1,821	3,472	6,848	13,481	13,330	1
2002	38,933	138	9	16	75	235	681	1,725	3,171	6,710	13,322	12,851	-
2001	37,511	113	14	14	76	198	691	1,677	3,036	6,866	12,538	12,288	-
2000	35,509	134	10	22	73	217	686	1,536	2,854	6,567	12,000	11,410	-
1999	33,830	145	12	22	66	231	643	1,368	2,621	6,451	11,474	10,797	-
Non-Hispanic 1/, male													
2005	20,045	69	9	8	37	136	384	1,120	2,147	3,483	6,832	5,820	-
2004	19,220	85	7	7	42	104	406	1,096	1,936	3,415	6,684	5,438	-
2003	19,358	87	7	8	46	131	409	1,082	1,895	3,546	6,753	5,393	1
2002	18,682	88	6	9	36	117	367	992	1,677	3,569	6,620	5,201	-
2001	17,908	61	7	4	36	100	366	982	1,577	3,598	6,220	4,957	-
2000	16,958	84	4	15	35	118	392	908	1,529	3,357	5,995	4,521	-
1999	16,143	74	7	11	31	115	370	777	1,384	3,312	5,808	4,254	-
Non-Hispanic 1/, female													
2005	21,465	55	3	9	39	118	288	736	1,637	3,321	6,858	8,401	-
2004	20,972	52	3	3	33	119	323	780	1,701	3,164	6,708	8,085	1
2003	20,814	56	6	10	30	121	308	739	1,577	3,302	6,728	7,937	-
2002	20,251	50	3	7	39	118	314	733	1,494	3,141	6,702	7,650	-
2001	19,603	52	7	10	40	98	325	695	1,459	3,268	6,318	7,331	-
2000	18,551	50	6	7	38	99	294	628	1,325	3,210	6,005	6,889	-
1999	17,687	71	5	11	35	116	273	591	1,237	3,139	5,666	6,543	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)													
Non-Hispanic white, both sexes													
2005 2/	32,492	62	9	13	45	123	333	1,059	2,317	4,846	11,298	12,387	-
2004 2/	31,508	61	6	7	41	108	394	1,005	2,238	4,745	11,106	11,796	1
2003 2/	31,497	80	9	11	38	128	340	977	2,197	4,891	11,179	11,647	-
2002	30,669	77	5	10	33	119	353	922	1,978	4,870	11,110	11,192	-
2001	29,449	58	11	6	45	100	359	904	1,861	4,984	10,398	10,723	-
2000	27,924	61	4	17	48	110	359	829	1,776	4,741	10,026	9,953	-
1999	26,418	81	10	13	36	116	319	717	1,636	4,718	9,504	9,268	-
Non-Hispanic white, male													
2005 2/	15,953	32	6	6	24	67	201	656	1,345	2,625	5,802	5,189	-
2004 2/	15,247	37	4	4	21	47	228	586	1,234	2,540	5,721	4,825	-
2003 2/	15,323	52	5	6	24	64	205	577	1,216	2,614	5,750	4,810	-
2002	14,883	49	3	6	17	64	191	533	1,045	2,694	5,687	4,594	-
2001	14,331	31	4	3	21	45	211	543	1,012	2,729	5,310	4,422	-
2000	13,567	41	2	12	25	62	206	491	990	2,574	5,141	4,023	-
1999	12,793	46	5	8	17	60	188	406	903	2,532	4,907	3,721	-
Non-Hispanic white, female													
2005 2/	16,539	30	3	7	21	56	132	403	972	2,221	5,496	7,198	-
2004 2/	16,261	24	2	3	20	61	166	419	1,004	2,205	5,385	6,971	1
2003 2/	16,174	28	4	5	14	64	135	400	981	2,277	5,429	6,837	-
2002	15,786	28	2	4	16	55	162	389	933	2,176	5,423	6,598	-
2001	15,118	27	7	3	24	55	148	361	849	2,255	5,088	6,301	-
2000	14,357	20	2	5	23	48	153	338	786	2,167	4,885	5,930	-
1999	13,625	35	5	5	19	56	131	311	733	2,186	4,597	5,547	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)													
Non-Hispanic black, both sexes													
2005 2/	8,017	57	3	3	29	120	312	745	1,340	1,744	2,090	1,574	-
2004 2/	7,759	66	4	2	31	103	316	813	1,283	1,629	1,993	1,519	-
2003 2/	7,804	60	3	7	34	118	355	780	1,172	1,758	2,015	1,501	1
2002	7,410	55	4	5	36	108	306	745	1,072	1,670	1,941	1,468	-
2001	7,214	52	3	7	27	93	306	702	1,048	1,698	1,891	1,387	-
2000	6,853	69	5	4	24	99	305	664	980	1,637	1,776	1,290	-
1999	6,659	58	1	8	30	103	310	615	879	1,554	1,750	1,351	-
Non-Hispanic black, male													
2005 2/	3,612	36	3	1	11	60	164	438	746	750	891	512	-
2004 2/	3,505	40	3	2	19	51	166	474	640	771	827	512	-
2003 2/	3,614	32	1	2	20	64	191	467	627	853	862	494	1
2002	3,389	35	3	2	15	49	161	428	573	794	809	520	-
2001	3,158	28	3	-	12	54	139	399	509	782	784	448	-
2000	3,045	40	2	2	10	50	175	394	484	706	759	423	-
1999	2,984	25	1	3	14	47	173	353	426	691	803	448	-
Non-Hispanic black, female													
2005 2/	4,405	21	-	2	18	60	148	307	594	994	1,199	1,062	-
2004 2/	4,254	26	1	-	12	52	150	339	643	858	1,166	1,007	-
2003 2/	4,190	28	2	5	14	54	164	313	545	905	1,153	1,007	-
2002	4,021	20	1	3	21	59	145	317	499	876	1,132	948	-
2001	4,056	24	-	7	15	39	167	303	539	916	1,107	939	-
2000	3,808	29	3	2	14	49	130	270	496	931	1,017	867	-
1999	3,675	33	-	5	16	56	137	262	453	863	947	903	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)													
Origin not stated 3/, both sexes													
2005	62	1	-	1	-	1	2	7	6	10	14	20	-
2004	92	-	1	-	-	1	-	9	7	17	37	19	1
2003	111	3	-	-	-	-	3	6	16	13	34	36	-
2002	143	-	-	-	-	1	3	11	17	27	42	42	-
2001	112	2	-	-	-	2	3	13	11	20	35	25	1
2000	96	3	-	-	1	1	1	6	14	19	29	21	1
1999	111	1	-	-	-	-	2	5	13	21	27	42	-
Origin not stated 3/, male													
2005	37	1	-	1	-	1	2	5	4	6	7	10	-
2004	49	-	1	-	-	1	-	5	4	11	17	9	1
2003	54	-	-	-	-	-	1	4	10	9	16	14	-
2002	76	-	-	-	-	-	2	6	6	19	27	16	-
2001	52	2	-	-	-	2	2	7	5	11	14	9	-
2000	48	3	-	-	1	-	1	5	7	13	12	5	1
1999	59	1	-	-	-	-	1	5	9	14	11	18	-
Origin not stated 3/, female													
2005	25	-	-	-	-	-	-	2	2	4	7	10	-
2004	43	-	-	-	-	-	-	4	3	6	20	10	-
2003	57	3	-	-	-	-	2	2	6	4	18	22	-
2002	67	-	-	-	-	1	1	5	11	8	15	26	-
2001	60	-	-	-	-	-	1	6	6	9	21	16	1
2000	48	-	-	-	-	1	-	1	7	6	17	16	-
1999	52	-	-	-	-	-	1	-	4	7	16	24	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Acute and rapidly progressive nephritic and nephrotic syndrome (N00-N01,N04)													
All origins, both sexes													
2005	137	6	2	2	-	3	4	6	15	17	43	39	-
2004	157	7	5	1	2	1	2	16	9	27	49	38	-
2003	148	8	2	1	4	2	4	7	14	30	50	26	-
2002	166	6	5	2	3	1	4	9	16	27	51	42	-
2001	156	8	4	1	-	3	8	6	13	27	48	38	-
2000	177	8	3	2	1	4	3	16	10	29	52	49	-
1999	182	3	2	5	2	5	5	11	16	37	62	34	-
All origins, male													
2005	67	2	2	-	-	2	4	4	8	11	19	15	-
2004	73	2	4	-	2	1	1	10	4	14	24	11	-
2003	65	7	1	-	4	1	2	3	7	15	20	5	-
2002	70	3	2	2	1	1	-	5	11	14	17	14	-
2001	73	4	3	-	-	2	6	5	9	10	19	15	-
2000	91	3	2	2	1	3	1	11	8	12	26	22	-
1999	90	3	2	2	2	4	2	6	6	17	32	14	-
All origins, female													
2005	70	4	-	2	-	1	-	2	7	6	24	24	-
2004	84	5	1	1	-	-	1	6	5	13	25	27	-
2003	83	1	1	1	-	1	2	4	7	15	30	21	-
2002	96	3	3	-	2	-	4	4	5	13	34	28	-
2001	83	4	1	1	-	1	2	1	4	17	29	23	-
2000	86	5	1	-	-	1	2	5	2	17	26	27	-
1999	92	-	-	3	-	1	3	5	10	20	30	20	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Acute and rapidly progressive nephritic and nephrotic syndrome (N00-N01,N04)													
Hispanic, both sexes													
2005	11	-	2	1	-	-	-	-	1	1	2	4	-
2004	15	1	2	1	1	1	-	1	2	-	4	2	-
2003	13	1	-	1	3	-	-	-	1	3	2	2	-
2002	10	-	2	-	-	-	2	-	2	2	2	-	-
2001	13	1	-	-	-	1	1	2	2	3	1	2	-
2000	9	2	1	-	-	-	-	2	-	2	1	1	-
1999	12	1	-	2	-	-	-	1	2	4	1	1	-
Hispanic, male													
2005	8	-	2	-	-	-	-	-	1	1	1	3	-
2004	8	-	2	-	1	1	-	-	1	-	2	1	-
2003	7	1	-	-	3	-	-	-	-	-	2	1	-
2002	4	-	-	-	-	-	-	-	2	1	1	-	-
2001	7	-	-	-	-	-	1	2	2	2	-	-	-
2000	4	-	1	-	-	-	-	1	-	-	1	1	-
1999	4	1	-	1	-	-	-	-	1	-	-	1	-
Hispanic, female													
2005	3	-	-	1	-	-	-	-	-	-	1	1	-
2004	7	1	-	1	-	-	-	1	1	-	2	1	-
2003	6	-	-	1	-	-	-	-	1	3	-	1	-
2002	6	-	2	-	-	-	2	-	-	1	1	-	-
2001	6	1	-	-	-	1	-	-	-	1	1	2	-
2000	5	2	-	-	-	-	-	1	-	2	-	-	-
1999	8	-	-	1	-	-	-	1	1	4	1	-	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Acute and rapidly progressive nephritic and nephrotic syndrome (N00-N01,N04)													
Non-Hispanic 1/, both sexes													
2005	126	6	-	1	-	3	4	6	14	16	41	35	-
2004	142	6	3	-	1	-	2	15	7	27	45	36	-
2003	135	7	2	-	1	2	4	7	13	27	48	24	-
2002	155	6	3	2	3	1	2	9	14	25	49	41	-
2001	142	7	4	1	-	2	7	4	11	24	46	36	-
2000	168	6	2	2	1	4	3	14	10	27	51	48	-
1999	170	2	2	3	2	5	5	10	14	33	61	33	-
Non-Hispanic 1/, male													
2005	59	2	-	-	-	2	4	4	7	10	18	12	-
2004	65	2	2	-	1	-	1	10	3	14	22	10	-
2003	58	6	1	-	1	1	2	3	7	15	18	4	-
2002	66	3	2	2	1	1	-	5	9	13	16	14	-
2001	65	4	3	-	-	2	5	3	7	8	18	15	-
2000	87	3	1	2	1	3	1	10	8	12	25	21	-
1999	86	2	2	1	2	4	2	6	5	17	32	13	-
Non-Hispanic 1/, female													
2005	67	4	-	1	-	1	-	2	7	6	23	23	-
2004	77	4	1	-	-	-	1	5	4	13	23	26	-
2003	77	1	1	-	-	1	2	4	6	12	30	20	-
2002	89	3	1	-	2	-	2	4	5	12	33	27	-
2001	77	3	1	1	-	-	2	1	4	16	28	21	-
2000	81	3	1	-	-	1	2	4	2	15	26	27	-
1999	84	-	-	2	-	1	3	4	9	16	29	20	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Acute and rapidly progressive nephritic and nephrotic syndrome (N00-N01,N04)													
Non-Hispanic white, both sexes													
2005 2/	103	4	-	-	-	-	3	4	14	14	36	28	-
2004 2/	113	2	2	-	-	-	-	12	6	22	39	30	-
2003 2/	107	5	1	-	-	1	3	7	8	23	40	19	-
2002	119	5	2	2	1	-	1	5	12	19	40	32	-
2001	116	6	3	-	-	-	7	3	9	18	41	29	-
2000	132	4	-	1	1	3	1	4	10	20	45	43	-
1999	132	2	2	2	1	2	3	6	7	28	51	28	-
Non-Hispanic white, male													
2005 2/	52	1	-	-	-	-	3	3	7	9	18	11	-
2004 2/	52	-	2	-	-	-	-	8	2	14	18	8	-
2003 2/	40	4	-	-	-	-	2	3	3	12	12	4	-
2002	52	2	1	2	1	-	-	4	8	10	13	11	-
2001	52	3	2	-	-	-	5	3	5	4	18	12	-
2000	68	3	-	1	1	2	1	4	8	9	22	17	-
1999	63	2	2	1	1	1	1	4	2	14	26	9	-
Non-Hispanic white, female													
2005 2/	51	3	-	-	-	-	-	1	7	5	18	17	-
2004 2/	61	2	-	-	-	-	-	4	4	8	21	22	-
2003 2/	67	1	1	-	-	1	1	4	5	11	28	15	-
2002	67	3	1	-	-	-	1	1	4	9	27	21	-
2001	64	3	1	-	-	-	2	-	4	14	23	17	-
2000	64	1	-	-	-	1	-	-	2	11	23	26	-
1999	69	-	-	1	-	1	2	2	5	14	25	19	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Acute and rapidly progressive nephritic and nephrotic syndrome (N00-N01,N04)													
Non-Hispanic black, both sexes													
2005 2/	18	1	-	1	-	3	1	2	-	2	4	4	-
2004 2/	24	2	1	-	1	-	2	3	1	5	5	4	-
2003 2/	18	1	-	-	-	1	1	-	3	4	6	2	-
2002	29	1	1	-	2	1	1	4	1	5	6	7	-
2001	21	-	1	1	-	2	-	1	1	6	5	4	-
2000	30	2	2	-	-	1	2	10	-	6	5	2	-
1999	28	-	-	1	1	2	2	4	5	3	6	4	-
Non-Hispanic black, male													
2005 2/	6	-	-	-	-	2	1	1	-	1	-	1	-
2004 2/	10	1	-	-	1	-	1	2	1	-	3	1	-
2003 2/	12	1	-	-	-	1	-	-	2	3	5	-	-
2002	11	1	1	-	-	1	-	1	1	2	2	2	-
2001	9	-	1	-	-	2	-	-	1	4	-	1	-
2000	13	-	1	-	-	1	-	6	-	2	2	1	-
1999	16	-	-	-	1	2	1	2	2	2	3	3	-
Non-Hispanic black, female													
2005 2/	12	1	-	1	-	1	-	1	-	1	4	3	-
2004 2/	14	1	1	-	-	-	1	1	-	5	2	3	-
2003 2/	6	-	-	-	-	-	1	-	1	1	1	2	-
2002	18	-	-	-	2	-	1	3	-	3	4	5	-
2001	12	-	-	1	-	-	-	1	-	2	5	3	-
2000	17	2	1	-	-	-	2	4	-	4	3	1	-
1999	12	-	-	1	-	-	1	2	3	1	3	1	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Acute and rapidly progressive nephritic and nephrotic syndrome (N00-N01,N04)													
Origin not stated 3/, both sexes													
2005	-	-	-	-	-	-	-	-	-	-	-	-	-
2004	-	-	-	-	-	-	-	-	-	-	-	-	-
2003	-	-	-	-	-	-	-	-	-	-	-	-	-
2002	1	-	-	-	-	-	-	-	-	-	-	1	-
2001	1	-	-	-	-	-	-	-	-	-	1	-	-
2000	-	-	-	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-	-	-	-
Origin not stated 3/, male													
2005	-	-	-	-	-	-	-	-	-	-	-	-	-
2004	-	-	-	-	-	-	-	-	-	-	-	-	-
2003	-	-	-	-	-	-	-	-	-	-	-	-	-
2002	-	-	-	-	-	-	-	-	-	-	-	-	-
2001	1	-	-	-	-	-	-	-	-	-	1	-	-
2000	-	-	-	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-	-	-	-
Origin not stated 3/, female													
2005	-	-	-	-	-	-	-	-	-	-	-	-	-
2004	-	-	-	-	-	-	-	-	-	-	-	-	-
2003	-	-	-	-	-	-	-	-	-	-	-	-	-
2002	1	-	-	-	-	-	-	-	-	-	-	1	-
2001	-	-	-	-	-	-	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-	-	-	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Chronic glomerulonephritis, nephritis and nephropathy not specified as acute or chronic, and renal sclerosis unspecified (N02-N03,N05-N07,N26)													
All origins, both sexes													
2005	867	2	1	2	11	21	30	48	93	132	252	275	-
2004	553	1	-	2	5	21	30	39	64	88	168	135	-
2003	537	3	-	-	8	14	20	45	50	102	165	130	-
2002	553	2	-	3	4	23	26	53	52	85	178	127	-
2001	511	-	2	1	8	12	27	41	40	111	147	122	-
2000	556	1	-	4	9	17	34	39	59	103	172	118	-
1999	589	-	1	1	7	15	28	44	60	101	200	132	-
All origins, male													
2005	444	-	1	-	4	15	23	32	53	67	129	120	-
2004	275	1	-	-	1	6	17	20	40	46	84	60	-
2003	269	1	-	-	5	5	14	35	22	48	80	59	-
2002	278	1	-	1	3	13	14	25	27	49	92	53	-
2001	264	-	1	-	5	1	13	24	25	63	84	48	-
2000	284	-	-	1	2	11	20	27	31	54	83	55	-
1999	304	-	-	1	6	10	18	25	32	45	107	60	-
All origins, female													
2005	423	2	-	2	7	6	7	16	40	65	123	155	-
2004	278	-	-	2	4	15	13	19	24	42	84	75	-
2003	268	2	-	-	3	9	6	10	28	54	85	71	-
2002	275	1	-	2	1	10	12	28	25	36	86	74	-
2001	247	-	1	1	3	11	14	17	15	48	63	74	-
2000	272	1	-	3	7	6	14	12	28	49	89	63	-
1999	285	-	1	-	1	5	10	19	28	56	93	72	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Chronic glomerulonephritis, nephritis and nephropathy not specified as acute or chronic, and renal sclerosis unspecified (N02-N03,N05-N07,N26)													
Hispanic, both sexes													
2005	63	-	-	2	4	6	6	4	6	10	15	10	-
2004	30	1	-	1	-	1	1	3	5	9	2	7	-
2003	35	1	-	-	1	2	2	5	7	5	9	3	-
2002	32	-	-	2	-	3	2	3	5	3	9	5	-
2001	34	-	1	-	1	1	6	4	4	6	6	5	-
2000	28	-	-	-	-	2	1	4	4	5	7	5	-
1999	33	-	1	-	1	3	2	2	6	8	5	5	-
Hispanic, male													
2005	27	-	-	-	1	2	3	4	4	5	3	5	-
2004	18	1	-	-	-	1	-	1	5	5	1	4	-
2003	17	-	-	-	-	1	1	5	3	3	4	-	-
2002	15	-	-	-	-	3	-	-	4	-	6	2	-
2001	17	-	-	-	1	-	5	-	3	2	4	2	-
2000	17	-	-	-	-	1	-	3	3	2	3	5	-
1999	18	-	-	-	1	3	2	1	4	4	1	2	-
Hispanic, female													
2005	36	-	-	2	3	4	3	-	2	5	12	5	-
2004	12	-	-	1	-	-	1	2	-	4	1	3	-
2003	18	1	-	-	1	1	1	-	4	2	5	3	-
2002	17	-	-	2	-	-	2	3	1	3	3	3	-
2001	17	-	1	-	-	1	1	4	1	4	2	3	-
2000	11	-	-	-	-	1	1	1	1	3	4	-	-
1999	15	-	1	-	-	-	-	1	2	4	4	3	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Chronic glomerulonephritis, nephritis and nephropathy not specified as acute or chronic, and renal sclerosis unspecified (N02-N03,N05-N07,N26)													
Non-Hispanic 1/, both sexes													
2005	804	2	1	-	7	15	24	44	87	122	237	265	-
2004	522	-	-	1	5	20	29	36	59	79	166	127	-
2003	501	2	-	-	7	12	17	40	43	97	156	127	-
2002	520	2	-	1	4	19	24	50	47	82	169	122	-
2001	477	-	1	1	7	11	21	37	36	105	141	117	-
2000	528	1	-	4	9	15	33	35	55	98	165	113	-
1999	555	-	-	1	6	12	26	42	53	93	195	127	-
Non-Hispanic 1/, male													
2005	417	-	1	-	3	13	20	28	49	62	126	115	-
2004	256	-	-	-	1	5	17	19	35	41	83	55	-
2003	251	1	-	-	5	4	12	30	19	45	76	59	-
2002	263	1	-	1	3	10	14	25	23	49	86	51	-
2001	247	-	1	-	4	1	8	24	22	61	80	46	-
2000	267	-	-	1	2	10	20	24	28	52	80	50	-
1999	285	-	-	1	5	7	16	24	27	41	106	58	-
Non-Hispanic 1/, female													
2005	387	2	-	-	4	2	4	16	38	60	111	150	-
2004	266	-	-	1	4	15	12	17	24	38	83	72	-
2003	250	1	-	-	2	8	5	10	24	52	80	68	-
2002	257	1	-	-	1	9	10	25	24	33	83	71	-
2001	230	-	-	1	3	10	13	13	14	44	61	71	-
2000	261	1	-	3	7	5	13	11	27	46	85	63	-
1999	270	-	-	-	1	5	10	18	26	52	89	69	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Chronic glomerulonephritis, nephritis and nephropathy not specified as acute or chronic, and renal sclerosis unspecified (N02-N03,N05-N07,N26)													
Non-Hispanic white, both sexes													
2005 2/	623	1	-	-	4	7	12	24	59	88	200	228	-
2004 2/	416	-	-	1	3	12	19	23	42	63	142	111	-
2003 2/	394	2	-	-	2	6	13	26	35	74	128	108	-
2002	419	-	-	-	3	12	20	31	34	76	141	102	-
2001	384	-	1	-	6	8	9	17	24	87	123	109	-
2000	411	-	-	3	7	9	19	22	39	80	143	89	-
1999	436	-	-	1	3	8	13	30	36	72	169	104	-
Non-Hispanic white, male													
2005 2/	323	-	-	-	3	7	10	16	33	49	105	100	-
2004 2/	202	-	-	-	-	4	12	14	23	31	72	46	-
2003 2/	196	1	-	-	1	2	10	20	16	35	61	50	-
2002	218	-	-	-	2	7	11	16	19	45	72	46	-
2001	200	-	1	-	3	1	3	10	14	52	71	45	-
2000	209	-	-	1	1	6	12	15	20	43	72	39	-
1999	227	-	-	1	2	5	8	18	19	33	93	48	-
Non-Hispanic white, female													
2005 2/	300	1	-	-	1	-	2	8	26	39	95	128	-
2004 2/	214	-	-	1	3	8	7	9	19	32	70	65	-
2003 2/	198	1	-	-	1	4	3	6	19	39	67	58	-
2002	201	-	-	-	1	5	9	15	15	31	69	56	-
2001	184	-	-	-	3	7	6	7	10	35	52	64	-
2000	202	-	-	2	6	3	7	7	19	37	71	50	-
1999	209	-	-	-	1	3	5	12	17	39	76	56	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Chronic glomerulonephritis, nephritis and nephropathy not specified as acute or chronic, and renal sclerosis unspecified (N02-N03,N05-N07,N26)													
Non-Hispanic black, both sexes													
2005 2/	161	1	1	-	3	6	10	17	27	28	34	34	-
2004 2/	90	-	-	-	2	8	10	10	15	15	18	12	-
2003 2/	76	-	-	-	4	4	4	12	6	15	19	12	-
2002	80	2	-	-	1	6	3	18	8	5	20	17	-
2001	68	-	-	1	-	2	8	15	8	14	13	7	-
2000	90	1	-	1	2	4	10	12	12	14	19	15	-
1999	92	-	-	-	3	2	11	10	11	17	19	19	-
Non-Hispanic black, male													
2005 2/	81	-	1	-	-	4	8	9	16	9	21	13	-
2004 2/	42	-	-	-	1	1	5	4	10	9	7	5	-
2003 2/	39	-	-	-	3	1	2	9	2	7	10	5	-
2002	37	1	-	-	1	3	2	8	3	4	11	4	-
2001	36	-	-	-	-	-	5	12	6	7	5	1	-
2000	44	-	-	-	1	2	5	8	7	8	6	7	-
1999	46	-	-	-	3	1	6	6	6	6	10	8	-
Non-Hispanic black, female													
2005 2/	80	1	-	-	3	2	2	8	11	19	13	21	-
2004 2/	48	-	-	-	1	7	5	6	5	6	11	7	-
2003 2/	37	-	-	-	1	3	2	3	4	8	9	7	-
2002	43	1	-	-	-	3	1	10	5	1	9	13	-
2001	32	-	-	1	-	2	3	3	2	7	8	6	-
2000	46	1	-	1	1	2	5	4	5	6	13	8	-
1999	46	-	-	-	-	1	5	4	5	11	9	11	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Chronic glomerulonephritis, nephritis and nephropathy not specified as acute or chronic, and renal sclerosis unspecified (N02-N03,N05-N07,N26)													
Origin not stated 3/, both sexes													
2005	-	-	-	-	-	-	-	-	-	-	-	-	-
2004	1	-	-	-	-	-	-	-	-	-	-	1	-
2003	1	-	-	-	-	-	1	-	-	-	-	-	-
2002	1	-	-	-	-	1	-	-	-	-	-	-	-
2001	-	-	-	-	-	-	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-	-	-	-
1999	1	-	-	-	-	-	-	-	1	-	-	-	-
Origin not stated 3/, male													
2005	-	-	-	-	-	-	-	-	-	-	-	-	-
2004	1	-	-	-	-	-	-	-	-	-	-	1	-
2003	1	-	-	-	-	-	1	-	-	-	-	-	-
2002	-	-	-	-	-	-	-	-	-	-	-	-	-
2001	-	-	-	-	-	-	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-	-	-	-
1999	1	-	-	-	-	-	-	-	1	-	-	-	-
Origin not stated 3/, female													
2005	-	-	-	-	-	-	-	-	-	-	-	-	-
2004	-	-	-	-	-	-	-	-	-	-	-	-	-
2003	-	-	-	-	-	-	-	-	-	-	-	-	-
2002	1	-	-	-	-	1	-	-	-	-	-	-	-
2001	-	-	-	-	-	-	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-	-	-	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Renal failure (N17-N19)													
All origins, both sexes													
2005	42,868	151	13	19	79	261	705	1,973	4,025	7,165	14,102	14,375	-
2004	41,732	165	9	18	79	223	767	2,009	3,883	6,997	13,830	13,750	2
2003	41,737	170	11	21	82	266	771	1,944	3,741	7,206	13,870	13,653	2
2002	40,222	165	9	15	77	243	718	1,829	3,383	7,046	13,657	13,080	-
2001	38,784	125	9	16	82	209	732	1,770	3,226	7,211	12,898	12,505	1
2000	36,471	152	11	19	78	221	701	1,590	3,023	6,848	12,223	11,603	2
1999	34,719	163	12	25	63	230	673	1,422	2,784	6,697	11,652	10,998	-
All origins, male													
2005	20,745	86	9	10	41	136	395	1,201	2,288	3,650	7,027	5,902	-
2004	20,006	106	6	11	46	105	433	1,175	2,072	3,616	6,895	5,540	1
2003	20,135	98	5	9	49	141	443	1,151	2,036	3,719	6,947	5,535	2
2002	19,341	105	5	8	37	114	393	1,055	1,786	3,739	6,808	5,291	-
2001	18,509	72	3	6	40	109	394	1,022	1,680	3,752	6,381	5,050	-
2000	17,416	100	5	14	40	115	407	941	1,607	3,492	6,102	4,591	2
1999	16,611	87	7	15	25	110	382	814	1,492	3,440	5,888	4,351	-
All origins, female													
2005	22,123	65	4	9	38	125	310	772	1,737	3,515	7,075	8,473	-
2004	21,726	59	3	7	33	118	334	834	1,811	3,381	6,935	8,210	1
2003	21,602	72	6	12	33	125	328	793	1,705	3,487	6,923	8,118	-
2002	20,881	60	4	7	40	129	325	774	1,597	3,307	6,849	7,789	-
2001	20,275	53	6	10	42	100	338	748	1,546	3,459	6,517	7,455	1
2000	19,055	52	6	5	38	106	294	649	1,416	3,356	6,121	7,012	-
1999	18,108	76	5	10	38	120	291	608	1,292	3,257	5,764	6,647	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Renal failure (N17-N19)													
Hispanic, both sexes													
2005	2,254	34	2	2	11	24	62	160	344	495	682	438	-
2004	2,149	35	1	9	10	20	71	178	311	514	623	377	-
2003	2,121	33	1	3	15	28	74	168	310	476	567	445	1
2002	1,856	35	3	3	9	30	61	154	260	422	521	358	-
2001	1,806	17	1	4	15	23	66	125	230	460	517	348	-
2000	1,605	22	3	3	14	24	51	103	227	397	420	340	1
1999	1,535	19	2	7	5	18	60	104	222	356	417	325	-
Hispanic, male													
2005	1,151	18	1	1	7	14	34	108	196	238	334	200	-
2004	1,073	24	-	4	6	6	45	106	173	246	303	160	-
2003	1,044	18	-	1	10	15	48	101	157	227	274	192	1
2002	918	21	1	3	5	9	38	88	135	213	266	139	-
2001	867	13	-	2	8	10	39	62	127	213	247	146	-
2000	782	16	2	2	7	10	35	65	111	192	202	139	1
1999	790	14	2	6	1	7	29	63	132	174	209	153	-
Hispanic, female													
2005	1,103	16	1	1	4	10	28	52	148	257	348	238	-
2004	1,076	11	1	5	4	14	26	72	138	268	320	217	-
2003	1,077	15	1	2	5	13	26	67	153	249	293	253	-
2002	938	14	2	-	4	21	23	66	125	209	255	219	-
2001	939	4	1	2	7	13	27	63	103	247	270	202	-
2000	823	6	1	1	7	14	16	38	116	205	218	201	-
1999	745	5	-	1	4	11	31	41	90	182	208	172	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Renal failure (N17-N19)													
Non-Hispanic 1/, both sexes													
2005	40,552	116	11	16	68	236	641	1,806	3,675	6,660	13,406	13,917	-
2004	39,492	130	7	9	69	202	696	1,822	3,565	6,466	13,170	13,355	1
2003	39,506	134	10	18	67	238	695	1,770	3,415	6,717	13,269	13,172	1
2002	38,225	130	6	12	68	213	654	1,664	3,106	6,597	13,094	12,681	-
2001	36,867	106	8	12	67	184	663	1,632	2,985	6,731	12,347	12,132	-
2000	34,771	127	8	16	63	196	649	1,481	2,783	6,432	11,774	11,242	-
1999	33,074	143	10	18	58	212	611	1,313	2,550	6,320	11,208	10,631	-
Non-Hispanic 1/, male													
2005	19,557	67	8	8	34	121	359	1,088	2,088	3,406	6,686	5,692	-
2004	18,885	82	5	7	40	98	388	1,064	1,895	3,359	6,575	5,372	-
2003	19,038	80	5	8	39	126	395	1,046	1,869	3,483	6,657	5,329	1
2002	18,347	84	4	5	32	105	353	961	1,645	3,507	6,515	5,136	-
2001	17,591	57	3	4	32	97	353	953	1,548	3,528	6,121	4,895	-
2000	16,586	81	3	12	32	105	371	871	1,489	3,287	5,888	4,447	-
1999	15,763	72	5	9	24	103	352	746	1,352	3,252	5,668	4,180	-
Non-Hispanic 1/, female													
2005	20,995	49	3	8	34	115	282	718	1,587	3,254	6,720	8,225	-
2004	20,607	48	2	2	29	104	308	758	1,670	3,107	6,595	7,983	1
2003	20,468	54	5	10	28	112	300	724	1,546	3,234	6,612	7,843	-
2002	19,878	46	2	7	36	108	301	703	1,461	3,090	6,579	7,545	-
2001	19,276	49	5	8	35	87	310	679	1,437	3,203	6,226	7,237	-
2000	18,185	46	5	4	31	91	278	610	1,294	3,145	5,886	6,795	-
1999	17,311	71	5	9	34	109	259	567	1,198	3,068	5,540	6,451	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Renal failure (N17-N19)													
Non-Hispanic white, both sexes													
2005 2/	31,743	57	9	13	40	116	316	1,031	2,236	4,739	11,057	12,129	-
2004 2/	30,948	58	4	6	38	95	374	969	2,184	4,653	10,916	11,650	1
2003 2/	30,973	73	7	11	36	121	323	941	2,153	4,791	11,004	11,513	-
2002	30,104	72	3	8	29	105	331	885	1,928	4,769	10,922	11,052	-
2001	28,932	52	6	6	38	91	343	881	1,825	4,873	10,233	10,584	-
2000	27,350	57	4	13	40	97	339	800	1,722	4,632	9,830	9,816	-
1999	25,831	79	8	10	32	104	302	679	1,592	4,614	9,279	9,132	-
Non-Hispanic white, male													
2005 2/	15,566	31	6	6	21	60	187	637	1,302	2,562	5,677	5,077	-
2004 2/	14,981	36	2	4	21	42	216	563	1,206	2,494	5,627	4,770	-
2003 2/	15,080	47	4	6	23	62	193	552	1,197	2,566	5,675	4,755	-
2002	14,610	47	2	4	14	56	180	513	1,018	2,639	5,600	4,537	-
2001	14,076	28	1	3	18	44	203	528	993	2,672	5,221	4,365	-
2000	13,275	38	2	10	23	54	193	471	959	2,516	5,045	3,964	-
1999	12,496	44	3	6	14	53	179	383	882	2,484	4,787	3,661	-
Non-Hispanic white, female													
2005 2/	16,177	26	3	7	19	56	129	394	934	2,177	5,380	7,052	-
2004 2/	15,967	22	2	2	17	53	158	406	978	2,159	5,289	6,880	1
2003 2/	15,893	26	3	5	13	59	130	389	956	2,225	5,329	6,758	-
2002	15,494	25	1	4	15	49	151	372	910	2,130	5,322	6,515	-
2001	14,856	24	5	3	20	47	140	353	832	2,201	5,012	6,219	-
2000	14,075	19	2	3	17	43	146	329	763	2,116	4,785	5,852	-
1999	13,335	35	5	4	18	51	123	296	710	2,130	4,492	5,471	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Renal failure (N17-N19)													
Non-Hispanic black, both sexes													
2005 2/	7,834	55	2	2	26	111	301	726	1,313	1,713	2,051	1,534	-
2004 2/	7,640	64	3	2	28	95	303	798	1,267	1,609	1,968	1,503	-
2003 2/	7,704	59	3	7	29	113	350	767	1,163	1,736	1,989	1,487	1
2002	7,297	52	3	4	33	101	302	722	1,063	1,660	1,914	1,443	-
2001	7,119	52	2	5	26	89	298	686	1,038	1,678	1,871	1,374	-
2000	6,724	66	3	3	22	93	292	639	968	1,616	1,751	1,271	-
1999	6,527	58	1	7	26	99	297	600	860	1,533	1,720	1,326	-
Non-Hispanic black, male													
2005 2/	3,525	36	2	1	11	54	155	428	730	740	870	498	-
2004 2/	3,451	39	3	2	17	50	160	466	629	762	817	506	-
2003 2/	3,559	31	1	2	16	62	189	457	623	841	847	489	1
2002	3,339	33	2	1	14	45	159	418	569	788	796	514	-
2001	3,111	28	2	-	12	52	134	387	502	771	778	445	-
2000	2,986	40	1	2	9	47	170	378	477	696	751	415	-
1999	2,920	25	1	3	10	44	166	345	418	682	789	437	-
Non-Hispanic black, female													
2005 2/	4,309	19	-	1	15	57	146	298	583	973	1,181	1,036	-
2004 2/	4,189	25	-	-	11	45	143	332	638	847	1,151	997	-
2003 2/	4,145	28	2	5	13	51	161	310	540	895	1,142	998	-
2002	3,958	19	1	3	19	56	143	304	494	872	1,118	929	-
2001	4,008	24	-	5	14	37	164	299	536	907	1,093	929	-
2000	3,738	26	2	1	13	46	122	261	491	920	1,000	856	-
1999	3,607	33	-	4	16	55	131	255	442	851	931	889	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Renal failure (N17-N19)													
Origin not stated 3/, both sexes													
2005	62	1	-	1	-	1	2	7	6	10	14	20	-
2004	91	-	1	-	-	1	-	9	7	17	37	18	1
2003	110	3	-	-	-	-	2	6	16	13	34	36	-
2002	141	-	-	-	-	-	3	11	17	27	42	41	-
2001	111	2	-	-	-	2	3	13	11	20	34	25	1
2000	95	3	-	-	1	1	1	6	13	19	29	21	1
1999	110	1	-	-	-	-	2	5	12	21	27	42	-
Origin not stated 3/, male													
2005	37	1	-	1	-	1	2	5	4	6	7	10	-
2004	48	-	1	-	-	1	-	5	4	11	17	8	1
2003	53	-	-	-	-	-	-	4	10	9	16	14	-
2002	76	-	-	-	-	-	2	6	6	19	27	16	-
2001	51	2	-	-	-	2	2	7	5	11	13	9	-
2000	48	3	-	-	1	-	1	5	7	13	12	5	1
1999	58	1	-	-	-	-	1	5	8	14	11	18	-
Origin not stated 3/, female													
2005	25	-	-	-	-	-	-	2	2	4	7	10	-
2004	43	-	-	-	-	-	-	4	3	6	20	10	-
2003	57	3	-	-	-	-	2	2	6	4	18	22	-
2002	65	-	-	-	-	-	1	5	11	8	15	25	-
2001	60	-	-	-	-	-	1	6	6	9	21	16	1
2000	47	-	-	-	-	1	-	1	6	6	17	16	-
1999	52	-	-	-	-	-	1	-	4	7	16	24	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Other disorders of kidney (N25,N27)													
All origins, both sexes													
2005	29	-	-	-	1	-	3	1	8	6	6	4	-
2004	38	1	-	-	1	1	2	3	7	7	11	5	-
2003	31	-	1	-	1	-	1	5	1	7	8	7	-
2002	33	-	-	1	-	2	1	2	4	6	10	7	-
2001	29	-	1	-	2	1	1	5	5	7	4	3	-
2000	47	1	-	-	-	2	2	6	8	10	11	7	-
1999	35	-	-	-	-	2	1	3	4	6	13	6	-
All origins, male													
2005	12	-	-	-	-	-	1	-	3	5	2	1	-
2004	16	1	-	-	1	1	-	3	4	1	4	1	-
2003	12	-	1	-	1	-	-	4	-	3	2	1	-
2002	6	-	-	1	-	1	-	1	-	-	3	-	-
2001	6	-	-	-	-	-	1	2	-	1	1	1	-
2000	20	-	-	-	-	-	1	3	5	6	2	3	-
1999	11	-	-	-	-	1	-	1	-	2	4	3	-
All origins, female													
2005	17	-	-	-	1	-	2	1	5	1	4	3	-
2004	22	-	-	-	-	-	2	-	3	6	7	4	-
2003	19	-	-	-	-	-	1	1	1	4	6	6	-
2002	27	-	-	-	-	1	1	1	4	6	7	7	-
2001	23	-	1	-	2	1	-	3	5	6	3	2	-
2000	27	1	-	-	-	2	1	3	3	4	9	4	-
1999	24	-	-	-	-	1	1	2	4	4	9	3	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Other disorders of kidney (N25,N27)													
Hispanic, both sexes													
2005	1	-	-	-	-	-	-	1	-	-	-	-	-
2004	2	-	-	-	1	-	-	-	1	-	-	-	-
2003	1	-	-	-	-	-	-	1	-	-	-	-	-
2002	-	-	-	-	-	-	-	-	-	-	-	-	-
2001	4	-	-	-	-	-	1	1	1	1	-	-	-
2000	4	1	-	-	-	-	1	-	1	-	1	-	-
1999	4	-	-	-	-	-	-	-	-	1	3	-	-
Hispanic, male													
2005	-	-	-	-	-	-	-	-	-	-	-	-	-
2004	2	-	-	-	1	-	-	-	1	-	-	-	-
2003	1	-	-	-	-	-	-	1	-	-	-	-	-
2002	-	-	-	-	-	-	-	-	-	-	-	-	-
2001	1	-	-	-	-	-	1	-	-	-	-	-	-
2000	2	-	-	-	-	-	1	-	1	-	-	-	-
1999	2	-	-	-	-	-	-	-	-	-	2	-	-
Hispanic, female													
2005	1	-	-	-	-	-	-	1	-	-	-	-	-
2004	-	-	-	-	-	-	-	-	-	-	-	-	-
2003	-	-	-	-	-	-	-	-	-	-	-	-	-
2002	-	-	-	-	-	-	-	-	-	-	-	-	-
2001	3	-	-	-	-	-	-	1	1	1	-	-	-
2000	2	1	-	-	-	-	-	-	-	-	1	-	-
1999	2	-	-	-	-	-	-	-	-	1	1	-	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Other disorders of kidney (N25,N27)													
Non-Hispanic 1/, both sexes													
2005	28	-	-	-	1	-	3	-	8	6	6	4	-
2004	36	1	-	-	-	1	2	3	6	7	11	5	-
2003	30	-	1	-	1	-	1	4	1	7	8	7	-
2002	33	-	-	1	-	2	1	2	4	6	10	7	-
2001	25	-	1	-	2	1	-	4	4	6	4	3	-
2000	42	-	-	-	-	2	1	6	6	10	10	7	-
1999	31	-	-	-	-	2	1	3	4	5	10	6	-
Non-Hispanic 1/, male													
2005	12	-	-	-	-	-	1	-	3	5	2	1	-
2004	14	1	-	-	-	1	-	3	3	1	4	1	-
2003	11	-	1	-	1	-	-	3	-	3	2	1	-
2002	6	-	-	1	-	1	-	1	-	-	3	-	-
2001	5	-	-	-	-	-	-	2	-	1	1	1	-
2000	18	-	-	-	-	-	-	3	4	6	2	3	-
1999	9	-	-	-	-	1	-	1	-	2	2	3	-
Non-Hispanic 1/, female													
2005	16	-	-	-	1	-	2	-	5	1	4	3	-
2004	22	-	-	-	-	-	2	-	3	6	7	4	-
2003	19	-	-	-	-	-	1	1	1	4	6	6	-
2002	27	-	-	-	-	1	1	1	4	6	7	7	-
2001	20	-	1	-	2	1	-	2	4	5	3	2	-
2000	24	-	-	-	-	2	1	3	2	4	8	4	-
1999	22	-	-	-	-	1	1	2	4	3	8	3	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Other disorders of kidney (N25,N27)													
Non-Hispanic white, both sexes													
2005 2/	23	-	-	-	1	-	2	-	8	5	5	2	-
2004 2/	31	1	-	-	-	1	1	1	6	7	9	5	-
2003 2/	23	-	1	-	-	-	1	3	1	3	7	7	-
2002	27	-	-	-	-	2	1	1	4	6	7	6	-
2001	17	-	1	-	1	1	-	3	3	6	1	1	-
2000	31	-	-	-	-	1	-	3	5	9	8	5	-
1999	19	-	-	-	-	2	1	2	1	4	5	4	-
Non-Hispanic white, male													
2005 2/	12	-	-	-	-	-	1	-	3	5	2	1	-
2004 2/	12	1	-	-	-	1	-	1	3	1	4	1	-
2003 2/	7	-	1	-	-	-	-	2	-	1	2	1	-
2002	3	-	-	-	-	1	-	-	-	-	2	-	-
2001	3	-	-	-	-	-	-	2	-	1	-	-	-
2000	15	-	-	-	-	-	-	1	3	6	2	3	-
1999	7	-	-	-	-	1	-	1	-	1	1	3	-
Non-Hispanic white, female													
2005 2/	11	-	-	-	1	-	1	-	5	-	3	1	-
2004 2/	19	-	-	-	-	-	1	-	3	6	5	4	-
2003 2/	16	-	-	-	-	-	1	1	1	2	5	6	-
2002	24	-	-	-	-	1	1	1	4	6	5	6	-
2001	14	-	1	-	1	1	-	1	3	5	1	1	-
2000	16	-	-	-	-	1	-	2	2	3	6	2	-
1999	12	-	-	-	-	1	1	1	1	3	4	1	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Other disorders of kidney (N25,N27)													
Non-Hispanic black, both sexes													
2005 2/	4	-	-	-	-	-	-	-	-	1	1	2	-
2004 2/	5	-	-	-	-	-	1	2	-	-	2	-	-
2003 2/	6	-	-	-	1	-	-	1	-	3	1	-	-
2002	4	-	-	1	-	-	-	1	-	-	1	1	-
2001	6	-	-	-	1	-	-	-	1	-	2	2	-
2000	9	-	-	-	-	1	1	3	-	1	1	2	-
1999	12	-	-	-	-	-	-	1	3	1	5	2	-
Non-Hispanic black, male													
2005 2/	-	-	-	-	-	-	-	-	-	-	-	-	-
2004 2/	2	-	-	-	-	-	-	2	-	-	-	-	-
2003 2/	4	-	-	-	1	-	-	1	-	2	-	-	-
2002	2	-	-	1	-	-	-	1	-	-	-	-	-
2001	2	-	-	-	-	-	-	-	-	-	1	1	-
2000	2	-	-	-	-	-	-	2	-	-	-	-	-
1999	2	-	-	-	-	-	-	-	-	1	1	-	-
Non-Hispanic black, female													
2005 2/	4	-	-	-	-	-	-	-	-	1	1	2	-
2004 2/	3	-	-	-	-	-	1	-	-	-	2	-	-
2003 2/	2	-	-	-	-	-	-	-	-	1	1	-	-
2002	2	-	-	-	-	-	-	-	-	-	1	1	-
2001	4	-	-	-	1	-	-	-	1	-	1	1	-
2000	7	-	-	-	-	1	1	1	-	1	1	2	-
1999	10	-	-	-	-	-	-	1	3	-	4	2	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Other disorders of kidney (N25,N27)													
Origin not stated 3/, both sexes													
2005	-	-	-	-	-	-	-	-	-	-	-	-	-
2004	-	-	-	-	-	-	-	-	-	-	-	-	-
2003	-	-	-	-	-	-	-	-	-	-	-	-	-
2002	-	-	-	-	-	-	-	-	-	-	-	-	-
2001	-	-	-	-	-	-	-	-	-	-	-	-	-
2000	1	-	-	-	-	-	-	-	1	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-	-	-	-
Origin not stated 3/, male													
2005	-	-	-	-	-	-	-	-	-	-	-	-	-
2004	-	-	-	-	-	-	-	-	-	-	-	-	-
2003	-	-	-	-	-	-	-	-	-	-	-	-	-
2002	-	-	-	-	-	-	-	-	-	-	-	-	-
2001	-	-	-	-	-	-	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-	-	-	-
Origin not stated 3/, female													
2005	-	-	-	-	-	-	-	-	-	-	-	-	-
2004	-	-	-	-	-	-	-	-	-	-	-	-	-
2003	-	-	-	-	-	-	-	-	-	-	-	-	-
2002	-	-	-	-	-	-	-	-	-	-	-	-	-
2001	-	-	-	-	-	-	-	-	-	-	-	-	-
2000	1	-	-	-	-	-	-	-	1	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-	-	-	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Infections of kidney (N10-N12,N13.6,N15.1)													
All origins, both sexes													
2005	767	6	4	2	9	14	26	51	53	100	239	263	-
2004	764	3	4	-	4	15	34	47	65	104	237	251	-
2003	823	4	2	1	10	16	36	62	62	105	247	278	-
2002	788	2	-	3	5	13	28	40	59	110	238	290	-
2001	748	2	4	-	5	11	33	46	46	87	249	265	-
2000	813	5	3	-	2	9	21	42	59	104	248	320	-
1999	846	3	1	1	4	9	28	55	59	133	242	311	-
All origins, male													
2005	236	3	1	-	2	5	7	18	22	37	83	58	-
2004	237	2	1	-	1	3	7	12	32	34	77	68	-
2003	246	3	-	1	2	1	10	25	21	31	74	78	-
2002	251	2	-	1	1	6	12	15	21	47	72	74	-
2001	213	1	2	-	2	3	10	11	12	40	76	56	-
2000	214	3	-	-	1	3	8	15	15	26	74	69	-
1999	263	1	-	-	1	-	7	19	27	44	83	81	-
All origins, female													
2005	531	3	3	2	7	9	19	33	31	63	156	205	-
2004	527	1	3	-	3	12	27	35	33	70	160	183	-
2003	577	1	2	-	8	15	26	37	41	74	173	200	-
2002	537	-	-	2	4	7	16	25	38	63	166	216	-
2001	535	1	2	-	3	8	23	35	34	47	173	209	-
2000	599	2	3	-	1	6	13	27	44	78	174	251	-
1999	583	2	1	1	3	9	21	36	32	89	159	230	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Infections of kidney (N10-N12,N13.6,N15.1)													
Hispanic, both sexes													
2005	50	2	1	-	-	3	2	3	6	6	15	12	-
2004	52	-	-	-	1	2	5	5	8	10	15	6	-
2003	46	-	-	-	1	-	2	2	8	11	13	9	-
2002	47	1	-	2	-	3	2	3	2	12	14	8	-
2001	33	-	-	-	-	1	3	5	1	8	9	6	-
2000	39	1	-	-	-	-	2	4	4	7	10	11	-
1999	55	-	-	-	-	1	7	6	9	8	15	9	-
Hispanic, male													
2005	20	1	-	-	-	1	-	-	3	3	8	4	-
2004	17	-	-	-	1	-	1	1	4	2	6	2	-
2003	11	-	-	-	-	-	-	1	1	1	4	4	-
2002	19	1	-	-	-	2	1	2	1	4	5	3	-
2001	10	-	-	-	-	-	-	3	-	3	1	3	-
2000	11	1	-	-	-	-	-	3	1	3	2	1	-
1999	21	-	-	-	-	-	2	2	5	4	6	2	-
Hispanic, female													
2005	30	1	1	-	-	2	2	3	3	3	7	8	-
2004	35	-	-	-	-	2	4	4	4	8	9	4	-
2003	35	-	-	-	1	-	2	1	7	10	9	5	-
2002	28	-	-	2	-	1	1	1	1	8	9	5	-
2001	23	-	-	-	-	1	3	2	1	5	8	3	-
2000	28	-	-	-	-	-	2	1	3	4	8	10	-
1999	34	-	-	-	-	1	5	4	4	4	9	7	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Infections of kidney (N10-N12,N13.6,N15.1)													
Non-Hispanic 1/, both sexes													
2005	716	4	3	2	9	11	24	48	47	93	224	251	-
2004	711	3	4	-	3	12	29	42	57	94	222	245	-
2003	774	4	2	1	9	16	33	59	54	94	233	269	-
2002	740	1	-	1	5	10	26	37	57	98	224	281	-
2001	714	2	4	-	5	10	30	41	45	79	239	259	-
2000	771	4	3	-	2	9	19	38	55	96	236	309	-
1999	786	3	1	1	4	8	20	49	49	125	225	301	-
Non-Hispanic 1/, male													
2005	215	2	1	-	2	4	7	18	19	33	75	54	-
2004	220	2	1	-	-	3	6	11	28	32	71	66	-
2003	233	3	-	1	2	1	9	23	20	30	70	74	-
2002	232	1	-	1	1	4	11	13	20	43	67	71	-
2001	203	1	2	-	2	3	10	8	12	37	75	53	-
2000	201	2	-	-	1	3	8	12	14	22	71	68	-
1999	240	1	-	-	1	-	4	17	22	40	76	79	-
Non-Hispanic 1/, female													
2005	501	2	2	2	7	7	17	30	28	60	149	197	-
2004	491	1	3	-	3	9	23	31	29	62	151	179	-
2003	541	1	2	-	7	15	24	36	34	64	163	195	-
2002	508	-	-	-	4	6	15	24	37	55	157	210	-
2001	511	1	2	-	3	7	20	33	33	42	164	206	-
2000	570	2	3	-	1	6	11	26	41	74	165	241	-
1999	546	2	1	1	3	8	16	32	27	85	149	222	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Infections of kidney (N10-N12,N13.6,N15.1)													
Non-Hispanic white, both sexes													
2005 2/.....	610	3	3	2	7	6	14	36	36	77	200	226	-
2004 2/.....	606	2	3	-	3	8	24	29	41	82	191	223	-
2003 2/.....	656	1	1	1	7	11	25	44	38	82	201	245	-
2002.....	645	1	-	-	1	7	20	23	44	89	203	257	-
2001.....	614	1	2	-	2	7	23	29	36	65	210	239	-
2000.....	647	4	3	-	1	7	10	26	44	74	198	280	-
1999.....	689	1	1	1	2	7	12	38	38	102	206	281	-
Non-Hispanic white, male													
2005 2/.....	184	1	1	-	2	3	5	14	14	28	69	47	-
2004 2/.....	187	1	-	-	-	1	5	8	21	25	64	62	-
2003 2/.....	202	-	-	1	2	1	8	19	14	27	58	72	-
2002.....	200	1	-	-	1	4	9	10	16	36	58	65	-
2001.....	166	-	1	-	1	2	9	6	9	27	64	47	-
2000.....	169	2	-	-	-	3	4	9	12	13	62	64	-
1999.....	209	1	-	-	1	-	2	12	17	30	70	76	-
Non-Hispanic white, female													
2005 2/.....	426	2	2	2	5	3	9	22	22	49	131	179	-
2004 2/.....	419	1	3	-	3	7	19	21	20	57	127	161	-
2003 2/.....	454	1	1	-	5	10	17	25	24	55	143	173	-
2002.....	445	-	-	-	-	3	11	13	28	53	145	192	-
2001.....	448	1	1	-	1	5	14	23	27	38	146	192	-
2000.....	478	2	3	-	1	4	6	17	32	61	136	216	-
1999.....	480	-	1	1	1	7	10	26	21	72	136	205	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Infections of kidney (N10-N12,N13.6,N15.1)													
Non-Hispanic black, both sexes													
2005 2/	89	1	-	-	2	4	10	10	10	14	19	19	-
2004 2/	83	1	1	-	-	4	4	10	12	9	25	17	-
2003 2/	95	3	1	-	2	2	7	12	14	10	24	20	-
2002	73	-	-	-	3	2	6	8	10	8	15	21	-
2001	80	1	-	-	1	3	7	10	7	13	22	16	-
2000	101	-	-	-	1	2	7	11	8	19	28	25	-
1999	85	1	-	-	2	1	6	10	11	20	16	18	-
Non-Hispanic black, male													
2005 2/	27	1	-	-	-	1	2	3	5	5	5	5	-
2004 2/	27	1	1	-	-	2	1	2	6	6	5	3	-
2003 2/	29	3	-	-	-	-	1	4	5	3	11	2	-
2002	22	-	-	-	-	-	2	1	3	6	5	5	-
2001	28	1	-	-	1	1	1	2	3	9	6	4	-
2000	27	-	-	-	1	-	2	3	2	8	7	4	-
1999	29	-	-	-	-	-	2	5	5	10	4	3	-
Non-Hispanic black, female													
2005 2/	62	-	-	-	2	3	8	7	5	9	14	14	-
2004 2/	56	-	-	-	-	2	3	8	6	3	20	14	-
2003 2/	66	-	1	-	2	2	6	8	9	7	13	18	-
2002	51	-	-	-	3	2	4	7	7	2	10	16	-
2001	52	-	-	-	-	2	6	8	4	4	16	12	-
2000	74	-	-	-	-	2	5	8	6	11	21	21	-
1999	56	1	-	-	2	1	4	5	6	10	12	15	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Infections of kidney (N10-N12,N13.6,N15.1)													
Origin not stated 3/, both sexes													
2005	1	-	-	-	-	-	-	-	-	1	-	-	-
2004	1	-	-	-	-	1	-	-	-	-	-	-	-
2003	3	-	-	-	-	-	1	1	-	-	1	-	-
2002	1	-	-	-	-	-	-	-	-	-	-	1	-
2001	1	-	-	-	-	-	-	-	-	-	1	-	-
2000	3	-	-	-	-	-	-	-	-	1	2	-	-
1999	5	-	-	-	-	-	1	-	1	-	2	1	-
Origin not stated 3/, male													
2005	1	-	-	-	-	-	-	-	-	1	-	-	-
2004	-	-	-	-	-	-	-	-	-	-	-	-	-
2003	2	-	-	-	-	-	1	1	-	-	-	-	-
2002	-	-	-	-	-	-	-	-	-	-	-	-	-
2001	-	-	-	-	-	-	-	-	-	-	-	-	-
2000	2	-	-	-	-	-	-	-	-	1	1	-	-
1999	2	-	-	-	-	-	1	-	-	-	1	-	-
Origin not stated 3/, female													
2005	-	-	-	-	-	-	-	-	-	-	-	-	-
2004	1	-	-	-	-	1	-	-	-	-	-	-	-
2003	1	-	-	-	-	-	-	-	-	-	1	-	-
2002	1	-	-	-	-	-	-	-	-	-	-	1	-
2001	1	-	-	-	-	-	-	-	-	-	1	-	-
2000	1	-	-	-	-	-	-	-	-	-	1	-	-
1999	3	-	-	-	-	-	-	-	1	-	1	1	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Hyperplasia of prostate (N40)													
All origins, both sexes													
2005	525	-	-	-	-	-	-	-	10	52	171	292	-
2004	507	-	-	-	-	-	-	4	13	45	177	268	-
2003	477	1	-	-	-	-	-	1	12	50	165	248	-
2002	437	-	-	-	-	-	1	-	10	41	164	220	1
2001	439	-	-	-	-	-	-	2	9	45	141	242	-
2000	433	-	-	-	-	-	1	1	11	38	159	222	1
1999	430	-	-	-	-	-	-	1	13	47	160	209	-
All origins, male													
2005	525	-	-	-	-	-	-	-	10	52	171	292	-
2004	507	-	-	-	-	-	-	4	13	45	177	268	-
2003	477	1	-	-	-	-	-	1	12	50	165	248	-
2002	437	-	-	-	-	-	1	-	10	41	164	220	1
2001	439	-	-	-	-	-	-	2	9	45	141	242	-
2000	433	-	-	-	-	-	1	1	11	38	159	222	1
1999	430	-	-	-	-	-	-	1	13	47	160	209	-
All origins, female													
2005	...	...	...	...	...	...	...	...	...	...	...	...	...
2004	...	...	...	...	...	...	...	...	...	...	...	...	...
2003	...	...	...	...	...	...	...	...	...	...	...	...	...
2002	...	...	...	...	...	...	...	...	...	...	...	...	...
2001	...	...	...	...	...	...	...	...	...	...	...	...	...
2000	...	...	...	...	...	...	...	...	...	...	...	...	...
1999	...	...	...	...	...	...	...	...	...	...	...	...	...

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Hyperplasia of prostate (N40)													
Hispanic, both sexes													
2005	18	-	-	-	-	-	-	-	2	4	3	9	-
2004	19	-	-	-	-	-	-	-	1	5	4	9	-
2003	24	-	-	-	-	-	-	-	-	6	11	7	-
2002	14	-	-	-	-	-	-	-	1	3	9	1	-
2001	11	-	-	-	-	-	-	-	1	4	4	2	-
2000	19	-	-	-	-	-	-	-	1	5	3	10	-
1999	11	-	-	-	-	-	-	-	1	3	4	3	-
Hispanic, male													
2005	18	-	-	-	-	-	-	-	2	4	3	9	-
2004	19	-	-	-	-	-	-	-	1	5	4	9	-
2003	24	-	-	-	-	-	-	-	-	6	11	7	-
2002	14	-	-	-	-	-	-	-	1	3	9	1	-
2001	11	-	-	-	-	-	-	-	1	4	4	2	-
2000	19	-	-	-	-	-	-	-	1	5	3	10	-
1999	11	-	-	-	-	-	-	-	1	3	4	3	-
Hispanic, female													
2005	...	...	...	...	...	...	...	...	...	...	...	...	...
2004	...	...	...	...	...	...	...	...	...	...	...	...	...
2003	...	...	...	...	...	...	...	...	...	...	...	...	...
2002	...	...	...	...	...	...	...	...	...	...	...	...	...
2001	...	...	...	...	...	...	...	...	...	...	...	...	...
2000	...	...	...	...	...	...	...	...	...	...	...	...	...
1999	...	...	...	...	...	...	...	...	...	...	...	...	...

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Hyperplasia of prostate (N40)													
Non-Hispanic 1/, both sexes													
2005	506	-	-	-	-	-	-	-	8	48	168	282	-
2004	486	-	-	-	-	-	-	4	12	40	172	258	-
2003	449	1	-	-	-	-	-	1	11	42	153	241	-
2002	417	-	-	-	-	-	1	-	8	36	155	216	1
2001	423	-	-	-	-	-	-	2	8	39	134	240	-
2000	414	-	-	-	-	-	1	1	10	33	156	212	1
1999	417	-	-	-	-	-	-	-	12	43	156	206	-
Non-Hispanic 1/, male													
2005	506	-	-	-	-	-	-	-	8	48	168	282	-
2004	486	-	-	-	-	-	-	4	12	40	172	258	-
2003	449	1	-	-	-	-	-	1	11	42	153	241	-
2002	417	-	-	-	-	-	1	-	8	36	155	216	1
2001	423	-	-	-	-	-	-	2	8	39	134	240	-
2000	414	-	-	-	-	-	1	1	10	33	156	212	1
1999	417	-	-	-	-	-	-	-	12	43	156	206	-
Non-Hispanic 1/, female													
2005	...	...	...	...	...	...	...	...	...	...	...	...	...
2004	...	...	...	...	...	...	...	...	...	...	...	...	...
2003	...	...	...	...	...	...	...	...	...	...	...	...	...
2002	...	...	...	...	...	...	...	...	...	...	...	...	...
2001	...	...	...	...	...	...	...	...	...	...	...	...	...
2000	...	...	...	...	...	...	...	...	...	...	...	...	...
1999	...	...	...	...	...	...	...	...	...	...	...	...	...

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Hyperplasia of prostate (N40)													
Non-Hispanic white, both sexes													
2005 2/	464	-	-	-	-	-	-	-	5	41	156	262	-
2004 2/	438	-	-	-	-	-	-	2	11	35	151	239	-
2003 2/	414	-	-	-	-	-	-	-	9	37	138	230	-
2002	370	-	-	-	-	-	-	-	5	29	137	199	-
2001	388	-	-	-	-	-	-	1	4	35	122	226	-
2000	370	-	-	-	-	-	-	1	6	30	142	191	-
1999	377	-	-	-	-	-	-	-	8	36	143	190	-
Non-Hispanic white, male													
2005 2/	464	-	-	-	-	-	-	-	5	41	156	262	-
2004 2/	438	-	-	-	-	-	-	2	11	35	151	239	-
2003 2/	414	-	-	-	-	-	-	-	9	37	138	230	-
2002	370	-	-	-	-	-	-	-	5	29	137	199	-
2001	388	-	-	-	-	-	-	1	4	35	122	226	-
2000	370	-	-	-	-	-	-	1	6	30	142	191	-
1999	377	-	-	-	-	-	-	-	8	36	143	190	-
Non-Hispanic white, female													
2005 2/	...	...	...	...	...	...	...	...	...	...	...	...	...
2004 2/	...	...	...	...	...	...	...	...	...	...	...	...	...
2003 2/	...	...	...	...	...	...	...	...	...	...	...	...	...
2002	...	...	...	...	...	...	...	...	...	...	...	...	...
2001	...	...	...	...	...	...	...	...	...	...	...	...	...
2000	...	...	...	...	...	...	...	...	...	...	...	...	...
1999	...	...	...	...	...	...	...	...	...	...	...	...	...

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Hyperplasia of prostate (N40)													
Non-Hispanic black, both sexes													
2005 2/	30	-	-	-	-	-	-	-	3	5	8	14	-
2004 2/	34	-	-	-	-	-	-	2	1	5	13	13	-
2003 2/	28	1	-	-	-	-	-	1	2	4	11	9	-
2002	38	-	-	-	-	-	1	-	3	7	13	13	1
2001	31	-	-	-	-	-	-	1	4	3	12	11	-
2000	37	-	-	-	-	-	1	-	4	2	14	15	1
1999	38	-	-	-	-	-	-	-	4	6	13	15	-
Non-Hispanic black, male													
2005 2/	30	-	-	-	-	-	-	-	3	5	8	14	-
2004 2/	34	-	-	-	-	-	-	2	1	5	13	13	-
2003 2/	28	1	-	-	-	-	-	1	2	4	11	9	-
2002	38	-	-	-	-	-	1	-	3	7	13	13	1
2001	31	-	-	-	-	-	-	1	4	3	12	11	-
2000	37	-	-	-	-	-	1	-	4	2	14	15	1
1999	38	-	-	-	-	-	-	-	4	6	13	15	-
Non-Hispanic black, female													
2005 2/	...	...	...	...	...	...	...	...	...	...	...	...	...
2004 2/	...	...	...	...	...	...	...	...	...	...	...	...	...
2003 2/	...	...	...	...	...	...	...	...	...	...	...	...	...
2002	...	...	...	...	...	...	...	...	...	...	...	...	...
2001	...	...	...	...	...	...	...	...	...	...	...	...	...
2000	...	...	...	...	...	...	...	...	...	...	...	...	...
1999	...	...	...	...	...	...	...	...	...	...	...	...	...

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Hyperplasia of prostate (N40)													
Origin not stated 3/, both sexes													
2005	1	-	-	-	-	-	-	-	-	-	-	1	-
2004	2	-	-	-	-	-	-	-	-	-	1	1	-
2003	4	-	-	-	-	-	-	-	1	2	1	-	-
2002	6	-	-	-	-	-	-	-	1	2	-	3	-
2001	5	-	-	-	-	-	-	-	-	2	3	-	-
2000	-	-	-	-	-	-	-	-	-	-	-	-	-
1999	2	-	-	-	-	-	-	1	-	1	-	-	-
Origin not stated 3/, male													
2005	1	-	-	-	-	-	-	-	-	-	-	1	-
2004	2	-	-	-	-	-	-	-	-	-	1	1	-
2003	4	-	-	-	-	-	-	-	1	2	1	-	-
2002	6	-	-	-	-	-	-	-	1	2	-	3	-
2001	5	-	-	-	-	-	-	-	-	2	3	-	-
2000	-	-	-	-	-	-	-	-	-	-	-	-	-
1999	2	-	-	-	-	-	-	1	-	1	-	-	-
Origin not stated 3/, female													
2005	...	...	...	...	...	...	...	...	...	...	...	...	...
2004	...	...	...	...	...	...	...	...	...	...	...	...	...
2003	...	...	...	...	...	...	...	...	...	...	...	...	...
2002	...	...	...	...	...	...	...	...	...	...	...	...	...
2001	...	...	...	...	...	...	...	...	...	...	...	...	...
2000	...	...	...	...	...	...	...	...	...	...	...	...	...
1999	...	...	...	...	...	...	...	...	...	...	...	...	...

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Inflammatory diseases of female pelvic organs (N70-N76)													
All origins, both sexes													
2005	120	-	-	-	1	5	7	13	9	23	33	29	-
2004	112	1	-	-	2	4	8	11	15	16	21	34	-
2003	125	1	-	1	6	3	14	14	10	16	29	31	-
2002	114	1	-	-	2	4	17	11	6	8	29	36	-
2001	127	1	-	-	2	3	10	12	12	18	31	38	-
2000	120	2	-	1	2	6	15	9	8	21	29	27	-
1999	108	2	1	-	3	6	12	6	13	15	32	18	-
All origins, male													
2005	...	...	...	...	...	...	...	...	...	...	...	...	...
2004	...	...	...	...	...	...	...	...	...	...	...	...	...
2003	...	...	...	...	...	...	...	...	...	...	...	...	...
2002	...	...	...	...	...	...	...	...	...	...	...	...	...
2001	...	...	...	...	...	...	...	...	...	...	...	...	...
2000	...	...	...	...	...	...	...	...	...	...	...	...	...
1999	...	...	...	...	...	...	...	...	...	...	...	...	...
All origins, female													
2005	120	-	-	-	1	5	7	13	9	23	33	29	-
2004	112	1	-	-	2	4	8	11	15	16	21	34	-
2003	125	1	-	1	6	3	14	14	10	16	29	31	-
2002	114	1	-	-	2	4	17	11	6	8	29	36	-
2001	127	1	-	-	2	3	10	12	12	18	31	38	-
2000	120	2	-	1	2	6	15	9	8	21	29	27	-
1999	108	2	1	-	3	6	12	6	13	15	32	18	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Inflammatory diseases of female pelvic organs (N70-N76)													
Hispanic, both sexes													
2005	9	-	-	-	-	2	4	-	-	2	1	-	-
2004	9	1	-	-	1	1	-	-	-	2	3	1	-
2003	9	-	-	-	1	1	1	1	-	3	1	1	-
2002	3	-	-	-	-	1	2	-	-	-	-	-	-
2001	5	-	-	-	-	-	-	2	-	2	-	1	-
2000	7	-	-	-	-	1	1	-	2	2	-	1	-
1999	7	-	-	-	-	-	1	2	1	-	1	2	-
Hispanic, male													
2005	...	...	...	...	...	...	...	...	...	...	...	...	...
2004	...	...	...	...	...	...	...	...	...	...	...	...	...
2003	...	...	...	...	...	...	...	...	...	...	...	...	...
2002	...	...	...	...	...	...	...	...	...	...	...	...	...
2001	...	...	...	...	...	...	...	...	...	...	...	...	...
2000	...	...	...	...	...	...	...	...	...	...	...	...	...
1999	...	...	...	...	...	...	...	...	...	...	...	...	...
Hispanic, female													
2005	9	-	-	-	-	2	4	-	-	2	1	-	-
2004	9	1	-	-	1	1	-	-	-	2	3	1	-
2003	9	-	-	-	1	1	1	1	-	3	1	1	-
2002	3	-	-	-	-	1	2	-	-	-	-	-	-
2001	5	-	-	-	-	-	-	2	-	2	-	1	-
2000	7	-	-	-	-	1	1	-	2	2	-	1	-
1999	7	-	-	-	-	-	1	2	1	-	1	2	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Inflammatory diseases of female pelvic organs (N70-N76)													
Non-Hispanic 1/, both sexes													
2005	111	-	-	-	1	3	3	13	9	21	32	29	-
2004	102	-	-	-	1	3	8	11	14	14	18	33	-
2003	115	1	-	1	5	2	13	12	10	13	28	30	-
2002	110	1	-	-	2	3	15	11	6	8	28	36	-
2001	122	1	-	-	2	3	10	10	12	16	31	37	-
2000	111	2	-	1	2	5	14	7	6	19	29	26	-
1999	101	2	1	-	3	6	11	4	12	15	31	16	-
Non-Hispanic 1/, male													
2005	...	...	...	...	...	...	...	...	...	...	...	...	...
2004	...	...	...	...	...	...	...	...	...	...	...	...	...
2003	...	...	...	...	...	...	...	...	...	...	...	...	...
2002	...	...	...	...	...	...	...	...	...	...	...	...	...
2001	...	...	...	...	...	...	...	...	...	...	...	...	...
2000	...	...	...	...	...	...	...	...	...	...	...	...	...
1999	...	...	...	...	...	...	...	...	...	...	...	...	...
Non-Hispanic 1/, female													
2005	111	-	-	-	1	3	3	13	9	21	32	29	-
2004	102	-	-	-	1	3	8	11	14	14	18	33	-
2003	115	1	-	1	5	2	13	12	10	13	28	30	-
2002	110	1	-	-	2	3	15	11	6	8	28	36	-
2001	122	1	-	-	2	3	10	10	12	16	31	37	-
2000	111	2	-	1	2	5	14	7	6	19	29	26	-
1999	101	2	1	-	3	6	11	4	12	15	31	16	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Inflammatory diseases of female pelvic organs (N70-N76)													
Non-Hispanic white, both sexes													
2005 2/	89	-	-	-	-	1	2	7	7	17	30	25	-
2004 2/	83	-	-	-	1	2	6	6	13	12	13	30	-
2003 2/	93	-	-	1	3	1	9	8	8	10	24	29	-
2002	88	1	-	-	-	2	7	7	5	7	27	32	-
2001	98	1	-	-	2	3	6	6	8	12	27	33	-
2000	87	1	-	-	2	3	6	6	5	15	26	23	-
1999	73	1	1	-	2	3	3	2	11	10	25	15	-
Non-Hispanic white, male													
2005 2/	...	...	...	...	...	...	...	...	...	...	...	...	...
2004 2/	...	...	...	...	...	...	...	...	...	...	...	...	...
2003 2/	...	...	...	...	...	...	...	...	...	...	...	...	...
2002	...	...	...	...	...	...	...	...	...	...	...	...	...
2001	...	...	...	...	...	...	...	...	...	...	...	...	...
2000	...	...	...	...	...	...	...	...	...	...	...	...	...
1999	...	...	...	...	...	...	...	...	...	...	...	...	...
Non-Hispanic white, female													
2005 2/	89	-	-	-	-	1	2	7	7	17	30	25	-
2004 2/	83	-	-	-	1	2	6	6	13	12	13	30	-
2003 2/	93	-	-	1	3	1	9	8	8	10	24	29	-
2002	88	1	-	-	-	2	7	7	5	7	27	32	-
2001	98	1	-	-	2	3	6	6	8	12	27	33	-
2000	87	1	-	-	2	3	6	6	5	15	26	23	-
1999	73	1	1	-	2	3	3	2	11	10	25	15	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Inflammatory diseases of female pelvic organs (N70-N76)													
Non-Hispanic black, both sexes													
2005 2/	19	-	-	-	1	1	-	6	2	3	2	4	-
2004 2/	19	-	-	-	-	1	2	5	1	2	5	3	-
2003 2/	19	1	-	-	2	1	4	4	2	2	2	1	-
2002	20	-	-	-	2	1	8	3	1	1	1	3	-
2001	23	-	-	-	-	-	4	4	4	4	4	3	-
2000	21	1	-	1	-	1	7	1	1	4	3	2	-
1999	26	1	-	-	1	2	7	2	1	5	6	1	-
Non-Hispanic black, male													
2005 2/	...	...	...	...	...	...	...	...	...	...	...	...	...
2004 2/	...	...	...	...	...	...	...	...	...	...	...	...	...
2003 2/	...	...	...	...	...	...	...	...	...	...	...	...	...
2002	...	...	...	...	...	...	...	...	...	...	...	...	...
2001	...	...	...	...	...	...	...	...	...	...	...	...	...
2000	...	...	...	...	...	...	...	...	...	...	...	...	...
1999	...	...	...	...	...	...	...	...	...	...	...	...	...
Non-Hispanic black, female													
2005 2/	19	-	-	-	1	1	-	6	2	3	2	4	-
2004 2/	19	-	-	-	-	1	2	5	1	2	5	3	-
2003 2/	19	1	-	-	2	1	4	4	2	2	2	1	-
2002	20	-	-	-	2	1	8	3	1	1	1	3	-
2001	23	-	-	-	-	-	4	4	4	4	4	3	-
2000	21	1	-	1	-	1	7	1	1	4	3	2	-
1999	26	1	-	-	1	2	7	2	1	5	6	1	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Inflammatory diseases of female pelvic organs (N70-N76)													
Origin not stated 3/, both sexes													
2005	-	-	-	-	-	-	-	-	-	-	-	-	-
2004	1	-	-	-	-	-	-	-	1	-	-	-	-
2003	1	-	-	-	-	-	-	1	-	-	-	-	-
2002	1	-	-	-	-	-	-	-	-	-	1	-	-
2001	-	-	-	-	-	-	-	-	-	-	-	-	-
2000	2	-	-	-	-	-	-	2	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-	-	-	-
Origin not stated 3/, male													
2005	...	...	...	...	...	...	...	...	...	...	...	...	...
2004	...	...	...	...	...	...	...	...	...	...	...	...	...
2003	...	...	...	...	...	...	...	...	...	...	...	...	...
2002	...	...	...	...	...	...	...	...	...	...	...	...	...
2001	...	...	...	...	...	...	...	...	...	...	...	...	...
2000	...	...	...	...	...	...	...	...	...	...	...	...	...
1999	...	...	...	...	...	...	...	...	...	...	...	...	...
Origin not stated 3/, female													
2005	-	-	-	-	-	-	-	-	-	-	-	-	-
2004	1	-	-	-	-	-	-	-	1	-	-	-	-
2003	1	-	-	-	-	-	-	1	-	-	-	-	-
2002	1	-	-	-	-	-	-	-	-	-	1	-	-
2001	-	-	-	-	-	-	-	-	-	-	-	-	-
2000	2	-	-	-	-	-	-	2	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-	-	-	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Pregnancy, childbirth and the puerperium (O00-O99)													
All origins, both sexes													
2005	760	...	...	-	183	312	207	54	3	1	-	-	-
2004	697	...	...	2	168	287	157	79	1	1	2	-	-
2003	545	...	...	1	116	229	166	30	2	1	-	-	-
2002	379	...	...	1	89	175	105	8	-	-	-	1	-
2001	416	...	...	3	110	190	106	6	-	-	-	1	-
2000	404	...	...	-	95	182	121	6	-	-	-	-	-
1999	406	...	...	-	97	181	121	7	-	-	-	-	-
All origins, male													
2005	...	...	...	...	...	...	...	...	...	...	...	...	...
2004	...	...	...	...	...	...	...	...	...	...	...	...	...
2003	...	...	...	...	...	...	...	...	...	...	...	...	...
2002	...	...	...	...	...	...	...	...	...	...	...	...	...
2001	...	...	...	...	...	...	...	...	...	...	...	...	...
2000	...	...	...	...	...	...	...	...	...	...	...	...	...
1999	...	...	...	...	...	...	...	...	...	...	...	...	...
All origins, female													
2005	760	...	...	-	183	312	207	54	3	1	-	-	-
2004	697	...	...	2	168	287	157	79	1	1	2	-	-
2003	545	...	...	1	116	229	166	30	2	1	-	-	-
2002	379	...	...	1	89	175	105	8	-	-	-	1	-
2001	416	...	...	3	110	190	106	6	-	-	-	1	-
2000	404	...	...	-	95	182	121	6	-	-	-	-	-
1999	406	...	...	-	97	181	121	7	-	-	-	-	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Pregnancy, childbirth and the puerperium (O00-O99)													
Hispanic, both sexes													
2005	129	...	...	-	35	60	32	2	-	-	-	-	-
2004	115	...	...	-	29	49	26	11	-	-	-	-	-
2003	104	...	...	-	24	46	28	4	1	1	-	-	-
2002	70	...	...	-	19	28	21	2	-	-	-	-	-
2001	83	...	...	1	26	38	17	1	-	-	-	-	-
2000	84	...	...	-	24	36	23	1	-	-	-	-	-
1999	70	...	...	-	19	26	25	-	-	-	-	-	-
Hispanic, male													
2005	...	...	...	...	...	...	...	...	...	...	...	...	...
2004	...	...	...	...	...	...	...	...	...	...	...	...	...
2003	...	...	...	...	...	...	...	...	...	...	...	...	...
2002	...	...	...	...	...	...	...	...	...	...	...	...	...
2001	...	...	...	...	...	...	...	...	...	...	...	...	...
2000	...	...	...	...	...	...	...	...	...	...	...	...	...
1999	...	...	...	...	...	...	...	...	...	...	...	...	...
Hispanic, female													
2005	129	...	...	-	35	60	32	2	-	-	-	-	-
2004	115	...	...	-	29	49	26	11	-	-	-	-	-
2003	104	...	...	-	24	46	28	4	1	1	-	-	-
2002	70	...	...	-	19	28	21	2	-	-	-	-	-
2001	83	...	...	1	26	38	17	1	-	-	-	-	-
2000	84	...	...	-	24	36	23	1	-	-	-	-	-
1999	70	...	...	-	19	26	25	-	-	-	-	-	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Pregnancy, childbirth and the puerperium (O00-O99)													
Non-Hispanic 1/, both sexes													
2005	630	...	...	-	148	252	175	51	3	1	-	-	-
2004	581	...	...	2	139	238	130	68	1	1	2	-	-
2003	437	...	...	1	89	182	138	26	1	-	-	-	-
2002	305	...	...	1	69	145	83	6	-	-	-	1	-
2001	332	...	...	2	84	151	89	5	-	-	-	1	-
2000	319	...	...	-	71	146	97	5	-	-	-	-	-
1999	333	...	...	-	76	154	96	7	-	-	-	-	-
Non-Hispanic 1/, male													
2005	...	...	...	...	...	...	...	...	...	...	...	...	...
2004	...	...	...	...	...	...	...	...	...	...	...	...	...
2003	...	...	...	...	...	...	...	...	...	...	...	...	...
2002	...	...	...	...	...	...	...	...	...	...	...	...	...
2001	...	...	...	...	...	...	...	...	...	...	...	...	...
2000	...	...	...	...	...	...	...	...	...	...	...	...	...
1999	...	...	...	...	...	...	...	...	...	...	...	...	...
Non-Hispanic 1/, female													
2005	630	...	...	-	148	252	175	51	3	1	-	-	-
2004	581	...	...	2	139	238	130	68	1	1	2	-	-
2003	437	...	...	1	89	182	138	26	1	-	-	-	-
2002	305	...	...	1	69	145	83	6	-	-	-	1	-
2001	332	...	...	2	84	151	89	5	-	-	-	1	-
2000	319	...	...	-	71	146	97	5	-	-	-	-	-
1999	333	...	...	-	76	154	96	7	-	-	-	-	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Pregnancy, childbirth and the puerperium (O00-O99)													
Non-Hispanic white, both sexes													
2005 2/	327	...	...	-	76	121	94	33	2	1	-	-	-
2004 2/	302	...	...	-	60	122	74	43	1	1	1	-	-
2003 2/	213	...	...	-	41	87	66	19	-	-	-	-	-
2002	133	...	...	-	21	69	39	3	-	-	-	1	-
2001	156	...	...	1	38	77	37	2	-	-	-	1	-
2000	161	...	...	-	33	72	54	2	-	-	-	-	-
1999	152	...	...	-	26	74	48	4	-	-	-	-	-
Non-Hispanic white, male													
2005 2/	...	...	...	...	...	...	...	...	...	...	...	...	...
2004 2/	...	...	...	...	...	...	...	...	...	...	...	...	...
2003 2/	...	...	...	...	...	...	...	...	...	...	...	...	...
2002	...	...	...	...	...	...	...	...	...	...	...	...	...
2001	...	...	...	...	...	...	...	...	...	...	...	...	...
2000	...	...	...	...	...	...	...	...	...	...	...	...	...
1999	...	...	...	...	...	...	...	...	...	...	...	...	...
Non-Hispanic white, female													
2005 2/	327	...	...	-	76	121	94	33	2	1	-	-	-
2004 2/	302	...	...	-	60	122	74	43	1	1	1	-	-
2003 2/	213	...	...	-	41	87	66	19	-	-	-	-	-
2002	133	...	...	-	21	69	39	3	-	-	-	1	-
2001	156	...	...	1	38	77	37	2	-	-	-	1	-
2000	161	...	...	-	33	72	54	2	-	-	-	-	-
1999	152	...	...	-	26	74	48	4	-	-	-	-	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Pregnancy, childbirth and the puerperium (O00-O99)													
Non-Hispanic black, both sexes													
2005 2/	265	...	...	-	69	115	65	15	1	-	-	-	-
2004 2/	245	...	...	-	73	106	44	22	-	-	-	-	-
2003 2/	190	...	...	1	42	82	58	7	-	-	-	-	-
2002	153	...	...	1	45	64	41	2	-	-	-	-	-
2001	154	...	...	1	45	65	40	3	-	-	-	-	-
2000	139	...	...	-	34	65	37	3	-	-	-	-	-
1999	158	...	...	-	46	70	40	2	-	-	-	-	-
Non-Hispanic black, male													
2005 2/	...	...	...	...	...	...	...	...	...	...	...	...	...
2004 2/	...	...	...	...	...	...	...	...	...	...	...	...	...
2003 2/	...	...	...	...	...	...	...	...	...	...	...	...	...
2002	...	...	...	...	...	...	...	...	...	...	...	...	...
2001	...	...	...	...	...	...	...	...	...	...	...	...	...
2000	...	...	...	...	...	...	...	...	...	...	...	...	...
1999	...	...	...	...	...	...	...	...	...	...	...	...	...
Non-Hispanic black, female													
2005 2/	265	...	...	-	69	115	65	15	1	-	-	-	-
2004 2/	245	...	...	-	73	106	44	22	-	-	-	-	-
2003 2/	190	...	...	1	42	82	58	7	-	-	-	-	-
2002	153	...	...	1	45	64	41	2	-	-	-	-	-
2001	154	...	...	1	45	65	40	3	-	-	-	-	-
2000	139	...	...	-	34	65	37	3	-	-	-	-	-
1999	158	...	...	-	46	70	40	2	-	-	-	-	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Pregnancy, childbirth and the puerperium (O00-O99)													
Origin not stated 3/, both sexes													
2005	1	...	...	-	-	-	-	1	-	-	-	-	-
2004	1	...	...	-	-	-	1	-	-	-	-	-	-
2003	4	...	...	-	3	1	-	-	-	-	-	-	-
2002	4	...	...	-	1	2	1	-	-	-	-	-	-
2001	1	...	...	-	-	1	-	-	-	-	-	-	-
2000	1	...	...	-	-	-	1	-	-	-	-	-	-
1999	3	...	...	-	2	1	-	-	-	-	-	-	-
Origin not stated 3/, male													
2005	...	...	...	...	...	...	...	...	...	...	...	...	...
2004	...	...	...	...	...	...	...	...	...	...	...	...	...
2003	...	...	...	...	...	...	...	...	...	...	...	...	...
2002	...	...	...	...	...	...	...	...	...	...	...	...	...
2001	...	...	...	...	...	...	...	...	...	...	...	...	...
2000	...	...	...	...	...	...	...	...	...	...	...	...	...
1999	...	...	...	...	...	...	...	...	...	...	...	...	...
Origin not stated 3/, female													
2005	1	...	...	-	-	-	-	1	-	-	-	-	-
2004	1	...	...	-	-	-	1	-	-	-	-	-	-
2003	4	...	...	-	3	1	-	-	-	-	-	-	-
2002	4	...	...	-	1	2	1	-	-	-	-	-	-
2001	1	...	...	-	-	1	-	-	-	-	-	-	-
2000	1	...	...	-	-	-	1	-	-	-	-	-	-
1999	3	...	...	-	2	1	-	-	-	-	-	-	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Pregnancy with abortive outcome (O00-O07)													
All origins, both sexes													
2005	33	...	...	-	6	14	11	2	-	-	-	-	-
2004	32	...	...	-	14	11	6	1	-	-	-	-	-
2003	40	...	...	-	11	19	9	1	-	-	-	-	-
2002	22	...	...	1	4	9	7	1	-	-	-	-	-
2001	38	...	...	2	13	14	9	-	-	-	-	-	-
2000	37	...	...	-	6	17	14	-	-	-	-	-	-
1999	31	...	...	-	8	15	8	-	-	-	-	-	-
All origins, male													
2005	...	...	...	...	...	...	...	...	...	...	...	...	...
2004	...	...	...	...	...	...	...	...	...	...	...	...	...
2003	...	...	...	...	...	...	...	...	...	...	...	...	...
2002	...	...	...	...	...	...	...	...	...	...	...	...	...
2001	...	...	...	...	...	...	...	...	...	...	...	...	...
2000	...	...	...	...	...	...	...	...	...	...	...	...	...
1999	...	...	...	...	...	...	...	...	...	...	...	...	...
All origins, female													
2005	33	...	...	-	6	14	11	2	-	-	-	-	-
2004	32	...	...	-	14	11	6	1	-	-	-	-	-
2003	40	...	...	-	11	19	9	1	-	-	-	-	-
2002	22	...	...	1	4	9	7	1	-	-	-	-	-
2001	38	...	...	2	13	14	9	-	-	-	-	-	-
2000	37	...	...	-	6	17	14	-	-	-	-	-	-
1999	31	...	...	-	8	15	8	-	-	-	-	-	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Pregnancy with abortive outcome (O00-O07)													
Hispanic, both sexes													
2005	2	...	...	-	1	1	-	-	-	-	-	-	-
2004	1	...	...	-	-	1	-	-	-	-	-	-	-
2003	8	...	...	-	1	6	1	-	-	-	-	-	-
2002	1	...	...	-	-	1	-	-	-	-	-	-	-
2001	1	...	...	1	-	-	-	-	-	-	-	-	-
2000	6	...	...	-	2	3	1	-	-	-	-	-	-
1999	4	...	...	-	2	-	2	-	-	-	-	-	-
Hispanic, male													
2005	...	...	...	...	...	...	...	...	...	...	...	...	...
2004	...	...	...	...	...	...	...	...	...	...	...	...	...
2003	...	...	...	...	...	...	...	...	...	...	...	...	...
2002	...	...	...	...	...	...	...	...	...	...	...	...	...
2001	...	...	...	...	...	...	...	...	...	...	...	...	...
2000	...	...	...	...	...	...	...	...	...	...	...	...	...
1999	...	...	...	...	...	...	...	...	...	...	...	...	...
Hispanic, female													
2005	2	...	...	-	1	1	-	-	-	-	-	-	-
2004	1	...	...	-	-	1	-	-	-	-	-	-	-
2003	8	...	...	-	1	6	1	-	-	-	-	-	-
2002	1	...	...	-	-	1	-	-	-	-	-	-	-
2001	1	...	...	1	-	-	-	-	-	-	-	-	-
2000	6	...	...	-	2	3	1	-	-	-	-	-	-
1999	4	...	...	-	2	-	2	-	-	-	-	-	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Pregnancy with abortive outcome (O00-O07)													
Non-Hispanic 1/, both sexes													
2005	31	...	...	-	5	13	11	2	-	-	-	-	-
2004	31	...	...	-	14	10	6	1	-	-	-	-	-
2003	32	...	...	-	10	13	8	1	-	-	-	-	-
2002	20	...	...	1	4	8	6	1	-	-	-	-	-
2001	37	...	...	1	13	14	9	-	-	-	-	-	-
2000	30	...	...	-	4	14	12	-	-	-	-	-	-
1999	26	...	...	-	5	15	6	-	-	-	-	-	-
Non-Hispanic 1/, male													
2005	...	...	...	...	...	...	...	...	...	...	...	...	...
2004	...	...	...	...	...	...	...	...	...	...	...	...	...
2003	...	...	...	...	...	...	...	...	...	...	...	...	...
2002	...	...	...	...	...	...	...	...	...	...	...	...	...
2001	...	...	...	...	...	...	...	...	...	...	...	...	...
2000	...	...	...	...	...	...	...	...	...	...	...	...	...
1999	...	...	...	...	...	...	...	...	...	...	...	...	...
Non-Hispanic 1/, female													
2005	31	...	...	-	5	13	11	2	-	-	-	-	-
2004	31	...	...	-	14	10	6	1	-	-	-	-	-
2003	32	...	...	-	10	13	8	1	-	-	-	-	-
2002	20	...	...	1	4	8	6	1	-	-	-	-	-
2001	37	...	...	1	13	14	9	-	-	-	-	-	-
2000	30	...	...	-	4	14	12	-	-	-	-	-	-
1999	26	...	...	-	5	15	6	-	-	-	-	-	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Pregnancy with abortive outcome (O00-O07)													
Non-Hispanic white, both sexes													
2005 2/	14	...	...	-	4	3	5	2	-	-	-	-	-
2004 2/	10	...	...	-	5	4	-	1	-	-	-	-	-
2003 2/	11	...	...	-	2	6	3	-	-	-	-	-	-
2002	6	...	...	-	1	4	1	-	-	-	-	-	-
2001	19	...	...	-	7	8	4	-	-	-	-	-	-
2000	12	...	...	-	-	5	7	-	-	-	-	-	-
1999	11	...	...	-	2	6	3	-	-	-	-	-	-
Non-Hispanic white, male													
2005 2/	...	...	...	...	...	...	...	...	...	...	...	...	...
2004 2/	...	...	...	...	...	...	...	...	...	...	...	...	...
2003 2/	...	...	...	...	...	...	...	...	...	...	...	...	...
2002	...	...	...	...	...	...	...	...	...	...	...	...	...
2001	...	...	...	...	...	...	...	...	...	...	...	...	...
2000	...	...	...	...	...	...	...	...	...	...	...	...	...
1999	...	...	...	...	...	...	...	...	...	...	...	...	...
Non-Hispanic white, female													
2005 2/	14	...	...	-	4	3	5	2	-	-	-	-	-
2004 2/	10	...	...	-	5	4	-	1	-	-	-	-	-
2003 2/	11	...	...	-	2	6	3	-	-	-	-	-	-
2002	6	...	...	-	1	4	1	-	-	-	-	-	-
2001	19	...	...	-	7	8	4	-	-	-	-	-	-
2000	12	...	...	-	-	5	7	-	-	-	-	-	-
1999	11	...	...	-	2	6	3	-	-	-	-	-	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Pregnancy with abortive outcome (O00-O07)													
Non-Hispanic black, both sexes													
2005 2/	13	...	...	-	1	8	4	-	-	-	-	-	-
2004 2/	19	...	...	-	8	6	5	-	-	-	-	-	-
2003 2/	18	...	...	-	8	5	4	1	-	-	-	-	-
2002	11	...	...	1	2	4	4	-	-	-	-	-	-
2001	16	...	...	1	6	5	4	-	-	-	-	-	-
2000	18	...	...	-	4	9	5	-	-	-	-	-	-
1999	14	...	...	-	3	8	3	-	-	-	-	-	-
Non-Hispanic black, male													
2005 2/	...	...	...	...	...	...	...	...	...	...	...	...	...
2004 2/	...	...	...	...	...	...	...	...	...	...	...	...	...
2003 2/	...	...	...	...	...	...	...	...	...	...	...	...	...
2002	...	...	...	...	...	...	...	...	...	...	...	...	...
2001	...	...	...	...	...	...	...	...	...	...	...	...	...
2000	...	...	...	...	...	...	...	...	...	...	...	...	...
1999	...	...	...	...	...	...	...	...	...	...	...	...	...
Non-Hispanic black, female													
2005 2/	13	...	...	-	1	8	4	-	-	-	-	-	-
2004 2/	19	...	...	-	8	6	5	-	-	-	-	-	-
2003 2/	18	...	...	-	8	5	4	1	-	-	-	-	-
2002	11	...	...	1	2	4	4	-	-	-	-	-	-
2001	16	...	...	1	6	5	4	-	-	-	-	-	-
2000	18	...	...	-	4	9	5	-	-	-	-	-	-
1999	14	...	...	-	3	8	3	-	-	-	-	-	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Pregnancy with abortive outcome (O00-O07)													
Origin not stated 3/, both sexes													
2005	-	...	...	-	-	-	-	-	-	-	-	-	-
2004	-	...	...	-	-	-	-	-	-	-	-	-	-
2003	-	...	...	-	-	-	-	-	-	-	-	-	-
2002	1	...	...	-	-	-	1	-	-	-	-	-	-
2001	-	...	...	-	-	-	-	-	-	-	-	-	-
2000	1	...	...	-	-	-	1	-	-	-	-	-	-
1999	1	...	...	-	1	-	-	-	-	-	-	-	-
Origin not stated 3/, male													
2005	...	...	...	...	...	...	...	...	...	...	...	...	...
2004	...	...	...	...	...	...	...	...	...	...	...	...	...
2003	...	...	...	...	...	...	...	...	...	...	...	...	...
2002	...	...	...	...	...	...	...	...	...	...	...	...	...
2001	...	...	...	...	...	...	...	...	...	...	...	...	...
2000	...	...	...	...	...	...	...	...	...	...	...	...	...
1999	...	...	...	...	...	...	...	...	...	...	...	...	...
Origin not stated 3/, female													
2005	-	...	...	-	-	-	-	-	-	-	-	-	-
2004	-	...	...	-	-	-	-	-	-	-	-	-	-
2003	-	...	...	-	-	-	-	-	-	-	-	-	-
2002	1	...	...	-	-	-	1	-	-	-	-	-	-
2001	-	...	...	-	-	-	-	-	-	-	-	-	-
2000	1	...	...	-	-	-	1	-	-	-	-	-	-
1999	1	...	...	-	1	-	-	-	-	-	-	-	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Other complications of pregnancy, childbirth and the puerperium (O10-O99)													
All origins, both sexes													
2005	727	...	...	-	177	298	196	52	3	1	-	-	-
2004	665	...	...	2	154	276	151	78	1	1	2	-	-
2003	505	...	...	1	105	210	157	29	2	1	-	-	-
2002	357	...	...	-	85	166	98	7	-	-	-	1	-
2001	378	...	...	1	97	176	97	6	-	-	-	1	-
2000	367	...	...	-	89	165	107	6	-	-	-	-	-
1999	375	...	...	-	89	166	113	7	-	-	-	-	-
All origins, male													
2005	...	...	...	...	...	...	...	...	...	...	...	...	...
2004	...	...	...	...	...	...	...	...	...	...	...	...	...
2003	...	...	...	...	...	...	...	...	...	...	...	...	...
2002	...	...	...	...	...	...	...	...	...	...	...	...	...
2001	...	...	...	...	...	...	...	...	...	...	...	...	...
2000	...	...	...	...	...	...	...	...	...	...	...	...	...
1999	...	...	...	...	...	...	...	...	...	...	...	...	...
All origins, female													
2005	727	...	...	-	177	298	196	52	3	1	-	-	-
2004	665	...	...	2	154	276	151	78	1	1	2	-	-
2003	505	...	...	1	105	210	157	29	2	1	-	-	-
2002	357	...	...	-	85	166	98	7	-	-	-	1	-
2001	378	...	...	1	97	176	97	6	-	-	-	1	-
2000	367	...	...	-	89	165	107	6	-	-	-	-	-
1999	375	...	...	-	89	166	113	7	-	-	-	-	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Other complications of pregnancy, childbirth and the puerperium (O10-O99)													
Hispanic, both sexes													
2005	127	...	...	-	34	59	32	2	-	-	-	-	-
2004	114	...	...	-	29	48	26	11	-	-	-	-	-
2003	96	...	...	-	23	40	27	4	1	1	-	-	-
2002	69	...	...	-	19	27	21	2	-	-	-	-	-
2001	82	...	...	-	26	38	17	1	-	-	-	-	-
2000	78	...	...	-	22	33	22	1	-	-	-	-	-
1999	66	...	...	-	17	26	23	-	-	-	-	-	-
Hispanic, male													
2005	...	...	...	...	...	...	...	...	...	...	...	...	...
2004	...	...	...	...	...	...	...	...	...	...	...	...	...
2003	...	...	...	...	...	...	...	...	...	...	...	...	...
2002	...	...	...	...	...	...	...	...	...	...	...	...	...
2001	...	...	...	...	...	...	...	...	...	...	...	...	...
2000	...	...	...	...	...	...	...	...	...	...	...	...	...
1999	...	...	...	...	...	...	...	...	...	...	...	...	...
Hispanic, female													
2005	127	...	...	-	34	59	32	2	-	-	-	-	-
2004	114	...	...	-	29	48	26	11	-	-	-	-	-
2003	96	...	...	-	23	40	27	4	1	1	-	-	-
2002	69	...	...	-	19	27	21	2	-	-	-	-	-
2001	82	...	...	-	26	38	17	1	-	-	-	-	-
2000	78	...	...	-	22	33	22	1	-	-	-	-	-
1999	66	...	...	-	17	26	23	-	-	-	-	-	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Other complications of pregnancy, childbirth and the puerperium (O10-O99)													
Non-Hispanic 1/, both sexes													
2005	599	...	...	-	143	239	164	49	3	1	-	-	-
2004	550	...	...	2	125	228	124	67	1	1	2	-	-
2003	405	...	...	1	79	169	130	25	1	-	-	-	-
2002	285	...	...	-	65	137	77	5	-	-	-	1	-
2001	295	...	...	1	71	137	80	5	-	-	-	1	-
2000	289	...	...	-	67	132	85	5	-	-	-	-	-
1999	307	...	...	-	71	139	90	7	-	-	-	-	-
Non-Hispanic 1/, male													
2005	...	...	...	...	...	...	...	...	...	...	...	...	...
2004	...	...	...	...	...	...	...	...	...	...	...	...	...
2003	...	...	...	...	...	...	...	...	...	...	...	...	...
2002	...	...	...	...	...	...	...	...	...	...	...	...	...
2001	...	...	...	...	...	...	...	...	...	...	...	...	...
2000	...	...	...	...	...	...	...	...	...	...	...	...	...
1999	...	...	...	...	...	...	...	...	...	...	...	...	...
Non-Hispanic 1/, female													
2005	599	...	...	-	143	239	164	49	3	1	-	-	-
2004	550	...	...	2	125	228	124	67	1	1	2	-	-
2003	405	...	...	1	79	169	130	25	1	-	-	-	-
2002	285	...	...	-	65	137	77	5	-	-	-	1	-
2001	295	...	...	1	71	137	80	5	-	-	-	1	-
2000	289	...	...	-	67	132	85	5	-	-	-	-	-
1999	307	...	...	-	71	139	90	7	-	-	-	-	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Other complications of pregnancy, childbirth and the puerperium (O10-O99)													
Non-Hispanic white, both sexes													
2005 2/	313	...	...	-	72	118	89	31	2	1	-	-	-
2004 2/	292	...	...	-	55	118	74	42	1	1	1	-	-
2003 2/	202	...	...	-	39	81	63	19	-	-	-	-	-
2002	127	...	...	-	20	65	38	3	-	-	-	1	-
2001	137	...	...	1	31	69	33	2	-	-	-	1	-
2000	149	...	...	-	33	67	47	2	-	-	-	-	-
1999	141	...	...	-	24	68	45	4	-	-	-	-	-
Non-Hispanic white, male													
2005 2/	...	...	...	...	...	...	...	...	...	...	...	...	...
2004 2/	...	...	...	...	...	...	...	...	...	...	...	...	...
2003 2/	...	...	...	...	...	...	...	...	...	...	...	...	...
2002	...	...	...	...	...	...	...	...	...	...	...	...	...
2001	...	...	...	...	...	...	...	...	...	...	...	...	...
2000	...	...	...	...	...	...	...	...	...	...	...	...	...
1999	...	...	...	...	...	...	...	...	...	...	...	...	...
Non-Hispanic white, female													
2005 2/	313	...	...	-	72	118	89	31	2	1	-	-	-
2004 2/	292	...	...	-	55	118	74	42	1	1	1	-	-
2003 2/	202	...	...	-	39	81	63	19	-	-	-	-	-
2002	127	...	...	-	20	65	38	3	-	-	-	1	-
2001	137	...	...	1	31	69	33	2	-	-	-	1	-
2000	149	...	...	-	33	67	47	2	-	-	-	-	-
1999	141	...	...	-	24	68	45	4	-	-	-	-	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Other complications of pregnancy, childbirth and the puerperium (O10-O99)													
Non-Hispanic black, both sexes													
2005 2/	252	...	...	-	68	107	61	15	1	-	-	-	-
2004 2/	226	...	...	-	65	100	39	22	-	-	-	-	-
2003 2/	172	...	...	1	34	77	54	6	-	-	-	-	-
2002	142	...	...	-	43	60	37	2	-	-	-	-	-
2001	138	...	...	-	39	60	36	3	-	-	-	-	-
2000	121	...	...	-	30	56	32	3	-	-	-	-	-
1999	144	...	...	-	43	62	37	2	-	-	-	-	-
Non-Hispanic black, male													
2005 2/	...	...	...	...	...	...	...	...	...	...	...	...	...
2004 2/	...	...	...	...	...	...	...	...	...	...	...	...	...
2003 2/	...	...	...	...	...	...	...	...	...	...	...	...	...
2002	...	...	...	...	...	...	...	...	...	...	...	...	...
2001	...	...	...	...	...	...	...	...	...	...	...	...	...
2000	...	...	...	...	...	...	...	...	...	...	...	...	...
1999	...	...	...	...	...	...	...	...	...	...	...	...	...
Non-Hispanic black, female													
2005 2/	252	...	...	-	68	107	61	15	1	-	-	-	-
2004 2/	226	...	...	-	65	100	39	22	-	-	-	-	-
2003 2/	172	...	...	1	34	77	54	6	-	-	-	-	-
2002	142	...	...	-	43	60	37	2	-	-	-	-	-
2001	138	...	...	-	39	60	36	3	-	-	-	-	-
2000	121	...	...	-	30	56	32	3	-	-	-	-	-
1999	144	...	...	-	43	62	37	2	-	-	-	-	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Other complications of pregnancy, childbirth and the puerperium (O10-O99)													
Origin not stated 3/, both sexes													
2005	1	...	...	-	-	-	-	1	-	-	-	-	-
2004	1	...	...	-	-	-	1	-	-	-	-	-	-
2003	4	...	...	-	3	1	-	-	-	-	-	-	-
2002	3	...	...	-	1	2	-	-	-	-	-	-	-
2001	1	...	...	-	-	1	-	-	-	-	-	-	-
2000	-	...	...	-	-	-	-	-	-	-	-	-	-
1999	2	...	...	-	1	1	-	-	-	-	-	-	-
Origin not stated 3/, male													
2005	...	...	...	...	...	...	...	...	...	...	...	...	...
2004	...	...	...	...	...	...	...	...	...	...	...	...	...
2003	...	...	...	...	...	...	...	...	...	...	...	...	...
2002	...	...	...	...	...	...	...	...	...	...	...	...	...
2001	...	...	...	...	...	...	...	...	...	...	...	...	...
2000	...	...	...	...	...	...	...	...	...	...	...	...	...
1999	...	...	...	...	...	...	...	...	...	...	...	...	...
Origin not stated 3/, female													
2005	1	...	...	-	-	-	-	1	-	-	-	-	-
2004	1	...	...	-	-	-	1	-	-	-	-	-	-
2003	4	...	...	-	3	1	-	-	-	-	-	-	-
2002	3	...	...	-	1	2	-	-	-	-	-	-	-
2001	1	...	...	-	-	1	-	-	-	-	-	-	-
2000	-	...	...	-	-	-	-	-	-	-	-	-	-
1999	2	...	...	-	1	1	-	-	-	-	-	-	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Certain conditions originating in the perinatal period (P00-P96)													
All origins, both sexes													
2005	14,549	14,423	58	26	18	6	6	6	2	-	1	-	3
2004	14,213	14,079	61	37	11	10	6	3	3	1	1	1	-
2003	14,378	14,254	79	19	16	3	3	2	-	2	-	-	-
2002	14,254	14,106	65	31	21	10	10	2	1	3	2	1	2
2001	13,887	13,734	72	36	17	6	9	5	2	3	1	1	1
2000	14,069	13,918	79	41	15	7	4	2	2	-	-	-	1
1999	14,259	14,084	92	43	16	7	3	2	3	2	1	1	5
All origins, male													
2005	8,187	8,113	36	11	12	3	4	5	1	-	1	-	1
2004	8,093	8,013	38	23	5	3	5	1	3	1	1	-	-
2003	8,197	8,129	43	11	8	2	1	1	-	2	-	-	-
2002	8,047	7,960	40	19	12	4	5	2	1	1	1	-	2
2001	7,868	7,775	44	24	7	2	6	5	1	2	1	-	1
2000	7,968	7,872	52	23	11	6	2	2	-	-	-	-	-
1999	8,044	7,941	56	23	10	2	2	2	1	1	1	1	4
All origins, female													
2005	6,362	6,310	22	15	6	3	2	1	1	-	-	-	2
2004	6,120	6,066	23	14	6	7	1	2	-	-	-	1	-
2003	6,181	6,125	36	8	8	1	2	1	-	-	-	-	-
2002	6,207	6,146	25	12	9	6	5	-	-	2	1	1	-
2001	6,019	5,959	28	12	10	4	3	-	1	1	-	1	-
2000	6,101	6,046	27	18	4	1	2	-	2	-	-	-	1
1999	6,215	6,143	36	20	6	5	1	-	2	1	-	-	1

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Certain conditions originating in the perinatal period (P00-P96)													
Hispanic, both sexes													
2005	2,816	2,802	6	3	4	1	-	-	-	-	-	-	-
2004	2,681	2,660	10	8	2	-	-	1	-	-	-	-	-
2003	2,628	2,599	15	8	3	1	1	-	-	1	-	-	-
2002	2,402	2,375	14	6	3	1	2	-	-	-	-	1	-
2001	2,227	2,203	14	3	2	4	-	1	-	-	-	-	-
2000	2,145	2,120	12	10	2	-	1	-	-	-	-	-	-
1999	2,153	2,127	18	2	4	1	-	-	-	-	-	-	1
Hispanic, male													
2005	1,612	1,604	2	2	3	1	-	-	-	-	-	-	-
2004	1,538	1,527	6	5	-	-	-	-	-	-	-	-	-
2003	1,459	1,442	8	4	2	1	1	-	-	1	-	-	-
2002	1,352	1,335	9	5	2	1	-	-	-	-	-	-	-
2001	1,268	1,258	6	2	-	1	-	1	-	-	-	-	-
2000	1,194	1,180	7	6	1	-	-	-	-	-	-	-	-
1999	1,192	1,181	5	2	3	-	-	-	-	-	-	-	1
Hispanic, female													
2005	1,204	1,198	4	1	1	-	-	-	-	-	-	-	-
2004	1,143	1,133	4	3	2	-	-	1	-	-	-	-	-
2003	1,169	1,157	7	4	1	-	-	-	-	-	-	-	-
2002	1,050	1,040	5	1	1	-	2	-	-	-	-	1	-
2001	959	945	8	1	2	3	-	-	-	-	-	-	-
2000	951	940	5	4	1	-	1	-	-	-	-	-	-
1999	961	946	13	-	1	1	-	-	-	-	-	-	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Certain conditions originating in the perinatal period (P00-P96)													
Non-Hispanic 1/, both sexes													
2005	11,594	11,482	52	23	14	5	6	6	2	-	1	-	3
2004	11,371	11,258	51	29	9	10	6	2	3	1	1	1	-
2003	11,559	11,464	64	11	13	2	2	2	-	1	-	-	-
2002	11,673	11,553	50	25	18	9	8	2	1	3	2	-	2
2001	11,500	11,371	58	33	15	2	9	4	2	3	1	1	1
2000	11,672	11,546	67	31	13	7	3	2	2	-	-	-	1
1999	11,815	11,668	73	41	12	6	3	2	3	2	1	1	3
Non-Hispanic 1/, male													
2005	6,503	6,437	34	9	9	2	4	5	1	-	1	-	1
2004	6,469	6,400	32	18	5	3	5	1	3	1	1	-	-
2003	6,622	6,571	35	7	6	1	-	1	-	1	-	-	-
2002	6,596	6,527	30	14	10	3	5	2	1	1	1	-	2
2001	6,511	6,428	38	22	7	1	6	4	1	2	1	-	1
2000	6,637	6,555	45	17	10	6	2	2	-	-	-	-	-
1999	6,680	6,589	50	21	7	2	2	2	1	1	1	1	3
Non-Hispanic 1/, female													
2005	5,091	5,045	18	14	5	3	2	1	1	-	-	-	2
2004	4,902	4,858	19	11	4	7	1	1	-	-	-	1	-
2003	4,937	4,893	29	4	7	1	2	1	-	-	-	-	-
2002	5,077	5,026	20	11	8	6	3	-	-	2	1	-	-
2001	4,989	4,943	20	11	8	1	3	-	1	1	-	1	-
2000	5,035	4,991	22	14	3	1	1	-	2	-	-	-	1
1999	5,135	5,079	23	20	5	4	1	-	2	1	-	-	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Certain conditions originating in the perinatal period (P00-P96)													
Non-Hispanic white, both sexes													
2005 2/	6,216	6,138	29	19	12	4	5	6	2	-	1	-	-
2004 2/	6,118	6,049	24	18	8	8	4	2	3	-	1	1	-
2003 2/	6,359	6,303	31	8	12	2	2	1	-	-	-	-	-
2002	6,384	6,313	22	15	14	6	6	2	1	2	2	-	1
2001	6,323	6,239	34	18	11	2	8	4	1	3	1	1	1
2000	6,318	6,242	34	20	11	3	3	2	2	-	-	-	1
1999	6,384	6,286	37	31	11	6	3	2	3	2	1	1	1
Non-Hispanic white, male													
2005 2/	3,495	3,448	18	9	8	2	3	5	1	-	1	-	-
2004 2/	3,529	3,485	17	11	5	3	3	1	3	-	1	-	-
2003 2/	3,673	3,643	17	6	6	1	-	-	-	-	-	-	-
2002	3,663	3,628	10	7	7	3	3	2	1	-	1	-	1
2001	3,574	3,518	22	14	6	1	5	4	-	2	1	-	1
2000	3,605	3,553	23	14	9	2	2	2	-	-	-	-	-
1999	3,627	3,570	24	15	7	2	2	2	1	1	1	1	1
Non-Hispanic white, female													
2005 2/	2,721	2,690	11	10	4	2	2	1	1	-	-	-	-
2004 2/	2,589	2,564	7	7	3	5	1	1	-	-	-	1	-
2003 2/	2,686	2,660	14	2	6	1	2	1	-	-	-	-	-
2002	2,721	2,685	12	8	7	3	3	-	-	2	1	-	-
2001	2,749	2,721	12	4	5	1	3	-	1	1	-	1	-
2000	2,713	2,689	11	6	2	1	1	-	2	-	-	-	1
1999	2,757	2,716	13	16	4	4	1	-	2	1	-	-	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Certain conditions originating in the perinatal period (P00-P96)													
Non-Hispanic black, both sexes													
2005 2/	4,837	4,806	23	3	2	-	1	-	-	-	-	-	2
2004 2/	4,740	4,703	23	9	1	2	2	-	-	-	-	-	-
2003 2/	4,716	4,682	29	2	1	-	-	1	-	1	-	-	-
2002	4,777	4,735	26	8	3	2	1	-	-	1	-	-	1
2001	4,713	4,675	22	11	4	-	1	-	-	-	-	-	-
2000	4,878	4,836	27	10	1	4	-	-	-	-	-	-	-
1999	5,000	4,963	29	5	1	-	-	-	-	-	-	-	2
Non-Hispanic black, male													
2005 2/	2,712	2,693	16	-	1	-	1	-	-	-	-	-	1
2004 2/	2,653	2,633	13	5	-	-	2	-	-	-	-	-	-
2003 2/	2,687	2,668	16	1	-	-	-	1	-	1	-	-	-
2002	2,640	2,611	18	6	2	-	1	-	-	1	-	-	1
2001	2,674	2,653	14	5	1	-	1	-	-	-	-	-	-
2000	2,760	2,734	18	3	1	4	-	-	-	-	-	-	-
1999	2,818	2,794	19	3	-	-	-	-	-	-	-	-	2
Non-Hispanic black, female													
2005 2/	2,125	2,113	7	3	1	-	-	-	-	-	-	-	1
2004 2/	2,087	2,070	10	4	1	2	-	-	-	-	-	-	-
2003 2/	2,029	2,014	13	1	1	-	-	-	-	-	-	-	-
2002	2,137	2,124	8	2	1	2	-	-	-	-	-	-	-
2001	2,039	2,022	8	6	3	-	-	-	-	-	-	-	-
2000	2,118	2,102	9	7	-	-	-	-	-	-	-	-	-
1999	2,182	2,169	10	2	1	-	-	-	-	-	-	-	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Certain conditions originating in the perinatal period (P00-P96)													
Origin not stated 3/, both sexes													
2005	139	139	-	-	-	-	-	-	-	-	-	-	-
2004	161	161	-	-	-	-	-	-	-	-	-	-	-
2003	191	191	-	-	-	-	-	-	-	-	-	-	-
2002	179	178	1	-	-	-	-	-	-	-	-	-	-
2001	160	160	-	-	-	-	-	-	-	-	-	-	-
2000	252	252	-	-	-	-	-	-	-	-	-	-	-
1999	291	289	1	-	-	-	-	-	-	-	-	-	1
Origin not stated 3/, male													
2005	72	72	-	-	-	-	-	-	-	-	-	-	-
2004	86	86	-	-	-	-	-	-	-	-	-	-	-
2003	116	116	-	-	-	-	-	-	-	-	-	-	-
2002	99	98	1	-	-	-	-	-	-	-	-	-	-
2001	89	89	-	-	-	-	-	-	-	-	-	-	-
2000	137	137	-	-	-	-	-	-	-	-	-	-	-
1999	172	171	1	-	-	-	-	-	-	-	-	-	-
Origin not stated 3/, female													
2005	67	67	-	-	-	-	-	-	-	-	-	-	-
2004	75	75	-	-	-	-	-	-	-	-	-	-	-
2003	75	75	-	-	-	-	-	-	-	-	-	-	-
2002	80	80	-	-	-	-	-	-	-	-	-	-	-
2001	71	71	-	-	-	-	-	-	-	-	-	-	-
2000	115	115	-	-	-	-	-	-	-	-	-	-	-
1999	119	118	-	-	-	-	-	-	-	-	-	-	1

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)													
All origins, both sexes													
2005	10,410	5,552	522	396	504	436	517	693	716	338	408	327	1
2004	10,498	5,622	569	389	483	420	493	692	634	428	472	296	-
2003	10,518	5,621	541	386	451	426	564	667	666	413	464	319	-
2002	10,687	5,623	530	417	492	475	572	709	634	440	489	305	1
2001	10,442	5,513	557	376	505	458	586	636	581	448	465	317	-
2000	10,578	5,743	495	399	441	477	569	612	569	464	488	320	1
1999	10,393	5,473	549	428	434	465	552	647	557	469	491	326	2
All origins, male													
2005	5,447	2,965	264	195	326	244	272	356	370	153	191	111	-
2004	5,423	2,898	318	211	282	235	279	375	320	208	195	102	-
2003	5,460	3,020	275	198	272	233	304	349	336	198	178	97	-
2002	5,515	2,942	282	215	282	284	296	381	321	213	201	97	1
2001	5,448	2,954	282	210	306	249	317	323	276	228	203	100	-
2000	5,517	3,028	259	207	268	279	316	347	264	222	197	130	-
1999	5,346	2,875	310	215	262	242	296	347	283	209	204	102	1
All origins, female													
2005	4,963	2,587	258	201	178	192	245	337	346	185	217	216	1
2004	5,075	2,724	251	178	201	185	214	317	314	220	277	194	-
2003	5,058	2,601	266	188	179	193	260	318	330	215	286	222	-
2002	5,172	2,681	248	202	210	191	276	328	313	227	288	208	-
2001	4,994	2,559	275	166	199	209	269	313	305	220	262	217	-
2000	5,061	2,715	236	192	173	198	253	265	305	242	291	190	1
1999	5,047	2,598	239	213	172	223	256	300	274	260	287	224	1

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)													
Hispanic, both sexes													
2005	1,896	1,382	119	85	74	52	47	51	41	17	23	5	-
2004	1,765	1,278	138	70	61	53	40	44	32	20	23	6	-
2003	1,787	1,324	122	65	57	58	44	38	35	17	13	14	-
2002	1,715	1,257	111	70	67	53	40	42	25	18	23	9	-
2001	1,632	1,195	103	58	73	45	42	36	24	22	24	10	-
2000	1,612	1,182	111	52	54	46	43	36	31	21	24	12	-
1999	1,498	1,072	101	57	56	54	34	42	27	30	16	9	-
Hispanic, male													
2005	996	720	65	47	43	30	26	24	23	6	11	1	-
2004	902	622	82	49	40	24	22	27	16	8	10	2	-
2003	937	693	62	28	30	37	25	22	19	10	6	5	-
2002	903	653	60	35	39	36	19	23	15	6	13	4	-
2001	892	639	54	36	39	26	27	25	14	14	13	5	-
2000	857	605	60	28	39	26	27	22	17	11	15	7	-
1999	756	534	63	29	30	25	15	24	9	17	7	3	-
Hispanic, female													
2005	900	662	54	38	31	22	21	27	18	11	12	4	-
2004	863	656	56	21	21	29	18	17	16	12	13	4	-
2003	850	631	60	37	27	21	19	16	16	7	7	9	-
2002	812	604	51	35	28	17	21	19	10	12	10	5	-
2001	740	556	49	22	34	19	15	11	10	8	11	5	-
2000	755	577	51	24	15	20	16	14	14	10	9	5	-
1999	742	538	38	28	26	29	19	18	18	13	9	6	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)													
Non-Hispanic 1/, both sexes													
2005	8,479	4,139	403	310	430	384	470	640	675	321	384	322	1
2004	8,686	4,309	430	316	421	366	452	647	599	408	449	289	-
2003	8,687	4,269	418	317	393	367	516	628	629	395	451	304	-
2002	8,904	4,316	415	347	422	422	530	662	608	421	465	295	1
2001	8,748	4,271	452	317	432	412	542	596	557	422	441	306	-
2000	8,900	4,514	384	345	386	427	525	574	532	443	462	307	1
1999	8,825	4,349	447	370	378	410	516	601	529	435	473	315	2
Non-Hispanic 1/, male													
2005	4,431	2,228	199	147	283	214	246	331	347	147	179	110	-
2004	4,493	2,256	235	159	242	210	256	347	303	200	185	100	-
2003	4,500	2,312	213	167	241	196	276	327	317	187	172	92	-
2002	4,577	2,262	221	180	241	248	276	355	306	206	188	93	1
2001	4,527	2,297	227	173	267	222	289	294	262	211	190	95	-
2000	4,623	2,400	199	177	229	250	288	323	243	211	181	122	-
1999	4,550	2,311	246	185	232	217	280	320	274	191	196	97	1
Non-Hispanic 1/, female													
2005	4,048	1,911	204	163	147	170	224	309	328	174	205	212	1
2004	4,193	2,053	195	157	179	156	196	300	296	208	264	189	-
2003	4,187	1,957	205	150	152	171	240	301	312	208	279	212	-
2002	4,327	2,054	194	167	181	174	254	307	302	215	277	202	-
2001	4,221	1,974	225	144	165	190	253	302	295	211	251	211	-
2000	4,277	2,114	185	168	157	177	237	251	289	232	281	185	1
1999	4,275	2,038	201	185	146	193	236	281	255	244	277	218	1

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)													
Non-Hispanic white, both sexes													
2005 2/	6,412	2,848	270	207	313	287	381	553	605	299	355	293	1
2004 2/	6,598	3,016	270	217	290	295	369	555	540	373	401	272	-
2003 2/	6,594	2,977	272	222	285	285	430	530	560	342	411	280	-
2002	6,830	3,068	281	253	315	331	423	562	527	374	420	275	1
2001	6,709	3,053	287	226	302	330	439	503	503	378	405	283	-
2000	6,792	3,199	240	240	279	344	419	501	460	390	431	288	1
1999	6,782	3,127	284	244	277	334	426	514	473	383	427	291	2
Non-Hispanic white, male													
2005 2/	3,352	1,540	139	101	210	163	195	287	310	140	169	98	-
2004 2/	3,375	1,572	145	111	156	168	205	303	271	184	164	96	-
2003 2/	3,353	1,596	136	117	175	152	233	271	279	158	151	85	-
2002	3,510	1,643	151	125	178	195	223	295	263	183	165	88	1
2001	3,487	1,664	144	135	183	186	236	251	235	196	174	83	-
2000	3,519	1,720	124	119	163	203	233	276	217	177	172	115	-
1999	3,498	1,693	154	119	175	178	230	271	248	169	171	89	1
Non-Hispanic white, female													
2005 2/	3,060	1,308	131	106	103	124	186	266	295	159	186	195	1
2004 2/	3,223	1,444	125	106	134	127	164	252	269	189	237	176	-
2003 2/	3,241	1,381	136	105	110	133	197	259	281	184	260	195	-
2002	3,320	1,425	130	128	137	136	200	267	264	191	255	187	-
2001	3,222	1,389	143	91	119	144	203	252	268	182	231	200	-
2000	3,273	1,479	116	121	116	141	186	225	243	213	259	173	1
1999	3,284	1,434	130	125	102	156	196	243	225	214	256	202	1

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)													
Non-Hispanic black, both sexes													
2005 2/	1,656	1,029	104	80	99	81	74	71	59	16	21	22	-
2004 2/	1,697	1,041	127	82	111	59	69	82	47	27	39	13	-
2003 2/	1,671	1,015	118	73	92	75	69	79	60	40	30	20	-
2002	1,716	1,015	113	81	98	75	92	83	69	40	32	18	-
2001	1,674	999	127	73	113	67	89	76	48	35	28	19	-
2000	1,739	1,062	123	89	91	74	91	67	58	43	25	16	-
1999	1,680	994	137	105	82	60	76	73	46	46	38	23	-
Non-Hispanic black, male													
2005 2/	856	537	46	37	63	43	43	33	31	5	9	9	-
2004 2/	914	551	74	38	73	36	42	39	25	14	18	4	-
2003 2/	905	558	62	38	55	38	32	42	33	25	16	6	-
2002	887	506	58	47	59	43	45	47	40	21	16	5	-
2001	861	525	65	33	72	30	46	34	23	12	12	9	-
2000	911	545	67	50	58	43	49	41	17	26	9	6	-
1999	880	518	79	53	50	32	41	42	22	16	19	8	-
Non-Hispanic black, female													
2005 2/	800	492	58	43	36	38	31	38	28	11	12	13	-
2004 2/	783	490	53	44	38	23	27	43	22	13	21	9	-
2003 2/	766	457	56	35	37	37	37	37	27	15	14	14	-
2002	829	509	55	34	39	32	47	36	29	19	16	13	-
2001	813	474	62	40	41	37	43	42	25	23	16	10	-
2000	828	517	56	39	33	31	42	26	41	17	16	10	-
1999	800	476	58	52	32	28	35	31	24	30	19	15	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)													
Origin not stated 3/, both sexes													
2005	35	31	-	1	-	-	-	2	-	-	1	-	-
2004	47	35	1	3	1	1	1	1	3	-	-	1	-
2003	44	28	1	4	1	1	4	1	2	1	-	1	-
2002	68	50	4	-	3	-	2	5	1	1	1	1	-
2001	62	47	2	1	-	1	2	4	-	4	-	1	-
2000	66	47	-	2	1	4	1	2	6	-	2	1	-
1999	70	52	1	1	-	1	2	4	1	4	2	2	-
Origin not stated 3/, male													
2005	20	17	-	1	-	-	-	1	-	-	1	-	-
2004	28	20	1	3	-	1	1	1	1	-	-	-	-
2003	23	15	-	3	1	-	3	-	-	1	-	-	-
2002	35	27	1	-	2	-	1	3	-	1	-	-	-
2001	29	18	1	1	-	1	1	4	-	3	-	-	-
2000	37	23	-	2	-	3	1	2	4	-	1	1	-
1999	40	30	1	1	-	-	1	3	-	1	1	2	-
Origin not stated 3/, female													
2005	15	14	-	-	-	-	-	1	-	-	-	-	-
2004	19	15	-	-	1	-	-	-	2	-	-	1	-
2003	21	13	1	1	-	1	1	1	2	-	-	1	-
2002	33	23	3	-	1	-	1	2	1	-	1	1	-
2001	33	29	1	-	-	-	1	-	-	1	-	1	-
2000	29	24	-	-	1	1	-	-	2	-	1	-	-
1999	30	22	-	-	-	1	1	1	1	3	1	-	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Symptoms, signs and abnormal clinical and laboratory findings, not elsewhere classified (R00-R99)													
All origins, both sexes													
2005	31,999	3,589	233	155	775	1,178	2,052	2,831	2,199	2,390	5,088	11,436	73
2004	30,126	3,361	189	120	658	1,092	1,900	2,402	2,041	2,323	5,159	10,762	119
2003	31,444	3,318	256	178	837	1,327	2,382	2,874	2,067	2,522	5,006	10,565	112
2002	29,975	3,456	248	139	740	1,319	2,402	2,672	2,002	2,398	4,936	9,554	109
2001	32,326	3,491	272	181	949	1,530	3,126	3,504	2,292	2,774	5,003	9,089	115
2000	31,876	3,639	248	177	868	1,554	2,987	3,142	2,253	3,040	5,029	8,824	115
1999	26,672	3,538	236	137	609	1,095	2,062	2,151	1,860	2,698	4,413	7,798	75
All origins, male													
2005	14,647	2,139	132	94	546	807	1,263	1,822	1,367	1,321	2,150	2,949	57
2004	13,712	2,021	100	59	452	732	1,235	1,546	1,268	1,264	2,181	2,768	86
2003	14,583	1,974	143	92	563	907	1,526	1,840	1,294	1,431	2,160	2,570	83
2002	14,269	2,032	145	82	501	905	1,540	1,723	1,273	1,324	2,228	2,439	77
2001	15,779	2,041	162	99	654	1,040	2,007	2,286	1,457	1,497	2,184	2,274	78
2000	15,863	2,156	137	99	581	1,060	1,994	2,123	1,429	1,708	2,244	2,242	90
1999	12,877	2,072	136	77	422	721	1,339	1,446	1,181	1,513	1,930	1,984	56
All origins, female													
2005	17,352	1,450	101	61	229	371	789	1,009	832	1,069	2,938	8,487	16
2004	16,414	1,340	89	61	206	360	665	856	773	1,059	2,978	7,994	33
2003	16,861	1,344	113	86	274	420	856	1,034	773	1,091	2,846	7,995	29
2002	15,706	1,424	103	57	239	414	862	949	729	1,074	2,708	7,115	32
2001	16,547	1,450	110	82	295	490	1,119	1,218	835	1,277	2,819	6,815	37
2000	16,013	1,483	111	78	287	494	993	1,019	824	1,332	2,785	6,582	25
1999	13,795	1,466	100	60	187	374	723	705	679	1,185	2,483	5,814	19

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Symptoms, signs and abnormal clinical and laboratory findings, not elsewhere classified (R00-R99)													
Hispanic, both sexes													
2005	1,857	555	44	27	108	184	201	153	86	103	136	246	14
2004	1,621	457	37	18	115	161	192	147	87	79	112	209	7
2003	1,623	492	30	20	98	170	181	141	73	75	118	214	11
2002	1,548	472	40	11	91	149	177	144	95	83	112	165	9
2001	2,190	496	47	36	184	286	327	283	127	112	107	173	12
2000	2,071	506	51	31	140	260	342	251	114	107	104	154	11
1999	1,281	415	30	16	83	138	148	93	63	73	93	124	5
Hispanic, male													
2005	1,135	328	23	17	76	153	146	119	58	60	63	80	12
2004	971	278	21	11	88	116	145	99	61	40	44	62	6
2003	998	309	16	12	65	135	135	108	55	44	49	62	8
2002	926	283	26	7	62	116	120	85	59	50	57	53	8
2001	1,410	280	28	15	139	221	239	212	92	71	56	46	11
2000	1,369	310	31	16	102	201	262	192	72	68	52	53	10
1999	768	237	14	10	64	99	110	68	42	38	46	36	4
Hispanic, female													
2005	722	227	21	10	32	31	55	34	28	43	73	166	2
2004	650	179	16	7	27	45	47	48	26	39	68	147	1
2003	625	183	14	8	33	35	46	33	18	31	69	152	3
2002	622	189	14	4	29	33	57	59	36	33	55	112	1
2001	780	216	19	21	45	65	88	71	35	41	51	127	1
2000	702	196	20	15	38	59	80	59	42	39	52	101	1
1999	513	178	16	6	19	39	38	25	21	35	47	88	1

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Symptoms, signs and abnormal clinical and laboratory findings, not elsewhere classified (R00-R99)													
Non-Hispanic 1/, both sexes													
2005	29,884	2,999	184	125	655	986	1,829	2,638	2,085	2,272	4,922	11,166	23
2004	28,297	2,881	150	101	538	923	1,690	2,230	1,935	2,235	5,028	10,541	45
2003	29,617	2,805	226	157	731	1,151	2,181	2,708	1,975	2,439	4,876	10,324	44
2002	28,211	2,968	207	126	643	1,162	2,206	2,504	1,890	2,307	4,804	9,351	43
2001	29,931	2,978	225	144	757	1,227	2,782	3,193	2,154	2,653	4,886	8,896	36
2000	29,514	3,108	196	144	720	1,279	2,604	2,849	2,109	2,908	4,908	8,654	35
1999	25,181	3,097	204	121	521	951	1,885	2,023	1,780	2,601	4,310	7,658	30
Non-Hispanic 1/, male													
2005	13,353	1,790	107	77	460	649	1,101	1,677	1,293	1,256	2,073	2,856	14
2004	12,607	1,730	78	48	361	609	1,076	1,431	1,192	1,218	2,130	2,701	33
2003	13,448	1,650	127	80	495	767	1,375	1,711	1,225	1,379	2,102	2,499	38
2002	13,206	1,739	119	74	434	783	1,408	1,619	1,198	1,267	2,156	2,376	33
2001	14,246	1,750	134	84	510	807	1,754	2,058	1,355	1,422	2,125	2,221	26
2000	14,288	1,834	106	82	474	848	1,700	1,901	1,331	1,621	2,178	2,183	30
1999	11,962	1,817	121	67	355	619	1,209	1,347	1,125	1,463	1,876	1,944	19
Non-Hispanic 1/, female													
2005	16,531	1,209	77	48	195	337	728	961	792	1,016	2,849	8,310	9
2004	15,690	1,151	72	53	177	314	614	799	743	1,017	2,898	7,840	12
2003	16,169	1,155	99	77	236	384	806	997	750	1,060	2,774	7,825	6
2002	15,005	1,229	88	52	209	379	798	885	692	1,040	2,648	6,975	10
2001	15,685	1,228	91	60	247	420	1,028	1,135	799	1,231	2,761	6,675	10
2000	15,226	1,274	90	62	246	431	904	948	778	1,287	2,730	6,471	5
1999	13,219	1,280	83	54	166	332	676	676	655	1,138	2,434	5,714	11

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Symptoms, signs and abnormal clinical and laboratory findings, not elsewhere classified (R00-R99)													
Non-Hispanic white, both sexes													
2005 2/	24,697	1,804	109	64	479	729	1,400	2,023	1,632	1,850	4,346	10,244	17
2004 2/	23,344	1,654	91	60	404	673	1,281	1,725	1,518	1,841	4,447	9,617	33
2003 2/	24,452	1,665	136	99	562	843	1,661	2,022	1,562	2,013	4,367	9,487	35
2002	23,099	1,777	124	71	468	849	1,664	1,822	1,493	1,934	4,275	8,591	31
2001	24,596	1,752	139	89	563	907	2,175	2,510	1,713	2,191	4,350	8,178	29
2000	24,064	1,805	134	88	518	918	2,027	2,173	1,652	2,394	4,358	7,975	22
1999	20,200	1,812	129	81	348	660	1,341	1,459	1,374	2,166	3,775	7,033	22
Non-Hispanic white, male													
2005 2/	10,691	1,122	66	42	340	491	863	1,304	1,008	1,022	1,823	2,598	12
2004 2/	10,064	984	47	31	275	458	832	1,103	930	1,015	1,893	2,470	26
2003 2/	10,752	1,025	75	51	389	570	1,050	1,297	958	1,143	1,881	2,282	31
2002	10,432	1,056	66	44	326	581	1,071	1,177	945	1,065	1,897	2,180	24
2001	11,404	1,046	79	52	376	614	1,387	1,634	1,058	1,184	1,920	2,032	22
2000	11,351	1,067	79	52	356	625	1,366	1,465	1,036	1,328	1,945	2,013	19
1999	9,249	1,083	82	45	244	431	884	963	870	1,222	1,638	1,771	16
Non-Hispanic white, female													
2005 2/	14,006	682	43	22	139	238	537	719	624	828	2,523	7,646	5
2004 2/	13,280	670	44	29	129	215	449	622	588	826	2,554	7,147	7
2003 2/	13,700	640	61	48	173	273	611	725	604	870	2,486	7,205	4
2002	12,667	721	58	27	142	268	593	645	548	869	2,378	6,411	7
2001	13,192	706	60	37	187	293	788	876	655	1,007	2,430	6,146	7
2000	12,713	738	55	36	162	293	661	708	616	1,066	2,413	5,962	3
1999	10,951	729	47	36	104	229	457	496	504	944	2,137	5,262	6

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Symptoms, signs and abnormal clinical and laboratory findings, not elsewhere classified (R00-R99)													
Non-Hispanic black, both sexes													
2005 2/	4,612	1,073	62	54	156	222	381	572	414	380	511	781	6
2004 2/	4,376	1,106	49	35	111	216	364	465	376	351	516	776	11
2003 2/	4,628	1,022	80	49	150	267	476	632	385	400	439	721	7
2002	4,556	1,070	66	50	152	275	490	626	356	335	471	655	10
2001	4,606	1,081	69	42	158	266	518	605	373	401	466	622	5
2000	4,892	1,186	54	46	179	308	514	609	411	472	505	595	13
1999	4,508	1,149	67	33	152	260	504	520	371	395	488	563	6
Non-Hispanic black, male													
2005 2/	2,366	601	34	30	109	136	209	345	260	211	223	206	2
2004 2/	2,244	673	28	16	72	132	210	304	237	178	208	180	6
2003 2/	2,404	554	44	25	92	173	290	383	247	223	183	185	5
2002	2,482	615	44	28	92	174	305	407	229	180	234	167	7
2001	2,414	614	44	26	109	150	300	364	262	207	174	161	3
2000	2,614	694	26	28	105	191	283	390	266	267	214	139	11
1999	2,460	658	35	20	100	172	299	352	238	218	214	152	2
Non-Hispanic black, female													
2005 2/	2,246	472	28	24	47	86	172	227	154	169	288	575	4
2004 2/	2,132	433	21	19	39	84	154	161	139	173	308	596	5
2003 2/	2,224	468	36	24	58	94	186	249	138	177	256	536	2
2002	2,074	455	22	22	60	101	185	219	127	155	237	488	3
2001	2,192	467	25	16	49	116	218	241	111	194	292	461	2
2000	2,278	492	28	18	74	117	231	219	145	205	291	456	2
1999	2,048	491	32	13	52	88	205	168	133	177	274	411	4

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Symptoms, signs and abnormal clinical and laboratory findings, not elsewhere classified (R00-R99)													
Origin not stated 3/, both sexes													
2005	258	35	5	3	12	8	22	40	28	15	30	24	36
2004	208	23	2	1	5	8	18	25	19	9	19	12	67
2003	204	21	-	1	8	6	20	25	19	8	12	27	57
2002	216	16	1	2	6	8	19	24	17	8	20	38	57
2001	205	17	-	1	8	17	17	28	11	9	10	20	67
2000	291	25	1	2	8	15	41	42	30	25	17	16	69
1999	210	26	2	-	5	6	29	35	17	24	10	16	40
Origin not stated 3/, male													
2005	159	21	2	-	10	5	16	26	16	5	14	13	31
2004	134	13	1	-	3	7	14	16	15	6	7	5	47
2003	137	15	-	-	3	5	16	21	14	8	9	9	37
2002	137	10	-	1	5	6	12	19	16	7	15	10	36
2001	123	11	-	-	5	12	14	16	10	4	3	7	41
2000	206	12	-	1	5	11	32	30	26	19	14	6	50
1999	147	18	1	-	3	3	20	31	14	12	8	4	33
Origin not stated 3/, female													
2005	99	14	3	3	2	3	6	14	12	10	16	11	5
2004	74	10	1	1	2	1	4	9	4	3	12	7	20
2003	67	6	-	1	5	1	4	4	5	-	3	18	20
2002	79	6	1	1	1	2	7	5	1	1	5	28	21
2001	82	6	-	1	3	5	3	12	1	5	7	13	26
2000	85	13	1	1	3	4	9	12	4	6	3	10	19
1999	63	8	1	-	2	3	9	4	3	12	2	12	7

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
All other diseases (Residual) (D65-E07,E15-E34,E65-F99,G04-G12,G23-G25,G31-H93,K00-K22,K29-K31,K50-K66,K71-K72,K75-K76,K83-M99,N13.0-N13.5,N13.7-N13.9,N14, N15.0,N15.8-N15.9,N20-N23,N28-N39,N41-N64,N80-N98)													
All origins, both sexes													
2005	217,632	1,198	612	857	2,060	3,159	8,118	17,122	21,108	27,236	56,700	79,449	13
2004	201,914	1,192	611	893	1,956	3,232	8,072	16,205	19,486	26,104	53,123	71,028	12
2003	201,676	1,190	635	927	2,023	3,330	8,407	15,883	18,734	26,475	52,955	71,100	17
2002	194,591	1,144	661	868	1,853	3,136	8,506	15,151	17,378	25,689	51,618	68,574	13
2001	186,406	1,131	646	860	1,799	3,072	8,084	14,203	16,048	25,803	49,711	65,032	17
2000	178,378	1,105	609	879	1,764	3,097	8,113	13,326	15,156	25,007	48,019	61,292	11
1999	171,683	1,148	647	921	1,771	3,062	7,989	12,383	14,390	25,165	45,598	58,599	10
All origins, male													
2005	88,869	732	352	474	1,245	1,821	4,837	10,253	11,828	13,311	23,071	20,934	11
2004	82,725	704	328	490	1,158	1,861	4,739	9,767	10,796	12,805	21,422	18,646	9
2003	81,907	711	342	541	1,189	1,885	4,932	9,456	10,274	12,993	21,393	18,178	13
2002	79,007	690	363	486	1,072	1,837	5,019	9,115	9,528	12,667	20,667	17,552	11
2001	75,162	649	358	454	1,064	1,729	4,774	8,397	8,652	12,732	19,817	16,520	16
2000	72,227	629	338	474	1,057	1,811	4,801	7,911	8,236	12,321	19,102	15,536	11
1999	69,784	680	354	483	1,045	1,790	4,835	7,272	7,870	12,332	18,495	14,623	5
All origins, female													
2005	128,763	466	260	383	815	1,338	3,281	6,869	9,280	13,925	33,629	58,515	2
2004	119,189	488	283	403	798	1,371	3,333	6,438	8,690	13,299	31,701	52,382	3
2003	119,769	479	293	386	834	1,445	3,475	6,427	8,460	13,482	31,562	52,922	4
2002	115,584	454	298	382	781	1,299	3,487	6,036	7,850	13,022	30,951	51,022	2
2001	111,244	482	288	406	735	1,343	3,310	5,806	7,396	13,071	29,894	48,512	1
2000	106,151	476	271	405	707	1,286	3,312	5,415	6,920	12,686	28,917	45,756	-
1999	101,899	468	293	438	726	1,272	3,154	5,111	6,520	12,833	27,103	43,976	5

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
All other diseases (Residual) (D65-E07,E15-E34,E65-F99,G04-G12,G23-G25,G31-H93,K00-K22,K29-K31,K50-K66,K71-K72,K75-K76,K83-M99,N13.0-N13.5,N13.7-N13.9,N14, N15.0,N15.8-N15.9,N20-N23,N28-N39,N41-N64,N80-N98)													
Hispanic, both sexes													
2005	10,250	271	146	125	291	480	851	1,209	1,249	1,420	2,120	2,082	6
2004	9,217	237	115	148	260	412	807	1,118	1,090	1,283	1,894	1,851	2
2003	9,218	235	140	159	287	452	829	1,156	1,117	1,306	1,822	1,712	3
2002	8,377	209	120	151	250	409	757	1,006	958	1,313	1,632	1,571	1
2001	7,847	187	112	122	246	354	671	892	918	1,218	1,539	1,585	3
2000	7,325	196	109	113	214	381	706	862	813	1,113	1,438	1,376	4
1999	7,291	200	106	107	218	409	690	837	842	1,133	1,265	1,482	2
Hispanic, male													
2005	5,110	165	78	65	184	309	561	790	738	682	898	634	6
2004	4,541	133	62	81	169	260	556	759	609	620	740	550	2
2003	4,571	144	75	87	165	295	519	734	640	631	787	492	2
2002	4,221	116	66	89	148	261	517	647	527	659	707	483	1
2001	3,849	106	66	65	147	222	451	584	513	587	612	493	3
2000	3,663	100	54	63	140	246	483	552	464	568	557	432	4
1999	3,655	124	50	54	128	275	479	541	481	564	530	428	1
Hispanic, female													
2005	5,140	106	68	60	107	171	290	419	511	738	1,222	1,448	-
2004	4,676	104	53	67	91	152	251	359	481	663	1,154	1,301	-
2003	4,647	91	65	72	122	157	310	422	477	675	1,035	1,220	1
2002	4,156	93	54	62	102	148	240	359	431	654	925	1,088	-
2001	3,998	81	46	57	99	132	220	308	405	631	927	1,092	-
2000	3,662	96	55	50	74	135	223	310	349	545	881	944	-
1999	3,636	76	56	53	90	134	211	296	361	569	735	1,054	1

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
All other diseases (Residual) (D65-E07,E15-E34,E65-F99,G04-G12,G23-G25,G31-H93,K00-K22,K29-K31,K50-K66,K71-K72,K75-K76,K83-M99,N13.0-N13.5,N13.7-N13.9,N14, N15.0,N15.8-N15.9,N20-N23,N28-N39,N41-N64,N80-N98)													
Non-Hispanic 1/, both sexes													
2005	207,024	917	463	731	1,766	2,670	7,247	15,865	19,807	25,770	54,509	77,274	5
2004	192,252	950	494	741	1,691	2,811	7,238	15,018	18,340	24,761	51,148	69,054	6
2003	191,961	946	493	765	1,727	2,865	7,544	14,662	17,558	25,109	51,044	69,241	7
2002	185,608	927	538	715	1,600	2,714	7,711	14,060	16,347	24,307	49,835	66,846	8
2001	177,985	934	532	736	1,545	2,709	7,370	13,230	15,063	24,506	48,059	63,290	11
2000	170,462	898	497	762	1,541	2,697	7,366	12,376	14,283	23,799	46,458	59,780	5
1999	163,832	935	537	814	1,547	2,645	7,247	11,467	13,479	23,956	44,232	56,968	5
Non-Hispanic 1/, male													
2005	83,561	560	271	408	1,060	1,508	4,261	9,427	11,049	12,599	22,139	20,275	4
2004	77,963	567	265	408	986	1,595	4,166	8,959	10,148	12,153	20,649	18,064	3
2003	77,081	561	266	452	1,016	1,580	4,393	8,680	9,595	12,326	20,567	17,640	5
2002	74,500	572	294	396	923	1,568	4,475	8,410	8,953	11,964	19,905	17,033	7
2001	71,014	539	291	388	912	1,499	4,293	7,751	8,087	12,104	19,155	15,985	10
2000	68,263	522	282	411	911	1,556	4,287	7,294	7,734	11,700	18,493	15,068	5
1999	65,835	546	302	429	913	1,509	4,316	6,676	7,341	11,723	17,924	14,153	3
Non-Hispanic 1/, female													
2005	123,463	357	192	323	706	1,162	2,986	6,438	8,758	13,171	32,370	56,999	1
2004	114,289	383	229	333	705	1,216	3,072	6,059	8,192	12,608	30,499	50,990	3
2003	114,880	385	227	313	711	1,285	3,151	5,982	7,963	12,783	30,477	51,601	2
2002	111,108	355	244	319	677	1,146	3,236	5,650	7,394	12,343	29,930	49,813	1
2001	106,971	395	241	348	633	1,210	3,077	5,479	6,976	12,402	28,904	47,305	1
2000	102,199	376	215	351	630	1,141	3,079	5,082	6,549	12,099	27,965	44,712	-
1999	97,997	389	235	385	634	1,136	2,931	4,791	6,138	12,233	26,308	42,815	2

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
All other diseases (Residual) (D65-E07,E15-E34,E65-F99,G04-G12,G23-G25,G31-H93,K00-K22,K29-K31,K50-K66,K71-K72,K75-K76,K83-M99,N13.0-N13.5,N13.7-N13.9,N14, N15.0,N15.8-N15.9,N20-N23,N28-N39,N41-N64,N80-N98)													
Non-Hispanic white, both sexes													
2005 2/	178,181	536	305	478	1,275	1,815	5,238	11,976	15,959	21,617	48,408	70,571	3
2004 2/	165,312	573	322	516	1,188	1,843	5,211	11,337	14,723	20,787	45,589	63,218	5
2003 2/	165,261	565	324	531	1,245	1,910	5,404	10,965	14,071	21,224	45,510	63,507	5
2002	159,945	578	352	505	1,102	1,831	5,545	10,567	13,116	20,530	44,489	61,325	5
2001	153,156	556	362	541	1,089	1,808	5,201	9,814	11,982	20,807	42,990	58,000	6
2000	146,518	544	316	537	1,113	1,807	5,202	9,186	11,332	20,188	41,562	54,729	2
1999	141,002	554	375	569	1,100	1,845	5,036	8,437	10,749	20,267	39,663	52,406	1
Non-Hispanic white, male													
2005 2/	71,058	328	180	275	797	1,086	3,238	7,332	8,987	10,621	19,668	18,544	2
2004 2/	66,039	350	176	296	722	1,102	3,156	6,942	8,221	10,233	18,367	16,472	2
2003 2/	65,306	357	168	319	788	1,119	3,280	6,649	7,742	10,416	18,274	16,191	3
2002	63,109	376	191	279	677	1,113	3,386	6,464	7,230	10,109	17,748	15,531	5
2001	60,022	311	209	286	677	1,040	3,130	5,873	6,476	10,312	17,140	14,563	5
2000	57,565	310	183	307	683	1,105	3,128	5,569	6,149	9,933	16,519	13,677	2
1999	55,517	321	214	291	681	1,110	3,144	5,029	5,906	9,977	16,013	12,830	1
Non-Hispanic white, female													
2005 2/	107,123	208	125	203	478	729	2,000	4,644	6,972	10,996	28,740	52,027	1
2004 2/	99,273	223	146	220	466	741	2,055	4,395	6,502	10,554	27,222	46,746	3
2003 2/	99,955	208	156	212	457	791	2,124	4,316	6,329	10,808	27,236	47,316	2
2002	96,836	202	161	226	425	718	2,159	4,103	5,886	10,421	26,741	45,794	-
2001	93,134	245	153	255	412	768	2,071	3,941	5,506	10,495	25,850	43,437	1
2000	88,953	234	133	230	430	702	2,074	3,617	5,183	10,255	25,043	41,052	-
1999	85,485	233	161	278	419	735	1,892	3,408	4,843	10,290	23,650	39,576	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
All other diseases (Residual) (D65-E07,E15-E34,E65-F99,G04-G12,G23-G25,G31-H93,K00-K22,K29-K31,K50-K66,K71-K72,K75-K76,K83-M99,N13.0-N13.5,N13.7-N13.9,N14, N15.0,N15.8-N15.9,N20-N23,N28-N39,N41-N64,N80-N98)													
Non-Hispanic black, both sexes													
2005 2/	24,604	321	123	214	418	733	1,736	3,425	3,368	3,512	5,166	5,586	2
2004 2/	23,295	320	134	181	428	836	1,750	3,306	3,185	3,443	4,758	4,953	1
2003 2/	22,997	333	133	182	401	813	1,880	3,308	3,048	3,335	4,682	4,880	2
2002	22,297	298	146	172	416	775	1,935	3,147	2,827	3,247	4,559	4,772	3
2001	21,752	329	131	165	386	758	1,944	3,106	2,691	3,236	4,415	4,586	5
2000	21,081	296	144	189	373	810	1,948	2,897	2,607	3,153	4,248	4,413	3
1999	20,137	332	129	201	385	704	1,997	2,728	2,437	3,251	3,963	4,006	4
Non-Hispanic black, male													
2005 2/	10,478	205	74	118	222	354	857	1,824	1,787	1,665	2,044	1,326	2
2004 2/	10,222	184	73	86	219	421	853	1,794	1,696	1,675	1,924	1,296	1
2003 2/	10,019	183	78	99	182	386	952	1,802	1,610	1,643	1,915	1,167	2
2002	9,722	167	80	95	205	388	948	1,733	1,501	1,591	1,795	1,217	2
2001	9,561	192	62	83	196	381	1,030	1,713	1,411	1,570	1,767	1,151	5
2000	9,351	180	77	87	197	400	1,033	1,562	1,411	1,548	1,711	1,142	3
1999	8,967	192	67	118	202	344	1,040	1,487	1,270	1,518	1,630	1,097	2
Non-Hispanic black, female													
2005 2/	14,126	116	49	96	196	379	879	1,601	1,581	1,847	3,122	4,260	-
2004 2/	13,073	136	61	95	209	415	897	1,512	1,489	1,768	2,834	3,657	-
2003 2/	12,978	150	55	83	219	427	928	1,506	1,438	1,692	2,767	3,713	-
2002	12,575	131	66	77	211	387	987	1,414	1,326	1,656	2,764	3,555	1
2001	12,191	137	69	82	190	377	914	1,393	1,280	1,666	2,648	3,435	-
2000	11,730	116	67	102	176	410	915	1,335	1,196	1,605	2,537	3,271	-
1999	11,170	140	62	83	183	360	957	1,241	1,167	1,733	2,333	2,909	2

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
All other diseases (Residual) (D65-E07,E15-E34,E65-F99,G04-G12,G23-G25,G31-H93,K00-K22,K29-K31,K50-K66,K71-K72,K75-K76,K83-M99,N13.0-N13.5,N13.7-N13.9,N14, N15.0,N15.8-N15.9,N20-N23,N28-N39,N41-N64,N80-N98)													
Origin not stated 3/, both sexes													
2005	358	10	3	1	3	9	20	48	52	46	71	93	2
2004	445	5	2	4	5	9	27	69	56	60	81	123	4
2003	497	9	2	3	9	13	34	65	59	60	89	147	7
2002	606	8	3	2	3	13	38	85	73	69	151	157	4
2001	574	10	2	2	8	9	43	81	67	79	113	157	3
2000	591	11	3	4	9	19	41	88	60	95	123	136	2
1999	560	13	4	-	6	8	52	79	69	76	101	149	3
Origin not stated 3/, male													
2005	198	7	3	1	1	4	15	36	41	30	34	25	1
2004	221	4	1	1	3	6	17	49	39	32	33	32	4
2003	255	6	1	2	8	10	20	42	39	36	39	46	6
2002	286	2	3	1	1	8	27	58	48	44	55	36	3
2001	299	4	1	1	5	8	30	62	52	41	50	42	3
2000	301	7	2	-	6	9	31	65	38	53	52	36	2
1999	294	10	2	-	4	6	40	55	48	45	41	42	1
Origin not stated 3/, female													
2005	160	3	-	-	2	5	5	12	11	16	37	68	1
2004	224	1	1	3	2	3	10	20	17	28	48	91	-
2003	242	3	1	1	1	3	14	23	20	24	50	101	1
2002	320	6	-	1	2	5	11	27	25	25	96	121	1
2001	275	6	1	1	3	1	13	19	15	38	63	115	-
2000	290	4	1	4	3	10	10	23	22	42	71	100	-
1999	266	3	2	-	2	2	12	24	21	31	60	107	2

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Accidents (unintentional injuries) (V01-X59,Y85-Y86)													
All origins, both sexes													
2005	117,809	1,083	1,664	2,415	15,753	13,997	16,919	18,339	10,853	8,632	13,854	14,243	57
2004	112,012	1,052	1,641	2,666	15,449	13,032	16,471	16,942	9,651	8,116	13,457	13,447	88
2003	109,277	945	1,717	2,618	15,272	12,541	16,766	15,837	9,170	8,081	13,108	13,146	76
2002	106,742	946	1,641	2,718	15,412	12,569	16,710	14,675	8,345	8,086	12,904	12,651	85
2001	101,537	976	1,714	2,836	14,411	11,839	15,945	13,344	7,658	7,835	12,688	12,171	120
2000	97,900	881	1,826	2,979	14,113	11,769	15,413	12,278	7,505	7,698	11,758	11,595	85
1999	97,860	845	1,898	3,091	13,656	11,890	15,231	11,639	7,285	8,208	12,282	11,729	106
All origins, male													
2005	76,375	578	1,027	1,493	11,827	10,665	12,026	13,002	7,507	5,343	7,337	5,521	49
2004	72,050	604	964	1,661	11,423	9,871	11,671	11,988	6,562	4,992	7,030	5,210	74
2003	70,532	539	1,048	1,662	11,291	9,534	12,003	11,196	6,324	4,921	6,851	5,093	70
2002	69,257	553	1,024	1,713	11,438	9,635	12,012	10,492	5,781	5,031	6,687	4,817	74
2001	66,060	541	1,045	1,803	10,832	9,092	11,612	9,743	5,228	4,751	6,652	4,657	104
2000	63,817	501	1,098	1,861	10,460	9,059	11,444	9,063	5,133	4,639	6,063	4,413	83
1999	63,535	494	1,127	1,974	10,040	9,097	11,248	8,602	5,015	4,992	6,372	4,480	94
All origins, female													
2005	41,434	505	637	922	3,926	3,332	4,893	5,337	3,346	3,289	6,517	8,722	8
2004	39,962	448	677	1,005	4,026	3,161	4,800	4,954	3,089	3,124	6,427	8,237	14
2003	38,745	406	669	956	3,981	3,007	4,763	4,641	2,846	3,160	6,257	8,053	6
2002	37,485	393	617	1,005	3,974	2,934	4,698	4,183	2,564	3,055	6,217	7,834	11
2001	35,477	435	669	1,033	3,579	2,747	4,333	3,601	2,430	3,084	6,036	7,514	16
2000	34,083	380	728	1,118	3,653	2,710	3,969	3,215	2,372	3,059	5,695	7,182	2
1999	34,325	351	771	1,117	3,616	2,793	3,983	3,037	2,270	3,216	5,910	7,249	12

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Accidents (unintentional injuries) (V01-X59,Y85-Y86)													
Hispanic, both sexes													
2005	11,464	172	380	370	2,670	2,302	1,920	1,455	720	520	581	359	15
2004	10,408	159	313	391	2,372	2,086	1,823	1,249	678	473	508	338	18
2003	10,418	131	325	431	2,366	2,167	1,877	1,254	639	466	454	293	15
2002	10,106	127	341	389	2,413	2,027	1,841	1,168	566	466	464	276	28
2001	9,523	154	310	379	2,182	1,938	1,760	1,078	564	454	409	266	29
2000	8,830	118	306	383	2,095	1,776	1,618	953	492	418	365	278	28
1999	8,650	134	341	362	1,946	1,790	1,607	926	466	440	334	260	44
Hispanic, male													
2005	8,612	87	244	222	2,133	1,930	1,513	1,141	530	345	318	135	14
2004	7,796	87	186	241	1,926	1,733	1,454	968	474	294	278	138	17
2003	7,899	77	200	289	1,905	1,827	1,488	966	449	289	267	127	15
2002	7,698	83	215	244	1,958	1,709	1,468	888	407	307	258	134	27
2001	7,157	75	195	224	1,773	1,619	1,418	822	391	275	235	104	26
2000	6,696	67	174	210	1,717	1,499	1,317	750	347	271	191	125	28
1999	6,578	80	197	227	1,575	1,494	1,307	734	334	288	182	120	40
Hispanic, female													
2005	2,852	85	136	148	537	372	407	314	190	175	263	224	1
2004	2,612	72	127	150	446	353	369	281	204	179	230	200	1
2003	2,519	54	125	142	461	340	389	288	190	177	187	166	-
2002	2,408	44	126	145	455	318	373	280	159	159	206	142	1
2001	2,366	79	115	155	409	319	342	256	173	179	174	162	3
2000	2,134	51	132	173	378	277	301	203	145	147	174	153	-
1999	2,072	54	144	135	371	296	300	192	132	152	152	140	4

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Accidents (unintentional injuries) (V01-X59,Y85-Y86)													
Non-Hispanic 1/, both sexes													
2005	105,967	905	1,281	2,043	13,056	11,645	14,933	16,797	10,087	8,087	13,250	13,860	23
2004	101,235	891	1,323	2,263	13,045	10,901	14,597	15,607	8,938	7,619	12,924	13,089	38
2003	98,406	812	1,386	2,183	12,866	10,326	14,813	14,488	8,485	7,580	12,620	12,824	23
2002	96,175	816	1,296	2,318	12,956	10,489	14,792	13,422	7,728	7,584	12,394	12,347	33
2001	91,522	817	1,399	2,445	12,187	9,857	14,089	12,184	7,042	7,347	12,245	11,874	36
2000	88,558	759	1,512	2,585	11,968	9,935	13,706	11,219	6,966	7,247	11,361	11,288	12
1999	88,669	710	1,550	2,721	11,658	10,034	13,529	10,622	6,753	7,738	11,896	11,432	26
Non-Hispanic 1/, male													
2005	67,494	488	781	1,271	9,674	8,696	10,468	11,795	6,942	4,980	7,005	5,377	17
2004	63,982	516	775	1,415	9,476	8,103	10,179	10,948	6,061	4,677	6,738	5,063	31
2003	62,290	460	844	1,369	9,355	7,670	10,454	10,158	5,839	4,604	6,564	4,953	20
2002	61,201	468	806	1,461	9,448	7,882	10,482	9,537	5,325	4,693	6,398	4,671	30
2001	58,528	462	845	1,570	9,024	7,438	10,117	8,852	4,795	4,455	6,397	4,544	29
2000	56,736	431	921	1,648	8,707	7,512	10,054	8,231	4,747	4,344	5,855	4,275	11
1999	56,551	414	926	1,739	8,428	7,550	9,871	7,791	4,627	4,681	6,158	4,343	23
Non-Hispanic 1/, female													
2005	38,473	417	500	772	3,382	2,949	4,465	5,002	3,145	3,107	6,245	8,483	6
2004	37,253	375	548	848	3,569	2,798	4,418	4,659	2,877	2,942	6,186	8,026	7
2003	36,116	352	542	814	3,511	2,656	4,359	4,330	2,646	2,976	6,056	7,871	3
2002	34,974	348	490	857	3,508	2,607	4,310	3,885	2,403	2,891	5,996	7,676	3
2001	32,994	355	554	875	3,163	2,419	3,972	3,332	2,247	2,892	5,848	7,330	7
2000	31,822	328	591	937	3,261	2,423	3,652	2,988	2,219	2,903	5,506	7,013	1
1999	32,118	296	624	982	3,230	2,484	3,658	2,831	2,126	3,057	5,738	7,089	3

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Accidents (unintentional injuries) (V01-X59,Y85-Y86)													
Non-Hispanic white, both sexes													
2005 2/	88,970	567	898	1,423	10,711	9,347	12,252	13,657	8,303	6,890	11,942	12,962	18
2004 2/	85,466	557	933	1,573	10,847	8,712	12,025	12,744	7,390	6,532	11,774	12,344	35
2003 2/	82,815	496	985	1,535	10,638	8,238	12,106	11,721	7,043	6,466	11,487	12,078	22
2002	80,605	462	865	1,581	10,664	8,344	12,035	10,859	6,393	6,481	11,316	11,579	26
2001	76,262	500	1,006	1,688	9,867	7,798	11,340	9,704	5,776	6,309	11,157	11,092	25
2000	73,536	455	1,033	1,843	9,735	7,730	11,008	8,882	5,753	6,181	10,356	10,552	8
1999	73,326	435	1,042	1,908	9,410	7,855	10,844	8,392	5,455	6,568	10,757	10,647	13
Non-Hispanic white, male													
2005 2/	56,021	303	557	886	7,942	6,966	8,585	9,562	5,649	4,232	6,311	5,014	14
2004 2/	53,465	335	553	991	7,885	6,458	8,412	8,935	4,955	4,027	6,140	4,745	29
2003 2/	51,888	283	610	954	7,698	6,138	8,574	8,219	4,841	3,952	5,957	4,643	19
2002	50,652	279	527	997	7,746	6,273	8,550	7,745	4,374	3,991	5,805	4,342	23
2001	48,180	281	624	1,066	7,297	5,906	8,188	7,034	3,905	3,837	5,800	4,220	22
2000	46,470	266	638	1,176	7,065	5,861	8,113	6,521	3,871	3,692	5,292	3,968	7
1999	46,113	258	617	1,205	6,749	5,932	7,949	6,142	3,730	3,962	5,551	4,006	12
Non-Hispanic white, female													
2005 2/	32,949	264	341	537	2,769	2,381	3,667	4,095	2,654	2,658	5,631	7,948	4
2004 2/	32,001	222	380	582	2,962	2,254	3,613	3,809	2,435	2,505	5,634	7,599	6
2003 2/	30,927	213	375	581	2,940	2,100	3,532	3,502	2,202	2,514	5,530	7,435	3
2002	29,953	183	338	584	2,918	2,071	3,485	3,114	2,019	2,490	5,511	7,237	3
2001	28,082	219	382	622	2,570	1,892	3,152	2,670	1,871	2,472	5,357	6,872	3
2000	27,066	189	395	667	2,670	1,869	2,895	2,361	1,882	2,489	5,064	6,584	1
1999	27,213	177	425	703	2,661	1,923	2,895	2,250	1,725	2,606	5,206	6,641	1

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Accidents (unintentional injuries) (V01-X59,Y85-Y86)													
Non-Hispanic black, both sexes													
2005 2/	13,393	306	316	497	1,734	1,754	2,161	2,647	1,441	921	964	647	5
2004 2/	12,446	298	317	557	1,614	1,704	2,072	2,418	1,251	818	857	539	1
2003 2/	12,154	283	333	517	1,590	1,602	2,195	2,289	1,152	850	798	544	1
2002	12,285	324	350	607	1,680	1,676	2,227	2,145	1,075	818	799	580	4
2001	12,239	272	331	630	1,713	1,600	2,282	2,103	1,045	836	830	589	8
2000	12,087	268	396	619	1,662	1,740	2,253	1,961	983	857	768	576	4
1999	12,533	251	428	701	1,694	1,738	2,245	1,903	1,047	943	925	646	12
Non-Hispanic black, male													
2005 2/	9,130	168	179	312	1,301	1,332	1,521	1,897	1,071	589	518	239	3
2004 2/	8,430	159	182	343	1,191	1,303	1,434	1,714	901	520	467	215	1
2003 2/	8,237	158	197	330	1,205	1,197	1,539	1,625	824	518	429	214	1
2002	8,428	172	231	391	1,260	1,279	1,562	1,523	791	527	448	240	4
2001	8,381	158	188	432	1,285	1,192	1,608	1,574	750	505	461	222	6
2000	8,380	145	232	406	1,233	1,316	1,649	1,454	736	537	432	236	4
1999	8,629	140	260	467	1,281	1,313	1,628	1,441	742	596	483	268	10
Non-Hispanic black, female													
2005 2/	4,263	138	137	185	433	422	640	750	370	332	446	408	2
2004 2/	4,016	139	135	214	423	401	638	704	350	298	390	324	-
2003 2/	3,917	125	136	187	385	405	656	664	328	332	369	330	-
2002	3,857	152	119	216	420	397	665	622	284	291	351	340	-
2001	3,858	114	143	198	428	408	674	529	295	331	369	367	2
2000	3,707	123	164	213	429	424	604	507	247	320	336	340	-
1999	3,904	111	168	234	413	425	617	462	305	347	442	378	2

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Accidents (unintentional injuries) (V01-X59,Y85-Y86)													
Origin not stated 3/, both sexes													
2005	378	6	3	2	27	50	66	87	46	25	23	24	19
2004	369	2	5	12	32	45	51	86	35	24	25	20	32
2003	453	2	6	4	40	48	76	95	46	35	34	29	38
2002	461	3	4	11	43	53	77	85	51	36	46	28	24
2001	492	5	5	12	42	44	96	82	52	34	34	31	55
2000	512	4	8	11	50	58	89	106	47	33	32	29	45
1999	541	1	7	8	52	66	95	91	66	30	52	37	36
Origin not stated 3/, male													
2005	269	3	2	-	20	39	45	66	35	18	14	9	18
2004	272	1	3	5	21	35	38	72	27	21	14	9	26
2003	343	2	4	4	31	37	61	72	36	28	20	13	35
2002	358	2	3	8	32	44	62	67	49	31	31	12	17
2001	375	4	5	9	35	35	77	69	42	21	20	9	49
2000	385	3	3	3	36	48	73	82	39	24	17	13	44
1999	406	-	4	8	37	53	70	77	54	23	32	17	31
Origin not stated 3/, female													
2005	109	3	1	2	7	11	21	21	11	7	9	15	1
2004	97	1	2	7	11	10	13	14	8	3	11	11	6
2003	110	-	2	-	9	11	15	23	10	7	14	16	3
2002	103	1	1	3	11	9	15	18	2	5	15	16	7
2001	117	1	-	3	7	9	19	13	10	13	14	22	6
2000	127	1	5	8	14	10	16	24	8	9	15	16	1
1999	135	1	3	-	15	13	25	14	12	7	20	20	5

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Transport accidents (V01-V99,Y85)													
All origins, both sexes													
2005	48,441	147	646	1,547	11,288	7,638	7,295	7,101	4,918	3,353	3,168	1,329	11
2004	48,053	145	659	1,755	11,377	7,521	7,247	6,953	4,510	3,229	3,310	1,320	27
2003	48,071	148	649	1,765	11,432	7,280	7,594	6,754	4,441	3,210	3,377	1,401	20
2002	48,366	127	640	1,738	11,816	7,484	7,688	6,563	4,165	3,351	3,433	1,322	39
2001	47,288	151	678	1,823	11,139	7,454	7,827	6,251	3,861	3,282	3,507	1,272	43
2000	46,749	174	691	1,922	11,028	7,400	7,661	5,965	3,883	3,315	3,342	1,335	33
1999	46,423	189	712	1,975	10,689	7,434	7,553	5,645	3,812	3,579	3,468	1,330	37
All origins, male													
2005	34,086	74	369	910	8,208	5,840	5,303	5,208	3,435	2,157	1,835	737	10
2004	33,311	81	359	1,039	8,113	5,641	5,297	5,042	3,114	1,998	1,905	701	21
2003	33,291	81	366	1,084	8,171	5,511	5,497	4,865	3,087	1,975	1,885	751	18
2002	33,533	72	381	1,039	8,471	5,676	5,499	4,727	2,877	2,101	1,917	737	36
2001	32,794	68	392	1,115	8,111	5,661	5,613	4,484	2,631	2,015	1,983	684	37
2000	32,166	94	391	1,138	7,890	5,570	5,560	4,260	2,617	2,021	1,858	734	33
1999	31,642	101	400	1,206	7,558	5,511	5,376	4,016	2,561	2,162	1,959	758	34
All origins, female													
2005	14,355	73	277	637	3,080	1,798	1,992	1,893	1,483	1,196	1,333	592	1
2004	14,742	64	300	716	3,264	1,880	1,950	1,911	1,396	1,231	1,405	619	6
2003	14,780	67	283	681	3,261	1,769	2,097	1,889	1,354	1,235	1,492	650	2
2002	14,833	55	259	699	3,345	1,808	2,189	1,836	1,288	1,250	1,516	585	3
2001	14,494	83	286	708	3,028	1,793	2,214	1,767	1,230	1,267	1,524	588	6
2000	14,583	80	300	784	3,138	1,830	2,101	1,705	1,266	1,294	1,484	601	-
1999	14,781	88	312	769	3,131	1,923	2,177	1,629	1,251	1,417	1,509	572	3

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Transport accidents (V01-V99,Y85)													
Hispanic, both sexes													
2005	6,358	44	192	285	2,046	1,431	950	637	341	215	176	36	5
2004	6,010	40	164	302	1,866	1,381	915	575	353	216	150	42	6
2003	6,045	39	139	308	1,834	1,412	973	577	329	227	151	48	8
2002	5,921	26	182	277	1,875	1,340	930	554	300	217	157	46	17
2001	5,774	43	159	265	1,740	1,293	942	557	331	229	167	38	10
2000	5,217	42	154	279	1,616	1,126	831	508	268	220	121	38	14
1999	4,880	40	157	250	1,457	1,125	758	445	253	228	120	36	11
Hispanic, male													
2005	4,787	21	119	168	1,588	1,177	717	479	238	152	106	17	5
2004	4,551	18	95	181	1,492	1,123	720	436	229	135	97	20	5
2003	4,574	25	77	209	1,437	1,175	740	427	212	142	92	30	8
2002	4,481	16	113	173	1,479	1,094	722	394	207	149	91	27	16
2001	4,334	16	92	155	1,383	1,067	731	390	214	141	114	23	8
2000	3,921	24	85	155	1,285	921	639	373	189	140	74	22	14
1999	3,630	22	85	151	1,136	904	590	330	161	144	77	19	11
Hispanic, female													
2005	1,571	23	73	117	458	254	233	158	103	63	70	19	-
2004	1,459	22	69	121	374	258	195	139	124	81	53	22	1
2003	1,471	14	62	99	397	237	233	150	117	85	59	18	-
2002	1,440	10	69	104	396	246	208	160	93	68	66	19	1
2001	1,440	27	67	110	357	226	211	167	117	88	53	15	2
2000	1,296	18	69	124	331	205	192	135	79	80	47	16	-
1999	1,250	18	72	99	321	221	168	115	92	84	43	17	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Transport accidents (V01-V99,Y85)													
Non-Hispanic 1/, both sexes													
2005	41,957	103	453	1,261	9,222	6,179	6,325	6,440	4,564	3,131	2,986	1,288	5
2004	41,892	104	494	1,444	9,488	6,117	6,312	6,346	4,146	3,004	3,151	1,277	9
2003	41,856	109	505	1,454	9,570	5,846	6,597	6,150	4,090	2,966	3,217	1,348	4
2002	42,268	101	457	1,456	9,912	6,120	6,730	5,975	3,844	3,120	3,264	1,274	15
2001	41,339	108	517	1,554	9,371	6,141	6,848	5,668	3,513	3,044	3,330	1,230	15
2000	41,341	132	533	1,637	9,375	6,244	6,801	5,430	3,595	3,082	3,214	1,295	3
1999	41,330	149	553	1,719	9,192	6,272	6,766	5,175	3,535	3,338	3,329	1,291	11
Non-Hispanic 1/, male													
2005	29,209	53	249	742	6,606	4,641	4,573	4,712	3,187	2,000	1,725	717	4
2004	28,652	63	263	854	6,608	4,501	4,564	4,580	2,876	1,855	1,801	680	7
2003	28,591	56	286	872	6,713	4,320	4,737	4,418	2,858	1,820	1,789	718	4
2002	28,906	56	267	862	6,970	4,562	4,755	4,305	2,651	1,940	1,816	708	14
2001	28,327	52	298	957	6,706	4,578	4,855	4,071	2,405	1,869	1,865	659	12
2000	28,096	70	304	982	6,580	4,624	4,896	3,865	2,411	1,871	1,779	711	3
1999	27,850	79	313	1,049	6,396	4,578	4,764	3,666	2,379	2,009	1,869	738	10
Non-Hispanic 1/, female													
2005	12,748	50	204	519	2,616	1,538	1,752	1,728	1,377	1,131	1,261	571	1
2004	13,240	41	231	590	2,880	1,616	1,748	1,766	1,270	1,149	1,350	597	2
2003	13,265	53	219	582	2,857	1,526	1,860	1,732	1,232	1,146	1,428	630	-
2002	13,362	45	190	594	2,942	1,558	1,975	1,670	1,193	1,180	1,448	566	1
2001	13,012	56	219	597	2,665	1,563	1,993	1,597	1,108	1,175	1,465	571	3
2000	13,245	62	229	655	2,795	1,620	1,905	1,565	1,184	1,211	1,435	584	-
1999	13,480	70	240	670	2,796	1,694	2,002	1,509	1,156	1,329	1,460	553	1

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Transport accidents (V01-V99,Y85)													
Non-Hispanic white, both sexes													
2005 2/	34,349	59	306	930	7,475	4,749	5,081	5,302	3,869	2,692	2,681	1,201	4
2004 2/	34,315	66	338	1,085	7,812	4,652	5,065	5,152	3,498	2,593	2,842	1,203	9
2003 2/	34,220	73	352	1,095	7,826	4,480	5,276	5,019	3,414	2,543	2,892	1,246	4
2002	34,689	59	300	1,076	8,146	4,703	5,389	4,909	3,246	2,695	2,974	1,182	10
2001	33,753	74	353	1,122	7,563	4,771	5,494	4,611	2,937	2,628	3,044	1,148	8
2000	33,808	79	357	1,246	7,669	4,759	5,471	4,402	3,039	2,649	2,937	1,197	3
1999	33,803	98	349	1,291	7,493	4,889	5,425	4,211	2,946	2,897	3,020	1,180	4
Non-Hispanic white, male													
2005 2/	23,830	33	163	549	5,340	3,541	3,662	3,902	2,708	1,714	1,548	667	3
2004 2/	23,403	39	177	656	5,437	3,396	3,677	3,732	2,418	1,603	1,620	641	7
2003 2/	23,332	39	208	648	5,439	3,319	3,796	3,620	2,411	1,584	1,606	658	4
2002	23,628	35	168	643	5,707	3,501	3,816	3,570	2,240	1,661	1,630	648	9
2001	23,047	36	208	686	5,401	3,561	3,887	3,326	2,009	1,617	1,698	611	7
2000	22,855	44	205	748	5,361	3,518	3,935	3,160	2,014	1,597	1,609	661	3
1999	22,682	58	192	782	5,181	3,561	3,827	2,992	1,986	1,745	1,686	668	4
Non-Hispanic white, female													
2005 2/	10,519	26	143	381	2,135	1,208	1,419	1,400	1,161	978	1,133	534	1
2004 2/	10,912	27	161	429	2,375	1,256	1,388	1,420	1,080	990	1,222	562	2
2003 2/	10,888	34	144	447	2,387	1,161	1,480	1,399	1,003	959	1,286	588	-
2002	11,061	24	132	433	2,439	1,202	1,573	1,339	1,006	1,034	1,344	534	1
2001	10,706	38	145	436	2,162	1,210	1,607	1,285	928	1,011	1,346	537	1
2000	10,953	35	152	498	2,308	1,241	1,536	1,242	1,025	1,052	1,328	536	-
1999	11,121	40	157	509	2,312	1,328	1,598	1,219	960	1,152	1,334	512	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Transport accidents (V01-V99,Y85)													
Non-Hispanic black, both sexes													
2005 2/	5,739	36	118	259	1,296	1,079	970	891	515	315	197	62	1
2004 2/	5,744	32	129	284	1,228	1,134	956	946	496	276	211	52	-
2003 2/	5,684	28	125	276	1,221	1,039	1,040	862	501	303	217	72	-
2002	5,665	36	121	302	1,275	1,089	1,033	810	444	284	206	62	3
2001	5,803	18	127	354	1,322	1,058	1,071	829	455	307	198	57	7
2000	5,757	40	139	309	1,262	1,152	1,059	809	417	322	179	69	-
1999	5,810	44	168	360	1,262	1,075	1,074	758	430	314	232	86	7
Non-Hispanic black, male													
2005 2/	4,158	16	65	153	953	852	730	643	371	216	122	36	1
2004 2/	4,109	20	68	158	884	880	704	693	353	185	135	29	-
2003 2/	4,053	12	63	173	917	786	755	637	354	183	132	41	-
2002	4,051	18	79	177	917	837	733	582	327	196	139	43	3
2001	4,143	7	69	234	966	797	779	615	332	185	123	31	5
2000	4,127	21	77	191	907	875	796	571	314	218	118	39	-
1999	4,097	18	96	229	909	817	769	557	303	196	139	58	6
Non-Hispanic black, female													
2005 2/	1,581	20	53	106	343	227	240	248	144	99	75	26	-
2004 2/	1,635	12	61	126	344	254	252	253	143	91	76	23	-
2003 2/	1,631	16	62	103	304	253	285	225	147	120	85	31	-
2002	1,614	18	42	125	358	252	300	228	117	88	67	19	-
2001	1,660	11	58	120	356	261	292	214	123	122	75	26	2
2000	1,630	19	62	118	355	277	263	238	103	104	61	30	-
1999	1,713	26	72	131	353	258	305	201	127	118	93	28	1

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Transport accidents (V01-V99,Y85)													
Origin not stated 3/, both sexes													
2005	126	-	1	1	20	28	20	24	13	7	6	5	1
2004	151	1	1	9	23	23	20	32	11	9	9	1	12
2003	170	-	5	3	28	22	24	27	22	17	9	5	8
2002	177	-	1	5	29	24	28	34	21	14	12	2	7
2001	175	-	2	4	28	20	37	26	17	9	10	4	18
2000	191	-	4	6	37	30	29	27	20	13	7	2	16
1999	213	-	2	6	40	37	29	25	24	13	19	3	15
Origin not stated 3/, male													
2005	90	-	1	-	14	22	13	17	10	5	4	3	1
2004	108	-	1	4	13	17	13	26	9	8	7	1	9
2003	126	-	3	3	21	16	20	20	17	13	4	3	6
2002	146	-	1	4	22	20	22	28	19	12	10	2	6
2001	133	-	2	3	22	16	27	23	12	5	4	2	17
2000	149	-	2	1	25	25	25	22	17	10	5	1	16
1999	162	-	2	6	26	29	22	20	21	9	13	1	13
Origin not stated 3/, female													
2005	36	-	-	1	6	6	7	7	3	2	2	2	-
2004	43	1	-	5	10	6	7	6	2	1	2	-	3
2003	44	-	2	-	7	6	4	7	5	4	5	2	2
2002	31	-	-	1	7	4	6	6	2	2	2	-	1
2001	42	-	-	1	6	4	10	3	5	4	6	2	1
2000	42	-	2	5	12	5	4	5	3	3	2	1	-
1999	51	-	-	-	14	8	7	5	3	4	6	2	2

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Motor vehicle accidents (V02-V04,V09.0,V09.2,V12-V14,V19.0-V19.2,V19.4-V19.6,V20-V79,V80.3-V80.5,V81.0-V81.1,V82.0-V82.1,V83-V86,V87.0-V87.8,V88.0-V88.8, V89.0,V89.2)													
All origins, both sexes													
2005	45,343	146	617	1,447	10,908	7,239	6,748	6,397	4,454	3,110	2,991	1,280	6
2004	44,933	143	635	1,653	10,987	7,036	6,663	6,276	4,093	2,974	3,180	1,267	26
2003	44,757	145	621	1,642	10,972	6,833	6,961	6,074	3,958	2,967	3,210	1,358	16
2002	45,380	123	610	1,614	11,459	7,092	7,077	5,937	3,764	3,113	3,271	1,285	35
2001	43,788	144	626	1,686	10,725	6,937	7,083	5,564	3,465	2,990	3,312	1,223	33
2000	43,354	168	651	1,772	10,560	6,884	6,927	5,361	3,506	3,038	3,173	1,288	26
1999	42,401	184	650	1,771	10,128	6,778	6,738	4,972	3,370	3,276	3,243	1,261	30
All origins, male													
2005	31,631	74	351	848	7,904	5,501	4,869	4,631	3,055	1,976	1,709	708	5
2004	30,837	80	344	975	7,814	5,245	4,826	4,486	2,770	1,799	1,811	667	20
2003	30,655	80	353	1,009	7,806	5,156	4,984	4,303	2,683	1,778	1,767	722	14
2002	31,064	68	361	963	8,178	5,349	4,984	4,189	2,540	1,888	1,793	718	33
2001	30,064	66	356	1,029	7,778	5,236	5,016	3,938	2,309	1,786	1,859	664	27
2000	29,451	90	368	1,043	7,516	5,135	4,940	3,780	2,306	1,802	1,739	706	26
1999	28,552	97	366	1,062	7,127	4,993	4,735	3,473	2,200	1,931	1,827	714	27
All origins, female													
2005	13,712	72	266	599	3,004	1,738	1,879	1,766	1,399	1,134	1,282	572	1
2004	14,096	63	291	678	3,173	1,791	1,837	1,790	1,323	1,175	1,369	600	6
2003	14,102	65	268	633	3,166	1,677	1,977	1,771	1,275	1,189	1,443	636	2
2002	14,316	55	249	651	3,281	1,743	2,093	1,748	1,224	1,225	1,478	567	2
2001	13,724	78	270	657	2,947	1,701	2,067	1,626	1,156	1,204	1,453	559	6
2000	13,903	78	283	729	3,044	1,749	1,987	1,581	1,200	1,236	1,434	582	-
1999	13,849	87	284	709	3,001	1,785	2,003	1,499	1,170	1,345	1,416	547	3

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Motor vehicle accidents (V02-V04,V09.0,V09.2,V12-V14,V19.0-V19.2,V19.4-V19.6,V20-V79,V80.3-V80.5,V81.0-V81.1,V82.0-V82.1,V83-V86,V87.0-V87.8,V88.0-V88.8, V89.0,V89.2)													
Hispanic, both sexes													
2005	6,103	44	187	271	1,995	1,384	904	589	322	205	164	35	3
2004	5,739	40	159	292	1,820	1,309	859	532	330	205	146	41	6
2003	5,753	39	134	295	1,782	1,332	911	541	304	213	148	47	7
2002	5,659	26	176	259	1,810	1,294	885	509	283	211	146	45	15
2001	5,272	40	142	246	1,677	1,197	834	470	279	201	146	32	8
2000	4,900	40	144	260	1,557	1,066	756	472	246	198	113	34	14
1999	4,525	40	148	231	1,384	1,035	684	405	238	204	113	33	10
Hispanic, male													
2005	4,570	21	115	159	1,542	1,131	681	440	220	145	96	17	3
2004	4,318	18	93	172	1,455	1,058	673	397	208	126	93	20	5
2003	4,340	25	73	198	1,397	1,109	685	397	196	134	89	30	7
2002	4,262	16	108	163	1,421	1,052	683	358	192	144	84	26	15
2001	3,987	15	82	144	1,330	992	653	340	184	123	99	19	6
2000	3,658	24	78	142	1,236	869	574	341	170	123	68	19	14
1999	3,337	22	82	136	1,077	828	530	294	146	124	71	17	10
Hispanic, female													
2005	1,533	23	72	112	453	253	223	149	102	60	68	18	-
2004	1,421	22	66	120	365	251	186	135	122	79	53	21	1
2003	1,413	14	61	97	385	223	226	144	108	79	59	17	-
2002	1,397	10	68	96	389	242	202	151	91	67	62	19	-
2001	1,285	25	60	102	347	205	181	130	95	78	47	13	2
2000	1,242	16	66	118	321	197	182	131	76	75	45	15	-
1999	1,188	18	66	95	307	207	154	111	92	80	42	16	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Motor vehicle accidents (V02-V04,V09.0,V09.2,V12-V14,V19.0-V19.2,V19.4-V19.6,V20-V79,V80.3-V80.5,V81.0-V81.1,V82.0-V82.1,V83-V86,V87.0-V87.8,V88.0-V88.8, V89.0,V89.2)													
Non-Hispanic 1/, both sexes													
2005	39,125	102	429	1,175	8,893	5,831	5,826	5,788	4,119	2,899	2,821	1,240	2
2004	39,059	102	475	1,352	9,146	5,706	5,785	5,720	3,754	2,760	3,025	1,225	9
2003	38,849	106	482	1,344	9,165	5,480	6,031	5,507	3,636	2,738	3,053	1,306	1
2002	39,567	97	433	1,350	9,621	5,777	6,169	5,400	3,464	2,890	3,114	1,238	14
2001	38,362	104	483	1,436	9,021	5,722	6,217	5,072	3,173	2,780	3,156	1,187	11
2000	38,295	128	505	1,507	8,969	5,792	6,147	4,866	3,243	2,829	3,054	1,252	3
1999	37,684	144	500	1,534	8,707	5,712	6,026	4,546	3,110	3,059	3,112	1,225	9
Non-Hispanic 1/, male													
2005	26,982	53	235	689	6,348	4,352	4,177	4,178	2,825	1,827	1,609	688	1
2004	26,427	62	250	799	6,348	4,172	4,141	4,071	2,555	1,665	1,711	646	7
2003	26,202	55	277	808	6,391	4,032	4,283	3,887	2,473	1,632	1,674	689	1
2002	26,677	52	252	796	6,735	4,280	4,284	3,808	2,333	1,734	1,700	690	13
2001	25,962	51	273	882	6,426	4,230	4,340	3,579	2,116	1,658	1,756	643	8
2000	25,670	66	288	900	6,257	4,243	4,346	3,421	2,122	1,671	1,667	686	3
1999	25,072	75	282	920	6,027	4,141	4,184	3,163	2,034	1,798	1,744	696	8
Non-Hispanic 1/, female													
2005	12,143	49	194	486	2,545	1,479	1,649	1,610	1,294	1,072	1,212	552	1
2004	12,632	40	225	553	2,798	1,534	1,644	1,649	1,199	1,095	1,314	579	2
2003	12,647	51	205	536	2,774	1,448	1,748	1,620	1,163	1,106	1,379	617	-
2002	12,890	45	181	554	2,886	1,497	1,885	1,592	1,131	1,156	1,414	548	1
2001	12,400	53	210	554	2,595	1,492	1,877	1,493	1,057	1,122	1,400	544	3
2000	12,625	62	217	607	2,712	1,549	1,801	1,445	1,121	1,158	1,387	566	-
1999	12,612	69	218	614	2,680	1,571	1,842	1,383	1,076	1,261	1,368	529	1

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Motor vehicle accidents (V02-V04,V09.0,V09.2,V12-V14,V19.0-V19.2,V19.4-V19.6,V20-V79,V80.3-V80.5,V81.0-V81.1,V82.0-V82.1,V83-V86,V87.0-V87.8,V88.0-V88.8, V89.0,V89.2)													
Non-Hispanic white, both sexes													
2005 2/	32,002	58	289	860	7,220	4,473	4,662	4,785	3,471	2,488	2,536	1,158	2
2004 2/	31,931	65	324	1,007	7,535	4,321	4,622	4,633	3,155	2,376	2,730	1,154	9
2003 2/	31,662	70	333	1,005	7,495	4,182	4,792	4,479	3,008	2,343	2,748	1,206	1
2002	32,405	56	280	1,000	7,897	4,412	4,923	4,421	2,918	2,496	2,844	1,149	9
2001	31,283	71	323	1,036	7,266	4,424	4,980	4,121	2,652	2,398	2,891	1,113	8
2000	31,219	75	340	1,136	7,330	4,379	4,923	3,925	2,728	2,431	2,791	1,158	3
1999	30,743	94	315	1,144	7,099	4,425	4,797	3,678	2,579	2,655	2,832	1,122	3
Non-Hispanic white, male													
2005 2/	21,992	33	152	508	5,148	3,310	3,332	3,475	2,386	1,562	1,446	639	1
2004 2/	21,540	39	168	608	5,235	3,137	3,325	3,307	2,137	1,430	1,538	609	7
2003 2/	21,309	38	200	602	5,177	3,085	3,408	3,176	2,067	1,419	1,505	631	1
2002	21,743	32	155	594	5,505	3,262	3,423	3,148	1,968	1,484	1,532	632	8
2001	21,073	35	185	631	5,164	3,270	3,468	2,915	1,764	1,436	1,600	598	7
2000	20,789	40	197	680	5,094	3,196	3,473	2,781	1,759	1,424	1,505	637	3
1999	20,362	55	175	683	4,884	3,201	3,338	2,565	1,689	1,561	1,575	633	3
Non-Hispanic white, female													
2005 2/	10,010	25	137	352	2,072	1,163	1,330	1,310	1,085	926	1,090	519	1
2004 2/	10,391	26	156	399	2,300	1,184	1,297	1,326	1,018	946	1,192	545	2
2003 2/	10,353	32	133	403	2,318	1,097	1,384	1,303	941	924	1,243	575	-
2002	10,662	24	125	406	2,392	1,150	1,500	1,273	950	1,012	1,312	517	1
2001	10,210	36	138	405	2,102	1,154	1,512	1,206	888	962	1,291	515	1
2000	10,430	35	143	456	2,236	1,183	1,450	1,144	969	1,007	1,286	521	-
1999	10,381	39	140	461	2,215	1,224	1,459	1,113	890	1,094	1,257	489	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Motor vehicle accidents (V02-V04,V09.0,V09.2,V12-V14,V19.0-V19.2,V19.4-V19.6,V20-V79,V80.3-V80.5,V81.0-V81.1,V82.0-V82.1,V83-V86,V87.0-V87.8,V88.0-V88.8, V89.0,V89.2)													
Non-Hispanic black, both sexes													
2005 2/	5,394	36	113	247	1,245	1,030	911	788	483	296	186	59	-
2004 2/	5,395	31	125	272	1,174	1,075	892	864	452	260	198	52	-
2003 2/	5,349	28	121	261	1,168	991	974	784	463	285	203	71	-
2002	5,338	35	118	278	1,243	1,049	959	734	404	263	191	61	3
2001	5,423	18	123	324	1,285	1,007	985	747	410	283	184	54	3
2000	5,424	40	132	294	1,216	1,095	977	750	388	298	169	65	-
1999	5,388	43	152	331	1,209	1,006	985	685	387	291	214	79	6
Non-Hispanic black, male													
2005 2/	3,883	16	63	143	908	813	682	563	345	200	115	35	-
2004 2/	3,821	19	65	152	836	827	649	627	317	173	127	29	-
2003 2/	3,771	12	62	159	871	747	702	569	319	167	123	40	-
2002	3,774	17	77	163	891	802	673	516	290	175	125	42	3
2001	3,842	7	67	214	935	757	706	550	295	163	116	31	1
2000	3,857	21	72	179	868	828	727	524	289	199	112	38	-
1999	3,763	17	84	203	869	761	696	499	267	178	132	52	5
Non-Hispanic black, female													
2005 2/	1,511	20	50	104	337	217	229	225	138	96	71	24	-
2004 2/	1,574	12	60	120	338	248	243	237	135	87	71	23	-
2003 2/	1,578	16	59	102	297	244	272	215	144	118	80	31	-
2002	1,564	18	41	115	352	247	286	218	114	88	66	19	-
2001	1,581	11	56	110	350	250	279	197	115	120	68	23	2
2000	1,567	19	60	115	348	267	250	226	99	99	57	27	-
1999	1,625	26	68	128	340	245	289	186	120	113	82	27	1

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Motor vehicle accidents (V02-V04,V09.0,V09.2,V12-V14,V19.0-V19.2,V19.4-V19.6,V20-V79,V80.3-V80.5,V81.0-V81.1,V82.0-V82.1,V83-V86,V87.0-V87.8,V88.0-V88.8, V89.0,V89.2)													
Origin not stated 3/, both sexes													
2005	115	-	1	1	20	24	18	20	13	6	6	5	1
2004	135	1	1	9	21	21	19	24	9	9	9	1	11
2003	155	-	5	3	25	21	19	26	18	16	9	5	8
2002	154	-	1	5	28	21	23	28	17	12	11	2	6
2001	154	-	1	4	27	18	32	22	13	9	10	4	14
2000	159	-	2	5	34	26	24	23	17	11	6	2	9
1999	192	-	2	6	37	31	28	21	22	13	18	3	11
Origin not stated 3/, male													
2005	79	-	1	-	14	18	11	13	10	4	4	3	1
2004	92	-	1	4	11	15	12	18	7	8	7	1	8
2003	113	-	3	3	18	15	16	19	14	12	4	3	6
2002	125	-	1	4	22	17	17	23	15	10	9	2	5
2001	115	-	1	3	22	14	23	19	9	5	4	2	13
2000	123	-	2	1	23	23	20	18	14	8	4	1	9
1999	143	-	2	6	23	24	21	16	20	9	12	1	9
Origin not stated 3/, female													
2005	36	-	-	1	6	6	7	7	3	2	2	2	-
2004	43	1	-	5	10	6	7	6	2	1	2	-	3
2003	42	-	2	-	7	6	3	7	4	4	5	2	2
2002	29	-	-	1	6	4	6	5	2	2	2	-	1
2001	39	-	-	1	5	4	9	3	4	4	6	2	1
2000	36	-	-	4	11	3	4	5	3	3	2	1	-
1999	49	-	-	-	14	7	7	5	2	4	6	2	2

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Other land transport accidents (V01,V05-V06,V09.1,V09.3-V09.9,V10-V11,V15-V18,V19.3,V19.8-V19.9,V80.0-V80.2,V80.6-V80.9,V81.2-V81.9,V82.2-V82.9,V87.9,V88.9, V89.1,V89.3,V89.9)													
All origins, both sexes													
2005	1,241	1	27	50	192	162	210	267	138	81	84	25	4
2004	1,199	1	19	57	190	184	211	222	132	92	63	27	1
2003	1,357	2	17	60	264	204	239	234	147	81	80	25	4
2002	1,086	2	17	70	159	138	223	211	109	70	66	17	4
2001	1,405	3	28	80	198	177	285	247	133	117	98	29	10
2000	1,492	2	31	98	243	212	317	234	122	100	91	35	7
1999	2,118	4	48	128	362	317	408	304	182	145	159	54	7
All origins, male													
2005	957	-	17	35	156	136	160	212	110	54	58	15	4
2004	926	1	12	38	152	150	168	182	102	58	46	16	1
2003	1,044	-	9	38	205	166	186	192	113	63	52	16	4
2002	882	2	10	47	134	108	187	185	92	56	50	8	3
2001	1,067	-	20	49	158	152	235	189	104	84	56	10	10
2000	1,125	2	17	61	194	171	258	166	98	75	57	19	7
1999	1,559	4	26	91	279	244	319	234	135	97	88	35	7
All origins, female													
2005	284	1	10	15	36	26	50	55	28	27	26	10	-
2004	273	-	7	19	38	34	43	40	30	34	17	11	-
2003	313	2	8	22	59	38	53	42	34	18	28	9	-
2002	204	-	7	23	25	30	36	26	17	14	16	9	1
2001	338	3	8	31	40	25	50	58	29	33	42	19	-
2000	367	-	14	37	49	41	59	68	24	25	34	16	-
1999	559	-	22	37	83	73	89	70	47	48	71	19	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Other land transport accidents (V01,V05-V06,V09.1,V09.3-V09.9,V10-V11,V15-V18,V19.3,V19.8-V19.9,V80.0-V80.2,V80.6-V80.9,V81.2-V81.9,V82.2-V82.9,V87.9,V88.9, V89.1,V89.3,V89.9)													
Hispanic, both sexes													
2005	153	-	5	10	34	28	27	21	11	5	9	1	2
2004	161	-	4	10	25	40	34	24	11	8	4	1	-
2003	167	-	3	10	32	48	34	17	13	7	1	1	1
2002	144	-	2	10	42	21	28	22	7	2	8	-	2
2001	200	-	8	7	39	38	42	29	14	5	13	3	2
2000	201	-	10	15	34	36	44	25	14	14	5	4	-
1999	264	-	8	14	55	60	57	31	8	21	7	2	1
Hispanic, male													
2005	134	-	4	7	33	28	23	16	10	4	7	-	2
2004	141	-	2	9	20	38	29	23	10	6	4	-	-
2003	139	-	2	8	24	43	32	15	9	4	1	-	1
2002	119	-	1	6	38	18	25	17	6	2	5	-	1
2001	173	-	6	4	37	35	39	22	12	4	10	2	2
2000	166	-	7	11	31	32	36	22	11	10	3	3	-
1999	220	-	3	12	45	53	47	27	8	17	6	1	1
Hispanic, female													
2005	19	-	1	3	1	-	4	5	1	1	2	1	-
2004	20	-	2	1	5	2	5	1	1	2	-	1	-
2003	28	-	1	2	8	5	2	2	4	3	-	1	-
2002	25	-	1	4	4	3	3	5	1	-	3	-	1
2001	27	-	2	3	2	3	3	7	2	1	3	1	-
2000	35	-	3	4	3	4	8	3	3	4	2	1	-
1999	44	-	5	2	10	7	10	4	-	4	1	1	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Other land transport accidents (V01,V05-V06,V09.1,V09.3-V09.9,V10-V11,V15-V18,V19.3,V19.8-V19.9,V80.0-V80.2,V80.6-V80.9,V81.2-V81.9,V82.2-V82.9,V87.9,V88.9, V89.1,V89.3,V89.9)													
Non-Hispanic 1/, both sexes													
2005	1,081	1	22	40	158	131	181	244	127	76	75	24	2
2004	1,028	1	15	47	165	143	176	192	120	84	59	26	-
2003	1,185	2	14	50	232	156	203	216	133	73	79	24	3
2002	929	2	15	60	116	117	192	184	100	68	57	17	1
2001	1,187	3	19	73	158	137	241	214	115	112	85	26	4
2000	1,262	2	19	82	206	172	268	207	105	85	85	31	-
1999	1,836	4	40	114	304	252	350	271	172	124	151	52	2
Non-Hispanic 1/, male													
2005	816	-	13	28	123	105	135	194	100	50	51	15	2
2004	775	1	10	29	132	111	138	153	91	52	42	16	-
2003	901	-	7	30	181	123	153	176	103	58	51	16	3
2002	752	2	9	41	96	90	159	164	84	54	44	8	1
2001	878	-	13	45	121	115	194	163	89	80	46	8	4
2000	936	2	10	50	161	137	217	142	84	64	53	16	-
1999	1,323	4	23	79	231	187	271	205	126	80	81	34	2
Non-Hispanic 1/, female													
2005	265	1	9	12	35	26	46	50	27	26	24	9	-
2004	253	-	5	18	33	32	38	39	29	32	17	10	-
2003	284	2	7	20	51	33	50	40	30	15	28	8	-
2002	177	-	6	19	20	27	33	20	16	14	13	9	-
2001	309	3	6	28	37	22	47	51	26	32	39	18	-
2000	326	-	9	32	45	35	51	65	21	21	32	15	-
1999	513	-	17	35	73	65	79	66	46	44	70	18	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Other land transport accidents (V01,V05-V06,V09.1,V09.3-V09.9,V10-V11,V15-V18,V19.3,V19.8-V19.9,V80.0-V80.2,V80.6-V80.9,V81.2-V81.9,V82.2-V82.9,V87.9,V88.9, V89.1,V89.3,V89.9)													
Non-Hispanic white, both sexes													
2005 2/	815	1	15	31	114	90	143	174	106	55	64	21	1
2004 2/	782	-	11	39	118	97	143	132	96	71	51	24	-
2003 2/	952	2	11	39	195	117	159	162	112	61	69	22	3
2002	719	1	11	44	95	94	140	141	74	55	48	15	1
2001	901	2	16	49	126	103	165	165	89	92	73	21	-
2000	997	2	9	67	168	132	201	161	84	70	75	28	-
1999	1,480	3	25	86	239	195	282	219	151	102	131	46	1
Non-Hispanic white, male													
2005 2/	601	-	10	21	84	71	105	136	80	35	44	14	1
2004 2/	580	-	7	25	91	72	110	104	73	46	37	15	-
2003 2/	715	-	6	21	153	91	122	128	85	48	44	14	3
2002	582	1	7	32	81	71	116	124	63	43	36	7	1
2001	673	-	12	30	99	89	133	129	73	63	40	5	-
2000	741	2	3	41	130	106	165	110	68	51	49	16	-
1999	1,051	3	12	57	177	143	218	163	110	67	71	29	1
Non-Hispanic white, female													
2005 2/	214	1	5	10	30	19	38	38	26	20	20	7	-
2004 2/	202	-	4	14	27	25	33	28	23	25	14	9	-
2003 2/	237	2	5	18	42	26	37	34	27	13	25	8	-
2002	137	-	4	12	14	23	24	17	11	12	12	8	-
2001	228	2	4	19	27	14	32	36	16	29	33	16	-
2000	256	-	6	26	38	26	36	51	16	19	26	12	-
1999	429	-	13	29	62	52	64	56	41	35	60	17	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Other land transport accidents (V01,V05-V06,V09.1,V09.3-V09.9,V10-V11,V15-V18,V19.3,V19.8-V19.9,V80.0-V80.2,V80.6-V80.9,V81.2-V81.9,V82.2-V82.9,V87.9,V88.9, V89.1,V89.3,V89.9)													
Non-Hispanic black, both sexes													
2005 2/.....	200	-	5	8	32	31	32	55	13	15	6	2	1
2004 2/.....	194	1	3	7	41	34	28	43	22	7	8	-	-
2003 2/.....	188	-	3	8	27	30	39	46	19	8	7	1	-
2002.....	168	1	3	13	17	19	40	41	18	9	7	-	-
2001.....	216	-	3	23	21	24	59	38	22	15	6	1	4
2000.....	203	-	7	11	30	32	54	34	15	10	7	3	-
1999.....	248	1	12	20	33	42	52	40	17	14	11	5	1
Non-Hispanic black, male													
2005 2/.....	160	-	2	6	27	25	25	45	12	12	4	1	1
2004 2/.....	158	1	2	4	36	30	25	34	16	5	5	-	-
2003 2/.....	154	-	1	7	21	25	28	42	17	8	4	1	-
2002.....	143	1	2	7	12	17	34	39	16	9	6	-	-
2001.....	159	-	1	15	15	18	48	28	14	13	3	-	4
2000.....	155	-	5	8	24	25	43	26	12	9	3	-	-
1999.....	192	1	9	19	27	34	41	32	12	9	3	4	1
Non-Hispanic black, female													
2005 2/.....	40	-	3	2	5	6	7	10	1	3	2	1	-
2004 2/.....	36	-	1	3	5	4	3	9	6	2	3	-	-
2003 2/.....	34	-	2	1	6	5	11	4	2	-	3	-	-
2002.....	25	-	1	6	5	2	6	2	2	-	1	-	-
2001.....	57	-	2	8	6	6	11	10	8	2	3	1	-
2000.....	48	-	2	3	6	7	11	8	3	1	4	3	-
1999.....	56	-	3	1	6	8	11	8	5	5	8	1	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Other land transport accidents (V01,V05-V06,V09.1,V09.3-V09.9,V10-V11,V15-V18,V19.3,V19.8-V19.9,V80.0-V80.2,V80.6-V80.9,V81.2-V81.9,V82.2-V82.9,V87.9,V88.9, V89.1,V89.3,V89.9)													
Origin not stated 3/, both sexes													
2005	7	-	-	-	-	3	2	2	-	-	-	-	-
2004	10	-	-	-	-	1	1	6	1	-	-	-	1
2003	5	-	-	-	-	-	2	1	1	1	-	-	-
2002	13	-	-	-	1	-	3	5	2	-	1	-	1
2001	18	-	1	-	1	2	2	4	4	-	-	-	4
2000	29	-	2	1	3	4	5	2	3	1	1	-	7
1999	18	-	-	-	3	5	1	2	2	-	1	-	4
Origin not stated 3/, male													
2005	7	-	-	-	-	3	2	2	-	-	-	-	-
2004	10	-	-	-	-	1	1	6	1	-	-	-	1
2003	4	-	-	-	-	-	1	1	1	1	-	-	-
2002	11	-	-	-	-	-	3	4	2	-	1	-	1
2001	16	-	1	-	-	2	2	4	3	-	-	-	4
2000	23	-	-	-	2	2	5	2	3	1	1	-	7
1999	16	-	-	-	3	4	1	2	1	-	1	-	4
Origin not stated 3/, female													
2005	-	-	-	-	-	-	-	-	-	-	-	-	-
2004	-	-	-	-	-	-	-	-	-	-	-	-	-
2003	1	-	-	-	-	-	1	-	-	-	-	-	-
2002	2	-	-	-	1	-	-	1	-	-	-	-	-
2001	2	-	-	-	1	-	-	-	1	-	-	-	-
2000	6	-	2	1	1	2	-	-	-	-	-	-	-
1999	2	-	-	-	-	1	-	-	1	-	-	-	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Water, air and space, and other and unspecified transport accidents and their sequelae (V90-V99,Y85)													
All origins, both sexes													
2005	1,857	-	2	50	188	237	337	437	326	162	93	24	1
2004	1,921	1	5	45	200	301	373	455	285	163	67	26	-
2003	1,957	1	11	63	196	243	394	446	336	162	87	18	-
2002	1,900	2	13	54	198	254	388	415	292	168	96	20	-
2001	2,095	4	24	57	216	340	459	440	263	175	97	20	-
2000	1,903	4	9	52	225	304	417	370	255	177	78	12	-
1999	1,904	1	14	76	199	339	407	369	260	158	66	15	-
All origins, male													
2005	1,498	-	1	27	148	203	274	365	270	127	68	14	1
2004	1,548	-	3	26	147	246	303	374	242	141	48	18	-
2003	1,592	1	4	37	160	189	327	370	291	134	66	13	-
2002	1,587	2	10	29	159	219	328	353	245	157	74	11	-
2001	1,663	2	16	37	175	273	362	357	218	145	68	10	-
2000	1,590	2	6	34	180	264	362	314	213	144	62	9	-
1999	1,531	-	8	53	152	274	322	309	226	134	44	9	-
All origins, female													
2005	359	-	1	23	40	34	63	72	56	35	25	10	-
2004	373	1	2	19	53	55	70	81	43	22	19	8	-
2003	365	-	7	26	36	54	67	76	45	28	21	5	-
2002	313	-	3	25	39	35	60	62	47	11	22	9	-
2001	432	2	8	20	41	67	97	83	45	30	29	10	-
2000	313	2	3	18	45	40	55	56	42	33	16	3	-
1999	373	1	6	23	47	65	85	60	34	24	22	6	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Water, air and space, and other and unspecified transport accidents and their sequelae (V90-V99,Y85)													
Hispanic, both sexes													
2005	102	-	-	4	17	19	19	27	8	5	3	-	-
2004	110	-	1	-	21	32	22	19	12	3	-	-	-
2003	125	-	2	3	20	32	28	19	12	7	2	-	-
2002	118	-	4	8	23	25	17	23	10	4	3	1	-
2001	302	3	9	12	24	58	66	58	38	23	8	3	-
2000	116	2	-	4	25	24	31	11	8	8	3	-	-
1999	91	-	1	5	18	30	17	9	7	3	-	1	-
Hispanic, male													
2005	83	-	-	2	13	18	13	23	8	3	3	-	-
2004	92	-	-	-	17	27	18	16	11	3	-	-	-
2003	95	-	2	3	16	23	23	15	7	4	2	-	-
2002	100	-	4	4	20	24	14	19	9	3	2	1	-
2001	174	1	4	7	16	40	39	28	18	14	5	2	-
2000	97	-	-	2	18	20	29	10	8	7	3	-	-
1999	73	-	-	3	14	23	13	9	7	3	-	1	-
Hispanic, female													
2005	19	-	-	2	4	1	6	4	-	2	-	-	-
2004	18	-	1	-	4	5	4	3	1	-	-	-	-
2003	30	-	-	-	4	9	5	4	5	3	-	-	-
2002	18	-	-	4	3	1	3	4	1	1	1	-	-
2001	128	2	5	5	8	18	27	30	20	9	3	1	-
2000	19	2	-	2	7	4	2	1	-	1	-	-	-
1999	18	-	1	2	4	7	4	-	-	-	-	-	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Water, air and space, and other and unspecified transport accidents and their sequelae (V90-V99,Y85)													
Non-Hispanic 1/, both sexes													
2005	1,751	-	2	46	171	217	318	408	318	156	90	24	1
2004	1,805	1	4	45	177	268	351	434	272	160	67	26	-
2003	1,822	1	9	60	173	210	363	427	321	155	85	18	-
2002	1,772	2	9	46	175	226	369	391	280	162	93	19	-
2001	1,790	1	15	45	192	282	390	382	225	152	89	17	-
2000	1,784	2	9	48	200	280	386	357	247	168	75	12	-
1999	1,810	1	13	71	181	308	390	358	253	155	66	14	-
Non-Hispanic 1/, male													
2005	1,411	-	1	25	135	184	261	340	262	123	65	14	1
2004	1,450	-	3	26	128	218	285	356	230	138	48	18	-
2003	1,488	1	2	34	141	165	301	355	282	130	64	13	-
2002	1,477	2	6	25	139	192	312	333	234	152	72	10	-
2001	1,487	1	12	30	159	233	321	329	200	131	63	8	-
2000	1,490	2	6	32	162	244	333	302	205	136	59	9	-
1999	1,455	-	8	50	138	250	309	298	219	131	44	8	-
Non-Hispanic 1/, female													
2005	340	-	1	21	36	33	57	68	56	33	25	10	-
2004	355	1	1	19	49	50	66	78	42	22	19	8	-
2003	334	-	7	26	32	45	62	72	39	25	21	5	-
2002	295	-	3	21	36	34	57	58	46	10	21	9	-
2001	303	-	3	15	33	49	69	53	25	21	26	9	-
2000	294	-	3	16	38	36	53	55	42	32	16	3	-
1999	355	1	5	21	43	58	81	60	34	24	22	6	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Water, air and space, and other and unspecified transport accidents and their sequelae (V90-V99,Y85)													
Non-Hispanic white, both sexes													
2005 2/	1,532	-	2	39	141	186	276	343	292	149	81	22	1
2004 2/	1,602	1	3	39	159	234	300	387	247	146	61	25	-
2003 2/	1,606	1	8	51	136	181	325	378	294	139	75	18	-
2002	1,565	2	9	32	154	197	326	347	254	144	82	18	-
2001	1,569	1	14	37	171	244	349	325	196	138	80	14	-
2000	1,592	2	8	43	171	248	347	316	227	148	71	11	-
1999	1,580	1	9	61	155	269	346	314	216	140	57	12	-
Non-Hispanic white, male													
2005 2/	1,237	-	1	20	108	160	225	291	242	117	58	14	1
2004 2/	1,283	-	2	23	111	187	242	321	208	127	45	17	-
2003 2/	1,308	1	2	25	109	143	266	316	259	117	57	13	-
2002	1,303	2	6	17	121	168	277	298	209	134	62	9	-
2001	1,301	1	11	25	138	202	286	282	172	118	58	8	-
2000	1,325	2	5	27	137	216	297	269	187	122	55	8	-
1999	1,269	-	5	42	120	217	271	264	187	117	40	6	-
Non-Hispanic white, female													
2005 2/	295	-	1	19	33	26	51	52	50	32	23	8	-
2004 2/	319	1	1	16	48	47	58	66	39	19	16	8	-
2003 2/	298	-	6	26	27	38	59	62	35	22	18	5	-
2002	262	-	3	15	33	29	49	49	45	10	20	9	-
2001	268	-	3	12	33	42	63	43	24	20	22	6	-
2000	267	-	3	16	34	32	50	47	40	26	16	3	-
1999	311	1	4	19	35	52	75	50	29	23	17	6	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Water, air and space, and other and unspecified transport accidents and their sequelae (V90-V99,Y85)													
Non-Hispanic black, both sexes													
2005 2/	145	-	-	4	19	18	27	48	19	4	5	1	-
2004 2/	155	-	1	5	13	25	36	39	22	9	5	-	-
2003 2/	147	-	1	7	26	18	27	32	19	10	7	-	-
2002	159	-	-	11	15	21	34	35	22	12	8	1	-
2001	164	-	1	7	16	27	27	44	23	9	8	2	-
2000	130	-	-	4	16	25	28	25	14	14	3	1	-
1999	174	-	4	9	20	27	37	33	26	9	7	2	-
Non-Hispanic black, male													
2005 2/	115	-	-	4	18	14	23	35	14	4	3	-	-
2004 2/	130	-	1	2	12	23	30	32	20	7	3	-	-
2003 2/	128	-	-	7	25	14	25	26	18	8	5	-	-
2002	134	-	-	7	14	18	26	27	21	12	8	1	-
2001	142	-	1	5	16	22	25	37	23	9	4	-	-
2000	115	-	-	4	15	22	26	21	13	10	3	1	-
1999	142	-	3	7	13	22	32	26	24	9	4	2	-
Non-Hispanic black, female													
2005 2/	30	-	-	-	1	4	4	13	5	-	2	1	-
2004 2/	25	-	-	3	1	2	6	7	2	2	2	-	-
2003 2/	19	-	1	-	1	4	2	6	1	2	2	-	-
2002	25	-	-	4	1	3	8	8	1	-	-	-	-
2001	22	-	-	2	-	5	2	7	-	-	4	2	-
2000	15	-	-	-	1	3	2	4	1	4	-	-	-
1999	32	-	1	2	7	5	5	7	2	-	3	-	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Water, air and space, and other and unspecified transport accidents and their sequelae (V90-V99,Y85)													
Origin not stated 3/, both sexes													
2005	4	-	-	-	-	1	-	2	-	1	-	-	-
2004	6	-	-	-	2	1	-	2	1	-	-	-	-
2003	10	-	-	-	3	1	3	-	3	-	-	-	-
2002	10	-	-	-	-	3	2	1	2	2	-	-	-
2001	3	-	-	-	-	-	3	-	-	-	-	-	-
2000	3	-	-	-	-	-	-	2	-	1	-	-	-
1999	3	-	-	-	-	1	-	2	-	-	-	-	-
Origin not stated 3/, male													
2005	4	-	-	-	-	1	-	2	-	1	-	-	-
2004	6	-	-	-	2	1	-	2	1	-	-	-	-
2003	9	-	-	-	3	1	3	-	2	-	-	-	-
2002	10	-	-	-	-	3	2	1	2	2	-	-	-
2001	2	-	-	-	-	-	2	-	-	-	-	-	-
2000	3	-	-	-	-	-	-	2	-	1	-	-	-
1999	3	-	-	-	-	1	-	2	-	-	-	-	-
Origin not stated 3/, female													
2005	-	-	-	-	-	-	-	-	-	-	-	-	-
2004	-	-	-	-	-	-	-	-	-	-	-	-	-
2003	1	-	-	-	-	-	-	-	1	-	-	-	-
2002	-	-	-	-	-	-	-	-	-	-	-	-	-
2001	1	-	-	-	-	-	1	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-	-	-	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Nontransport accidents (W00-X59,Y86)													
All origins, both sexes													
2005	69,368	936	1,018	868	4,465	6,359	9,624	11,238	5,935	5,279	10,686	12,914	46
2004	63,959	907	982	911	4,072	5,511	9,224	9,989	5,141	4,887	10,147	12,127	61
2003	61,206	797	1,068	853	3,840	5,261	9,172	9,083	4,729	4,871	9,731	11,745	56
2002	58,376	819	1,001	980	3,596	5,085	9,022	8,112	4,180	4,735	9,471	11,329	46
2001	54,249	825	1,036	1,013	3,272	4,385	8,118	7,093	3,797	4,553	9,181	10,899	77
2000	51,151	707	1,135	1,057	3,085	4,369	7,752	6,313	3,622	4,383	8,416	10,260	52
1999	51,437	656	1,186	1,116	2,967	4,456	7,678	5,994	3,473	4,629	8,814	10,399	69
All origins, male													
2005	42,289	504	658	583	3,619	4,825	6,723	7,794	4,072	3,186	5,502	4,784	39
2004	38,739	523	605	622	3,310	4,230	6,374	6,946	3,448	2,994	5,125	4,509	53
2003	37,241	458	682	578	3,120	4,023	6,506	6,331	3,237	2,946	4,966	4,342	52
2002	35,724	481	643	674	2,967	3,959	6,513	5,765	2,904	2,930	4,770	4,080	38
2001	33,266	473	653	688	2,721	3,431	5,999	5,259	2,597	2,736	4,669	3,973	67
2000	31,651	407	707	723	2,570	3,489	5,884	4,803	2,516	2,618	4,205	3,679	50
1999	31,893	393	727	768	2,482	3,586	5,872	4,586	2,454	2,830	4,413	3,722	60
All origins, female													
2005	27,079	432	360	285	846	1,534	2,901	3,444	1,863	2,093	5,184	8,130	7
2004	25,220	384	377	289	762	1,281	2,850	3,043	1,693	1,893	5,022	7,618	8
2003	23,965	339	386	275	720	1,238	2,666	2,752	1,492	1,925	4,765	7,403	4
2002	22,652	338	358	306	629	1,126	2,509	2,347	1,276	1,805	4,701	7,249	8
2001	20,983	352	383	325	551	954	2,119	1,834	1,200	1,817	4,512	6,926	10
2000	19,500	300	428	334	515	880	1,868	1,510	1,106	1,765	4,211	6,581	2
1999	19,544	263	459	348	485	870	1,806	1,408	1,019	1,799	4,401	6,677	9

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Nontransport accidents (W00-X59,Y86)													
Hispanic, both sexes													
2005	5,106	128	188	85	624	871	970	818	379	305	405	323	10
2004	4,398	119	149	89	506	705	908	674	325	257	358	296	12
2003	4,373	92	186	123	532	755	904	677	310	239	303	245	7
2002	4,185	101	159	112	538	687	911	614	266	249	307	230	11
2001	3,749	111	151	114	442	645	818	521	233	225	242	228	19
2000	3,613	76	152	104	479	650	787	445	224	198	244	240	14
1999	3,770	94	184	112	489	665	849	481	213	212	214	224	33
Hispanic, male													
2005	3,825	66	125	54	545	753	796	662	292	193	212	118	9
2004	3,245	69	91	60	434	610	734	532	245	159	181	118	12
2003	3,325	52	123	80	468	652	748	539	237	147	175	97	7
2002	3,217	67	102	71	479	615	746	494	200	158	167	107	11
2001	2,823	59	103	69	390	552	687	432	177	134	121	81	18
2000	2,775	43	89	55	432	578	678	377	158	131	117	103	14
1999	2,948	58	112	76	439	590	717	404	173	144	105	101	29
Hispanic, female													
2005	1,281	62	63	31	79	118	174	156	87	112	193	205	1
2004	1,153	50	58	29	72	95	174	142	80	98	177	178	-
2003	1,048	40	63	43	64	103	156	138	73	92	128	148	-
2002	968	34	57	41	59	72	165	120	66	91	140	123	-
2001	926	52	48	45	52	93	131	89	56	91	121	147	1
2000	838	33	63	49	47	72	109	68	66	67	127	137	-
1999	822	36	72	36	50	75	132	77	40	68	109	123	4

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Nontransport accidents (W00-X59,Y86)													
Non-Hispanic 1/, both sexes													
2005	64,010	802	828	782	3,834	5,466	8,608	10,357	5,523	4,956	10,264	12,572	18
2004	59,343	787	829	819	3,557	4,784	8,285	9,261	4,792	4,615	9,773	11,812	29
2003	56,550	703	881	729	3,296	4,480	8,216	8,338	4,395	4,614	9,403	11,476	19
2002	53,907	715	839	862	3,044	4,369	8,062	7,447	3,884	4,464	9,130	11,073	18
2001	50,183	709	882	891	2,816	3,716	7,241	6,516	3,529	4,303	8,915	10,644	21
2000	47,217	627	979	948	2,593	3,691	6,905	5,789	3,371	4,165	8,147	9,993	9
1999	47,339	561	997	1,002	2,466	3,762	6,763	5,447	3,218	4,400	8,567	10,141	15
Non-Hispanic 1/, male													
2005	38,285	435	532	529	3,068	4,055	5,895	7,083	3,755	2,980	5,280	4,660	13
2004	35,330	453	512	561	2,868	3,602	5,615	6,368	3,185	2,822	4,937	4,383	24
2003	33,699	404	558	497	2,642	3,350	5,717	5,740	2,981	2,784	4,775	4,235	16
2002	32,295	412	539	599	2,478	3,320	5,727	5,232	2,674	2,753	4,582	3,963	16
2001	30,201	410	547	613	2,318	2,860	5,262	4,781	2,390	2,586	4,532	3,885	17
2000	28,640	361	617	666	2,127	2,888	5,158	4,366	2,336	2,473	4,076	3,564	8
1999	28,701	335	613	690	2,032	2,972	5,107	4,125	2,248	2,672	4,289	3,605	13
Non-Hispanic 1/, female													
2005	25,725	367	296	253	766	1,411	2,713	3,274	1,768	1,976	4,984	7,912	5
2004	24,013	334	317	258	689	1,182	2,670	2,893	1,607	1,793	4,836	7,429	5
2003	22,851	299	323	232	654	1,130	2,499	2,598	1,414	1,830	4,628	7,241	3
2002	21,612	303	300	263	566	1,049	2,335	2,215	1,210	1,711	4,548	7,110	2
2001	19,982	299	335	278	498	856	1,979	1,735	1,139	1,717	4,383	6,759	4
2000	18,577	266	362	282	466	803	1,747	1,423	1,035	1,692	4,071	6,429	1
1999	18,638	226	384	312	434	790	1,656	1,322	970	1,728	4,278	6,536	2

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Nontransport accidents (W00-X59,Y86)													
Non-Hispanic white, both sexes													
2005 2/	54,621	508	592	493	3,236	4,598	7,171	8,355	4,434	4,198	9,261	11,761	14
2004 2/	51,151	491	595	488	3,035	4,060	6,960	7,592	3,892	3,939	8,932	11,141	26
2003 2/	48,595	423	633	440	2,812	3,758	6,830	6,702	3,629	3,923	8,595	10,832	18
2002	45,916	403	565	505	2,518	3,641	6,646	5,950	3,147	3,786	8,342	10,397	16
2001	42,509	426	653	566	2,304	3,027	5,846	5,093	2,839	3,681	8,113	9,944	17
2000	39,728	376	676	597	2,066	2,971	5,537	4,480	2,714	3,532	7,419	9,355	5
1999	39,523	337	693	617	1,917	2,966	5,419	4,181	2,509	3,671	7,737	9,467	9
Non-Hispanic white, male													
2005 2/	32,191	270	394	337	2,602	3,425	4,923	5,660	2,941	2,518	4,763	4,347	11
2004 2/	30,062	296	376	335	2,448	3,062	4,735	5,203	2,537	2,424	4,520	4,104	22
2003 2/	28,556	244	402	306	2,259	2,819	4,778	4,599	2,430	2,368	4,351	3,985	15
2002	27,024	244	359	354	2,039	2,772	4,734	4,175	2,134	2,330	4,175	3,694	14
2001	25,133	245	416	380	1,896	2,345	4,301	3,708	1,896	2,220	4,102	3,609	15
2000	23,615	222	433	428	1,704	2,343	4,178	3,361	1,857	2,095	3,683	3,307	4
1999	23,431	200	425	423	1,568	2,371	4,122	3,150	1,744	2,217	3,865	3,338	8
Non-Hispanic white, female													
2005 2/	22,430	238	198	156	634	1,173	2,248	2,695	1,493	1,680	4,498	7,414	3
2004 2/	21,089	195	219	153	587	998	2,225	2,389	1,355	1,515	4,412	7,037	4
2003 2/	20,039	179	231	134	553	939	2,052	2,103	1,199	1,555	4,244	6,847	3
2002	18,892	159	206	151	479	869	1,912	1,775	1,013	1,456	4,167	6,703	2
2001	17,376	181	237	186	408	682	1,545	1,385	943	1,461	4,011	6,335	2
2000	16,113	154	243	169	362	628	1,359	1,119	857	1,437	3,736	6,048	1
1999	16,092	137	268	194	349	595	1,297	1,031	765	1,454	3,872	6,129	1

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Nontransport accidents (W00-X59,Y86)													
Non-Hispanic black, both sexes													
2005 2/	7,654	270	198	238	438	675	1,191	1,756	926	606	767	585	4
2004 2/	6,702	266	188	273	386	570	1,116	1,472	755	542	646	487	1
2003 2/	6,470	255	208	241	369	563	1,155	1,427	651	547	581	472	1
2002	6,620	288	229	305	405	587	1,194	1,335	631	534	593	518	1
2001	6,436	254	204	276	391	542	1,211	1,274	590	529	632	532	1
2000	6,330	228	257	310	400	588	1,194	1,152	566	535	589	507	4
1999	6,723	207	260	341	432	663	1,171	1,145	617	629	693	560	5
Non-Hispanic black, male													
2005 2/	4,972	152	114	159	348	480	791	1,254	700	373	396	203	2
2004 2/	4,321	139	114	185	307	423	730	1,021	548	335	332	186	1
2003 2/	4,184	146	134	157	288	411	784	988	470	335	297	173	1
2002	4,377	154	152	214	343	442	829	941	464	331	309	197	1
2001	4,238	151	119	198	319	395	829	959	418	320	338	191	1
2000	4,253	124	155	215	326	441	853	883	422	319	314	197	4
1999	4,532	122	164	238	372	496	859	884	439	400	344	210	4
Non-Hispanic black, female													
2005 2/	2,682	118	84	79	90	195	400	502	226	233	371	382	2
2004 2/	2,381	127	74	88	79	147	386	451	207	207	314	301	-
2003 2/	2,286	109	74	84	81	152	371	439	181	212	284	299	-
2002	2,243	134	77	91	62	145	365	394	167	203	284	321	-
2001	2,198	103	85	78	72	147	382	315	172	209	294	341	-
2000	2,077	104	102	95	74	147	341	269	144	216	275	310	-
1999	2,191	85	96	103	60	167	312	261	178	229	349	350	1

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Nontransport accidents (W00-X59,Y86)													
Origin not stated 3/, both sexes													
2005	252	6	2	1	7	22	46	63	33	18	17	19	18
2004	218	1	4	3	9	22	31	54	24	15	16	19	20
2003	283	2	1	1	12	26	52	68	24	18	25	24	30
2002	284	3	3	6	14	29	49	51	30	22	34	26	17
2001	317	5	3	8	14	24	59	56	35	25	24	27	37
2000	321	4	4	5	13	28	60	79	27	20	25	27	29
1999	328	1	5	2	12	29	66	66	42	17	33	34	21
Origin not stated 3/, male													
2005	179	3	1	-	6	17	32	49	25	13	10	6	17
2004	164	1	2	1	8	18	25	46	18	13	7	8	17
2003	217	2	1	1	10	21	41	52	19	15	16	10	29
2002	212	2	2	4	10	24	40	39	30	19	21	10	11
2001	242	4	3	6	13	19	50	46	30	16	16	7	32
2000	236	3	1	2	11	23	48	60	22	14	12	12	28
1999	244	-	2	2	11	24	48	57	33	14	19	16	18
Origin not stated 3/, female													
2005	73	3	1	1	1	5	14	14	8	5	7	13	1
2004	54	-	2	2	1	4	6	8	6	2	9	11	3
2003	66	-	-	-	2	5	11	16	5	3	9	14	1
2002	72	1	1	2	4	5	9	12	-	3	13	16	6
2001	75	1	-	2	1	5	9	10	5	9	8	20	5
2000	85	1	3	3	2	5	12	19	5	6	13	15	1
1999	84	1	3	-	1	5	18	9	9	3	14	18	3

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Falls (W00-W19)													
All origins, both sexes													
2005	19,656	16	34	32	236	295	607	1,181	1,451	2,319	5,957	7,526	2
2004	18,807	23	47	37	241	320	659	1,184	1,393	2,255	5,682	6,962	4
2003	17,229	14	54	45	230	285	636	1,043	1,220	2,048	5,249	6,404	1
2002	16,257	16	37	42	247	307	664	1,017	1,089	1,967	4,880	5,990	1
2001	15,019	23	32	66	256	340	647	1,024	1,004	1,833	4,440	5,350	4
2000	13,322	8	36	37	237	303	608	871	949	1,660	3,841	4,772	-
1999	13,162	12	55	53	242	343	645	824	887	1,663	3,856	4,578	4
All origins, male													
2005	10,154	6	20	24	198	256	489	897	1,023	1,384	3,051	2,804	2
2004	9,856	13	31	22	206	274	526	912	987	1,394	2,858	2,630	3
2003	8,910	9	39	30	186	237	525	800	827	1,209	2,661	2,386	1
2002	8,463	9	28	27	211	269	522	772	771	1,202	2,444	2,208	-
2001	8,089	15	25	46	232	295	544	821	718	1,090	2,277	2,023	3
2000	7,122	6	23	30	203	272	496	690	680	991	1,947	1,784	-
1999	7,109	8	37	40	209	294	528	661	650	1,027	1,950	1,701	4
All origins, female													
2005	9,502	10	14	8	38	39	118	284	428	935	2,906	4,722	-
2004	8,951	10	16	15	35	46	133	272	406	861	2,824	4,332	1
2003	8,319	5	15	15	44	48	111	243	393	839	2,588	4,018	-
2002	7,794	7	9	15	36	38	142	245	318	765	2,436	3,782	1
2001	6,930	8	7	20	24	45	103	203	286	743	2,163	3,327	1
2000	6,200	2	13	7	34	31	112	181	269	669	1,894	2,988	-
1999	6,053	4	18	13	33	49	117	163	237	636	1,906	2,877	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Falls (W00-W19)													
Hispanic, both sexes													
2005	1,064	-	6	1	49	73	84	115	112	156	254	214	-
2004	978	6	7	2	44	80	112	94	109	130	211	183	-
2003	840	1	11	10	38	60	83	100	105	113	163	156	-
2002	832	4	9	6	51	69	94	90	87	118	175	129	-
2001	757	7	6	14	44	73	100	88	78	103	131	112	1
2000	693	3	9	3	46	73	66	77	74	86	134	122	-
1999	652	4	9	3	45	68	70	65	63	84	116	123	2
Hispanic, male													
2005	689	-	4	1	45	70	74	103	85	97	140	70	-
2004	653	4	4	1	42	74	96	84	86	79	107	76	-
2003	557	1	8	5	33	56	80	83	74	64	94	59	-
2002	563	3	7	3	49	65	81	70	65	72	90	58	-
2001	525	4	5	13	44	71	93	75	62	57	62	38	1
2000	485	2	8	3	44	70	62	69	53	60	62	52	-
1999	464	4	6	3	43	65	62	56	55	52	60	56	2
Hispanic, female													
2005	375	-	2	-	4	3	10	12	27	59	114	144	-
2004	325	2	3	1	2	6	16	10	23	51	104	107	-
2003	283	-	3	5	5	4	3	17	31	49	69	97	-
2002	269	1	2	3	2	4	13	20	22	46	85	71	-
2001	232	3	1	1	-	2	7	13	16	46	69	74	-
2000	208	1	1	-	2	3	4	8	21	26	72	70	-
1999	188	-	3	-	2	3	8	9	8	32	56	67	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Falls (W00-W19)													
Non-Hispanic 1/, both sexes													
2005	18,551	16	28	31	187	222	518	1,062	1,333	2,159	5,692	7,301	2
2004	17,780	17	40	34	196	237	546	1,079	1,279	2,120	5,464	6,764	4
2003	16,337	13	42	35	192	223	548	937	1,109	1,930	5,073	6,235	-
2002	15,365	12	28	36	196	238	568	916	995	1,842	4,685	5,849	-
2001	14,204	15	26	51	212	266	544	929	921	1,718	4,299	5,223	-
2000	12,585	5	27	34	191	229	537	784	869	1,569	3,700	4,640	-
1999	12,447	8	46	50	197	273	573	753	813	1,572	3,721	4,440	1
Non-Hispanic 1/, male													
2005	9,442	6	16	23	153	186	412	790	934	1,284	2,906	2,730	2
2004	9,170	9	27	21	163	198	429	818	896	1,312	2,748	2,546	3
2003	8,320	8	30	25	153	179	440	713	750	1,140	2,560	2,322	-
2002	7,856	6	21	24	162	204	439	691	699	1,123	2,342	2,145	-
2001	7,527	10	20	33	188	223	448	739	651	1,024	2,209	1,982	-
2000	6,603	4	15	27	159	201	429	612	622	928	1,881	1,725	-
1999	6,602	4	31	37	166	227	464	599	587	969	1,878	1,639	1
Non-Hispanic 1/, female													
2005	9,109	10	12	8	34	36	106	272	399	875	2,786	4,571	-
2004	8,610	8	13	13	33	39	117	261	383	808	2,716	4,218	1
2003	8,017	5	12	10	39	44	108	224	359	790	2,513	3,913	-
2002	7,509	6	7	12	34	34	129	225	296	719	2,343	3,704	-
2001	6,677	5	6	18	24	43	96	190	270	694	2,090	3,241	-
2000	5,982	1	12	7	32	28	108	172	247	641	1,819	2,915	-
1999	5,845	4	15	13	31	46	109	154	226	603	1,843	2,801	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Falls (W00-W19)													
Non-Hispanic white, both sexes													
2005 2/	17,042	13	14	23	170	178	437	880	1,136	1,930	5,298	6,961	2
2004 2/	16,379	12	26	28	173	191	443	896	1,081	1,891	5,147	6,487	4
2003 2/	14,995	9	28	29	167	190	446	744	943	1,711	4,747	5,981	-
2002	14,157	8	15	28	168	203	466	762	853	1,644	4,419	5,591	-
2001	13,037	8	23	42	193	226	435	768	780	1,540	4,044	4,978	-
2000	11,537	2	15	27	172	192	438	646	725	1,393	3,487	4,440	-
1999	11,295	4	34	39	168	216	477	593	653	1,379	3,500	4,231	1
Non-Hispanic white, male													
2005 2/	8,560	5	9	17	140	152	348	649	793	1,155	2,707	2,583	2
2004 2/	8,318	7	19	16	145	161	361	679	739	1,179	2,592	2,417	3
2003 2/	7,531	5	22	20	136	154	362	566	638	1,008	2,395	2,225	-
2002	7,107	3	11	19	137	175	359	574	596	1,002	2,199	2,032	-
2001	6,785	5	17	27	173	191	360	603	551	921	2,067	1,870	-
2000	5,949	1	6	22	144	169	354	505	519	829	1,764	1,636	-
1999	5,860	3	23	29	144	181	380	474	471	846	1,759	1,549	1
Non-Hispanic white, female													
2005 2/	8,482	8	5	6	30	26	89	231	343	775	2,591	4,378	-
2004 2/	8,061	5	7	12	28	30	82	217	342	712	2,555	4,070	1
2003 2/	7,464	4	6	9	31	36	84	178	305	703	2,352	3,756	-
2002	7,050	5	4	9	31	28	107	188	257	642	2,220	3,559	-
2001	6,252	3	6	15	20	35	75	165	229	619	1,977	3,108	-
2000	5,588	1	9	5	28	23	84	141	206	564	1,723	2,804	-
1999	5,435	1	11	10	24	35	97	119	182	533	1,741	2,682	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Falls (W00-W19)													
Non-Hispanic black, both sexes													
2005 2/	1,001	2	11	6	10	28	52	152	139	154	253	194	-
2004 2/	980	3	9	4	14	33	81	140	158	171	202	165	-
2003 2/	898	2	11	3	17	25	74	150	131	150	185	150	-
2002	821	3	7	6	17	23	75	128	110	135	154	163	-
2001	837	6	3	6	13	26	90	128	108	143	173	141	-
2000	745	2	12	7	12	22	71	111	114	134	137	123	-
1999	865	2	11	10	21	42	73	136	127	150	153	140	-
Non-Hispanic black, male													
2005 2/	583	-	4	4	8	19	42	120	102	90	127	67	-
2004 2/	614	-	7	3	9	27	56	105	128	102	105	72	-
2003 2/	519	1	7	3	10	18	57	112	86	87	85	53	-
2002	518	2	6	5	15	20	58	98	78	81	85	70	-
2001	537	4	3	4	9	19	72	109	76	85	97	59	-
2000	471	2	9	5	10	18	54	88	87	76	70	52	-
1999	554	1	8	7	16	34	65	107	90	100	77	49	-
Non-Hispanic black, female													
2005 2/	418	2	7	2	2	9	10	32	37	64	126	127	-
2004 2/	366	3	2	1	5	6	25	35	30	69	97	93	-
2003 2/	379	1	4	-	7	7	17	38	45	63	100	97	-
2002	303	1	1	1	2	3	17	30	32	54	69	93	-
2001	300	2	-	2	4	7	18	19	32	58	76	82	-
2000	274	-	3	2	2	4	17	23	27	58	67	71	-
1999	311	1	3	3	5	8	8	29	37	50	76	91	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Falls (W00-W19)													
Origin not stated 3/, both sexes													
2005	41	-	-	-	-	-	5	4	6	4	11	11	-
2004	49	-	-	1	1	3	1	11	5	5	7	15	-
2003	52	-	1	-	-	2	5	6	6	5	13	13	1
2002	60	-	-	-	-	-	2	11	7	7	20	12	1
2001	58	1	-	1	-	1	3	7	5	12	10	15	3
2000	44	-	-	-	-	1	5	10	6	5	7	10	-
1999	63	-	-	-	-	2	2	6	11	7	19	15	1
Origin not stated 3/, male													
2005	23	-	-	-	-	-	3	4	4	3	5	4	-
2004	33	-	-	-	1	2	1	10	5	3	3	8	-
2003	33	-	1	-	-	2	5	4	3	5	7	5	1
2002	44	-	-	-	-	-	2	11	7	7	12	5	-
2001	37	1	-	-	-	1	3	7	5	9	6	3	2
2000	34	-	-	-	-	1	5	9	5	3	4	7	-
1999	43	-	-	-	-	2	2	6	8	6	12	6	1
Origin not stated 3/, female													
2005	18	-	-	-	-	-	2	-	2	1	6	7	-
2004	16	-	-	1	-	1	-	1	-	2	4	7	-
2003	19	-	-	-	-	-	-	2	3	-	6	8	-
2002	16	-	-	-	-	-	-	-	-	-	8	7	1
2001	21	-	-	1	-	-	-	-	-	3	4	12	1
2000	10	-	-	-	-	-	-	1	1	2	3	3	-
1999	20	-	-	-	-	-	-	-	3	1	7	9	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Accidental discharge of firearms (W32-W34)													
All origins, both sexes													
2005	789	1	22	52	203	130	114	108	67	44	33	15	-
2004	649	1	14	48	172	100	97	106	35	40	28	8	-
2003	730	-	7	49	200	120	118	92	62	39	37	6	-
2002	762	1	11	48	210	143	123	95	56	36	32	7	-
2001	802	-	15	57	206	122	146	103	69	33	41	8	2
2000	776	1	18	67	202	131	153	78	50	41	24	11	-
1999	824	-	12	76	251	143	136	92	48	33	23	10	-
All origins, male													
2005	683	1	18	39	190	118	94	88	58	37	27	13	-
2004	573	-	11	40	160	92	87	90	29	33	23	8	-
2003	656	-	7	46	182	110	105	78	59	30	33	6	-
2002	667	1	7	39	195	129	98	84	50	32	26	6	-
2001	690	-	12	49	193	103	121	87	55	26	36	7	1
2000	671	-	11	62	183	109	125	69	44	34	23	11	-
1999	707	-	7	60	229	122	112	78	41	31	20	7	-
All origins, female													
2005	106	-	4	13	13	12	20	20	9	7	6	2	-
2004	76	1	3	8	12	8	10	16	6	7	5	-	-
2003	74	-	-	3	18	10	13	14	3	9	4	-	-
2002	95	-	4	9	15	14	25	11	6	4	6	1	-
2001	112	-	3	8	13	19	25	16	14	7	5	1	1
2000	105	1	7	5	19	22	28	9	6	7	1	-	-
1999	117	-	5	16	22	21	24	14	7	2	3	3	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Accidental discharge of firearms (W32-W34)													
Hispanic, both sexes													
2005	95	1	1	3	37	25	9	14	1	2	2	-	-
2004	56	-	-	5	23	13	5	9	1	-	-	-	-
2003	71	-	-	8	33	14	8	3	3	1	1	-	-
2002	60	-	3	5	24	18	5	3	1	-	1	-	-
2001	86	-	1	4	36	21	12	6	1	1	2	1	1
2000	59	-	2	6	25	12	11	2	1	-	-	-	-
1999	66	-	-	3	31	17	8	4	1	2	-	-	-
Hispanic, male													
2005	86	1	1	2	36	22	8	12	1	2	1	-	-
2004	51	-	-	4	23	11	5	7	1	-	-	-	-
2003	64	-	-	6	30	14	7	2	3	1	1	-	-
2002	51	-	1	3	21	18	3	3	1	-	1	-	-
2001	76	-	1	3	33	16	12	5	1	1	2	1	1
2000	50	-	1	4	24	10	9	2	-	-	-	-	-
1999	61	-	-	3	29	16	7	3	1	2	-	-	-
Hispanic, female													
2005	9	-	-	1	1	3	1	2	-	-	1	-	-
2004	5	-	-	1	-	2	-	2	-	-	-	-	-
2003	7	-	-	2	3	-	1	1	-	-	-	-	-
2002	9	-	2	2	3	-	2	-	-	-	-	-	-
2001	10	-	-	1	3	5	-	1	-	-	-	-	-
2000	9	-	1	2	1	2	2	-	1	-	-	-	-
1999	5	-	-	-	2	1	1	1	-	-	-	-	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Accidental discharge of firearms (W32-W34)													
Non-Hispanic 1/, both sexes													
2005	692	-	21	49	166	104	104	94	66	42	31	15	-
2004	593	1	14	43	149	87	92	97	34	40	28	8	-
2003	658	-	7	41	167	106	109	89	59	38	36	6	-
2002	700	1	8	43	185	125	118	92	55	36	30	7	-
2001	715	-	14	53	169	101	134	97	68	32	39	7	1
2000	717	1	16	61	177	119	142	76	49	41	24	11	-
1999	758	-	12	73	220	126	128	88	47	31	23	10	-
Non-Hispanic 1/, male													
2005	595	-	17	37	154	95	85	76	57	35	26	13	-
2004	522	-	11	36	137	81	82	83	28	33	23	8	-
2003	591	-	7	40	152	96	97	76	56	29	32	6	-
2002	614	1	6	36	173	111	95	81	49	32	24	6	-
2001	613	-	11	46	159	87	109	82	54	25	34	6	-
2000	621	-	10	58	159	99	116	67	44	34	23	11	-
1999	646	-	7	57	200	106	105	75	40	29	20	7	-
Non-Hispanic 1/, female													
2005	97	-	4	12	12	9	19	18	9	7	5	2	-
2004	71	1	3	7	12	6	10	14	6	7	5	-	-
2003	67	-	-	1	15	10	12	13	3	9	4	-	-
2002	86	-	2	7	12	14	23	11	6	4	6	1	-
2001	102	-	3	7	10	14	25	15	14	7	5	1	1
2000	96	1	6	3	18	20	26	9	5	7	1	-	-
1999	112	-	5	16	20	20	23	13	7	2	3	3	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Accidental discharge of firearms (W32-W34)													
Non-Hispanic white, both sexes													
2005 2/	508	-	9	34	103	57	82	79	64	39	29	12	-
2004 2/	451	1	6	32	97	52	76	90	30	36	24	7	-
2003 2/	515	-	4	30	115	69	95	77	56	33	32	4	-
2002	531	1	6	31	118	82	97	78	49	35	27	7	-
2001	591	-	9	41	117	82	116	92	61	29	36	7	1
2000	555	1	9	46	117	83	118	65	46	37	22	11	-
1999	585	-	5	54	145	98	108	77	40	28	20	10	-
Non-Hispanic white, male													
2005 2/	432	-	8	27	99	49	64	64	55	32	24	10	-
2004 2/	395	-	5	27	89	48	66	79	25	30	19	7	-
2003 2/	461	-	4	29	103	61	87	65	54	26	28	4	-
2002	459	1	4	25	110	70	77	69	45	31	21	6	-
2001	506	-	7	38	109	71	96	78	48	22	31	6	-
2000	483	-	5	44	106	66	99	59	41	31	21	11	-
1999	495	-	3	43	131	80	87	65	36	26	17	7	-
Non-Hispanic white, female													
2005 2/	76	-	1	7	4	8	18	15	9	7	5	2	-
2004 2/	56	1	1	5	8	4	10	11	5	6	5	-	-
2003 2/	54	-	-	1	12	8	8	12	2	7	4	-	-
2002	72	-	2	6	8	12	20	9	4	4	6	1	-
2001	85	-	2	3	8	11	20	14	13	7	5	1	1
2000	72	1	4	2	11	17	19	6	5	6	1	-	-
1999	90	-	2	11	14	18	21	12	4	2	3	3	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Accidental discharge of firearms (W32-W34)													
Non-Hispanic black, both sexes													
2005 2/	150	-	12	14	49	39	17	10	2	3	1	3	-
2004 2/	127	-	7	9	45	33	13	7	4	4	4	1	-
2003 2/	127	-	3	9	49	34	12	7	3	5	3	2	-
2002	153	-	2	9	60	40	20	13	6	-	3	-	-
2001	110	-	5	8	48	17	15	5	6	3	3	-	-
2000	137	-	5	12	52	32	19	10	3	3	1	-	-
1999	156	-	7	17	67	23	18	11	7	3	3	-	-
Non-Hispanic black, male													
2005 2/	134	-	9	9	44	38	17	8	2	3	1	3	-
2004 2/	115	-	6	7	43	31	13	4	3	3	4	1	-
2003 2/	115	-	3	9	46	33	8	6	2	3	3	2	-
2002	140	-	2	8	57	38	17	11	4	-	3	-	-
2001	97	-	4	6	46	14	12	4	5	3	3	-	-
2000	117	-	3	12	46	29	14	7	3	2	1	-	-
1999	135	-	4	12	61	22	16	10	4	3	3	-	-
Non-Hispanic black, female													
2005 2/	16	-	3	5	5	1	-	2	-	-	-	-	-
2004 2/	12	-	1	2	2	2	-	3	1	1	-	-	-
2003 2/	12	-	-	-	3	1	4	1	1	2	-	-	-
2002	13	-	-	1	3	2	3	2	2	-	-	-	-
2001	13	-	1	2	2	3	3	1	1	-	-	-	-
2000	20	-	2	-	6	3	5	3	-	1	-	-	-
1999	21	-	3	5	6	1	2	1	3	-	-	-	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Accidental discharge of firearms (W32-W34)													
Origin not stated 3/, both sexes													
2005	2	-	-	-	-	1	1	-	-	-	-	-	-
2004	-	-	-	-	-	-	-	-	-	-	-	-	-
2003	1	-	-	-	-	-	1	-	-	-	-	-	-
2002	2	-	-	-	1	-	-	-	-	-	1	-	-
2001	1	-	-	-	1	-	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-	-	-	-
Origin not stated 3/, male													
2005	2	-	-	-	-	1	1	-	-	-	-	-	-
2004	-	-	-	-	-	-	-	-	-	-	-	-	-
2003	1	-	-	-	-	-	1	-	-	-	-	-	-
2002	2	-	-	-	1	-	-	-	-	-	1	-	-
2001	1	-	-	-	1	-	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-	-	-	-
Origin not stated 3/, female													
2005	-	-	-	-	-	-	-	-	-	-	-	-	-
2004	-	-	-	-	-	-	-	-	-	-	-	-	-
2003	-	-	-	-	-	-	-	-	-	-	-	-	-
2002	-	-	-	-	-	-	-	-	-	-	-	-	-
2001	-	-	-	-	-	-	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-	-	-	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Accidental drowning and submersion (W65-W74)													
All origins, both sexes													
2005	3,582	64	493	253	649	385	497	492	266	193	180	92	18
2004	3,308	62	430	269	574	385	435	444	271	188	158	78	14
2003	3,306	58	456	268	567	356	462	397	265	202	162	83	30
2002	3,447	63	454	321	629	433	526	374	212	169	167	79	20
2001	3,281	68	458	333	596	374	462	359	206	159	157	75	34
2000	3,482	75	493	375	646	419	480	354	217	179	156	64	24
1999	3,529	68	490	369	647	446	475	370	219	185	158	68	34
All origins, male													
2005	2,818	27	336	184	588	328	410	393	218	135	121	62	16
2004	2,594	38	265	194	522	332	367	358	208	140	113	43	14
2003	2,632	34	301	199	525	323	392	320	211	145	104	48	30
2002	2,761	40	312	244	578	380	439	301	161	132	106	50	18
2001	2,560	30	291	237	544	328	376	299	155	124	102	45	29
2000	2,735	43	316	267	583	364	406	299	179	133	88	34	23
1999	2,794	39	310	275	592	397	402	311	170	129	111	28	30
All origins, female													
2005	764	37	157	69	61	57	87	99	48	58	59	30	2
2004	714	24	165	75	52	53	68	86	63	48	45	35	-
2003	674	24	155	69	42	33	70	77	54	57	58	35	-
2002	686	23	142	77	51	53	87	73	51	37	61	29	2
2001	721	38	167	96	52	46	86	60	51	35	55	30	5
2000	747	32	177	108	63	55	74	55	38	46	68	30	1
1999	735	29	180	94	55	49	73	59	49	56	47	40	4

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Accidental drowning and submersion (W65-W74)													
Hispanic, both sexes													
2005	518	15	104	29	159	77	54	34	22	9	3	4	8
2004	427	17	78	31	102	73	61	30	10	9	6	4	6
2003	458	12	82	45	112	85	51	27	15	14	6	2	7
2002	464	18	65	29	143	80	55	30	15	12	10	2	5
2001	421	16	78	40	101	74	57	25	7	5	4	2	12
2000	433	9	75	43	147	62	41	20	11	12	4	1	8
1999	480	16	78	46	145	79	45	25	8	7	6	-	25
Hispanic, male													
2005	430	6	73	19	145	71	47	33	16	7	2	4	7
2004	355	7	48	19	95	72	53	28	9	9	5	4	6
2003	399	7	67	33	109	79	47	25	14	7	3	1	7
2002	409	13	48	23	137	78	48	28	10	10	7	2	5
2001	348	9	53	28	99	65	48	21	6	3	3	2	11
2000	350	6	44	22	135	60	36	18	8	10	3	-	8
1999	405	12	43	35	140	75	38	23	7	7	4	-	21
Hispanic, female													
2005	88	9	31	10	14	6	7	1	6	2	1	-	1
2004	72	10	30	12	7	1	8	2	1	-	1	-	-
2003	59	5	15	12	3	6	4	2	1	7	3	1	-
2002	55	5	17	6	6	2	7	2	5	2	3	-	-
2001	73	7	25	12	2	9	9	4	1	2	1	-	1
2000	83	3	31	21	12	2	5	2	3	2	1	1	-
1999	75	4	35	11	5	4	7	2	1	-	2	-	4

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Accidental drowning and submersion (W65-W74)													
Non-Hispanic 1/, both sexes													
2005	3,047	49	388	223	488	307	443	456	244	184	176	88	1
2004	2,856	44	350	237	469	310	373	408	257	178	152	74	4
2003	2,816	46	374	222	452	271	407	364	248	187	156	81	8
2002	2,952	45	387	289	485	347	467	342	195	156	155	77	7
2001	2,832	51	380	291	493	300	403	329	198	153	153	73	8
2000	3,018	65	417	331	496	355	437	332	205	166	152	61	1
1999	3,029	52	409	323	500	363	429	343	210	178	152	67	3
Non-Hispanic 1/, male													
2005	2,375	21	263	165	441	256	363	359	202	128	119	58	-
2004	2,220	30	217	174	425	258	313	325	197	130	108	39	4
2003	2,202	27	234	165	413	244	342	289	195	137	101	47	8
2002	2,326	27	262	219	440	298	387	271	149	121	97	48	7
2001	2,188	21	238	207	443	263	326	275	148	120	99	43	5
2000	2,358	37	272	244	445	302	368	279	170	122	85	33	1
1999	2,370	27	265	240	450	318	363	286	162	122	107	27	3
Non-Hispanic 1/, female													
2005	672	28	125	58	47	51	80	97	42	56	57	30	1
2004	636	14	133	63	44	52	60	83	60	48	44	35	-
2003	614	19	140	57	39	27	65	75	53	50	55	34	-
2002	626	18	125	70	45	49	80	71	46	35	58	29	-
2001	644	30	142	84	50	37	77	54	50	33	54	30	3
2000	660	28	145	87	51	53	69	53	35	44	67	28	-
1999	659	25	144	83	50	45	66	57	48	56	45	40	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Accidental drowning and submersion (W65-W74)													
Non-Hispanic white, both sexes													
2005 2/	2,305	41	316	118	315	229	338	372	202	151	145	77	1
2004 2/	2,133	34	291	112	306	222	289	328	207	148	130	62	4
2003 2/	2,152	32	306	118	306	187	320	297	208	163	139	68	8
2002	2,143	32	303	134	316	245	360	260	165	126	131	64	7
2001	2,082	36	321	135	322	202	306	265	158	133	134	66	4
2000	2,235	38	342	176	316	258	344	255	175	144	134	53	-
1999	2,230	30	337	142	317	250	349	291	174	147	132	59	2
Non-Hispanic white, male													
2005 2/	1,763	15	215	79	285	187	270	290	163	105	101	53	-
2004 2/	1,646	21	179	84	275	181	242	261	158	111	96	34	4
2003 2/	1,656	17	187	86	279	166	267	230	160	121	93	42	8
2002	1,669	18	203	96	284	209	294	205	125	99	85	44	7
2001	1,577	16	205	87	284	171	245	220	114	105	88	39	3
2000	1,737	22	226	129	278	222	287	212	150	104	78	29	-
1999	1,708	14	217	99	279	216	290	241	133	99	93	25	2
Non-Hispanic white, female													
2005 2/	542	26	101	39	30	42	68	82	39	46	44	24	1
2004 2/	487	13	112	28	31	41	47	67	49	37	34	28	-
2003 2/	496	15	119	32	27	21	53	67	48	42	46	26	-
2002	474	14	100	38	32	36	66	55	40	27	46	20	-
2001	505	20	116	48	38	31	61	45	44	28	46	27	1
2000	498	16	116	47	38	36	57	43	25	40	56	24	-
1999	522	16	120	43	38	34	59	50	41	48	39	34	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Accidental drowning and submersion (W65-W74)													
Non-Hispanic black, both sexes													
2005 2/	543	8	56	87	131	58	72	58	31	25	12	5	-
2004 2/	519	8	46	103	115	59	61	64	28	19	9	7	-
2003 2/	466	13	52	87	100	59	48	45	33	16	8	5	-
2002	604	10	62	134	126	74	74	57	21	18	17	11	-
2001	570	11	47	133	133	62	70	51	32	13	13	4	1
2000	580	25	54	134	128	58	77	52	17	15	12	7	1
1999	624	21	59	158	137	86	58	36	28	21	15	5	-
Non-Hispanic black, male													
2005 2/	463	6	36	72	122	53	62	53	29	19	8	3	-
2004 2/	416	7	35	74	106	49	52	50	22	14	6	1	-
2003 2/	394	9	37	66	94	54	45	42	29	10	5	3	-
2002	502	6	43	107	122	65	66	47	18	15	9	4	-
2001	471	3	28	102	126	61	58	42	26	12	9	3	1
2000	470	14	33	101	123	52	67	45	13	13	4	4	1
1999	521	12	40	124	128	78	53	34	22	17	11	2	-
Non-Hispanic black, female													
2005 2/	80	2	20	15	9	5	10	5	2	6	4	2	-
2004 2/	103	1	11	29	9	10	9	14	6	5	3	6	-
2003 2/	72	4	15	21	6	5	3	3	4	6	3	2	-
2002	102	4	19	27	4	9	8	10	3	3	8	7	-
2001	99	8	19	31	7	1	12	9	6	1	4	1	-
2000	110	11	21	33	5	6	10	7	4	2	8	3	-
1999	103	9	19	34	9	8	5	2	6	4	4	3	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Accidental drowning and submersion (W65-W74)													
Origin not stated 3/, both sexes													
2005	17	-	1	1	2	1	-	2	-	-	1	-	9
2004	25	1	2	1	3	2	1	6	4	1	-	-	4
2003	32	-	-	1	3	-	4	6	2	1	-	-	15
2002	31	-	2	3	1	6	4	2	2	1	2	-	8
2001	28	1	-	2	2	-	2	5	1	1	-	-	14
2000	31	1	1	1	3	2	2	2	1	1	-	2	15
1999	20	-	3	-	2	4	1	2	1	-	-	1	6
Origin not stated 3/, male													
2005	13	-	-	-	2	1	-	1	-	-	-	-	9
2004	19	1	-	1	2	2	1	5	2	1	-	-	4
2003	31	-	-	1	3	-	3	6	2	1	-	-	15
2002	26	-	2	2	1	4	4	2	2	1	2	-	6
2001	24	-	-	2	2	-	2	3	1	1	-	-	13
2000	27	-	-	1	3	2	2	2	1	1	-	1	14
1999	19	-	2	-	2	4	1	2	1	-	-	1	6
Origin not stated 3/, female													
2005	4	-	1	1	-	-	-	1	-	-	1	-	-
2004	6	-	2	-	1	-	-	1	2	-	-	-	-
2003	1	-	-	-	-	-	1	-	-	-	-	-	-
2002	5	-	-	1	-	2	-	-	-	-	-	-	2
2001	4	1	-	-	-	-	-	2	-	-	-	-	1
2000	4	1	1	-	-	-	-	-	-	-	-	1	1
1999	1	-	1	-	-	-	-	-	-	-	-	-	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Accidental exposure to smoke, fire and flames (X00-X09)													
All origins, both sexes													
2005	3,197	34	204	222	168	224	334	497	398	390	468	254	4
2004	3,229	26	223	255	186	240	313	490	412	349	459	273	3
2003	3,369	30	224	215	199	252	395	526	392	419	460	253	4
2002	3,159	36	221	253	193	266	361	448	340	385	432	221	3
2001	3,309	50	225	252	209	247	441	419	378	411	423	251	3
2000	3,377	37	290	266	192	241	402	439	369	401	472	263	5
1999	3,348	41	302	262	201	282	421	368	319	416	487	245	4
All origins, male													
2005	1,886	16	124	125	103	137	226	325	244	224	245	113	4
2004	1,902	14	120	152	107	165	209	329	251	210	234	109	2
2003	1,948	8	131	112	121	154	257	329	254	244	228	107	3
2002	1,935	21	126	151	132	170	251	292	219	242	231	97	3
2001	1,998	30	142	152	137	161	308	288	236	229	204	108	3
2000	2,025	20	174	145	121	157	268	303	241	232	244	115	5
1999	2,025	21	173	147	133	197	300	254	194	244	247	111	4
All origins, female													
2005	1,311	18	80	97	65	87	108	172	154	166	223	141	-
2004	1,327	12	103	103	79	75	104	161	161	139	225	164	1
2003	1,421	22	93	103	78	98	138	197	138	175	232	146	1
2002	1,224	15	95	102	61	96	110	156	121	143	201	124	-
2001	1,311	20	83	100	72	86	133	131	142	182	219	143	-
2000	1,352	17	116	121	71	84	134	136	128	169	228	148	-
1999	1,323	20	129	115	68	85	121	114	125	172	240	134	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Accidental exposure to smoke, fire and flames (X00-X09)													
Hispanic, both sexes													
2005	211	6	30	17	16	25	24	24	16	19	25	8	1
2004	179	5	19	13	15	24	17	18	9	22	26	11	-
2003	203	4	27	26	15	25	24	32	14	12	15	9	-
2002	223	5	36	27	25	17	27	22	17	20	16	10	1
2001	176	9	18	23	19	21	28	17	13	9	9	10	-
2000	178	4	21	19	21	19	25	11	17	18	17	6	-
1999	192	5	41	23	24	17	21	16	14	11	15	5	-
Hispanic, male													
2005	125	3	18	7	13	16	15	16	11	11	10	4	1
2004	107	3	9	8	9	16	14	12	6	14	12	4	-
2003	128	2	11	12	9	18	17	26	10	9	9	5	-
2002	144	3	18	17	19	9	21	18	12	14	8	4	1
2001	109	5	13	11	10	16	19	11	9	6	5	4	-
2000	105	2	11	10	11	14	17	7	9	10	10	4	-
1999	127	3	23	12	16	12	16	12	11	8	9	5	-
Hispanic, female													
2005	86	3	12	10	3	9	9	8	5	8	15	4	-
2004	72	2	10	5	6	8	3	6	3	8	14	7	-
2003	75	2	16	14	6	7	7	6	4	3	6	4	-
2002	79	2	18	10	6	8	6	4	5	6	8	6	-
2001	67	4	5	12	9	5	9	6	4	3	4	6	-
2000	73	2	10	9	10	5	8	4	8	8	7	2	-
1999	65	2	18	11	8	5	5	4	3	3	6	-	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Accidental exposure to smoke, fire and flames (X00-X09)													
Non-Hispanic 1/, both sexes													
2005	2,971	25	173	205	152	199	307	472	380	369	442	246	1
2004	3,038	21	204	242	171	213	293	470	401	327	432	262	2
2003	3,147	26	197	189	183	227	367	491	375	404	443	243	2
2002	2,918	31	184	225	168	247	334	424	317	362	414	211	1
2001	3,106	41	205	226	190	223	410	398	361	398	412	240	2
2000	3,178	31	268	246	169	222	375	426	349	382	451	256	3
1999	3,133	36	260	239	173	264	395	347	302	404	470	240	3
Non-Hispanic 1/, male													
2005	1,751	11	105	118	90	121	210	309	231	212	234	109	1
2004	1,785	11	111	144	98	147	192	315	243	196	221	105	2
2003	1,809	6	120	100	111	136	240	302	241	233	218	101	1
2002	1,780	18	108	133	113	159	230	274	201	227	223	93	1
2001	1,873	25	127	139	127	145	286	274	224	222	199	103	2
2000	1,907	16	163	135	108	143	250	295	230	222	232	110	3
1999	1,879	18	150	135	113	184	279	238	181	235	237	106	3
Non-Hispanic 1/, female													
2005	1,220	14	68	87	62	78	97	163	149	157	208	137	-
2004	1,253	10	93	98	73	66	101	155	158	131	211	157	-
2003	1,338	20	77	89	72	91	127	189	134	171	225	142	1
2002	1,138	13	76	92	55	88	104	150	116	135	191	118	-
2001	1,233	16	78	87	63	78	124	124	137	176	213	137	-
2000	1,271	15	105	111	61	79	125	131	119	160	219	146	-
1999	1,254	18	110	104	60	80	116	109	121	169	233	134	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Accidental exposure to smoke, fire and flames (X00-X09)													
Non-Hispanic white, both sexes													
2005 2/	2,147	15	96	110	108	138	233	366	268	284	341	187	1
2004 2/	2,151	10	120	123	126	154	215	338	296	232	331	205	1
2003 2/	2,313	18	118	89	141	170	292	362	286	305	340	190	2
2002	2,055	16	90	112	113	171	258	317	227	268	325	157	1
2001	2,204	26	124	136	139	153	297	284	261	291	311	180	2
2000	2,194	17	139	139	120	151	267	320	242	278	328	190	3
1999	2,172	21	137	136	125	199	286	253	218	281	343	171	2
Non-Hispanic white, male													
2005 2/	1,270	6	64	68	65	89	156	237	157	160	184	83	1
2004 2/	1,265	6	67	72	74	108	137	222	182	139	173	84	1
2003 2/	1,344	5	74	46	91	102	197	227	183	178	159	81	1
2002	1,271	12	52	73	78	115	173	204	146	166	182	69	1
2001	1,322	14	82	82	96	105	207	190	160	157	149	78	2
2000	1,330	7	88	76	79	99	179	222	160	159	172	86	3
1999	1,325	11	80	76	83	145	204	173	136	160	174	81	2
Non-Hispanic white, female													
2005 2/	877	9	32	42	43	49	77	129	111	124	157	104	-
2004 2/	886	4	53	51	52	46	78	116	114	93	158	121	-
2003 2/	969	13	44	43	50	68	95	135	103	127	181	109	1
2002	784	4	38	39	35	56	85	113	81	102	143	88	-
2001	882	12	42	54	43	48	90	94	101	134	162	102	-
2000	864	10	51	63	41	52	88	98	82	119	156	104	-
1999	847	10	57	60	42	54	82	80	82	121	169	90	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Accidental exposure to smoke, fire and flames (X00-X09)													
Non-Hispanic black, both sexes													
2005 2/	731	9	69	81	36	52	63	98	102	75	93	53	-
2004 2/	791	11	73	105	38	51	65	118	98	84	94	53	1
2003 2/	767	7	75	94	38	47	66	122	79	90	99	50	-
2002	765	13	88	95	48	61	60	98	84	83	82	53	-
2001	829	14	80	85	42	63	103	105	89	99	92	57	-
2000	915	14	120	101	42	64	97	97	100	98	117	65	-
1999	887	15	110	102	40	52	100	86	80	113	120	68	1
Non-Hispanic black, male													
2005 2/	433	5	36	44	20	27	45	70	69	46	48	23	-
2004 2/	468	5	38	64	21	33	44	85	57	54	47	19	1
2003 2/	430	1	42	50	18	28	39	72	53	53	56	18	-
2002	454	6	53	53	31	35	45	64	52	53	38	24	-
2001	501	10	44	53	24	34	73	79	56	61	44	23	-
2000	539	9	70	57	25	39	63	67	67	60	58	24	-
1999	507	7	63	58	24	29	69	59	44	69	59	25	1
Non-Hispanic black, female													
2005 2/	298	4	33	37	16	25	18	28	33	29	45	30	-
2004 2/	323	6	35	41	17	18	21	33	41	30	47	34	-
2003 2/	337	6	33	44	20	19	27	50	26	37	43	32	-
2002	311	7	35	42	17	26	15	34	32	30	44	29	-
2001	328	4	36	32	18	29	30	26	33	38	48	34	-
2000	376	5	50	44	17	25	34	30	33	38	59	41	-
1999	380	8	47	44	16	23	31	27	36	44	61	43	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Accidental exposure to smoke, fire and flames (X00-X09)													
Origin not stated 3/, both sexes													
2005	15	3	1	-	-	-	3	1	2	2	1	-	2
2004	12	-	-	-	-	3	3	2	2	-	1	-	1
2003	19	-	-	-	1	-	4	3	3	3	2	1	2
2002	18	-	1	1	-	2	-	2	6	3	2	-	1
2001	27	-	2	3	-	3	3	4	4	4	2	1	1
2000	21	2	1	1	2	-	2	2	3	1	4	1	2
1999	23	-	1	-	4	1	5	5	3	1	2	-	1
Origin not stated 3/, male													
2005	10	2	1	-	-	-	1	-	2	1	1	-	2
2004	10	-	-	-	-	2	3	2	2	-	1	-	-
2003	11	-	-	-	1	-	-	1	3	2	1	1	2
2002	11	-	-	1	-	2	-	-	6	1	-	-	1
2001	16	-	2	2	-	-	3	3	3	1	-	1	1
2000	13	2	-	-	2	-	1	1	2	-	2	1	2
1999	19	-	-	-	4	1	5	4	2	1	1	-	1
Origin not stated 3/, female													
2005	5	1	-	-	-	-	2	1	-	1	-	-	-
2004	2	-	-	-	-	1	-	-	-	-	-	-	1
2003	8	-	-	-	-	-	4	2	-	1	1	-	-
2002	7	-	1	-	-	-	-	2	-	2	2	-	-
2001	11	-	-	1	-	3	-	1	1	3	2	-	-
2000	8	-	1	1	-	-	1	1	1	1	2	-	-
1999	4	-	1	-	-	-	-	1	1	-	1	-	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Accidental poisoning and exposure to noxious substances (X40-X49)													
All origins, both sexes													
2005	23,618	20	21	51	2,484	4,386	6,729	6,983	2,007	435	317	179	6
2004	20,950	13	18	55	2,259	3,641	6,444	6,033	1,577	421	315	165	9
2003	19,457	20	49	61	1,999	3,435	6,230	5,434	1,370	403	284	166	6
2002	17,550	26	31	43	1,679	3,116	6,007	4,682	1,098	383	295	181	9
2001	14,078	15	31	50	1,362	2,507	5,036	3,547	798	291	279	152	10
2000	12,757	14	32	45	1,160	2,380	4,663	3,061	688	278	245	184	7
1999	12,186	12	34	40	964	2,355	4,549	2,844	668	300	265	143	12
All origins, male													
2005	15,884	11	12	31	1,940	3,193	4,423	4,537	1,269	228	161	74	5
2004	13,934	5	13	30	1,765	2,673	4,176	3,910	929	222	143	64	4
2003	13,176	9	27	31	1,562	2,514	4,166	3,598	841	223	142	58	5
2002	12,059	21	14	28	1,306	2,319	4,122	3,136	690	209	136	71	7
2001	9,885	8	21	23	1,080	1,862	3,522	2,503	500	170	137	49	10
2000	9,138	11	21	27	920	1,820	3,384	2,223	401	147	102	76	6
1999	8,887	11	17	25	754	1,828	3,349	2,106	448	172	116	53	8
All origins, female													
2005	7,734	9	9	20	544	1,193	2,306	2,446	738	207	156	105	1
2004	7,016	8	5	25	494	968	2,268	2,123	648	199	172	101	5
2003	6,281	11	22	30	437	921	2,064	1,836	529	180	142	108	1
2002	5,491	5	17	15	373	797	1,885	1,546	408	174	159	110	2
2001	4,193	7	10	27	282	645	1,514	1,044	298	121	142	103	-
2000	3,619	3	11	18	240	560	1,279	838	287	131	143	108	1
1999	3,299	1	17	15	210	527	1,200	738	220	128	149	90	4

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Accidental poisoning and exposure to noxious substances (X40-X49)													
Hispanic, both sexes													
2005	2,018	3	3	7	243	482	635	467	130	29	11	8	-
2004	1,716	2	3	10	202	372	567	390	119	30	15	5	1
2003	1,802	4	13	5	219	421	610	401	91	22	10	6	-
2002	1,652	6	4	9	195	364	586	368	79	26	13	2	-
2001	1,324	1	5	10	144	334	475	263	64	19	7	1	1
2000	1,308	2	4	4	132	344	493	251	53	15	5	5	-
1999	1,502	4	3	6	146	383	585	289	58	18	5	2	3
Hispanic, male													
2005	1,603	1	2	5	197	399	505	361	106	19	5	3	-
2004	1,327	1	2	5	158	316	439	287	89	18	8	3	1
2003	1,428	2	5	3	179	347	483	313	73	17	3	3	-
2002	1,324	5	-	4	161	312	466	288	60	18	9	1	-
2001	1,061	1	3	2	111	274	385	218	50	13	3	-	1
2000	1,090	2	2	2	113	295	415	211	35	10	2	3	-
1999	1,271	4	3	4	123	332	493	240	51	15	2	1	3
Hispanic, female													
2005	415	2	1	2	46	83	130	106	24	10	6	5	-
2004	389	1	1	5	44	56	128	103	30	12	7	2	-
2003	374	2	8	2	40	74	127	88	18	5	7	3	-
2002	328	1	4	5	34	52	120	80	19	8	4	1	-
2001	263	-	2	8	33	60	90	45	14	6	4	1	-
2000	218	-	2	2	19	49	78	40	18	5	3	2	-
1999	231	-	-	2	23	51	92	49	7	3	3	1	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Accidental poisoning and exposure to noxious substances (X40-X49)													
Non-Hispanic 1/, both sexes													
2005	21,488	17	18	44	2,236	3,889	6,064	6,472	1,864	402	306	170	6
2004	19,153	11	14	45	2,054	3,259	5,852	5,615	1,450	390	299	160	4
2003	17,537	16	36	56	1,776	2,994	5,590	4,986	1,269	377	273	160	4
2002	15,789	20	27	34	1,476	2,735	5,382	4,290	1,011	351	280	179	4
2001	12,631	14	26	40	1,210	2,157	4,519	3,250	719	270	271	151	4
2000	11,311	12	27	38	1,022	2,018	4,131	2,758	624	259	240	179	3
1999	10,542	8	31	33	814	1,955	3,907	2,516	596	280	258	139	5
Non-Hispanic 1/, male													
2005	14,198	10	10	26	1,739	2,783	3,896	4,142	1,153	207	156	71	5
2004	12,541	4	10	25	1,604	2,349	3,718	3,599	834	203	134	61	-
2003	11,655	7	22	28	1,381	2,151	3,659	3,250	759	202	138	55	3
2002	10,649	16	14	24	1,140	1,993	3,626	2,831	622	185	125	70	3
2001	8,720	7	18	21	962	1,574	3,103	2,257	437	155	133	49	4
2000	7,944	9	18	24	802	1,512	2,938	1,975	357	134	100	73	2
1999	7,510	7	14	20	627	1,482	2,817	1,833	387	155	114	51	3
Non-Hispanic 1/, female													
2005	7,290	7	8	18	497	1,106	2,168	2,330	711	195	150	99	1
2004	6,612	7	4	20	450	910	2,134	2,016	616	187	165	99	4
2003	5,882	9	14	28	395	843	1,931	1,736	510	175	135	105	1
2002	5,140	4	13	10	336	742	1,756	1,459	389	166	155	109	1
2001	3,911	7	8	19	248	583	1,416	993	282	115	138	102	-
2000	3,367	3	9	14	220	506	1,193	783	267	125	140	106	1
1999	3,032	1	17	13	187	473	1,090	683	209	125	144	88	2

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Accidental poisoning and exposure to noxious substances (X40-X49)													
Non-Hispanic white, both sexes													
2005 2/	18,126	9	8	32	2,071	3,398	5,151	5,269	1,448	313	267	155	5
2004 2/	16,313	8	8	30	1,900	2,888	5,009	4,613	1,131	318	259	146	3
2003 2/	14,829	8	19	31	1,659	2,611	4,708	4,058	1,044	292	253	143	3
2002	13,165	7	15	28	1,357	2,375	4,496	3,402	772	299	250	162	2
2001	10,269	8	16	30	1,078	1,836	3,698	2,468	546	226	229	130	4
2000	9,124	8	15	24	916	1,683	3,331	2,071	482	216	224	152	2
1999	8,311	6	17	21	694	1,581	3,096	1,867	444	224	238	120	3
Non-Hispanic white, male													
2005 2/	11,949	6	3	21	1,626	2,448	3,334	3,327	838	142	132	67	5
2004 2/	10,684	2	4	20	1,493	2,099	3,204	2,937	602	155	112	56	-
2003 2/	9,855	6	11	17	1,292	1,892	3,098	2,625	596	144	127	45	2
2002	8,849	7	6	21	1,048	1,737	3,045	2,225	436	150	108	65	1
2001	7,111	4	12	17	864	1,358	2,579	1,688	304	129	109	43	4
2000	6,413	6	9	16	736	1,272	2,399	1,464	251	104	93	62	1
1999	5,921	5	6	13	536	1,217	2,265	1,337	263	124	105	48	2
Non-Hispanic white, female													
2005 2/	6,177	3	5	11	445	950	1,817	1,942	610	171	135	88	-
2004 2/	5,629	6	4	10	407	789	1,805	1,676	529	163	147	90	3
2003 2/	4,974	2	8	14	367	719	1,610	1,433	448	148	126	98	1
2002	4,316	-	9	7	309	638	1,451	1,177	336	149	142	97	1
2001	3,158	4	4	13	214	478	1,119	780	242	97	120	87	-
2000	2,711	2	6	8	180	411	932	607	231	112	131	90	1
1999	2,390	1	11	8	158	364	831	530	181	100	133	72	1

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Accidental poisoning and exposure to noxious substances (X40-X49)													
Non-Hispanic black, both sexes													
2005 2/	2,920	7	7	9	114	388	799	1,089	390	77	29	10	1
2004 2/	2,460	3	6	11	113	288	733	915	281	64	33	13	-
2003 2/	2,354	7	15	17	86	303	770	848	204	68	19	16	1
2002	2,326	12	11	6	86	299	786	825	221	41	26	12	1
2001	2,122	6	8	6	93	273	749	736	157	39	34	21	-
2000	1,985	4	11	8	76	294	732	645	132	41	15	26	1
1999	2,008	1	12	7	93	328	729	613	139	49	18	17	2
Non-Hispanic black, male													
2005 2/	1,959	4	6	4	77	264	488	739	297	59	18	3	-
2004 2/	1,607	2	6	3	81	195	439	606	207	45	18	5	-
2003 2/	1,579	1	9	7	64	204	497	570	154	52	10	10	1
2002	1,601	8	8	3	66	213	517	564	174	31	15	1	1
2001	1,452	3	5	2	67	186	477	540	121	25	20	6	-
2000	1,400	3	8	6	49	209	496	480	101	29	7	11	1
1999	1,450	1	7	5	76	231	503	471	114	29	9	3	1
Non-Hispanic black, female													
2005 2/	961	3	1	5	37	124	311	350	93	18	11	7	1
2004 2/	853	1	-	8	32	93	294	309	74	19	15	8	-
2003 2/	775	6	6	10	22	99	273	278	50	16	9	6	-
2002	725	4	3	3	20	86	269	261	47	10	11	11	-
2001	670	3	3	4	26	87	272	196	36	14	14	15	-
2000	585	1	3	2	27	85	236	165	31	12	8	15	-
1999	558	-	5	2	17	97	226	142	25	20	9	14	1

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Accidental poisoning and exposure to noxious substances (X40-X49)													
Origin not stated 3/, both sexes													
2005	112	-	-	-	5	15	30	44	13	4	-	1	-
2004	81	-	1	-	3	10	25	28	8	1	1	-	4
2003	118	-	-	-	4	20	30	47	10	4	1	-	2
2002	109	-	-	-	8	17	39	24	8	6	2	-	5
2001	123	-	-	-	8	16	42	34	15	2	1	-	5
2000	138	-	1	3	6	18	39	52	11	4	-	-	4
1999	142	-	-	1	4	17	57	39	14	2	2	2	4
Origin not stated 3/, male													
2005	83	-	-	-	4	11	22	34	10	2	-	-	-
2004	66	-	1	-	3	8	19	24	6	1	1	-	3
2003	93	-	-	-	2	16	24	35	9	4	1	-	2
2002	86	-	-	-	5	14	30	17	8	6	2	-	4
2001	104	-	-	-	7	14	34	28	13	2	1	-	5
2000	104	-	1	1	5	13	31	37	9	3	-	-	4
1999	106	-	-	1	4	14	39	33	10	2	-	1	2
Origin not stated 3/, female													
2005	29	-	-	-	1	4	8	10	3	2	-	1	-
2004	15	-	-	-	-	2	6	4	2	-	-	-	1
2003	25	-	-	-	2	4	6	12	1	-	-	-	-
2002	23	-	-	-	3	3	9	7	-	-	-	-	1
2001	19	-	-	-	1	2	8	6	2	-	-	-	-
2000	34	-	-	2	1	5	8	15	2	1	-	-	-
1999	36	-	-	-	-	3	18	6	4	-	2	1	2

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Other and unspecified nontransport accidents and their sequelae (W20-W31,W35-W64,W75-W99,X10-X39,X50-X59,Y86)													
All origins, both sexes													
2005	18,526	801	244	258	725	939	1,343	1,977	1,746	1,898	3,731	4,848	16
2004	17,016	782	250	247	640	825	1,276	1,732	1,453	1,634	3,505	4,641	31
2003	17,115	675	278	215	645	813	1,331	1,591	1,420	1,760	3,539	4,833	15
2002	17,201	677	247	273	638	820	1,341	1,496	1,385	1,795	3,665	4,851	13
2001	17,760	669	275	255	643	795	1,386	1,641	1,342	1,826	3,841	5,063	24
2000	17,437	572	266	267	648	895	1,446	1,510	1,349	1,824	3,678	4,966	16
1999	18,388	523	293	316	662	887	1,452	1,496	1,332	2,032	4,025	5,355	15
All origins, male													
2005	10,864	443	148	180	600	793	1,081	1,554	1,260	1,178	1,897	1,718	12
2004	9,880	453	165	184	550	694	1,009	1,347	1,044	995	1,754	1,655	30
2003	9,919	398	177	160	544	685	1,061	1,206	1,045	1,095	1,798	1,737	13
2002	9,839	389	156	185	545	692	1,081	1,180	1,013	1,113	1,827	1,648	10
2001	10,044	390	162	181	535	682	1,128	1,261	933	1,097	1,913	1,741	21
2000	9,960	327	162	192	560	767	1,205	1,219	971	1,081	1,801	1,659	16
1999	10,371	314	183	221	565	748	1,181	1,176	951	1,227	1,969	1,822	14
All origins, female													
2005	7,662	358	96	78	125	146	262	423	486	720	1,834	3,130	4
2004	7,136	329	85	63	90	131	267	385	409	639	1,751	2,986	1
2003	7,196	277	101	55	101	128	270	385	375	665	1,741	3,096	2
2002	7,362	288	91	88	93	128	260	316	372	682	1,838	3,203	3
2001	7,716	279	113	74	108	113	258	380	409	729	1,928	3,322	3
2000	7,477	245	104	75	88	128	241	291	378	743	1,877	3,307	-
1999	8,017	209	110	95	97	139	271	320	381	805	2,056	3,533	1

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Other and unspecified nontransport accidents and their sequelae (W20-W31,W35-W64,W75-W99,X10-X39,X50-X59,Y86)													
Hispanic, both sexes													
2005	1,200	103	44	28	120	189	164	164	98	90	110	89	1
2004	1,042	89	42	28	120	143	146	133	77	66	100	93	5
2003	999	71	53	29	115	150	128	114	82	77	108	72	-
2002	954	68	42	36	100	139	144	101	67	73	92	87	5
2001	985	78	43	23	98	122	146	122	70	88	89	102	4
2000	942	58	41	29	108	140	151	84	68	67	84	106	6
1999	878	65	53	31	98	101	120	82	69	90	72	94	3
Hispanic, male													
2005	892	55	27	20	109	175	147	137	73	57	54	37	1
2004	752	54	28	23	107	121	127	114	54	39	49	31	5
2003	749	40	32	21	108	138	114	90	63	49	65	29	-
2002	726	43	28	21	92	133	127	87	52	44	52	42	5
2001	704	40	28	12	93	110	130	102	49	54	46	36	4
2000	695	31	23	14	105	129	139	70	53	41	40	44	6
1999	620	35	37	19	88	90	101	70	48	60	30	39	3
Hispanic, female													
2005	308	48	17	8	11	14	17	27	25	33	56	52	-
2004	290	35	14	5	13	22	19	19	23	27	51	62	-
2003	250	31	21	8	7	12	14	24	19	28	43	43	-
2002	228	25	14	15	8	6	17	14	15	29	40	45	-
2001	281	38	15	11	5	12	16	20	21	34	43	66	-
2000	247	27	18	15	3	11	12	14	15	26	44	62	-
1999	258	30	16	12	10	11	19	12	21	30	42	55	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Other and unspecified nontransport accidents and their sequelae (W20-W31,W35-W64,W75-W99,X10-X39,X50-X59,Y86)													
Non-Hispanic 1/, both sexes													
2005	17,261	695	200	230	605	745	1,172	1,801	1,636	1,800	3,617	4,752	8
2004	15,923	693	207	218	518	678	1,129	1,592	1,371	1,560	3,398	4,544	15
2003	16,055	602	225	186	526	659	1,195	1,471	1,335	1,678	3,422	4,751	5
2002	16,183	606	205	235	534	677	1,193	1,383	1,311	1,717	3,566	4,750	6
2001	16,695	588	231	230	542	669	1,231	1,513	1,262	1,732	3,741	4,950	6
2000	16,408	513	224	238	538	748	1,283	1,413	1,275	1,748	3,580	4,846	2
1999	17,430	457	239	284	562	781	1,331	1,400	1,250	1,935	3,943	5,245	3
Non-Hispanic 1/, male													
2005	9,924	387	121	160	491	614	929	1,407	1,178	1,114	1,839	1,679	5
2004	9,092	399	136	161	441	569	881	1,228	987	948	1,703	1,624	15
2003	9,122	356	145	139	432	544	939	1,110	980	1,043	1,726	1,704	4
2002	9,070	344	128	163	450	555	950	1,084	954	1,065	1,771	1,601	5
2001	9,280	347	133	167	439	568	990	1,154	876	1,040	1,858	1,702	6
2000	9,207	295	139	178	454	631	1,057	1,138	913	1,033	1,755	1,612	2
1999	9,694	279	146	201	476	655	1,079	1,094	891	1,162	1,933	1,775	3
Non-Hispanic 1/, female													
2005	7,337	308	79	70	114	131	243	394	458	686	1,778	3,073	3
2004	6,831	294	71	57	77	109	248	364	384	612	1,695	2,920	-
2003	6,933	246	80	47	94	115	256	361	355	635	1,696	3,047	1
2002	7,113	262	77	72	84	122	243	299	357	652	1,795	3,149	1
2001	7,415	241	98	63	103	101	241	359	386	692	1,883	3,248	-
2000	7,201	218	85	60	84	117	226	275	362	715	1,825	3,234	-
1999	7,736	178	93	83	86	126	252	306	359	773	2,010	3,470	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Other and unspecified nontransport accidents and their sequelae (W20-W31,W35-W64,W75-W99,X10-X39,X50-X59,Y86)													
Non-Hispanic white, both sexes													
2005 2/	14,493	430	149	176	469	598	930	1,389	1,316	1,481	3,181	4,369	5
2004 2/	13,724	426	144	163	433	553	928	1,327	1,147	1,314	3,041	4,234	14
2003 2/	13,791	356	158	143	424	531	969	1,164	1,092	1,419	3,084	4,446	5
2002	13,865	339	136	172	446	565	969	1,131	1,081	1,414	3,190	4,416	6
2001	14,326	348	160	182	455	528	994	1,216	1,033	1,462	3,359	4,583	6
2000	14,083	310	156	185	425	604	1,039	1,123	1,044	1,464	3,224	4,509	-
1999	14,930	276	163	225	468	622	1,103	1,100	980	1,612	3,504	4,876	1
Non-Hispanic white, male													
2005 2/	8,217	238	95	125	387	500	751	1,093	935	924	1,615	1,551	3
2004 2/	7,754	260	102	116	372	465	725	1,025	831	810	1,528	1,506	14
2003 2/	7,709	211	104	108	358	444	767	886	799	891	1,549	1,588	4
2002	7,669	203	83	120	382	466	786	898	786	882	1,580	1,478	5
2001	7,832	206	93	129	370	449	814	929	719	886	1,658	1,573	6
2000	7,703	186	99	141	361	515	860	899	736	868	1,555	1,483	-
1999	8,122	167	96	163	395	532	896	860	705	962	1,717	1,628	1
Non-Hispanic white, female													
2005 2/	6,276	192	54	51	82	98	179	296	381	557	1,566	2,818	2
2004 2/	5,970	166	42	47	61	88	203	302	316	504	1,513	2,728	-
2003 2/	6,082	145	54	35	66	87	202	278	293	528	1,535	2,858	1
2002	6,196	136	53	52	64	99	183	233	295	532	1,610	2,938	1
2001	6,494	142	67	53	85	79	180	287	314	576	1,701	3,010	-
2000	6,380	124	57	44	64	89	179	224	308	596	1,669	3,026	-
1999	6,808	109	67	62	73	90	207	240	275	650	1,787	3,248	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Other and unspecified nontransport accidents and their sequelae (W20-W31,W35-W64,W75-W99,X10-X39,X50-X59,Y86)													
Non-Hispanic black, both sexes													
2005 2/	2,309	244	43	41	98	110	188	349	262	272	379	320	3
2004 2/	1,825	241	47	41	61	106	163	228	186	200	304	248	-
2003 2/	1,858	226	52	31	79	95	185	255	201	218	267	249	-
2002	1,951	250	59	55	68	90	179	214	189	257	311	279	-
2001	1,968	217	61	38	62	101	184	249	198	232	317	309	-
2000	1,968	183	55	48	90	118	198	237	200	244	307	286	2
1999	2,183	168	61	47	74	132	193	263	236	293	384	330	2
Non-Hispanic black, male													
2005 2/	1,400	137	23	26	77	79	137	264	201	156	194	104	2
2004 2/	1,101	125	22	34	47	88	126	171	131	117	152	88	-
2003 2/	1,147	134	36	22	56	74	138	186	146	130	138	87	-
2002	1,162	132	40	38	52	71	126	157	138	151	159	98	-
2001	1,180	131	35	31	47	81	137	185	134	134	165	100	-
2000	1,256	96	32	34	73	94	159	196	151	139	174	106	2
1999	1,365	101	42	32	67	102	153	203	165	182	185	131	2
Non-Hispanic black, female													
2005 2/	909	107	20	15	21	31	51	85	61	116	185	216	1
2004 2/	724	116	25	7	14	18	37	57	55	83	152	160	-
2003 2/	711	92	16	9	23	21	47	69	55	88	129	162	-
2002	789	118	19	17	16	19	53	57	51	106	152	181	-
2001	788	86	26	7	15	20	47	64	64	98	152	209	-
2000	712	87	23	14	17	24	39	41	49	105	133	180	-
1999	818	67	19	15	7	30	40	60	71	111	199	199	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Other and unspecified nontransport accidents and their sequelae (W20-W31,W35-W64,W75-W99,X10-X39,X50-X59,Y86)													
Origin not stated 3/, both sexes													
2005	65	3	-	-	-	5	7	12	12	8	4	7	7
2004	51	-	1	1	2	4	1	7	5	8	7	4	11
2003	61	2	-	-	4	4	8	6	3	5	9	10	10
2002	64	3	-	2	4	4	4	12	7	5	7	14	2
2001	80	3	1	2	3	4	9	6	10	6	11	11	14
2000	87	1	1	-	2	7	12	13	6	9	14	14	8
1999	80	1	1	1	2	5	1	14	13	7	10	16	9
Origin not stated 3/, male													
2005	48	1	-	-	-	4	5	10	9	7	4	2	6
2004	36	-	1	-	2	4	1	5	3	8	2	-	10
2003	48	2	-	-	4	3	8	6	2	3	7	4	9
2002	43	2	-	1	3	4	4	9	7	4	4	5	-
2001	60	3	1	2	3	4	8	5	8	3	9	3	11
2000	58	1	-	-	1	7	9	11	5	7	6	3	8
1999	57	-	-	1	1	3	1	12	12	5	6	8	8
Origin not stated 3/, female													
2005	17	2	-	-	-	1	2	2	3	1	-	5	1
2004	15	-	-	1	-	-	-	2	2	-	5	4	1
2003	13	-	-	-	-	1	-	-	1	2	2	6	1
2002	21	1	-	1	1	-	-	3	-	1	3	9	2
2001	20	-	-	-	-	-	1	1	2	3	2	8	3
2000	29	-	1	-	1	-	3	2	1	2	8	11	-
1999	23	1	1	-	1	2	-	2	1	2	4	8	1

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)													
All origins, both sexes													
2005	32,637	...	...	272	4,212	4,990	6,550	6,991	4,210	2,344	2,200	860	8
2004	32,439	...	...	285	4,316	5,074	6,638	6,906	4,011	2,279	2,120	799	11
2003	31,484	...	...	250	3,988	5,065	6,602	6,481	3,843	2,335	2,115	798	7
2002	31,655	...	...	264	4,010	5,046	6,851	6,308	3,618	2,463	2,259	826	10
2001 4/	30,622	...	...	279	3,971	5,070	6,635	5,942	3,317	2,432	2,192	769	15
2000	29,350	...	...	307	3,994	4,792	6,562	5,437	2,945	2,292	2,181	833	7
1999	29,199	...	...	244	3,901	5,106	6,466	5,081	2,896	2,471	2,217	801	16
All origins, male													
2005	25,907	...	...	204	3,498	4,066	5,063	5,269	3,250	1,940	1,889	721	7
2004	25,566	...	...	187	3,596	4,142	5,067	5,078	3,088	1,901	1,817	679	11
2003	25,203	...	...	194	3,381	4,156	5,138	4,887	2,987	1,954	1,809	690	7
2002	25,409	...	...	199	3,432	4,135	5,305	4,796	2,837	2,053	1,938	704	10
2001 4/	24,672	...	...	214	3,409	4,199	5,180	4,504	2,563	2,040	1,885	664	14
2000	23,618	...	...	244	3,424	3,938	5,114	4,149	2,265	1,888	1,884	705	7
1999	23,458	...	...	194	3,326	4,194	5,019	3,854	2,255	2,051	1,868	681	16
All origins, female													
2005	6,730	...	...	68	714	924	1,487	1,722	960	404	311	139	1
2004	6,873	...	...	98	720	932	1,571	1,828	923	378	303	120	-
2003	6,281	...	...	56	607	909	1,464	1,594	856	381	306	108	-
2002	6,246	...	...	65	578	911	1,546	1,512	781	410	321	122	-
2001 4/	5,950	...	...	65	562	871	1,455	1,438	754	392	307	105	1
2000	5,732	...	...	63	570	854	1,448	1,288	680	404	297	128	-
1999	5,741	...	...	50	575	912	1,447	1,227	641	420	349	120	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)													
Hispanic, both sexes													
2005	2,188	...	...	39	553	534	472	283	142	78	64	21	2
2004	2,207	...	...	39	571	545	421	304	156	87	62	19	3
2003	2,007	...	...	43	491	499	402	275	145	80	51	21	-
2002	1,954	...	...	37	455	484	407	260	142	89	55	23	2
2001 4/	1,850	...	...	37	414	490	397	239	114	86	49	22	2
2000	1,787	...	...	29	447	420	385	244	102	84	59	16	1
1999	1,695	...	...	30	391	423	369	210	112	73	63	22	2
Hispanic, male													
2005	1,841	...	...	27	462	461	399	237	116	63	56	18	2
2004	1,840	...	...	24	487	467	351	242	120	77	55	14	3
2003	1,711	...	...	31	421	434	341	220	129	71	47	17	-
2002	1,651	...	...	21	388	426	340	206	123	77	47	21	2
2001 4/	1,576	...	...	26	343	419	350	192	100	77	45	22	2
2000	1,525	...	...	22	387	370	320	194	89	74	53	15	1
1999	1,430	...	...	21	338	364	307	169	94	63	54	18	2
Hispanic, female													
2005	347	...	...	12	91	73	73	46	26	15	8	3	-
2004	367	...	...	15	84	78	70	62	36	10	7	5	-
2003	296	...	...	12	70	65	61	55	16	9	4	4	-
2002	303	...	...	16	67	58	67	54	19	12	8	2	-
2001 4/	274	...	...	11	71	71	47	47	14	9	4	-	-
2000	262	...	...	7	60	50	65	50	13	10	6	1	-
1999	265	...	...	9	53	59	62	41	18	10	9	4	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)													
Non-Hispanic 1/, both sexes													
2005	30,338	...	...	232	3,647	4,445	6,058	6,685	4,044	2,263	2,127	836	1
2004	30,111	...	...	245	3,728	4,516	6,186	6,576	3,841	2,184	2,054	779	2
2003	29,367	...	...	206	3,483	4,554	6,171	6,184	3,682	2,247	2,063	775	2
2002	29,543	...	...	226	3,536	4,538	6,413	6,015	3,455	2,360	2,199	797	4
2001 4/	28,647	...	...	242	3,545	4,560	6,213	5,682	3,184	2,334	2,138	744	5
2000	27,426	...	...	276	3,529	4,350	6,145	5,165	2,827	2,198	2,119	813	4
1999	27,379	...	...	212	3,500	4,665	6,068	4,842	2,773	2,388	2,146	776	9
Non-Hispanic 1/, male													
2005	23,974	...	...	176	3,026	3,595	4,649	5,014	3,115	1,874	1,825	700	-
2004	23,633	...	...	162	3,095	3,666	4,692	4,818	2,956	1,818	1,760	664	2
2003	23,406	...	...	162	2,950	3,712	4,775	4,651	2,844	1,878	1,761	671	2
2002	23,623	...	...	177	3,026	3,689	4,941	4,563	2,696	1,963	1,887	677	4
2001 4/	22,983	...	...	188	3,055	3,762	4,809	4,292	2,446	1,952	1,835	639	5
2000	21,981	...	...	221	3,023	3,553	4,767	3,931	2,161	1,805	1,828	688	4
1999	21,929	...	...	171	2,983	3,816	4,689	3,663	2,152	1,978	1,807	661	9
Non-Hispanic 1/, female													
2005	6,364	...	...	56	621	850	1,409	1,671	929	389	302	136	1
2004	6,478	...	...	83	633	850	1,494	1,758	885	366	294	115	-
2003	5,961	...	...	44	533	842	1,396	1,533	838	369	302	104	-
2002	5,920	...	...	49	510	849	1,472	1,452	759	397	312	120	-
2001 4/	5,664	...	...	54	490	798	1,404	1,390	738	382	303	105	-
2000	5,445	...	...	55	506	797	1,378	1,234	666	393	291	125	-
1999	5,450	...	...	41	517	849	1,379	1,179	621	410	339	115	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)													
Non-Hispanic white, both sexes													
2005 2/	27,305	...	...	170	2,992	3,708	5,434	6,208	3,822	2,137	2,037	796	1
2004 2/	27,001	...	...	180	3,045	3,747	5,598	6,101	3,584	2,043	1,954	748	1
2003 2/	26,434	...	...	151	2,832	3,811	5,563	5,774	3,462	2,110	1,980	749	2
2002	26,691	...	...	177	2,951	3,809	5,795	5,603	3,254	2,228	2,105	765	4
2001 4/	25,813	...	...	183	2,926	3,871	5,604	5,292	2,979	2,187	2,049	718	4
2000	24,616	...	...	225	2,868	3,658	5,540	4,774	2,659	2,075	2,028	786	3
1999	24,534	...	...	174	2,830	3,927	5,469	4,487	2,580	2,249	2,058	754	6
Non-Hispanic white, male													
2005 2/	21,605	...	...	133	2,487	2,995	4,164	4,656	2,953	1,785	1,756	676	-
2004 2/	21,214	...	...	119	2,535	3,064	4,249	4,465	2,754	1,701	1,682	644	1
2003 2/	21,085	...	...	122	2,412	3,099	4,296	4,357	2,678	1,769	1,696	654	2
2002	21,323	...	...	138	2,527	3,065	4,462	4,252	2,546	1,865	1,812	652	4
2001 4/	20,710	...	...	144	2,522	3,190	4,329	4,016	2,284	1,836	1,763	622	4
2000	19,711	...	...	179	2,467	2,989	4,287	3,627	2,031	1,701	1,761	666	3
1999	19,620	...	...	141	2,416	3,205	4,210	3,388	2,001	1,869	1,740	644	6
Non-Hispanic white, female													
2005 2/	5,700	...	...	37	505	713	1,270	1,552	869	352	281	120	1
2004 2/	5,787	...	...	61	510	683	1,349	1,636	830	342	272	104	-
2003 2/	5,349	...	...	29	420	712	1,267	1,417	784	341	284	95	-
2002	5,368	...	...	39	424	744	1,333	1,351	708	363	293	113	-
2001 4/	5,103	...	...	39	404	681	1,275	1,276	695	351	286	96	-
2000	4,905	...	...	46	401	669	1,253	1,147	628	374	267	120	-
1999	4,914	...	...	33	414	722	1,259	1,099	579	380	318	110	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)													
Non-Hispanic black, both sexes													
2005 2/	1,953	...	...	41	428	474	415	309	140	77	53	16	-
2004 2/	1,970	...	...	49	450	510	369	279	164	76	60	12	1
2003 2/	1,918	...	...	43	438	498	409	266	132	77	43	12	-
2002	1,896	...	...	35	392	497	432	265	125	80	51	19	-
2001 4/	1,904	...	...	40	425	472	418	268	126	90	52	13	-
2000	1,923	...	...	39	471	469	427	266	101	85	53	12	-
1999	1,924	...	...	29	463	497	430	226	116	94	52	15	2
Non-Hispanic black, male													
2005 2/	1,592	...	...	27	375	394	333	238	106	60	45	14	-
2004 2/	1,614	...	...	30	382	420	293	214	140	71	54	9	1
2003 2/	1,566	...	...	34	377	424	324	199	102	59	39	8	-
2002	1,595	...	...	29	342	441	345	212	97	66	46	17	-
2001 4/	1,578	...	...	31	388	406	329	196	102	73	44	9	-
2000	1,601	...	...	34	407	397	346	210	80	76	40	11	-
1999	1,630	...	...	22	408	432	353	183	95	78	45	12	2
Non-Hispanic black, female													
2005 2/	361	...	...	14	53	80	82	71	34	17	8	2	-
2004 2/	356	...	...	19	68	90	76	65	24	5	6	3	-
2003 2/	352	...	...	9	61	74	85	67	30	18	4	4	-
2002	301	...	...	6	50	56	87	53	28	14	5	2	-
2001 4/	326	...	...	9	37	66	89	72	24	17	8	4	-
2000	322	...	...	5	64	72	81	56	21	9	13	1	-
1999	294	...	...	7	55	65	77	43	21	16	7	3	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)													
Origin not stated 3/, both sexes													
2005	111	...	...	1	12	11	20	23	24	3	9	3	5
2004	121	...	...	1	17	13	31	26	14	8	4	1	6
2003	110	...	...	1	14	12	29	22	16	8	1	2	5
2002	158	...	...	1	19	24	31	33	21	14	5	6	4
2001 4/	125	...	...	-	12	20	25	21	19	12	5	3	8
2000	137	...	...	2	18	22	32	28	16	10	3	4	2
1999	125	...	...	2	10	18	29	29	11	10	8	3	5
Origin not stated 3/, male													
2005	92	...	...	1	10	10	15	18	19	3	8	3	5
2004	93	...	...	1	14	9	24	18	12	6	2	1	6
2003	86	...	...	1	10	10	22	16	14	5	1	2	5
2002	135	...	...	1	18	20	24	27	18	13	4	6	4
2001 4/	113	...	...	-	11	18	21	20	17	11	5	3	7
2000	112	...	...	1	14	15	27	24	15	9	3	2	2
1999	99	...	...	2	5	14	23	22	9	10	7	2	5
Origin not stated 3/, female													
2005	19	...	...	-	2	1	5	5	5	-	1	-	-
2004	28	...	...	-	3	4	7	8	2	2	2	-	-
2003	24	...	...	-	4	2	7	6	2	3	-	-	-
2002	23	...	...	-	1	4	7	6	3	1	1	-	-
2001 4/	12	...	...	-	1	2	4	1	2	1	-	-	1
2000	25	...	...	1	4	7	5	4	1	1	-	2	-
1999	26	...	...	-	5	4	6	7	2	-	1	1	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Intentional self-harm (suicide) by discharge of firearms (X72-X74)													
All origins, both sexes													
2005	17,002	...	...	84	1,962	2,269	2,855	3,472	2,470	1,669	1,649	571	1
2004	16,750	...	...	59	2,104	2,283	2,868	3,349	2,328	1,631	1,588	537	3
2003	16,907	...	...	74	2,075	2,381	2,927	3,279	2,317	1,700	1,595	559	-
2002	17,108	...	...	86	2,088	2,399	3,157	3,136	2,234	1,776	1,680	550	2
2001	16,869	...	...	90	2,130	2,564	3,030	3,023	2,083	1,758	1,648	537	6
2000	16,586	...	...	110	2,267	2,453	3,191	2,858	1,833	1,700	1,611	558	5
1999	16,599	...	...	103	2,315	2,621	3,097	2,677	1,860	1,791	1,606	524	5
All origins, male													
2005	14,916	...	...	72	1,771	1,981	2,416	2,945	2,140	1,501	1,548	542	-
2004	14,523	...	...	50	1,889	1,978	2,376	2,761	1,994	1,475	1,479	518	3
2003	14,827	...	...	63	1,879	2,099	2,470	2,768	2,001	1,538	1,480	529	-
2002	15,045	...	...	68	1,897	2,078	2,672	2,677	1,930	1,620	1,576	525	2
2001	14,758	...	...	69	1,935	2,246	2,551	2,551	1,771	1,592	1,531	506	6
2000	14,454	...	...	90	2,043	2,123	2,657	2,409	1,568	1,528	1,505	526	5
1999	14,479	...	...	80	2,083	2,270	2,586	2,251	1,594	1,615	1,499	496	5
All origins, female													
2005	2,086	...	...	12	191	288	439	527	330	168	101	29	1
2004	2,227	...	...	9	215	305	492	588	334	156	109	19	-
2003	2,080	...	...	11	196	282	457	511	316	162	115	30	-
2002	2,063	...	...	18	191	321	485	459	304	156	104	25	-
2001	2,111	...	...	21	195	318	479	472	312	166	117	31	-
2000	2,132	...	...	20	224	330	534	449	265	172	106	32	-
1999	2,120	...	...	23	232	351	511	426	266	176	107	28	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Intentional self-harm (suicide) by discharge of firearms (X72-X74)													
Hispanic, both sexes													
2005	824	...	...	6	224	203	165	100	48	33	33	12	-
2004	888	...	...	2	266	206	147	122	60	45	34	6	-
2003	835	...	...	9	228	208	145	113	57	38	26	11	-
2002	834	...	...	10	224	198	145	98	67	52	28	12	-
2001	798	...	...	5	193	224	135	102	59	45	26	8	1
2000	813	...	...	7	224	181	144	124	47	49	27	9	1
1999	794	...	...	13	219	182	148	96	48	37	39	11	1
Hispanic, male													
2005	756	...	...	5	206	183	148	92	47	31	33	11	-
2004	799	...	...	2	246	181	132	108	53	40	32	5	-
2003	774	...	...	7	210	194	133	102	55	38	25	10	-
2002	763	...	...	5	208	183	128	88	62	49	28	12	-
2001	731	...	...	4	172	203	126	90	57	44	26	8	1
2000	737	...	...	7	205	165	126	106	46	46	26	9	1
1999	705	...	...	7	196	164	127	81	45	35	38	11	1
Hispanic, female													
2005	68	...	...	1	18	20	17	8	1	2	-	1	-
2004	89	...	...	-	20	25	15	14	7	5	2	1	-
2003	61	...	...	2	18	14	12	11	2	-	1	1	-
2002	71	...	...	5	16	15	17	10	5	3	-	-	-
2001	67	...	...	1	21	21	9	12	2	1	-	-	-
2000	76	...	...	-	19	16	18	18	1	3	1	-	-
1999	89	...	...	6	23	18	21	15	3	2	1	-	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Intentional self-harm (suicide) by discharge of firearms (X72-X74)													
Non-Hispanic 1/, both sexes													
2005	16,124	...	...	78	1,732	2,061	2,680	3,363	2,409	1,634	1,610	556	1
2004	15,823	...	...	57	1,830	2,075	2,711	3,223	2,261	1,583	1,552	530	1
2003	16,035	...	...	65	1,843	2,171	2,772	3,158	2,254	1,657	1,569	546	-
2002	16,213	...	...	76	1,855	2,195	3,002	3,024	2,159	1,716	1,649	536	1
2001	16,020	...	...	85	1,933	2,335	2,887	2,916	2,011	1,706	1,617	526	4
2000	15,717	...	...	103	2,035	2,264	3,036	2,722	1,779	1,645	1,583	547	3
1999	15,755	...	...	90	2,092	2,431	2,937	2,573	1,806	1,749	1,563	511	3
Non-Hispanic 1/, male													
2005	14,110	...	...	67	1,560	1,794	2,260	2,844	2,080	1,468	1,509	528	-
2004	13,691	...	...	48	1,636	1,795	2,235	2,651	1,936	1,432	1,445	512	1
2003	14,025	...	...	56	1,667	1,903	2,330	2,660	1,940	1,497	1,455	517	-
2002	14,226	...	...	63	1,680	1,890	2,536	2,576	1,861	1,563	1,545	511	1
2001	13,978	...	...	65	1,759	2,039	2,417	2,457	1,701	1,541	1,500	495	4
2000	13,664	...	...	83	1,831	1,951	2,520	2,291	1,515	1,476	1,478	516	3
1999	13,732	...	...	73	1,884	2,101	2,448	2,164	1,543	1,575	1,457	484	3
Non-Hispanic 1/, female													
2005	2,014	...	...	11	172	267	420	519	329	166	101	28	1
2004	2,132	...	...	9	194	280	476	572	325	151	107	18	-
2003	2,010	...	...	9	176	268	442	498	314	160	114	29	-
2002	1,987	...	...	13	175	305	466	448	298	153	104	25	-
2001	2,042	...	...	20	174	296	470	459	310	165	117	31	-
2000	2,053	...	...	20	204	313	516	431	264	169	105	31	-
1999	2,023	...	...	17	208	330	489	409	263	174	106	27	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Intentional self-harm (suicide) by discharge of firearms (X72-X74)													
Non-Hispanic white, both sexes													
2005 2/	14,829	...	...	69	1,414	1,741	2,440	3,171	2,315	1,568	1,563	547	1
2004 2/	14,507	...	...	47	1,509	1,733	2,484	3,053	2,154	1,507	1,498	521	1
2003 2/	14,737	...	...	56	1,503	1,813	2,564	2,996	2,158	1,589	1,520	538	-
2002	14,865	...	...	63	1,521	1,853	2,749	2,846	2,063	1,653	1,595	521	1
2001	14,648	...	...	69	1,565	2,001	2,629	2,760	1,919	1,628	1,560	514	3
2000	14,333	...	...	96	1,655	1,916	2,780	2,552	1,679	1,572	1,545	536	2
1999	14,316	...	...	81	1,684	2,071	2,676	2,413	1,703	1,672	1,514	499	3
Non-Hispanic white, male													
2005 2/	12,947	...	...	60	1,270	1,504	2,043	2,674	1,999	1,414	1,464	519	-
2004 2/	12,534	...	...	39	1,349	1,491	2,042	2,511	1,844	1,360	1,393	504	1
2003 2/	12,865	...	...	49	1,361	1,573	2,145	2,521	1,857	1,438	1,411	510	-
2002	13,014	...	...	53	1,375	1,573	2,319	2,423	1,774	1,507	1,493	496	1
2001	12,771	...	...	55	1,414	1,739	2,199	2,333	1,619	1,474	1,451	484	3
2000	12,448	...	...	79	1,489	1,648	2,296	2,151	1,427	1,405	1,446	505	2
1999	12,441	...	...	65	1,514	1,776	2,213	2,029	1,451	1,505	1,412	473	3
Non-Hispanic white, female													
2005 2/	1,882	...	...	9	144	237	397	497	316	154	99	28	1
2004 2/	1,973	...	...	8	160	242	442	542	310	147	105	17	-
2003 2/	1,872	...	...	7	142	240	419	475	301	151	109	28	-
2002	1,851	...	...	10	146	280	430	423	289	146	102	25	-
2001	1,877	...	...	14	151	262	430	427	300	154	109	30	-
2000	1,885	...	...	17	166	268	484	401	252	167	99	31	-
1999	1,875	...	...	16	170	295	463	384	252	167	102	26	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Intentional self-harm (suicide) by discharge of firearms (X72-X74)													
Non-Hispanic black, both sexes													
2005 2/	997	...	...	6	244	247	187	137	72	56	39	9	-
2004 2/	995	...	...	8	251	264	154	114	87	61	48	8	-
2003 2/	993	...	...	7	264	279	150	117	75	56	38	7	-
2002	1,041	...	...	9	250	272	199	130	71	54	41	15	-
2001	1,069	...	...	12	293	258	198	122	67	62	47	10	-
2000	1,073	...	...	4	304	268	193	131	71	62	32	8	-
1999	1,112	...	...	5	323	268	197	118	81	65	43	12	-
Non-Hispanic black, male													
2005 2/	901	...	...	5	227	224	170	122	61	46	37	9	-
2004 2/	891	...	...	7	223	238	137	98	75	60	46	7	-
2003 2/	892	...	...	5	243	260	131	103	63	47	34	6	-
2002	945	...	...	8	229	258	170	115	63	48	39	15	-
2001	949	...	...	9	277	233	169	98	61	53	40	9	-
2000	949	...	...	4	277	237	168	107	61	60	27	8	-
1999	1,007	...	...	4	294	247	178	102	73	59	39	11	-
Non-Hispanic black, female													
2005 2/	96	...	...	1	17	23	17	15	11	10	2	-	-
2004 2/	104	...	...	1	28	26	17	16	12	1	2	1	-
2003 2/	101	...	...	2	21	19	19	14	12	9	4	1	-
2002	96	...	...	1	21	14	29	15	8	6	2	-	-
2001	120	...	...	3	16	25	29	24	6	9	7	1	-
2000	124	...	...	-	27	31	25	24	10	2	5	-	-
1999	105	...	...	1	29	21	19	16	8	6	4	1	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Intentional self-harm (suicide) by discharge of firearms (X72-X74)													
Origin not stated 3/, both sexes													
2005	54	...	...	-	6	5	10	9	13	2	6	3	-
2004	39	...	...	-	8	2	10	4	7	3	2	1	2
2003	37	...	...	-	4	2	10	8	6	5	-	2	-
2002	61	...	...	-	9	6	10	14	8	8	3	2	1
2001	51	...	...	-	4	5	8	5	13	7	5	3	1
2000	56	...	...	-	8	8	11	12	7	6	1	2	1
1999	50	...	...	-	4	8	12	8	6	5	4	2	1
Origin not stated 3/, male													
2005	50	...	...	-	5	4	8	9	13	2	6	3	-
2004	33	...	...	-	7	2	9	2	5	3	2	1	2
2003	28	...	...	-	2	2	7	6	6	3	-	2	-
2002	56	...	...	-	9	5	8	13	7	8	3	2	1
2001	49	...	...	-	4	4	8	4	13	7	5	3	1
2000	53	...	...	-	7	7	11	12	7	6	1	1	1
1999	42	...	...	-	3	5	11	6	6	5	4	1	1
Origin not stated 3/, female													
2005	4	...	...	-	1	1	2	-	-	-	-	-	-
2004	6	...	...	-	1	-	1	2	2	-	-	-	-
2003	9	...	...	-	2	-	3	2	-	2	-	-	-
2002	5	...	...	-	-	1	2	1	1	-	-	-	-
2001	2	...	...	-	-	1	-	1	-	-	-	-	-
2000	3	...	...	-	1	1	-	-	-	-	-	1	-
1999	8	...	...	-	1	3	1	2	-	-	-	1	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Intentional self-harm (suicide) by other and unspecified means and their sequelae (*U03,X60-X71,X75-X84,Y87.0)													
All origins, both sexes													
2005	15,635	...	...	188	2,250	2,721	3,695	3,519	1,740	675	551	289	7
2004	15,689	...	...	226	2,212	2,791	3,770	3,557	1,683	648	532	262	8
2003	14,577	...	...	176	1,913	2,684	3,675	3,202	1,526	635	520	239	7
2002	14,547	...	...	178	1,922	2,647	3,694	3,172	1,384	687	579	276	8
2001 4/	13,753	...	...	189	1,841	2,506	3,605	2,919	1,234	674	544	232	9
2000	12,764	...	...	197	1,727	2,339	3,371	2,579	1,112	592	570	275	2
1999	12,600	...	...	141	1,586	2,485	3,369	2,404	1,036	680	611	277	11
All origins, male													
2005	10,991	...	...	132	1,727	2,085	2,647	2,324	1,110	439	341	179	7
2004	11,043	...	...	137	1,707	2,164	2,691	2,317	1,094	426	338	161	8
2003	10,376	...	...	131	1,502	2,057	2,668	2,119	986	416	329	161	7
2002	10,364	...	...	131	1,535	2,057	2,633	2,119	907	433	362	179	8
2001 4/	9,914	...	...	145	1,474	1,953	2,629	1,953	792	448	354	158	8
2000	9,164	...	...	154	1,381	1,815	2,457	1,740	697	360	379	179	2
1999	8,979	...	...	114	1,243	1,924	2,433	1,603	661	436	369	185	11
All origins, female													
2005	4,644	...	...	56	523	636	1,048	1,195	630	236	210	110	-
2004	4,646	...	...	89	505	627	1,079	1,240	589	222	194	101	-
2003	4,201	...	...	45	411	627	1,007	1,083	540	219	191	78	-
2002	4,183	...	...	47	387	590	1,061	1,053	477	254	217	97	-
2001 4/	3,839	...	...	44	367	553	976	966	442	226	190	74	1
2000	3,600	...	...	43	346	524	914	839	415	232	191	96	-
1999	3,621	...	...	27	343	561	936	801	375	244	242	92	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Intentional self-harm (suicide) by other and unspecified means and their sequelae (*U03,X60-X71,X75-X84,Y87.0)													
Hispanic, both sexes													
2005	1,364	...	...	33	329	331	307	183	94	45	31	9	2
2004	1,319	...	...	37	305	339	274	182	96	42	28	13	3
2003	1,172	...	...	34	263	291	257	162	88	42	25	10	-
2002	1,120	...	...	27	231	286	262	162	75	37	27	11	2
2001 4/	1,052	...	...	32	221	266	262	137	55	41	23	14	1
2000	974	...	...	22	223	239	241	120	55	35	32	7	-
1999	901	...	...	17	172	241	221	114	64	36	24	11	1
Hispanic, male													
2005	1,085	...	...	22	256	278	251	145	69	32	23	7	2
2004	1,041	...	...	22	241	286	219	134	67	37	23	9	3
2003	937	...	...	24	211	240	208	118	74	33	22	7	-
2002	888	...	...	16	180	243	212	118	61	28	19	9	2
2001 4/	845	...	...	22	171	216	224	102	43	33	19	14	1
2000	788	...	...	15	182	205	194	88	43	28	27	6	-
1999	725	...	...	14	142	200	180	88	49	28	16	7	1
Hispanic, female													
2005	279	...	...	11	73	53	56	38	25	13	8	2	-
2004	278	...	...	15	64	53	55	48	29	5	5	4	-
2003	235	...	...	10	52	51	49	44	14	9	3	3	-
2002	232	...	...	11	51	43	50	44	14	9	8	2	-
2001 4/	207	...	...	10	50	50	38	35	12	8	4	-	-
2000	186	...	...	7	41	34	47	32	12	7	5	1	-
1999	176	...	...	3	30	41	41	26	15	8	8	4	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Intentional self-harm (suicide) by other and unspecified means and their sequelae (*U03,X60-X71,X75-X84,Y87.0)													
Non-Hispanic 1/, both sexes													
2005	14,214	...	...	154	1,915	2,384	3,378	3,322	1,635	629	517	280	-
2004	14,288	...	...	188	1,898	2,441	3,475	3,353	1,580	601	502	249	1
2003	13,332	...	...	141	1,640	2,383	3,399	3,026	1,428	590	494	229	2
2002	13,330	...	...	150	1,681	2,343	3,411	2,991	1,296	644	550	261	3
2001 4/	12,627	...	...	157	1,612	2,225	3,326	2,766	1,173	628	521	218	1
2000	11,709	...	...	173	1,494	2,086	3,109	2,443	1,048	553	536	266	1
1999	11,624	...	...	122	1,408	2,234	3,131	2,269	967	639	583	265	6
Non-Hispanic 1/, male													
2005	9,864	...	...	109	1,466	1,801	2,389	2,170	1,035	406	316	172	-
2004	9,942	...	...	114	1,459	1,871	2,457	2,167	1,020	386	315	152	1
2003	9,381	...	...	106	1,283	1,809	2,445	1,991	904	381	306	154	2
2002	9,397	...	...	114	1,346	1,799	2,405	1,987	835	400	342	166	3
2001 4/	9,005	...	...	123	1,296	1,723	2,392	1,835	745	411	335	144	1
2000	8,317	...	...	138	1,192	1,602	2,247	1,640	646	329	350	172	1
1999	8,197	...	...	98	1,099	1,715	2,241	1,499	609	403	350	177	6
Non-Hispanic 1/, female													
2005	4,350	...	...	45	449	583	989	1,152	600	223	201	108	-
2004	4,346	...	...	74	439	570	1,018	1,186	560	215	187	97	-
2003	3,951	...	...	35	357	574	954	1,035	524	209	188	75	-
2002	3,933	...	...	36	335	544	1,006	1,004	461	244	208	95	-
2001 4/	3,622	...	...	34	316	502	934	931	428	217	186	74	-
2000	3,392	...	...	35	302	484	862	803	402	224	186	94	-
1999	3,427	...	...	24	309	519	890	770	358	236	233	88	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Intentional self-harm (suicide) by other and unspecified means and their sequelae (*U03,X60-X71,X75-X84,Y87.0)													
Non-Hispanic white, both sexes													
2005 2/	12,476	...	...	101	1,578	1,967	2,994	3,037	1,507	569	474	249	-
2004 2/	12,494	...	...	133	1,536	2,014	3,114	3,048	1,430	536	456	227	-
2003 2/	11,697	...	...	95	1,329	1,998	2,999	2,778	1,304	521	460	211	2
2002	11,826	...	...	114	1,430	1,956	3,046	2,757	1,191	575	510	244	3
2001 4/	11,165	...	...	114	1,361	1,870	2,975	2,532	1,060	559	489	204	1
2000	10,283	...	...	129	1,213	1,742	2,760	2,222	980	503	483	250	1
1999	10,218	...	...	93	1,146	1,856	2,793	2,074	877	577	544	255	3
Non-Hispanic white, male													
2005 2/	8,658	...	...	73	1,217	1,491	2,121	1,982	954	371	292	157	-
2004 2/	8,680	...	...	80	1,186	1,573	2,207	1,954	910	341	289	140	-
2003 2/	8,220	...	...	73	1,051	1,526	2,151	1,836	821	331	285	144	2
2002	8,309	...	...	85	1,152	1,492	2,143	1,829	772	358	319	156	3
2001 4/	7,939	...	...	89	1,108	1,451	2,130	1,683	665	362	312	138	1
2000	7,263	...	...	100	978	1,341	1,991	1,476	604	296	315	161	1
1999	7,179	...	...	76	902	1,429	1,997	1,359	550	364	328	171	3
Non-Hispanic white, female													
2005 2/	3,818	...	...	28	361	476	873	1,055	553	198	182	92	-
2004 2/	3,814	...	...	53	350	441	907	1,094	520	195	167	87	-
2003 2/	3,477	...	...	22	278	472	848	942	483	190	175	67	-
2002	3,517	...	...	29	278	464	903	928	419	217	191	88	-
2001 4/	3,226	...	...	25	253	419	845	849	395	197	177	66	-
2000	3,020	...	...	29	235	401	769	746	376	207	168	89	-
1999	3,039	...	...	17	244	427	796	715	327	213	216	84	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Intentional self-harm (suicide) by other and unspecified means and their sequelae (*U03,X60-X71,X75-X84,Y87.0)													
Non-Hispanic black, both sexes													
2005 2/	956	...	...	35	184	227	228	172	68	21	14	7	-
2004 2/	975	...	...	41	199	246	215	165	77	15	12	4	1
2003 2/	925	...	...	36	174	219	259	149	57	21	5	5	-
2002	855	...	...	26	142	225	233	135	54	26	10	4	-
2001 4/	835	...	...	28	132	214	220	146	59	28	5	3	-
2000	850	...	...	35	167	201	234	135	30	23	21	4	-
1999	812	...	...	24	140	229	233	108	35	29	9	3	2
Non-Hispanic black, male													
2005 2/	691	...	...	22	148	170	163	116	45	14	8	5	-
2004 2/	723	...	...	23	159	182	156	116	65	11	8	2	1
2003 2/	674	...	...	29	134	164	193	96	39	12	5	2	-
2002	650	...	...	21	113	183	175	97	34	18	7	2	-
2001 4/	629	...	...	22	111	173	160	98	41	20	4	-	-
2000	652	...	...	30	130	160	178	103	19	16	13	3	-
1999	623	...	...	18	114	185	175	81	22	19	6	1	2
Non-Hispanic black, female													
2005 2/	265	...	...	13	36	57	65	56	23	7	6	2	-
2004 2/	252	...	...	18	40	64	59	49	12	4	4	2	-
2003 2/	251	...	...	7	40	55	66	53	18	9	-	3	-
2002	205	...	...	5	29	42	58	38	20	8	3	2	-
2001 4/	206	...	...	6	21	41	60	48	18	8	1	3	-
2000	198	...	...	5	37	41	56	32	11	7	8	1	-
1999	189	...	...	6	26	44	58	27	13	10	3	2	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Intentional self-harm (suicide) by other and unspecified means and their sequelae (*U03,X60-X71,X75-X84,Y87.0)													
Origin not stated 3/, both sexes													
2005	57	...	...	1	6	6	10	14	11	1	3	-	5
2004	82	...	...	1	9	11	21	22	7	5	2	-	4
2003	73	...	...	1	10	10	19	14	10	3	1	-	5
2002	97	...	...	1	10	18	21	19	13	6	2	4	3
2001 4/	74	...	...	-	8	15	17	16	6	5	-	-	7
2000	81	...	...	2	10	14	21	16	9	4	2	2	1
1999	75	...	...	2	6	10	17	21	5	5	4	1	4
Origin not stated 3/, male													
2005	42	...	...	1	5	6	7	9	6	1	2	-	5
2004	60	...	...	1	7	7	15	16	7	3	-	-	4
2003	58	...	...	1	8	8	15	10	8	2	1	-	5
2002	79	...	...	1	9	15	16	14	11	5	1	4	3
2001 4/	64	...	...	-	7	14	13	16	4	4	-	-	6
2000	59	...	...	1	7	8	16	12	8	3	2	1	1
1999	57	...	...	2	2	9	12	16	3	5	3	1	4
Origin not stated 3/, female													
2005	15	...	...	-	1	-	3	5	5	-	1	-	-
2004	22	...	...	-	2	4	6	6	-	2	2	-	-
2003	15	...	...	-	2	2	4	4	2	1	-	-	-
2002	18	...	...	-	1	3	5	5	2	1	1	-	-
2001 4/	10	...	...	-	1	1	4	-	2	1	-	-	1
2000	22	...	...	1	3	6	5	4	1	1	-	1	-
1999	18	...	...	-	4	1	5	5	2	-	1	-	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)													
All origins, both sexes													
2005	18,124	306	375	341	5,466	4,752	3,109	2,060	862	440	283	109	21
2004	17,357	325	377	329	5,085	4,495	2,984	2,008	879	446	289	104	36
2003	17,732	341	376	324	5,368	4,516	3,110	2,017	786	432	317	104	41
2002	17,638	303	423	356	5,219	4,489	3,239	1,915	841	421	296	95	41
2001 4/	20,308	332	415	326	5,297	5,204	4,268	2,467	1,018	532	312	105	32
2000	16,765	349	356	371	4,939	4,164	3,219	1,755	738	443	301	100	30
1999	16,889	331	376	432	4,998	4,231	3,206	1,668	722	473	304	99	49
All origins, male													
2005	14,376	173	214	204	4,765	4,008	2,317	1,579	636	282	138	44	16
2004	13,578	167	204	200	4,380	3,764	2,224	1,492	637	302	138	41	29
2003	13,882	205	202	215	4,624	3,741	2,308	1,488	568	285	171	46	29
2002	13,640	164	234	197	4,474	3,669	2,386	1,394	614	277	158	41	32
2001 4/	15,555	197	238	193	4,541	4,157	3,122	1,832	723	332	155	42	23
2000	12,820	203	198	227	4,203	3,361	2,319	1,284	537	279	154	40	15
1999	12,785	187	194	250	4,191	3,346	2,307	1,229	541	304	162	41	33
All origins, female													
2005	3,748	133	161	137	701	744	792	481	226	158	145	65	5
2004	3,779	158	173	129	705	731	760	516	242	144	151	63	7
2003	3,850	136	174	109	744	775	802	529	218	147	146	58	12
2002	3,998	139	189	159	745	820	853	521	227	144	138	54	9
2001 4/	4,753	135	177	133	756	1,047	1,146	635	295	200	157	63	9
2000	3,945	146	158	144	736	803	900	471	201	164	147	60	15
1999	4,104	144	182	182	807	885	899	439	181	169	142	58	16

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)													
Hispanic, both sexes													
2005	3,520	60	82	62	1,307	1,085	520	264	71	36	23	3	7
2004	3,271	73	80	67	1,235	962	477	238	84	29	9	11	6
2003	3,355	60	82	53	1,286	971	502	246	75	46	23	5	6
2002	3,129	52	81	64	1,200	903	464	216	87	29	19	3	11
2001 4/	3,331	55	69	56	1,222	960	567	248	82	45	16	3	8
2000	2,917	54	57	76	1,126	810	447	211	77	36	16	4	3
1999	2,864	53	52	89	1,119	801	432	197	69	33	11	3	5
Hispanic, male													
2005	3,008	30	46	40	1,187	956	428	218	53	26	16	1	7
2004	2,782	42	45	40	1,125	844	384	193	69	23	5	6	6
2003	2,809	35	36	36	1,139	852	401	194	59	37	13	2	5
2002	2,635	28	49	37	1,081	789	367	166	72	25	9	2	10
2001 4/	2,756	33	38	38	1,099	800	451	190	59	33	8	2	5
2000	2,431	26	30	56	1,014	697	359	158	57	22	8	2	2
1999	2,399	26	25	59	990	698	354	150	56	26	8	2	5
Hispanic, female													
2005	512	30	36	22	120	129	92	46	18	10	7	2	-
2004	489	31	35	27	110	118	93	45	15	6	4	5	-
2003	546	25	46	17	147	119	101	52	16	9	10	3	1
2002	494	24	32	27	119	114	97	50	15	4	10	1	1
2001 4/	575	22	31	18	123	160	116	58	23	12	8	1	3
2000	486	28	27	20	112	113	88	53	20	14	8	2	1
1999	465	27	27	30	129	103	78	47	13	7	3	1	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)													
Non-Hispanic 1/, both sexes													
2005	14,504	245	292	278	4,132	3,646	2,567	1,785	786	404	260	105	4
2004	13,965	247	297	261	3,830	3,506	2,488	1,745	793	415	279	93	11
2003	14,245	278	293	271	4,045	3,520	2,588	1,757	705	381	291	97	19
2002	14,346	246	340	289	3,983	3,548	2,746	1,682	747	388	273	92	12
2001 4/	16,726	270	343	269	4,028	4,178	3,652	2,182	916	480	296	101	11
2000	13,717	291	299	293	3,776	3,337	2,749	1,526	652	402	285	95	12
1999	13,873	270	323	341	3,860	3,402	2,743	1,452	642	437	292	95	16
Non-Hispanic 1/, male													
2005	11,288	143	168	164	3,555	3,033	1,870	1,352	578	256	122	43	4
2004	10,699	121	159	159	3,240	2,898	1,824	1,278	566	277	133	35	9
2003	10,969	168	166	179	3,455	2,868	1,894	1,282	503	243	156	42	13
2002	10,881	131	184	157	3,362	2,852	2,002	1,213	536	248	148	39	9
2001 4/	12,608	161	199	154	3,401	3,309	2,632	1,612	651	294	147	39	9
2000	10,290	174	168	170	3,159	2,650	1,939	1,115	473	254	146	37	5
1999	10,275	158	168	191	3,188	2,624	1,930	1,063	476	275	153	39	10
Non-Hispanic 1/, female													
2005	3,216	102	124	114	577	613	697	433	208	148	138	62	-
2004	3,266	126	138	102	590	608	664	467	227	138	146	58	2
2003	3,276	110	127	92	590	652	694	475	202	138	135	55	6
2002	3,465	115	156	132	621	696	744	469	211	140	125	53	3
2001 4/	4,118	109	144	115	627	869	1,020	570	265	186	149	62	2
2000	3,427	117	131	123	617	687	810	411	179	148	139	58	7
1999	3,598	112	155	150	672	778	813	389	166	162	139	56	6

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)													
Non-Hispanic white, both sexes													
2005 2/	5,325	139	138	111	895	973	1,099	920	502	260	200	84	4
2004 2/	5,427	137	131	119	917	976	1,124	933	504	292	210	75	9
2003 2/	5,367	156	123	109	907	953	1,147	947	455	254	220	80	16
2002	5,571	117	169	133	958	977	1,268	898	496	275	197	73	10
2001 4/	7,928	139	160	164	1,058	1,642	2,081	1,373	652	342	225	85	7
2000	5,429	141	144	164	874	1,023	1,273	834	415	264	211	78	8
1999	5,748	143	156	176	962	1,135	1,327	814	418	319	214	76	8
Non-Hispanic white, male													
2005 2/	3,515	82	74	61	639	664	710	664	355	151	84	27	4
2004 2/	3,561	72	73	70	624	672	751	649	343	180	95	25	7
2003 2/	3,550	94	61	72	665	648	747	645	303	153	115	36	11
2002	3,661	70	89	61	685	660	874	586	332	166	99	32	7
2001 4/	5,453	84	95	83	729	1,177	1,498	990	454	197	110	31	5
2000	3,510	96	86	93	588	677	836	560	288	153	101	28	4
1999	3,747	94	83	93	644	753	881	579	293	188	109	27	3
Non-Hispanic white, female													
2005 2/	1,810	57	64	50	256	309	389	256	147	109	116	57	-
2004 2/	1,866	65	58	49	293	304	373	284	161	112	115	50	2
2003 2/	1,817	62	62	37	242	305	400	302	152	101	105	44	5
2002	1,910	47	80	72	273	317	394	312	164	109	98	41	3
2001 4/	2,475	55	65	81	329	465	583	383	198	145	115	54	2
2000	1,919	45	58	71	286	346	437	274	127	111	110	50	4
1999	2,001	49	73	83	318	382	446	235	125	131	105	49	5

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)													
Non-Hispanic black, both sexes													
2005 2/	8,522	91	138	145	3,030	2,538	1,351	785	244	124	58	18	-
2004 2/	8,000	100	154	130	2,771	2,405	1,276	732	247	104	62	17	2
2003 2/	8,276	106	155	147	2,955	2,443	1,335	731	218	105	63	16	2
2002	8,147	114	156	132	2,854	2,425	1,357	710	216	95	69	17	2
2001 4/	8,077	119	166	92	2,817	2,335	1,427	704	219	117	63	14	4
2000	7,748	139	143	122	2,771	2,193	1,359	615	201	123	64	14	4
1999	7,536	114	151	145	2,752	2,122	1,305	557	194	103	67	18	8
Non-Hispanic black, male													
2005 2/	7,288	51	88	93	2,750	2,270	1,075	624	193	93	37	14	-
2004 2/	6,730	45	79	81	2,495	2,130	1,005	574	190	83	36	10	2
2003 2/	6,980	60	96	98	2,642	2,128	1,078	576	178	78	39	6	1
2002	6,780	54	89	84	2,540	2,092	1,042	572	180	72	47	6	2
2001 4/	6,658	68	100	66	2,551	1,998	1,036	544	165	85	34	7	4
2000	6,382	73	73	70	2,459	1,883	1,027	498	159	93	39	7	1
1999	6,131	58	74	89	2,433	1,771	980	431	159	79	39	11	7
Non-Hispanic black, female													
2005 2/	1,234	40	50	52	280	268	276	161	51	31	21	4	-
2004 2/	1,270	55	75	49	276	275	271	158	57	21	26	7	-
2003 2/	1,296	46	59	49	313	315	257	155	40	27	24	10	1
2002	1,367	60	67	48	314	333	315	138	36	23	22	11	-
2001 4/	1,419	51	66	26	266	337	391	160	54	32	29	7	-
2000	1,366	66	70	52	312	310	332	117	42	30	25	7	3
1999	1,405	56	77	56	319	351	325	126	35	24	28	7	1

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)													
Origin not stated 3/, both sexes													
2005	100	1	1	1	27	21	22	11	5	-	-	1	10
2004	121	5	-	1	20	27	19	25	2	2	1	-	19
2003	132	3	1	-	37	25	20	14	6	5	3	2	16
2002	163	5	2	3	36	38	29	17	7	4	4	-	18
2001 4/	251	7	3	1	47	66	49	37	20	7	-	1	13
2000	131	4	-	2	37	17	23	18	9	5	-	1	15
1999	152	8	1	2	19	28	31	19	11	3	1	1	28
Origin not stated 3/, male													
2005	80	-	-	-	23	19	19	9	5	-	-	-	5
2004	97	4	-	1	15	22	16	21	2	2	-	-	14
2003	104	2	-	-	30	21	13	12	6	5	2	2	11
2002	124	5	1	3	31	28	17	15	6	4	1	-	13
2001 4/	191	3	1	1	41	48	39	30	13	5	-	1	9
2000	99	3	-	1	30	14	21	11	7	3	-	1	8
1999	111	3	1	-	13	24	23	16	9	3	1	-	18
Origin not stated 3/, female													
2005	20	1	1	1	4	2	3	2	-	-	-	1	5
2004	24	1	-	-	5	5	3	4	-	-	1	-	5
2003	28	1	1	-	7	4	7	2	-	-	1	-	5
2002	39	-	1	-	5	10	12	2	1	-	3	-	5
2001 4/	60	4	2	-	6	18	10	7	7	2	-	-	4
2000	32	1	-	1	7	3	2	7	2	2	-	-	7
1999	41	5	-	2	6	4	8	3	2	-	-	1	10

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Assault (homicide) by discharge of firearms (*U01.4,X93-X95)													
All origins, both sexes													
2005	12,352	6	37	187	4,499	3,780	2,010	1,097	405	201	97	24	9
2004	11,624	6	36	184	4,127	3,503	1,895	1,062	468	193	100	29	21
2003	11,920	8	40	187	4,410	3,540	1,941	1,110	394	161	88	23	18
2002	11,829	9	49	205	4,317	3,465	2,042	1,025	409	161	115	19	13
2001	11,348	11	55	180	4,200	3,308	1,978	934	364	192	99	16	11
2000	10,801	12	28	187	3,963	3,074	1,906	959	367	184	81	28	12
1999	10,828	8	50	224	4,038	3,048	1,868	893	373	193	95	23	15
All origins, male													
2005	10,561	6	22	131	4,096	3,344	1,599	859	312	131	44	10	7
2004	9,921	2	18	134	3,718	3,111	1,556	821	347	138	41	15	20
2003	10,126	7	20	134	3,977	3,125	1,535	860	292	115	35	10	16
2002	9,899	3	30	127	3,851	3,017	1,623	761	301	101	64	9	12
2001	9,532	7	27	125	3,796	2,864	1,528	726	264	123	54	7	11
2000	9,006	9	17	138	3,548	2,650	1,487	730	266	108	37	9	7
1999	8,944	5	26	155	3,569	2,609	1,414	676	294	124	48	10	14
All origins, female													
2005	1,791	-	15	56	403	436	411	238	93	70	53	14	2
2004	1,703	4	18	50	409	392	339	241	121	55	59	14	1
2003	1,794	1	20	53	433	415	406	250	102	46	53	13	2
2002	1,930	6	19	78	466	448	419	264	108	60	51	10	1
2001	1,816	4	28	55	404	444	450	208	100	69	45	9	-
2000	1,795	3	11	49	415	424	419	229	101	76	44	19	5
1999	1,884	3	24	69	469	439	454	217	79	69	47	13	1

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Assault (homicide) by discharge of firearms (*U01.4,X93-X95)													
Hispanic, both sexes													
2005	2,453	1	6	38	1,040	829	338	143	31	20	4	-	3
2004	2,241	3	7	38	991	715	310	110	44	11	3	5	4
2003	2,316	1	9	31	1,044	730	313	138	26	17	3	-	4
2002	2,168	2	12	40	976	679	294	116	31	9	4	1	4
2001	2,123	4	12	30	974	654	306	102	20	14	3	1	3
2000	1,958	1	5	41	904	581	249	126	34	14	1	-	2
1999	1,939	2	5	52	908	550	258	112	35	9	5	1	2
Hispanic, male													
2005	2,211	1	4	30	976	755	281	118	26	15	2	-	3
2004	2,031	1	4	26	924	660	268	92	37	10	1	4	4
2003	2,069	1	5	22	957	673	259	112	22	13	2	-	3
2002	1,942	-	8	29	909	619	245	91	26	7	3	1	4
2001	1,892	3	5	23	898	596	257	81	15	10	-	1	3
2000	1,738	-	3	34	837	524	207	99	26	7	-	-	1
1999	1,709	2	3	39	824	499	218	84	27	6	4	1	2
Hispanic, female													
2005	242	-	2	8	64	74	57	25	5	5	2	-	-
2004	210	2	3	12	67	55	42	18	7	1	2	1	-
2003	247	-	4	9	87	57	54	26	4	4	1	-	1
2002	226	2	4	11	67	60	49	25	5	2	1	-	-
2001	231	1	7	7	76	58	49	21	5	4	3	-	-
2000	220	1	2	7	67	57	42	27	8	7	1	-	1
1999	230	-	2	13	84	51	40	28	8	3	1	-	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Assault (homicide) by discharge of firearms (*U01.4,X93-X95)													
Non-Hispanic 1/, both sexes													
2005	9,835	5	31	148	3,441	2,933	1,658	947	372	181	93	24	2
2004	9,323	3	29	145	3,123	2,771	1,574	947	424	181	97	24	5
2003	9,536	7	31	156	3,339	2,792	1,616	970	365	144	84	23	9
2002	9,575	7	37	164	3,309	2,760	1,735	902	376	151	110	18	6
2001	9,134	7	43	149	3,186	2,631	1,662	827	341	175	96	14	3
2000	8,767	11	23	144	3,032	2,480	1,640	828	330	168	80	28	3
1999	8,821	6	45	172	3,115	2,476	1,597	776	335	184	89	22	4
Non-Hispanic 1/, male													
2005	8,296	5	18	101	3,105	2,572	1,305	736	284	116	42	10	2
2004	7,837	1	14	107	2,784	2,436	1,278	724	310	127	40	11	5
2003	7,997	6	15	112	2,996	2,436	1,267	746	267	102	32	10	8
2002	7,884	3	22	97	2,913	2,378	1,367	664	274	93	60	8	5
2001	7,559	4	22	101	2,863	2,246	1,264	641	246	110	54	5	3
2000	7,205	9	14	103	2,686	2,115	1,264	628	237	101	37	9	2
1999	7,177	3	23	116	2,734	2,091	1,184	588	264	118	43	9	4
Non-Hispanic 1/, female													
2005	1,539	-	13	47	336	361	353	211	88	65	51	14	-
2004	1,486	2	15	38	339	335	296	223	114	54	57	13	-
2003	1,539	1	16	44	343	356	349	224	98	42	52	13	1
2002	1,691	4	15	67	396	382	368	238	102	58	50	10	1
2001	1,575	3	21	48	323	385	398	186	95	65	42	9	-
2000	1,562	2	9	41	346	365	376	200	93	67	43	19	1
1999	1,644	3	22	56	381	385	413	188	71	66	46	13	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Assault (homicide) by discharge of firearms (*U01.4,X93-X95)													
Non-Hispanic white, both sexes													
2005 2/	2,871	3	17	51	563	633	655	489	241	120	78	19	2
2004 2/	2,921	-	7	56	561	640	650	498	273	132	81	19	4
2003 2/	2,883	1	11	51	610	613	649	516	242	99	65	20	6
2002	3,052	1	22	66	642	636	735	483	253	107	87	16	4
2001	3,085	4	21	90	618	681	759	460	235	126	75	14	2
2000	2,861	4	12	72	544	624	729	445	224	118	63	24	2
1999	2,995	2	21	80	591	694	749	416	215	140	67	20	-
Non-Hispanic white, male													
2005 2/	2,018	3	9	33	437	462	444	351	174	68	30	5	2
2004 2/	2,083	-	3	39	416	492	471	353	186	80	32	7	4
2003 2/	2,068	-	3	37	489	456	445	366	166	68	23	9	6
2002	2,139	1	12	34	488	465	531	321	174	58	44	8	3
2001	2,172	2	12	56	468	495	536	335	155	69	37	5	2
2000	1,967	2	6	47	401	444	510	311	149	63	25	8	1
1999	2,119	2	11	50	443	515	514	308	157	84	27	8	-
Non-Hispanic white, female													
2005 2/	853	-	8	18	126	171	211	138	67	52	48	14	-
2004 2/	838	-	4	17	145	148	179	145	87	52	49	12	-
2003 2/	815	1	8	14	121	157	204	150	76	31	42	11	-
2002	913	-	10	32	154	171	204	162	79	49	43	8	1
2001	913	2	9	34	150	186	223	125	80	57	38	9	-
2000	894	2	6	25	143	180	219	134	75	55	38	16	1
1999	876	-	10	30	148	179	235	108	58	56	40	12	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Assault (homicide) by discharge of firearms (*U01.4,X93-X95)													
Non-Hispanic black, both sexes													
2005 2/	6,600	2	13	86	2,726	2,216	949	426	108	54	15	5	-
2004 2/	6,119	3	19	80	2,474	2,063	879	408	132	41	14	5	1
2003 2/	6,319	6	19	95	2,606	2,100	909	415	106	42	16	3	2
2002	6,181	6	14	80	2,547	2,048	931	385	107	39	20	2	2
2001	5,790	2	18	56	2,490	1,874	864	332	92	43	18	-	1
2000	5,622	7	8	68	2,396	1,790	854	347	85	45	17	4	1
1999	5,508	2	18	81	2,426	1,695	796	320	104	39	21	2	4
Non-Hispanic black, male													
2005 2/	5,977	2	9	63	2,537	2,039	816	359	91	44	12	5	-
2004 2/	5,518	1	9	61	2,292	1,889	767	341	107	39	7	4	1
2003 2/	5,666	6	11	67	2,404	1,914	780	351	89	33	9	1	1
2002	5,482	2	9	54	2,325	1,854	784	317	88	32	15	-	2
2001	5,190	1	9	43	2,329	1,697	700	280	78	37	15	-	1
2000	5,011	7	5	52	2,206	1,616	715	290	69	37	12	1	1
1999	4,826	1	7	60	2,208	1,512	636	256	94	31	16	1	4
Non-Hispanic black, female													
2005 2/	623	-	4	23	189	177	133	67	17	10	3	-	-
2004 2/	601	2	10	19	182	174	112	67	25	2	7	1	-
2003 2/	653	-	8	28	202	186	129	64	17	9	7	2	1
2002	699	4	5	26	222	194	147	68	19	7	5	2	-
2001	600	1	9	13	161	177	164	52	14	6	3	-	-
2000	611	-	3	16	190	174	139	57	16	8	5	3	-
1999	682	1	11	21	218	183	160	64	10	8	5	1	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Assault (homicide) by discharge of firearms (*U01.4,X93-X95)													
Origin not stated 3/, both sexes													
2005	64	-	-	1	18	18	14	7	2	-	-	-	4
2004	60	-	-	1	13	17	11	5	-	1	-	-	12
2003	68	-	-	-	27	18	12	2	3	-	1	-	5
2002	86	-	-	1	32	26	13	7	2	1	1	-	3
2001	91	-	-	1	40	23	10	5	3	3	-	1	5
2000	76	-	-	2	27	13	17	5	3	2	-	-	7
1999	68	-	-	-	15	22	13	5	3	-	1	-	9
Origin not stated 3/, male													
2005	54	-	-	-	15	17	13	5	2	-	-	-	2
2004	53	-	-	1	10	15	10	5	-	1	-	-	11
2003	60	-	-	-	24	16	9	2	3	-	1	-	5
2002	73	-	-	1	29	20	11	6	1	1	1	-	3
2001	81	-	-	1	35	22	7	4	3	3	-	1	5
2000	63	-	-	1	25	11	16	3	3	-	-	-	4
1999	58	-	-	-	11	19	12	4	3	-	1	-	8
Origin not stated 3/, female													
2005	10	-	-	1	3	1	1	2	-	-	-	-	2
2004	7	-	-	-	3	2	1	-	-	-	-	-	1
2003	8	-	-	-	3	2	3	-	-	-	-	-	-
2002	13	-	-	-	3	6	2	1	1	-	-	-	-
2001	10	-	-	-	5	1	3	1	-	-	-	-	-
2000	13	-	-	1	2	2	1	2	-	2	-	-	3
1999	10	-	-	-	4	3	1	1	-	-	-	-	1

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Assault (homicide) by other and unspecified means and their sequelae (*U01.0-*U01.3,*U01.5-*U01.9,*U02,X85-X92,X96-Y09,Y87.1)													
All origins, both sexes													
2005	5,772	300	338	154	967	972	1,099	963	457	239	186	85	12
2004	5,733	319	341	145	958	992	1,089	946	411	253	189	75	15
2003	5,812	333	336	137	958	976	1,169	907	392	271	229	81	23
2002	5,809	294	374	151	902	1,024	1,197	890	432	260	181	76	28
2001 4/	8,960	321	360	146	1,097	1,896	2,290	1,533	654	340	213	89	21
2000	5,964	337	328	184	976	1,090	1,313	796	371	259	220	72	18
1999	6,061	323	326	208	960	1,183	1,338	775	349	280	209	76	34
All origins, male													
2005	3,815	167	192	73	669	664	718	720	324	151	94	34	9
2004	3,657	165	186	66	662	653	668	671	290	164	97	26	9
2003	3,756	198	182	81	647	616	773	628	276	170	136	36	13
2002	3,741	161	204	70	623	652	763	633	313	176	94	32	20
2001 4/	6,023	190	211	68	745	1,293	1,594	1,106	459	209	101	35	12
2000	3,814	194	181	89	655	711	832	554	271	171	117	31	8
1999	3,841	182	168	95	622	737	893	553	247	180	114	31	19
All origins, female													
2005	1,957	133	146	81	298	308	381	243	133	88	92	51	3
2004	2,076	154	155	79	296	339	421	275	121	89	92	49	6
2003	2,056	135	154	56	311	360	396	279	116	101	93	45	10
2002	2,068	133	170	81	279	372	434	257	119	84	87	44	8
2001 4/	2,937	131	149	78	352	603	696	427	195	131	112	54	9
2000	2,150	143	147	95	321	379	481	242	100	88	103	41	10
1999	2,220	141	158	113	338	446	445	222	102	100	95	45	15

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Assault (homicide) by other and unspecified means and their sequelae (*U01.0-*U01.3,*U01.5-*U01.9,*U02,X85-X92,X96-Y09,Y87.1)													
Hispanic, both sexes													
2005	1,067	59	76	24	267	256	182	121	40	16	19	3	4
2004	1,030	70	73	29	244	247	167	128	40	18	6	6	2
2003	1,039	59	73	22	242	241	189	108	49	29	20	5	2
2002	961	50	69	24	224	224	170	100	56	20	15	2	7
2001 4/	1,208	51	57	26	248	306	261	146	62	31	13	2	5
2000	959	53	52	35	222	229	198	85	43	22	15	4	1
1999	925	51	47	37	211	251	174	85	34	24	6	2	3
Hispanic, male													
2005	797	29	42	10	211	201	147	100	27	11	14	1	4
2004	751	41	41	14	201	184	116	101	32	13	4	2	2
2003	740	34	31	14	182	179	142	82	37	24	11	2	2
2002	693	28	41	8	172	170	122	75	46	18	6	1	6
2001 4/	864	30	33	15	201	204	194	109	44	23	8	1	2
2000	693	26	27	22	177	173	152	59	31	15	8	2	1
1999	690	24	22	20	166	199	136	66	29	20	4	1	3
Hispanic, female													
2005	270	30	34	14	56	55	35	21	13	5	5	2	-
2004	279	29	32	15	43	63	51	27	8	5	2	4	-
2003	299	25	42	8	60	62	47	26	12	5	9	3	-
2002	268	22	28	16	52	54	48	25	10	2	9	1	1
2001 4/	344	21	24	11	47	102	67	37	18	8	5	1	3
2000	266	27	25	13	45	56	46	26	12	7	7	2	-
1999	235	27	25	17	45	52	38	19	5	4	2	1	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Assault (homicide) by other and unspecified means and their sequelae (*U01.0-*U01.3,*U01.5-*U01.9,*U02,X85-X92,X96-Y09,Y87.1)													
Non-Hispanic 1/, both sexes													
2005	4,669	240	261	130	691	713	909	838	414	223	167	81	2
2004	4,642	244	268	116	707	735	914	798	369	234	182	69	6
2003	4,709	271	262	115	706	728	972	787	340	237	207	74	10
2002	4,771	239	303	125	674	788	1,011	780	371	237	163	74	6
2001 4/	7,592	263	300	120	842	1,547	1,990	1,355	575	305	200	87	8
2000	4,950	280	276	149	744	857	1,109	698	322	234	205	67	9
1999	5,052	264	278	169	745	926	1,146	676	307	253	203	73	12
Non-Hispanic 1/, male													
2005	2,992	138	150	63	450	461	565	616	294	140	80	33	2
2004	2,862	120	145	52	456	462	546	554	256	150	93	24	4
2003	2,972	162	151	67	459	432	627	536	236	141	124	32	5
2002	2,997	128	162	60	449	474	635	549	262	155	88	31	4
2001 4/	5,049	157	177	53	538	1,063	1,368	971	405	184	93	34	6
2000	3,085	165	154	67	473	535	675	487	236	153	109	28	3
1999	3,098	155	145	75	454	533	746	475	212	157	110	30	6
Non-Hispanic 1/, female													
2005	1,677	102	111	67	241	252	344	222	120	83	87	48	-
2004	1,780	124	123	64	251	273	368	244	113	84	89	45	2
2003	1,737	109	111	48	247	296	345	251	104	96	83	42	5
2002	1,774	111	141	65	225	314	376	231	109	82	75	43	2
2001 4/	2,543	106	123	67	304	484	622	384	170	121	107	53	2
2000	1,865	115	122	82	271	322	434	211	86	81	96	39	6
1999	1,954	109	133	94	291	393	400	201	95	96	93	43	6

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Assault (homicide) by other and unspecified means and their sequelae (*U01.0-*U01.3,*U01.5-*U01.9,*U02,X85-X92,X96-Y09,Y87.1)													
Non-Hispanic white, both sexes													
2005 2/	2,454	136	121	60	332	340	444	431	261	140	122	65	2
2004 2/	2,506	137	124	63	356	336	474	435	231	160	129	56	5
2003 2/	2,484	155	112	58	297	340	498	431	213	155	155	60	10
2002	2,519	116	147	67	316	341	533	415	243	168	110	57	6
2001 4/	4,843	135	139	74	440	961	1,322	913	417	216	150	71	5
2000	2,568	137	132	92	330	399	544	389	191	146	148	54	6
1999	2,753	141	135	96	371	441	578	398	203	179	147	56	8
Non-Hispanic white, male													
2005 2/	1,497	79	65	28	202	202	266	313	181	83	54	22	2
2004 2/	1,478	72	70	31	208	180	280	296	157	100	63	18	3
2003 2/	1,482	94	58	35	176	192	302	279	137	85	92	27	5
2002	1,522	69	77	27	197	195	343	265	158	108	55	24	4
2001 4/	3,281	82	83	27	261	682	962	655	299	128	73	26	3
2000	1,543	94	80	46	187	233	326	249	139	90	76	20	3
1999	1,628	92	72	43	201	238	367	271	136	104	82	19	3
Non-Hispanic white, female													
2005 2/	957	57	56	32	130	138	178	118	80	57	68	43	-
2004 2/	1,028	65	54	32	148	156	194	139	74	60	66	38	2
2003 2/	1,002	61	54	23	121	148	196	152	76	70	63	33	5
2002	997	47	70	40	119	146	190	150	85	60	55	33	2
2001 4/	1,562	53	56	47	179	279	360	258	118	88	77	45	2
2000	1,025	43	52	46	143	166	218	140	52	56	72	34	3
1999	1,125	49	63	53	170	203	211	127	67	75	65	37	5

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Assault (homicide) by other and unspecified means and their sequelae (*U01.0-*U01.3,*U01.5-*U01.9,*U02,X85-X92,X96-Y09,Y87.1)													
Non-Hispanic black, both sexes													
2005 2/	1,922	89	125	59	304	322	402	359	136	70	43	13	-
2004 2/	1,881	97	135	50	297	342	397	324	115	63	48	12	1
2003 2/	1,957	100	136	52	349	343	426	316	112	63	47	13	-
2002	1,966	108	142	52	307	377	426	325	109	56	49	15	-
2001 4/	2,287	117	148	36	327	461	563	372	127	74	45	14	3
2000	2,126	132	135	54	375	403	505	268	116	78	47	10	3
1999	2,028	112	133	64	326	427	509	237	90	64	46	16	4
Non-Hispanic black, male													
2005 2/	1,311	49	79	30	213	231	259	265	102	49	25	9	-
2004 2/	1,212	44	70	20	203	241	238	233	83	44	29	6	1
2003 2/	1,314	54	85	31	238	214	298	225	89	45	30	5	-
2002	1,298	52	80	30	215	238	258	255	92	40	32	6	-
2001 4/	1,468	67	91	23	222	301	336	264	87	48	19	7	3
2000	1,371	66	68	18	253	267	312	208	90	56	27	6	-
1999	1,305	57	67	29	225	259	344	175	65	48	23	10	3
Non-Hispanic black, female													
2005 2/	611	40	46	29	91	91	143	94	34	21	18	4	-
2004 2/	669	53	65	30	94	101	159	91	32	19	19	6	-
2003 2/	643	46	51	21	111	129	128	91	23	18	17	8	-
2002	668	56	62	22	92	139	168	70	17	16	17	9	-
2001 4/	819	50	57	13	105	160	227	108	40	26	26	7	-
2000	755	66	67	36	122	136	193	60	26	22	20	4	3
1999	723	55	66	35	101	168	165	62	25	16	23	6	1

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Assault (homicide) by other and unspecified means and their sequelae (*U01.0-*U01.3,*U01.5-*U01.9,*U02,X85-X92,X96-Y09,Y87.1)													
Origin not stated 3/, both sexes													
2005	36	1	1	-	9	3	8	4	3	-	-	1	6
2004	61	5	-	-	7	10	8	20	2	1	1	-	7
2003	64	3	1	-	10	7	8	12	3	5	2	2	11
2002	77	5	2	2	4	12	16	10	5	3	3	-	15
2001 4/	160	7	3	-	7	43	39	32	17	4	-	-	8
2000	55	4	-	-	10	4	6	13	6	3	-	1	8
1999	84	8	1	2	4	6	18	14	8	3	-	1	19
Origin not stated 3/, male													
2005	26	-	-	-	8	2	6	4	3	-	-	-	3
2004	44	4	-	-	5	7	6	16	2	1	-	-	3
2003	44	2	-	-	6	5	4	10	3	5	1	2	6
2002	51	5	1	2	2	8	6	9	5	3	-	-	10
2001 4/	110	3	1	-	6	26	32	26	10	2	-	-	4
2000	36	3	-	-	5	3	5	8	4	3	-	1	4
1999	53	3	1	-	2	5	11	12	6	3	-	-	10
Origin not stated 3/, female													
2005	10	1	1	-	1	1	2	-	-	-	-	1	3
2004	17	1	-	-	2	3	2	4	-	-	1	-	4
2003	20	1	1	-	4	2	4	2	-	-	1	-	5
2002	26	-	1	-	2	4	10	1	-	-	3	-	5
2001 4/	50	4	2	-	1	17	7	6	7	2	-	-	4
2000	19	1	-	-	5	1	1	5	2	-	-	-	4
1999	31	5	-	2	2	1	7	2	2	-	-	1	9

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Legal intervention (Y35,Y89.0)													
All origins, both sexes													
2005	414	-	-	1	83	110	124	56	30	7	3	-	-
2004	372	-	-	-	80	106	104	58	12	7	5	-	-
2003	423	-	-	2	93	135	117	51	19	4	2	-	-
2002	384	-	-	3	91	102	109	56	16	6	1	-	-
2001	396	-	-	-	88	107	115	51	24	6	4	1	-
2000	359	-	-	3	73	110	101	55	10	5	1	1	-
1999	398	-	-	1	67	136	121	50	20	2	1	-	-
All origins, male													
2005	401	-	-	1	82	109	119	54	28	5	3	-	-
2004	363	-	-	-	79	104	103	55	11	6	5	-	-
2003	405	-	-	1	89	131	114	46	18	4	2	-	-
2002	369	-	-	3	87	99	105	52	16	6	1	-	-
2001	380	-	-	-	84	104	108	51	22	6	4	1	-
2000	348	-	-	3	71	108	96	54	9	5	1	1	-
1999	392	-	-	1	67	135	117	49	20	2	1	-	-
All origins, female													
2005	13	-	-	-	1	1	5	2	2	2	-	-	-
2004	9	-	-	-	1	2	1	3	1	1	-	-	-
2003	18	-	-	1	4	4	3	5	1	-	-	-	-
2002	15	-	-	-	4	3	4	4	-	-	-	-	-
2001	16	-	-	-	4	3	7	-	2	-	-	-	-
2000	11	-	-	-	2	2	5	1	1	-	-	-	-
1999	6	-	-	-	-	1	4	1	-	-	-	-	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Legal intervention (Y35,Y89.0)													
Hispanic, both sexes													
2005	81	-	-	-	32	24	20	3	2	-	-	-	-
2004	77	-	-	-	30	30	13	4	-	-	-	-	-
2003	74	-	-	1	28	32	10	2	-	1	-	-	-
2002	68	-	-	1	22	22	14	7	1	1	-	-	-
2001	65	-	-	-	23	24	13	4	1	-	-	-	-
2000	48	-	-	1	18	15	9	4	1	-	-	-	-
1999	58	-	-	-	13	23	14	5	2	1	-	-	-
Hispanic, male													
2005	81	-	-	-	32	24	20	3	2	-	-	-	-
2004	77	-	-	-	30	30	13	4	-	-	-	-	-
2003	73	-	-	-	28	32	10	2	-	1	-	-	-
2002	67	-	-	1	22	21	14	7	1	1	-	-	-
2001	65	-	-	-	23	24	13	4	1	-	-	-	-
2000	47	-	-	1	17	15	9	4	1	-	-	-	-
1999	58	-	-	-	13	23	14	5	2	1	-	-	-
Hispanic, female													
2005	-	-	-	-	-	-	-	-	-	-	-	-	-
2004	-	-	-	-	-	-	-	-	-	-	-	-	-
2003	1	-	-	1	-	-	-	-	-	-	-	-	-
2002	1	-	-	-	-	1	-	-	-	-	-	-	-
2001	-	-	-	-	-	-	-	-	-	-	-	-	-
2000	1	-	-	-	1	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-	-	-	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Legal intervention (Y35,Y89.0)													
Non-Hispanic 1/, both sexes													
2005	333	-	-	1	51	86	104	53	28	7	3	-	-
2004	293	-	-	-	50	75	90	54	12	7	5	-	-
2003	347	-	-	1	64	103	107	49	18	3	2	-	-
2002	315	-	-	2	69	80	94	49	15	5	1	-	-
2001	330	-	-	-	65	83	102	46	23	6	4	1	-
2000	308	-	-	2	54	95	90	51	9	5	1	1	-
1999	338	-	-	1	54	112	106	45	18	1	1	-	-
Non-Hispanic 1/, male													
2005	320	-	-	1	50	85	99	51	26	5	3	-	-
2004	284	-	-	-	49	73	89	51	11	6	5	-	-
2003	330	-	-	1	60	99	104	44	17	3	2	-	-
2002	301	-	-	2	65	78	90	45	15	5	1	-	-
2001	314	-	-	-	61	80	95	46	21	6	4	1	-
2000	298	-	-	2	53	93	85	50	8	5	1	1	-
1999	332	-	-	1	54	111	102	44	18	1	1	-	-
Non-Hispanic 1/, female													
2005	13	-	-	-	1	1	5	2	2	2	-	-	-
2004	9	-	-	-	1	2	1	3	1	1	-	-	-
2003	17	-	-	-	4	4	3	5	1	-	-	-	-
2002	14	-	-	-	4	2	4	4	-	-	-	-	-
2001	16	-	-	-	4	3	7	-	2	-	-	-	-
2000	10	-	-	-	1	2	5	1	1	-	-	-	-
1999	6	-	-	-	-	1	4	1	-	-	-	-	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Legal intervention (Y35,Y89.0)													
Non-Hispanic white, both sexes													
2005 2/	208	-	-	-	27	44	65	40	24	6	2	-	-
2004 2/	187	-	-	-	27	37	67	38	8	5	5	-	-
2003 2/	211	-	-	-	27	45	80	39	16	3	1	-	-
2002	191	-	-	-	28	48	64	34	12	4	1	-	-
2001	230	-	-	-	37	56	74	36	17	6	4	-	-
2000	187	-	-	-	26	57	52	39	8	3	1	1	-
1999	197	-	-	-	20	62	65	35	13	1	1	-	-
Non-Hispanic white, male													
2005 2/	196	-	-	-	26	43	61	38	22	4	2	-	-
2004 2/	181	-	-	-	26	36	66	36	8	4	5	-	-
2003 2/	199	-	-	-	25	43	77	35	15	3	1	-	-
2002	181	-	-	-	25	47	62	30	12	4	1	-	-
2001	221	-	-	-	35	55	69	36	16	6	4	-	-
2000	182	-	-	-	25	55	51	39	7	3	1	1	-
1999	192	-	-	-	20	62	61	34	13	1	1	-	-
Non-Hispanic white, female													
2005 2/	12	-	-	-	1	1	4	2	2	2	-	-	-
2004 2/	6	-	-	-	1	1	1	2	-	1	-	-	-
2003 2/	12	-	-	-	2	2	3	4	1	-	-	-	-
2002	10	-	-	-	3	1	2	4	-	-	-	-	-
2001	9	-	-	-	2	1	5	-	1	-	-	-	-
2000	5	-	-	-	1	2	1	-	1	-	-	-	-
1999	5	-	-	-	-	-	4	1	-	-	-	-	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Legal intervention (Y35,Y89.0)													
Non-Hispanic black, both sexes													
2005 2/	112	-	-	1	24	39	31	13	2	1	1	-	-
2004 2/	90	-	-	-	19	33	19	13	4	2	-	-	-
2003 2/	117	-	-	1	33	52	22	7	1	-	1	-	-
2002	109	-	-	2	37	28	28	11	2	1	-	-	-
2001	82	-	-	-	26	21	22	9	3	-	-	1	-
2000	110	-	-	2	27	33	35	11	-	2	-	-	-
1999	125	-	-	1	31	42	36	10	5	-	-	-	-
Non-Hispanic black, male													
2005 2/	111	-	-	1	24	39	30	13	2	1	1	-	-
2004 2/	88	-	-	-	19	33	19	12	3	2	-	-	-
2003 2/	114	-	-	1	31	51	22	7	1	-	1	-	-
2002	105	-	-	2	36	27	26	11	2	1	-	-	-
2001	77	-	-	-	24	20	20	9	3	-	-	1	-
2000	107	-	-	2	27	33	33	10	-	2	-	-	-
1999	124	-	-	1	31	41	36	10	5	-	-	-	-
Non-Hispanic black, female													
2005 2/	1	-	-	-	-	-	1	-	-	-	-	-	-
2004 2/	2	-	-	-	-	-	-	1	1	-	-	-	-
2003 2/	3	-	-	-	2	1	-	-	-	-	-	-	-
2002	4	-	-	-	1	1	2	-	-	-	-	-	-
2001	5	-	-	-	2	1	2	-	-	-	-	-	-
2000	3	-	-	-	-	-	2	1	-	-	-	-	-
1999	1	-	-	-	-	1	-	-	-	-	-	-	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Legal intervention (Y35,Y89.0)													
Origin not stated 3/, both sexes													
2005	-	-	-	-	-	-	-	-	-	-	-	-	-
2004	2	-	-	-	-	1	1	-	-	-	-	-	-
2003	2	-	-	-	1	-	-	-	1	-	-	-	-
2002	1	-	-	-	-	-	1	-	-	-	-	-	-
2001	1	-	-	-	-	-	-	1	-	-	-	-	-
2000	3	-	-	-	1	-	2	-	-	-	-	-	-
1999	2	-	-	-	-	1	1	-	-	-	-	-	-
Origin not stated 3/, male													
2005	-	-	-	-	-	-	-	-	-	-	-	-	-
2004	2	-	-	-	-	1	1	-	-	-	-	-	-
2003	2	-	-	-	1	-	-	-	1	-	-	-	-
2002	1	-	-	-	-	-	1	-	-	-	-	-	-
2001	1	-	-	-	-	-	-	1	-	-	-	-	-
2000	3	-	-	-	1	-	2	-	-	-	-	-	-
1999	2	-	-	-	-	1	1	-	-	-	-	-	-
Origin not stated 3/, female													
2005	-	-	-	-	-	-	-	-	-	-	-	-	-
2004	-	-	-	-	-	-	-	-	-	-	-	-	-
2003	-	-	-	-	-	-	-	-	-	-	-	-	-
2002	-	-	-	-	-	-	-	-	-	-	-	-	-
2001	-	-	-	-	-	-	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-	-	-	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Events of undetermined intent (Y10-Y34,Y87.2,Y89.9)													
All origins, both sexes													
2005	4,742	104	69	64	484	742	1,194	1,314	424	157	106	74	10
2004	4,976	104	47	64	501	806	1,389	1,302	399	156	119	80	9
2003	5,072	75	31	55	563	835	1,581	1,267	368	131	96	52	18
2002	4,830	101	57	68	487	828	1,516	1,140	320	139	95	65	14
2001	4,198	83	44	51	412	728	1,396	990	225	99	108	44	18
2000	3,819	65	39	60	370	679	1,286	841	228	104	79	57	11
1999	3,917	55	43	49	430	743	1,295	793	222	114	96	58	19
All origins, male													
2005	2,861	67	44	43	355	478	695	758	245	91	49	26	10
2004	3,164	65	26	43	380	559	872	808	231	91	52	28	9
2003	3,295	50	17	41	412	580	1,013	811	222	65	49	21	14
2002	3,114	52	28	38	358	585	981	696	203	83	46	31	13
2001	2,833	49	19	38	316	515	931	662	146	64	57	20	16
2000	2,635	27	26	50	285	489	885	570	152	70	49	21	11
1999	2,692	33	21	38	345	523	887	532	148	72	55	25	13
All origins, female													
2005	1,881	37	25	21	129	264	499	556	179	66	57	48	-
2004	1,812	39	21	21	121	247	517	494	168	65	67	52	-
2003	1,777	25	14	14	151	255	568	456	146	66	47	31	4
2002	1,716	49	29	30	129	243	535	444	117	56	49	34	1
2001	1,365	34	25	13	96	213	465	328	79	35	51	24	2
2000	1,184	38	13	10	85	190	401	271	76	34	30	36	-
1999	1,225	22	22	11	85	220	408	261	74	42	41	33	6

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Events of undetermined intent (Y10-Y34,Y87.2,Y89.9)													
Hispanic, both sexes													
2005	325	17	11	4	60	59	74	67	18	5	9	-	1
2004	329	13	5	6	51	75	80	57	24	7	4	6	1
2003	339	6	6	13	51	85	80	66	19	6	5	2	-
2002	320	14	11	4	51	73	90	49	15	9	2	-	2
2001	283	7	6	4	34	63	92	54	9	6	2	4	2
2000	215	5	5	1	34	61	64	25	12	5	2	-	1
1999	249	6	9	4	52	69	61	28	13	3	3	-	1
Hispanic, male													
2005	219	9	8	2	40	44	53	39	14	4	5	-	1
2004	234	11	5	4	37	55	55	38	16	6	3	3	1
2003	258	1	4	9	36	69	65	50	13	5	4	2	-
2002	242	6	7	1	45	58	64	40	14	4	1	-	2
2001	216	4	3	4	26	52	73	36	7	5	1	3	2
2000	165	2	2	1	30	49	47	20	9	2	2	-	1
1999	200	3	6	3	46	54	49	23	11	2	2	-	1
Hispanic, female													
2005	106	8	3	2	20	15	21	28	4	1	4	-	-
2004	95	2	-	2	14	20	25	19	8	1	1	3	-
2003	81	5	2	4	15	16	15	16	6	1	1	-	-
2002	78	8	4	3	6	15	26	9	1	5	1	-	-
2001	67	3	3	-	8	11	19	18	2	1	1	1	-
2000	50	3	3	-	4	12	17	5	3	3	-	-	-
1999	49	3	3	1	6	15	12	5	2	1	1	-	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Events of undetermined intent (Y10-Y34,Y87.2,Y89.9)													
Non-Hispanic 1/, both sexes													
2005	4,388	87	58	60	423	680	1,115	1,237	403	150	97	74	4
2004	4,625	89	42	58	449	729	1,305	1,239	375	149	114	74	2
2003	4,701	68	25	42	512	748	1,491	1,196	346	124	91	50	8
2002	4,470	87	44	63	432	749	1,420	1,086	301	128	93	62	5
2001	3,876	76	38	47	377	661	1,298	927	210	91	105	40	6
2000	3,571	58	34	59	336	614	1,214	812	211	96	77	56	4
1999	3,630	49	34	45	376	666	1,230	757	207	107	93	56	10
Non-Hispanic 1/, male													
2005	2,621	58	36	41	314	432	641	711	228	86	44	26	4
2004	2,910	53	21	39	342	503	813	764	215	85	48	25	2
2003	3,016	48	13	32	376	510	942	759	206	59	45	19	7
2002	2,836	46	21	36	310	521	912	651	185	77	45	28	4
2001	2,586	45	16	34	289	459	853	622	134	57	55	17	5
2000	2,445	25	24	49	255	436	834	546	140	65	47	20	4
1999	2,463	30	15	35	297	462	834	503	135	67	53	24	8
Non-Hispanic 1/, female													
2005	1,767	29	22	19	109	248	474	526	175	64	53	48	-
2004	1,715	36	21	19	107	226	492	475	160	64	66	49	-
2003	1,685	20	12	10	136	238	549	437	140	65	46	31	1
2002	1,634	41	23	27	122	228	508	435	116	51	48	34	1
2001	1,290	31	22	13	88	202	445	305	76	34	50	23	1
2000	1,126	33	10	10	81	178	380	266	71	31	30	36	-
1999	1,167	19	19	10	79	204	396	254	72	40	40	32	2

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Events of undetermined intent (Y10-Y34,Y87.2,Y89.9)													
Non-Hispanic white, both sexes													
2005 2/	3,583	52	33	43	349	570	898	1,021	341	128	79	67	2
2004 2/	3,811	54	25	30	380	605	1,083	1,030	319	116	101	67	1
2003 2/	3,898	45	15	24	439	635	1,247	986	277	96	80	48	6
2002	3,627	43	33	44	370	607	1,160	873	253	105	77	58	4
2001	3,155	40	25	35	318	549	1,054	749	176	75	93	37	4
2000	2,823	35	22	37	271	490	965	647	161	76	70	47	2
1999	2,848	30	15	38	290	531	960	578	172	92	86	50	6
Non-Hispanic white, male													
2005 2/	2,099	36	21	30	261	356	513	569	182	71	33	25	2
2004 2/	2,365	32	11	20	291	417	667	619	175	66	42	24	1
2003 2/	2,441	28	7	20	316	422	780	604	155	45	41	18	5
2002	2,265	23	16	24	264	419	739	504	151	57	38	27	3
2001	2,054	25	10	25	238	377	673	485	106	45	51	16	3
2000	1,900	16	15	31	201	341	661	425	101	48	43	16	2
1999	1,887	19	8	30	232	357	646	363	101	57	47	22	5
Non-Hispanic white, female													
2005 2/	1,484	16	12	13	88	214	385	452	159	57	46	42	-
2004 2/	1,446	22	14	10	89	188	416	411	144	50	59	43	-
2003 2/	1,457	17	8	4	123	213	467	382	122	51	39	30	1
2002	1,362	20	17	20	106	188	421	369	102	48	39	31	1
2001	1,101	15	15	10	80	172	381	264	70	30	42	21	1
2000	923	19	7	6	70	149	304	222	60	28	27	31	-
1999	961	11	7	8	58	174	314	215	71	35	39	28	1

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Events of undetermined intent (Y10-Y34,Y87.2,Y89.9)													
Non-Hispanic black, both sexes													
2005 2/	675	31	22	11	55	90	188	188	50	18	13	7	2
2004 2/	691	31	15	26	56	97	189	189	50	22	10	5	1
2003 2/	675	21	9	14	50	93	213	190	53	21	7	2	2
2002	738	37	7	15	52	118	240	195	39	16	16	3	-
2001	629	36	13	11	43	90	222	163	30	12	5	3	1
2000	654	14	11	21	50	103	230	153	42	15	6	7	2
1999	686	18	15	7	70	119	239	166	28	12	5	3	4
Non-Hispanic black, male													
2005 2/	447	19	13	8	41	62	114	129	38	12	8	1	2
2004 2/	472	18	9	18	43	71	124	131	38	14	5	-	1
2003 2/	487	19	5	8	42	74	141	141	40	12	2	1	2
2002	504	18	3	11	39	84	161	136	30	15	7	-	-
2001	467	20	6	8	37	65	165	127	26	10	1	1	1
2000	478	3	8	17	45	78	160	113	32	13	4	3	2
1999	514	10	7	5	55	91	170	132	27	9	4	1	3
Non-Hispanic black, female													
2005 2/	228	12	9	3	14	28	74	59	12	6	5	6	-
2004 2/	219	13	6	8	13	26	65	58	12	8	5	5	-
2003 2/	188	2	4	6	8	19	72	49	13	9	5	1	-
2002	234	19	4	4	13	34	79	59	9	1	9	3	-
2001	162	16	7	3	6	25	57	36	4	2	4	2	-
2000	176	11	3	4	5	25	70	40	10	2	2	4	-
1999	172	8	8	2	15	28	69	34	1	3	1	2	1

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Events of undetermined intent (Y10-Y34,Y87.2,Y89.9)													
Origin not stated 3/, both sexes													
2005	29	-	-	-	1	3	5	10	3	2	-	-	5
2004	22	2	-	-	1	2	4	6	-	-	1	-	6
2003	32	1	-	-	-	2	10	5	3	1	-	-	10
2002	40	-	2	1	4	6	6	5	4	2	-	3	7
2001	39	-	-	-	1	4	6	9	6	2	1	-	10
2000	33	2	-	-	-	4	8	4	5	3	-	1	6
1999	38	-	-	-	2	8	4	8	2	4	-	2	8
Origin not stated 3/, male													
2005	21	-	-	-	1	2	1	8	3	1	-	-	5
2004	20	1	-	-	1	1	4	6	-	-	1	-	6
2003	21	1	-	-	-	1	6	2	3	1	-	-	7
2002	36	-	-	1	3	6	5	5	4	2	-	3	7
2001	31	-	-	-	1	4	5	4	5	2	1	-	9
2000	25	-	-	-	-	4	4	4	3	3	-	1	6
1999	29	-	-	-	2	7	4	6	2	3	-	1	4
Origin not stated 3/, female													
2005	8	-	-	-	-	1	4	2	-	1	-	-	-
2004	2	1	-	-	-	1	-	-	-	-	-	-	-
2003	11	-	-	-	-	1	4	3	-	-	-	-	3
2002	4	-	2	-	1	-	1	-	-	-	-	-	-
2001	8	-	-	-	-	-	1	5	1	-	-	-	1
2000	8	2	-	-	-	-	4	-	2	-	-	-	-
1999	9	-	-	-	-	1	-	2	-	1	-	1	4

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Discharge of firearms, undetermined intent (Y22-Y24)													
All origins, both sexes													
2005	221	-	3	12	77	35	28	28	23	10	3	2	-
2004	235	-	1	9	58	45	44	29	20	16	8	2	3
2003	232	-	1	12	69	38	43	31	20	8	9	1	-
2002	243	-	1	6	79	46	50	29	12	15	5	-	-
2001	231	-	-	6	64	48	50	29	12	10	11	1	-
2000	230	-	-	10	71	52	39	23	15	11	6	2	1
1999	324	-	3	12	127	51	57	28	14	15	13	4	-
All origins, male													
2005	178	-	2	10	68	29	21	22	14	7	3	2	-
2004	179	-	-	7	50	33	30	22	14	13	6	1	3
2003	186	-	1	12	60	33	26	22	16	7	8	1	-
2002	199	-	1	5	71	39	39	23	7	10	4	-	-
2001	190	-	-	6	59	42	33	21	11	8	9	1	-
2000	190	-	-	9	62	43	30	18	11	9	5	2	1
1999	277	-	3	12	111	44	45	21	11	14	12	4	-
All origins, female													
2005	43	-	1	2	9	6	7	6	9	3	-	-	-
2004	56	-	1	2	8	12	14	7	6	3	2	1	-
2003	46	-	-	-	9	5	17	9	4	1	1	-	-
2002	44	-	-	1	8	7	11	6	5	5	1	-	-
2001	41	-	-	-	5	6	17	8	1	2	2	-	-
2000	40	-	-	1	9	9	9	5	4	2	1	-	-
1999	47	-	-	-	16	7	12	7	3	1	1	-	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Discharge of firearms, undetermined intent (Y22-Y24)													
Hispanic, both sexes													
2005	27	-	-	-	17	8	2	-	-	-	-	-	-
2004	20	-	-	1	5	8	2	3	1	-	-	-	-
2003	31	-	-	3	13	6	5	3	-	1	-	-	-
2002	23	-	-	-	14	6	3	-	-	-	-	-	-
2001	21	-	-	1	8	4	4	3	1	-	-	-	-
2000	20	-	-	-	13	4	2	-	-	1	-	-	-
1999	31	-	1	1	18	3	3	2	1	2	-	-	-
Hispanic, male													
2005	21	-	-	-	13	6	2	-	-	-	-	-	-
2004	19	-	-	1	5	8	2	3	-	-	-	-	-
2003	26	-	-	3	9	6	5	2	-	1	-	-	-
2002	21	-	-	-	13	6	2	-	-	-	-	-	-
2001	16	-	-	1	7	3	2	2	1	-	-	-	-
2000	17	-	-	-	13	2	1	-	-	1	-	-	-
1999	26	-	1	1	16	2	2	1	1	2	-	-	-
Hispanic, female													
2005	6	-	-	-	4	2	-	-	-	-	-	-	-
2004	1	-	-	-	-	-	-	-	1	-	-	-	-
2003	5	-	-	-	4	-	-	1	-	-	-	-	-
2002	2	-	-	-	1	-	1	-	-	-	-	-	-
2001	5	-	-	-	1	1	2	1	-	-	-	-	-
2000	3	-	-	-	-	2	1	-	-	-	-	-	-
1999	5	-	-	-	2	1	1	1	-	-	-	-	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Discharge of firearms, undetermined intent (Y22-Y24)													
Non-Hispanic 1/, both sexes													
2005	192	-	3	12	60	27	26	26	23	10	3	2	-
2004	212	-	1	8	53	37	42	26	19	16	8	2	-
2003	201	-	1	9	56	32	38	28	20	7	9	1	-
2002	215	-	1	5	63	39	47	29	11	15	5	-	-
2001	208	-	-	5	56	43	46	26	11	9	11	1	-
2000	209	-	-	10	58	48	37	23	15	10	6	2	-
1999	293	-	2	11	109	48	54	26	13	13	13	4	-
Non-Hispanic 1/, male													
2005	156	-	2	10	55	23	19	21	14	7	3	2	-
2004	157	-	-	6	45	25	28	19	14	13	6	1	-
2003	160	-	1	9	51	27	21	20	16	6	8	1	-
2002	173	-	1	4	56	32	37	23	6	10	4	-	-
2001	172	-	-	5	52	38	31	19	10	7	9	1	-
2000	172	-	-	9	49	41	29	18	11	8	5	2	-
1999	251	-	2	11	95	42	43	20	10	12	12	4	-
Non-Hispanic 1/, female													
2005	36	-	1	2	5	4	7	5	9	3	-	-	-
2004	55	-	1	2	8	12	14	7	5	3	2	1	-
2003	41	-	-	-	5	5	17	8	4	1	1	-	-
2002	42	-	-	1	7	7	10	6	5	5	1	-	-
2001	36	-	-	-	4	5	15	7	1	2	2	-	-
2000	37	-	-	1	9	7	8	5	4	2	1	-	-
1999	42	-	-	-	14	6	11	6	3	1	1	-	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Discharge of firearms, undetermined intent (Y22-Y24)													
Non-Hispanic white, both sexes													
2005 2/	148	-	-	10	42	20	20	22	21	9	2	2	-
2004 2/	166	-	-	3	37	24	37	22	18	15	8	2	-
2003 2/	152	-	-	4	34	21	33	26	17	7	9	1	-
2002	169	-	-	3	47	27	38	27	9	14	4	-	-
2001	166	-	-	3	36	35	41	23	8	9	10	1	-
2000	148	-	-	5	36	30	30	20	11	8	6	2	-
1999	217	-	1	10	67	33	47	21	11	11	12	4	-
Non-Hispanic white, male													
2005 2/	121	-	-	8	38	16	15	19	14	7	2	2	-
2004 2/	122	-	-	3	32	15	24	15	14	12	6	1	-
2003 2/	118	-	-	4	31	17	17	19	15	6	8	1	-
2002	132	-	-	2	40	21	30	21	6	9	3	-	-
2001	133	-	-	3	33	31	27	16	7	7	8	1	-
2000	120	-	-	5	30	26	24	15	7	6	5	2	-
1999	187	-	1	10	60	29	39	15	8	10	11	4	-
Non-Hispanic white, female													
2005 2/	27	-	-	2	4	4	5	3	7	2	-	-	-
2004 2/	44	-	-	-	5	9	13	7	4	3	2	1	-
2003 2/	34	-	-	-	3	4	16	7	2	1	1	-	-
2002	37	-	-	1	7	6	8	6	3	5	1	-	-
2001	33	-	-	-	3	4	14	7	1	2	2	-	-
2000	28	-	-	-	6	4	6	5	4	2	1	-	-
1999	30	-	-	-	7	4	8	6	3	1	1	-	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Discharge of firearms, undetermined intent (Y22-Y24)													
Non-Hispanic black, both sexes													
2005 2/	36	-	2	1	16	6	5	3	2	1	-	-	-
2004 2/	37	-	1	4	15	8	4	3	1	1	-	-	-
2003 2/	35	-	1	4	17	7	3	2	1	-	-	-	-
2002	37	-	1	2	14	10	6	1	1	1	1	-	-
2001	31	-	-	2	15	5	2	3	3	-	1	-	-
2000	53	-	-	5	19	15	7	2	3	2	-	-	-
1999	65	-	1	1	35	14	6	4	1	2	1	-	-
Non-Hispanic black, male													
2005 2/	28	-	1	1	15	6	4	1	-	-	-	-	-
2004 2/	29	-	-	3	12	7	3	3	-	1	-	-	-
2003 2/	32	-	1	4	16	7	2	1	1	-	-	-	-
2002	35	-	1	2	14	10	5	1	-	1	1	-	-
2001	28	-	-	2	14	4	1	3	3	-	1	-	-
2000	46	-	-	4	18	12	5	2	3	2	-	-	-
1999	55	-	1	1	29	12	4	4	1	2	1	-	-
Non-Hispanic black, female													
2005 2/	8	-	1	-	1	-	1	2	2	1	-	-	-
2004 2/	8	-	1	1	3	1	1	-	1	-	-	-	-
2003 2/	3	-	-	-	1	-	1	1	-	-	-	-	-
2002	2	-	-	-	-	-	1	-	1	-	-	-	-
2001	3	-	-	-	1	1	1	-	-	-	-	-	-
2000	7	-	-	1	1	3	2	-	-	-	-	-	-
1999	10	-	-	-	6	2	2	-	-	-	-	-	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Discharge of firearms, undetermined intent (Y22-Y24)													
Origin not stated 3/, both sexes													
2005	2	-	-	-	-	-	-	2	-	-	-	-	-
2004	3	-	-	-	-	-	-	-	-	-	-	-	3
2003	-	-	-	-	-	-	-	-	-	-	-	-	-
2002	5	-	-	1	2	1	-	-	1	-	-	-	-
2001	2	-	-	-	-	1	-	-	-	1	-	-	-
2000	1	-	-	-	-	-	-	-	-	-	-	-	1
1999	-	-	-	-	-	-	-	-	-	-	-	-	-
Origin not stated 3/, male													
2005	1	-	-	-	-	-	-	1	-	-	-	-	-
2004	3	-	-	-	-	-	-	-	-	-	-	-	3
2003	-	-	-	-	-	-	-	-	-	-	-	-	-
2002	5	-	-	1	2	1	-	-	1	-	-	-	-
2001	2	-	-	-	-	1	-	-	-	1	-	-	-
2000	1	-	-	-	-	-	-	-	-	-	-	-	1
1999	-	-	-	-	-	-	-	-	-	-	-	-	-
Origin not stated 3/, female													
2005	1	-	-	-	-	-	-	1	-	-	-	-	-
2004	-	-	-	-	-	-	-	-	-	-	-	-	-
2003	-	-	-	-	-	-	-	-	-	-	-	-	-
2002	-	-	-	-	-	-	-	-	-	-	-	-	-
2001	-	-	-	-	-	-	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-	-	-	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Other and unspecified events of undetermined intent and their sequelae (Y10-Y21,Y25-Y34,Y87.2,Y89.9)													
All origins, both sexes													
2005	4,521	104	66	52	407	707	1,166	1,286	401	147	103	72	10
2004	4,741	104	46	55	443	761	1,345	1,273	379	140	111	78	6
2003	4,840	75	30	43	494	797	1,538	1,236	348	123	87	51	18
2002	4,587	101	56	62	408	782	1,466	1,111	308	124	90	65	14
2001	3,967	83	44	45	348	680	1,346	961	213	89	97	43	18
2000	3,589	65	39	50	299	627	1,247	818	213	93	73	55	10
1999	3,593	55	40	37	303	692	1,238	765	208	99	83	54	19
All origins, male													
2005	2,683	67	42	33	287	449	674	736	231	84	46	24	10
2004	2,985	65	26	36	330	526	842	786	217	78	46	27	6
2003	3,109	50	16	29	352	547	987	789	206	58	41	20	14
2002	2,915	52	27	33	287	546	942	673	196	73	42	31	13
2001	2,643	49	19	32	257	473	898	641	135	56	48	19	16
2000	2,445	27	26	41	223	446	855	552	141	61	44	19	10
1999	2,415	33	18	26	234	479	842	511	137	58	43	21	13
All origins, female													
2005	1,838	37	24	19	120	258	492	550	170	63	57	48	-
2004	1,756	39	20	19	113	235	503	487	162	62	65	51	-
2003	1,731	25	14	14	142	250	551	447	142	65	46	31	4
2002	1,672	49	29	29	121	236	524	438	112	51	48	34	1
2001	1,324	34	25	13	91	207	448	320	78	33	49	24	2
2000	1,144	38	13	9	76	181	392	266	72	32	29	36	-
1999	1,178	22	22	11	69	213	396	254	71	41	40	33	6

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Other and unspecified events of undetermined intent and their sequelae (Y10-Y21,Y25-Y34,Y87.2,Y89.9)													
Hispanic, both sexes													
2005	298	17	11	4	43	51	72	67	18	5	9	-	1
2004	309	13	5	5	46	67	78	54	23	7	4	6	1
2003	308	6	6	10	38	79	75	63	19	5	5	2	-
2002	297	14	11	4	37	67	87	49	15	9	2	-	2
2001	262	7	6	3	26	59	88	51	8	6	2	4	2
2000	195	5	5	1	21	57	62	25	12	4	2	-	1
1999	218	6	8	3	34	66	58	26	12	1	3	-	1
Hispanic, male													
2005	198	9	8	2	27	38	51	39	14	4	5	-	1
2004	215	11	5	3	32	47	53	35	16	6	3	3	1
2003	232	1	4	6	27	63	60	48	13	4	4	2	-
2002	221	6	7	1	32	52	62	40	14	4	1	-	2
2001	200	4	3	3	19	49	71	34	6	5	1	3	2
2000	148	2	2	1	17	47	46	20	9	1	2	-	1
1999	174	3	5	2	30	52	47	22	10	-	2	-	1
Hispanic, female													
2005	100	8	3	2	16	13	21	28	4	1	4	-	-
2004	94	2	-	2	14	20	25	19	7	1	1	3	-
2003	76	5	2	4	11	16	15	15	6	1	1	-	-
2002	76	8	4	3	5	15	25	9	1	5	1	-	-
2001	62	3	3	-	7	10	17	17	2	1	1	1	-
2000	47	3	3	-	4	10	16	5	3	3	-	-	-
1999	44	3	3	1	4	14	11	4	2	1	1	-	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Other and unspecified events of undetermined intent and their sequelae (Y10-Y21,Y25-Y34,Y87.2,Y89.9)													
Non-Hispanic 1/, both sexes													
2005	4,196	87	55	48	363	653	1,089	1,211	380	140	94	72	4
2004	4,413	89	41	50	396	692	1,263	1,213	356	133	106	72	2
2003	4,500	68	24	33	456	716	1,453	1,168	326	117	82	49	8
2002	4,255	87	43	58	369	710	1,373	1,057	290	113	88	62	5
2001	3,668	76	38	42	321	618	1,252	901	199	82	94	39	6
2000	3,362	58	34	49	278	566	1,177	789	196	86	71	54	4
1999	3,337	49	32	34	267	618	1,176	731	194	94	80	52	10
Non-Hispanic 1/, male													
2005	2,465	58	34	31	259	409	622	690	214	79	41	24	4
2004	2,753	53	21	33	297	478	785	745	201	72	42	24	2
2003	2,856	48	12	23	325	483	921	739	190	53	37	18	7
2002	2,663	46	20	32	254	489	875	628	179	67	41	28	4
2001	2,414	45	16	29	237	421	822	603	124	50	46	16	5
2000	2,273	25	24	40	206	395	805	528	129	57	42	18	4
1999	2,212	30	13	24	202	420	791	483	125	55	41	20	8
Non-Hispanic 1/, female													
2005	1,731	29	21	17	104	244	467	521	166	61	53	48	-
2004	1,660	36	20	17	99	214	478	468	155	61	64	48	-
2003	1,644	20	12	10	131	233	532	429	136	64	45	31	1
2002	1,592	41	23	26	115	221	498	429	111	46	47	34	1
2001	1,254	31	22	13	84	197	430	298	75	32	48	23	1
2000	1,089	33	10	9	72	171	372	261	67	29	29	36	-
1999	1,125	19	19	10	65	198	385	248	69	39	39	32	2

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Other and unspecified events of undetermined intent and their sequelae (Y10-Y21,Y25-Y34,Y87.2,Y89.9)													
Non-Hispanic white, both sexes													
2005 2/	3,435	52	33	33	307	550	878	999	320	119	77	65	2
2004 2/	3,645	54	25	27	343	581	1,046	1,008	301	101	93	65	1
2003 2/	3,746	45	15	20	405	614	1,214	960	260	89	71	47	6
2002	3,458	43	33	41	323	580	1,122	846	244	91	73	58	4
2001	2,989	40	25	32	282	514	1,013	726	168	66	83	36	4
2000	2,675	35	22	32	235	460	935	627	150	68	64	45	2
1999	2,631	30	14	28	223	498	913	557	161	81	74	46	6
Non-Hispanic white, male													
2005 2/	1,978	36	21	22	223	340	498	550	168	64	31	23	2
2004 2/	2,243	32	11	17	259	402	643	604	161	54	36	23	1
2003 2/	2,323	28	7	16	285	405	763	585	140	39	33	17	5
2002	2,133	23	16	22	224	398	709	483	145	48	35	27	3
2001	1,921	25	10	22	205	346	646	469	99	38	43	15	3
2000	1,780	16	15	26	171	315	637	410	94	42	38	14	2
1999	1,700	19	7	20	172	328	607	348	93	47	36	18	5
Non-Hispanic white, female													
2005 2/	1,457	16	12	11	84	210	380	449	152	55	46	42	-
2004 2/	1,402	22	14	10	84	179	403	404	140	47	57	42	-
2003 2/	1,423	17	8	4	120	209	451	375	120	50	38	30	1
2002	1,325	20	17	19	99	182	413	363	99	43	38	31	1
2001	1,068	15	15	10	77	168	367	257	69	28	40	21	1
2000	895	19	7	6	64	145	298	217	56	26	26	31	-
1999	931	11	7	8	51	170	306	209	68	34	38	28	1

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Other and unspecified events of undetermined intent and their sequelae (Y10-Y21,Y25-Y34,Y87.2,Y89.9)													
Non-Hispanic black, both sexes													
2005 2/	639	31	20	10	39	84	183	185	48	17	13	7	2
2004 2/	654	31	14	22	41	89	185	186	49	21	10	5	1
2003 2/	640	21	8	10	33	86	210	188	52	21	7	2	2
2002	701	37	6	13	38	108	234	194	38	15	15	3	-
2001	598	36	13	9	28	85	220	160	27	12	4	3	1
2000	601	14	11	16	31	88	223	151	39	13	6	7	2
1999	621	18	14	6	35	105	233	162	27	10	4	3	4
Non-Hispanic black, male													
2005 2/	419	19	12	7	26	56	110	128	38	12	8	1	2
2004 2/	443	18	9	15	31	64	121	128	38	13	5	-	1
2003 2/	455	19	4	4	26	67	139	140	39	12	2	1	2
2002	469	18	2	9	25	74	156	135	30	14	6	-	-
2001	439	20	6	6	23	61	164	124	23	10	-	1	1
2000	432	3	8	13	27	66	155	111	29	11	4	3	2
1999	459	10	6	4	26	79	166	128	26	7	3	1	3
Non-Hispanic black, female													
2005 2/	220	12	8	3	13	28	73	57	10	5	5	6	-
2004 2/	211	13	5	7	10	25	64	58	11	8	5	5	-
2003 2/	185	2	4	6	7	19	71	48	13	9	5	1	-
2002	232	19	4	4	13	34	78	59	8	1	9	3	-
2001	159	16	7	3	5	24	56	36	4	2	4	2	-
2000	169	11	3	3	4	22	68	40	10	2	2	4	-
1999	162	8	8	2	9	26	67	34	1	3	1	2	1

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Other and unspecified events of undetermined intent and their sequelae (Y10-Y21,Y25-Y34,Y87.2,Y89.9)													
Origin not stated 3/, both sexes													
2005	27	-	-	-	1	3	5	8	3	2	-	-	5
2004	19	2	-	-	1	2	4	6	-	-	1	-	3
2003	32	1	-	-	-	2	10	5	3	1	-	-	10
2002	35	-	2	-	2	5	6	5	3	2	-	3	7
2001	37	-	-	-	1	3	6	9	6	1	1	-	10
2000	32	2	-	-	-	4	8	4	5	3	-	1	5
1999	38	-	-	-	2	8	4	8	2	4	-	2	8
Origin not stated 3/, male													
2005	20	-	-	-	1	2	1	7	3	1	-	-	5
2004	17	1	-	-	1	1	4	6	-	-	1	-	3
2003	21	1	-	-	-	1	6	2	3	1	-	-	7
2002	31	-	-	-	1	5	5	5	3	2	-	3	7
2001	29	-	-	-	1	3	5	4	5	1	1	-	9
2000	24	-	-	-	-	4	4	4	3	3	-	1	5
1999	29	-	-	-	2	7	4	6	2	3	-	1	4
Origin not stated 3/, female													
2005	7	-	-	-	-	1	4	1	-	1	-	-	-
2004	2	1	-	-	-	1	-	-	-	-	-	-	-
2003	11	-	-	-	-	1	4	3	-	-	-	-	3
2002	4	-	2	-	1	-	1	-	-	-	-	-	-
2001	8	-	-	-	-	-	1	5	1	-	-	-	1
2000	8	2	-	-	-	-	4	-	2	-	-	-	-
1999	9	-	-	-	-	1	-	2	-	1	-	1	4

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Operations of war and their sequelae (Y36,Y89.1)													
All origins, both sexes													
2005	27	-	-	-	3	3	4	1	3	2	10	1	-
2004	28	-	-	-	4	1	3	-	4	1	14	1	-
2003	14	-	-	-	1	-	-	1	5	2	5	-	-
2002	20	-	-	-	-	1	1	1	3	3	10	1	-
2001	17	-	-	-	1	-	1	2	-	4	8	1	-
2000	16	-	-	-	-	-	-	2	1	3	9	1	-
1999	23	-	-	-	-	-	1	2	1	8	8	3	-
All origins, male													
2005	27	-	-	-	3	3	4	1	3	2	10	1	-
2004	28	-	-	-	4	1	3	-	4	1	14	1	-
2003	14	-	-	-	1	-	-	1	5	2	5	-	-
2002	20	-	-	-	-	1	1	1	3	3	10	1	-
2001	16	-	-	-	1	-	1	2	-	4	7	1	-
2000	16	-	-	-	-	-	-	2	1	3	9	1	-
1999	22	-	-	-	-	-	1	2	1	8	7	3	-
All origins, female													
2005	-	-	-	-	-	-	-	-	-	-	-	-	-
2004	-	-	-	-	-	-	-	-	-	-	-	-	-
2003	-	-	-	-	-	-	-	-	-	-	-	-	-
2002	-	-	-	-	-	-	-	-	-	-	-	-	-
2001	1	-	-	-	-	-	-	-	-	-	1	-	-
2000	-	-	-	-	-	-	-	-	-	-	-	-	-
1999	1	-	-	-	-	-	-	-	-	-	1	-	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Operations of war and their sequelae (Y36,Y89.1)													
Hispanic, both sexes													
2005	2	-	-	-	1	-	1	-	-	-	-	-	-
2004	1	-	-	-	-	-	-	-	-	-	1	-	-
2003	-	-	-	-	-	-	-	-	-	-	-	-	-
2002	1	-	-	-	-	1	-	-	-	-	-	-	-
2001	2	-	-	-	1	-	-	-	-	-	1	-	-
2000	1	-	-	-	-	-	-	-	-	1	-	-	-
1999	2	-	-	-	-	-	-	-	-	-	1	1	-
Hispanic, male													
2005	2	-	-	-	1	-	1	-	-	-	-	-	-
2004	1	-	-	-	-	-	-	-	-	-	1	-	-
2003	-	-	-	-	-	-	-	-	-	-	-	-	-
2002	1	-	-	-	-	1	-	-	-	-	-	-	-
2001	2	-	-	-	1	-	-	-	-	-	1	-	-
2000	1	-	-	-	-	-	-	-	-	1	-	-	-
1999	2	-	-	-	-	-	-	-	-	-	1	1	-
Hispanic, female													
2005	-	-	-	-	-	-	-	-	-	-	-	-	-
2004	-	-	-	-	-	-	-	-	-	-	-	-	-
2003	-	-	-	-	-	-	-	-	-	-	-	-	-
2002	-	-	-	-	-	-	-	-	-	-	-	-	-
2001	-	-	-	-	-	-	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-	-	-	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Operations of war and their sequelae (Y36,Y89.1)													
Non-Hispanic 1/, both sexes													
2005	25	-	-	-	2	3	3	1	3	2	10	1	-
2004	27	-	-	-	4	1	3	-	4	1	13	1	-
2003	14	-	-	-	1	-	-	1	5	2	5	-	-
2002	19	-	-	-	-	-	1	1	3	3	10	1	-
2001	15	-	-	-	-	-	1	2	-	4	7	1	-
2000	15	-	-	-	-	-	-	2	1	2	9	1	-
1999	21	-	-	-	-	-	1	2	1	8	7	2	-
Non-Hispanic 1/, male													
2005	25	-	-	-	2	3	3	1	3	2	10	1	-
2004	27	-	-	-	4	1	3	-	4	1	13	1	-
2003	14	-	-	-	1	-	-	1	5	2	5	-	-
2002	19	-	-	-	-	-	1	1	3	3	10	1	-
2001	14	-	-	-	-	-	1	2	-	4	6	1	-
2000	15	-	-	-	-	-	-	2	1	2	9	1	-
1999	20	-	-	-	-	-	1	2	1	8	6	2	-
Non-Hispanic 1/, female													
2005	-	-	-	-	-	-	-	-	-	-	-	-	-
2004	-	-	-	-	-	-	-	-	-	-	-	-	-
2003	-	-	-	-	-	-	-	-	-	-	-	-	-
2002	-	-	-	-	-	-	-	-	-	-	-	-	-
2001	1	-	-	-	-	-	-	-	-	-	1	-	-
2000	-	-	-	-	-	-	-	-	-	-	-	-	-
1999	1	-	-	-	-	-	-	-	-	-	1	-	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Operations of war and their sequelae (Y36,Y89.1)													
Non-Hispanic white, both sexes													
2005 2/	19	-	-	-	2	1	2	-	2	2	9	1	-
2004 2/	25	-	-	-	3	1	2	-	4	1	13	1	-
2003 2/	13	-	-	-	1	-	-	1	4	2	5	-	-
2002	14	-	-	-	-	-	-	-	2	2	9	1	-
2001	14	-	-	-	-	-	1	2	-	3	7	1	-
2000	14	-	-	-	-	-	-	1	1	2	9	1	-
1999	20	-	-	-	-	-	1	2	1	7	7	2	-
Non-Hispanic white, male													
2005 2/	19	-	-	-	2	1	2	-	2	2	9	1	-
2004 2/	25	-	-	-	3	1	2	-	4	1	13	1	-
2003 2/	13	-	-	-	1	-	-	1	4	2	5	-	-
2002	14	-	-	-	-	-	-	-	2	2	9	1	-
2001	13	-	-	-	-	-	1	2	-	3	6	1	-
2000	14	-	-	-	-	-	-	1	1	2	9	1	-
1999	19	-	-	-	-	-	1	2	1	7	6	2	-
Non-Hispanic white, female													
2005 2/	-	-	-	-	-	-	-	-	-	-	-	-	-
2004 2/	-	-	-	-	-	-	-	-	-	-	-	-	-
2003 2/	-	-	-	-	-	-	-	-	-	-	-	-	-
2002	-	-	-	-	-	-	-	-	-	-	-	-	-
2001	1	-	-	-	-	-	-	-	-	-	1	-	-
2000	-	-	-	-	-	-	-	-	-	-	-	-	-
1999	1	-	-	-	-	-	-	-	-	-	1	-	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Operations of war and their sequelae (Y36,Y89.1)													
Non-Hispanic black, both sexes													
2005 2/	6	-	-	-	-	2	1	1	1	-	1	-	-
2004 2/	2	-	-	-	1	-	1	-	-	-	-	-	-
2003 2/	1	-	-	-	-	-	-	-	1	-	-	-	-
2002	5	-	-	-	-	-	1	1	1	1	1	-	-
2001	1	-	-	-	-	-	-	-	-	1	-	-	-
2000	1	-	-	-	-	-	-	1	-	-	-	-	-
1999	1	-	-	-	-	-	-	-	-	1	-	-	-
Non-Hispanic black, male													
2005 2/	6	-	-	-	-	2	1	1	1	-	1	-	-
2004 2/	2	-	-	-	1	-	1	-	-	-	-	-	-
2003 2/	1	-	-	-	-	-	-	-	1	-	-	-	-
2002	5	-	-	-	-	-	1	1	1	1	1	-	-
2001	1	-	-	-	-	-	-	-	-	1	-	-	-
2000	1	-	-	-	-	-	-	1	-	-	-	-	-
1999	1	-	-	-	-	-	-	-	-	1	-	-	-
Non-Hispanic black, female													
2005 2/	-	-	-	-	-	-	-	-	-	-	-	-	-
2004 2/	-	-	-	-	-	-	-	-	-	-	-	-	-
2003 2/	-	-	-	-	-	-	-	-	-	-	-	-	-
2002	-	-	-	-	-	-	-	-	-	-	-	-	-
2001	-	-	-	-	-	-	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-	-	-	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Operations of war and their sequelae (Y36,Y89.1)													
Origin not stated 3/, both sexes													
2005	-	-	-	-	-	-	-	-	-	-	-	-	-
2004	-	-	-	-	-	-	-	-	-	-	-	-	-
2003	-	-	-	-	-	-	-	-	-	-	-	-	-
2002	-	-	-	-	-	-	-	-	-	-	-	-	-
2001	-	-	-	-	-	-	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-	-	-	-
Origin not stated 3/, male													
2005	-	-	-	-	-	-	-	-	-	-	-	-	-
2004	-	-	-	-	-	-	-	-	-	-	-	-	-
2003	-	-	-	-	-	-	-	-	-	-	-	-	-
2002	-	-	-	-	-	-	-	-	-	-	-	-	-
2001	-	-	-	-	-	-	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-	-	-	-
Origin not stated 3/, female													
2005	-	-	-	-	-	-	-	-	-	-	-	-	-
2004	-	-	-	-	-	-	-	-	-	-	-	-	-
2003	-	-	-	-	-	-	-	-	-	-	-	-	-
2002	-	-	-	-	-	-	-	-	-	-	-	-	-
2001	-	-	-	-	-	-	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-	-	-	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Complications of medical and surgical care (Y40-Y84,Y88)													
All origins, both sexes													
2005	2,653	19	16	16	29	70	144	287	364	487	759	462	-
2004	2,883	16	19	19	48	71	157	293	403	557	787	513	-
2003	2,855	15	24	12	44	69	179	273	380	534	794	530	1
2002	2,843	15	19	35	35	79	171	265	351	560	792	521	-
2001	3,021	26	24	14	43	79	172	283	384	567	864	564	1
2000	3,059	19	22	31	32	85	149	295	376	647	825	578	-
1999	2,823	54	30	18	36	78	152	214	315	586	811	529	-
All origins, male													
2005	1,220	16	10	9	16	26	62	131	192	250	335	173	-
2004	1,364	6	10	12	24	31	68	148	220	280	385	180	-
2003	1,333	8	16	4	26	36	96	146	195	264	355	186	1
2002	1,236	8	11	26	19	47	82	127	169	256	320	171	-
2001	1,357	15	14	8	19	42	83	143	196	265	383	189	-
2000	1,430	12	11	15	16	47	76	143	181	331	402	196	-
1999	1,308	31	12	9	21	43	78	98	165	279	375	197	-
All origins, female													
2005	1,433	3	6	7	13	44	82	156	172	237	424	289	-
2004	1,519	10	9	7	24	40	89	145	183	277	402	333	-
2003	1,522	7	8	8	18	33	83	127	185	270	439	344	-
2002	1,607	7	8	9	16	32	89	138	182	304	472	350	-
2001	1,664	11	10	6	24	37	89	140	188	302	481	375	1
2000	1,629	7	11	16	16	38	73	152	195	316	423	382	-
1999	1,515	23	18	9	15	35	74	116	150	307	436	332	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Complications of medical and surgical care (Y40-Y84,Y88)													
Hispanic, both sexes													
2005	160	7	5	3	3	7	12	27	19	30	36	11	-
2004	172	6	5	6	5	15	15	25	24	29	32	10	-
2003	155	3	4	1	5	9	22	14	21	38	24	13	1
2002	137	2	7	5	6	8	13	14	19	28	29	6	-
2001	145	3	6	4	8	7	12	22	13	25	29	15	1
2000	136	3	4	6	7	4	11	13	11	35	26	16	-
1999	100	5	6	3	4	8	8	11	16	15	12	12	-
Hispanic, male													
2005	61	5	2	1	-	3	6	9	9	12	12	2	-
2004	79	1	2	3	3	6	5	13	13	14	18	1	-
2003	76	1	4	-	3	2	15	10	9	17	10	4	1
2002	57	2	4	4	3	1	5	8	6	11	13	-	-
2001	65	-	3	3	3	3	4	11	8	8	14	8	-
2000	70	2	2	4	6	2	5	4	8	16	15	6	-
1999	52	3	3	2	2	3	7	6	9	5	8	4	-
Hispanic, female													
2005	99	2	3	2	3	4	6	18	10	18	24	9	-
2004	93	5	3	3	2	9	10	12	11	15	14	9	-
2003	79	2	-	1	2	7	7	4	12	21	14	9	-
2002	80	-	3	1	3	7	8	6	13	17	16	6	-
2001	80	3	3	1	5	4	8	11	5	17	15	7	1
2000	66	1	2	2	1	2	6	9	3	19	11	10	-
1999	48	2	3	1	2	5	1	5	7	10	4	8	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Complications of medical and surgical care (Y40-Y84,Y88)													
Non-Hispanic 1/, both sexes													
2005	2,484	12	11	13	26	63	132	260	344	455	719	449	-
2004	2,708	10	14	13	43	56	142	268	379	528	753	502	-
2003	2,693	12	19	11	38	60	156	257	359	496	769	516	-
2002	2,695	13	12	30	29	71	158	249	332	530	758	513	-
2001	2,861	23	18	10	34	72	158	261	369	537	831	548	-
2000	2,914	16	18	25	25	81	137	281	363	610	797	561	-
1999	2,714	49	24	15	32	70	141	202	298	570	798	515	-
Non-Hispanic 1/, male													
2005	1,152	11	8	8	16	23	56	122	182	237	319	170	-
2004	1,285	5	8	9	21	25	63	135	207	266	367	179	-
2003	1,252	7	11	4	22	34	80	135	186	247	344	182	-
2002	1,175	6	7	22	16	46	77	117	163	244	306	171	-
2001	1,286	15	11	5	15	39	78	132	188	254	368	181	-
2000	1,354	10	9	11	10	45	70	139	172	313	385	190	-
1999	1,253	28	9	7	19	40	70	91	156	274	366	193	-
Non-Hispanic 1/, female													
2005	1,332	1	3	5	10	40	76	138	162	218	400	279	-
2004	1,423	5	6	4	22	31	79	133	172	262	386	323	-
2003	1,441	5	8	7	16	26	76	122	173	249	425	334	-
2002	1,520	7	5	8	13	25	81	132	169	286	452	342	-
2001	1,575	8	7	5	19	33	80	129	181	283	463	367	-
2000	1,560	6	9	14	15	36	67	142	191	297	412	371	-
1999	1,461	21	15	8	13	30	71	111	142	296	432	322	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Complications of medical and surgical care (Y40-Y84,Y88)													
Non-Hispanic white, both sexes													
2005 2/	2,009	4	9	11	21	47	100	178	264	347	620	408	-
2004 2/	2,219	5	11	10	28	42	98	201	300	431	649	444	-
2003 2/	2,202	6	9	7	31	39	119	178	276	410	671	456	-
2002	2,188	6	7	16	21	49	107	169	252	440	664	457	-
2001	2,294	14	10	6	25	51	105	185	279	417	735	467	-
2000	2,378	12	8	17	17	56	103	218	272	498	691	486	-
1999	2,212	30	16	11	26	47	103	148	224	478	688	441	-
Non-Hispanic white, male													
2005 2/	952	3	8	7	13	20	47	86	136	194	279	159	-
2004 2/	1,084	3	6	7	13	20	49	107	167	227	326	159	-
2003 2/	1,044	4	5	3	18	23	66	88	148	214	312	163	-
2002	966	3	5	12	13	34	60	85	129	200	276	149	-
2001	1,016	9	7	2	12	26	50	91	137	197	333	152	-
2000	1,135	9	6	7	7	31	58	111	135	255	347	169	-
1999	1,025	19	8	5	15	29	52	65	117	230	321	164	-
Non-Hispanic white, female													
2005 2/	1,057	1	1	4	8	27	53	92	128	153	341	249	-
2004 2/	1,135	2	5	3	15	22	49	94	133	204	323	285	-
2003 2/	1,158	2	4	4	13	16	53	90	128	196	359	293	-
2002	1,222	3	2	4	8	15	47	84	123	240	388	308	-
2001	1,278	5	3	4	13	25	55	94	142	220	402	315	-
2000	1,243	3	2	10	10	25	45	107	137	243	344	317	-
1999	1,187	11	8	6	11	18	51	83	107	248	367	277	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Complications of medical and surgical care (Y40-Y84,Y88)													
Non-Hispanic black, both sexes													
2005 2/	428	8	2	2	4	15	29	76	75	94	89	34	-
2004 2/	435	2	3	3	13	10	43	60	74	86	92	49	-
2003 2/	425	5	9	3	6	16	33	66	74	76	86	51	-
2002	445	7	5	14	5	17	48	75	70	77	80	47	-
2001	515	9	7	3	7	21	48	70	84	104	86	76	-
2000	484	4	8	5	7	23	31	57	86	101	95	67	-
1999	442	18	8	4	6	18	35	51	61	84	94	63	-
Non-Hispanic black, male													
2005 2/	177	8	-	1	2	3	8	34	41	36	34	10	-
2004 2/	172	1	2	2	6	3	14	23	37	33	36	15	-
2003 2/	174	3	5	1	3	8	11	40	35	27	27	14	-
2002	182	3	2	10	2	9	16	29	30	37	25	19	-
2001	247	6	4	2	2	13	26	38	47	53	30	26	-
2000	192	1	2	2	2	13	12	25	34	52	33	16	-
1999	195	8	1	2	4	10	16	26	30	40	38	20	-
Non-Hispanic black, female													
2005 2/	251	-	2	1	2	12	21	42	34	58	55	24	-
2004 2/	263	1	1	1	7	7	29	37	37	53	56	34	-
2003 2/	251	2	4	2	3	8	22	26	39	49	59	37	-
2002	263	4	3	4	3	8	32	46	40	40	55	28	-
2001	268	3	3	1	5	8	22	32	37	51	56	50	-
2000	292	3	6	3	5	10	19	32	52	49	62	51	-
1999	247	10	7	2	2	8	19	25	31	44	56	43	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Complications of medical and surgical care (Y40-Y84,Y88)													
Origin not stated 3/, both sexes													
2005	9	-	-	-	-	-	-	-	1	2	4	2	-
2004	3	-	-	-	-	-	-	-	-	-	2	1	-
2003	7	-	1	-	1	-	1	2	-	-	1	1	-
2002	11	-	-	-	-	-	-	2	-	2	5	2	-
2001	15	-	-	-	1	-	2	-	2	5	4	1	-
2000	9	-	-	-	-	-	1	1	2	2	2	1	-
1999	9	-	-	-	-	-	3	1	1	1	1	2	-
Origin not stated 3/, male													
2005	7	-	-	-	-	-	-	-	1	1	4	1	-
2004	-	-	-	-	-	-	-	-	-	-	-	-	-
2003	5	-	1	-	1	-	1	1	-	-	1	-	-
2002	4	-	-	-	-	-	-	2	-	1	1	-	-
2001	6	-	-	-	1	-	1	-	-	3	1	-	-
2000	6	-	-	-	-	-	1	-	1	2	2	-	-
1999	3	-	-	-	-	-	1	1	-	-	1	-	-
Origin not stated 3/, female													
2005	2	-	-	-	-	-	-	-	-	1	-	1	-
2004	3	-	-	-	-	-	-	-	-	-	2	1	-
2003	2	-	-	-	-	-	-	1	-	-	-	1	-
2002	7	-	-	-	-	-	-	-	-	1	4	2	-
2001	9	-	-	-	-	-	1	-	2	2	3	1	-
2000	3	-	-	-	-	-	-	1	1	-	-	1	-
1999	6	-	-	-	-	-	2	-	1	1	-	2	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Alcohol-induced causes (E24.4,F10,G31.2,G62.1,G72.1,I42.6,K29.2,K70,K86.0,R78.0,X45,X65,Y15) 5/													
All origins, both sexes													
2005	21,634	-	-	2	149	541	3,281	7,463	5,930	2,822	1,201	236	9
2004	21,081	-	-	6	139	610	3,359	7,215	5,442	2,892	1,200	211	7
2003	20,687	-	-	5	124	596	3,572	7,063	5,168	2,771	1,183	194	11
2002	20,218	1	2	3	105	601	3,636	6,748	4,888	2,835	1,190	201	8
2001	20,114	2	-	2	106	645	3,746	6,727	4,599	2,846	1,204	221	16
2000	19,643	3	1	4	95	650	3,837	6,143	4,542	2,910	1,223	227	8
1999	19,469	1	-	2	100	641	3,829	6,002	4,439	2,926	1,291	230	8
All origins, male													
2005	16,238	-	-	1	127	405	2,336	5,569	4,616	2,133	895	149	7
2004	15,906	-	-	5	107	468	2,434	5,401	4,224	2,232	894	135	6
2003	15,630	-	-	3	101	472	2,534	5,414	3,997	2,120	866	113	10
2002	15,272	-	1	2	86	458	2,571	5,171	3,837	2,158	845	135	8
2001	15,149	2	-	-	88	466	2,665	5,116	3,607	2,181	858	151	15
2000	14,993	2	-	4	76	504	2,808	4,792	3,489	2,249	915	146	8
1999	14,894	1	-	1	80	459	2,816	4,702	3,481	2,270	936	141	7
All origins, female													
2005	5,396	-	-	1	22	136	945	1,894	1,314	689	306	87	2
2004	5,175	-	-	1	32	142	925	1,814	1,218	660	306	76	1
2003	5,057	-	-	2	23	124	1,038	1,649	1,171	651	317	81	1
2002	4,946	1	1	1	19	143	1,065	1,577	1,051	677	345	66	-
2001	4,965	-	-	2	18	179	1,081	1,611	992	665	346	70	1
2000	4,650	1	1	-	19	146	1,029	1,351	1,053	661	308	81	-
1999	4,575	-	-	1	20	182	1,013	1,300	958	656	355	89	1

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Alcohol-induced causes (E24.4,F10,G31.2,G62.1,G72.1,I42.6,K29.2,K70,K86.0,R78.0,X45,X65,Y15) 5/													
Hispanic, both sexes													
2005	2,658	-	-	-	46	159	534	854	595	342	105	19	4
2004	2,406	-	-	3	41	144	498	829	501	263	117	8	2
2003	2,422	-	-	1	39	133	493	772	576	281	114	10	3
2002	2,408	-	-	1	33	147	506	774	521	319	87	19	1
2001	2,381	-	-	-	31	146	493	766	522	296	113	10	4
2000	2,323	-	-	1	32	136	531	698	504	311	96	11	3
1999	2,184	-	-	1	27	131	505	668	470	271	90	20	1
Hispanic, male													
2005	2,265	-	-	-	42	137	442	727	512	294	89	18	4
2004	2,056	-	-	2	38	132	425	728	416	213	93	7	2
2003	2,048	-	-	-	34	113	410	673	500	219	91	6	2
2002	2,065	-	-	-	30	134	431	661	454	276	65	13	1
2001	2,026	-	-	-	25	122	417	652	450	262	85	9	4
2000	2,024	-	-	1	29	129	445	606	440	278	86	7	3
1999	1,864	-	-	1	25	114	426	575	406	233	69	14	1
Hispanic, female													
2005	393	-	-	-	4	22	92	127	83	48	16	1	-
2004	350	-	-	1	3	12	73	101	85	50	24	1	-
2003	374	-	-	1	5	20	83	99	76	62	23	4	1
2002	343	-	-	1	3	13	75	113	67	43	22	6	-
2001	355	-	-	-	6	24	76	114	72	34	28	1	-
2000	299	-	-	-	3	7	86	92	64	33	10	4	-
1999	320	-	-	-	2	17	79	93	64	38	21	6	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Alcohol-induced causes (E24.4,F10,G31.2,G62.1,G72.1,I42.6,K29.2,K70,K86.0,R78.0,X45,X65,Y15) 5/													
Non-Hispanic 1/, both sexes													
2005	18,877	-	-	2	103	380	2,733	6,576	5,305	2,463	1,096	216	3
2004	18,567	-	-	3	98	464	2,850	6,336	4,913	2,616	1,081	203	3
2003	18,160	-	-	4	84	460	3,066	6,259	4,556	2,475	1,067	184	5
2002	17,661	1	2	2	71	450	3,102	5,921	4,330	2,500	1,097	181	4
2001	17,593	2	-	2	75	495	3,234	5,909	4,041	2,530	1,087	211	7
2000	17,177	3	1	3	62	509	3,280	5,389	4,012	2,579	1,120	215	4
1999	17,143	1	-	1	71	505	3,291	5,288	3,938	2,637	1,196	210	5
Non-Hispanic 1/, male													
2005	13,890	-	-	1	85	267	1,884	4,815	4,075	1,824	806	131	2
2004	13,761	-	-	3	69	334	2,000	4,632	3,784	2,010	799	128	2
2003	13,490	-	-	3	66	356	2,114	4,714	3,463	1,889	773	107	5
2002	13,078	-	1	2	55	320	2,120	4,463	3,351	1,866	775	121	4
2001	13,009	2	-	-	63	341	2,231	4,422	3,128	1,904	770	142	6
2000	12,843	2	-	3	46	371	2,338	4,137	3,027	1,955	822	138	4
1999	12,905	1	-	-	53	340	2,363	4,087	3,047	2,020	862	127	5
Non-Hispanic 1/, female													
2005	4,987	-	-	1	18	113	849	1,761	1,230	639	290	85	1
2004	4,806	-	-	-	29	130	850	1,704	1,129	606	282	75	1
2003	4,670	-	-	1	18	104	952	1,545	1,093	586	294	77	-
2002	4,583	1	1	-	16	130	982	1,458	979	634	322	60	-
2001	4,584	-	-	2	12	154	1,003	1,487	913	626	317	69	1
2000	4,334	1	1	-	16	138	942	1,252	985	624	298	77	-
1999	4,238	-	-	1	18	165	928	1,201	891	617	334	83	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Alcohol-induced causes (E24.4,F10,G31.2,G62.1,G72.1,I42.6,K29.2,K70,K86.0,R78.0,X45,X65,Y15) 5/													
Non-Hispanic white, both sexes													
2005 2/	15,729	-	-	1	73	276	2,206	5,433	4,477	2,100	967	194	2
2004 2/	15,418	-	-	1	70	337	2,323	5,267	4,068	2,190	972	188	2
2003 2/	14,977	-	-	3	57	347	2,465	5,105	3,746	2,124	959	167	4
2002	14,494	-	1	1	53	327	2,505	4,834	3,518	2,119	972	161	3
2001	14,186	1	-	2	47	360	2,533	4,691	3,267	2,131	967	182	5
2000	13,815	3	-	2	44	365	2,555	4,259	3,219	2,182	992	192	2
1999	13,633	1	-	1	49	380	2,507	4,142	3,129	2,200	1,038	181	5
Non-Hispanic white, male													
2005 2/	11,610	-	-	1	64	194	1,542	3,983	3,451	1,547	708	119	1
2004 2/	11,428	-	-	1	51	242	1,633	3,860	3,133	1,668	719	120	1
2003 2/	11,133	-	-	3	48	267	1,718	3,837	2,845	1,628	687	96	4
2002	10,783	-	1	1	43	236	1,730	3,678	2,736	1,564	687	104	3
2001	10,497	1	-	-	41	254	1,740	3,553	2,512	1,585	685	122	4
2000	10,408	2	-	2	34	269	1,840	3,317	2,444	1,648	729	121	2
1999	10,337	1	-	-	37	261	1,830	3,243	2,431	1,669	751	109	5
Non-Hispanic white, female													
2005 2/	4,119	-	-	-	9	82	664	1,450	1,026	553	259	75	1
2004 2/	3,990	-	-	-	19	95	690	1,407	935	522	253	68	1
2003 2/	3,844	-	-	-	9	80	747	1,268	901	496	272	71	-
2002	3,711	-	-	-	10	91	775	1,156	782	555	285	57	-
2001	3,689	-	-	2	6	106	793	1,138	755	546	282	60	1
2000	3,407	1	-	-	10	96	715	942	775	534	263	71	-
1999	3,296	-	-	1	12	119	677	899	698	531	287	72	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Alcohol-induced causes (E24.4,F10,G31.2,G62.1,G72.1,I42.6,K29.2,K70,K86.0,R78.0,X45,X65,Y15) 5/													
Non-Hispanic black, both sexes													
2005 2/	2,282	-	-	-	16	46	315	848	652	286	98	20	1
2004 2/	2,318	-	-	-	13	52	319	830	674	343	76	10	1
2003 2/	2,367	-	-	1	7	60	388	867	657	285	87	14	1
2002	2,396	1	1	-	10	70	384	853	653	303	104	16	1
2001	2,677	1	-	-	17	70	496	986	645	337	104	19	2
2000	2,672	-	1	1	8	92	516	946	639	343	104	20	2
1999	2,794	-	-	-	10	72	559	933	687	377	132	24	-
Non-Hispanic black, male													
2005 2/	1,670	-	-	-	11	31	201	626	493	222	75	10	1
2004 2/	1,754	-	-	-	8	42	231	605	529	281	52	5	1
2003 2/	1,787	-	-	-	5	48	258	677	507	214	67	10	1
2002	1,768	-	-	-	8	52	252	623	499	246	74	13	1
2001	2,016	1	-	-	13	47	355	725	515	273	74	11	2
2000	1,959	-	-	1	7	63	356	693	474	269	79	15	2
1999	2,066	-	-	-	8	45	380	690	533	304	92	14	-
Non-Hispanic black, female													
2005 2/	612	-	-	-	5	15	114	222	159	64	23	10	-
2004 2/	564	-	-	-	5	10	88	225	145	62	24	5	-
2003 2/	580	-	-	1	2	12	130	190	150	71	20	4	-
2002	628	1	1	-	2	18	132	230	154	57	30	3	-
2001	661	-	-	-	4	23	141	261	130	64	30	8	-
2000	713	-	1	-	1	29	160	253	165	74	25	5	-
1999	728	-	-	-	2	27	179	243	154	73	40	10	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Alcohol-induced causes (E24.4,F10,G31.2,G62.1,G72.1,I42.6,K29.2,K70,K86.0,R78.0,X45,X65,Y15) 5/													
Origin not stated 3/, both sexes													
2005	99	-	-	-	-	2	14	33	30	17	-	1	2
2004	108	-	-	-	-	2	11	50	28	13	2	-	2
2003	105	-	-	-	1	3	13	32	36	15	2	-	3
2002	149	-	-	-	1	4	28	53	37	16	6	1	3
2001	140	-	-	-	-	4	19	52	36	20	4	-	5
2000	143	-	-	-	1	5	26	56	26	20	7	1	1
1999	142	-	-	-	2	5	33	46	31	18	5	-	2
Origin not stated 3/, male													
2005	83	-	-	-	-	1	10	27	29	15	-	-	1
2004	89	-	-	-	-	2	9	41	24	9	2	-	2
2003	92	-	-	-	1	3	10	27	34	12	2	-	3
2002	129	-	-	-	1	4	20	47	32	16	5	1	3
2001	114	-	-	-	-	3	17	42	29	15	3	-	5
2000	126	-	-	-	1	4	25	49	22	16	7	1	1
1999	125	-	-	-	2	5	27	40	28	17	5	-	1
Origin not stated 3/, female													
2005	16	-	-	-	-	1	4	6	1	2	-	1	1
2004	19	-	-	-	-	-	2	9	4	4	-	-	-
2003	13	-	-	-	-	-	3	5	2	3	-	-	-
2002	20	-	-	-	-	-	8	6	5	-	1	-	-
2001	26	-	-	-	-	1	2	10	7	5	1	-	-
2000	17	-	-	-	-	1	1	7	4	4	-	-	-
1999	17	-	-	-	-	-	6	6	3	1	-	-	1

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Drug-induced causes (D52.1,D59.0,D59.2,D61.1,D64.2,E06.4,E16.0,E23.1,E24.2,E27.3,E66.1,F11.0-F11.5,F11.7-F11.9,F12.0-F12.5,F12.7-F12.9,F13.0-F13.5,F13.7-F13.9,F14.0-F14.5,F14.7-F14.9,F15.0-F15.5,F15.7-F15.9,F16.0-F16.5,F16.7- F16.9,F17.0,F17.3-F17.5,F17.7-F17.9,F18.0-F18.5,F18.7-F18.9,F19.0-F19.5,F19.7-F19.9,G21.1,G24.0,G25.1,G25.4,G25.6,G44.4,G62.0,G72.0,I95.2,J70.2-J70.4,L10.5,L27.0-L27.1,M10.2,M32.0,M80.4,M81.4,M83.5,M87.1,R78.1-R78.5,X40-X44,X60-X64,X85,Y10-Y14) 5/													
All origins, both sexes													
2005	33,541	35	30	51	3,095	5,730	9,333	10,041	3,240	887	706	388	5
2004	30,711	30	24	63	2,913	5,061	9,228	9,021	2,651	785	619	304	12
2003	28,723	25	35	41	2,619	4,845	9,158	8,174	2,239	755	537	283	12
2002	26,040	29	30	42	2,215	4,449	8,858	7,182	1,828	653	484	260	10
2001	21,705	22	26	45	1,813	3,739	7,669	5,796	1,361	553	439	226	16
2000	19,720	18	15	44	1,567	3,527	7,206	4,986	1,187	486	430	240	14
1999	19,128	21	25	30	1,351	3,579	7,095	4,624	1,161	556	467	201	18
All origins, male													
2005	21,208	18	23	24	2,319	3,970	5,829	6,143	1,912	452	354	159	5
2004	19,362	11	16	25	2,187	3,554	5,737	5,528	1,496	393	278	130	7
2003	18,426	17	21	20	1,955	3,361	5,873	5,153	1,266	384	253	112	11
2002	16,734	20	9	24	1,622	3,139	5,728	4,517	1,040	311	215	102	7
2001	14,253	14	21	22	1,350	2,640	5,079	3,793	760	293	191	74	16
2000	13,137	12	13	25	1,166	2,515	4,890	3,358	631	235	187	92	13
1999	12,885	14	12	16	981	2,579	4,907	3,133	688	286	184	74	11
All origins, female													
2005	12,333	17	7	27	776	1,760	3,504	3,898	1,328	435	352	229	-
2004	11,349	19	8	38	726	1,507	3,491	3,493	1,155	392	341	174	5
2003	10,297	8	14	21	664	1,484	3,285	3,021	973	371	284	171	1
2002	9,306	9	21	18	593	1,310	3,130	2,665	788	342	269	158	3
2001	7,452	8	5	23	463	1,099	2,590	2,003	601	260	248	152	-
2000	6,583	6	2	19	401	1,012	2,316	1,628	556	251	243	148	1
1999	6,243	7	13	14	370	1,000	2,188	1,491	473	270	283	127	7

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Drug-induced causes (D52.1,D59.0,D59.2,D61.1,D64.2,E06.4,E16.0,E23.1,E24.2,E27.3,E66.1,F11.0-F11.5,F11.7-F11.9,F12.0-F12.5,F12.7-F12.9,F13.0-F13.5,F13.7-F13.9,F14.0-F14.5,F14.7-F14.9,F15.0-F15.5,F15.7-F15.9,F16.0-F16.5,F16.7- F16.9,F17.0,F17.3-F17.5,F17.7-F17.9,F18.0-F18.5,F18.7-F18.9,F19.0-F19.5,F19.7-F19.9,G21.1,G24.0,G25.1,G25.4,G25.6,G44.4,G62.0,G72.0,I95.2,J70.2-J70.4,L10.5,L27.0-L27.1,M10.2,M32.0,M80.4,M81.4,M83.5,M87.1,R78.1-R78.5,X40-X44,X60-X64,X85,Y10-Y14) 5/													
Hispanic, both sexes													
2005	2,596	5	4	5	284	578	800	644	186	49	29	12	-
2004	2,257	-	1	7	233	482	746	527	185	39	24	12	1
2003	2,358	4	5	7	243	533	789	570	135	38	24	10	-
2002	2,137	5	4	4	217	456	777	505	111	36	20	2	-
2001	1,731	-	4	5	162	408	644	373	88	29	12	4	2
2000	1,700	2	2	4	148	449	651	332	79	20	8	5	-
1999	1,965	7	4	3	172	514	741	398	84	28	7	2	5
Hispanic, male													
2005	1,969	1	3	2	220	462	606	478	148	29	14	6	-
2004	1,671	-	1	3	175	386	556	370	132	27	13	7	1
2003	1,800	3	3	1	194	421	606	422	104	28	13	5	-
2002	1,647	4	-	1	172	371	601	385	81	18	13	1	-
2001	1,335	-	3	1	119	320	504	291	69	18	6	2	2
2000	1,348	2	2	-	121	367	521	265	52	11	3	4	-
1999	1,605	4	4	2	142	431	606	320	69	21	1	1	4
Hispanic, female													
2005	627	4	1	3	64	116	194	166	38	20	15	6	-
2004	586	-	-	4	58	96	190	157	53	12	11	5	-
2003	558	1	2	6	49	112	183	148	31	10	11	5	-
2002	490	1	4	3	45	85	176	120	30	18	7	1	-
2001	396	-	1	4	43	88	140	82	19	11	6	2	-
2000	352	-	-	4	27	82	130	67	27	9	5	1	-
1999	360	3	-	1	30	83	135	78	15	7	6	1	1

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Drug-induced causes (D52.1,D59.0,D59.2,D61.1,D64.2,E06.4,E16.0,E23.1,E24.2,E27.3,E66.1,F11.0-F11.5,F11.7-F11.9,F12.0-F12.5,F12.7-F12.9,F13.0-F13.5,F13.7-F13.9,F14.0-F14.5,F14.7-F14.9,F15.0-F15.5,F15.7-F15.9,F16.0-F16.5,F16.7- F16.9,F17.0,F17.3-F17.5,F17.7-F17.9,F18.0-F18.5,F18.7-F18.9,F19.0-F19.5,F19.7-F19.9,G21.1,G24.0,G25.1,G25.4,G25.6,G44.4,G62.0,G72.0,I95.2,J70.2-J70.4,L10.5,L27.0-L27.1,M10.2,M32.0,M80.4,M81.4,M83.5,M87.1,R78.1-R78.5,X40-X44,X60-X64,X85,Y10-Y14) 5/													
Non-Hispanic 1/, both sexes													
2005	30,809	30	26	46	2,804	5,136	8,498	9,341	3,037	834	677	375	5
2004	28,339	29	22	56	2,674	4,568	8,447	8,451	2,458	744	592	292	6
2003	26,199	21	30	34	2,371	4,287	8,321	7,546	2,087	711	511	273	7
2002	23,756	24	26	38	1,989	3,973	8,029	6,639	1,705	608	461	257	7
2001	19,799	22	22	40	1,642	3,313	6,969	5,367	1,252	520	426	221	5
2000	17,835	16	12	39	1,410	3,053	6,509	4,578	1,091	462	422	235	8
1999	16,966	14	21	27	1,174	3,046	6,279	4,168	1,052	524	458	197	6
Non-Hispanic 1/, male													
2005	19,140	17	20	22	2,093	3,497	5,199	5,623	1,751	421	340	152	5
2004	17,605	11	14	22	2,008	3,160	5,154	5,124	1,358	365	264	123	2
2003	16,497	14	18	19	1,758	2,921	5,231	4,689	1,146	350	238	107	6
2002	14,978	16	9	23	1,445	2,751	5,089	4,107	948	285	200	100	5
2001	12,778	14	18	21	1,224	2,304	4,532	3,459	673	272	184	72	5
2000	11,656	10	10	24	1,038	2,133	4,334	3,041	565	222	184	88	7
1999	11,136	10	8	14	835	2,132	4,249	2,768	600	261	183	72	4
Non-Hispanic 1/, female													
2005	11,669	13	6	24	711	1,639	3,299	3,718	1,286	413	337	223	-
2004	10,734	18	8	34	666	1,408	3,293	3,327	1,100	379	328	169	4
2003	9,702	7	12	15	613	1,366	3,090	2,857	941	361	273	166	1
2002	8,778	8	17	15	544	1,222	2,940	2,532	757	323	261	157	2
2001	7,021	8	4	19	418	1,009	2,437	1,908	579	248	242	149	-
2000	6,179	6	2	15	372	920	2,175	1,537	526	240	238	147	1
1999	5,830	4	13	13	339	914	2,030	1,400	452	263	275	125	2

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Drug-induced causes (D52.1,D59.0,D59.2,D61.1,D64.2,E06.4,E16.0,E23.1,E24.2,E27.3,E66.1,F11.0-F11.5,F11.7-F11.9,F12.0-F12.5,F12.7-F12.9,F13.0-F13.5,F13.7-F13.9,F14.0-F14.5,F14.7-F14.9,F15.0-F15.5,F15.7-F15.9,F16.0-F16.5,F16.7- F16.9,F17.0,F17.3-F17.5,F17.7-F17.9,F18.0-F18.5,F18.7-F18.9,F19.0-F19.5,F19.7-F19.9,G21.1,G24.0,G25.1,G25.4,G25.6,G44.4,G62.0,G72.0,I95.2,J70.2-J70.4,L10.5,L27.0-L27.1,M10.2,M32.0,M80.4,M81.4,M83.5,M87.1,R78.1-R78.5,X40-X44,X60-X64,X85,Y10-Y14) 5/													
Non-Hispanic white, both sexes													
2005 2/	26,186	20	11	33	2,600	4,490	7,221	7,634	2,473	726	619	354	5
2004 2/	24,201	18	13	41	2,477	4,026	7,225	6,942	1,993	650	544	268	4
2003 2/	22,245	12	17	25	2,208	3,739	6,994	6,158	1,730	612	488	258	4
2002	19,949	9	14	32	1,819	3,429	6,722	5,328	1,385	542	428	236	5
2001	16,367	10	13	33	1,484	2,835	5,730	4,190	1,019	456	394	198	5
2000	14,585	9	5	33	1,266	2,554	5,264	3,532	900	410	399	209	4
1999	13,644	6	10	22	1,019	2,493	5,001	3,179	840	461	433	178	2
Non-Hispanic white, male													
2005 2/	16,170	11	8	19	1,960	3,083	4,445	4,501	1,337	347	308	146	5
2004 2/	14,952	7	6	20	1,871	2,808	4,419	4,155	1,020	296	237	112	1
2003 2/	13,959	11	11	16	1,651	2,578	4,414	3,767	900	286	225	97	3
2002	12,478	7	3	19	1,325	2,393	4,276	3,230	708	241	179	94	3
2001	10,465	4	12	18	1,112	1,992	3,750	2,614	498	232	164	64	5
2000	9,439	5	3	21	944	1,790	3,525	2,284	428	185	175	76	3
1999	8,831	5	4	12	727	1,750	3,401	2,040	431	223	169	68	1
Non-Hispanic white, female													
2005 2/	10,016	9	3	14	640	1,407	2,776	3,133	1,136	379	311	208	-
2004 2/	9,249	11	7	21	606	1,218	2,806	2,787	973	354	307	156	3
2003 2/	8,286	1	6	9	557	1,161	2,580	2,391	830	326	263	161	1
2002	7,471	2	11	13	494	1,036	2,446	2,098	677	301	249	142	2
2001	5,902	6	1	15	372	843	1,980	1,576	521	224	230	134	-
2000	5,146	4	2	12	322	764	1,739	1,248	472	225	224	133	1
1999	4,813	1	6	10	292	743	1,600	1,139	409	238	264	110	1

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Drug-induced causes (D52.1,D59.0,D59.2,D61.1,D64.2,E06.4,E16.0,E23.1,E24.2,E27.3,E66.1,F11.0-F11.5,F11.7-F11.9,F12.0-F12.5,F12.7-F12.9,F13.0-F13.5,F13.7-F13.9,F14.0-F14.5,F14.7-F14.9,F15.0-F15.5,F15.7-F15.9,F16.0-F16.5,F16.7- F16.9,F17.0,F17.3-F17.5,F17.7-F17.9,F18.0-F18.5,F18.7-F18.9,F19.0-F19.5,F19.7-F19.9,G21.1,G24.0,G25.1,G25.4,G25.6,G44.4,G62.0,G72.0,I95.2,J70.2-J70.4,L10.5,L27.0-L27.1,M10.2,M32.0,M80.4,M81.4,M83.5,M87.1,R78.1-R78.5,X40-X44,X60-X64,X85,Y10-Y14) 5/													
Non-Hispanic black, both sexes													
2005 2/	4,019	10	14	10	136	515	1,118	1,544	526	89	42	15	-
2004 2/	3,577	11	9	12	137	429	1,062	1,366	414	80	37	19	1
2003 2/	3,466	9	13	8	112	431	1,181	1,278	315	82	21	13	3
2002	3,404	13	10	5	127	451	1,181	1,229	301	48	25	13	1
2001	3,099	11	8	5	112	404	1,140	1,108	214	52	25	20	-
2000	2,977	6	5	4	113	440	1,143	990	178	48	22	24	4
1999	3,030	8	11	4	123	488	1,170	940	190	55	20	17	4
Non-Hispanic black, male													
2005 2/	2,612	6	11	3	92	335	658	1,024	392	64	23	4	-
2004 2/	2,309	4	8	2	96	286	633	877	309	61	23	9	1
2003 2/	2,256	3	7	3	77	270	736	851	227	58	12	9	3
2002	2,264	8	6	3	90	300	740	833	231	34	16	2	1
2001	2,113	9	5	2	85	269	721	802	163	35	15	7	-
2000	2,050	4	5	1	76	303	749	723	131	34	9	11	4
1999	2,134	5	4	2	93	341	786	695	155	35	11	4	3
Non-Hispanic black, female													
2005 2/	1,407	4	3	7	44	180	460	520	134	25	19	11	-
2004 2/	1,268	7	1	10	41	143	429	489	105	19	14	10	-
2003 2/	1,210	6	6	5	35	161	445	427	88	24	9	4	-
2002	1,140	5	4	2	37	151	441	396	70	14	9	11	-
2001	986	2	3	3	27	135	419	306	51	17	10	13	-
2000	927	2	-	3	37	137	394	267	47	14	13	13	-
1999	896	3	7	2	30	147	384	245	35	20	9	13	1

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Drug-induced causes (D52.1,D59.0,D59.2,D61.1,D64.2,E06.4,E16.0,E23.1,E24.2,E27.3,E66.1,F11.0-F11.5,F11.7-F11.9,F12.0-F12.5,F12.7-F12.9,F13.0-F13.5,F13.7-F13.9,F14.0-F14.5,F14.7-F14.9,F15.0-F15.5,F15.7-F15.9,F16.0-F16.5,F16.7- F16.9,F17.0,F17.3-F17.5,F17.7-F17.9,F18.0-F18.5,F18.7-F18.9,F19.0-F19.5,F19.7-F19.9,G21.1,G24.0,G25.1,G25.4,G25.6,G44.4,G62.0,G72.0,I95.2,J70.2-J70.4,L10.5,L27.0-L27.1,M10.2,M32.0,M80.4,M81.4,M83.5,M87.1,R78.1-R78.5,X40-X44,X60-X64,X85,Y10-Y14) 5/													
Origin not stated 3/, both sexes													
2005	136	-	-	-	7	16	35	56	17	4	-	1	-
2004	115	1	1	-	6	11	35	43	8	2	3	-	5
2003	166	-	-	-	5	25	48	58	17	6	2	-	5
2002	147	-	-	-	9	20	52	38	12	9	3	1	3
2001	175	-	-	-	9	18	56	56	21	4	1	1	9
2000	185	-	1	1	9	25	46	76	17	4	-	-	6
1999	197	-	-	-	5	19	75	58	25	4	2	2	7
Origin not stated 3/, male													
2005	99	-	-	-	6	11	24	42	13	2	-	1	-
2004	86	-	1	-	4	8	27	34	6	1	1	-	4
2003	129	-	-	-	3	19	36	42	16	6	2	-	5
2002	109	-	-	-	5	17	38	25	11	8	2	1	2
2001	140	-	-	-	7	16	43	43	18	3	1	-	9
2000	133	-	1	1	7	15	35	52	14	2	-	-	6
1999	144	-	-	-	4	16	52	45	19	4	-	1	3
Origin not stated 3/, female													
2005	37	-	-	-	1	5	11	14	4	2	-	-	-
2004	29	1	-	-	2	3	8	9	2	1	2	-	1
2003	37	-	-	-	2	6	12	16	1	-	-	-	-
2002	38	-	-	-	4	3	14	13	1	1	1	-	1
2001	35	-	-	-	2	2	13	13	3	1	-	1	-
2000	52	-	-	-	2	10	11	24	3	2	-	-	-
1999	53	-	-	-	1	3	23	13	6	-	2	1	4

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Injury by firearms (*U01.4,W32-W34,X72-X74,X93-X95,Y22-Y24,Y35.0) 5/													
All origins, both sexes													
2005	30,694	7	62	335	6,821	6,302	5,101	4,744	2,987	1,929	1,784	612	10
2004	29,569	7	51	300	6,540	6,021	4,986	4,588	2,859	1,885	1,729	576	27
2003	30,136	8	48	324	6,846	6,188	5,118	4,550	2,804	1,912	1,731	589	18
2002	30,242	10	61	348	6,780	6,140	5,446	4,321	2,719	1,993	1,833	576	15
2001	29,573	11	70	333	6,687	6,137	5,288	4,120	2,544	1,998	1,803	563	19
2000	28,663	13	46	377	6,575	5,789	5,358	3,951	2,272	1,941	1,723	600	18
1999	28,874	8	65	416	6,795	5,975	5,234	3,722	2,306	2,034	1,738	561	20
All origins, male													
2005	26,657	7	42	252	6,204	5,559	4,221	3,951	2,544	1,679	1,624	567	7
2004	25,498	2	29	231	5,895	5,302	4,130	3,733	2,391	1,663	1,554	542	26
2003	26,124	7	28	256	6,186	5,472	4,222	3,761	2,378	1,694	1,558	546	16
2002	26,098	4	38	242	6,096	5,347	4,503	3,579	2,296	1,768	1,671	540	14
2001	25,480	7	39	249	6,066	5,347	4,312	3,416	2,116	1,754	1,634	522	18
2000	24,582	9	28	302	5,906	5,003	4,363	3,258	1,896	1,684	1,571	549	13
1999	24,700	5	36	308	6,056	5,156	4,229	3,057	1,951	1,786	1,580	517	19
All origins, female													
2005	4,037	-	20	83	617	743	880	793	443	250	160	45	3
2004	4,071	5	22	69	645	719	856	855	468	222	175	34	1
2003	4,012	1	20	68	660	716	896	789	426	218	173	43	2
2002	4,144	6	23	106	684	793	943	742	423	225	162	36	1
2001	4,093	4	31	84	621	790	976	704	428	244	169	41	1
2000	4,081	4	18	75	669	786	995	693	376	257	152	51	5
1999	4,174	3	29	108	739	819	1,005	665	355	248	158	44	1

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Injury by firearms (*U01.4,W32-W34,X72-X74,X93-X95,Y22-Y24,Y35.0) 5/													
Hispanic, both sexes													
2005	3,469	2	7	47	1,350	1,084	529	259	82	55	39	12	3
2004	3,278	3	7	46	1,315	970	475	248	106	56	37	11	4
2003	3,319	1	9	52	1,346	985	478	259	86	58	30	11	4
2002	3,143	2	15	56	1,260	920	459	220	99	62	33	13	4
2001	3,087	4	13	40	1,233	926	468	215	82	60	31	10	5
2000	2,891	1	7	55	1,184	788	414	255	83	64	28	9	3
1999	2,878	2	6	69	1,189	770	427	218	87	51	44	12	3
Hispanic, male													
2005	3,144	2	5	37	1,263	985	454	224	76	48	36	11	3
2004	2,973	1	4	33	1,228	888	418	214	91	50	33	9	4
2003	2,998	1	5	38	1,234	914	411	220	80	54	28	10	3
2002	2,834	-	9	38	1,173	844	390	185	89	57	32	13	4
2001	2,774	3	6	31	1,132	841	408	180	75	55	28	10	5
2000	2,582	-	4	46	1,096	711	351	210	73	54	26	9	2
1999	2,549	2	4	50	1,078	699	364	173	76	46	42	12	3
Hispanic, female													
2005	325	-	2	10	87	99	75	35	6	7	3	1	-
2004	305	2	3	13	87	82	57	34	15	6	4	2	-
2003	321	-	4	14	112	71	67	39	6	4	2	1	1
2002	309	2	6	18	87	76	69	35	10	5	1	-	-
2001	313	1	7	9	101	85	60	35	7	5	3	-	-
2000	309	1	3	9	88	77	63	45	10	10	2	-	1
1999	329	-	2	19	111	71	63	45	11	5	2	-	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Injury by firearms (*U01.4,W32-W34,X72-X74,X93-X95,Y22-Y24,Y35.0) 5/													
Non-Hispanic 1/, both sexes													
2005	27,103	5	55	287	5,447	5,194	4,547	4,467	2,890	1,872	1,739	597	3
2004	26,189	4	44	253	5,204	5,032	4,490	4,331	2,746	1,825	1,690	564	6
2003	26,710	7	39	272	5,468	5,183	4,617	4,281	2,709	1,849	1,700	576	9
2002	26,944	8	46	290	5,476	5,187	4,963	4,080	2,609	1,922	1,795	561	7
2001	26,341	7	57	292	5,409	5,182	4,802	3,895	2,446	1,927	1,767	549	8
2000	25,637	12	39	320	5,355	4,980	4,915	3,679	2,179	1,869	1,694	589	6
1999	25,877	6	59	347	5,587	5,174	4,782	3,491	2,210	1,978	1,689	547	7
Non-Hispanic 1/, male													
2005	23,406	5	37	215	4,921	4,552	3,745	3,712	2,453	1,629	1,582	553	2
2004	22,436	1	25	197	4,650	4,397	3,693	3,512	2,295	1,609	1,519	532	6
2003	23,036	6	23	218	4,925	4,540	3,794	3,533	2,289	1,637	1,529	534	8
2002	23,127	4	29	202	4,882	4,477	4,093	3,375	2,198	1,702	1,634	525	6
2001	22,573	4	33	217	4,894	4,479	3,889	3,228	2,025	1,688	1,601	508	7
2000	21,881	9	24	255	4,777	4,274	3,984	3,033	1,813	1,624	1,544	539	5
1999	22,050	3	32	258	4,964	4,432	3,842	2,874	1,866	1,735	1,533	504	7
Non-Hispanic 1/, female													
2005	3,697	-	18	72	526	642	802	755	437	243	157	44	1
2004	3,753	3	19	56	554	635	797	819	451	216	171	32	-
2003	3,674	1	16	54	543	643	823	748	420	212	171	42	1
2002	3,817	4	17	88	594	710	870	705	411	220	161	36	1
2001	3,768	3	24	75	515	703	913	667	421	239	166	41	1
2000	3,756	3	15	65	578	706	931	646	366	245	150	50	1
1999	3,827	3	27	89	623	742	940	617	344	243	156	43	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Injury by firearms (*U01.4,W32-W34,X72-X74,X93-X95,Y22-Y24,Y35.0) 5/													
Non-Hispanic white, both sexes													
2005 2/.....	18,521	3	26	164	2,147	2,487	3,248	3,792	2,657	1,741	1,673	580	3
2004 2/.....	18,200	1	13	138	2,231	2,480	3,302	3,691	2,480	1,694	1,616	549	5
2003 2/.....	18,457	1	15	141	2,289	2,552	3,404	3,645	2,483	1,731	1,627	563	6
2002.....	18,762	2	28	163	2,356	2,639	3,662	3,457	2,380	1,812	1,714	544	5
2001.....	18,676	4	30	203	2,373	2,847	3,601	3,361	2,233	1,797	1,685	536	6
2000.....	18,042	5	21	219	2,377	2,702	3,694	3,106	1,965	1,738	1,637	574	4
1999.....	18,260	2	27	225	2,504	2,949	3,624	2,951	1,976	1,852	1,614	533	3
Non-Hispanic white, male													
2005 2/.....	15,672	3	17	128	1,868	2,066	2,614	3,137	2,256	1,524	1,521	536	2
2004 2/.....	15,283	-	8	108	1,912	2,076	2,657	2,984	2,074	1,485	1,455	519	5
2003 2/.....	15,670	-	7	119	2,009	2,141	2,754	2,997	2,101	1,541	1,471	524	6
2002.....	15,881	2	16	114	2,038	2,169	2,998	2,855	2,005	1,608	1,562	510	4
2001.....	15,760	2	19	152	2,059	2,383	2,910	2,788	1,838	1,577	1,531	496	5
2000.....	15,160	2	11	175	2,050	2,232	2,965	2,560	1,629	1,508	1,498	527	3
1999.....	15,384	2	15	168	2,165	2,453	2,893	2,440	1,659	1,626	1,468	492	3
Non-Hispanic white, female													
2005 2/.....	2,849	-	9	36	279	421	634	655	401	217	152	44	1
2004 2/.....	2,917	1	5	30	319	404	645	707	406	209	161	30	-
2003 2/.....	2,787	1	8	22	280	411	650	648	382	190	156	39	-
2002.....	2,881	-	12	49	318	470	664	602	375	204	152	34	1
2001.....	2,916	2	11	51	314	464	691	573	395	220	154	40	1
2000.....	2,882	3	10	44	327	470	729	546	336	230	139	47	1
1999.....	2,876	-	12	57	339	496	731	511	317	226	146	41	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Injury by firearms (*U01.4,W32-W34,X72-X74,X93-X95,Y22-Y24,Y35.0) 5/													
Non-Hispanic black, both sexes													
2005 2/	7,865	2	27	107	3,058	2,538	1,178	582	186	114	56	17	-
2004 2/	7,347	3	27	101	2,803	2,394	1,063	540	227	108	66	14	1
2003 2/	7,566	6	23	116	2,968	2,460	1,088	544	186	103	58	12	2
2002	7,494	6	17	102	2,904	2,393	1,172	535	186	95	65	17	2
2001	7,063	2	23	78	2,872	2,172	1,092	464	171	108	69	11	1
2000	6,958	7	13	91	2,798	2,121	1,093	496	162	114	50	12	1
1999	6,933	2	26	105	2,882	2,034	1,037	457	195	109	68	14	4
Non-Hispanic black, male													
2005 2/	7,122	2	19	78	2,846	2,337	1,027	496	156	93	51	17	-
2004 2/	6,620	1	15	78	2,588	2,191	933	453	187	104	57	12	1
2003 2/	6,794	6	15	86	2,739	2,253	935	464	156	83	47	9	1
2002	6,681	2	12	74	2,657	2,182	991	450	156	82	58	15	2
2001	6,323	1	13	60	2,690	1,965	894	387	150	93	59	10	1
2000	6,193	7	8	74	2,574	1,910	920	411	136	103	40	9	1
1999	6,114	1	12	78	2,623	1,826	854	376	174	95	59	12	4
Non-Hispanic black, female													
2005 2/	743	-	8	29	212	201	151	86	30	21	5	-	-
2004 2/	727	2	12	23	215	203	130	87	40	4	9	2	-
2003 2/	772	-	8	30	229	207	153	80	30	20	11	3	1
2002	813	4	5	28	247	211	181	85	30	13	7	2	-
2001	740	1	10	18	182	207	198	77	21	15	10	1	-
2000	765	-	5	17	224	211	173	85	26	11	10	3	-
1999	819	1	14	27	259	208	183	81	21	14	9	2	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Injury by firearms (*U01.4,W32-W34,X72-X74,X93-X95,Y22-Y24,Y35.0) 5/													
Origin not stated 3/, both sexes													
2005	122	-	-	1	24	24	25	18	15	2	6	3	4
2004	102	-	-	1	21	19	21	9	7	4	2	1	17
2003	107	-	-	-	32	20	23	10	9	5	1	2	5
2002	155	-	-	2	44	33	24	21	11	9	5	2	4
2001	145	-	-	1	45	29	18	10	16	11	5	4	6
2000	135	-	-	2	36	21	29	17	10	8	1	2	9
1999	119	-	-	-	19	31	25	13	9	5	5	2	10
Origin not stated 3/, male													
2005	107	-	-	-	20	22	22	15	15	2	6	3	2
2004	89	-	-	1	17	17	19	7	5	4	2	1	16
2003	90	-	-	-	27	18	17	8	9	3	1	2	5
2002	137	-	-	2	41	26	20	19	9	9	5	2	4
2001	133	-	-	1	40	27	15	8	16	11	5	4	6
2000	119	-	-	1	33	18	28	15	10	6	1	1	6
1999	101	-	-	-	14	25	23	10	9	5	5	1	9

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2005

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Injury by firearms (*U01.4,W32-W34,X72-X74,X93-X95,Y22-Y24,Y35.0) 5/													
Origin not stated 3/, female													
2005	15	-	-	1	4	2	3	3	-	-	-	-	2
2004	13	-	-	-	4	2	2	2	2	-	-	-	1
2003	17	-	-	-	5	2	6	2	-	2	-	-	-
2002	18	-	-	-	3	7	4	2	2	-	-	-	-
2001	12	-	-	-	5	2	3	2	-	-	-	-	-
2000	16	-	-	1	3	3	1	2	-	2	-	1	3
1999	18	-	-	-	5	6	2	3	-	-	-	1	1

- Quantity zero.
... Category not applicable.
1/ Includes races other than white and black.
2/ Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming in 2005, by California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and by California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005."
3/ Includes deaths for which Hispanic origin was not reported on the death certificate.
4/ Figures include September 11, 2001 related deaths for which death certificates were filed as of 10/24/02; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001."
5/ Included in selected categories.

Source: CDC/NCHS, National Vital Statistics System, Mortality.