

Worktable 250R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race, and sex: United States, 2007

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2007. Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Shigellosis and amebiasis (A03,A06)												
All races 3/, both sexes..	*	*	*	*	*	*	*	*	*	*	*	*
Male.....	*	*	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*	*	*
White, both sexes.....	*	*	*	*	*	*	*	*	*	*	*	*
Male.....	*	*	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*	*	*
Black, both sexes.....	*	*	*	*	*	*	*	*	*	*	*	*
Male.....	*	*	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*	*	*
Certain other intestinal infections (A04,A07-A09)												
All races 3/, both sexes..	2.2	*	*	*	*	*	0.1	0.3	1.0	4.7	18.8	52.0
Male.....	1.7	*	*	*	*	*	*	0.2	1.1	4.6	18.3	50.4
Female.....	2.8	*	*	*	*	*	0.1	0.3	0.9	4.9	19.1	52.7
White, both sexes.....	2.6	*	*	*	*	*	0.1	0.3	1.0	4.9	19.8	54.9
Male.....	1.9	*	*	*	*	*	*	0.3	1.1	4.7	19.1	53.5
Female.....	3.2	*	*	*	*	*	*	0.3	0.9	5.1	20.4	55.6
Black, both sexes.....	1.1	*	*	*	*	*	*	*	1.3	4.5	13.1	31.7
Male.....	0.8	*	*	*	*	*	*	*	1.5	5.0	14.4	28.0
Female.....	1.3	*	*	*	*	*	*	*	1.1	4.2	12.3	33.2

Worktable 250R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race, and sex: United States, 2007

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2007. Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Measles (B05)												
All races 3/, both sexes..	*	*	*	*	*	*	*	*	*	*	*	*
Male.....	*	*	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*	*	*
White, both sexes.....	*	*	*	*	*	*	*	*	*	*	*	*
Male.....	*	*	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*	*	*
Black, both sexes.....	*	*	*	*	*	*	*	*	*	*	*	*
Male.....	*	*	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*	*	*
Viral hepatitis (B15-B19)												
All races 3/, both sexes..	2.5	*	*	*	*	0.2	1.1	6.4	7.4	4.5	4.4	2.9
Male.....	3.3	*	*	*	*	0.2	1.5	9.2	11.2	5.2	4.7	3.0
Female.....	1.6	*	*	*	*	0.1	0.8	3.7	3.8	3.9	4.2	2.9
White, both sexes.....	2.4	*	*	*	*	0.2	1.1	6.5	6.8	4.1	4.0	2.7
Male.....	3.3	*	*	*	*	0.2	1.5	9.4	10.3	4.8	4.4	3.0
Female.....	1.6	*	*	*	*	*	0.7	3.6	3.5	3.5	3.7	2.6
Black, both sexes.....	2.6	*	*	*	*	*	1.1	7.2	12.1	5.8	5.4	*
Male.....	3.6	*	*	*	*	*	1.5	10.4	19.7	7.9	6.1	*
Female.....	1.7	*	*	*	*	*	0.8	4.5	6.1	4.3	5.0	*

Worktable 250R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race, and sex: United States, 2007

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2007. Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Other and unspecified infectious and parasitic diseases and their sequelae (A00,A05,A20-A36,A42-A44,A48-A49,A54-A79,A81-A82,A85.0-A85.1,A85.8,A86-B04,B06-B09,B25-B49,B55-B99)												
All races 3/, both sexes..	1.9	3.6	0.4	0.1	0.2	0.3	0.5	1.2	2.8	5.7	11.2	20.0
Male.....	2.0	3.9	0.4	0.2	0.3	0.4	0.6	1.4	3.2	6.6	13.5	25.3
Female.....	1.9	3.4	0.4	*	0.2	0.3	0.5	0.9	2.5	4.9	9.7	17.5
White, both sexes.....	2.0	3.3	0.4	0.1	0.2	0.3	0.4	1.0	2.7	5.6	11.2	20.5
Male.....	2.0	3.4	0.3	0.2	0.2	0.3	0.5	1.3	3.1	6.3	13.5	25.8
Female.....	1.9	3.1	0.4	*	0.2	0.3	0.3	0.8	2.3	5.0	9.6	18.0
Black, both sexes.....	1.9	5.7	*	*	0.4	0.7	1.2	2.1	4.2	6.7	12.6	17.2
Male.....	2.1	6.6	*	*	*	1.0	1.1	2.7	4.7	9.8	14.2	23.8
Female.....	1.8	*	*	*	*	*	1.2	1.7	3.9	4.6	11.7	14.5
Malignant neoplasms (C00-C97)												
All races 3/, both sexes..	186.6	1.7	2.2	2.4	3.9	8.5	30.8	114.3	315.4	715.5	1,256.3	1,590.2
Male.....	197.0	1.8	2.3	2.4	4.5	8.2	26.4	117.5	358.5	854.3	1,617.4	2,249.2
Female.....	176.5	1.6	2.2	2.3	3.2	8.9	35.2	111.3	275.2	597.6	1,007.4	1,276.7
White, both sexes.....	198.7	1.8	2.2	2.5	3.9	8.4	29.9	109.6	308.3	715.1	1,264.7	1,602.3
Male.....	208.8	1.8	2.2	2.5	4.6	8.1	25.9	112.0	346.7	845.4	1,617.4	2,253.2
Female.....	188.8	1.8	2.3	2.4	3.2	8.6	33.9	107.1	271.8	602.3	1,017.7	1,291.6
Black, both sexes.....	160.0	*	2.0	2.2	3.9	10.8	41.7	166.3	438.0	872.2	1,385.8	1,668.2
Male.....	172.9	*	2.4	2.2	4.2	9.5	34.0	178.0	544.1	1,139.5	1,936.9	2,637.1
Female.....	148.2	*	1.7	2.2	3.5	12.0	48.5	156.1	352.5	681.0	1,071.7	1,265.2

Worktable 250R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race, and sex: United States, 2007

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2007. Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Malignant neoplasms of lip, oral cavity and pharynx (C00-C14)												
All races 3/, both sexes..	2.7	*	*	*	*	0.1	0.6	2.6	6.2	9.9	13.4	17.1
Male.....	3.7	*	*	*	*	0.1	0.8	4.0	10.0	15.5	20.1	24.0
Female.....	1.7	*	*	*	*	0.1	0.4	1.3	2.6	5.2	8.7	13.8
White, both sexes.....	2.8	*	*	*	*	0.1	0.5	2.4	5.9	9.7	13.5	17.6
Male.....	3.8	*	*	*	*	*	0.7	3.7	9.6	14.7	20.0	24.4
Female.....	1.8	*	*	*	*	*	0.3	1.1	2.4	5.3	8.9	14.3
Black, both sexes.....	2.7	*	*	*	*	*	1.0	4.1	9.7	13.7	13.5	14.2
Male.....	4.0	*	*	*	*	*	1.4	6.5	16.3	25.8	23.5	23.8
Female.....	1.4	*	*	*	*	*	*	2.0	4.4	5.0	7.8	10.3
Malignant neoplasm of esophagus (C15)												
All races 3/, both sexes..	4.5	*	*	*	*	0.1	0.6	3.3	10.3	19.3	25.9	25.0
Male.....	7.2	*	*	*	*	0.1	1.0	5.8	18.2	33.8	47.8	48.4
Female.....	1.9	*	*	*	*	*	0.2	0.9	3.0	6.9	10.9	13.9
White, both sexes.....	4.9	*	*	*	*	0.1	0.6	3.3	10.3	19.7	26.5	25.8
Male.....	7.9	*	*	*	*	0.1	1.0	5.9	18.2	34.6	48.7	49.8
Female.....	1.9	*	*	*	*	*	0.2	0.8	2.8	6.7	10.9	14.3
Black, both sexes.....	3.6	*	*	*	*	*	0.6	4.2	13.9	21.3	25.4	18.7
Male.....	5.5	*	*	*	*	*	0.8	6.7	23.7	37.9	46.5	39.9
Female.....	1.9	*	*	*	*	*	*	2.0	5.9	9.4	13.4	9.9

Worktable 250R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race, and sex: United States, 2007

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2007. Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Malignant neoplasm of stomach (C16)												
All races 3/, both sexes..	3.8	*	*	*	*	0.3	1.1	2.6	5.7	13.2	25.0	35.4
Male.....	4.5	*	*	*	*	0.3	1.1	3.3	8.0	19.1	35.2	51.1
Female.....	3.0	*	*	*	*	0.3	1.0	2.0	3.7	8.2	17.9	27.9
White, both sexes.....	3.5	*	*	*	*	0.3	1.0	2.2	4.8	11.4	21.9	31.9
Male.....	4.2	*	*	*	*	0.3	1.1	2.7	6.7	16.7	30.5	46.4
Female.....	2.8	*	*	*	*	0.3	0.8	1.7	3.0	6.9	15.8	25.0
Black, both sexes.....	5.0	*	*	*	*	*	1.4	4.9	11.9	25.0	49.7	65.4
Male.....	6.2	*	*	*	*	*	1.6	7.2	17.6	38.4	78.3	96.0
Female.....	3.9	*	*	*	*	*	1.3	3.0	7.3	15.4	33.4	52.7
Malignant neoplasms of colon, rectum and anus (C18-C21)												
All races 3/, both sexes..	17.8	*	*	*	0.1	0.7	3.0	10.9	27.7	60.1	118.4	200.8
Male.....	18.2	*	*	*	0.1	0.7	3.3	12.3	33.8	74.5	144.0	223.5
Female.....	17.3	*	*	*	*	0.6	2.8	9.6	22.0	47.9	100.7	190.0
White, both sexes.....	18.5	*	*	*	0.1	0.6	2.9	10.2	25.9	58.3	117.0	200.9
Male.....	19.0	*	*	*	*	0.7	3.1	11.5	31.8	72.2	142.8	223.1
Female.....	18.1	*	*	*	*	0.6	2.7	8.9	20.3	46.3	99.0	190.3
Black, both sexes.....	17.3	*	*	*	*	0.9	4.2	17.9	46.8	86.9	154.4	225.4
Male.....	18.0	*	*	*	*	1.1	4.9	20.8	57.9	112.0	192.5	266.9
Female.....	16.6	*	*	*	*	0.8	3.6	15.4	37.9	69.0	132.7	208.2

Worktable 250R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race, and sex: United States, 2007

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2007. Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Malignant neoplasms of liver and intrahepatic bile ducts (C22)												
All races 3/, both sexes..	5.7	*	*	0.1	0.1	0.2	0.9	5.7	12.8	20.1	32.8	32.0
Male.....	7.6	*	*	*	*	0.3	1.2	9.1	20.5	28.7	46.0	43.3
Female.....	3.8	*	*	*	0.1	0.2	0.5	2.4	5.5	12.7	23.6	26.7
White, both sexes.....	5.6	*	*	0.1	0.1	0.2	0.7	5.1	11.1	18.8	31.7	31.1
Male.....	7.4	*	*	*	*	0.2	0.9	8.1	17.6	26.6	44.6	41.8
Female.....	3.8	*	*	*	*	0.1	0.5	2.2	5.0	12.1	22.6	26.0
Black, both sexes.....	5.7	*	*	*	*	0.5	1.4	8.4	22.8	24.3	33.7	34.2
Male.....	8.3	*	*	*	*	0.7	2.2	14.2	41.2	38.7	48.4	50.1
Female.....	3.2	*	*	*	*	*	*	3.4	8.1	14.1	25.3	27.6
Malignant neoplasm of pancreas (C25)												
All races 3/, both sexes..	11.3	*	*	*	*	0.1	1.2	6.4	19.9	44.8	79.2	94.6
Male.....	11.5	*	*	*	*	0.2	1.5	7.8	24.5	52.5	88.1	104.9
Female.....	11.1	*	*	*	*	0.1	1.0	5.0	15.6	38.3	73.1	89.8
White, both sexes.....	11.9	*	*	*	*	0.1	1.2	6.1	19.5	44.1	78.7	93.6
Male.....	12.2	*	*	*	*	0.1	1.5	7.5	24.0	52.3	88.4	105.4
Female.....	11.7	*	*	*	*	*	0.9	4.8	15.3	37.1	71.8	88.0
Black, both sexes.....	10.2	*	*	*	*	*	1.6	9.6	27.2	58.6	97.7	114.1
Male.....	9.9	*	*	*	*	*	1.9	12.0	36.0	65.7	99.2	107.1
Female.....	10.4	*	*	*	*	*	1.3	7.5	20.1	53.6	96.8	117.0

Worktable 250R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race, and sex: United States, 2007

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2007. Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Malignant neoplasm of larynx (C32)												
All races 3/, both sexes..	1.2	*	*	*	*	*	0.1	1.0	2.8	5.1	6.7	6.0
Male.....	1.9	*	*	*	*	*	0.2	1.7	4.9	8.6	12.9	13.9
Female.....	0.5	*	*	*	*	*	*	0.4	0.9	2.1	2.4	2.3
White, both sexes.....	1.2	*	*	*	*	*	0.1	0.9	2.6	4.8	6.4	5.9
Male.....	1.9	*	*	*	*	*	0.2	1.5	4.4	7.9	12.2	13.3
Female.....	0.5	*	*	*	*	*	*	0.4	0.8	2.1	2.4	2.3
Black, both sexes.....	1.6	*	*	*	*	*	*	2.1	6.1	9.7	11.4	8.2
Male.....	2.9	*	*	*	*	*	*	4.0	11.8	18.1	25.7	22.9
Female.....	0.5	*	*	*	*	*	*	*	1.4	3.7	3.2	*
Malignant neoplasms of trachea, bronchus and lung (C33-C34)												
All races 3/, both sexes..	52.6	*	*	*	0.1	0.3	4.3	28.4	95.4	248.8	371.3	299.8
Male.....	59.4	*	*	*	*	0.4	4.2	32.1	116.2	310.2	498.3	453.0
Female.....	46.0	*	*	*	*	0.3	4.4	24.9	76.1	196.7	283.8	227.0
White, both sexes.....	57.0	*	*	*	0.1	0.3	4.3	27.4	95.2	253.8	379.4	303.7
Male.....	63.4	*	*	*	*	0.4	4.2	30.2	113.1	310.4	502.7	453.3
Female.....	50.7	*	*	*	*	0.3	4.5	24.7	78.3	204.7	293.0	232.3
Black, both sexes.....	41.2	*	*	*	*	*	5.1	42.5	122.8	270.2	357.1	283.1
Male.....	51.5	*	*	*	*	*	5.5	54.1	177.7	395.6	546.2	504.8
Female.....	31.8	*	*	*	*	*	4.8	32.5	78.6	180.4	249.3	190.9

Worktable 250R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race, and sex: United States, 2007

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2007. Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Malignant melanoma of skin (C43)												
All races 3/, both sexes..	2.8	*	*	*	0.1	0.4	1.1	2.7	5.3	9.0	15.6	19.9
Male.....	3.7	*	*	*	0.1	0.5	1.2	3.5	7.7	13.5	24.8	35.8
Female.....	1.9	*	*	*	*	0.4	0.9	1.9	3.1	5.2	9.3	12.4
White, both sexes.....	3.4	*	*	*	0.1	0.5	1.3	3.2	6.2	10.2	17.3	21.6
Male.....	4.5	*	*	*	0.1	0.6	1.5	4.1	8.8	15.3	27.4	38.9
Female.....	2.3	*	*	*	*	0.5	1.1	2.2	3.6	5.8	10.1	13.4
Black, both sexes.....	0.3	*	*	*	*	*	*	*	0.8	1.2	3.2	5.7
Male.....	0.3	*	*	*	*	*	*	*	*	*	*	*
Female.....	0.4	*	*	*	*	*	*	*	*	*	3.2	*
Malignant neoplasm of breast (C50)												
All races 3/, both sexes..	13.6	*	*	*	*	0.8	5.1	13.7	26.8	42.3	69.7	116.6
Male.....	0.2	*	*	*	*	*	*	0.2	0.4	1.0	2.0	3.4
Female.....	26.5	*	*	*	*	1.7	10.1	26.7	51.3	77.3	116.3	170.4
White, both sexes.....	14.0	*	*	*	*	0.7	4.5	12.5	25.5	41.9	69.8	117.6
Male.....	0.3	*	*	*	*	*	*	0.2	0.4	1.0	1.9	3.4
Female.....	27.6	*	*	*	*	1.4	9.1	24.7	49.4	77.3	117.3	172.1
Black, both sexes.....	14.6	*	*	*	*	1.9	9.7	24.0	42.4	56.7	87.2	136.8
Male.....	0.3	*	*	*	*	*	*	*	*	*	*	*
Female.....	27.7	*	*	*	*	3.6	18.2	44.3	75.7	96.0	135.2	191.9

Worktable 250R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race, and sex: United States, 2007

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2007. Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Malignant neoplasm of cervix uteri (C53)												
All races 3/, both sexes..	1.3	*	*	*	*	0.5	1.5	2.1	2.7	3.2	3.9	4.7
Male.....
Female.....	2.6	*	*	*	*	0.9	3.0	4.1	5.3	5.8	6.6	7.0
White, both sexes.....	1.2	*	*	*	*	0.4	1.4	1.9	2.5	2.8	3.4	4.0
Male.....
Female.....	2.5	*	*	*	*	0.8	2.8	3.8	4.8	5.2	5.8	5.9
Black, both sexes.....	2.0	*	*	*	*	0.9	2.4	3.5	4.7	6.8	9.3	13.0
Male.....
Female.....	3.9	*	*	*	*	1.7	4.6	6.4	8.6	11.7	14.6	18.4
Malignant neoplasms of corpus uteri and uterus, part unspecified (C54-C55)												
All races 3/, both sexes..	2.5	*	*	*	*	0.1	0.4	1.3	4.8	10.4	15.0	20.1
Male.....
Female.....	4.9	*	*	*	*	0.2	0.8	2.6	9.3	19.3	25.4	29.6
White, both sexes.....	2.4	*	*	*	*	0.1	0.3	1.3	4.5	9.1	14.0	19.9
Male.....
Female.....	4.9	*	*	*	*	0.2	0.7	2.6	8.8	17.0	23.7	29.4
Black, both sexes.....	3.2	*	*	*	*	*	0.6	1.8	8.5	24.6	30.3	27.0
Male.....
Female.....	6.2	*	*	*	*	*	1.1	3.4	15.3	42.1	47.5	38.2

Worktable 250R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race, and sex: United States, 2007

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2007. Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Malignant neoplasms of kidney and renal pelvis (C64-C65)												
All races 3/, both sexes..	4.2	*	*	0.1	0.1	0.1	0.6	2.8	7.9	16.3	27.2	33.0
Male.....	5.4	*	*	*	0.1	0.2	0.7	4.1	11.5	23.0	39.2	50.9
Female.....	3.1	*	*	*	*	*	0.4	1.6	4.5	10.6	18.9	24.5
White, both sexes.....	4.6	*	*	*	*	0.1	0.5	2.8	8.1	16.8	27.7	33.4
Male.....	5.8	*	*	*	*	0.1	0.7	4.1	11.6	23.4	40.1	52.2
Female.....	3.4	*	*	*	*	*	0.4	1.6	4.7	11.1	19.0	24.5
Black, both sexes.....	3.1	*	*	*	*	*	0.8	3.4	8.5	15.0	26.9	30.2
Male.....	3.9	*	*	*	*	*	1.1	5.1	13.2	22.9	36.9	41.6
Female.....	2.3	*	*	*	*	*	*	1.8	4.7	9.3	21.2	25.5
Malignant neoplasm of bladder (C67)												
All races 3/, both sexes..	4.6	*	*	*	*	*	0.2	1.3	4.8	14.6	38.5	68.6
Male.....	6.5	*	*	*	*	*	0.3	1.9	7.5	23.7	66.6	132.7
Female.....	2.7	*	*	*	*	*	0.2	0.7	2.3	6.8	19.1	38.1
White, both sexes.....	5.2	*	*	*	*	*	0.2	1.3	5.0	15.4	40.1	71.2
Male.....	7.4	*	*	*	*	*	0.3	2.0	7.7	25.1	70.2	140.0
Female.....	3.0	*	*	*	*	*	0.1	0.7	2.3	7.0	19.0	38.4
Black, both sexes.....	2.6	*	*	*	*	*	*	1.6	4.9	12.0	32.0	53.9
Male.....	3.0	*	*	*	*	*	*	1.9	7.2	18.1	44.9	79.0
Female.....	2.2	*	*	*	*	*	*	1.3	3.0	7.5	24.7	43.5

Worktable 250R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race, and sex: United States, 2007

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2007. Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Malignant neoplasms of meninges, brain and other parts of central nervous system (C70-C72)												
All races 3/, both sexes..	4.4	*	0.7	0.8	0.5	0.9	2.0	4.4	9.1	15.5	20.0	16.1
Male.....	4.9	*	0.7	0.7	0.5	1.1	2.4	5.3	11.3	18.5	24.1	21.1
Female.....	3.9	*	0.6	0.8	0.4	0.7	1.5	3.4	7.0	13.0	17.2	13.8
White, both sexes.....	5.0	*	0.7	0.8	0.5	1.0	2.2	4.8	9.9	16.8	21.6	16.9
Male.....	5.6	*	0.8	0.8	0.6	1.2	2.7	5.8	12.2	19.9	26.0	22.5
Female.....	4.4	*	0.6	0.9	0.4	0.8	1.7	3.7	7.7	14.1	18.5	14.2
Black, both sexes.....	2.0	*	*	0.6	0.3	0.6	1.0	2.9	5.3	9.5	8.7	9.0
Male.....	2.1	*	*	*	*	0.8	1.3	3.5	6.9	11.3	9.4	*
Female.....	1.9	*	*	0.8	*	*	0.7	2.3	4.0	8.1	8.4	9.2
Malignant neoplasms of lymphoid, hematopoietic and related tissue (C81-C96)												
All races 3/, both sexes..	18.2	0.6	0.7	0.9	1.5	1.9	3.4	8.2	23.5	63.2	137.3	185.5
Male.....	20.3	*	0.7	1.0	1.7	2.2	4.1	10.2	29.4	80.7	182.2	253.4
Female.....	16.2	*	0.7	0.7	1.2	1.6	2.7	6.3	18.1	48.2	106.4	153.3
White, both sexes.....	19.9	0.6	0.7	0.9	1.5	1.9	3.2	7.9	23.2	64.2	141.6	192.7
Male.....	22.2	*	0.6	1.0	1.7	2.2	3.9	9.8	29.3	82.3	188.1	263.7
Female.....	17.6	*	0.7	0.8	1.2	1.5	2.5	6.0	17.5	48.5	109.0	158.9
Black, both sexes.....	13.1	*	0.7	0.8	1.5	2.3	5.2	11.7	30.6	68.3	116.0	138.6
Male.....	13.9	*	*	0.9	1.6	2.6	6.3	14.6	36.5	84.7	147.0	181.9
Female.....	12.4	*	*	0.6	1.3	2.1	4.2	9.2	25.8	56.6	98.3	120.5

Worktable 250R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race, and sex: United States, 2007

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2007. Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Hodgkin's disease (C81)												
All races 3/, both sexes..	0.4	*	*	*	0.2	0.3	0.3	0.3	0.5	1.1	2.2	2.4
Male.....	0.5	*	*	*	0.2	0.4	0.4	0.5	0.6	1.3	2.6	2.8
Female.....	0.4	*	*	*	0.1	0.2	0.2	0.2	0.5	0.9	1.9	2.2
White, both sexes.....	0.5	*	*	*	0.2	0.3	0.3	0.3	0.5	1.2	2.4	2.6
Male.....	0.5	*	*	*	0.2	0.4	0.4	0.5	0.6	1.4	2.8	2.9
Female.....	0.4	*	*	*	0.1	0.2	0.2	0.2	0.5	1.0	2.1	2.4
Black, both sexes.....	0.3	*	*	*	*	*	0.4	0.5	0.8	*	*	*
Male.....	0.4	*	*	*	*	*	*	0.9	*	*	*	*
Female.....	0.3	*	*	*	*	*	*	*	*	*	*	*
Non-Hodgkin's lymphoma (C82-C85)												
All races 3/, both sexes..	6.8	*	*	0.1	0.3	0.5	1.2	3.2	8.9	23.1	52.7	72.1
Male.....	7.4	*	*	0.1	0.4	0.6	1.6	4.1	11.3	29.6	65.7	91.6
Female.....	6.2	*	*	*	0.2	0.4	0.8	2.3	6.7	17.6	43.8	62.8
White, both sexes.....	7.6	*	*	0.1	0.3	0.5	1.1	3.1	9.1	24.4	55.7	76.6
Male.....	8.2	*	*	*	0.4	0.6	1.5	4.0	11.6	30.9	69.2	97.0
Female.....	7.0	*	*	*	0.2	0.3	0.8	2.2	6.8	18.8	46.3	66.8
Black, both sexes.....	3.5	*	*	*	0.4	0.9	1.9	3.9	9.1	16.4	25.6	31.2
Male.....	3.9	*	*	*	*	1.0	2.5	5.1	11.6	23.1	29.9	36.5
Female.....	3.1	*	*	*	*	0.7	1.2	2.9	7.1	11.6	23.2	29.0

Worktable 250R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race, and sex: United States, 2007

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2007. Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Leukemia (C91-C95)												
All races 3/, both sexes..	7.2	0.5	0.6	0.8	1.0	1.1	1.5	3.1	8.6	23.8	52.7	76.7
Male.....	8.3	*	0.6	0.9	1.1	1.2	1.7	3.6	10.9	31.6	75.7	109.4
Female.....	6.2	*	0.7	0.7	0.9	0.9	1.4	2.6	6.4	17.2	36.8	61.2
White, both sexes.....	8.0	*	0.6	0.8	1.0	1.1	1.5	3.1	8.7	24.6	55.1	80.2
Male.....	9.2	*	0.6	0.9	1.2	1.2	1.6	3.6	11.2	32.6	79.2	114.5
Female.....	6.8	*	0.7	0.7	0.9	1.0	1.3	2.6	6.4	17.6	38.1	63.9
Black, both sexes.....	4.6	*	*	0.7	0.9	1.1	2.0	3.6	9.1	22.5	39.3	51.9
Male.....	5.0	*	*	0.8	1.1	1.2	2.3	4.5	10.6	29.6	52.1	76.5
Female.....	4.2	*	*	*	0.8	1.0	1.8	2.8	7.8	17.5	32.0	41.7
Multiple myeloma and immunoproliferative neoplasms (C88,C90)												
All races 3/, both sexes..	3.7	*	*	*	*	*	0.4	1.6	5.5	15.1	29.6	34.1
Male.....	4.1	*	*	*	*	*	0.4	2.0	6.5	18.2	38.0	49.4
Female.....	3.4	*	*	*	*	*	0.3	1.3	4.5	12.4	23.8	26.8
White, both sexes.....	3.8	*	*	*	*	*	0.3	1.3	4.8	14.0	28.3	33.1
Male.....	4.2	*	*	*	*	*	0.4	1.7	5.9	17.4	36.7	49.1
Female.....	3.4	*	*	*	*	*	0.2	1.0	3.9	11.1	22.4	25.5
Black, both sexes.....	4.7	*	*	*	*	*	0.9	3.7	11.5	28.7	50.0	53.7
Male.....	4.6	*	*	*	*	*	0.9	4.2	13.1	31.4	63.9	65.4
Female.....	4.8	*	*	*	*	*	0.9	3.2	10.3	26.7	42.1	48.8

Worktable 250R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race, and sex: United States, 2007

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2007. Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
---	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Other and unspecified malignant neoplasms of lymphoid, hematopoietic and related tissue (C96)

All races 3/, both sexes..	0.0	*	*	*	*	*	*	*	*	*	*	*
Male.....	0.0	*	*	*	*	*	*	*	*	*	*	*
Female.....	0.0	*	*	*	*	*	*	*	*	*	*	*
White, both sexes.....	0.0	*	*	*	*	*	*	*	*	*	*	*
Male.....	0.0	*	*	*	*	*	*	*	*	*	*	*
Female.....	0.0	*	*	*	*	*	*	*	*	*	*	*
Black, both sexes.....	*	*	*	*	*	*	*	*	*	*	*	*
Male.....	*	*	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*	*	*

All other and unspecified malignant neoplasms (C17, C23-C24, C26-C31, C37-C41, C44-C49, C51-C52, C57-C60, C62-C63, C66, C68-C69, C73-C80, C97)

All races 3/, both sexes..	21.0	0.5	0.7	0.6	1.3	1.7	4.0	12.4	33.4	71.6	139.7	206.7
Male.....	22.1	*	0.7	0.6	1.7	2.1	4.2	14.2	40.1	86.7	174.0	260.9
Female.....	19.8	*	0.7	0.7	0.9	1.3	3.9	10.6	27.2	58.8	116.0	180.9
White, both sexes.....	22.5	*	0.7	0.6	1.3	1.7	3.9	11.9	32.7	71.8	142.2	212.4
Male.....	23.7	*	0.6	0.6	1.7	2.1	4.0	13.6	39.3	86.9	177.7	269.8
Female.....	21.3	*	0.7	0.7	0.9	1.3	3.8	10.1	26.5	58.7	117.3	184.9
Black, both sexes.....	16.7	*	0.7	0.6	1.3	2.0	5.4	18.3	45.9	82.5	133.9	170.8
Male.....	17.5	*	*	*	1.4	2.6	5.9	21.9	58.0	100.6	163.4	205.7
Female.....	15.9	*	*	0.7	1.1	1.5	5.0	15.3	36.2	69.6	117.0	156.2

Worktable 250R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race, and sex: United States, 2007

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2007. Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)												
All races 3/, both sexes..	4.7	1.4	0.4	0.2	0.2	0.3	0.8	1.6	4.3	12.4	36.2	76.4
Male.....	4.9	1.1	0.3	0.2	0.2	0.4	0.8	1.8	5.3	15.2	48.4	95.7
Female.....	4.5	1.7	0.4	0.2	0.2	0.2	0.7	1.4	3.4	10.0	27.8	67.2
White, both sexes.....	5.2	1.4	0.4	0.2	0.2	0.3	0.8	1.6	4.3	12.5	37.4	79.4
Male.....	5.4	*	0.4	0.2	0.2	0.4	0.8	1.8	5.2	15.3	50.0	99.4
Female.....	5.0	1.8	0.4	0.2	0.2	0.2	0.7	1.4	3.4	10.2	28.5	69.9
Black, both sexes.....	3.0	*	*	0.3	*	0.5	1.1	2.0	5.2	13.6	30.1	49.4
Male.....	3.1	*	*	*	*	*	1.2	2.2	7.0	17.7	38.0	62.0
Female.....	2.9	*	*	*	*	*	1.1	1.9	3.7	10.7	25.6	44.2
Anemias (D50-D64)												
All races 3/, both sexes..	1.6	*	*	0.1	0.2	0.3	0.4	0.6	1.0	2.9	9.4	36.1
Male.....	1.3	*	*	0.1	0.2	0.3	0.5	0.6	1.0	3.0	10.1	34.3
Female.....	1.9	*	*	*	0.2	0.3	0.4	0.6	1.0	2.8	8.9	36.9
White, both sexes.....	1.5	*	*	*	0.1	0.1	0.1	0.3	0.8	2.6	9.2	36.4
Male.....	1.2	*	*	*	*	*	*	0.3	0.8	2.7	9.9	34.5
Female.....	1.9	*	*	*	*	*	0.1	0.3	0.8	2.5	8.7	37.3
Black, both sexes.....	2.4	*	*	*	1.0	1.8	2.5	2.6	3.0	6.0	13.0	35.7
Male.....	2.3	*	*	*	1.1	2.0	2.9	2.8	3.1	5.9	14.2	33.1
Female.....	2.5	*	*	*	0.9	1.6	2.1	2.5	2.9	6.1	12.3	36.8

Worktable 250R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race, and sex: United States, 2007

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2007. Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Diabetes mellitus (E10-E14)												
All races 3/, both sexes..	23.7	*	*	0.1	0.4	1.5	4.6	13.1	34.6	78.1	162.7	276.2
Male.....	23.9	*	*	*	0.4	1.7	5.7	16.3	42.1	93.2	185.9	307.4
Female.....	23.5	*	*	*	0.3	1.3	3.5	10.1	27.5	65.3	146.7	261.4
White, both sexes.....	23.2	*	*	*	0.3	1.2	3.9	11.4	30.3	70.1	151.2	262.3
Male.....	23.8	*	*	*	0.4	1.4	4.9	14.4	37.7	86.3	177.7	300.4
Female.....	22.5	*	*	*	0.2	0.9	2.8	8.4	23.2	56.1	132.7	244.1
Black, both sexes.....	31.1	*	*	*	0.8	3.7	9.9	27.0	73.8	155.4	297.8	442.1
Male.....	28.7	*	*	*	0.6	4.2	11.8	32.4	86.0	171.6	312.5	393.5
Female.....	33.3	*	*	*	1.0	3.3	8.3	22.4	63.9	143.9	289.4	462.4
Nutritional deficiencies (E40-E64)												
All races 3/, both sexes..	0.9	*	*	*	*	*	0.1	0.2	0.6	1.7	6.2	24.4
Male.....	0.7	*	*	*	*	*	*	0.2	0.6	1.9	6.3	22.1
Female.....	1.2	*	*	*	*	*	0.1	0.2	0.6	1.6	6.2	25.5
White, both sexes.....	1.0	*	*	*	*	*	0.1	0.2	0.6	1.5	5.9	23.3
Male.....	0.7	*	*	*	*	*	*	0.2	0.5	1.6	5.9	21.2
Female.....	1.2	*	*	*	*	*	*	0.2	0.6	1.4	5.9	24.4
Black, both sexes.....	1.0	*	*	*	*	*	*	*	0.9	3.8	10.4	36.4
Male.....	0.9	*	*	*	*	*	*	*	1.4	4.7	12.6	33.1
Female.....	1.1	*	*	*	*	*	*	*	*	3.2	9.1	37.8

Worktable 250R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race, and sex: United States, 2007

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2007. Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Meningitis (G00,G03)												
All races 3/, both sexes..	0.2	1.9	*	0.1	0.1	0.1	0.2	0.3	0.3	0.4	0.7	0.7
Male.....	0.2	1.6	*	*	*	*	0.2	0.3	0.3	0.5	0.8	*
Female.....	0.2	2.3	*	*	*	*	0.1	0.2	0.2	0.3	0.6	0.7
White, both sexes.....	0.2	1.6	*	*	0.1	0.1	0.2	0.2	0.2	0.3	0.7	0.7
Male.....	0.2	*	*	*	*	*	0.2	0.3	0.3	0.4	0.8	*
Female.....	0.2	2.0	*	*	*	*	0.1	0.2	0.2	0.3	0.6	0.7
Black, both sexes.....	0.3	3.2	*	*	*	*	*	0.5	0.7	*	*	*
Male.....	0.4	*	*	*	*	*	*	*	*	*	*	*
Female.....	0.3	*	*	*	*	*	*	*	*	*	*	*
Parkinson's disease (G20-G21)												
All races 3/, both sexes..	6.7	*	*	*	*	*	*	0.1	1.2	11.9	71.9	143.5
Male.....	7.8	*	*	*	*	*	*	0.2	1.6	17.0	107.2	228.0
Female.....	5.6	*	*	*	*	*	*	0.1	0.9	7.7	47.6	103.3
White, both sexes.....	7.7	*	*	*	*	*	*	0.1	1.3	12.7	76.7	151.6
Male.....	9.0	*	*	*	*	*	*	0.2	1.7	17.8	113.9	240.9
Female.....	6.5	*	*	*	*	*	*	*	0.9	8.3	50.7	109.0
Black, both sexes.....	2.0	*	*	*	*	*	*	*	1.1	7.3	35.0	65.9
Male.....	2.3	*	*	*	*	*	*	*	1.3	12.4	53.7	99.4
Female.....	1.7	*	*	*	*	*	*	*	*	3.7	24.4	52.0

Worktable 250R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race, and sex: United States, 2007

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2007. Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Alzheimer's disease (G30)												
All races 3/, both sexes..	24.7	*	*	*	*	*	*	0.2	2.2	20.6	176.7	849.1
Male.....	14.7	*	*	*	*	*	*	0.2	2.1	19.1	155.3	646.0
Female.....	34.5	*	*	*	*	*	*	0.2	2.3	21.9	191.4	945.7
White, both sexes.....	28.3	*	*	*	*	*	*	0.2	2.4	21.4	183.4	879.4
Male.....	16.7	*	*	*	*	*	*	0.2	2.2	19.7	160.5	670.4
Female.....	39.7	*	*	*	*	*	*	0.3	2.6	22.9	199.5	979.2
Black, both sexes.....	11.9	*	*	*	*	*	*	*	1.1	19.6	157.9	680.1
Male.....	6.8	*	*	*	*	*	*	*	*	19.8	144.1	500.6
Female.....	16.5	*	*	*	*	*	*	*	1.1	19.4	165.8	754.7
Major cardiovascular diseases (I00-I78)												
All races 3/, both sexes..	267.3	13.4	1.4	0.8	3.2	9.7	34.4	105.5	247.0	597.5	1,758.7	5,679.0
Male.....	261.0	14.8	1.2	0.9	4.0	12.4	46.8	148.4	346.4	781.3	2,127.2	5,894.7
Female.....	273.3	11.9	1.6	0.8	2.4	6.9	22.1	63.9	154.4	441.4	1,504.9	5,576.4
White, both sexes.....	283.8	11.0	1.2	0.8	2.7	8.0	29.9	93.6	225.5	565.7	1,744.9	5,777.1
Male.....	274.6	12.0	1.0	0.8	3.3	10.3	41.2	135.0	319.9	745.1	2,116.2	6,037.2
Female.....	292.8	9.9	1.4	0.7	2.1	5.5	18.3	52.7	135.7	410.3	1,484.9	5,652.9
Black, both sexes.....	240.0	24.5	2.2	1.2	6.2	21.3	70.0	210.5	477.3	994.3	2,215.7	5,428.0
Male.....	244.2	29.0	2.3	1.2	8.2	26.9	93.3	278.9	662.1	1,306.3	2,695.4	5,181.6
Female.....	236.1	19.7	2.0	1.2	4.1	16.2	49.3	151.6	328.4	771.1	1,942.2	5,530.5

Worktable 250R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race, and sex: United States, 2007

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2007. Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Diseases of heart (I00-I09,I11,I13,I20-I51)												
All races 3/, both sexes..	204.3	10.0	1.1	0.6	2.6	7.9	27.4	85.3	200.3	462.9	1,315.0	4,267.7
Male.....	208.4	10.9	1.0	0.6	3.2	10.5	38.6	124.6	288.8	624.9	1,656.5	4,621.8
Female.....	200.2	9.0	1.1	0.6	1.9	5.3	16.2	47.2	117.9	325.4	1,079.7	4,099.3
White, both sexes.....	218.3	7.7	0.9	0.5	2.1	6.5	24.4	77.9	186.3	443.0	1,312.1	4,359.1
Male.....	221.1	8.5	0.7	0.6	2.6	8.7	34.7	116.2	271.4	603.0	1,659.3	4,756.1
Female.....	215.5	6.9	1.0	0.5	1.6	4.2	13.8	40.0	105.3	304.4	1,068.9	4,169.6
Black, both sexes.....	177.9	19.4	1.6	1.0	5.2	17.4	52.8	157.6	364.6	741.1	1,607.3	3,937.8
Male.....	186.5	21.6	2.0	1.1	6.9	22.8	73.2	216.3	516.3	989.4	1,999.2	3,879.6
Female.....	170.0	17.1	*	0.9	3.4	12.4	34.6	107.0	242.5	563.5	1,384.0	3,962.0
Acute rheumatic fever and chronic rheumatic heart diseases (I00-I09)												
All races 3/, both sexes..	1.1	*	*	*	*	0.1	0.2	0.4	1.0	2.7	8.1	17.9
Male.....	0.7	*	*	*	*	*	0.1	0.4	0.8	2.0	6.3	14.4
Female.....	1.4	*	*	*	*	*	0.2	0.4	1.2	3.4	9.4	19.6
White, both sexes.....	1.2	*	*	*	*	*	0.1	0.3	0.9	2.8	8.6	19.1
Male.....	0.8	*	*	*	*	*	0.1	0.3	0.8	2.0	6.5	15.4
Female.....	1.6	*	*	*	*	*	0.2	0.4	1.1	3.4	10.1	20.8
Black, both sexes.....	0.6	*	*	*	*	*	*	0.8	1.7	2.3	4.5	8.5
Male.....	0.4	*	*	*	*	*	*	*	1.6	*	*	*
Female.....	0.8	*	*	*	*	*	*	0.9	1.8	3.0	4.1	9.5

Worktable 250R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race, and sex: United States, 2007

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2007. Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Hypertensive heart disease (I11)												
All races 3/, both sexes..	10.2	*	*	*	0.1	0.8	3.2	8.2	13.7	20.7	48.6	192.3
Male.....	9.6	*	*	*	0.2	1.1	4.4	11.5	19.1	25.5	50.3	146.3
Female.....	10.8	*	*	*	*	0.5	1.9	5.1	8.7	16.7	47.3	214.1
White, both sexes.....	9.4	*	*	*	*	0.5	2.2	6.2	10.5	16.7	42.7	184.6
Male.....	8.5	*	*	*	*	0.7	3.2	9.0	14.9	20.6	43.6	138.5
Female.....	10.4	*	*	*	*	0.3	1.2	3.5	6.3	13.3	42.1	206.5
Black, both sexes.....	17.8	*	*	*	0.4	2.8	9.9	23.9	43.5	61.5	119.9	311.8
Male.....	19.2	*	*	*	*	3.9	13.6	32.2	61.3	81.0	141.7	266.9
Female.....	16.4	*	*	*	*	1.8	6.7	16.8	29.2	47.5	107.4	330.5
Hypertensive heart and renal disease (I13)												
All races 3/, both sexes..	1.0	*	*	*	*	0.1	0.2	0.4	0.9	2.0	6.1	22.3
Male.....	0.9	*	*	*	*	*	0.3	0.6	1.1	2.1	6.5	23.0
Female.....	1.1	*	*	*	*	*	0.1	0.3	0.7	1.8	5.8	22.0
White, both sexes.....	0.8	*	*	*	*	*	0.1	0.2	0.5	1.3	5.1	21.0
Male.....	0.7	*	*	*	*	*	*	0.3	0.6	1.4	5.5	21.5
Female.....	1.0	*	*	*	*	*	*	*	0.4	1.1	4.7	20.7
Black, both sexes.....	2.1	*	*	*	*	*	1.0	2.0	4.5	8.3	17.4	39.4
Male.....	2.2	*	*	*	*	*	1.6	2.6	6.6	9.5	19.2	42.5
Female.....	2.0	*	*	*	*	*	*	1.5	2.9	7.4	16.3	38.2

Worktable 250R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race, and sex: United States, 2007

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2007. Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Ischemic heart diseases (I20-I25)												
All races 3/, both sexes..	134.7	0.6	*	0.1	0.4	2.6	14.4	55.6	141.1	325.7	891.9	2,705.3
Male.....	145.3	*	*	*	0.5	3.9	21.9	85.8	212.2	458.8	1,180.1	3,089.2
Female.....	124.4	*	*	*	0.2	1.2	6.9	26.3	74.9	212.6	693.3	2,522.7
White, both sexes.....	145.5	*	*	*	0.3	2.3	13.9	53.4	135.3	316.4	893.9	2,762.1
Male.....	156.6	*	*	*	0.4	3.5	21.3	83.5	205.3	449.8	1,189.4	3,183.0
Female.....	134.7	*	*	*	0.2	1.1	6.4	23.5	68.7	200.9	686.9	2,561.2
Black, both sexes.....	106.6	*	*	*	0.7	4.4	20.3	84.3	222.6	473.2	1,033.9	2,466.1
Male.....	113.8	*	*	*	1.0	6.9	29.8	122.1	324.8	648.9	1,321.2	2,497.7
Female.....	100.0	*	*	*	*	2.1	11.8	51.7	140.3	347.6	870.1	2,452.9
Acute myocardial infarction (I21-I22)												
All races 3/, both sexes..	44.1	*	*	*	0.1	1.0	5.6	21.6	54.5	121.1	288.9	756.7
Male.....	48.2	*	*	*	0.2	1.4	8.2	33.0	80.8	166.8	370.8	855.8
Female.....	40.0	*	*	*	0.1	0.5	2.9	10.5	30.1	82.3	232.5	709.5
White, both sexes.....	47.6	*	*	*	0.1	0.9	5.5	21.3	53.7	119.4	289.4	766.9
Male.....	52.2	*	*	*	0.1	1.3	8.2	33.1	80.5	165.9	373.7	876.2
Female.....	43.1	*	*	*	*	0.5	2.8	9.6	28.2	79.2	230.3	714.8
Black, both sexes.....	35.2	*	*	*	*	1.7	7.3	28.8	75.8	163.0	337.5	750.5
Male.....	36.6	*	*	*	*	2.5	10.3	40.2	105.2	215.3	409.8	753.8
Female.....	34.0	*	*	*	*	0.9	4.6	19.0	52.1	125.5	296.3	749.1

Worktable 250R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race, and sex: United States, 2007

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2007. Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Other acute ischemic heart diseases (I24)												
All races 3/, both sexes..	1.4	*	*	*	*	*	0.3	0.9	2.1	3.8	7.8	20.8
Male.....	1.5	*	*	*	*	*	0.4	1.2	3.1	5.4	9.4	18.7
Female.....	1.3	*	*	*	*	*	0.2	0.5	1.1	2.5	6.8	21.8
White, both sexes.....	1.4	*	*	*	*	*	0.2	0.8	1.9	3.6	7.7	20.9
Male.....	1.5	*	*	*	*	*	0.3	1.1	2.8	5.1	9.2	18.8
Female.....	1.3	*	*	*	*	*	*	0.4	1.1	2.3	6.7	21.9
Black, both sexes.....	1.3	*	*	*	*	*	0.4	1.4	3.5	6.1	11.1	21.7
Male.....	1.6	*	*	*	*	*	*	2.1	5.6	9.0	14.2	20.4
Female.....	1.1	*	*	*	*	*	*	0.8	1.8	4.1	9.3	22.3
Other forms of chronic ischemic heart disease (I20,I25)												
All races 3/, both sexes..	89.3	*	*	*	0.2	1.6	8.6	33.2	84.5	200.7	595.1	1,927.9
Male.....	95.6	*	*	*	0.3	2.4	13.4	51.5	128.3	286.6	799.9	2,214.7
Female.....	83.1	*	*	*	0.1	0.7	3.8	15.3	43.7	127.8	454.0	1,791.4
White, both sexes.....	96.5	*	*	*	0.2	1.4	8.2	31.3	79.6	193.3	596.8	1,974.2
Male.....	102.9	*	*	*	0.3	2.2	12.8	49.2	121.9	278.8	806.5	2,287.9
Female.....	90.3	*	*	*	*	0.6	3.5	13.5	39.4	119.3	449.8	1,824.5
Black, both sexes.....	70.1	*	*	*	0.4	2.6	12.5	54.1	143.4	304.1	685.3	1,693.9
Male.....	75.7	*	*	*	0.7	4.2	18.9	79.8	214.1	424.6	897.2	1,723.5
Female.....	64.9	*	*	*	*	1.2	6.9	31.9	86.4	217.9	564.5	1,681.6

Worktable 250R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race, and sex: United States, 2007

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2007. Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Atherosclerotic cardiovascular disease, so described (I25.0)												
All races 3/, both sexes..	19.6	*	*	*	*	0.5	3.2	13.5	30.7	51.1	109.2	314.7
Male.....	22.5	*	*	*	*	0.9	5.1	21.4	48.1	73.7	142.2	326.1
Female.....	16.8	*	*	*	*	0.2	1.3	5.8	14.6	31.9	86.5	309.3
White, both sexes.....	20.0	*	*	*	*	0.5	3.0	12.4	28.1	46.6	104.2	312.5
Male.....	22.7	*	*	*	*	0.7	4.9	20.0	44.1	67.7	135.4	325.4
Female.....	17.3	*	*	*	*	0.2	1.2	5.0	12.8	28.3	82.4	306.3
Black, both sexes.....	22.3	*	*	*	*	0.9	4.9	23.6	59.7	102.0	179.9	395.7
Male.....	27.0	*	*	*	*	1.7	7.6	36.2	94.0	151.9	254.5	408.8
Female.....	18.0	*	*	*	*	*	2.5	12.9	32.1	66.3	137.3	390.3
All other forms of chronic ischemic heart disease (I20,I25.1-I25.9)												
All races 3/, both sexes..	69.7	*	*	*	0.2	1.0	5.4	19.7	53.8	149.7	485.9	1,613.2
Male.....	73.2	*	*	*	0.3	1.6	8.2	30.2	80.3	213.0	657.7	1,888.6
Female.....	66.3	*	*	*	*	0.5	2.5	9.5	29.1	95.9	367.5	1,482.2
White, both sexes.....	76.6	*	*	*	0.2	1.0	5.1	18.8	51.6	146.8	492.5	1,661.8
Male.....	80.2	*	*	*	0.2	1.5	7.9	29.2	77.8	211.1	671.2	1,962.6
Female.....	72.9	*	*	*	*	0.4	2.3	8.5	26.6	91.1	367.4	1,518.2
Black, both sexes.....	47.8	*	*	*	0.4	1.7	7.6	30.4	83.7	202.1	505.4	1,298.2
Male.....	48.7	*	*	*	*	2.5	11.2	43.6	120.1	272.7	642.7	1,314.7
Female.....	46.9	*	*	*	*	0.9	4.4	19.1	54.4	151.7	427.2	1,291.3

Worktable 250R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race, and sex: United States, 2007

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2007. Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Other heart diseases (I26-I51)												
All races 3/, both sexes..	57.3	9.3	1.0	0.5	2.1	4.4	9.5	20.7	43.5	111.8	360.4	1,329.8
Male.....	51.9	10.3	0.9	0.6	2.5	5.3	11.9	26.5	55.5	136.4	413.4	1,348.9
Female.....	62.5	8.3	1.1	0.5	1.6	3.4	7.1	15.1	32.4	91.0	323.9	1,320.8
White, both sexes.....	61.3	7.1	0.8	0.5	1.8	3.6	8.0	17.8	39.1	105.8	361.8	1,372.4
Male.....	54.6	7.9	0.7	0.5	2.1	4.4	9.9	23.0	49.9	129.1	414.3	1,397.6
Female.....	67.9	6.3	1.0	0.4	1.4	2.7	6.0	12.6	28.8	85.6	325.1	1,360.4
Black, both sexes.....	50.8	19.0	1.5	0.9	3.9	9.8	21.3	46.6	92.2	195.8	431.8	1,112.0
Male.....	50.9	21.4	2.0	1.0	5.2	11.6	28.0	58.8	122.0	248.6	512.0	1,066.6
Female.....	50.8	16.5	*	0.8	2.6	8.1	15.3	36.1	68.2	158.0	386.0	1,130.8
Acute and subacute endocarditis (I33)												
All races 3/, both sexes..	0.4	*	*	*	*	0.1	0.2	0.5	0.6	1.2	2.4	2.4
Male.....	0.5	*	*	*	*	0.1	0.3	0.6	0.8	1.4	3.2	3.4
Female.....	0.3	*	*	*	*	*	0.1	0.3	0.4	1.0	1.8	2.0
White, both sexes.....	0.4	*	*	*	*	0.1	0.2	0.4	0.5	1.1	2.4	2.4
Male.....	0.5	*	*	*	*	0.1	0.3	0.4	0.7	1.2	3.2	3.4
Female.....	0.3	*	*	*	*	*	0.1	0.3	0.4	1.0	1.8	1.9
Black, both sexes.....	0.6	*	*	*	*	*	0.4	1.1	1.5	2.7	2.3	*
Male.....	0.8	*	*	*	*	*	*	1.7	2.3	3.8	*	*
Female.....	0.4	*	*	*	*	*	*	*	*	2.0	*	*

Worktable 250R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race, and sex: United States, 2007

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2007. Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Diseases of pericardium and acute myocarditis (I30-I31,I40)												
All races 3/, both sexes..	0.3	0.6	*	0.1	0.1	0.1	0.2	0.3	0.4	0.6	1.1	2.1
Male.....	0.3	*	*	*	0.1	0.2	0.3	0.3	0.5	0.7	1.1	2.3
Female.....	0.3	*	*	*	0.1	0.1	0.1	0.3	0.3	0.5	1.0	2.0
White, both sexes.....	0.3	*	*	*	0.1	0.1	0.2	0.2	0.3	0.6	1.1	2.1
Male.....	0.3	*	*	*	*	0.2	0.2	0.2	0.4	0.7	1.1	2.3
Female.....	0.3	*	*	*	*	*	0.1	0.2	0.3	0.5	1.1	2.0
Black, both sexes.....	0.4	*	*	*	*	*	*	0.5	0.8	*	*	*
Male.....	0.4	*	*	*	*	*	*	*	*	*	*	*
Female.....	0.3	*	*	*	*	*	*	*	*	*	*	*
Heart failure (I50)												
All races 3/, both sexes..	18.8	0.5	*	*	0.1	0.2	0.7	2.4	8.4	29.7	122.4	554.4
Male.....	15.4	*	*	*	0.1	0.3	1.0	3.0	10.4	36.0	137.1	552.8
Female.....	22.0	*	*	*	*	0.2	0.5	1.9	6.6	24.3	112.2	555.1
White, both sexes.....	20.7	*	*	*	0.1	0.1	0.5	1.9	7.3	28.5	123.7	574.5
Male.....	16.8	*	*	*	*	0.2	0.6	2.4	9.0	34.8	138.6	575.8
Female.....	24.5	*	*	*	*	*	0.4	1.4	5.7	23.0	113.3	573.8
Black, both sexes.....	13.7	*	*	*	*	0.7	2.4	7.1	20.1	50.0	143.0	460.4
Male.....	12.2	*	*	*	*	0.8	3.3	9.0	26.7	60.8	163.4	433.4
Female.....	15.1	*	*	*	*	*	1.6	5.5	14.8	42.2	131.4	471.6

Worktable 250R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race, and sex: United States, 2007

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2007. Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
All other forms of heart disease (I26-I28,I34-I38,I42-I49,I51)												
All races 3/, both sexes..	37.8	8.2	0.8	0.5	1.8	4.0	8.3	17.5	34.1	80.3	234.6	770.9
Male.....	35.7	9.0	0.7	0.5	2.3	4.8	10.4	22.6	43.8	98.2	271.9	790.5
Female.....	39.9	7.3	0.9	0.4	1.3	3.1	6.3	12.7	25.0	65.1	208.9	761.7
White, both sexes.....	39.9	6.3	0.7	0.4	1.5	3.2	7.1	15.3	30.9	75.7	234.6	793.5
Male.....	37.0	7.0	0.5	0.4	1.9	4.0	8.7	20.0	39.7	92.4	271.3	816.1
Female.....	42.8	5.7	0.8	0.4	1.2	2.4	5.4	10.6	22.4	61.2	208.9	782.7
Black, both sexes.....	36.2	16.2	1.3	0.8	3.5	8.8	18.2	37.9	69.8	142.2	285.7	646.4
Male.....	37.5	18.1	1.6	0.8	4.6	10.4	23.8	47.6	92.1	182.9	344.1	627.2
Female.....	35.0	14.3	*	0.8	2.4	7.3	13.2	29.4	51.9	113.0	252.4	654.3
Essential hypertension and hypertensive renal disease (I10,I12,I15)												
All races 3/, both sexes..	7.9	*	*	*	0.1	0.2	0.9	2.8	6.5	16.2	49.5	191.1
Male.....	6.3	*	*	*	*	0.2	1.2	3.5	8.1	18.0	50.2	157.9
Female.....	9.5	*	*	*	*	0.2	0.6	2.1	5.0	14.6	49.0	207.0
White, both sexes.....	7.6	*	*	*	*	0.1	0.6	1.9	4.7	13.2	44.4	183.0
Male.....	5.9	*	*	*	*	0.1	0.8	2.4	6.1	14.7	45.1	150.4
Female.....	9.3	*	*	*	*	*	0.3	1.3	3.4	11.9	44.0	198.6
Black, both sexes.....	11.6	*	*	*	*	0.9	3.1	10.1	22.3	45.1	106.3	297.6
Male.....	10.3	*	*	*	*	0.8	3.8	12.5	27.0	53.5	116.6	245.6
Female.....	12.7	*	*	*	*	0.9	2.5	8.0	18.5	39.0	100.4	319.2

Worktable 250R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race, and sex: United States, 2007

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2007. Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Cerebrovascular diseases (I60-I69)												
All races 3/, both sexes..	45.1	3.1	0.3	0.2	0.5	1.2	4.9	14.6	32.1	93.0	322.3	1,015.5
Male.....	36.4	3.5	0.2	0.2	0.5	1.2	5.3	16.2	38.0	105.2	333.2	895.7
Female.....	53.5	2.6	0.4	0.2	0.4	1.3	4.6	12.9	26.6	82.7	314.9	1,072.4
White, both sexes.....	47.1	2.9	0.3	0.2	0.4	1.0	3.9	11.5	26.9	84.4	315.7	1,026.1
Male.....	37.0	3.0	*	0.2	0.4	1.0	4.3	13.0	31.4	94.3	323.1	905.0
Female.....	57.0	2.7	*	0.2	0.4	1.1	3.6	10.0	22.5	75.8	310.5	1,083.8
Black, both sexes.....	42.7	4.8	*	*	0.7	2.6	11.5	36.7	76.9	176.9	425.6	996.6
Male.....	39.5	6.9	*	*	0.9	2.6	12.6	41.0	99.8	223.3	491.9	866.9
Female.....	45.6	*	*	*	*	2.6	10.6	33.0	58.4	143.8	387.9	1,050.6
Atherosclerosis (I70)												
All races 3/, both sexes..	2.7	*	*	*	*	*	0.1	0.3	1.1	4.3	17.6	83.3
Male.....	2.2	*	*	*	*	*	*	0.4	1.5	5.3	19.3	76.5
Female.....	3.3	*	*	*	*	*	*	0.2	0.6	3.4	16.5	86.5
White, both sexes.....	3.1	*	*	*	*	*	0.1	0.3	1.0	4.2	18.0	86.0
Male.....	2.4	*	*	*	*	*	*	0.4	1.4	5.3	20.0	79.2
Female.....	3.7	*	*	*	*	*	*	0.2	0.7	3.4	16.7	89.2
Black, both sexes.....	1.6	*	*	*	*	*	*	0.6	1.6	5.4	17.2	69.9
Male.....	1.4	*	*	*	*	*	*	0.9	2.8	7.0	16.8	62.9
Female.....	1.9	*	*	*	*	*	*	*	*	4.3	17.4	72.8

Worktable 250R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race, and sex: United States, 2007

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2007. Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Other diseases of circulatory system (I71-I78)												
All races 3/, both sexes..	7.3	*	*	*	0.2	0.3	1.1	2.5	7.0	21.0	54.3	121.4
Male.....	7.7	*	*	*	0.2	0.4	1.6	3.5	10.0	27.8	68.0	142.8
Female.....	6.8	*	*	*	*	0.2	0.6	1.5	4.2	15.2	44.9	111.3
White, both sexes.....	7.8	*	*	*	0.1	0.3	1.0	2.1	6.6	20.9	54.7	122.9
Male.....	8.2	*	*	*	0.2	0.4	1.4	3.0	9.5	27.9	68.7	146.5
Female.....	7.3	*	*	*	*	0.2	0.5	1.2	3.7	14.9	44.9	111.7
Black, both sexes.....	6.2	*	*	*	*	0.5	2.5	5.6	11.9	25.7	59.2	126.1
Male.....	6.5	*	*	*	*	*	3.7	8.1	16.3	33.0	70.8	126.6
Female.....	5.9	*	*	*	*	*	1.5	3.4	8.3	20.6	52.6	125.8
Aortic aneurysm and dissection (I71)												
All races 3/, both sexes..	4.3	*	*	*	0.1	0.2	0.8	1.7	4.5	13.5	33.4	59.5
Male.....	5.1	*	*	*	0.1	0.4	1.3	2.6	6.9	19.1	44.1	81.8
Female.....	3.6	*	*	*	*	0.1	0.4	0.8	2.3	8.8	26.0	48.9
White, both sexes.....	4.7	*	*	*	0.1	0.2	0.7	1.4	4.3	13.8	34.4	61.6
Male.....	5.5	*	*	*	0.1	0.3	1.0	2.2	6.7	19.6	45.5	85.8
Female.....	3.9	*	*	*	*	*	0.3	0.6	2.0	8.8	26.6	50.1
Black, both sexes.....	3.1	*	*	*	*	*	2.1	3.9	6.9	12.9	26.3	40.4
Male.....	3.7	*	*	*	*	*	3.3	6.1	9.9	16.8	31.5	41.6
Female.....	2.6	*	*	*	*	*	1.0	2.1	4.4	10.2	23.3	39.9

Worktable 250R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race, and sex: United States, 2007

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2007. Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Other diseases of arteries, arterioles and capillaries (I72-I78)												
All races 3/, both sexes..	3.0	*	*	*	0.1	0.1	0.3	0.8	2.5	7.5	20.9	62.0
Male.....	2.6	*	*	*	*	*	0.3	0.9	3.1	8.7	23.9	61.0
Female.....	3.3	*	*	*	*	0.1	0.2	0.6	1.9	6.4	18.9	62.4
White, both sexes.....	3.1	*	*	*	0.1	0.1	0.3	0.7	2.3	7.1	20.3	61.3
Male.....	2.8	*	*	*	*	*	0.3	0.8	2.8	8.2	23.2	60.7
Female.....	3.5	*	*	*	*	*	0.2	0.5	1.7	6.1	18.3	61.6
Black, both sexes.....	3.0	*	*	*	*	*	0.4	1.6	5.0	12.8	32.9	85.6
Male.....	2.8	*	*	*	*	*	*	2.0	6.4	16.2	39.3	85.0
Female.....	3.2	*	*	*	*	*	*	1.3	3.9	10.4	29.3	85.9
Other disorders of circulatory system (I80-I99)												
All races 3/, both sexes..	1.4	1.2	*	*	0.1	0.3	0.7	1.3	1.8	2.9	6.8	17.4
Male.....	1.3	1.4	*	*	0.1	0.3	0.9	1.6	2.1	3.1	6.9	15.8
Female.....	1.4	1.1	*	*	*	0.3	0.6	1.1	1.5	2.6	6.8	18.2
White, both sexes.....	1.4	1.2	*	*	0.1	0.2	0.6	1.1	1.6	2.8	6.8	17.4
Male.....	1.3	1.3	*	*	*	0.3	0.7	1.4	1.8	3.1	6.8	16.0
Female.....	1.5	*	*	*	*	0.2	0.5	0.9	1.4	2.5	6.8	18.0
Black, both sexes.....	1.8	*	*	*	*	0.7	1.8	2.7	3.7	4.5	8.9	23.0
Male.....	1.8	*	*	*	*	0.7	2.0	3.3	5.1	4.8	9.6	19.5
Female.....	1.7	*	*	*	*	0.7	1.6	2.2	2.6	4.3	8.5	24.4

Worktable 250R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race, and sex: United States, 2007

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2007. Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Emphysema (J43)												
All races 3/, both sexes..	4.2	*	*	*	*	*	0.1	1.1	4.9	17.0	37.1	45.5
Male.....	4.4	*	*	*	*	*	0.2	1.4	5.7	20.1	46.2	61.6
Female.....	4.0	*	*	*	*	*	*	0.8	4.1	14.4	30.8	37.9
White, both sexes.....	4.9	*	*	*	*	*	0.1	1.1	5.2	18.4	39.4	48.2
Male.....	5.0	*	*	*	*	*	0.2	1.4	5.9	21.4	47.9	64.0
Female.....	4.8	*	*	*	*	*	*	0.8	4.5	15.9	33.4	40.6
Black, both sexes.....	1.8	*	*	*	*	*	*	1.3	3.8	10.4	22.6	24.0
Male.....	2.4	*	*	*	*	*	*	1.7	5.9	16.0	37.2	45.9
Female.....	1.3	*	*	*	*	*	*	0.9	2.2	6.4	14.3	14.8
Asthma (J45-J46)												
All races 3/, both sexes..	1.1	*	0.2	0.3	0.3	0.5	0.7	1.2	1.4	2.1	4.4	12.0
Male.....	0.9	*	0.2	0.3	0.4	0.6	0.6	1.0	1.2	1.5	3.2	7.1
Female.....	1.4	*	0.2	0.2	0.2	0.4	0.9	1.5	1.6	2.6	5.1	14.2
White, both sexes.....	1.0	*	*	0.1	0.2	0.4	0.5	0.9	1.1	1.8	4.0	11.8
Male.....	0.7	*	*	0.2	0.3	0.4	0.4	0.7	0.9	1.2	2.7	6.7
Female.....	1.3	*	*	*	0.1	0.3	0.7	1.0	1.3	2.3	4.9	14.2
Black, both sexes.....	2.3	*	1.0	0.9	0.9	1.4	2.1	3.8	4.3	5.1	6.4	11.7
Male.....	2.1	*	*	1.2	1.2	1.8	2.1	2.9	4.4	4.8	6.7	*
Female.....	2.4	*	*	*	*	0.9	2.2	4.7	4.2	5.3	6.2	14.1

Worktable 250R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race, and sex: United States, 2007

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2007. Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Pneumonitis due to solids and liquids (J69)												
All races 3/, both sexes..	5.6	*	*	*	0.1	0.2	0.4	1.0	2.7	8.9	39.8	153.3
Male.....	6.0	*	*	*	0.1	0.2	0.5	1.3	3.5	11.7	55.1	222.0
Female.....	5.3	*	*	*	*	0.1	0.3	0.7	2.0	6.5	29.3	120.7
White, both sexes.....	6.3	*	*	*	0.1	0.2	0.4	1.0	2.6	8.9	40.4	157.6
Male.....	6.6	*	*	*	0.1	0.2	0.5	1.3	3.3	11.6	55.3	229.4
Female.....	5.9	*	*	*	*	0.1	0.3	0.7	1.9	6.6	29.9	123.3
Black, both sexes.....	3.6	*	*	*	*	*	0.4	1.3	4.7	11.1	42.5	125.3
Male.....	3.9	*	*	*	*	*	*	1.6	6.7	16.8	65.2	160.6
Female.....	3.3	*	*	*	*	*	*	1.1	3.1	7.1	29.6	110.6
Other diseases of respiratory system (J00-J06,J30-J39,J67,J70-J98)												
All races 3/, both sexes..	9.5	7.5	0.6	0.2	0.4	0.5	1.1	3.1	9.3	28.7	73.2	138.9
Male.....	9.6	8.7	0.8	0.2	0.4	0.6	1.2	3.7	10.8	35.1	92.7	169.2
Female.....	9.3	6.2	0.4	0.2	0.3	0.5	1.1	2.6	7.9	23.3	59.8	124.5
White, both sexes.....	10.3	5.8	0.5	0.1	0.3	0.5	1.0	2.8	9.0	29.2	76.1	143.8
Male.....	10.5	6.9	0.6	*	0.4	0.5	1.0	3.3	10.5	35.7	96.3	175.8
Female.....	10.2	4.6	0.4	0.2	0.3	0.4	1.0	2.4	7.7	23.5	62.0	128.5
Black, both sexes.....	6.5	15.7	1.1	*	0.6	1.0	2.1	5.8	13.4	29.1	52.8	95.1
Male.....	6.6	17.8	1.7	*	0.6	0.9	2.2	6.9	16.2	34.1	65.8	113.9
Female.....	6.4	13.4	*	*	*	1.1	1.9	4.8	11.1	25.4	45.4	87.3

Worktable 250R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race, and sex: United States, 2007

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2007. Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Hernia (K40-K46)												
All races 3/, both sexes..	0.6	0.8	*	*	*	*	0.1	0.2	0.5	1.3	3.6	11.3
Male.....	0.5	1.0	*	*	*	*	*	0.3	0.6	1.2	3.6	12.0
Female.....	0.6	*	*	*	*	*	*	0.2	0.5	1.3	3.7	11.0
White, both sexes.....	0.6	0.9	*	*	*	*	0.1	0.2	0.5	1.2	3.7	12.1
Male.....	0.5	*	*	*	*	*	*	0.2	0.5	1.2	3.4	12.6
Female.....	0.7	*	*	*	*	*	*	0.2	0.5	1.2	3.8	11.8
Black, both sexes.....	0.4	*	*	*	*	*	*	0.4	0.7	2.0	4.0	6.0
Male.....	0.4	*	*	*	*	*	*	*	*	*	5.9	*
Female.....	0.4	*	*	*	*	*	*	*	*	2.1	*	*
Chronic liver disease and cirrhosis (K70,K73-K74)												
All races 3/, both sexes..	9.7	*	*	*	0.1	0.9	6.0	18.7	24.5	26.7	28.4	19.8
Male.....	12.9	*	*	*	*	1.3	8.0	26.8	36.3	36.2	36.7	25.8
Female.....	6.5	*	*	*	*	0.6	3.9	10.9	13.4	18.6	22.6	17.0
White, both sexes.....	10.5	*	*	*	0.1	1.0	6.3	19.4	24.8	27.8	29.8	20.6
Male.....	13.9	*	*	*	*	1.3	8.5	27.9	36.8	37.3	38.2	27.2
Female.....	7.1	*	*	*	*	0.6	4.2	11.0	13.5	19.7	24.0	17.5
Black, both sexes.....	6.4	*	*	*	*	0.5	3.6	16.1	25.7	21.9	15.2	9.7
Male.....	8.8	*	*	*	*	*	4.6	22.8	40.3	35.4	24.1	*
Female.....	4.2	*	*	*	*	*	2.6	10.3	14.0	12.3	10.2	9.9

Worktable 250R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race, and sex: United States, 2007

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2007. Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Alcoholic liver disease (K70)												
All races 3/, both sexes..	4.8	*	*	*	*	0.7	4.2	11.7	13.2	10.1	5.7	2.1
Male.....	7.1	*	*	*	*	1.0	5.6	17.0	20.9	16.9	10.4	4.7
Female.....	2.5	*	*	*	*	0.5	2.9	6.5	6.0	4.3	2.5	0.9
White, both sexes.....	5.1	*	*	*	*	0.7	4.5	12.3	13.6	10.4	6.0	2.2
Male.....	7.7	*	*	*	*	1.0	6.0	17.9	21.5	17.2	10.8	4.9
Female.....	2.7	*	*	*	*	0.4	3.0	6.7	6.1	4.5	2.6	0.9
Black, both sexes.....	3.0	*	*	*	*	0.3	2.2	8.2	11.8	9.5	3.8	*
Male.....	4.5	*	*	*	*	*	2.9	12.1	19.5	18.1	8.0	*
Female.....	1.7	*	*	*	*	*	1.7	4.8	5.6	3.4	*	*
Other chronic liver disease and cirrhosis (K73-K74)												
All races 3/, both sexes..	4.9	*	*	*	*	0.2	1.7	7.0	11.3	16.6	22.6	17.7
Male.....	5.8	*	*	*	*	0.3	2.3	9.8	15.5	19.3	26.3	21.0
Female.....	4.0	*	*	*	*	0.2	1.1	4.4	7.4	14.3	20.1	16.1
White, both sexes.....	5.3	*	*	*	*	0.2	1.8	7.1	11.3	17.4	23.9	18.4
Male.....	6.2	*	*	*	*	0.3	2.4	9.9	15.2	20.1	27.4	22.3
Female.....	4.4	*	*	*	*	0.2	1.1	4.3	7.5	15.1	21.4	16.6
Black, both sexes.....	3.4	*	*	*	*	*	1.3	7.9	13.9	12.4	11.5	8.2
Male.....	4.3	*	*	*	*	*	1.8	10.7	20.8	17.3	16.0	*
Female.....	2.5	*	*	*	*	*	1.0	5.5	8.4	8.9	8.8	8.8

Worktable 250R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race, and sex: United States, 2007

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2007. Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Cholelithiasis and other disorders of gallbladder (K80-K82)												
All races 3/, both sexes..	1.1	*	*	*	*	0.1	0.1	0.3	0.7	2.5	7.8	23.3
Male.....	1.0	*	*	*	*	*	0.1	0.3	0.8	3.0	9.5	23.8
Female.....	1.2	*	*	*	*	*	0.1	0.3	0.6	2.0	6.5	23.1
White, both sexes.....	1.2	*	*	*	*	*	0.1	0.3	0.7	2.5	7.7	24.0
Male.....	1.1	*	*	*	*	*	*	0.3	0.8	3.1	9.7	24.4
Female.....	1.3	*	*	*	*	*	*	0.2	0.5	1.9	6.4	23.8
Black, both sexes.....	0.7	*	*	*	*	*	*	0.5	1.3	2.8	8.5	18.0
Male.....	0.6	*	*	*	*	*	*	*	*	2.9	10.2	19.5
Female.....	0.9	*	*	*	*	*	*	*	1.4	2.8	7.6	17.3
Nephritis, nephrotic syndrome and nephrosis (N00-N07, N17-N19, N25-N27)												
All races 3/, both sexes..	15.4	3.4	0.1	0.1	0.2	0.6	1.7	5.1	13.6	40.1	113.0	290.6
Male.....	15.2	3.6	*	*	0.2	0.7	2.0	5.8	15.7	45.5	139.9	376.3
Female.....	15.6	3.2	*	*	0.2	0.6	1.5	4.4	11.6	35.4	94.4	249.9
White, both sexes.....	15.1	2.9	*	*	0.2	0.4	1.2	3.7	10.4	34.0	106.7	282.5
Male.....	15.1	2.9	*	*	0.2	0.5	1.4	4.4	12.3	39.2	135.0	373.5
Female.....	15.2	2.9	*	*	0.2	0.4	1.0	3.1	8.7	29.5	86.9	239.0
Black, both sexes.....	21.0	6.6	*	*	0.4	2.0	5.8	15.5	41.3	100.0	199.9	437.1
Male.....	19.7	7.9	*	*	*	2.2	6.6	17.7	48.6	114.2	230.7	493.8
Female.....	22.1	*	*	*	*	1.8	5.1	13.6	35.4	89.8	182.3	413.6

Worktable 250R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race, and sex: United States, 2007

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2007. Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Acute and rapidly progressive nephritic and nephrotic syndrome (N00-N01,N04)												
All races 3/, both sexes..	0.1	*	*	*	*	*	*	*	0.1	0.2	0.5	0.9
Male.....	0.1	*	*	*	*	*	*	*	*	*	0.7	1.2
Female.....	0.1	*	*	*	*	*	*	*	*	*	0.3	0.7
White, both sexes.....	0.1	*	*	*	*	*	*	*	*	0.2	0.5	0.9
Male.....	0.1	*	*	*	*	*	*	*	*	*	0.7	*
Female.....	0.1	*	*	*	*	*	*	*	*	*	0.3	0.7
Black, both sexes.....	0.1	*	*	*	*	*	*	*	*	*	*	*
Male.....	*	*	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*	*	*
Chronic glomerulonephritis, nephritis and nephropathy not specified as acute or chronic, and renal sclerosis unspecified (N02-N03,N05-N07,N26)												
All races 3/, both sexes..	1.0	*	*	*	*	*	0.1	0.3	0.6	2.0	7.1	22.3
Male.....	1.0	*	*	*	*	*	0.1	0.3	0.8	2.4	8.7	29.9
Female.....	1.0	*	*	*	*	*	0.1	0.2	0.5	1.6	6.0	18.7
White, both sexes.....	1.0	*	*	*	*	*	0.1	0.2	0.5	1.7	6.6	21.0
Male.....	1.0	*	*	*	*	*	*	0.3	0.7	2.1	8.2	28.7
Female.....	1.0	*	*	*	*	*	*	0.2	0.4	1.3	5.4	17.4
Black, both sexes.....	1.3	*	*	*	*	*	0.4	0.8	1.7	5.1	14.1	38.7
Male.....	1.2	*	*	*	*	*	*	0.9	2.1	6.1	16.0	45.0
Female.....	1.4	*	*	*	*	*	*	0.7	1.3	4.3	13.0	36.1

Worktable 250R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race, and sex: United States, 2007

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2007. Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Inflammatory diseases of female pelvic organs (N70-N76)												
All races 3/, both sexes..	0.0	*	*	*	*	*	*	*	*	0.1	0.2	0.4
Male.....
Female.....	0.1	*	*	*	*	*	*	*	*	0.2	0.3	0.6
White, both sexes.....	0.0	*	*	*	*	*	*	*	*	*	0.2	0.4
Male.....
Female.....	0.1	*	*	*	*	*	*	*	*	*	0.3	0.6
Black, both sexes.....	0.0	*	*	*	*	*	*	*	*	*	*	*
Male.....
Female.....	0.1	*	*	*	*	*	*	*	*	*	*	*
Pregnancy, childbirth and the puerperium (000-099)												
All races 3/, both sexes..	0.3	*	0.4	0.8	0.5	0.1	*	*	*	*
Male.....
Female.....	0.5	*	0.8	1.6	1.0	0.3	*	*	*	*
White, both sexes.....	0.2	*	0.3	0.6	0.4	0.1	*	*	*	*
Male.....
Female.....	0.4	*	0.5	1.3	0.8	0.2	*	*	*	*
Black, both sexes.....	0.6	*	1.0	1.9	1.1	*	*	*	*	*
Male.....
Female.....	1.2	*	2.1	3.7	2.0	*	*	*	*	*

Worktable 250R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race, and sex: United States, 2007

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2007. Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Pregnancy with abortive outcome (000-007)												
All races 3/, both sexes..	0.0	*	*	*	*	*	*	*	*	*
Male.....
Female.....	0.0	*	*	*	*	*	*	*	*	*
White, both sexes.....	*	*	*	*	*	*	*	*	*	*
Male.....
Female.....	*	*	*	*	*	*	*	*	*	*
Black, both sexes.....	*	*	*	*	*	*	*	*	*	*
Male.....
Female.....	*	*	*	*	*	*	*	*	*	*
Other complications of pregnancy, childbirth and the puerperium (010-099)												
All races 3/, both sexes..	0.2	*	0.4	0.8	0.5	0.1	*	*	*	*
Male.....
Female.....	0.5	*	0.7	1.6	1.0	0.3	*	*	*	*
White, both sexes.....	0.2	*	0.3	0.6	0.4	0.1	*	*	*	*
Male.....
Female.....	0.4	*	0.5	1.2	0.8	0.2	*	*	*	*
Black, both sexes.....	0.6	*	0.9	1.7	1.0	*	*	*	*	*
Male.....
Female.....	1.1	*	1.9	3.4	1.9	*	*	*	*	*

Worktable 250R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race, and sex: United States, 2007

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2007. Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Certain conditions originating in the perinatal period (P00-P96)												
All races 3/, both sexes..	4.8	339.8	0.4	0.1	*	*	*	*	*	*	*	*
Male.....	5.5	373.6	0.4	*	*	*	*	*	*	*	*	*
Female.....	4.2	304.4	0.4	*	*	*	*	*	*	*	*	*
White, both sexes.....	3.6	268.4	0.3	*	*	*	*	*	*	*	*	*
Male.....	4.1	294.5	*	*	*	*	*	*	*	*	*	*
Female.....	3.2	241.1	0.3	*	*	*	*	*	*	*	*	*
Black, both sexes.....	12.6	700.6	1.0	*	*	*	*	*	*	*	*	*
Male.....	14.9	774.6	*	*	*	*	*	*	*	*	*	*
Female.....	10.6	623.6	*	*	*	*	*	*	*	*	*	*
Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)												
All races 3/, both sexes..	3.5	135.9	3.3	0.9	0.9	1.0	1.2	1.6	2.1	1.8	3.1	4.8
Male.....	3.7	140.7	3.4	0.9	1.1	1.2	1.2	1.7	2.4	1.9	3.4	5.0
Female.....	3.2	130.8	3.2	0.9	0.8	0.9	1.2	1.5	1.8	1.7	3.0	4.7
White, both sexes.....	3.3	132.7	3.1	0.9	0.9	1.0	1.2	1.6	2.2	1.8	3.2	4.9
Male.....	3.6	138.8	3.2	0.9	1.0	1.1	1.2	1.7	2.5	2.0	3.4	5.1
Female.....	3.1	126.4	3.0	0.8	0.8	0.9	1.2	1.5	1.9	1.6	3.0	4.8
Black, both sexes.....	4.5	158.9	4.4	1.2	1.2	1.5	1.4	1.5	1.8	1.9	3.3	*
Male.....	4.8	159.5	4.2	1.1	1.5	1.9	1.5	1.7	2.1	*	*	*
Female.....	4.2	158.3	4.6	1.2	0.9	1.0	1.3	1.4	1.5	2.2	3.0	*

Worktable 250R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race, and sex: United States, 2007

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2007. Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Symptoms, signs and abnormal clinical and laboratory findings, not elsewhere classified (R00-R99)												
All races 3/, both sexes..	11.1	85.0	1.4	0.3	1.4	2.2	3.3	5.0	6.8	12.9	44.8	251.4
Male.....	9.6	96.3	1.6	0.3	1.8	2.8	4.1	6.3	8.8	15.5	45.8	197.6
Female.....	12.6	73.1	1.3	0.2	0.9	1.5	2.5	3.7	4.9	10.7	44.1	277.0
White, both sexes.....	11.5	71.6	1.3	0.3	1.3	2.0	3.1	4.6	6.4	12.4	44.7	256.8
Male.....	9.6	82.8	1.5	0.3	1.8	2.7	3.9	5.7	8.3	14.9	45.6	201.3
Female.....	13.4	59.8	1.2	0.2	0.8	1.3	2.4	3.4	4.7	10.3	44.1	283.4
Black, both sexes.....	11.6	154.6	1.8	0.4	1.7	3.7	5.3	9.3	11.7	20.7	57.9	239.9
Male.....	11.8	166.1	1.8	*	2.3	4.8	6.9	12.3	16.8	27.5	63.6	182.7
Female.....	11.4	142.6	1.8	*	1.2	2.7	3.9	6.7	7.7	15.8	54.7	263.7
All other diseases (Residual) (D65-E07, E15-E34, E65-F99, G04-G12, G23-G25, G31-H93, K00-K22, K29-K31, K50-K66, K71-K72, K75-K76, K83-M99, N13.0-N13.5, N13.7-N13.9, N14, N15.0, N15.8-N15.9, N20-N23, N28-N39, N41-N64, N80-N98)												
All races 3/, both sexes..	79.0	31.4	3.6	2.2	4.6	7.8	16.4	37.6	70.0	149.0	469.6	1,701.9
Male.....	64.9	35.8	3.9	2.5	5.3	8.6	18.7	45.3	80.6	161.6	474.7	1,449.1
Female.....	92.7	26.9	3.4	1.8	3.9	6.8	14.1	30.1	60.1	138.3	466.0	1,822.1
White, both sexes.....	85.0	25.5	3.1	2.1	4.4	7.0	15.1	35.5	67.4	146.6	473.2	1,736.2
Male.....	68.7	28.9	3.4	2.3	5.2	8.1	17.6	43.3	77.3	157.7	475.5	1,479.5
Female.....	100.9	21.9	2.9	1.7	3.6	5.9	12.5	27.8	57.9	137.0	471.6	1,858.7
Black, both sexes.....	65.9	58.8	5.8	2.9	6.0	13.5	28.8	61.2	109.9	204.7	538.3	1,629.2
Male.....	59.0	67.1	5.8	3.6	6.3	13.8	30.7	71.5	132.9	242.2	589.5	1,363.2
Female.....	72.2	50.2	5.7	2.3	5.7	13.2	27.1	52.4	91.3	177.9	509.2	1,739.9

Worktable 250R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race, and sex: United States, 2007

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2007. Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Accidents (unintentional injuries) (V01-X59,Y85-Y86)												
All races 3/, both sexes..	41.0	30.2	9.6	5.5	37.4	36.9	39.2	46.3	37.3	45.2	105.5	286.7
Male.....	53.7	32.8	11.7	6.6	54.5	55.0	55.6	64.9	53.0	62.9	136.8	349.0
Female.....	28.7	27.4	7.5	4.3	19.3	18.0	22.8	28.3	22.6	30.2	83.9	257.0
White, both sexes.....	43.6	24.5	9.2	5.2	40.6	39.8	41.2	47.1	36.9	45.3	109.3	302.0
Male.....	56.4	27.1	11.3	6.2	59.1	58.7	57.7	65.2	51.7	62.2	140.5	366.1
Female.....	31.1	21.8	6.9	4.1	20.9	19.6	24.3	29.2	22.8	30.7	87.5	271.4
Black, both sexes.....	33.9	59.8	12.1	7.0	25.9	31.1	38.0	52.1	47.4	49.5	74.2	154.0
Male.....	48.5	60.8	13.8	8.4	38.3	49.5	58.6	79.5	76.6	78.7	106.4	198.0
Female.....	20.5	58.8	10.3	5.6	13.2	13.9	19.8	28.6	24.0	28.6	55.8	135.7
Transport accidents (V01-V99,Y85)												
All races 3/, both sexes..	15.5	3.0	3.5	3.4	25.7	18.4	15.8	16.4	14.8	16.5	22.9	24.1
Male.....	22.5	2.7	3.9	4.0	36.3	27.6	23.5	24.8	21.9	24.1	33.8	41.7
Female.....	8.8	3.3	3.1	2.9	14.5	8.8	8.1	8.3	8.1	10.1	15.4	15.7
White, both sexes.....	16.0	2.8	3.3	3.5	27.5	18.9	16.1	16.7	14.9	16.6	23.3	24.6
Male.....	23.1	2.7	3.8	4.0	38.7	28.0	23.6	25.0	21.9	24.0	33.9	41.9
Female.....	9.1	2.9	2.9	2.9	15.5	9.1	8.4	8.5	8.2	10.2	15.9	16.3
Black, both sexes.....	14.6	3.9	4.5	3.6	19.5	19.4	17.7	17.6	16.2	16.1	18.5	17.7
Male.....	22.7	*	4.5	4.3	28.5	31.8	29.3	28.8	26.7	26.6	34.2	39.1
Female.....	7.3	*	4.5	3.0	10.2	7.8	7.4	8.0	7.8	8.7	9.6	8.8

Worktable 250R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race, and sex: United States, 2007

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2007. Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Water, air and space, and other and unspecified transport accidents and their sequelae (V90-V99,Y85)												
All races 3/, both sexes..	0.6	*	*	0.1	0.4	0.6	0.6	1.0	1.1	0.9	0.6	0.6
Male.....	1.0	*	*	0.1	0.7	0.9	1.1	1.7	1.9	1.6	1.2	1.1
Female.....	0.2	*	*	*	0.2	0.2	0.2	0.3	0.4	0.3	*	*
White, both sexes.....	0.6	*	*	0.1	0.5	0.6	0.7	1.0	1.1	0.9	0.6	0.6
Male.....	1.1	*	*	*	0.7	0.9	1.2	1.7	1.9	1.6	1.2	1.3
Female.....	0.2	*	*	*	0.2	0.3	0.2	0.4	0.4	0.3	*	*
Black, both sexes.....	0.4	*	*	*	0.4	0.4	*	0.8	1.0	*	*	*
Male.....	0.8	*	*	*	0.6	*	*	1.7	1.7	*	*	*
Female.....	0.1	*	*	*	*	*	*	*	*	*	*	*
Nontransport accidents (W00-X59,Y86)												
All races 3/, both sexes..	25.5	27.2	6.1	2.0	11.7	18.5	23.4	29.9	22.5	28.7	82.6	262.6
Male.....	31.2	30.2	7.8	2.6	18.1	27.4	32.1	40.1	31.0	38.9	103.0	307.3
Female.....	19.9	24.1	4.3	1.5	4.9	9.3	14.7	20.0	14.5	20.1	68.5	241.4
White, both sexes.....	27.6	21.7	5.8	1.7	13.2	20.9	25.1	30.4	22.0	28.7	86.0	277.5
Male.....	33.3	24.4	7.5	2.3	20.4	30.7	34.0	40.3	29.8	38.2	106.6	324.1
Female.....	22.0	18.8	4.1	1.2	5.4	10.5	15.9	20.7	14.6	20.5	71.6	255.2
Black, both sexes.....	19.2	55.9	7.6	3.4	6.4	11.7	20.3	34.5	31.2	33.4	55.6	136.3
Male.....	25.8	57.8	9.3	4.2	9.8	17.6	29.3	50.7	49.9	52.1	72.2	158.9
Female.....	13.2	53.9	5.8	2.6	3.0	6.1	12.4	20.6	16.2	19.9	46.2	126.9

Worktable 250R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race, and sex: United States, 2007

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2007. Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Accidental drowning and submersion (W65-W74)												
All races 3/, both sexes..	1.1	1.3	2.8	0.6	1.5	0.9	1.0	1.1	1.0	1.0	1.4	1.6
Male.....	1.8	1.4	3.7	0.8	2.5	1.6	1.5	1.8	1.6	1.7	2.2	3.0
Female.....	0.5	1.3	1.8	0.3	0.4	0.3	0.4	0.4	0.4	0.4	0.9	0.9
White, both sexes.....	1.1	1.2	3.0	0.4	1.4	0.9	1.0	1.1	1.0	1.0	1.4	1.6
Male.....	1.7	*	3.8	0.6	2.4	1.5	1.5	1.8	1.6	1.6	2.3	3.2
Female.....	0.5	1.3	2.1	0.2	0.4	0.2	0.4	0.5	0.5	0.4	0.9	0.9
Black, both sexes.....	1.3	*	1.9	1.1	1.9	1.3	0.9	1.0	1.0	1.1	*	*
Male.....	2.2	*	3.0	1.4	3.4	2.3	1.6	1.8	2.0	*	*	*
Female.....	0.4	*	*	0.8	*	*	*	*	*	*	*	*
Accidental exposure to smoke, fire and flames (X00-X09)												
All races 3/, both sexes..	1.1	0.9	1.2	0.5	0.5	0.5	0.7	1.1	1.5	2.2	3.3	4.9
Male.....	1.3	0.9	1.5	0.6	0.6	0.7	0.9	1.4	2.1	2.8	4.0	6.5
Female.....	0.9	*	0.9	0.5	0.3	0.4	0.5	0.8	1.0	1.6	2.9	4.2
White, both sexes.....	1.0	0.7	0.8	0.4	0.5	0.5	0.6	1.0	1.4	1.9	3.0	4.1
Male.....	1.2	*	1.2	0.4	0.6	0.6	0.8	1.3	1.9	2.4	3.6	5.3
Female.....	0.8	*	0.5	0.4	0.3	0.4	0.5	0.8	0.9	1.5	2.6	3.5
Black, both sexes.....	1.9	*	3.0	1.3	0.5	0.9	1.3	1.9	2.7	4.9	7.9	15.5
Male.....	2.4	*	3.4	1.5	*	1.1	1.9	2.6	4.3	7.9	10.2	22.9
Female.....	1.5	*	2.6	1.2	*	0.8	0.8	1.3	1.4	2.8	6.6	12.4

Worktable 250R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race, and sex: United States, 2007

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2007. Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Accidental poisoning and exposure to noxious substances (X40-X49)												
All races 3/, both sexes..	9.9	*	0.2	0.2	7.4	14.0	17.6	20.5	9.5	3.1	2.7	3.5
Male.....	13.2	*	*	0.2	11.1	20.2	22.9	25.6	12.1	4.0	2.9	3.7
Female.....	6.7	*	*	0.2	3.5	7.7	12.2	15.6	7.2	2.4	2.6	3.4
White, both sexes.....	10.7	*	*	0.2	8.8	16.3	19.3	21.1	9.1	3.0	2.7	3.6
Male.....	14.2	*	*	0.2	13.3	23.3	25.0	25.9	11.0	3.6	2.9	3.9
Female.....	7.2	*	*	0.2	4.1	8.9	13.4	16.3	7.3	2.4	2.6	3.4
Black, both sexes.....	8.2	*	*	*	2.3	6.8	13.6	23.5	15.8	5.2	3.1	*
Male.....	11.2	*	*	*	3.0	9.8	18.1	32.8	25.3	9.0	*	*
Female.....	5.4	*	*	*	1.4	4.0	9.6	15.6	8.1	2.5	*	*
Other and unspecified nontransport accidents and their sequelae (W20-W31,W35-W64,W75-W99,X10-X39,X50-X59,Y86)												
All races 3/, both sexes..	5.7	23.9	1.6	0.6	1.4	2.0	2.6	4.0	5.0	8.9	24.5	85.8
Male.....	6.7	26.5	1.9	0.7	2.3	3.2	4.2	6.4	7.4	12.3	30.6	97.1
Female.....	4.6	21.2	1.2	0.4	0.4	0.6	1.0	1.6	2.7	5.9	20.3	80.4
White, both sexes.....	6.0	18.9	1.6	0.6	1.6	2.1	2.6	3.9	4.8	8.7	25.0	89.5
Male.....	7.0	21.5	1.9	0.7	2.6	3.4	4.1	6.3	7.3	12.0	31.2	101.4
Female.....	4.9	16.2	1.2	0.4	0.4	0.6	1.0	1.6	2.4	5.8	20.7	83.8
Black, both sexes.....	5.0	49.8	1.8	0.5	0.8	1.7	3.1	5.1	7.1	12.7	23.7	60.9
Male.....	6.2	50.7	2.1	*	1.1	2.7	5.4	8.5	10.6	18.2	30.5	67.1
Female.....	3.9	48.8	*	*	*	0.8	1.2	2.3	4.3	8.8	19.8	58.3

Worktable 250R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race, and sex: United States, 2007

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2007. Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)												
All races 3/, both sexes..	11.5	0.5	9.7	13.0	15.6	17.7	15.5	12.6	16.3	15.6
Male.....	18.3	0.6	15.9	20.7	23.8	27.0	24.3	22.5	34.3	41.8
Female.....	4.8	0.3	3.2	5.0	7.3	8.8	7.3	4.2	3.8	3.1
White, both sexes.....	12.9	0.4	10.4	14.2	17.5	20.1	17.3	13.9	17.5	16.9
Male.....	20.5	0.6	16.9	22.4	26.5	30.3	26.8	24.7	36.9	45.4
Female.....	5.4	0.2	3.4	5.5	8.3	10.1	8.3	4.7	3.9	3.3
Black, both sexes.....	4.9	0.6	6.0	8.4	7.5	6.1	4.7	3.9	4.7	*
Male.....	8.4	0.9	10.3	14.8	12.5	10.0	8.4	8.3	11.2	*
Female.....	1.7	*	1.6	2.4	3.0	2.7	1.7	*	*	*
Intentional self-harm (suicide) by discharge of firearms (X72-X74)												
All races 3/, both sexes..	5.8	0.1	4.5	5.7	6.7	8.0	8.5	8.8	12.2	11.0
Male.....	10.2	0.2	7.9	9.8	11.1	13.8	15.2	17.3	27.9	32.9
Female.....	1.4	*	0.8	1.4	2.3	2.5	2.3	1.5	1.4	0.6
White, both sexes.....	6.6	0.1	4.7	6.3	7.7	9.2	9.6	9.8	13.4	12.1
Male.....	11.6	0.2	8.2	10.7	12.5	15.6	16.9	19.2	30.3	36.3
Female.....	1.6	*	0.9	1.6	2.7	2.9	2.6	1.7	1.5	0.6
Black, both sexes.....	2.5	*	3.5	4.4	3.1	2.7	2.5	2.5	3.8	*
Male.....	4.6	*	6.5	8.1	5.7	4.8	5.0	5.7	9.4	*
Female.....	0.5	*	*	0.9	0.8	0.8	*	*	*	*

Worktable 250R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race, and sex: United States, 2007

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2007. Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Intentional self-harm (suicide) by other and unspecified means and their sequelae (*U03,X60-X71,X75-X84,Y87.0)												
All races 3/, both sexes..	5.7	0.3	5.3	7.3	8.9	9.7	7.0	3.8	4.1	4.6
Male.....	8.1	0.4	8.0	10.9	12.8	13.2	9.1	5.2	6.4	8.8
Female.....	3.4	0.2	2.4	3.6	5.0	6.3	5.0	2.7	2.5	2.5
White, both sexes.....	6.3	0.3	5.7	8.0	9.8	10.9	7.7	4.1	4.2	4.7
Male.....	8.9	0.3	8.7	11.7	14.0	14.6	9.9	5.5	6.6	9.1
Female.....	3.7	0.2	2.5	4.0	5.6	7.2	5.6	2.9	2.5	2.7
Black, both sexes.....	2.4	0.5	2.5	4.0	4.4	3.4	2.2	1.4	*	*
Male.....	3.8	0.7	3.8	6.7	6.8	5.2	3.4	2.6	*	*
Female.....	1.2	*	1.2	1.5	2.2	1.9	1.3	*	*	*
Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)												
All races 3/, both sexes..	6.1	8.3	2.4	0.9	13.1	11.7	7.1	4.9	3.0	2.1	2.1	1.5
Male.....	9.8	9.5	2.5	1.0	22.1	19.4	10.6	7.3	4.6	3.1	2.8	1.5
Female.....	2.5	7.0	2.3	0.7	3.5	3.7	3.5	2.5	1.5	1.3	1.5	1.4
White, both sexes.....	3.7	7.0	1.7	0.6	6.6	6.4	4.6	3.5	2.2	1.7	1.7	1.2
Male.....	5.4	7.7	1.8	0.7	10.5	9.9	6.4	5.0	3.2	2.5	2.1	*
Female.....	1.9	6.1	1.6	0.6	2.5	2.8	2.7	2.0	1.3	1.1	1.4	1.3
Black, both sexes.....	22.2	15.4	6.2	2.1	47.6	44.8	24.1	15.1	9.5	5.8	6.4	*
Male.....	39.7	19.2	6.5	2.9	85.3	82.5	41.2	25.4	17.8	10.3	13.1	*
Female.....	6.2	11.4	6.0	1.4	8.9	9.4	8.9	6.3	2.7	2.5	*	*

Worktable 250R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race, and sex: United States, 2007

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2007. Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Assault (homicide) by discharge of firearms (*U01.4,X93-X95)												
All races 3/, both sexes..	4.2	*	0.3	0.5	11.0	9.2	4.7	2.6	1.4	1.0	0.7	0.4
Male.....	7.2	*	0.3	0.7	19.2	16.1	7.6	4.2	2.2	1.4	0.9	*
Female.....	1.2	*	0.3	0.3	2.3	2.1	1.8	1.2	0.6	0.6	0.5	0.5
White, both sexes.....	2.2	*	0.2	0.3	5.0	4.6	2.8	1.8	1.0	0.8	0.6	*
Male.....	3.5	*	*	0.4	8.2	7.4	4.3	2.7	1.4	1.0	0.7	*
Female.....	0.9	*	*	0.2	1.6	1.6	1.4	1.0	0.6	0.6	0.5	*
Black, both sexes.....	17.4	*	0.7	1.4	43.2	38.8	17.8	8.8	4.8	2.5	1.9	*
Male.....	32.8	*	*	2.1	79.2	73.9	32.1	16.1	9.4	4.7	*	*
Female.....	3.2	*	*	0.7	6.3	5.8	5.0	2.6	1.2	*	*	*
Assault (homicide) by other and unspecified means and their sequelae (*U01.0-*U01.3,*U01.5-*U01.9,*U02,X85-X92,X96-Y09,Y87.1)												
All races 3/, both sexes..	1.9	7.9	2.1	0.4	2.1	2.5	2.3	2.2	1.6	1.2	1.4	1.0
Male.....	2.5	8.9	2.3	0.3	2.9	3.4	3.0	3.1	2.4	1.8	1.9	1.3
Female.....	1.3	6.8	2.0	0.4	1.2	1.6	1.7	1.4	0.9	0.6	1.0	0.9
White, both sexes.....	1.5	6.6	1.5	0.3	1.6	1.9	1.7	1.7	1.2	0.9	1.1	0.8
Male.....	1.9	7.3	1.6	0.2	2.2	2.5	2.1	2.4	1.8	1.4	1.4	*
Female.....	1.0	6.0	1.4	0.3	0.9	1.2	1.4	1.1	0.8	0.5	0.9	0.8
Black, both sexes.....	4.8	14.7	5.5	0.7	4.4	6.0	6.3	6.3	4.6	3.3	4.5	*
Male.....	6.8	18.4	5.9	0.8	6.1	8.6	9.1	9.3	8.4	5.6	8.6	*
Female.....	2.9	10.8	5.1	0.7	2.6	3.6	3.9	3.7	1.6	*	*	*

Worktable 250R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race, and sex: United States, 2007

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2007. Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Legal intervention (Y35,Y89.0)												
All races 3/, both sexes..	0.1	*	*	*	0.2	0.3	0.2	0.1	*	*	*	*
Male.....	0.3	*	*	*	0.4	0.6	0.4	0.3	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*	*	*
White, both sexes.....	0.1	*	*	*	0.1	0.3	0.2	0.1	*	*	*	*
Male.....	0.2	*	*	*	0.3	0.5	0.4	0.3	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*	*	*
Black, both sexes.....	0.3	*	*	*	0.7	0.8	*	*	*	*	*	*
Male.....	0.6	*	*	*	1.3	1.6	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*	*	*
Events of undetermined intent (Y10-Y34,Y87.2,Y89.9)												
All races 3/, both sexes..	1.8	2.1	0.3	0.1	1.4	2.1	2.9	3.4	1.8	0.9	1.0	1.3
Male.....	2.3	2.5	0.3	0.2	2.0	2.8	3.6	4.0	2.3	1.2	1.5	1.6
Female.....	1.3	1.6	0.3	*	0.6	1.5	2.2	2.9	1.4	0.7	0.7	1.1
White, both sexes.....	1.8	1.6	0.3	0.1	1.4	2.4	3.0	3.4	1.7	0.9	0.9	1.3
Male.....	2.3	2.0	*	0.1	2.1	3.0	3.7	3.8	2.1	1.1	1.2	1.7
Female.....	1.4	1.3	*	*	0.7	1.7	2.3	3.0	1.4	0.7	0.7	1.1
Black, both sexes.....	1.9	4.1	*	*	1.0	1.6	3.0	4.3	2.9	1.5	1.9	*
Male.....	2.8	*	*	*	1.8	2.2	4.3	6.1	4.7	2.6	*	*
Female.....	1.1	*	*	*	*	1.0	1.8	2.8	1.5	*	*	*

Worktable 250R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race, and sex: United States, 2007

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2007. Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Discharge of firearms, undetermined intent (Y22-Y24)												
All races 3/, both sexes..	0.1	*	*	*	0.2	0.1	0.1	0.1	0.1	*	*	*
Male.....	0.2	*	*	*	0.3	0.2	0.1	0.1	0.2	*	*	*
Female.....	0.0	*	*	*	*	*	*	*	*	*	*	*
White, both sexes.....	0.1	*	*	*	0.1	0.1	0.1	0.1	0.1	*	*	*
Male.....	0.1	*	*	*	0.2	0.2	0.1	0.2	0.2	*	*	*
Female.....	0.0	*	*	*	*	*	*	*	*	*	*	*
Black, both sexes.....	0.1	*	*	*	0.5	*	*	*	*	*	*	*
Male.....	0.2	*	*	*	0.8	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*	*	*
Other and unspecified events of undetermined intent and their sequelae (Y10-Y21, Y25-Y34, Y87.2, Y89.9)												
All races 3/, both sexes..	1.7	2.1	0.3	0.1	1.2	2.0	2.8	3.3	1.7	0.9	1.0	1.3
Male.....	2.1	2.5	0.3	0.1	1.7	2.6	3.5	3.8	2.1	1.1	1.4	1.5
Female.....	1.3	1.6	0.3	*	0.6	1.5	2.1	2.8	1.4	0.7	0.7	1.1
White, both sexes.....	1.7	1.6	0.3	0.1	1.3	2.3	2.9	3.3	1.6	0.8	0.9	1.3
Male.....	2.1	2.0	*	*	1.9	2.8	3.6	3.7	1.9	1.0	1.2	1.6
Female.....	1.4	1.3	*	*	0.7	1.7	2.3	3.0	1.4	0.7	0.7	1.1
Black, both sexes.....	1.8	4.1	*	*	0.5	1.4	2.9	4.2	2.8	1.4	*	*
Male.....	2.5	*	*	*	0.9	1.9	4.2	6.0	4.6	*	*	*
Female.....	1.1	*	*	*	*	1.0	1.8	2.7	1.4	*	*	*

Worktable 250R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race, and sex: United States, 2007

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2007. Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Operations of war and their sequelae (Y36,Y89.1)												
All races 3/, both sexes..	0.0	*	*	*	*	*	*	*	*	*	*	*
Male.....	0.0	*	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*	*	*
White, both sexes.....	0.0	*	*	*	*	*	*	*	*	*	*	*
Male.....	0.0	*	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*	*	*
Black, both sexes.....	*	*	*	*	*	*	*	*	*	*	*	*
Male.....	*	*	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*	*	*
Complications of medical and surgical care (Y40-Y84,Y88)												
All races 3/, both sexes..	0.9	0.5	0.1	0.1	0.1	0.2	0.3	0.6	1.2	2.5	5.5	7.9
Male.....	0.8	*	*	*	*	0.1	0.3	0.6	1.3	2.7	6.4	9.8
Female.....	0.9	*	*	*	*	0.2	0.4	0.6	1.2	2.3	4.9	6.9
White, both sexes.....	0.9	*	*	*	0.1	0.2	0.3	0.5	1.2	2.4	5.3	7.6
Male.....	0.8	*	*	*	*	0.1	0.3	0.5	1.2	2.6	6.3	9.5
Female.....	0.9	*	*	*	*	0.2	0.3	0.5	1.1	2.2	4.5	6.7
Black, both sexes.....	1.1	*	*	*	*	*	0.6	1.2	2.0	4.6	8.7	12.2
Male.....	1.0	*	*	*	*	*	*	0.9	2.5	5.3	9.9	*
Female.....	1.2	*	*	*	*	*	0.7	1.4	1.7	4.0	8.1	10.6

Worktable 250R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race, and sex: United States, 2007

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2007. Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Alcohol-induced causes (E24.4,F10,G31.2,G62.1,G72.1,I42.6,K29.2,K70,K85.2,K86.0,R78.0,X45,X65,Y15) 4/												
All races 3/, both sexes..	7.7	*	*	*	0.5	1.8	7.3	18.3	20.2	15.5	9.7	4.6
Male.....	11.7	*	*	*	0.7	2.7	10.2	27.4	32.5	26.4	17.8	9.6
Female.....	3.8	*	*	*	0.2	0.9	4.3	9.5	8.6	6.2	4.1	2.2
White, both sexes.....	8.2	*	*	*	0.5	1.9	7.7	19.0	20.5	15.7	9.8	4.6
Male.....	12.4	*	*	*	0.7	2.8	10.7	28.3	32.9	26.3	17.8	9.8
Female.....	4.0	*	*	*	0.2	0.9	4.5	9.8	8.8	6.4	4.3	2.1
Black, both sexes.....	5.6	*	*	*	*	1.0	4.6	14.8	20.9	16.9	8.9	5.5
Male.....	8.9	*	*	*	*	1.4	6.6	23.5	36.6	34.1	20.6	*
Female.....	2.6	*	*	*	*	*	2.9	7.3	8.2	4.6	*	*
Drug-induced causes (D52.1,D59.0,D59.2,D61.1,D64.2,E06.4,E16.0,E23.1,E24.2,E27.3,E66.1,F11.0-F11.5,F11.7-F11.9,F12.0-F12.5,F12.7-F12.9,F13.0-F13.5,F13.7-F13.9,F14.0-F14.5,F14.7-F14.9,F15.0-F15.5,F15.7-F15.9,F16.0-F16.5,F16.7-F16.9,F17.0-F17.5,F17.7-F17.9,F18.0-F18.5,F18.7-F18.9,F19.0-F19.5,F19.7-F19.9,G21.1,G24.0,G25.1,G25.4,G25.6,G44.4,G62.0,G72.0,I95.2,J70.2-J70.4,K85.3,L10.5,L27.0-L27.1,M10.2,M32.0,M80.4,M81.4,M83.5,M87.1,R50.2,R78.1-R78.5,X40-X44,X60-X64,X85,Y10-Y14) 4/												
All races 3/, both sexes..	12.7	0.8	0.3	0.2	8.6	17.1	22.4	26.9	13.6	4.7	3.9	4.8
Male.....	16.1	*	0.2	0.2	12.6	23.5	27.9	31.7	16.1	5.3	3.8	5.5
Female.....	9.5	*	0.3	0.2	4.4	10.5	16.9	22.2	11.3	4.1	3.9	4.4
White, both sexes.....	13.7	*	0.2	0.2	10.3	19.9	24.5	27.7	13.3	4.7	4.0	5.0
Male.....	17.2	*	*	0.2	15.1	27.2	30.2	32.0	14.8	5.0	3.9	5.8
Female.....	10.3	*	*	0.2	5.2	12.2	18.6	23.6	11.8	4.5	4.0	4.5
Black, both sexes.....	10.5	*	0.7	*	2.4	8.4	18.3	30.5	20.4	6.0	3.6	*
Male.....	14.2	*	*	*	3.4	11.5	23.8	41.8	32.6	10.9	*	*
Female.....	7.1	*	*	*	1.5	5.5	13.4	20.8	10.7	2.5	3.8	*

Worktable 250R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race, and sex: United States, 2007

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2007. Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Injury by firearms (*U01.4,W32-W34,X72-X74,X93-X95,Y22-Y24,Y35.0) 4/												
All races 3/, both sexes..	10.4	*	0.4	0.8	16.2	15.5	11.9	11.1	10.2	10.0	13.2	11.6
Male.....	18.2	*	0.5	1.2	28.5	27.0	19.5	18.6	17.9	19.1	29.5	33.5
Female.....	2.7	*	0.4	0.4	3.2	3.6	4.2	3.7	3.0	2.3	2.0	1.1
White, both sexes.....	9.2	*	0.3	0.6	10.3	11.3	11.0	11.4	10.9	10.9	14.3	12.7
Male.....	15.8	*	0.4	0.9	17.5	18.9	17.6	19.0	19.0	20.7	31.7	36.9
Female.....	2.7	*	*	0.3	2.6	3.3	4.2	4.0	3.3	2.4	2.1	1.1
Black, both sexes.....	20.6	*	1.0	1.8	48.6	44.5	21.4	11.8	7.5	5.3	5.8	*
Male.....	38.9	*	*	2.7	89.1	84.8	38.9	21.5	14.8	10.8	14.2	*
Female.....	3.9	*	*	0.8	7.1	6.9	5.9	3.5	1.7	*	*	*

* Figure does not meet standards of reliability or precision.

0.0 Quantity more than zero but less than 0.05.

... Category not applicable.

1/ Figures for age not stated included in "All ages" but not distributed among age groups.

2/ Death rates for "Under 1 year" (based on population estimates) differ from infant mortality rates (based on live births).

3/ Includes races other than white and black.

4/ Included in selected categories.

Source: CDC/NCHS, National Vital Statistics System, Mortality.