

HIST293. AGE-ADJUSTED DEATH RATES FOR APPROXIMATELY 64 SELECTED CAUSES, BY RACE AND SEX: UNITED STATES, 1950-59

[Based on age-specific death rates per 100,000 population in specified group. Computed by the direct method, using as the standard population the age distribution of the total population of the United States as projected for 2000. Numbers after causes of death are category numbers of the seventh revision, International Classification of Diseases]

CAUSE OF DEATH CODE AND YEAR	-- ALL RACES --			-- WHITE --			--- ALL OTHER ---		
	BOTH	MALE	FEMALE	BOTH	MALE	FEMALE	BOTH	MALE	FEMALE
ALL CAUSES									
1959	1,317.3	1,573.4	1,093.5	1,291.9	1,552.1	1,065.3	1,504.4	1,724.7	1,305.0
1958	1,343.4	1,596.3	1,120.9	1,316.3	1,573.1	1,090.8	1,552.1	1,779.2	1,354.6
1957	1,356.7	1,606.7	1,134.5	1,328.7	1,581.7	1,104.1	1,571.7	1,797.0	1,374.6
1956	1,333.7	1,575.5	1,117.5	1,309.0	1,554.4	1,089.9	1,519.2	1,724.1	1,331.9
1955	1,332.3	1,565.2	1,121.9	1,308.3	1,544.7	1,095.4	1,510.2	1,707.5	1,326.7
1954	1,314.8	1,540.7	1,109.7	1,288.6	1,516.3	1,081.8	1,514.8	1,726.6	1,326.3
1953	1,385.6	1,619.0	1,172.6	1,354.6	1,589.0	1,141.0	1,641.2	1,863.1	1,434.3
1952	1,394.6	1,620.8	1,186.7	1,361.7	1,588.5	1,153.3	1,671.3	1,894.1	1,465.0
1951	1,423.5	1,651.7	1,213.4	1,391.0	1,621.7	1,179.1	1,698.2	1,903.4	1,508.3
1950	1,446.0	1,674.2	1,236.0	1,410.8	1,642.5	1,198.0	1,757.6	1,949.5	1,574.1
TUBERCULOSIS, ALL FORMS (001-019)									
1959	8.0	12.2	4.3	6.7	10.5	3.3	20.6	29.1	12.9
1958	8.7	13.2	4.6	7.2	11.2	3.6	23.2	33.9	13.5
1957	9.5	14.5	5.0	8.0	12.5	4.0	24.3	34.5	14.9
1956	10.2	15.2	5.7	8.5	13.1	4.5	26.3	36.8	16.5
1955	10.9	16.0	6.2	9.1	13.7	4.9	28.7	39.2	18.7
1954	12.1	17.8	7.0	10.1	15.2	5.4	31.9	43.4	21.0
1953	14.6	21.0	8.6	11.9	17.7	6.6	39.4	53.4	26.0
1952	18.2	25.9	11.1	14.8	21.5	8.5	50.8	68.2	34.0
1951	23.0	32.2	14.4	18.7	26.9	11.0	63.4	83.4	44.1
1950	25.5	35.1	16.4	20.5	29.3	12.1	72.5	91.0	54.0

Source: CDC/NCHS, National Vital Statistics System, Mortality.

HIST293. AGE-ADJUSTED DEATH RATES FOR APPROXIMATELY 64 SELECTED CAUSES, BY RACE AND SEX: UNITED STATES, 1950-59

[Based on age-specific death rates per 100,000 population in specified group. Computed by the direct method, using as the standard population the age distribution of the total population of the United States as projected for 2000. Numbers after causes of death are category numbers of the seventh revision, International Classification of Diseases]

CAUSE OF DEATH CODE AND YEAR	-- ALL RACES --			-- WHITE --			--- ALL OTHER ---		
	BOTH	MALE	FEMALE	BOTH	MALE	FEMALE	BOTH	MALE	FEMALE
TUBERCULOSIS OF RESPIRATORY SYSTEM (001-008)									
1959	7.4	11.6	3.8	6.3	10.1	3.1	18.6	26.6	11.2
1958	8.1	12.5	4.2	6.8	10.7	3.4	20.9	30.9	11.8
1957	8.9	13.7	4.6	7.6	11.9	3.7	21.9	31.4	13.2
1956	9.5	14.4	5.2	8.0	12.4	4.1	23.6	33.7	14.2
1955	10.1	15.1	5.6	8.5	13.0	4.4	25.4	35.3	16.0
1954	11.2	16.7	6.3	9.5	14.4	5.0	28.6	39.7	18.1
1953	13.4	19.7	7.7	11.2	16.8	6.0	35.2	48.1	22.7
1952	16.9	24.2	10.0	13.9	20.4	7.7	45.3	61.5	29.6
1951	21.3	30.2	13.0	17.5	25.4	10.0	57.5	76.0	39.5
1950	23.7	32.9	14.8	19.2	27.7	11.0	65.7	83.0	48.5
TUBERCULOSIS, OTHER FORMS (010-019)									
1959	0.5	0.7	0.4	0.4	0.5	0.3	2.0	2.5	1.7
1958	0.5	0.7	0.4	0.3	0.5	0.2	2.4	2.9	1.8
1957	0.6	0.8	0.4	0.4	0.5	0.3	2.4	3.2	1.7
1956	0.7	0.9	0.5	0.5	0.6	0.3	2.7	3.0	2.3
1955	0.8	1.0	0.6	0.6	0.7	0.4	3.3	3.9	2.7
1954	0.9	1.1	0.7	0.6	0.8	0.4	3.3	3.7	2.9
1953	1.1	1.4	0.9	0.8	0.9	0.6	4.3	5.4	3.3
1952	1.4	1.6	1.1	0.9	1.1	0.7	5.5	6.7	4.3
1951	1.7	2.0	1.4	1.2	1.4	1.0	6.0	7.4	4.6
1950	1.9	2.2	1.5	1.3	1.6	1.1	6.8	8.0	5.6

Source: CDC/NCHS, National Vital Statistics System, Mortality.

HIST293. AGE-ADJUSTED DEATH RATES FOR APPROXIMATELY 64 SELECTED CAUSES, BY RACE AND SEX: UNITED STATES, 1950-59

[Based on age-specific death rates per 100,000 population in specified group. Computed by the direct method, using as the standard population the age distribution of the total population of the United States as projected for 2000. Numbers after causes of death are category numbers of the seventh revision, International Classification of Diseases]

CAUSE OF DEATH CODE AND YEAR	-- ALL RACES --			-- WHITE --			--- ALL OTHER ---		
	BOTH	MALE	FEMALE	BOTH	MALE	FEMALE	BOTH	MALE	FEMALE
SYPHILIS AND ITS SEQUELAE (020-029)									
1959	2.1	3.2	1.2	1.6	2.5	0.9	7.4	10.5	4.6
1958	2.4	3.7	1.3	1.7	2.7	0.9	9.9	14.4	5.8
1957	2.7	4.1	1.5	2.0	3.1	1.0	10.6	15.3	6.3
1956	2.8	4.2	1.5	2.0	3.2	1.0	10.6	15.2	6.3
1955	2.8	4.3	1.4	2.0	3.1	1.0	11.7	17.2	6.5
1954	3.7	5.5	1.9	2.7	4.2	1.4	13.4	19.5	7.6
1953	4.0	6.1	2.1	2.9	4.5	1.5	15.7	22.7	8.9
1952	4.4	6.8	2.2	3.3	5.1	1.5	16.8	24.4	9.4
1951	4.9	7.4	2.5	3.6	5.6	1.7	18.7	26.8	10.8
1950	6.1	9.2	3.3	4.6	7.0	2.3	22.6	32.4	13.0
DYSENTERY, ALL FORMS (045-048)									
1959	0.2	0.2	0.2	0.2	0.2	0.2	0.6	0.7	0.5
1958	0.2	0.2	0.2	0.1	0.2	0.1	0.6	0.7	0.4
1957	0.2	0.3	0.2	0.2	0.2	0.2	0.7	0.8	0.6
1956	0.3	0.3	0.2	0.2	0.2	0.2	0.8	1.0	0.6
1955	0.3	0.3	0.3	0.2	0.3	0.2	1.0	1.0	1.0
1954	0.4	0.4	0.3	0.3	0.3	0.3	1.2	1.7	0.9
1953	0.5	0.5	0.5	0.4	0.4	0.4	1.4	1.6	1.2
1952	0.6	0.6	0.5	0.4	0.5	0.4	1.5	1.8	1.3
1951	0.6	0.6	0.6	0.5	0.6	0.5	1.4	1.4	1.5
1950	0.5	0.6	0.5	0.5	0.5	0.4	1.4	1.8	1.0

Source: CDC/NCHS, National Vital Statistics System, Mortality.

HIST293. AGE-ADJUSTED DEATH RATES FOR APPROXIMATELY 64 SELECTED CAUSES, BY RACE AND SEX: UNITED STATES, 1950-59

[Based on age-specific death rates per 100,000 population in specified group. Computed by the direct method, using as the standard population the age distribution of the total population of the United States as projected for 2000. Numbers after causes of death are category numbers of the seventh revision, International Classification of Diseases]

CAUSE OF DEATH CODE AND YEAR	-- ALL RACES --			-- WHITE --			--- ALL OTHER ---		
	BOTH	MALE	FEMALE	BOTH	MALE	FEMALE	BOTH	MALE	FEMALE
SCARLET FEVER AND STREPTOCOCCAL SORE THROAT (050,051)									
1959	0.1	0.1	0.1	0.1	0.1	0.1	0.1	*	*
1958	0.1	0.1	0.1	0.1	0.1	0.1	0.1	*	*
1957	0.1	0.1	0.1	0.1	0.1	0.2	0.2	0.3	*
1956	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2	*
1955	0.2	0.2	0.1	0.2	0.1	0.1	0.2	0.2	*
1954	0.2	0.2	0.2	0.2	0.2	0.2	0.2	*	0.2
1953	0.2	0.2	0.2	0.2	0.2	0.2	0.2	*	*
1952	0.2	0.2	0.2	0.2	0.2	0.2	0.3	0.3	0.2
1951	0.2	0.3	0.2	0.2	0.3	0.2	0.3	0.3	0.2
1950	0.2	0.2	0.3	0.2	0.2	0.2	0.3	0.2	0.4
SEPTICEMIA (053)									
1959	1.1	1.3	0.9	1.0	1.1	0.8	2.0	2.4	1.6
1958	1.0	1.2	0.9	1.0	1.1	0.8	1.7	2.0	1.4
1957	0.8	1.0	0.6	0.7	0.9	0.5	1.3	1.7	1.0
1956	0.7	0.8	0.6	0.6	0.7	0.5	1.1	1.2	1.0
1955	0.6	0.7	0.5	0.5	0.6	0.4	0.9	0.9	0.9
1954	0.5	0.6	0.5	0.5	0.6	0.5	0.8	0.9	0.8
1953	0.5	0.6	0.4	0.4	0.5	0.3	0.9	1.2	0.6
1952	0.4	0.5	0.4	0.4	0.5	0.3	0.7	0.8	0.7
1951	0.4	0.5	0.4	0.4	0.5	0.3	0.8	0.8	0.9
1950	0.4	0.5	0.4	0.4	0.4	0.3	0.8	0.7	0.9

\* Figure does not meet standard of reliability or precision

Source: CDC/NCHS, National Vital Statistics System, Mortality.

HIST293. AGE-ADJUSTED DEATH RATES FOR APPROXIMATELY 64 SELECTED CAUSES, BY RACE AND SEX: UNITED STATES, 1950-59

[Based on age-specific death rates per 100,000 population in specified group. Computed by the direct method, using as the standard population the age distribution of the total population of the United States as projected for 2000. Numbers after causes of death are category numbers of the seventh revision, International Classification of Diseases]

CAUSE OF DEATH CODE AND YEAR	-- ALL RACES --			-- WHITE --			--- ALL OTHER ---		
	BOTH	MALE	FEMALE	BOTH	MALE	FEMALE	BOTH	MALE	FEMALE
DIPHTHERIA (055)									
1959	0.0	0.0	0.0	0.0	0.0	0.0	0.1	*	*
1958	0.0	0.0	0.0	0.0	0.0	0.0	0.1	*	*
1957	0.0	0.0	0.0	0.0	0.0	0.0	0.1	*	*
1956	0.0	0.1	0.0	0.0	0.0	0.0	0.1	0.2	*
1955	0.1	0.1	0.1	0.0	0.1	0.0	0.2	0.2	0.2
1954	0.1	0.1	0.1	0.0	0.0	0.0	0.2	0.2	0.2
1953	0.1	0.1	0.1	0.0	0.1	0.0	0.2	0.2	0.2
1952	0.1	0.1	0.1	0.1	0.1	0.1	0.2	0.2	0.2
1951	0.2	0.2	0.1	0.1	0.2	0.1	0.3	0.4	0.2
1950	0.2	0.2	0.2	0.2	0.2	0.2	0.3	0.3	0.3
WHOOPING COUGH (056)									
1959	0.1	0.1	0.1	0.0	0.0	0.1	0.3	0.3	0.4
1958	0.1	0.1	0.1	0.0	0.0	0.0	0.2	0.2	0.2
1957	0.1	0.1	0.1	0.0	0.0	0.0	0.2	0.2	0.3
1956	0.1	0.1	0.1	0.0	0.0	0.0	0.4	0.4	0.5
1955	0.2	0.1	0.2	0.1	0.1	0.1	0.7	0.6	0.7
1954	0.1	0.1	0.2	0.1	0.1	0.1	0.4	0.4	0.5
1953	0.1	0.1	0.1	0.1	0.1	0.1	0.3	0.3	0.3
1952	0.2	0.2	0.2	0.1	0.1	0.1	0.6	0.6	0.6
1951	0.4	0.4	0.4	0.2	0.2	0.2	1.6	1.6	1.6
1950	0.5	0.4	0.6	0.3	0.3	0.4	1.4	1.3	1.6

\* Figure does not meet standard of reliability or precision

Source: CDC/NCHS, National Vital Statistics System, Mortality.

HIST293. AGE-ADJUSTED DEATH RATES FOR APPROXIMATELY 64 SELECTED CAUSES, BY RACE AND SEX: UNITED STATES, 1950-59

[Based on age-specific death rates per 100,000 population in specified group. Computed by the direct method, using as the standard population the age distribution of the total population of the United States as projected for 2000. Numbers after causes of death are category numbers of the seventh revision, International Classification of Diseases]

CAUSE OF DEATH CODE AND YEAR	-- ALL RACES --			-- WHITE --			--- ALL OTHER ---		
	BOTH	MALE	FEMALE	BOTH	MALE	FEMALE	BOTH	MALE	FEMALE
MENINGOCOCCAL INFECTIONS (057)									
1959	0.3	0.3	0.3	0.3	0.3	0.2	0.5	0.4	0.5
1958	0.3	0.3	0.3	0.3	0.3	0.3	0.5	0.5	0.4
1957	0.3	0.4	0.3	0.3	0.4	0.3	0.5	0.5	0.5
1956	0.4	0.4	0.3	0.4	0.4	0.3	0.5	0.5	0.4
1955	0.4	0.5	0.4	0.4	0.4	0.3	0.7	0.6	0.7
1954	0.5	0.5	0.4	0.4	0.5	0.4	0.6	0.8	0.5
1953	0.6	0.7	0.5	0.6	0.7	0.5	0.9	1.1	0.8
1952	0.7	0.7	0.6	0.6	0.7	0.6	0.9	1.0	0.9
1951	0.6	0.6	0.5	0.5	0.6	0.5	0.8	1.0	0.6
1950	0.5	0.6	0.4	0.5	0.6	0.4	0.7	0.9	0.5
ACUTE POLIOMYELITIS (080)									
1959	0.2	0.3	0.1	0.2	0.3	0.1	0.3	0.5	0.2
1958	0.1	0.2	0.1	0.1	0.2	0.1	0.2	0.2	*
1957	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.2	*
1956	0.3	0.3	0.3	0.3	0.3	0.3	0.3	0.4	0.3
1955	0.6	0.6	0.5	0.6	0.7	0.5	0.2	0.2	0.2
1954	0.8	0.9	0.6	0.8	0.9	0.7	0.3	0.3	0.2
1953	0.8	1.0	0.7	0.9	1.0	0.7	0.3	0.4	0.3
1952	1.8	2.1	1.5	1.9	2.3	1.6	0.6	0.8	0.5
1951	0.9	1.0	0.7	0.9	1.1	0.7	0.4	0.5	0.4
1950	1.1	1.3	0.9	1.2	1.5	0.9	0.5	0.5	0.4

\* Figure does not meet standard of reliability or precision

Source: CDC/NCHS, National Vital Statistics System, Mortality.

HIST293. AGE-ADJUSTED DEATH RATES FOR APPROXIMATELY 64 SELECTED CAUSES, BY RACE AND SEX: UNITED STATES, 1950-59

[Based on age-specific death rates per 100,000 population in specified group. Computed by the direct method, using as the standard population the age distribution of the total population of the United States as projected for 2000. Numbers after causes of death are category numbers of the seventh revision, International Classification of Diseases]

CAUSE OF DEATH CODE AND YEAR	-- ALL RACES --			-- WHITE --			--- ALL OTHER ---		
	BOTH	MALE	FEMALE	BOTH	MALE	FEMALE	BOTH	MALE	FEMALE
MEASLES (085)									
1959	0.1	0.1	0.2	0.1	0.1	0.1	0.2	0.2	0.3
1958	0.2	0.2	0.2	0.2	0.2	0.2	0.3	0.4	0.3
1957	0.2	0.2	0.1	0.1	0.2	0.1	0.2	0.2	0.2
1956	0.2	0.2	0.2	0.2	0.2	0.2	0.4	0.4	0.3
1955	0.1	0.1	0.1	0.1	0.1	0.1	0.3	0.2	0.3
1954	0.2	0.2	0.2	0.2	0.2	0.2	0.3	0.4	0.4
1953	0.2	0.2	0.2	0.2	0.2	0.2	0.3	0.2	0.4
1952	0.3	0.3	0.3	0.3	0.2	0.3	0.5	0.5	0.6
1951	0.3	0.3	0.3	0.3	0.3	0.3	0.5	0.5	0.5
1950	0.2	0.2	0.2	0.2	0.2	0.2	0.4	0.3	0.4
OTHER INFECTIVE AND PARASITIC DISEASES									
1959	3.3	3.7	3.0	3.0	3.3	2.7	6.2	7.0	5.5
1958	3.2	3.6	2.9	3.0	3.3	2.7	5.4	5.8	4.9
1957	2.9	3.3	2.6	2.7	3.1	2.3	5.0	5.6	4.4
1956	2.8	3.2	2.5	2.6	2.9	2.3	5.0	5.5	4.5
1955	2.7	3.0	2.5	2.5	2.7	2.3	4.8	5.2	4.4
1954	2.8	3.1	2.5	2.5	2.8	2.3	5.2	6.1	4.3
1953	2.8	3.3	2.4	2.5	3.0	2.1	5.3	5.9	4.7
1952	3.0	3.3	2.7	2.7	3.0	2.4	5.3	5.9	4.7
1951	2.9	3.2	2.6	2.6	2.9	2.3	5.2	5.9	4.6
1950	2.8	3.1	2.4	2.5	2.9	2.1	5.4	5.8	5.0

NOTE: Other infective and parasitic diseases include ICD Nos. 030-044, 049, 052-054, 058-074, 081-084, and 086-138 in 1959; and ICD Nos. 030-039, 041, 042, 044, 049, 052-.054, 059-074, 081-083, 086-096, 120-138 in 1950-1958

Source: CDC/NCHS, National Vital Statistics System, Mortality.

HIST293. AGE-ADJUSTED DEATH RATES FOR APPROXIMATELY 64 SELECTED CAUSES, BY RACE AND SEX: UNITED STATES, 1950-59

[Based on age-specific death rates per 100,000 population in specified group. Computed by the direct method, using as the standard population the age distribution of the total population of the United States as projected for 2000. Numbers after causes of death are category numbers of the seventh revision, International Classification of Diseases]

CAUSE OF DEATH CODE AND YEAR	-- ALL RACES --			-- WHITE --			--- ALL OTHER ---		
	BOTH	MALE	FEMALE	BOTH	MALE	FEMALE	BOTH	MALE	FEMALE
MALIGNANT NEOPLASMS (ICD NOS. 140-205)									
1959	191.6	221.8	167.3	191.1	221.5	166.8	191.5	221.7	164.8
1958	192.0	219.2	169.8	191.7	219.7	169.0	188.5	209.7	169.9
1957	195.4	222.6	173.0	195.2	223.0	172.5	190.5	213.7	170.0
1956	195.6	222.0	173.9	195.5	222.7	173.2	190.2	209.4	172.8
1955	194.5	217.7	175.2	194.9	218.9	175.1	182.3	198.3	167.7
1954	194.3	216.3	176.1	194.6	217.2	175.9	183.6	200.7	168.5
1953	194.6	214.4	178.0	195.4	216.1	178.1	178.2	188.6	168.4
1952	194.2	211.8	179.5	195.0	213.5	179.6	177.5	186.5	169.4
1951	192.0	207.6	179.1	193.1	209.8	179.3	172.2	177.4	167.9
1950	193.9	208.1	182.3	194.6	210.0	182.0	177.4	180.0	174.8

MALIGNANT NEOPLASM OF BUCCAL CAVITY AND PHARYNX (140-148)

1959	4.5	7.5	1.9	4.5	7.5	1.9	4.2	6.7	1.9
1958	4.4	7.3	2.0	4.4	7.3	2.0	4.0	6.2	1.9
1957	4.6	7.7	1.9	4.6	7.8	1.9	4.4	7.1	2.0
1956	4.5	7.5	1.9	4.5	7.7	1.8	4.0	6.0	2.1
1955	4.5	7.6	1.9	4.6	7.8	1.9	3.8	5.6	2.0
1954	4.6	7.6	2.0	4.6	7.7	1.9	3.7	5.7	1.8
1953	4.5	7.4	1.9	4.5	7.6	1.9	3.4	5.3	1.6
1952	4.5	7.4	2.0	4.5	7.5	2.0	3.8	5.5	2.1
1951	4.5	7.4	1.9	4.5	7.5	1.9	3.7	5.5	1.9
1950	4.9	8.2	2.0	5.0	8.4	2.0	4.1	5.8	2.4

Source: CDC/NCHS, National Vital Statistics System, Mortality.



HIST293. AGE-ADJUSTED DEATH RATES FOR APPROXIMATELY 64 SELECTED CAUSES, BY RACE AND SEX: UNITED STATES, 1950-59

[Based on age-specific death rates per 100,000 population in specified group. Computed by the direct method, using as the standard population the age distribution of the total population of the United States as projected for 2000. Numbers after causes of death are category numbers of the seventh revision, International Classification of Diseases]

CAUSE OF DEATH CODE AND YEAR	-- ALL RACES --			-- WHITE --			--- ALL OTHER ---		
	BOTH	MALE	FEMALE	BOTH	MALE	FEMALE	BOTH	MALE	FEMALE
MALIGNANT NEOPLASM OF DIGESTIVE ORGANS AND PERITONEUM, NOT SPECIFIED AS SECONDARY (150-156a, 157-159)									
1959	69.1	80.9	59.0	68.9	80.2	59.3	68.6	86.7	52.2
1958	69.7	80.9	60.0	69.5	80.5	60.1	68.2	82.9	54.9
1957	71.3	82.9	61.2	71.2	82.4	61.6	69.1	86.4	53.5
1956	72.8	84.6	62.4	72.8	84.2	62.9	69.7	87.2	53.4
1955	72.9	84.2	62.9	73.1	84.0	63.6	67.3	83.6	51.9
1954	74.3	85.3	64.5	74.4	84.7	65.2	70.6	88.7	53.9
1953	75.4	86.3	65.5	75.7	86.3	66.4	67.4	83.2	52.4
1952	76.1	86.6	66.7	76.5	86.5	67.5	68.8	85.3	53.2
1951	76.5	86.9	67.1	77.0	87.2	67.9	66.8	80.1	54.0
1950	79.2	89.7	69.6	79.5	89.8	70.2	71.6	85.8	57.7

MALIGNANT NEOPLASM OF RESPIRATORY SYSTEM, NOT SPECIFIED AS SECONDARY (160-164)

1959	25.2	45.3	7.4	25.3	45.6	7.4	23.9	41.6	7.6
1958	24.4	43.4	7.4	24.5	43.7	7.4	22.1	38.5	6.9
1957	23.8	42.3	7.2	23.9	42.6	7.2	21.3	37.2	6.4
1956	23.1	40.7	7.1	23.3	41.2	7.1	20.0	34.3	6.4
1955	21.9	38.3	7.0	22.2	38.9	6.9	18.5	30.6	7.0
1954	20.7	35.8	6.8	20.9	36.2	6.9	17.4	29.8	5.6
1953	20.2	34.7	6.8	20.4	35.1	6.9	17.0	28.5	5.8
1952	19.1	32.1	6.9	19.3	32.6	7.0	15.6	26.0	5.6
1951	18.1	30.0	6.8	18.3	30.5	6.9	14.7	23.8	5.8
1950	17.6	28.9	6.9	17.9	29.5	7.0	13.4	21.1	5.6

Source: CDC/NCHS, National Vital Statistics System, Mortality.

HIST293. AGE-ADJUSTED DEATH RATES FOR APPROXIMATELY 64 SELECTED CAUSES, BY RACE AND SEX: UNITED STATES, 1950-59

[Based on age-specific death rates per 100,000 population in specified group. Computed by the direct method, using as the standard population the age distribution of the total population of the United States as projected for 2000. Numbers after causes of death are category numbers of the seventh revision, International Classification of Diseases]

CAUSE OF DEATH CODE AND YEAR	-- ALL RACES --			-- WHITE --			--- ALL OTHER ---		
	BOTH	MALE	FEMALE	BOTH	MALE	FEMALE	BOTH	MALE	FEMALE
MALIGNANT NEOPLASM OF BREAST (170)									
1959	16.7	0.3	31.0	16.9	0.3	31.4	13.7	0.5	25.7
1958	16.7	0.4	31.2	17.0	0.4	31.6	13.4	0.4	25.4
1957	17.2	0.4	32.0	17.4	0.4	32.4	13.9	0.6	26.3
1956	17.2	0.4	32.1	17.4	0.4	32.5	14.4	0.6	27.2
1955	17.3	0.4	32.4	17.6	0.4	32.9	13.5	*	25.8
1954	16.8	0.4	31.6	17.1	0.4	32.0	13.7	0.8	25.7
1953	17.0	0.4	32.0	17.3	0.4	32.5	13.2	0.5	25.1
1952	16.7	0.4	31.4	16.9	0.4	31.9	13.1	0.6	25.0
1951	16.6	0.4	31.4	16.9	0.4	31.9	12.9	0.4	24.9
1950	16.8	0.5	31.9	17.1	0.4	32.4	13.0	0.5	25.2
MALIGNANT NEOPLASM OF GENITAL ORGANS (171-179)									
1959	29.3	29.7	30.6	28.0	28.8	29.0	42.6	40.3	45.5
1958	30.2	30.5	31.6	29.0	29.8	29.9	42.9	38.0	48.4
1957	30.9	31.5	32.0	29.6	30.8	30.3	43.5	38.7	48.7
1956	31.2	31.8	32.2	29.9	31.3	30.3	44.4	37.3	51.7
1955	31.4	31.1	33.2	30.3	30.7	31.4	43.3	36.3	50.4
1954	31.8	31.5	33.5	30.7	31.1	31.8	42.9	35.6	50.5
1953	31.9	30.7	34.4	30.8	30.5	32.5	42.9	32.3	53.6
1952	31.9	30.6	34.5	30.8	30.4	32.6	42.8	32.8	53.0
1951	31.6	29.6	34.8	30.6	29.4	33.0	41.8	31.4	52.5
1950	32.8	30.3	36.5	31.7	30.1	34.6	43.8	31.9	55.8

\* Figure does not meet standard of reliability or precision

Source: CDC/NCHS, National Vital Statistics System, Mortality.

HIST293. AGE-ADJUSTED DEATH RATES FOR APPROXIMATELY 64 SELECTED CAUSES, BY RACE AND SEX: UNITED STATES, 1950-59

[Based on age-specific death rates per 100,000 population in specified group. Computed by the direct method, using as the standard population the age distribution of the total population of the United States as projected for 2000. Numbers after causes of death are category numbers of the seventh revision, International Classification of Diseases]

CAUSE OF DEATH CODE AND YEAR	-- ALL RACES --			-- WHITE --			--- ALL OTHER ---		
	BOTH	MALE	FEMALE	BOTH	MALE	FEMALE	BOTH	MALE	FEMALE
MALIGNANT NEOPLASM OF URINARY ORGANS (180, 181)									
1959	9.6	13.8	6.0	9.7	14.2	6.0	7.1	9.2	5.3
1958	9.5	13.6	6.1	9.7	13.9	6.2	7.1	9.2	5.3
1957	9.8	13.7	6.4	9.9	14.1	6.4	7.3	9.1	5.6
1956	9.7	13.6	6.3	9.8	14.0	6.3	7.1	9.0	5.3
1955	9.7	13.3	6.5	9.9	13.8	6.6	6.2	7.6	4.9
1954	9.6	13.5	6.3	9.8	13.9	6.3	6.9	8.6	5.4
1953	9.6	13.2	6.4	9.8	13.7	6.4	6.8	8.0	5.6
1952	9.6	13.3	6.3	9.8	13.7	6.4	6.6	8.1	5.1
1951	9.5	13.0	6.3	9.6	13.3	6.4	6.4	8.2	4.7
1950	9.5	13.0	6.3	9.7	13.4	6.4	6.8	8.3	5.3

MALIGNANT NEOPLASM OF OTHER AND UNSPECIFIED SITES (156b, 165, 190-199)

1959	20.6	23.5	18.1	20.7	23.8	18.2	19.1	21.1	17.4
1958	20.6	23.0	18.5	20.8	23.3	18.5	18.7	19.8	17.6
1957	21.5	23.9	19.3	21.6	24.3	19.3	19.3	20.0	18.7
1956	21.1	23.6	19.0	21.3	23.9	19.0	18.7	19.9	17.6
1955	21.2	23.6	19.0	21.3	23.8	19.1	19.4	21.2	17.9
1954	21.4	23.8	19.2	21.6	24.3	19.2	17.9	17.8	18.0
1953	21.5	23.7	19.5	21.7	24.1	19.6	18.0	18.8	17.2
1952	22.0	24.1	20.2	22.4	24.7	20.3	17.4	16.7	18.0
1951	21.6	23.6	19.9	21.9	24.1	20.0	17.6	17.6	17.6
1950	19.9	21.6	18.4	20.2	22.0	18.5	16.5	16.6	16.2

Source: CDC/NCHS, National Vital Statistics System, Mortality.

HIST293. AGE-ADJUSTED DEATH RATES FOR APPROXIMATELY 64 SELECTED CAUSES, BY RACE AND SEX: UNITED STATES, 1950-59

[Based on age-specific death rates per 100,000 population in specified group. Computed by the direct method, using as the standard population the age distribution of the total population of the United States as projected for 2000. Numbers after causes of death are category numbers of the seventh revision, International Classification of Diseases]

CAUSE OF DEATH CODE AND YEAR	-- ALL RACES --			-- WHITE --			--- ALL OTHER ---		
	BOTH	MALE	FEMALE	BOTH	MALE	FEMALE	BOTH	MALE	FEMALE
LEUKEMIA AND ALEUKEMIA (204)									
1959	8.2	10.2	6.5	8.5	10.6	6.7	5.3	6.3	4.3
1958	8.3	10.4	6.5	8.5	10.7	6.6	5.3	6.3	4.4
1957	8.1	9.9	6.6	8.4	10.2	6.8	5.2	6.6	3.9
1956	8.1	9.8	6.6	8.3	10.1	6.7	5.3	6.5	4.2
1955	7.8	9.6	6.2	8.1	10.0	6.5	4.6	5.7	3.7
1954	7.7	9.4	6.2	7.9	9.6	6.4	4.8	6.3	3.4
1953	7.4	9.1	6.0	7.7	9.4	6.2	4.4	5.3	3.6
1952	7.4	8.8	6.2	7.7	9.2	6.3	4.4	5.1	3.8
1951	7.2	8.7	5.8	7.4	9.0	6.0	4.2	5.3	3.1
1950	6.9	8.2	5.7	7.2	8.5	5.9	3.7	4.3	3.2
LYMPHOSARCOMA AND OTHER NEOPLASMS OF LYMPHATIC AND HEMATOPOIETIC TISSUES (200-203, 205)									
1959	8.4	10.4	6.7	8.6	10.4	6.9	6.9	9.2	4.9
1958	8.2	9.9	6.6	8.3	10.1	6.7	6.7	8.5	5.0
1957	8.3	10.3	6.4	8.4	10.5	6.6	6.5	8.1	5.0
1956	8.1	10.0	6.4	8.2	10.1	6.5	6.6	8.5	4.9
1955	7.6	9.4	6.0	7.8	9.6	6.2	5.7	7.3	4.3
1954	7.5	9.1	5.9	7.6	9.2	6.1	5.7	7.5	4.0
1953	7.2	8.9	5.6	7.3	9.1	5.7	5.1	6.7	3.5
1952	6.9	8.4	5.4	7.0	8.5	5.6	5.0	6.4	3.7
1951	6.5	8.1	5.0	6.7	8.4	5.2	4.2	5.1	3.3
1950	6.2	7.6	4.8	6.3	7.7	5.0	4.5	5.6	3.5

Source: CDC/NCHS, National Vital Statistics System, Mortality.

HIST293. AGE-ADJUSTED DEATH RATES FOR APPROXIMATELY 64 SELECTED CAUSES, BY RACE AND SEX: UNITED STATES, 1950-59

[Based on age-specific death rates per 100,000 population in specified group. Computed by the direct method, using as the standard population the age distribution of the total population of the United States as projected for 2000. Numbers after causes of death are category numbers of the seventh revision, International Classification of Diseases]

CAUSE OF DEATH CODE AND YEAR	-- ALL RACES --			-- WHITE --			--- ALL OTHER ---		
	BOTH	MALE	FEMALE	BOTH	MALE	FEMALE	BOTH	MALE	FEMALE
BENIGN NEOPLASMS AND NEOPLASMS OF UNSPECIFIED NATURE (210-239)									
1959	3.4	3.0	3.6	3.2	3.0	3.3	4.9	3.7	6.1
1958	3.4	3.2	3.6	3.2	3.2	3.3	4.9	3.7	6.0
1957	3.6	3.3	3.8	3.4	3.3	3.5	5.2	3.7	6.6
1956	3.5	3.1	3.8	3.3	3.1	3.5	5.1	3.3	6.9
1955	3.8	3.3	4.1	3.5	3.3	3.7	5.9	3.6	8.0
1954	3.7	3.3	4.2	3.5	3.2	3.8	5.9	4.1	7.5
1953	3.9	3.5	4.4	3.7	3.4	4.0	5.7	3.6	7.7
1952	3.9	3.2	4.5	3.6	3.2	4.0	6.5	3.6	9.3
1951	4.1	3.3	4.7	3.8	3.2	4.3	6.1	3.5	8.5
1950	4.3	3.4	5.1	4.1	3.4	4.7	6.4	3.3	9.4
ASTHMA (241)									
1959	3.5	4.9	2.3	3.4	4.8	2.1	4.5	5.1	4.0
1958	3.6	5.2	2.3	3.5	5.2	2.1	4.4	5.1	3.8
1957	5.0	7.1	3.1	4.8	7.0	2.9	5.9	7.5	4.4
1956	4.6	6.4	2.9	4.5	6.4	2.8	4.6	5.9	3.5
1955	4.7	6.6	3.0	4.6	6.6	2.9	4.5	5.8	3.3
1954	4.9	6.8	3.2	4.9	6.8	3.2	4.6	5.6	3.6
1953	5.5	7.6	3.6	5.6	7.7	3.6	4.8	5.6	3.9
1952	5.8	7.9	3.8	5.7	8.0	3.7	5.8	6.4	5.2
1951	6.0	8.1	4.1	6.0	8.3	4.0	5.0	5.6	4.5
1950	3.7	4.9	2.7	3.7	4.9	2.6	3.8	4.4	3.1

Source: CDC/NCHS, National Vital Statistics System, Mortality.

HIST293. AGE-ADJUSTED DEATH RATES FOR APPROXIMATELY 64 SELECTED CAUSES, BY RACE AND SEX: UNITED STATES, 1950-59

[Based on age-specific death rates per 100,000 population in specified group. Computed by the direct method, using as the standard population the age distribution of the total population of the United States as projected for 2000. Numbers after causes of death are category numbers of the seventh revision, International Classification of Diseases]

CAUSE OF DEATH CODE AND YEAR	-- ALL RACES --			-- WHITE --			--- ALL OTHER ---		
	BOTH	MALE	FEMALE	BOTH	MALE	FEMALE	BOTH	MALE	FEMALE
DIABETES (260)									
1959	21.3	18.6	23.6	20.7	18.4	22.5	27.3	19.9	34.0
1958	21.4	18.6	23.9	20.9	18.4	22.9	26.6	19.2	33.6
1957	21.4	18.1	24.3	21.0	18.1	23.5	24.9	17.7	31.8
1956	21.3	18.1	24.1	21.0	18.2	23.4	23.4	16.3	30.1
1955	21.2	18.0	24.1	21.0	18.1	23.5	22.9	15.9	29.4
1954	21.4	18.0	24.5	21.2	18.1	24.0	22.4	16.1	28.6
1953	22.5	18.5	26.0	22.2	18.6	25.4	23.9	16.6	30.9
1952	22.7	19.0	26.0	22.5	19.2	25.5	22.8	16.6	28.9
1951	22.8	18.3	26.8	22.6	18.4	26.5	22.8	16.7	28.7
1950	23.1	18.8	27.0	22.9	18.9	26.5	23.7	16.7	30.5
MENINGITIS, EXCEPT MENINGOCOCCAL AND TUBERCULOUS (340)									
1959	1.0	1.2	0.9	0.9	1.0	0.7	2.5	2.8	2.2
1958	1.1	1.3	0.9	0.9	1.1	0.7	2.5	2.8	2.2
1957	1.0	1.1	0.8	0.8	0.9	0.7	2.5	2.8	2.3
1956	1.0	1.1	0.8	0.8	1.0	0.6	2.5	2.9	2.1
1955	1.0	1.1	0.8	0.8	0.9	0.6	2.6	3.0	2.3
1954	1.0	1.1	0.8	0.8	0.9	0.6	2.7	3.0	2.4
1953	1.1	1.3	0.9	0.9	1.2	0.7	2.5	2.9	2.2
1952	1.1	1.2	0.9	0.9	1.0	0.7	2.7	3.0	2.4
1951	1.1	1.3	0.8	0.9	1.1	0.6	2.7	3.1	2.3
1950	1.1	1.3	0.8	0.9	1.1	0.7	2.8	3.4	2.1

Source: CDC/NCHS, National Vital Statistics System, Mortality.

HIST293. AGE-ADJUSTED DEATH RATES FOR APPROXIMATELY 64 SELECTED CAUSES, BY RACE AND SEX: UNITED STATES, 1950-59

[Based on age-specific death rates per 100,000 population in specified group. Computed by the direct method, using as the standard population the age distribution of the total population of the United States as projected for 2000. Numbers after causes of death are category numbers of the seventh revision, International Classification of Diseases]

CAUSE OF DEATH CODE AND YEAR	-- ALL RACES --			-- WHITE --			--- ALL OTHER ---		
	BOTH	MALE	FEMALE	BOTH	MALE	FEMALE	BOTH	MALE	FEMALE
MAJOR CARDIOVASCULAR-RENAL DISEASES (330-334, 400-468, 592-594)									
1959	807.6	946.9	685.4	800.8	946.7	673.2	849.4	926.4	780.2
1958	826.4	964.1	705.0	818.6	962.6	691.7	881.0	964.6	811.5
1957	828.6	962.9	708.8	820.2	960.3	695.2	886.0	965.2	820.6
1956	815.0	945.0	698.3	808.1	944.0	686.1	859.7	929.6	796.6
1955	813.0	939.9	698.1	806.5	939.0	686.8	853.1	921.8	789.1
1954	795.8	916.9	685.2	788.0	913.9	673.0	847.1	921.3	783.8
1953	834.9	959.4	720.6	825.4	954.2	707.5	906.9	986.7	832.5
1952	837.6	959.0	725.7	826.9	952.5	711.0	927.0	1,005.9	855.3
1951	853.9	977.6	739.8	843.9	973.0	725.2	935.5	1,004.5	873.4
1950	867.6	994.4	751.3	856.7	989.7	734.8	964.9	1,024.7	908.1
DISEASES OF CARDIOVASCULAR SYSTEM (330-334, 400-468)									
1959	797.7	935.2	677.0	792.1	936.3	665.9	825.9	900.0	759.1
1958	814.9	950.5	695.1	808.5	950.6	683.1	854.3	934.8	787.4
1957	815.9	948.5	697.7	809.0	947.4	685.5	856.8	933.3	793.2
1956	801.6	929.7	686.5	796.2	930.3	675.8	829.3	896.0	768.7
1955	798.8	923.7	685.7	793.8	924.3	675.9	822.1	889.4	759.4
1954	780.2	899.1	671.5	774.1	898.0	660.7	812.4	880.8	753.9
1953	817.1	939.1	704.9	809.5	935.9	693.7	866.9	943.5	795.3
1952	817.4	935.8	708.0	809.1	931.9	695.6	877.7	951.6	810.4
1951	830.9	952.0	719.2	823.5	950.1	706.9	883.1	947.0	825.0
1950	841.4	965.0	727.8	833.2	963.2	714.0	905.5	960.5	852.9

Source: CDC/NCHS, National Vital Statistics System, Mortality.

HIST293. AGE-ADJUSTED DEATH RATES FOR APPROXIMATELY 64 SELECTED CAUSES, BY RACE AND SEX: UNITED STATES, 1950-59

[Based on age-specific death rates per 100,000 population in specified group. Computed by the direct method, using as the standard population the age distribution of the total population of the United States as projected for 2000. Numbers after causes of death are category numbers of the seventh revision, International Classification of Diseases]

CAUSE OF DEATH CODE AND YEAR	-- ALL RACES --			-- WHITE --			--- ALL OTHER ---		
	BOTH	MALE	FEMALE	BOTH	MALE	FEMALE	BOTH	MALE	FEMALE
CEREBROVASCULAR DISEASES (ICD 330-334)									
1959	179.0	185.8	172.9	173.9	181.6	167.2	228.3	227.2	229.2
1958	183.0	189.0	177.4	178.0	184.3	172.0	232.4	237.7	229.3
1957	183.0	189.7	176.8	177.9	185.4	171.1	231.0	231.8	231.5
1956	177.4	182.8	172.3	172.7	178.8	166.9	223.5	221.8	225.0
1955	177.3	182.8	172.1	172.7	178.8	167.0	222.3	222.8	221.9
1954	173.4	177.4	169.4	168.5	173.0	163.9	221.7	221.1	223.7
1953	179.2	183.2	175.2	174.2	178.7	169.9	227.8	226.5	228.6
1952	180.2	184.9	175.8	174.9	180.1	170.0	234.1	233.7	235.0
1951	181.4	185.1	177.8	176.3	180.7	172.1	233.5	230.3	237.3
1950	180.7	186.4	175.8	175.5	182.1	169.7	236.0	231.3	240.6
DISEASES OF HEART (ICD 400-402, 410-413)									
1959	552.1	674.9	444.2	552.2	680.8	439.6	527.5	595.0	466.7
1958	563.6	684.7	456.6	563.1	690.0	451.0	547.3	614.9	489.2
1957	568.5	687.0	462.8	567.3	690.8	457.0	557.2	626.3	498.4
1956	560.6	675.9	456.8	560.4	680.8	452.1	540.0	603.3	482.9
1955	556.7	668.6	455.1	556.9	673.7	451.1	531.8	592.0	475.5
1954	544.6	653.7	444.9	544.1	657.7	440.4	523.9	587.4	467.8
1953	570.1	681.8	467.6	568.2	683.8	462.1	567.7	636.7	503.2
1952	568.8	676.7	468.9	566.3	678.2	462.8	573.0	640.8	510.7
1951	577.9	689.0	475.3	575.8	691.7	469.0	578.1	638.5	522.2
1950	588.8	699.0	486.6	586.0	701.4	479.2	599.1	653.2	547.4

Source: CDC/NCHS, National Vital Statistics System, Mortality.



HIST293. AGE-ADJUSTED DEATH RATES FOR APPROXIMATELY 64 SELECTED CAUSES, BY RACE AND SEX: UNITED STATES, 1950-59

[Based on age-specific death rates per 100,000 population in specified group. Computed by the direct method, using as the standard population the age distribution of the total population of the United States as projected for 2000. Numbers after causes of death are category numbers of the seventh revision, International Classification of Diseases]

CAUSE OF DEATH CODE AND YEAR	-- ALL RACES --			-- WHITE --			--- ALL OTHER ---		
	BOTH	MALE	FEMALE	BOTH	MALE	FEMALE	BOTH	MALE	FEMALE
RHEUMATIC FEVER AND CHRONIC RHEUMATIC HEART DISEASE (400-402, 410-416)									
1959	12.5	12.5	12.5	12.6	12.5	12.6	11.0	11.6	10.4
1958	13.2	13.2	13.1	13.3	13.3	13.2	11.5	11.7	11.4
1957	14.4	14.5	14.2	14.4	14.5	14.3	13.5	13.7	13.3
1956	14.7	15.0	14.4	14.7	15.0	14.4	13.5	14.5	12.7
1955	14.7	15.0	14.5	14.7	15.0	14.4	14.0	13.8	14.0
1954	14.9	15.1	14.6	14.8	15.0	14.6	15.0	15.7	14.4
1953	16.4	16.7	16.1	16.3	16.6	15.8	17.2	17.1	17.3
1952	17.0	17.7	16.4	16.8	17.5	16.2	19.0	19.9	18.3
1951	17.7	18.2	17.3	17.5	18.0	17.1	19.2	19.4	19.0
1950	19.0	19.6	18.4	18.7	19.3	18.1	22.5	23.2	21.6
ARTERIOSCLEROTIC HEART DISEASE, INCLUDING CORONARY DISEASE (420)									
1959	400.5	518.0	297.9	409.0	532.3	301.6	288.2	348.7	233.6
1958	399.8	516.9	297.0	408.4	531.7	300.5	285.9	343.8	235.0
1957	401.1	513.8	301.1	409.5	527.7	304.8	287.7	346.2	236.3
1956	388.2	497.4	290.5	397.2	512.1	294.6	269.1	319.2	223.1
1955	376.6	482.1	281.4	385.8	496.9	285.9	254.5	300.6	211.2
1954	359.0	461.0	266.2	367.9	474.9	270.8	237.9	288.0	192.5
1953	361.9	464.9	267.8	370.3	478.2	272.1	246.5	296.5	199.9
1952	348.3	445.9	258.3	356.9	459.6	262.7	230.4	275.1	188.5
1951	342.1	441.0	251.0	351.1	455.2	255.4	220.8	263.1	181.4
1950	337.0	434.0	247.1	345.6	448.0	251.0	218.5	256.7	181.5

Source: CDC/NCHS, National Vital Statistics System, Mortality.

HIST293. AGE-ADJUSTED DEATH RATES FOR APPROXIMATELY 64 SELECTED CAUSES, BY RACE AND SEX: UNITED STATES, 1950-59

[Based on age-specific death rates per 100,000 population in specified group. Computed by the direct method, using as the standard population the age distribution of the total population of the United States as projected for 2000. Numbers after causes of death are category numbers of the seventh revision, International Classification of Diseases]

CAUSE OF DEATH CODE AND YEAR	-- ALL RACES --			-- WHITE --			--- ALL OTHER ---		
	BOTH	MALE	FEMALE	BOTH	MALE	FEMALE	BOTH	MALE	FEMALE
NONRHEUMATIC CHRONIC ENDOCARDITIS AND OTHER MYOCARDIAL DEGENERATION (421, 422)									
1959	58.2	62.8	54.3	57.5	62.1	53.6	63.7	69.4	58.7
1958	62.3	66.4	58.8	61.4	65.2	58.0	71.5	78.7	65.7
1957	67.8	72.8	63.3	66.9	71.7	62.5	75.6	82.5	70.2
1956	70.4	75.6	65.8	69.6	74.5	65.2	77.9	85.6	70.9
1955	75.3	80.4	70.6	74.7	79.6	70.2	79.5	87.9	71.7
1954	77.7	83.5	72.4	76.7	82.4	71.6	84.9	93.9	77.2
1953	89.0	95.5	82.9	88.1	94.3	82.5	95.0	106.6	83.9
1952	95.2	102.4	88.6	94.2	101.1	87.9	103.4	114.5	93.2
1951	104.0	113.2	95.7	103.1	112.0	95.1	111.3	123.8	99.8
1950	114.1	123.7	105.5	113.2	122.8	104.6	122.7	132.2	113.6
OTHER DISEASES OF HEART (430-434)									
1959	20.7	25.5	16.5	19.4	24.1	15.4	33.7	40.2	28.0
1958	21.5	26.6	17.1	20.1	24.9	16.0	35.9	45.0	28.0
1957	19.1	23.7	15.0	17.6	22.0	13.7	34.8	41.4	29.0
1956	18.9	23.6	14.8	17.5	22.0	13.6	33.5	40.6	27.1
1955	18.9	23.8	14.7	17.5	22.1	13.4	35.0	42.1	28.7
1954	19.7	24.5	15.3	18.2	22.9	14.0	34.9	41.6	28.9
1953	21.8	27.2	17.0	20.1	25.2	15.5	40.8	49.0	33.2
1952	22.3	27.7	17.4	20.5	25.6	15.8	42.8	51.9	34.3
1951	23.6	29.2	18.6	21.7	27.1	16.8	45.1	52.1	38.7
1950	25.4	31.4	20.0	23.4	29.3	18.1	47.5	54.8	40.8

Source: CDC/NCHS, National Vital Statistics System, Mortality.

HIST293. AGE-ADJUSTED DEATH RATES FOR APPROXIMATELY 64 SELECTED CAUSES, BY RACE AND SEX: UNITED STATES, 1950-59

[Based on age-specific death rates per 100,000 population in specified group. Computed by the direct method, using as the standard population the age distribution of the total population of the United States as projected for 2000. Numbers after causes of death are category numbers of the seventh revision, International Classification of Diseases]

CAUSE OF DEATH CODE AND YEAR	-- ALL RACES --			-- WHITE --			--- ALL OTHER ---		
	BOTH	MALE	FEMALE	BOTH	MALE	FEMALE	BOTH	MALE	FEMALE
HYPERTENSIVE HEART DISEASE (440-443)									
1959	60.2	56.1	63.1	53.7	49.7	56.4	130.9	125.2	136.0
1958	66.8	61.6	70.6	59.9	54.8	63.4	142.3	135.8	149.1
1957	66.2	62.2	69.0	58.9	54.9	61.7	145.6	142.6	149.6
1956	68.4	64.3	71.3	61.3	57.2	64.3	146.1	143.4	149.0
1955	71.2	67.4	74.0	64.3	60.1	67.2	148.8	147.6	149.9
1954	73.5	69.6	76.4	66.5	62.6	69.4	151.2	148.2	154.8
1953	81.1	77.5	83.8	73.4	69.5	76.2	168.2	167.5	169.0
1952	85.9	82.9	88.2	77.9	74.4	80.3	177.5	179.4	176.3
1951	90.4	87.4	92.7	82.4	79.3	84.7	181.5	180.1	183.4
1950	93.3	90.2	95.6	85.1	81.9	87.5	188.0	186.4	189.9
OTHER HYPERTENSIVE DISEASE (444-447)									
1959	12.0	13.3	11.0	11.0	12.1	10.0	22.0	24.5	19.8
1958	12.8	14.1	11.7	11.7	12.9	10.7	23.9	26.4	21.9
1957	10.7	11.6	10.0	9.7	10.5	9.0	21.5	22.8	20.4
1956	10.9	11.9	10.0	9.9	10.9	9.0	21.6	22.7	20.5
1955	11.4	12.4	10.5	10.3	11.3	9.4	22.5	24.5	20.9
1954	11.6	12.5	10.9	10.5	11.4	9.7	24.0	24.6	23.6
1953	13.0	14.0	12.0	11.7	12.6	10.9	26.2	28.9	23.7
1952	13.5	14.2	12.8	12.2	13.0	11.6	26.5	28.1	25.2
1951	14.8	15.4	14.2	13.5	14.2	13.0	28.3	29.4	27.3
1950	14.7	15.6	13.8	13.6	14.7	12.7	26.0	26.3	25.8

Source: CDC/NCHS, National Vital Statistics System, Mortality.

HIST293. AGE-ADJUSTED DEATH RATES FOR APPROXIMATELY 64 SELECTED CAUSES, BY RACE AND SEX: UNITED STATES, 1950-59

[Based on age-specific death rates per 100,000 population in specified group. Computed by the direct method, using as the standard population the age distribution of the total population of the United States as projected for 2000. Numbers after causes of death are category numbers of the seventh revision, International Classification of Diseases]

CAUSE OF DEATH CODE AND YEAR	-- ALL RACES --			-- WHITE --			--- ALL OTHER ---		
	BOTH	MALE	FEMALE	BOTH	MALE	FEMALE	BOTH	MALE	FEMALE
GENERAL ARTERIOSCLEROSIS (450)									
1959	40.5	43.2	38.3	41.1	43.9	38.9	32.1	34.6	29.7
1958	41.9	45.0	39.3	42.4	45.6	39.8	35.0	38.5	32.4
1957	41.5	44.3	39.2	42.1	44.9	39.8	33.4	36.2	31.4
1956	41.5	44.3	39.1	42.2	45.0	39.9	31.4	34.1	29.2
1955	43.4	46.8	40.6	44.2	47.6	41.3	33.8	37.0	30.9
1954	41.9	44.4	39.8	42.6	44.9	40.5	32.9	36.7	29.9
1953	46.5	49.8	43.8	47.4	50.4	44.7	35.7	40.3	31.7
1952	47.2	50.1	44.6	48.1	50.9	45.6	35.3	39.5	31.6
1951	49.4	53.4	46.0	50.5	54.5	47.2	34.5	38.8	30.6
1950	50.4	55.7	46.0	51.5	56.9	47.1	36.3	40.8	32.1
OTHER DISEASES OF CIRCULATORY SYSTEM (451-468)									
1959	14.1	18.1	10.5	13.8	18.0	10.2	16.0	18.6	13.8
1958	13.6	17.8	10.1	13.4	17.8	9.6	15.8	17.3	14.5
1957	12.1	15.9	8.9	11.9	15.8	8.6	13.8	16.3	11.5
1956	11.3	14.8	8.2	11.1	14.8	7.9	12.6	14.3	11.1
1955	10.0	12.9	7.4	9.8	12.9	7.1	11.6	13.1	10.2
1954	8.6	11.0	6.4	8.4	10.9	6.2	9.9	11.0	8.9
1953	8.2	10.3	6.3	8.1	10.2	6.1	9.6	11.1	8.1
1952	7.8	9.9	5.9	7.6	9.8	5.6	8.7	9.5	8.0
1951	7.4	9.2	5.9	7.3	9.0	5.7	8.8	10.0	7.7
1950	6.8	8.3	5.5	6.7	8.2	5.3	7.9	8.8	7.0

Source: CDC/NCHS, National Vital Statistics System, Mortality.

HIST293. AGE-ADJUSTED DEATH RATES FOR APPROXIMATELY 64 SELECTED CAUSES, BY RACE AND SEX: UNITED STATES, 1950-59

[Based on age-specific death rates per 100,000 population in specified group. Computed by the direct method, using as the standard population the age distribution of the total population of the United States as projected for 2000. Numbers after causes of death are category numbers of the seventh revision, International Classification of Diseases]

CAUSE OF DEATH CODE AND YEAR	-- ALL RACES --			-- WHITE --			--- ALL OTHER ---		
	BOTH	MALE	FEMALE	BOTH	MALE	FEMALE	BOTH	MALE	FEMALE
CHRONIC AND UNSPECIFIED NEPHRITIS AND OTHER RENAL SCLEROSIS (592-594)									
1959	9.9	11.7	8.5	8.7	10.3	7.3	23.5	26.4	21.1
1958	11.5	13.5	9.9	10.1	12.0	8.6	26.7	29.9	24.2
1957	12.7	14.4	11.2	11.2	12.9	9.7	29.2	31.8	27.3
1956	13.4	15.4	11.7	11.8	13.8	10.3	30.4	33.6	27.9
1955	14.2	16.2	12.5	12.7	14.7	10.9	31.0	32.4	29.8
1954	15.6	17.9	13.7	13.9	15.9	12.3	34.7	40.5	29.9
1953	17.8	20.3	15.7	15.9	18.3	13.8	40.0	43.3	37.2
1952	20.2	23.2	17.7	17.7	20.5	15.3	49.3	54.3	44.9
1951	23.0	25.6	20.7	20.4	22.9	18.2	52.5	57.5	48.4
1950	26.2	29.4	23.5	23.5	26.5	20.8	59.4	64.3	55.2
INFLUENZA AND PNEUMONIA, EXCEPT PNEUMONIA OF NEWBORN (480-493)									
1959	44.3	53.7	36.5	42.0	51.0	34.6	61.2	74.6	49.0
1958	47.4	57.7	38.7	44.5	54.2	36.5	71.1	88.6	55.9
1957	50.3	60.7	41.5	47.1	57.2	38.7	76.9	91.0	64.4
1956	40.8	48.6	34.2	38.6	46.0	32.2	58.5	69.1	48.7
1955	39.2	45.8	33.3	36.8	42.9	31.5	58.7	71.5	47.1
1954	36.2	42.7	30.5	33.8	39.8	28.4	56.6	67.7	46.6
1953	49.2	57.7	41.6	45.4	53.4	38.4	84.7	98.5	71.8
1952	43.1	49.7	37.1	39.9	46.0	34.4	71.2	82.9	60.1
1951	47.4	54.2	41.3	44.3	50.7	38.5	76.1	86.6	66.3
1950	48.1	55.0	41.9	44.8	51.4	38.8	80.5	90.0	71.5

Source: CDC/NCHS, National Vital Statistics System, Mortality.

HIST293. AGE-ADJUSTED DEATH RATES FOR APPROXIMATELY 64 SELECTED CAUSES, BY RACE AND SEX: UNITED STATES, 1950-59

[Based on age-specific death rates per 100,000 population in specified group. Computed by the direct method, using as the standard population the age distribution of the total population of the United States as projected for 2000. Numbers after causes of death are category numbers of the seventh revision, International Classification of Diseases]

CAUSE OF DEATH CODE AND YEAR	-- ALL RACES --			-- WHITE --			--- ALL OTHER ---		
	BOTH	MALE	FEMALE	BOTH	MALE	FEMALE	BOTH	MALE	FEMALE
INFLUENZA (480-483)									
1959	2.4	2.6	2.2	2.2	2.5	2.1	3.9	4.1	3.7
1958	4.0	4.4	3.6	3.5	3.8	3.2	9.1	11.1	7.4
1957	6.2	6.9	5.5	5.6	6.4	5.0	11.8	12.2	11.5
1956	2.2	2.4	2.1	2.0	2.2	1.8	4.8	4.9	4.8
1955	2.6	2.6	2.6	2.3	2.3	2.2	5.9	6.0	5.8
1954	2.5	2.7	2.4	2.2	2.3	2.0	5.7	5.8	5.6
1953	9.9	10.9	9.2	8.9	9.8	8.2	21.3	22.1	20.6
1952	5.7	6.0	5.5	5.1	5.3	4.9	12.2	12.6	11.9
1951	7.7	8.0	7.4	7.0	7.3	6.7	14.3	14.1	14.6
1950	7.5	7.8	7.2	6.7	7.1	6.4	15.6	15.3	15.9
PNEUMONIA, EXCEPT PNEUMONIA OF NEWBORN (490-493)									
1959	41.9	51.1	34.3	39.8	48.5	32.6	57.2	70.5	45.2
1958	43.4	53.3	35.2	41.1	50.4	33.3	62.1	77.5	48.5
1957	44.1	53.7	36.0	41.6	50.7	33.7	65.1	78.8	52.9
1956	38.6	46.2	32.1	36.6	43.9	30.4	53.7	64.2	44.0
1955	36.6	43.2	30.7	34.5	40.6	29.2	52.9	65.5	41.3
1954	33.7	40.0	28.1	31.6	37.5	26.4	50.9	61.9	41.0
1953	39.2	46.8	32.5	36.5	43.5	30.3	63.3	76.3	51.2
1952	37.4	43.7	31.6	34.8	40.7	29.5	58.9	70.3	48.2
1951	39.8	46.3	34.0	37.3	43.3	31.9	61.8	72.5	51.7
1950	40.6	47.2	34.7	38.0	44.3	32.4	64.8	74.7	55.6

Source: CDC/NCHS, National Vital Statistics System, Mortality.

HIST293. AGE-ADJUSTED DEATH RATES FOR APPROXIMATELY 64 SELECTED CAUSES, BY RACE AND SEX: UNITED STATES, 1950-59

[Based on age-specific death rates per 100,000 population in specified group. Computed by the direct method, using as the standard population the age distribution of the total population of the United States as projected for 2000. Numbers after causes of death are category numbers of the seventh revision, International Classification of Diseases]

CAUSE OF DEATH CODE AND YEAR	-- ALL RACES --			-- WHITE --			--- ALL OTHER ---		
	BOTH	MALE	FEMALE	BOTH	MALE	FEMALE	BOTH	MALE	FEMALE
BRONCHITIS (500-502)									
1959	2.6	3.8	1.6	2.6	3.9	1.5	2.1	2.7	1.6
1958	2.7	3.9	1.7	2.7	3.9	1.7	2.5	3.1	2.0
1957	2.5	3.5	1.7	2.5	3.4	1.6	2.6	3.1	2.1
1956	2.3	3.1	1.6	2.3	3.2	1.5	2.1	2.5	1.7
1955	2.3	3.0	1.7	2.3	3.0	1.7	2.1	2.5	1.8
1954	2.2	2.9	1.6	2.2	2.9	1.6	1.9	2.2	1.7
1953	2.5	3.2	1.8	2.5	3.2	1.8	2.5	3.0	2.0
1952	2.4	3.0	1.8	2.4	3.0	1.8	2.4	2.9	2.0
1951	2.6	3.3	2.0	2.6	3.3	1.9	2.6	2.9	2.4
1950	2.9	3.5	2.3	2.9	3.5	2.3	2.9	3.0	2.8
ULCER OF STOMACH AND DUODEMA (540, 541)									
1959	8.1	12.8	4.0	8.2	13.0	4.1	6.4	10.1	3.1
1958	8.4	13.2	4.1	8.4	13.5	4.1	7.0	10.1	4.1
1957	8.1	12.9	4.0	8.2	13.1	4.0	6.6	10.3	3.2
1956	8.1	13.0	3.8	8.2	13.3	3.8	6.1	9.0	3.5
1955	7.8	12.6	3.6	7.9	12.8	3.6	6.4	9.7	3.4
1954	7.7	12.5	3.3	7.7	12.8	3.3	6.3	9.6	3.2
1953	7.7	12.7	3.1	7.7	12.9	3.1	6.6	10.2	3.1
1952	7.3	12.1	3.0	7.4	12.3	3.0	6.0	9.4	2.8
1951	7.2	11.9	2.9	7.3	12.1	2.9	6.0	9.1	3.1
1950	7.2	11.8	2.8	7.2	11.9	2.8	6.5	10.2	2.8

Source: CDC/NCHS, National Vital Statistics System, Mortality.

HIST293. AGE-ADJUSTED DEATH RATES FOR APPROXIMATELY 64 SELECTED CAUSES, BY RACE AND SEX: UNITED STATES, 1950-59

[Based on age-specific death rates per 100,000 population in specified group. Computed by the direct method, using as the standard population the age distribution of the total population of the United States as projected for 2000. Numbers after causes of death are category numbers of the seventh revision, International Classification of Diseases]

CAUSE OF DEATH CODE AND YEAR	-- ALL RACES --			-- WHITE --			--- ALL OTHER ---		
	BOTH	MALE	FEMALE	BOTH	MALE	FEMALE	BOTH	MALE	FEMALE
APPENDICITIS (550-553)									
1959	1.3	1.8	0.9	1.2	1.7	0.8	2.3	2.8	1.9
1958	1.4	1.8	1.0	1.3	1.7	0.9	2.1	2.6	1.6
1957	1.5	1.9	1.0	1.4	1.9	0.9	2.2	2.5	1.9
1956	1.6	2.1	1.2	1.5	2.0	1.1	2.3	2.7	1.9
1955	1.7	2.3	1.3	1.7	2.2	1.2	2.5	3.1	2.0
1954	1.8	2.4	1.3	1.7	2.2	1.2	2.9	3.5	2.2
1953	1.8	2.5	1.2	1.7	2.4	1.1	2.6	3.2	2.0
1952	2.1	2.8	1.6	2.0	2.6	1.4	3.6	4.5	2.7
1951	2.3	2.9	1.7	2.1	2.7	1.5	3.7	4.5	2.9
1950	2.5	3.2	1.7	2.3	3.1	1.6	3.7	4.7	2.8
HERNIA AND INTESTINAL OBSTRUCTION (560, 561, 570)									
1959	7.3	7.7	6.9	7.1	7.6	6.8	8.1	8.7	7.6
1958	7.1	7.6	6.7	6.9	7.4	6.6	8.7	9.8	7.8
1957	7.1	7.7	6.6	6.9	7.5	6.5	8.2	9.6	7.1
1956	7.1	7.6	6.6	6.9	7.4	6.5	8.4	9.8	7.1
1955	7.5	8.2	6.9	7.3	8.0	6.8	8.6	10.2	7.1
1954	7.4	8.2	6.7	7.2	7.9	6.6	9.0	10.6	7.6
1953	7.8	8.4	7.2	7.5	8.2	7.0	9.7	10.7	8.7
1952	7.8	8.6	7.0	7.6	8.3	6.9	9.3	10.9	7.9
1951	8.0	8.7	7.3	7.7	8.5	7.1	9.9	11.1	8.7
1950	8.7	9.4	8.0	8.4	9.1	7.8	10.9	12.4	9.5

Source: CDC/NCHS, National Vital Statistics System, Mortality.



HIST293. AGE-ADJUSTED DEATH RATES FOR APPROXIMATELY 64 SELECTED CAUSES, BY RACE AND SEX: UNITED STATES, 1950-59

[Based on age-specific death rates per 100,000 population in specified group. Computed by the direct method, using as the standard population the age distribution of the total population of the United States as projected for 2000. Numbers after causes of death are category numbers of the seventh revision, International Classification of Diseases]

CAUSE OF DEATH CODE AND YEAR	-- ALL RACES --			-- WHITE --			--- ALL OTHER ---		
	BOTH	MALE	FEMALE	BOTH	MALE	FEMALE	BOTH	MALE	FEMALE
GASTRITIS, DUODENITIS, ENTERITIS, AND COLITIS, EXCEPT DIARRHEA OF NEWBORN (543, 571, 572)									
1959	4.9	4.9	4.9	4.6	4.5	4.6	6.7	7.3	6.2
1958	4.9	5.1	4.8	4.6	4.7	4.5	6.8	7.2	6.4
1957	5.0	5.1	4.8	4.5	4.6	4.5	7.2	7.8	6.5
1956	4.9	5.2	4.7	4.6	4.8	4.3	6.7	7.1	6.2
1955	4.9	5.0	4.8	4.5	4.6	4.4	7.3	7.5	7.1
1954	5.0	5.2	4.8	4.6	4.7	4.4	7.9	8.7	7.1
1953	5.3	5.5	5.1	4.8	5.0	4.6	8.9	9.1	8.6
1952	5.5	5.8	5.2	4.8	5.0	4.6	9.9	10.8	9.1
1951	5.3	5.3	5.1	4.8	4.8	4.7	8.6	9.1	8.0
1950	5.2	5.4	5.0	4.8	4.8	4.6	8.9	9.7	8.1
CIRRHOSIS OF LIVER (581)									
1959	12.9	17.6	8.5	12.8	17.7	8.3	13.1	16.2	10.2
1958	12.7	17.5	8.4	12.7	17.7	8.3	11.7	15.2	8.5
1957	13.4	18.5	8.7	13.4	18.6	8.6	12.8	16.2	9.6
1956	12.7	17.2	8.6	12.8	17.5	8.5	11.5	14.3	8.9
1955	12.2	16.6	8.1	12.3	16.9	8.1	10.4	12.9	8.0
1954	12.0	16.4	8.0	12.2	16.7	8.0	9.8	12.1	7.5
1953	12.5	16.7	8.5	12.7	17.1	8.5	9.8	11.9	7.6
1952	12.3	16.8	8.1	12.5	17.1	8.2	9.9	13.1	6.8
1951	12.0	15.9	8.3	12.2	16.2	8.3	9.9	12.2	7.7
1950	11.3	14.9	7.8	11.5	15.3	7.8	9.3	11.2	7.3

Source: CDC/NCHS, National Vital Statistics System, Mortality.

HIST293. AGE-ADJUSTED DEATH RATES FOR APPROXIMATELY 64 SELECTED CAUSES, BY RACE AND SEX: UNITED STATES, 1950-59

[Based on age-specific death rates per 100,000 population in specified group. Computed by the direct method, using as the standard population the age distribution of the total population of the United States as projected for 2000. Numbers after causes of death are category numbers of the seventh revision, International Classification of Diseases]

CAUSE OF DEATH CODE AND YEAR	-- ALL RACES --			-- WHITE --			--- ALL OTHER ---		
	BOTH	MALE	FEMALE	BOTH	MALE	FEMALE	BOTH	MALE	FEMALE
CHOLELITHIASIS, CHOLECYSTITIS, AND CHOLANGITIS (584, 585)									
1959	3.8	3.4	4.1	3.9	3.5	4.2	2.3	1.6	2.9
1958	4.1	3.6	4.5	4.2	3.8	4.6	2.2	1.8	2.6
1957	4.3	3.7	4.8	4.5	3.9	5.0	2.3	1.7	2.8
1956	4.7	3.9	5.4	4.9	4.1	5.6	2.3	1.8	2.8
1955	5.3	4.3	6.1	5.5	4.5	6.3	2.3	1.5	3.0
1954	5.2	4.2	6.0	5.4	4.4	6.2	2.4	1.4	3.3
1953	5.1	4.2	6.0	5.3	4.3	6.2	2.5	1.8	3.2
1952	5.5	4.2	6.7	5.8	4.4	6.9	2.3	1.5	3.1
1951	5.5	4.3	6.6	5.8	4.5	6.9	2.2	1.7	2.7
1950	5.8	4.5	7.0	6.0	4.7	7.3	2.2	1.4	3.1
NEPHRITIS AND NEPHROSIS (590-594)									
1959	11.2	13.3	9.5	9.8	11.7	8.2	26.5	29.8	23.8
1958	13.2	15.4	11.3	11.6	13.7	9.8	30.3	33.8	27.4
1957	14.4	16.5	12.6	12.6	14.7	10.9	33.2	36.1	31.0
1956	15.2	17.5	13.4	13.4	15.6	11.6	34.9	38.5	31.9
1955	16.0	18.3	14.0	14.2	16.6	12.2	35.5	37.2	33.8
1954	17.6	20.2	15.5	15.7	18.0	13.8	39.2	45.5	33.9
1953	20.0	22.9	17.7	17.8	20.6	15.4	45.7	49.6	42.3
1952	22.8	26.0	20.0	19.9	23.0	17.3	55.3	60.6	50.7
1951	25.7	28.7	23.1	22.8	25.6	20.3	59.8	65.3	55.4
1950	29.3	32.9	26.1	26.1	29.6	23.0	66.8	72.5	61.9

Source: CDC/NCHS, National Vital Statistics System, Mortality.

HIST293. AGE-ADJUSTED DEATH RATES FOR APPROXIMATELY 64 SELECTED CAUSES, BY RACE AND SEX: UNITED STATES, 1950-59

[Based on age-specific death rates per 100,000 population in specified group. Computed by the direct method, using as the standard population the age distribution of the total population of the United States as projected for 2000. Numbers after causes of death are category numbers of the seventh revision, International Classification of Diseases]

CAUSE OF DEATH CODE AND YEAR	-- ALL RACES --			-- WHITE --			--- ALL OTHER ---		
	BOTH	MALE	FEMALE	BOTH	MALE	FEMALE	BOTH	MALE	FEMALE
ACUTE NEPHRITIS, AND NEPHRITIS WITH EDEMA INCLUDING NEPHROSIS (590, 591)									
1959	1.3	1.5	1.1	1.1	1.4	0.9	3.0	3.5	2.7
1958	1.6	1.9	1.4	1.4	1.7	1.2	3.6	3.9	3.3
1957	1.7	2.0	1.4	1.5	1.8	1.2	4.0	4.3	3.6
1956	1.8	2.1	1.6	1.6	1.8	1.4	4.4	4.9	4.0
1955	1.8	2.2	1.6	1.6	1.9	1.3	4.5	4.8	4.1
1954	2.0	2.4	1.7	1.8	2.1	1.5	4.5	5.1	4.0
1953	2.3	2.6	2.0	1.9	2.2	1.6	5.7	6.4	5.2
1952	2.6	2.8	2.3	2.2	2.5	1.9	6.0	6.3	5.7
1951	2.8	3.1	2.5	2.3	2.6	2.1	7.3	7.8	7.0
1950	3.1	3.6	2.6	2.6	3.2	2.2	7.4	8.2	6.6
INFECTIONS OF KIDNEY (600)									
1959	5.5	6.2	5.0	5.0	5.6	4.5	10.9	12.2	10.0
1958	5.8	6.6	5.1	5.3	6.1	4.7	10.7	12.1	9.5
1957	5.4	6.1	5.0	5.0	5.6	4.5	10.5	11.2	9.8
1956	5.1	5.8	4.5	4.7	5.4	4.2	8.7	10.0	7.7
1955	4.4	4.9	4.0	4.1	4.7	3.7	7.5	7.9	7.2
1954	3.9	4.2	3.6	3.6	3.9	3.4	6.6	7.2	6.0
1953	3.8	4.4	3.4	3.5	4.0	3.1	7.1	8.6	5.9
1952	3.5	3.9	3.1	3.2	3.7	2.9	5.8	6.6	5.0
1951	3.1	3.6	2.8	2.9	3.4	2.6	5.2	6.0	4.6
1950	3.2	3.5	2.9	3.0	3.3	2.7	5.1	5.6	4.7

Source: CDC/NCHS, National Vital Statistics System, Mortality.

HIST293. AGE-ADJUSTED DEATH RATES FOR APPROXIMATELY 64 SELECTED CAUSES, BY RACE AND SEX: UNITED STATES, 1950-59

[Based on age-specific death rates per 100,000 population in specified group. Computed by the direct method, using as the standard population the age distribution of the total population of the United States as projected for 2000. Numbers after causes of death are category numbers of the seventh revision, International Classification of Diseases]

CAUSE OF DEATH CODE AND YEAR	-- ALL RACES --			-- WHITE --			--- ALL OTHER ---		
	BOTH	MALE	FEMALE	BOTH	MALE	FEMALE	BOTH	MALE	FEMALE
HYPERPLASIA OF PROSTATE (610)									
1959	4.6	10.7	...	4.4	10.4	...	6.4	13.8	...
1958	4.8	11.3	...	4.7	11.0	...	6.7	14.5	...
1957	5.4	12.3	...	5.2	11.9	...	7.8	16.8	...
1956	5.8	13.2	...	5.6	12.9	...	7.7	16.5	...
1955	6.7	15.1	...	6.6	14.9	...	8.0	17.0	...
1954	6.9	15.6	...	6.8	15.4	...	8.0	17.1	...
1953	7.3	16.4	...	7.2	16.2	...	9.1	19.2	...
1952	7.7	17.3	...	7.6	17.1	...	9.3	19.7	...
1951	8.2	18.4	...	8.1	18.1	...	10.2	21.6	...
1950	8.6	19.2	...	8.4	18.9	...	10.9	22.7	...

DELIVERIES AND COMPLICATIONS OF PREGNANCY, CHILDBIRTH AND THE PUERPERIUM (640-689)

1959	1.0	...	1.9	0.6	...	1.3	3.6	...	6.8
1958	1.0	...	1.9	0.6	...	1.3	3.6	...	6.8
1957	1.1	...	2.2	0.7	...	1.4	4.2	...	8.0
1956	1.1	...	2.1	0.7	...	1.4	3.9	...	7.4
1955	1.2	...	2.4	0.8	...	1.6	4.4	...	8.4
1954	1.3	...	2.6	0.9	...	1.8	4.8	...	9.0
1953	1.5	...	3.0	1.1	...	2.1	5.3	...	10.0
1952	1.7	...	3.2	1.2	...	2.3	5.7	...	10.9
1951	1.8	...	3.5	1.3	...	2.5	6.1	...	11.5
1950	1.9	...	3.7	1.3	...	2.6	6.4	...	12.2

... Not applicable

Source: CDC/NCHS, National Vital Statistics System, Mortality.

HIST293. AGE-ADJUSTED DEATH RATES FOR APPROXIMATELY 64 SELECTED CAUSES, BY RACE AND SEX: UNITED STATES, 1950-59

[Based on age-specific death rates per 100,000 population in specified group. Computed by the direct method, using as the standard population the age distribution of the total population of the United States as projected for 2000. Numbers after causes of death are category numbers of the seventh revision, International Classification of Diseases]

CAUSE OF DEATH CODE AND YEAR	-- ALL RACES --			-- WHITE --			--- ALL OTHER ---		
	BOTH	MALE	FEMALE	BOTH	MALE	FEMALE	BOTH	MALE	FEMALE
ABORTION (650-652)									
1959	0.2	...	0.3	0.1	...	0.2	0.8	...	1.5
1958	0.2	...	0.3	0.1	...	0.2	0.7	...	1.2
1957	0.2	...	0.3	0.1	...	0.2	0.7	...	1.4
1956	0.1	...	0.3	0.1	...	0.2	0.5	...	0.9
1955	0.2	...	0.3	0.1	...	0.2	0.7	...	1.2
1954	0.2	...	0.4	0.1	...	0.2	0.7	...	1.4
1953	0.2	...	0.4	0.1	...	0.2	0.8	...	1.4
1952	0.2	...	0.4	0.1	...	0.3	0.7	...	1.4
1951	0.2	...	0.4	0.1	...	0.2	0.8	...	1.5
1950	0.2	...	0.4	0.1	...	0.3	0.7	...	1.3

OTHER COMPLICATIONS OF PREGNANCY, CHILDBIRTH, AND THE PUERPERIUM (640-649, 660-689)

1959	0.8	...	1.6	0.6	...	1.1	2.8	...	5.4
1958	0.8	...	1.6	0.6	...	1.1	2.9	...	5.6
1957	0.9	...	1.8	0.6	...	1.2	3.5	...	6.6
1956	0.9	...	1.8	0.6	...	1.2	3.4	...	6.5
1955	1.0	...	2.0	0.7	...	1.4	3.8	...	7.1
1954	1.2	...	2.3	0.8	...	1.6	4.0	...	7.6
1953	1.3	...	2.6	1.0	...	1.9	4.5	...	8.6
1952	1.5	...	2.9	1.0	...	2.0	5.0	...	9.6
1951	1.6	...	3.1	1.2	...	2.3	5.3	...	10.1
1950	1.7	...	3.3	1.2	...	2.4	5.7	...	10.9

... Not applicable

Source: CDC/NCHS, National Vital Statistics System, Mortality.

HIST293. AGE-ADJUSTED DEATH RATES FOR APPROXIMATELY 64 SELECTED CAUSES, BY RACE AND SEX: UNITED STATES, 1950-59

[Based on age-specific death rates per 100,000 population in specified group. Computed by the direct method, using as the standard population the age distribution of the total population of the United States as projected for 2000. Numbers after causes of death are category numbers of the seventh revision, International Classification of Diseases]

CAUSE OF DEATH CODE AND YEAR	-- ALL RACES --			-- WHITE --			--- ALL OTHER ---		
	BOTH	MALE	FEMALE	BOTH	MALE	FEMALE	BOTH	MALE	FEMALE
CONGENITAL MALFORMATIONS (750-759)									
1959	8.4	9.1	7.7	8.5	9.3	7.8	7.5	8.3	6.8
1958	8.3	9.0	7.7	8.4	9.0	7.8	8.0	8.9	7.1
1957	8.6	9.3	7.9	8.7	9.3	8.1	7.8	8.6	7.0
1956	8.5	9.2	7.8	8.7	9.4	8.0	7.4	8.3	6.5
1955	8.5	9.1	7.8	8.6	9.2	8.0	7.4	8.1	6.6
1954	8.5	9.2	7.8	8.7	9.3	8.1	7.1	8.2	6.0
1953	8.7	9.5	7.9	8.9	9.6	8.2	7.1	8.3	6.0
1952	8.8	9.5	8.1	9.0	9.7	8.4	6.8	7.8	5.9
1951	8.6	9.2	7.9	8.8	9.4	8.3	6.4	7.4	5.4
1950	8.8	9.5	8.1	9.1	9.7	8.4	7.1	7.9	6.4
CERTAIN DISEASES OF EARLY INFANCY (760-776)									
1959	23.0	26.5	19.5	20.6	23.9	17.2	37.1	41.7	32.4
1958	23.6	26.9	20.1	21.0	24.2	17.7	39.0	43.6	34.5
1957	23.0	26.3	19.5	20.6	23.8	17.3	37.1	41.6	32.6
1956	23.3	26.9	19.6	21.0	24.3	17.5	36.9	42.3	31.5
1955	23.6	27.0	20.0	21.4	24.8	17.9	36.8	41.5	32.2
1954	24.1	27.7	20.3	21.9	25.4	18.3	37.4	42.2	32.7
1953	24.9	28.7	20.9	22.8	26.6	18.9	38.3	42.4	34.2
1952	25.8	29.6	21.8	23.8	27.5	19.9	39.3	44.5	34.1
1951	26.4	30.6	22.1	24.6	28.7	20.4	37.9	43.4	32.4
1950	26.8	30.9	22.5	24.9	28.9	20.7	39.4	44.2	34.6

Source: CDC/NCHS, National Vital Statistics System, Mortality.

HIST293. AGE-ADJUSTED DEATH RATES FOR APPROXIMATELY 64 SELECTED CAUSES, BY RACE AND SEX: UNITED STATES, 1950-59

[Based on age-specific death rates per 100,000 population in specified group. Computed by the direct method, using as the standard population the age distribution of the total population of the United States as projected for 2000. Numbers after causes of death are category numbers of the seventh revision, International Classification of Diseases]

CAUSE OF DEATH CODE AND YEAR	-- ALL RACES --			-- WHITE --			--- ALL OTHER ---		
	BOTH	MALE	FEMALE	BOTH	MALE	FEMALE	BOTH	MALE	FEMALE
BIRTH INJURIES, POSTNATAL ASPHYXIA, AND ATELECTASIS (760-762)									
1959	10.3	12.1	8.4	9.6	11.4	7.8	13.8	15.8	11.9
1958	10.8	12.6	8.9	10.1	11.9	8.2	15.0	17.0	13.0
1957	10.4	12.3	8.5	9.8	11.5	7.9	14.5	16.9	12.2
1956	10.7	12.7	8.6	10.1	12.0	8.0	14.3	16.7	11.9
1955	10.8	12.7	8.8	10.2	12.2	8.2	14.1	16.0	12.2
1954	10.9	12.9	8.8	10.4	12.4	8.3	13.9	15.9	11.9
1953	11.1	13.2	8.9	10.6	12.8	8.4	13.9	15.7	12.1
1952	11.1	13.2	8.9	10.7	12.8	8.5	13.7	16.0	11.3
1951	11.3	13.6	8.9	11.1	13.3	8.7	12.8	15.0	10.6
1950	11.0	13.1	8.8	10.8	12.9	8.5	12.7	14.7	10.7
INFECTIONS OF NEWBORN (763-768)									
1959	1.7	2.0	1.4	1.3	1.5	1.1	4.3	5.0	3.6
1958	1.9	2.2	1.6	1.4	1.6	1.2	4.7	5.4	4.0
1957	1.8	2.0	1.5	1.3	1.6	1.1	4.3	4.7	3.8
1956	1.7	1.9	1.4	1.3	1.5	1.1	4.1	4.7	3.4
1955	1.5	1.8	1.3	1.2	1.4	1.0	3.8	4.4	3.2
1954	1.7	1.9	1.4	1.3	1.5	1.1	4.0	4.5	3.4
1953	1.6	1.9	1.4	1.3	1.5	1.1	4.0	4.5	3.6
1952	1.7	1.9	1.4	1.3	1.5	1.1	4.0	4.7	3.3
1951	1.6	1.9	1.4	1.3	1.5	1.1	3.6	4.1	3.0
1950	1.7	1.9	1.5	1.4	1.6	1.2	3.6	4.0	3.3

Source: CDC/NCHS, National Vital Statistics System, Mortality.

HIST293. AGE-ADJUSTED DEATH RATES FOR APPROXIMATELY 64 SELECTED CAUSES, BY RACE AND SEX: UNITED STATES, 1950-59

[Based on age-specific death rates per 100,000 population in specified group. Computed by the direct method, using as the standard population the age distribution of the total population of the United States as projected for 2000. Numbers after causes of death are category numbers of the seventh revision, International Classification of Diseases]

CAUSE OF DEATH CODE AND YEAR	-- ALL RACES --			-- WHITE --			--- ALL OTHER ---		
	BOTH	MALE	FEMALE	BOTH	MALE	FEMALE	BOTH	MALE	FEMALE
OTHER DISEASES PECULIAR TO EARLY INFANCY, AND IMMATURITY UNQUALIFIED (769-776)									
1959	11.1	12.4	9.6	9.7	11.0	8.3	18.9	20.9	16.9
1958	10.9	12.2	9.6	9.5	10.7	8.2	19.3	21.1	17.5
1957	10.8	12.0	9.5	9.5	10.7	8.3	18.3	20.0	16.5
1956	10.9	12.3	9.5	9.7	10.9	8.4	18.5	20.9	16.1
1955	11.3	12.5	9.9	10.0	11.2	8.8	18.9	21.1	16.8
1954	11.5	12.9	10.2	10.3	11.5	9.0	19.6	21.9	17.4
1953	12.2	13.6	10.6	10.9	12.3	9.4	20.4	22.2	18.5
1952	13.0	14.5	11.5	11.8	13.2	10.3	21.6	23.8	19.4
1951	13.5	15.2	11.7	12.2	13.8	10.6	21.5	24.2	18.8
1950	14.1	15.8	12.3	12.7	14.4	11.0	23.1	25.5	20.6
SYMPTOMS, SENILITY, AND ILL-DEFINED CONDITIONS (780-795)									
1959	15.2	18.7	12.2	11.7	14.7	9.1	52.8	60.9	45.6
1958	16.3	19.9	13.2	12.5	15.4	10.0	56.5	67.0	48.0
1957	16.6	20.2	13.4	12.8	15.9	10.2	57.0	66.8	48.8
1956	16.7	20.4	13.4	12.9	16.2	10.1	56.9	64.5	49.8
1955	17.9	21.8	14.5	14.0	17.5	10.9	60.7	68.2	53.7
1954	19.2	23.2	15.6	15.0	18.4	12.0	64.9	75.8	55.6
1953	21.8	25.8	18.3	16.7	20.0	13.6	80.1	90.9	70.4
1952	22.8	26.5	19.5	17.5	20.7	14.5	82.9	91.5	74.9
1951	24.9	29.2	21.0	19.2	22.7	15.9	89.5	101.4	78.9
1950	26.0	30.3	22.2	20.0	23.4	16.9	95.8	108.5	84.1

Source: CDC/NCHS, National Vital Statistics System, Mortality.



HIST293. AGE-ADJUSTED DEATH RATES FOR APPROXIMATELY 64 SELECTED CAUSES, BY RACE AND SEX: UNITED STATES, 1950-59

[Based on age-specific death rates per 100,000 population in specified group. Computed by the direct method, using as the standard population the age distribution of the total population of the United States as projected for 2000. Numbers after causes of death are category numbers of the seventh revision, International Classification of Diseases]

CAUSE OF DEATH CODE AND YEAR	-- ALL RACES --			-- WHITE --			--- ALL OTHER ---		
	BOTH	MALE	FEMALE	BOTH	MALE	FEMALE	BOTH	MALE	FEMALE
ALL OTHER DISEASES (RESIDUAL)									
1959	47.1	60.6	35.6	45.9	59.4	34.3	55.7	68.6	44.0
1958	46.4	59.6	34.9	45.2	58.5	33.7	54.5	66.5	43.6
1957	58.4	72.6	46.1	56.7	70.9	44.4	71.8	86.5	58.5
1956	55.9	68.3	44.9	54.5	67.0	43.5	65.8	77.7	54.8
1955	54.7	65.9	44.7	53.4	64.7	43.4	63.5	74.6	53.1
1954	53.5	63.8	44.3	52.2	62.6	43.0	63.0	72.6	53.9
1953	55.7	66.5	46.1	54.3	64.8	44.8	67.1	80.0	55.3
1952	56.7	66.4	47.9	55.3	64.8	46.6	67.6	79.2	56.6
1951	57.3	67.2	48.4	56.0	66.0	47.0	67.4	76.0	59.4
1950	55.7	63.7	48.5	54.4	62.4	47.2	66.4	75.0	58.0
ACCIDENTS AND ADVERSE EFFECTS (E800-E962)									
1959	63.2	86.8	40.4	61.5	84.2	39.7	73.8	108.0	42.3
1958	63.6	86.9	41.1	61.9	84.2	40.2	75.3	109.1	44.4
1957	69.1	93.7	45.3	67.4	90.9	44.5	80.4	116.3	47.5
1956	70.2	95.6	45.7	68.6	92.6	45.2	81.1	119.5	45.7
1955	70.9	95.5	47.0	69.4	92.9	46.4	80.7	115.7	48.1
1954	69.9	94.0	46.5	68.4	91.4	46.0	79.7	115.6	47.0
1953	75.4	101.6	50.0	73.6	98.5	49.4	87.3	127.5	49.9
1952	78.0	103.9	52.9	76.4	100.7	52.4	89.4	130.3	51.5
1951	79.7	105.8	54.0	78.3	103.2	53.6	88.8	127.3	52.6
1950	78.4	102.2	54.5	77.4	100.1	54.4	82.9	118.6	48.9

Source: CDC/NCHS, National Vital Statistics System, Mortality.

HIST293. AGE-ADJUSTED DEATH RATES FOR APPROXIMATELY 64 SELECTED CAUSES, BY RACE AND SEX: UNITED STATES, 1950-59

[Based on age-specific death rates per 100,000 population in specified group. Computed by the direct method, using as the standard population the age distribution of the total population of the United States as projected for 2000. Numbers after causes of death are category numbers of the seventh revision, International Classification of Diseases]

CAUSE OF DEATH CODE AND YEAR	-- ALL RACES --			-- WHITE --			--- ALL OTHER ---		
	BOTH	MALE	FEMALE	BOTH	MALE	FEMALE	BOTH	MALE	FEMALE
MOTOR VEHICLE ACCIDENTS (E810-E835)									
1959	23.4	35.9	11.6	23.1	35.3	11.6	25.7	41.6	11.1
1958	23.2	35.9	11.4	23.0	35.3	11.5	25.2	41.2	10.6
1957	24.8	38.5	12.0	24.4	37.8	11.9	28.4	45.5	12.9
1956	25.8	40.3	12.3	25.3	39.4	12.2	30.3	49.4	12.7
1955	25.4	39.5	12.2	25.0	38.8	12.1	29.0	46.3	13.1
1954	24.0	37.4	11.5	23.6	36.6	11.4	27.8	44.6	12.4
1953	25.9	40.5	12.4	25.5	39.5	12.4	30.6	50.1	12.7
1952	26.1	40.7	12.4	25.6	39.8	12.3	30.3	49.2	12.9
1951	25.9	40.5	12.2	25.5	39.7	12.0	29.7	47.6	13.1
1950	24.6	38.5	11.5	24.4	37.9	11.4	27.1	43.5	11.7
OTHER ACCIDENTS (E800-E802, E840-E962)									
1959	39.8	50.9	28.8	38.4	48.9	28.0	48.2	66.3	31.3
1958	40.3	51.0	29.7	38.9	48.9	28.8	50.1	67.9	33.8
1957	44.3	55.2	33.3	42.9	53.1	32.6	52.0	70.8	34.6
1956	44.4	55.2	33.4	43.2	53.3	32.9	50.8	70.1	33.0
1955	45.5	56.0	34.8	44.4	54.2	34.4	51.7	69.4	34.9
1954	45.9	56.6	35.0	44.8	54.8	34.6	52.0	70.9	34.7
1953	49.4	61.1	37.6	48.2	58.9	37.1	56.7	77.5	37.2
1952	52.0	63.2	40.4	50.7	60.9	40.1	59.1	81.1	38.6
1951	53.9	65.3	41.9	52.8	63.4	41.6	59.1	79.7	39.6
1950	53.8	63.8	43.0	53.0	62.2	43.0	55.8	75.2	37.2

Source: CDC/NCHS, National Vital Statistics System, Mortality.

HIST293. AGE-ADJUSTED DEATH RATES FOR APPROXIMATELY 64 SELECTED CAUSES, BY RACE AND SEX: UNITED STATES, 1950-59

[Based on age-specific death rates per 100,000 population in specified group. Computed by the direct method, using as the standard population the age distribution of the total population of the United States as projected for 2000. Numbers after causes of death are category numbers of the seventh revision, International Classification of Diseases]

CAUSE OF DEATH CODE AND YEAR	-- ALL RACES --			-- WHITE --			--- ALL OTHER ---		
	BOTH	MALE	FEMALE	BOTH	MALE	FEMALE	BOTH	MALE	FEMALE
SUICIDE (E963, E970-E979)									
1959	12.3	20.2	5.2	13.0	21.2	5.5	6.0	10.0	2.4
1958	12.4	20.4	5.3	13.1	21.4	5.6	5.6	9.2	2.2
1957	11.4	18.7	4.8	12.0	19.7	5.1	5.2	8.9	1.8
1956	11.6	19.0	4.9	12.3	20.0	5.2	4.7	7.7	2.0
1955	11.8	19.2	5.1	12.5	20.3	5.4	4.7	7.8	1.7
1954	11.7	19.4	4.6	12.3	20.4	4.9	5.0	8.4	1.8
1953	11.7	19.3	4.7	12.3	20.3	5.0	4.8	8.4	1.4
1952	11.6	18.9	4.9	12.3	20.0	5.2	4.5	7.7	1.5
1951	12.0	19.4	5.2	12.7	20.4	5.4	4.9	8.0	1.9
1950	13.2	21.2	5.6	13.9	22.3	6.0	5.2	8.6	1.9
HOMICIDE (E964, E980-E985)									
1959	5.0	7.7	2.5	2.7	3.8	1.5	25.4	41.8	10.4
1958	4.9	7.5	2.5	2.6	3.7	1.5	24.9	41.0	10.2
1957	4.8	7.4	2.3	2.4	3.5	1.4	25.5	42.5	9.9
1956	4.9	7.5	2.4	2.4	3.5	1.3	25.9	42.4	10.9
1955	4.8	7.5	2.3	2.5	3.6	1.3	25.2	41.7	10.1
1954	5.1	7.9	2.3	2.6	3.7	1.4	26.9	45.3	10.1
1953	5.1	7.9	2.3	2.6	3.7	1.4	27.0	45.5	10.0
1952	5.4	8.5	2.4	2.6	3.9	1.4	29.2	49.2	10.9
1951	5.0	7.8	2.4	2.5	3.7	1.4	26.8	44.4	10.6
1950	5.3	8.3	2.4	2.7	4.0	1.4	28.6	47.6	10.9

Source: CDC/NCHS, National Vital Statistics System, Mortality.