

LCWK4. Infant, neonatal, and postneonatal deaths, percent of total deaths, and mortality rates for the 15 leading causes of infant death, by race and Hispanic origin, and sex: United States, 2015

[Rates are infant (under 1 year), neonatal (under 28 days), and postneonatal (28 days through 11 months) deaths per 100,000 live births in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of inconsistencies between reporting Hispanic origin on birth and death certificates and because deaths are based on race of decedent whereas live births are based on race of mother.]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race and sex	Number	Percent of total deaths	Mortality rate
Infants, all races and origins, both sexes				
...	All causes	23,455	100.0	589.5
1	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	4,825	20.6	121.3
2	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	4,084	17.4	102.7
3	Sudden infant death syndrome (R95)	1,568	6.7	39.4
4	Newborn affected by maternal complications of pregnancy (P01)	1,522	6.5	38.3
5	Accidents (unintentional injuries) (V01-X59)	1,291	5.5	32.4
6	Newborn affected by complications of placenta, cord and membranes (P02)	910	3.9	22.9
7	Bacterial sepsis of newborn (P36)	599	2.6	15.1
8	Respiratory distress of newborn (P22)	462	2.0	11.6
9	Diseases of the circulatory system (I00-I99)	428	1.8	10.8
10	Neonatal hemorrhage (P50-P52,P54)	406	1.7	10.2
11	Necrotizing enterocolitis of newborn (P77)	363	1.5	9.1
12	Intrauterine hypoxia and birth asphyxia (P20-P21)	314	1.3	7.9
13	Atelectasis (P28.0-P28.1)	273	1.2	6.9
14	Assault (homicide) (*U01,X85-Y09)	263	1.1	6.6
15	Diarrhea and gastroenteritis of infectious origin (A09)	210	0.9	5.3
...	All other causes (residual)	5,937	25.3	149.2
Infants, all races and origins, male				
...	All causes	13,008	100.0	638.8
1	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	2,551	19.6	125.3
2	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	2,249	17.3	110.5
3	Sudden infant death syndrome (R95)	923	7.1	45.3
4	Newborn affected by maternal complications of pregnancy (P01)	854	6.6	41.9
5	Accidents (unintentional injuries) (V01-X59)	755	5.8	37.1
6	Newborn affected by complications of placenta, cord and membranes (P02)	514	4.0	25.2
7	Bacterial sepsis of newborn (P36)	313	2.4	15.4
8	Respiratory distress of newborn (P22)	282	2.2	13.8
9	Neonatal hemorrhage (P50-P52,P54)	235	1.8	11.5
10	Diseases of the circulatory system (I00-I99)	226	1.7	11.1
11	Necrotizing enterocolitis of newborn (P77)	194	1.5	9.5
12	Intrauterine hypoxia and birth asphyxia (P20-P21)	162	1.2	8.0
12	Atelectasis (P28.0-P28.1)	162	1.2	8.0
14	Assault (homicide) (*U01,X85-Y09)	144	1.1	7.1
15	Influenza and pneumonia (J09-J18)	111	0.9	5.5
...	All other causes (residual)	3,333	25.6	163.7

... Category not applicable

\* Figure does not meet standards of reliability or precision

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015

LCWK4. Infant, neonatal, and postneonatal deaths, percent of total deaths, and mortality rates for the 15 leading causes of infant death, by race and Hispanic origin, and sex: United States, 2015

[Rates are infant (under 1 year), neonatal (under 28 days), and postneonatal (28 days through 11 months) deaths per 100,000 live births in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of inconsistencies between reporting Hispanic origin on birth and death certificates and because deaths are based on race of decedent whereas live births are based on race of mother.]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race and sex	Number	Percent of total deaths	Mortality rate
Infants, all races and origins, female				
...	All causes	10,447	100.0	537.9
1	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	2,274	21.8	117.1
2	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	1,835	17.6	94.5
3	Newborn affected by maternal complications of pregnancy (P01)	668	6.4	34.4
4	Sudden infant death syndrome (R95)	645	6.2	33.2
5	Accidents (unintentional injuries) (V01-X59)	536	5.1	27.6
6	Newborn affected by complications of placenta, cord and membranes (P02)	396	3.8	20.4
7	Bacterial sepsis of newborn (P36)	286	2.7	14.7
8	Diseases of the circulatory system (I00-I99)	202	1.9	10.4
9	Respiratory distress of newborn (P22)	180	1.7	9.3
10	Neonatal hemorrhage (P50-P52,P54)	171	1.6	8.8
11	Necrotizing enterocolitis of newborn (P77)	169	1.6	8.7
12	Intrauterine hypoxia and birth asphyxia (P20-P21)	152	1.5	7.8
13	Assault (homicide) (*U01,X85-Y09)	119	1.1	6.1
14	Atelectasis (P28.0-P28.1)	111	1.1	5.7
15	Diarrhea and gastroenteritis of infectious origin (A09)	100	1.0	5.1
...	All other causes (residual)	2,603	24.9	134.0
Infants, Hispanic, both sexes				
...	All causes	4,805	100.0	520.0
1	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	1,216	25.3	131.6
2	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	824	17.1	89.2
3	Newborn affected by maternal complications of pregnancy (P01)	339	7.1	36.7
4	Sudden infant death syndrome (R95)	202	4.2	21.9
5	Newborn affected by complications of placenta, cord and membranes (P02)	194	4.0	21.0
6	Accidents (unintentional injuries) (V01-X59)	187	3.9	20.2
7	Bacterial sepsis of newborn (P36)	127	2.6	13.7
8	Neonatal hemorrhage (P50-P52,P54)	99	2.1	10.7
9	Respiratory distress of newborn (P22)	87	1.8	9.4
10	Necrotizing enterocolitis of newborn (P77)	82	1.7	8.9
11	Diseases of the circulatory system (I00-I99)	74	1.5	8.0
12	Atelectasis (P28.0-P28.1)	62	1.3	6.7
13	Intrauterine hypoxia and birth asphyxia (P20-P21)	56	1.2	6.1
14	Hydrops fetalis not due to hemolytic disease (P83.2)	47	1.0	5.1
15	Diarrhea and gastroenteritis of infectious origin (A09)	42	0.9	4.5
...	All other causes (residual)	1,167	24.3	126.3

... Category not applicable

\* Figure does not meet standards of reliability or precision

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015

LCWK4. Infant, neonatal, and postneonatal deaths, percent of total deaths, and mortality rates for the 15 leading causes of infant death, by race and Hispanic origin, and sex: United States, 2015

[Rates are infant (under 1 year), neonatal (under 28 days), and postneonatal (28 days through 11 months) deaths per 100,000 live births in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of inconsistencies between reporting Hispanic origin on birth and death certificates and because deaths are based on race of decedent whereas live births are based on race of mother.]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race and sex	Number	Percent of total deaths	Mortality rate
Infants, Hispanic, male				
...	All causes	2,619	100.0	555.5
1	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	635	24.2	134.7
2	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	464	17.7	98.4
3	Newborn affected by maternal complications of pregnancy (P01)	208	7.9	44.1
4	Sudden infant death syndrome (R95)	115	4.4	24.4
5	Newborn affected by complications of placenta, cord and membranes (P02)	104	4.0	22.1
5	Accidents (unintentional injuries) (V01-X59)	104	4.0	22.1
7	Bacterial sepsis of newborn (P36)	70	2.7	14.8
8	Neonatal hemorrhage (P50-P52,P54)	54	2.1	11.5
9	Respiratory distress of newborn (P22)	51	1.9	10.8
10	Necrotizing enterocolitis of newborn (P77)	43	1.6	9.1
11	Atelectasis (P28.0-P28.1)	35	1.3	7.4
12	Diseases of the circulatory system (I00-I99)	29	1.1	6.2
13	Intrauterine hypoxia and birth asphyxia (P20-P21)	25	1.0	5.3
14	Septicemia (A40-A41)	20	0.8	4.2
14	Influenza and pneumonia (J09-J18)	20	0.8	4.2
14	Hydrops fetalis not due to hemolytic disease (P83.2)	20	0.8	4.2
...	All other causes (residual)	622	23.7	131.9
Infants, Hispanic, female				
...	All causes	2,186	100.0	483.0
1	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	581	26.6	128.4
2	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	360	16.5	79.5
3	Newborn affected by maternal complications of pregnancy (P01)	131	6.0	28.9
4	Newborn affected by complications of placenta, cord and membranes (P02)	90	4.1	19.9
5	Sudden infant death syndrome (R95)	87	4.0	19.2
6	Accidents (unintentional injuries) (V01-X59)	83	3.8	18.3
7	Bacterial sepsis of newborn (P36)	57	2.6	12.6
8	Diseases of the circulatory system (I00-I99)	45	2.1	9.9
8	Neonatal hemorrhage (P50-P52,P54)	45	2.1	9.9
10	Necrotizing enterocolitis of newborn (P77)	39	1.8	8.6
11	Respiratory distress of newborn (P22)	36	1.6	8.0
12	Intrauterine hypoxia and birth asphyxia (P20-P21)	31	1.4	6.8
13	Atelectasis (P28.0-P28.1)	27	1.2	6.0
13	Hydrops fetalis not due to hemolytic disease (P83.2)	27	1.2	6.0
15	Diarrhea and gastroenteritis of infectious origin (A09)	25	1.1	5.5
...	All other causes (residual)	522	23.9	115.3

... Category not applicable

\* Figure does not meet standards of reliability or precision

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015

LCWK4. Infant, neonatal, and postneonatal deaths, percent of total deaths, and mortality rates for the 15 leading causes of infant death, by race and Hispanic origin, and sex: United States, 2015

[Rates are infant (under 1 year), neonatal (under 28 days), and postneonatal (28 days through 11 months) deaths per 100,000 live births in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of inconsistencies between reporting Hispanic origin on birth and death certificates and because deaths are based on race of decedent whereas live births are based on race of mother.]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race and sex	Number	Percent of total deaths	Mortality rate
Infants, non-Hispanic, white, both sexes				
...	All causes	10,277	100.0	482.4
1	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	2,373	23.1	111.4
2	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	1,485	14.4	69.7
3	Sudden infant death syndrome (R95)	771	7.5	36.2
4	Accidents (unintentional injuries) (V01-X59)	632	6.1	29.7
5	Newborn affected by maternal complications of pregnancy (P01)	606	5.9	28.4
6	Newborn affected by complications of placenta, cord and membranes (P02)	379	3.7	17.8
7	Bacterial sepsis of newborn (P36)	230	2.2	10.8
8	Diseases of the circulatory system (I00-I99)	196	1.9	9.2
9	Respiratory distress of newborn (P22)	184	1.8	8.6
10	Neonatal hemorrhage (P50-P52,P54)	177	1.7	8.3
11	Intrauterine hypoxia and birth asphyxia (P20-P21)	150	1.5	7.0
12	Necrotizing enterocolitis of newborn (P77)	133	1.3	6.2
13	Assault (homicide) (*U01,X85-Y09)	119	1.2	5.6
14	Atelectasis (P28.0-P28.1)	118	1.1	5.5
15	Hydrops fetalis not due to hemolytic disease (P83.2)	95	0.9	4.5
...	All other causes (residual)	2,629	25.6	123.4
Infants, non-Hispanic, white, male				
...	All causes	5,758	100.0	526.9
1	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	1,263	21.9	115.6
2	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	784	13.6	71.7
3	Sudden infant death syndrome (R95)	468	8.1	42.8
4	Accidents (unintentional injuries) (V01-X59)	395	6.9	36.1
5	Newborn affected by maternal complications of pregnancy (P01)	322	5.6	29.5
6	Newborn affected by complications of placenta, cord and membranes (P02)	216	3.8	19.8
7	Bacterial sepsis of newborn (P36)	123	2.1	11.3
8	Diseases of the circulatory system (I00-I99)	117	2.0	10.7
9	Respiratory distress of newborn (P22)	116	2.0	10.6
10	Neonatal hemorrhage (P50-P52,P54)	107	1.9	9.8
11	Intrauterine hypoxia and birth asphyxia (P20-P21)	85	1.5	7.8
12	Atelectasis (P28.0-P28.1)	75	1.3	6.9
13	Necrotizing enterocolitis of newborn (P77)	74	1.3	6.8
13	Assault (homicide) (*U01,X85-Y09)	74	1.3	6.8
15	Hydrops fetalis not due to hemolytic disease (P83.2)	54	0.9	4.9
...	All other causes (residual)	1,485	25.8	135.9

... Category not applicable

\* Figure does not meet standards of reliability or precision

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015

LCWK4. Infant, neonatal, and postneonatal deaths, percent of total deaths, and mortality rates for the 15 leading causes of infant death, by race and Hispanic origin, and sex: United States, 2015

[Rates are infant (under 1 year), neonatal (under 28 days), and postneonatal (28 days through 11 months) deaths per 100,000 live births in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of inconsistencies between reporting Hispanic origin on birth and death certificates and because deaths are based on race of decedent whereas live births are based on race of mother.]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race and sex	Number	Percent of total deaths	Mortality rate
Infants, non-Hispanic, white, female				
...	All causes	4,519	100.0	435.6
1	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	1,110	24.6	107.0
2	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	701	15.5	67.6
3	Sudden infant death syndrome (R95)	303	6.7	29.2
4	Newborn affected by maternal complications of pregnancy (P01)	284	6.3	27.4
5	Accidents (unintentional injuries) (V01-X59)	237	5.2	22.8
6	Newborn affected by complications of placenta, cord and membranes (P02)	163	3.6	15.7
7	Bacterial sepsis of newborn (P36)	107	2.4	10.3
8	Diseases of the circulatory system (I00-I99)	79	1.7	7.6
9	Neonatal hemorrhage (P50-P52,P54)	70	1.5	6.7
10	Respiratory distress of newborn (P22)	68	1.5	6.6
11	Intrauterine hypoxia and birth asphyxia (P20-P21)	65	1.4	6.3
12	Necrotizing enterocolitis of newborn (P77)	59	1.3	5.7
13	Assault (homicide) (*U01,X85-Y09)	45	1.0	4.3
14	Atelectasis (P28.0-P28.1)	43	1.0	4.1
15	Hydrops fetalis not due to hemolytic disease (P83.2)	41	0.9	4.0
...	All other causes (residual)	1,144	25.3	110.3
Infants, non-Hispanic, black, both sexes				
...	All causes	6,907	100.0	1,172.6
1	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	1,513	21.9	256.9
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	930	13.5	157.9
3	Sudden infant death syndrome (R95)	513	7.4	87.1
4	Newborn affected by maternal complications of pregnancy (P01)	458	6.6	77.8
5	Accidents (unintentional injuries) (V01-X59)	410	5.9	69.6
6	Newborn affected by complications of placenta, cord and membranes (P02)	270	3.9	45.8
7	Bacterial sepsis of newborn (P36)	208	3.0	35.3
8	Respiratory distress of newborn (P22)	168	2.4	28.5
9	Diseases of the circulatory system (I00-I99)	128	1.9	21.7
10	Necrotizing enterocolitis of newborn (P77)	126	1.8	21.4
11	Neonatal hemorrhage (P50-P52,P54)	99	1.4	16.8
12	Intrauterine hypoxia and birth asphyxia (P20-P21)	93	1.3	15.8
13	Assault (homicide) (*U01,X85-Y09)	92	1.3	15.6
14	Atelectasis (P28.0-P28.1)	84	1.2	14.3
15	Diarrhea and gastroenteritis of infectious origin (A09)	80	1.2	13.6
...	All other causes (residual)	1,735	25.1	294.5

... Category not applicable

\* Figure does not meet standards of reliability or precision

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015

LCWK4. Infant, neonatal, and postneonatal deaths, percent of total deaths, and mortality rates for the 15 leading causes of infant death, by race and Hispanic origin, and sex: United States, 2015

[Rates are infant (under 1 year), neonatal (under 28 days), and postneonatal (28 days through 11 months) deaths per 100,000 live births in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of inconsistencies between reporting Hispanic origin on birth and death certificates and because deaths are based on race of decedent whereas live births are based on race of mother.]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race and sex	Number	Percent of total deaths	Mortality rate
Infants, non-Hispanic, black, male				
...	All causes	3,815	100.0	1,274.6
1	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	844	22.1	282.0
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	493	12.9	164.7
3	Sudden infant death syndrome (R95)	289	7.6	96.6
4	Newborn affected by maternal complications of pregnancy (P01)	268	7.0	89.5
5	Accidents (unintentional injuries) (V01-X59)	222	5.8	74.2
6	Newborn affected by complications of placenta, cord and membranes (P02)	157	4.1	52.5
7	Bacterial sepsis of newborn (P36)	106	2.8	35.4
8	Respiratory distress of newborn (P22)	99	2.6	33.1
9	Necrotizing enterocolitis of newborn (P77)	65	1.7	21.7
10	Diseases of the circulatory system (I00-I99)	63	1.7	21.0
11	Neonatal hemorrhage (P50-P52,P54)	55	1.4	18.4
12	Diarrhea and gastroenteritis of infectious origin (A09)	46	1.2	15.4
12	Atelectasis (P28.0-P28.1)	46	1.2	15.4
14	Intrauterine hypoxia and birth asphyxia (P20-P21)	44	1.2	14.7
14	Assault (homicide) (*U01,X85-Y09)	44	1.2	14.7
...	All other causes (residual)	974	25.5	325.4
Infants, non-Hispanic, black, female				
...	All causes	3,092	100.0	1,067.2
1	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	669	21.6	230.9
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	437	14.1	150.8
3	Sudden infant death syndrome (R95)	224	7.2	77.3
4	Newborn affected by maternal complications of pregnancy (P01)	190	6.1	65.6
5	Accidents (unintentional injuries) (V01-X59)	188	6.1	64.9
6	Newborn affected by complications of placenta, cord and membranes (P02)	113	3.7	39.0
7	Bacterial sepsis of newborn (P36)	102	3.3	35.2
8	Respiratory distress of newborn (P22)	69	2.2	23.8
9	Diseases of the circulatory system (I00-I99)	65	2.1	22.4
10	Necrotizing enterocolitis of newborn (P77)	61	2.0	21.1
11	Intrauterine hypoxia and birth asphyxia (P20-P21)	49	1.6	16.9
12	Assault (homicide) (*U01,X85-Y09)	48	1.6	16.6
13	Neonatal hemorrhage (P50-P52,P54)	44	1.4	15.2
14	Atelectasis (P28.0-P28.1)	38	1.2	13.1
15	Diarrhea and gastroenteritis of infectious origin (A09)	34	1.1	11.7
...	All other causes (residual)	761	24.6	262.7

... Category not applicable

\* Figure does not meet standards of reliability or precision

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015

LCWK4. Infant, neonatal, and postneonatal deaths, percent of total deaths, and mortality rates for the 15 leading causes of infant death, by race and Hispanic origin, and sex: United States, 2015

[Rates are infant (under 1 year), neonatal (under 28 days), and postneonatal (28 days through 11 months) deaths per 100,000 live births in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of inconsistencies between reporting Hispanic origin on birth and death certificates and because deaths are based on race of decedent whereas live births are based on race of mother.]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race and sex	Number	Percent of total deaths	Mortality rate
Infants, non-Hispanic, American Indian or Alaska Native, both sexes				
...	All causes	300	100.0	814.7
1	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	58	19.3	157.5
2	Sudden infant death syndrome (R95)	49	16.3	133.1
3	Accidents (unintentional injuries) (V01-X59)	32	10.7	86.9
4	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	25	8.3	67.9
5	Newborn affected by maternal complications of pregnancy (P01)	15	5.0	*
6	Newborn affected by complications of placenta, cord and membranes (P02)	11	3.7	*
7	Bacterial sepsis of newborn (P36)	7	2.3	*
8	Diseases of the circulatory system (I00-I99)	5	1.7	*
9	Influenza and pneumonia (J09-J18)	4	1.3	*
9	Intrauterine hypoxia and birth asphyxia (P20-P21)	4	1.3	*
11	Acute bronchitis and acute bronchiolitis (J20-J21)	3	1.0	*
11	Newborn affected by maternal hypertensive disorders (P00.0)	3	1.0	*
11	Newborn affected by other complications of labor and delivery (P03)	3	1.0	*
11	Atelectasis (P28.0-P28.1)	3	1.0	*
11	Necrotizing enterocolitis of newborn (P77)	3	1.0	*
11	Assault (homicide) (*U01,X85-Y09)	3	1.0	*
...	All other causes (residual)	72	24.0	195.5
Infants, non-Hispanic, American Indian or Alaska Native, male				
...	All causes	161	100.0	853.4
1	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	31	19.3	164.3
2	Sudden infant death syndrome (R95)	28	17.4	148.4
3	Accidents (unintentional injuries) (V01-X59)	17	10.6	*
4	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	14	8.7	*
5	Newborn affected by maternal complications of pregnancy (P01)	10	6.2	*
6	Newborn affected by complications of placenta, cord and membranes (P02)	6	3.7	*
7	Bacterial sepsis of newborn (P36)	4	2.5	*
8	Diseases of the circulatory system (I00-I99)	2	1.2	*
8	Influenza and pneumonia (J09-J18)	2	1.2	*
8	Acute bronchitis and acute bronchiolitis (J20-J21)	2	1.2	*
8	Newborn affected by other complications of labor and delivery (P03)	2	1.2	*
8	Intrauterine hypoxia and birth asphyxia (P20-P21)	2	1.2	*
8	Respiratory distress of newborn (P22)	2	1.2	*
8	Atelectasis (P28.0-P28.1)	2	1.2	*
8	Assault (homicide) (*U01,X85-Y09)	2	1.2	*
...	All other causes (residual)	35	21.7	185.5

... Category not applicable

\* Figure does not meet standards of reliability or precision

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015

LCWK4. Infant, neonatal, and postneonatal deaths, percent of total deaths, and mortality rates for the 15 leading causes of infant death, by race and Hispanic origin, and sex: United States, 2015

[Rates are infant (under 1 year), neonatal (under 28 days), and postneonatal (28 days through 11 months) deaths per 100,000 live births in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of inconsistencies between reporting Hispanic origin on birth and death certificates and because deaths are based on race of decedent whereas live births are based on race of mother.]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race and sex	Number	Percent of total deaths	Mortality rate
Infants, non-Hispanic, American Indian or Alaska Native, female				
...	All causes	139	100.0	773.9
1	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	27	19.4	150.3
2	Sudden infant death syndrome (R95)	21	15.1	116.9
3	Accidents (unintentional injuries) (V01-X59)	15	10.8	*
4	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	11	7.9	*
5	Newborn affected by maternal complications of pregnancy (P01)	5	3.6	*
5	Newborn affected by complications of placenta, cord and membranes (P02)	5	3.6	*
7	Diseases of the circulatory system (I00-I99)	3	2.2	*
7	Bacterial sepsis of newborn (P36)	3	2.2	*
7	Necrotizing enterocolitis of newborn (P77)	3	2.2	*
10	Meningitis (G00,G03)	2	1.4	*
10	Influenza and pneumonia (J09-J18)	2	1.4	*
10	Newborn affected by maternal hypertensive disorders (P00.0)	2	1.4	*
10	Intrauterine hypoxia and birth asphyxia (P20-P21)	2	1.4	*
10	Pulmonary hemorrhage originating in the perinatal period (P26)	2	1.4	*
15	Septicemia (A40-A41)	1	0.7	*
15	Diseases of the blood and blood-forming organs and certain disorders involving the immune mechanism (D50-D89)	1	0.7	*
15	Short stature, not elsewhere classified (E34.3)	1	0.7	*
15	Acute bronchitis and acute bronchiolitis (J20-J21)	1	0.7	*
15	Newborn affected by other maternal conditions which may be unrelated to present pregnancy (P00.1-P00.9)	1	0.7	*
15	Newborn affected by other complications of labor and delivery (P03)	1	0.7	*
15	Slow fetal growth and fetal malnutrition (P05)	1	0.7	*
15	Congenital pneumonia (P23)	1	0.7	*
15	Interstitial emphysema and related conditions originating in the perinatal period (P25)	1	0.7	*
15	Atelectasis (P28.0-P28.1)	1	0.7	*
15	Neonatal hemorrhage (P50-P52,P54)	1	0.7	*
15	Hematological disorders (P60-P61)	1	0.7	*
15	Hydrops fetalis not due to hemolytic disease (P83.2)	1	0.7	*
15	Assault (homicide) (*U01,X85-Y09)	1	0.7	*
...	All other causes (residual)	22	15.8	122.5

... Category not applicable

\* Figure does not meet standards of reliability or precision

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015

LCWK4. Infant, neonatal, and postneonatal deaths, percent of total deaths, and mortality rates for the 15 leading causes of infant death, by race and Hispanic origin, and sex: United States, 2015

[Rates are infant (under 1 year), neonatal (under 28 days), and postneonatal (28 days through 11 months) deaths per 100,000 live births in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of inconsistencies between reporting Hispanic origin on birth and death certificates and because deaths are based on race of decedent whereas live births are based on race of mother.]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race and sex	Number	Percent of total deaths	Mortality rate
Infants, non-Hispanic, Asian or Pacific Islander, both sexes				
...	All causes	892	100.0	335.5
1	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	211	23.7	79.4
2	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	142	15.9	53.4
3	Newborn affected by maternal complications of pregnancy (P01)	83	9.3	31.2
4	Newborn affected by complications of placenta, cord and membranes (P02)	39	4.4	14.7
5	Neonatal hemorrhage (P50-P52,P54)	26	2.9	9.8
6	Sudden infant death syndrome (R95)	25	2.8	9.4
7	Bacterial sepsis of newborn (P36)	23	2.6	8.7
7	Accidents (unintentional injuries) (V01-X59)	23	2.6	8.7
9	Diseases of the circulatory system (I00-I99)	20	2.2	7.5
10	Respiratory distress of newborn (P22)	19	2.1	*
11	Diarrhea and gastroenteritis of infectious origin (A09)	13	1.5	*
11	Necrotizing enterocolitis of newborn (P77)	13	1.5	*
13	Newborn affected by other complications of labor and delivery (P03)	12	1.3	*
14	Hydrops fetalis not due to hemolytic disease (P83.2)	11	1.2	*
15	Intrauterine hypoxia and birth asphyxia (P20-P21)	9	1.0	*
...	All other causes (residual)	223	25.0	83.9
Infants, non-Hispanic, Asian or Pacific Islander, male				
...	All causes	501	100.0	365.7
1	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	105	21.0	76.6
2	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	86	17.2	62.8
3	Newborn affected by maternal complications of pregnancy (P01)	37	7.4	27.0
4	Newborn affected by complications of placenta, cord and membranes (P02)	20	4.0	14.6
5	Neonatal hemorrhage (P50-P52,P54)	17	3.4	*
5	Sudden infant death syndrome (R95)	17	3.4	*
7	Accidents (unintentional injuries) (V01-X59)	15	3.0	*
8	Diseases of the circulatory system (I00-I99)	13	2.6	*
8	Respiratory distress of newborn (P22)	13	2.6	*
10	Diarrhea and gastroenteritis of infectious origin (A09)	8	1.6	*
10	Newborn affected by other complications of labor and delivery (P03)	8	1.6	*
10	Bacterial sepsis of newborn (P36)	8	1.6	*
10	Necrotizing enterocolitis of newborn (P77)	8	1.6	*
14	Diseases of the blood and blood-forming organs and certain disorders involving the immune mechanism (D50-D89)	5	1.0	*
14	Intrauterine hypoxia and birth asphyxia (P20-P21)	5	1.0	*
14	Hydrops fetalis not due to hemolytic disease (P83.2)	5	1.0	*
14	Assault (homicide) (*U01,X85-Y09)	5	1.0	*
...	All other causes (residual)	126	25.1	92.0

... Category not applicable

\* Figure does not meet standards of reliability or precision

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015

LCWK4. Infant, neonatal, and postneonatal deaths, percent of total deaths, and mortality rates for the 15 leading causes of infant death, by race and Hispanic origin, and sex: United States, 2015

[Rates are infant (under 1 year), neonatal (under 28 days), and postneonatal (28 days through 11 months) deaths per 100,000 live births in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of inconsistencies between reporting Hispanic origin on birth and death certificates and because deaths are based on race of decedent whereas live births are based on race of mother.]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race and sex	Number	Percent of total deaths	Mortality rate
Infants, non-Hispanic, Asian or Pacific Islander, female				
...	All causes	391	100.0	303.5
1	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	106	27.1	82.3
2	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	56	14.3	43.5
3	Newborn affected by maternal complications of pregnancy (P01)	46	11.8	35.7
4	Newborn affected by complications of placenta, cord and membranes (P02)	19	4.9	*
5	Bacterial sepsis of newborn (P36)	15	3.8	*
6	Neonatal hemorrhage (P50-P52,P54)	9	2.3	*
7	Sudden infant death syndrome (R95)	8	2.0	*
7	Accidents (unintentional injuries) (V01-X59)	8	2.0	*
9	Diseases of the circulatory system (I00-I99)	7	1.8	*
10	Respiratory distress of newborn (P22)	6	1.5	*
10	Hydrops fetalis not due to hemolytic disease (P83.2)	6	1.5	*
12	Diarrhea and gastroenteritis of infectious origin (A09)	5	1.3	*
12	Necrotizing enterocolitis of newborn (P77)	5	1.3	*
14	Newborn affected by other complications of labor and delivery (P03)	4	1.0	*
14	Intrauterine hypoxia and birth asphyxia (P20-P21)	4	1.0	*
...	All other causes (residual)	87	22.3	67.5
Neonates, all races and origins, both sexes				
...	All causes	15,652	100.0	393.4
1	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	3,981	25.4	100.1
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	3,463	22.1	87.0
3	Newborn affected by maternal complications of pregnancy (P01)	1,509	9.6	37.9
4	Newborn affected by complications of placenta, cord and membranes (P02)	898	5.7	22.6
5	Bacterial sepsis of newborn (P36)	581	3.7	14.6
6	Respiratory distress of newborn (P22)	452	2.9	11.4
7	Neonatal hemorrhage (P50-P52,P54)	400	2.6	10.1
8	Necrotizing enterocolitis of newborn (P77)	323	2.1	8.1
9	Intrauterine hypoxia and birth asphyxia (P20-P21)	307	2.0	7.7
10	Atelectasis (P28.0-P28.1)	253	1.6	6.4
11	Hydrops fetalis not due to hemolytic disease (P83.2)	167	1.1	4.2
12	Sudden infant death syndrome (R95)	153	1.0	3.8
13	Pulmonary hemorrhage originating in the perinatal period (P26)	152	1.0	3.8
14	Newborn affected by other complications of labor and delivery (P03)	124	0.8	3.1
15	Accidents (unintentional injuries) (V01-X59)	120	0.8	3.0
...	All other causes (residual)	2,769	17.7	69.6

... Category not applicable

\* Figure does not meet standards of reliability or precision

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015

LCWK4. Infant, neonatal, and postneonatal deaths, percent of total deaths, and mortality rates for the 15 leading causes of infant death, by race and Hispanic origin, and sex: United States, 2015

[Rates are infant (under 1 year), neonatal (under 28 days), and postneonatal (28 days through 11 months) deaths per 100,000 live births in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of inconsistencies between reporting Hispanic origin on birth and death certificates and because deaths are based on race of decedent whereas live births are based on race of mother.]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race and sex	Number	Percent of total deaths	Mortality rate
Neonates, all races and origins, male				
...	All causes	8,591	100.0	421.9
1	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	2,186	25.4	107.4
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	1,860	21.7	91.3
3	Newborn affected by maternal complications of pregnancy (P01)	845	9.8	41.5
4	Newborn affected by complications of placenta, cord and membranes (P02)	509	5.9	25.0
5	Bacterial sepsis of newborn (P36)	302	3.5	14.8
6	Respiratory distress of newborn (P22)	274	3.2	13.5
7	Neonatal hemorrhage (P50-P52,P54)	233	2.7	11.4
8	Necrotizing enterocolitis of newborn (P77)	175	2.0	8.6
9	Intrauterine hypoxia and birth asphyxia (P20-P21)	156	1.8	7.7
10	Atelectasis (P28.0-P28.1)	151	1.8	7.4
11	Pulmonary hemorrhage originating in the perinatal period (P26)	89	1.0	4.4
12	Hydrops fetalis not due to hemolytic disease (P83.2)	84	1.0	4.1
13	Newborn affected by other complications of labor and delivery (P03)	69	0.8	3.4
14	Sudden infant death syndrome (R95)	67	0.8	3.3
15	Accidents (unintentional injuries) (V01-X59)	61	0.7	3.0
...	All other causes (residual)	1,530	17.8	75.1
Neonates, all races and origins, female				
...	All causes	7,061	100.0	363.5
1	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	1,795	25.4	92.4
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	1,603	22.7	82.5
3	Newborn affected by maternal complications of pregnancy (P01)	664	9.4	34.2
4	Newborn affected by complications of placenta, cord and membranes (P02)	389	5.5	20.0
5	Bacterial sepsis of newborn (P36)	279	4.0	14.4
6	Respiratory distress of newborn (P22)	178	2.5	9.2
7	Neonatal hemorrhage (P50-P52,P54)	167	2.4	8.6
8	Intrauterine hypoxia and birth asphyxia (P20-P21)	151	2.1	7.8
9	Necrotizing enterocolitis of newborn (P77)	148	2.1	7.6
10	Atelectasis (P28.0-P28.1)	102	1.4	5.3
11	Sudden infant death syndrome (R95)	86	1.2	4.4
12	Hydrops fetalis not due to hemolytic disease (P83.2)	83	1.2	4.3
13	Pulmonary hemorrhage originating in the perinatal period (P26)	63	0.9	3.2
14	Accidents (unintentional injuries) (V01-X59)	59	0.8	3.0
15	Newborn affected by other complications of labor and delivery (P03)	55	0.8	2.8
15	Slow fetal growth and fetal malnutrition (P05)	55	0.8	2.8
...	All other causes (residual)	1,184	16.8	61.0

... Category not applicable

\* Figure does not meet standards of reliability or precision

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015

LCWK4. Infant, neonatal, and postneonatal deaths, percent of total deaths, and mortality rates for the 15 leading causes of infant death, by race and Hispanic origin, and sex: United States, 2015

[Rates are infant (under 1 year), neonatal (under 28 days), and postneonatal (28 days through 11 months) deaths per 100,000 live births in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of inconsistencies between reporting Hispanic origin on birth and death certificates and because deaths are based on race of decedent whereas live births are based on race of mother.]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race and sex	Number	Percent of total deaths	Mortality rate
Neonates, Hispanic, both sexes				
...	All causes	3,443	100.0	372.6
1	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	904	26.3	97.8
2	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	811	23.6	87.8
3	Newborn affected by maternal complications of pregnancy (P01)	338	9.8	36.6
4	Newborn affected by complications of placenta, cord and membranes (P02)	190	5.5	20.6
5	Bacterial sepsis of newborn (P36)	123	3.6	13.3
6	Neonatal hemorrhage (P50-P52,P54)	97	2.8	10.5
7	Respiratory distress of newborn (P22)	86	2.5	9.3
8	Necrotizing enterocolitis of newborn (P77)	74	2.1	8.0
9	Atelectasis (P28.0-P28.1)	58	1.7	6.3
10	Intrauterine hypoxia and birth asphyxia (P20-P21)	55	1.6	6.0
11	Hydrops fetalis not due to hemolytic disease (P83.2)	47	1.4	5.1
12	Pulmonary hemorrhage originating in the perinatal period (P26)	28	0.8	3.0
13	Newborn affected by other complications of labor and delivery (P03)	26	0.8	2.8
14	Hematological disorders (P60-P61)	23	0.7	2.5
15	Diseases of the circulatory system (I00-I99)	20	0.6	2.2
15	Accidents (unintentional injuries) (V01-X59)	20	0.6	2.2
...	All other causes (residual)	543	15.8	58.8
Neonates, Hispanic, male				
...	All causes	1,893	100.0	401.5
1	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	479	25.3	101.6
2	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	455	24.0	96.5
3	Newborn affected by maternal complications of pregnancy (P01)	207	10.9	43.9
4	Newborn affected by complications of placenta, cord and membranes (P02)	102	5.4	21.6
5	Bacterial sepsis of newborn (P36)	68	3.6	14.4
6	Neonatal hemorrhage (P50-P52,P54)	53	2.8	11.2
7	Respiratory distress of newborn (P22)	50	2.6	10.6
8	Necrotizing enterocolitis of newborn (P77)	41	2.2	8.7
9	Atelectasis (P28.0-P28.1)	33	1.7	7.0
10	Intrauterine hypoxia and birth asphyxia (P20-P21)	24	1.3	5.1
11	Hydrops fetalis not due to hemolytic disease (P83.2)	20	1.1	4.2
12	Pulmonary hemorrhage originating in the perinatal period (P26)	16	0.8	*
13	Newborn affected by other complications of labor and delivery (P03)	14	0.7	*
14	Accidents (unintentional injuries) (V01-X59)	11	0.6	*
15	Congenital pneumonia (P23)	10	0.5	*
...	All other causes (residual)	310	16.4	65.8

... Category not applicable

\* Figure does not meet standards of reliability or precision

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015

LCWK4. Infant, neonatal, and postneonatal deaths, percent of total deaths, and mortality rates for the 15 leading causes of infant death, by race and Hispanic origin, and sex: United States, 2015

[Rates are infant (under 1 year), neonatal (under 28 days), and postneonatal (28 days through 11 months) deaths per 100,000 live births in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of inconsistencies between reporting Hispanic origin on birth and death certificates and because deaths are based on race of decedent whereas live births are based on race of mother.]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race and sex	Number	Percent of total deaths	Mortality rate
Neonates, Hispanic, female				
...	All causes	1,550	100.0	342.5
1	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	425	27.4	93.9
2	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	356	23.0	78.7
3	Newborn affected by maternal complications of pregnancy (P01)	131	8.5	28.9
4	Newborn affected by complications of placenta, cord and membranes (P02)	88	5.7	19.4
5	Bacterial sepsis of newborn (P36)	55	3.5	12.2
6	Neonatal hemorrhage (P50-P52,P54)	44	2.8	9.7
7	Respiratory distress of newborn (P22)	36	2.3	8.0
8	Necrotizing enterocolitis of newborn (P77)	33	2.1	7.3
9	Intrauterine hypoxia and birth asphyxia (P20-P21)	31	2.0	6.8
10	Hydrops fetalis not due to hemolytic disease (P83.2)	27	1.7	6.0
11	Atelectasis (P28.0-P28.1)	25	1.6	5.5
12	Hematological disorders (P60-P61)	14	0.9	*
13	Newborn affected by other complications of labor and delivery (P03)	12	0.8	*
13	Pulmonary hemorrhage originating in the perinatal period (P26)	12	0.8	*
15	Diseases of the circulatory system (I00-I99)	11	0.7	*
...	All other causes (residual)	250	16.1	55.2
Neonates, non-Hispanic, white, both sexes				
...	All causes	6,722	100.0	315.5
1	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	1,744	25.9	81.9
2	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	1,452	21.6	68.2
3	Newborn affected by maternal complications of pregnancy (P01)	598	8.9	28.1
4	Newborn affected by complications of placenta, cord and membranes (P02)	375	5.6	17.6
5	Bacterial sepsis of newborn (P36)	223	3.3	10.5
6	Respiratory distress of newborn (P22)	181	2.7	8.5
7	Neonatal hemorrhage (P50-P52,P54)	176	2.6	8.3
8	Intrauterine hypoxia and birth asphyxia (P20-P21)	148	2.2	6.9
9	Necrotizing enterocolitis of newborn (P77)	115	1.7	5.4
10	Atelectasis (P28.0-P28.1)	108	1.6	5.1
11	Hydrops fetalis not due to hemolytic disease (P83.2)	90	1.3	4.2
12	Sudden infant death syndrome (R95)	82	1.2	3.8
13	Accidents (unintentional injuries) (V01-X59)	65	1.0	3.1
14	Pulmonary hemorrhage originating in the perinatal period (P26)	58	0.9	2.7
15	Diseases of the circulatory system (I00-I99)	52	0.8	2.4
...	All other causes (residual)	1,255	18.7	58.9

... Category not applicable

\* Figure does not meet standards of reliability or precision

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015

LCWK4. Infant, neonatal, and postneonatal deaths, percent of total deaths, and mortality rates for the 15 leading causes of infant death, by race and Hispanic origin, and sex: United States, 2015

[Rates are infant (under 1 year), neonatal (under 28 days), and postneonatal (28 days through 11 months) deaths per 100,000 live births in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of inconsistencies between reporting Hispanic origin on birth and death certificates and because deaths are based on race of decedent whereas live births are based on race of mother.]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race and sex	Number	Percent of total deaths	Mortality rate
Neonates, non-Hispanic, white, male				
...	All causes	3,688	100.0	337.5
1	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	951	25.8	87.0
2	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	762	20.7	69.7
3	Newborn affected by maternal complications of pregnancy (P01)	317	8.6	29.0
4	Newborn affected by complications of placenta, cord and membranes (P02)	214	5.8	19.6
5	Bacterial sepsis of newborn (P36)	118	3.2	10.8
6	Respiratory distress of newborn (P22)	115	3.1	10.5
7	Neonatal hemorrhage (P50-P52,P54)	107	2.9	9.8
8	Intrauterine hypoxia and birth asphyxia (P20-P21)	83	2.3	7.6
9	Atelectasis (P28.0-P28.1)	68	1.8	6.2
10	Necrotizing enterocolitis of newborn (P77)	65	1.8	5.9
11	Hydrops fetalis not due to hemolytic disease (P83.2)	50	1.4	4.6
12	Sudden infant death syndrome (R95)	40	1.1	3.7
13	Pulmonary hemorrhage originating in the perinatal period (P26)	35	0.9	3.2
13	Accidents (unintentional injuries) (V01-X59)	35	0.9	3.2
15	Diseases of the circulatory system (I00-I99)	31	0.8	2.8
...	All other causes (residual)	697	18.9	63.8
Neonates, non-Hispanic, white, female				
...	All causes	3,034	100.0	292.4
1	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	793	26.1	76.4
2	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	690	22.7	66.5
3	Newborn affected by maternal complications of pregnancy (P01)	281	9.3	27.1
4	Newborn affected by complications of placenta, cord and membranes (P02)	161	5.3	15.5
5	Bacterial sepsis of newborn (P36)	105	3.5	10.1
6	Neonatal hemorrhage (P50-P52,P54)	69	2.3	6.7
7	Respiratory distress of newborn (P22)	66	2.2	6.4
8	Intrauterine hypoxia and birth asphyxia (P20-P21)	65	2.1	6.3
9	Necrotizing enterocolitis of newborn (P77)	50	1.6	4.8
10	Sudden infant death syndrome (R95)	42	1.4	4.0
11	Atelectasis (P28.0-P28.1)	40	1.3	3.9
11	Hydrops fetalis not due to hemolytic disease (P83.2)	40	1.3	3.9
13	Slow fetal growth and fetal malnutrition (P05)	33	1.1	3.2
14	Accidents (unintentional injuries) (V01-X59)	30	1.0	2.9
15	Pulmonary hemorrhage originating in the perinatal period (P26)	23	0.8	2.2
...	All other causes (residual)	546	18.0	52.6

... Category not applicable

\* Figure does not meet standards of reliability or precision

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015

LCWK4. Infant, neonatal, and postneonatal deaths, percent of total deaths, and mortality rates for the 15 leading causes of infant death, by race and Hispanic origin, and sex: United States, 2015

[Rates are infant (under 1 year), neonatal (under 28 days), and postneonatal (28 days through 11 months) deaths per 100,000 live births in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of inconsistencies between reporting Hispanic origin on birth and death certificates and because deaths are based on race of decedent whereas live births are based on race of mother.]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race and sex	Number	Percent of total deaths	Mortality rate
Neonates, non-Hispanic, black, both sexes				
...	All causes	4,477	100.0	760.0
1	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	1,464	32.7	248.5
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	613	13.7	104.1
3	Newborn affected by maternal complications of pregnancy (P01)	455	10.2	77.2
4	Newborn affected by complications of placenta, cord and membranes (P02)	266	5.9	45.2
5	Bacterial sepsis of newborn (P36)	202	4.5	34.3
6	Respiratory distress of newborn (P22)	162	3.6	27.5
7	Necrotizing enterocolitis of newborn (P77)	113	2.5	19.2
8	Neonatal hemorrhage (P50-P52,P54)	96	2.1	16.3
9	Intrauterine hypoxia and birth asphyxia (P20-P21)	90	2.0	15.3
10	Atelectasis (P28.0-P28.1)	78	1.7	13.2
11	Pulmonary hemorrhage originating in the perinatal period (P26)	55	1.2	9.3
12	Sudden infant death syndrome (R95)	45	1.0	7.6
13	Newborn affected by maternal hypertensive disorders (P00.0)	36	0.8	6.1
13	Slow fetal growth and fetal malnutrition (P05)	36	0.8	6.1
13	Interstitial emphysema and related conditions originating in the perinatal period (P25)	36	0.8	6.1
...	All other causes (residual)	730	16.3	123.9
Neonates, non-Hispanic, black, male				
...	All causes	2,442	100.0	815.9
1	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	817	33.5	273.0
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	320	13.1	106.9
3	Newborn affected by maternal complications of pregnancy (P01)	266	10.9	88.9
4	Newborn affected by complications of placenta, cord and membranes (P02)	156	6.4	52.1
5	Bacterial sepsis of newborn (P36)	102	4.2	34.1
6	Respiratory distress of newborn (P22)	93	3.8	31.1
7	Necrotizing enterocolitis of newborn (P77)	58	2.4	19.4
8	Neonatal hemorrhage (P50-P52,P54)	54	2.2	18.0
9	Atelectasis (P28.0-P28.1)	44	1.8	14.7
10	Intrauterine hypoxia and birth asphyxia (P20-P21)	42	1.7	14.0
11	Pulmonary hemorrhage originating in the perinatal period (P26)	33	1.4	11.0
12	Interstitial emphysema and related conditions originating in the perinatal period (P25)	22	0.9	7.4
13	Slow fetal growth and fetal malnutrition (P05)	20	0.8	6.7
14	Newborn affected by maternal hypertensive disorders (P00.0)	18	0.7	*
15	Newborn affected by other complications of labor and delivery (P03)	17	0.7	*
15	Sudden infant death syndrome (R95)	17	0.7	*
...	All other causes (residual)	363	14.9	121.3

... Category not applicable

\* Figure does not meet standards of reliability or precision

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015

LCWK4. Infant, neonatal, and postneonatal deaths, percent of total deaths, and mortality rates for the 15 leading causes of infant death, by race and Hispanic origin, and sex: United States, 2015

[Rates are infant (under 1 year), neonatal (under 28 days), and postneonatal (28 days through 11 months) deaths per 100,000 live births in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of inconsistencies between reporting Hispanic origin on birth and death certificates and because deaths are based on race of decedent whereas live births are based on race of mother.]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race and sex	Number	Percent of total deaths	Mortality rate
Neonates, non-Hispanic, black, female				
...	All causes	2,035	100.0	702.4
1	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	647	31.8	223.3
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	293	14.4	101.1
3	Newborn affected by maternal complications of pregnancy (P01)	189	9.3	65.2
4	Newborn affected by complications of placenta, cord and membranes (P02)	110	5.4	38.0
5	Bacterial sepsis of newborn (P36)	100	4.9	34.5
6	Respiratory distress of newborn (P22)	69	3.4	23.8
7	Necrotizing enterocolitis of newborn (P77)	55	2.7	19.0
8	Intrauterine hypoxia and birth asphyxia (P20-P21)	48	2.4	16.6
9	Neonatal hemorrhage (P50-P52,P54)	42	2.1	14.5
10	Atelectasis (P28.0-P28.1)	34	1.7	11.7
11	Sudden infant death syndrome (R95)	28	1.4	9.7
12	Pulmonary hemorrhage originating in the perinatal period (P26)	22	1.1	7.6
13	Newborn affected by maternal hypertensive disorders (P00.0)	18	0.9	*
14	Newborn affected by other complications of labor and delivery (P03)	17	0.8	*
14	Accidents (unintentional injuries) (V01-X59)	17	0.8	*
...	All other causes (residual)	346	17.0	119.4
Neonates, non-Hispanic, American Indian or Alaska Native, both sexes				
...	All causes	159	100.0	431.8
1	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	38	23.9	103.2
2	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	23	14.5	62.5
3	Newborn affected by maternal complications of pregnancy (P01)	15	9.4	*
4	Newborn affected by complications of placenta, cord and membranes (P02)	11	6.9	*
5	Sudden infant death syndrome (R95)	9	5.7	*
6	Bacterial sepsis of newborn (P36)	7	4.4	*
7	Intrauterine hypoxia and birth asphyxia (P20-P21)	4	2.5	*
7	Accidents (unintentional injuries) (V01-X59)	4	2.5	*
9	Newborn affected by maternal hypertensive disorders (P00.0)	3	1.9	*
9	Newborn affected by other complications of labor and delivery (P03)	3	1.9	*
9	Atelectasis (P28.0-P28.1)	3	1.9	*
9	Necrotizing enterocolitis of newborn (P77)	3	1.9	*
13	Diseases of the circulatory system (I00-I99)	2	1.3	*
13	Newborn affected by other maternal conditions which may be unrelated to present pregnancy (P00.1-P00.9)	2	1.3	*
13	Respiratory distress of newborn (P22)	2	1.3	*
13	Congenital pneumonia (P23)	2	1.3	*
13	Interstitial emphysema and related conditions originating in the perinatal period (P25)	2	1.3	*
13	Pulmonary hemorrhage originating in the perinatal period (P26)	2	1.3	*
13	Neonatal hemorrhage (P50-P52,P54)	2	1.3	*
13	Hematological disorders (P60-P61)	2	1.3	*
...	All other causes (residual)	20	12.6	54.3

... Category not applicable

\* Figure does not meet standards of reliability or precision

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015

LCWK4. Infant, neonatal, and postneonatal deaths, percent of total deaths, and mortality rates for the 15 leading causes of infant death, by race and Hispanic origin, and sex: United States, 2015

[Rates are infant (under 1 year), neonatal (under 28 days), and postneonatal (28 days through 11 months) deaths per 100,000 live births in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of inconsistencies between reporting Hispanic origin on birth and death certificates and because deaths are based on race of decedent whereas live births are based on race of mother.]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race and sex	Number	Percent of total deaths	Mortality rate
Neonates, non-Hispanic, American Indian or Alaska Native, male				
...	All causes	84	100.0	445.3
1	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	20	23.8	106.0
2	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	13	15.5	*
3	Newborn affected by maternal complications of pregnancy (P01)	10	11.9	*
4	Newborn affected by complications of placenta, cord and membranes (P02)	6	7.1	*
5	Bacterial sepsis of newborn (P36)	4	4.8	*
6	Sudden infant death syndrome (R95)	3	3.6	*
7	Newborn affected by other complications of labor and delivery (P03)	2	2.4	*
7	Intrauterine hypoxia and birth asphyxia (P20-P21)	2	2.4	*
7	Respiratory distress of newborn (P22)	2	2.4	*
7	Atelectasis (P28.0-P28.1)	2	2.4	*
7	Accidents (unintentional injuries) (V01-X59)	2	2.4	*
12	Newborn affected by maternal hypertensive disorders (P00.0)	1	1.2	*
12	Newborn affected by other maternal conditions which may be unrelated to present pregnancy (P00.1-P00.9)	1	1.2	*
12	Congenital pneumonia (P23)	1	1.2	*
12	Neonatal aspiration syndromes (P24)	1	1.2	*
12	Interstitial emphysema and related conditions originating in the perinatal period (P25)	1	1.2	*
12	Neonatal hemorrhage (P50-P52,P54)	1	1.2	*
12	Hematological disorders (P60-P61)	1	1.2	*
12	Assault (homicide) (*U01,X85-Y09)	1	1.2	*
...	All other causes (residual)	10	11.9	*
Neonates, non-Hispanic, American Indian or Alaska Native, female				
...	All causes	75	100.0	417.6
1	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	18	24.0	*
2	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	10	13.3	*
3	Sudden infant death syndrome (R95)	6	8.0	*
4	Newborn affected by maternal complications of pregnancy (P01)	5	6.7	*
4	Newborn affected by complications of placenta, cord and membranes (P02)	5	6.7	*
6	Bacterial sepsis of newborn (P36)	3	4.0	*
6	Necrotizing enterocolitis of newborn (P77)	3	4.0	*
8	Diseases of the circulatory system (I00-I99)	2	2.7	*
8	Newborn affected by maternal hypertensive disorders (P00.0)	2	2.7	*
8	Intrauterine hypoxia and birth asphyxia (P20-P21)	2	2.7	*
8	Pulmonary hemorrhage originating in the perinatal period (P26)	2	2.7	*
8	Accidents (unintentional injuries) (V01-X59)	2	2.7	*
13	Diseases of the blood and blood-forming organs and certain disorders involving the immune mechanism (D50-D89)	1	1.3	*
13	Short stature, not elsewhere classified (E34.3)	1	1.3	*
13	Newborn affected by other maternal conditions which may be unrelated to present pregnancy (P00.1-P00.9)	1	1.3	*
13	Newborn affected by other complications of labor and delivery (P03)	1	1.3	*
13	Congenital pneumonia (P23)	1	1.3	*
13	Interstitial emphysema and related conditions originating in the perinatal period (P25)	1	1.3	*
13	Atelectasis (P28.0-P28.1)	1	1.3	*
13	Neonatal hemorrhage (P50-P52,P54)	1	1.3	*
13	Hematological disorders (P60-P61)	1	1.3	*
...	All other causes (residual)	6	8.0	*

... Category not applicable

\* Figure does not meet standards of reliability or precision

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015

LCWK4. Infant, neonatal, and postneonatal deaths, percent of total deaths, and mortality rates for the 15 leading causes of infant death, by race and Hispanic origin, and sex: United States, 2015

[Rates are infant (under 1 year), neonatal (under 28 days), and postneonatal (28 days through 11 months) deaths per 100,000 live births in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of inconsistencies between reporting Hispanic origin on birth and death certificates and because deaths are based on race of decedent whereas live births are based on race of mother.]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race and sex	Number	Percent of total deaths	Mortality rate
Neonates, non-Hispanic, Asian or Pacific Islander, both sexes				
...	All causes	635	100.0	238.9
1	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	139	21.9	52.3
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	136	21.4	51.2
3	Newborn affected by maternal complications of pregnancy (P01)	82	12.9	30.8
4	Newborn affected by complications of placenta, cord and membranes (P02)	39	6.1	14.7
5	Neonatal hemorrhage (P50-P52,P54)	26	4.1	9.8
6	Bacterial sepsis of newborn (P36)	22	3.5	8.3
7	Respiratory distress of newborn (P22)	19	3.0	*
8	Necrotizing enterocolitis of newborn (P77)	13	2.0	*
9	Newborn affected by other complications of labor and delivery (P03)	12	1.9	*
10	Hydrops fetalis not due to hemolytic disease (P83.2)	10	1.6	*
11	Intrauterine hypoxia and birth asphyxia (P20-P21)	8	1.3	*
12	Pulmonary hemorrhage originating in the perinatal period (P26)	7	1.1	*
12	Hematological disorders (P60-P61)	7	1.1	*
14	Atelectasis (P28.0-P28.1)	5	0.8	*
15	Diseases of the blood and blood-forming organs and certain disorders involving the immune mechanism (D50-D89)	3	0.5	*
15	Diseases of the circulatory system (I00-I99)	3	0.5	*
15	Newborn affected by other maternal conditions which may be unrelated to present pregnancy (P00.1-P00.9)	3	0.5	*
15	Slow fetal growth and fetal malnutrition (P05)	3	0.5	*
...	All other causes (residual)	98	15.4	36.9
Neonates, non-Hispanic, Asian or Pacific Islander, male				
...	All causes	359	100.0	262.0
1	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	84	23.4	61.3
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	73	20.3	53.3
3	Newborn affected by maternal complications of pregnancy (P01)	36	10.0	26.3
4	Newborn affected by complications of placenta, cord and membranes (P02)	20	5.6	14.6
5	Neonatal hemorrhage (P50-P52,P54)	17	4.7	*
6	Respiratory distress of newborn (P22)	13	3.6	*
7	Newborn affected by other complications of labor and delivery (P03)	8	2.2	*
7	Bacterial sepsis of newborn (P36)	8	2.2	*
7	Necrotizing enterocolitis of newborn (P77)	8	2.2	*
10	Intrauterine hypoxia and birth asphyxia (P20-P21)	4	1.1	*
10	Pulmonary hemorrhage originating in the perinatal period (P26)	4	1.1	*
10	Hematological disorders (P60-P61)	4	1.1	*
10	Hydrops fetalis not due to hemolytic disease (P83.2)	4	1.1	*
14	Slow fetal growth and fetal malnutrition (P05)	3	0.8	*
14	Atelectasis (P28.0-P28.1)	3	0.8	*
...	All other causes (residual)	70	19.5	51.1

... Category not applicable

\* Figure does not meet standards of reliability or precision

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015

LCWK4. Infant, neonatal, and postneonatal deaths, percent of total deaths, and mortality rates for the 15 leading causes of infant death, by race and Hispanic origin, and sex: United States, 2015

[Rates are infant (under 1 year), neonatal (under 28 days), and postneonatal (28 days through 11 months) deaths per 100,000 live births in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of inconsistencies between reporting Hispanic origin on birth and death certificates and because deaths are based on race of decedent whereas live births are based on race of mother.]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race and sex	Number	Percent of total deaths	Mortality rate
Neonates, non-Hispanic, Asian or Pacific Islander, female				
...	All causes	276	100.0	214.2
1	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	63	22.8	48.9
2	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	55	19.9	42.7
3	Newborn affected by maternal complications of pregnancy (P01)	46	16.7	35.7
4	Newborn affected by complications of placenta, cord and membranes (P02)	19	6.9	*
5	Bacterial sepsis of newborn (P36)	14	5.1	*
6	Neonatal hemorrhage (P50-P52,P54)	9	3.3	*
7	Respiratory distress of newborn (P22)	6	2.2	*
7	Hydrops fetalis not due to hemolytic disease (P83.2)	6	2.2	*
9	Necrotizing enterocolitis of newborn (P77)	5	1.8	*
10	Newborn affected by other complications of labor and delivery (P03)	4	1.4	*
10	Intrauterine hypoxia and birth asphyxia (P20-P21)	4	1.4	*
12	Pulmonary hemorrhage originating in the perinatal period (P26)	3	1.1	*
12	Hematological disorders (P60-P61)	3	1.1	*
14	Congenital pneumonia (P23)	2	0.7	*
14	Atelectasis (P28.0-P28.1)	2	0.7	*
...	All other causes (residual)	35	12.7	27.2
Postneonates, all races and origins, both sexes				
...	All causes	7,803	100.0	196.1
1	Sudden infant death syndrome (R95)	1,415	18.1	35.6
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	1,362	17.5	34.2
3	Accidents (unintentional injuries) (V01-X59)	1,171	15.0	29.4
4	Diseases of the circulatory system (I00-I99)	331	4.2	8.3
5	Assault (homicide) (*U01,X85-Y09)	244	3.1	6.1
6	Diarrhea and gastroenteritis of infectious origin (A09)	210	2.7	5.3
7	Septicemia (A40-A41)	177	2.3	4.4
8	Influenza and pneumonia (J09-J18)	173	2.2	4.3
9	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	103	1.3	2.6
10	Chronic respiratory disease originating in the perinatal period (P27)	102	1.3	2.6
11	Anoxic brain damage, not elsewhere classified (G93.1)	49	0.6	1.2
11	Acute bronchitis and acute bronchiolitis (J20-J21)	49	0.6	1.2
13	Diseases of the blood and blood-forming organs and certain disorders involving the immune mechanism (D50-D89)	47	0.6	1.2
14	Malignant neoplasms (C00-C97)	43	0.6	1.1
15	Necrotizing enterocolitis of newborn (P77)	40	0.5	1.0
...	All other causes (residual)	2,287	29.3	57.5

... Category not applicable

\* Figure does not meet standards of reliability or precision

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015

LCWK4. Infant, neonatal, and postneonatal deaths, percent of total deaths, and mortality rates for the 15 leading causes of infant death, by race and Hispanic origin, and sex: United States, 2015

[Rates are infant (under 1 year), neonatal (under 28 days), and postneonatal (28 days through 11 months) deaths per 100,000 live births in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of inconsistencies between reporting Hispanic origin on birth and death certificates and because deaths are based on race of decedent whereas live births are based on race of mother.]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race and sex	Number	Percent of total deaths	Mortality rate
Postneonates, all races and origins, male				
...	All causes	4,417	100.0	216.9
1	Sudden infant death syndrome (R95)	856	19.4	42.0
2	Accidents (unintentional injuries) (V01-X59)	694	15.7	34.1
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	691	15.6	33.9
4	Diseases of the circulatory system (I00-I99)	173	3.9	8.5
5	Assault (homicide) (*U01,X85-Y09)	137	3.1	6.7
6	Diarrhea and gastroenteritis of infectious origin (A09)	110	2.5	5.4
6	Influenza and pneumonia (J09-J18)	110	2.5	5.4
8	Septicemia (A40-A41)	94	2.1	4.6
9	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	63	1.4	3.1
10	Chronic respiratory disease originating in the perinatal period (P27)	51	1.2	2.5
11	Diseases of the blood and blood-forming organs and certain disorders involving the immune mechanism (D50-D89)	32	0.7	1.6
12	Acute bronchitis and acute bronchiolitis (J20-J21)	29	0.7	1.4
13	Malignant neoplasms (C00-C97)	24	0.5	1.2
14	Anoxic brain damage, not elsewhere classified (G93.1)	19	0.4	*
14	Necrotizing enterocolitis of newborn (P77)	19	0.4	*
...	All other causes (residual)	1,315	29.8	64.6
Postneonates, all races and origins, female				
...	All causes	3,386	100.0	174.3
1	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	671	19.8	34.5
2	Sudden infant death syndrome (R95)	559	16.5	28.8
3	Accidents (unintentional injuries) (V01-X59)	477	14.1	24.6
4	Diseases of the circulatory system (I00-I99)	158	4.7	8.1
5	Assault (homicide) (*U01,X85-Y09)	107	3.2	5.5
6	Diarrhea and gastroenteritis of infectious origin (A09)	100	3.0	5.1
7	Septicemia (A40-A41)	83	2.5	4.3
8	Influenza and pneumonia (J09-J18)	63	1.9	3.2
9	Chronic respiratory disease originating in the perinatal period (P27)	51	1.5	2.6
10	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	40	1.2	2.1
11	Anoxic brain damage, not elsewhere classified (G93.1)	30	0.9	1.5
12	Necrotizing enterocolitis of newborn (P77)	21	0.6	1.1
13	Meningitis (G00,G03)	20	0.6	1.0
13	Acute bronchitis and acute bronchiolitis (J20-J21)	20	0.6	1.0
15	Malignant neoplasms (C00-C97)	19	0.6	*
...	All other causes (residual)	967	28.6	49.8

... Category not applicable

\* Figure does not meet standards of reliability or precision

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015

LCWK4. Infant, neonatal, and postneonatal deaths, percent of total deaths, and mortality rates for the 15 leading causes of infant death, by race and Hispanic origin, and sex: United States, 2015

[Rates are infant (under 1 year), neonatal (under 28 days), and postneonatal (28 days through 11 months) deaths per 100,000 live births in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of inconsistencies between reporting Hispanic origin on birth and death certificates and because deaths are based on race of decedent whereas live births are based on race of mother.]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race and sex	Number	Percent of total deaths	Mortality rate
Postneonates, Hispanic, both sexes				
...	All causes	1,362	100.0	147.4
1	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	312	22.9	33.8
2	Sudden infant death syndrome (R95)	187	13.7	20.2
3	Accidents (unintentional injuries) (V01-X59)	167	12.3	18.1
4	Diseases of the circulatory system (I00-I99)	54	4.0	5.8
5	Diarrhea and gastroenteritis of infectious origin (A09)	42	3.1	4.5
6	Septicemia (A40-A41)	41	3.0	4.4
7	Assault (homicide) (*U01,X85-Y09)	37	2.7	4.0
8	Influenza and pneumonia (J09-J18)	30	2.2	3.2
9	Chronic respiratory disease originating in the perinatal period (P27)	16	1.2	*
10	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	13	1.0	*
11	Malignant neoplasms (C00-C97)	12	0.9	*
12	Diseases of the blood and blood-forming organs and certain disorders involving the immune mechanism (D50-D89)	9	0.7	*
13	Meningitis (G00,G03)	8	0.6	*
13	Necrotizing enterocolitis of newborn (P77)	8	0.6	*
15	Anoxic brain damage, not elsewhere classified (G93.1)	6	0.4	*
15	Acute bronchitis and acute bronchiolitis (J20-J21)	6	0.4	*
...	All other causes (residual)	414	30.4	44.8
Postneonates, Hispanic, male				
...	All causes	726	100.0	154.0
1	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	156	21.5	33.1
2	Sudden infant death syndrome (R95)	110	15.2	23.3
3	Accidents (unintentional injuries) (V01-X59)	93	12.8	19.7
4	Septicemia (A40-A41)	20	2.8	4.2
4	Diseases of the circulatory system (I00-I99)	20	2.8	4.2
4	Influenza and pneumonia (J09-J18)	20	2.8	4.2
7	Diarrhea and gastroenteritis of infectious origin (A09)	17	2.3	*
7	Assault (homicide) (*U01,X85-Y09)	17	2.3	*
9	Chronic respiratory disease originating in the perinatal period (P27)	10	1.4	*
10	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	9	1.2	*
11	Diseases of the blood and blood-forming organs and certain disorders involving the immune mechanism (D50-D89)	8	1.1	*
12	Malignant neoplasms (C00-C97)	7	1.0	*
13	Acute bronchitis and acute bronchiolitis (J20-J21)	5	0.7	*
14	Volume depletion, disorders of fluid, electrolyte and acid-base balance (E86-E87)	4	0.6	*
15	Meningitis (G00,G03)	3	0.4	*
15	Bronchitis, chronic and unspecified (J40-J42)	3	0.4	*
15	Hernia of abdominal cavity and intestinal obstruction without hernia (K40-K46,K56)	3	0.4	*
15	Renal failure and other disorders of kidney (N17-N19,N25,N27)	3	0.4	*
...	All other causes (residual)	218	30.0	46.2

... Category not applicable

\* Figure does not meet standards of reliability or precision

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015

LCWK4. Infant, neonatal, and postneonatal deaths, percent of total deaths, and mortality rates for the 15 leading causes of infant death, by race and Hispanic origin, and sex: United States, 2015

[Rates are infant (under 1 year), neonatal (under 28 days), and postneonatal (28 days through 11 months) deaths per 100,000 live births in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of inconsistencies between reporting Hispanic origin on birth and death certificates and because deaths are based on race of decedent whereas live births are based on race of mother.]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race and sex	Number	Percent of total deaths	Mortality rate
Postneonates, Hispanic, female				
...	All causes	636	100.0	140.5
1	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	156	24.5	34.5
2	Sudden infant death syndrome (R95)	77	12.1	17.0
3	Accidents (unintentional injuries) (V01-X59)	74	11.6	16.4
4	Diseases of the circulatory system (I00-I99)	34	5.3	7.5
5	Diarrhea and gastroenteritis of infectious origin (A09)	25	3.9	5.5
6	Septicemia (A40-A41)	21	3.3	4.6
7	Assault (homicide) (*U01,X85-Y09)	20	3.1	4.4
8	Influenza and pneumonia (J09-J18)	10	1.6	*
9	Chronic respiratory disease originating in the perinatal period (P27)	6	0.9	*
9	Necrotizing enterocolitis of newborn (P77)	6	0.9	*
11	Malignant neoplasms (C00-C97)	5	0.8	*
11	Meningitis (G00,G03)	5	0.8	*
11	Anoxic brain damage, not elsewhere classified (G93.1)	5	0.8	*
14	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	4	0.6	*
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2	0.3	*
15	Gastritis, duodenitis, and noninfective enteritis and colitis (K29,K50-K55)	2	0.3	*
15	Newborn affected by complications of placenta, cord and membranes (P02)	2	0.3	*
15	Slow fetal growth and fetal malnutrition (P05)	2	0.3	*
15	Atelectasis (P28.0-P28.1)	2	0.3	*
15	Bacterial sepsis of newborn (P36)	2	0.3	*
...	All other causes (residual)	176	27.7	38.9
Postneonates, non-Hispanic, white, both sexes				
...	All causes	3,555	100.0	166.9
1	Sudden infant death syndrome (R95)	689	19.4	32.3
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	629	17.7	29.5
3	Accidents (unintentional injuries) (V01-X59)	567	15.9	26.6
4	Diseases of the circulatory system (I00-I99)	144	4.1	6.8
5	Assault (homicide) (*U01,X85-Y09)	111	3.1	5.2
6	Influenza and pneumonia (J09-J18)	76	2.1	3.6
7	Diarrhea and gastroenteritis of infectious origin (A09)	73	2.1	3.4
8	Septicemia (A40-A41)	68	1.9	3.2
9	Chronic respiratory disease originating in the perinatal period (P27)	41	1.2	1.9
10	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	33	0.9	1.5
11	Anoxic brain damage, not elsewhere classified (G93.1)	31	0.9	1.5
12	Diseases of the blood and blood-forming organs and certain disorders involving the immune mechanism (D50-D89)	21	0.6	1.0
12	Acute bronchitis and acute bronchiolitis (J20-J21)	21	0.6	1.0
14	Malignant neoplasms (C00-C97)	20	0.6	0.9
15	Necrotizing enterocolitis of newborn (P77)	18	0.5	*
...	All other causes (residual)	1,013	28.5	47.6

... Category not applicable

\* Figure does not meet standards of reliability or precision

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015

LCWK4. Infant, neonatal, and postneonatal deaths, percent of total deaths, and mortality rates for the 15 leading causes of infant death, by race and Hispanic origin, and sex: United States, 2015

[Rates are infant (under 1 year), neonatal (under 28 days), and postneonatal (28 days through 11 months) deaths per 100,000 live births in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of inconsistencies between reporting Hispanic origin on birth and death certificates and because deaths are based on race of decedent whereas live births are based on race of mother.]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race and sex	Number	Percent of total deaths	Mortality rate
Postneonates, non-Hispanic, white, male				
...	All causes	2,070	100.0	189.4
1	Sudden infant death syndrome (R95)	428	20.7	39.2
2	Accidents (unintentional injuries) (V01-X59)	360	17.4	32.9
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	312	15.1	28.6
4	Diseases of the circulatory system (I00-I99)	86	4.2	7.9
5	Assault (homicide) (*U01,X85-Y09)	71	3.4	6.5
6	Influenza and pneumonia (J09-J18)	46	2.2	4.2
7	Diarrhea and gastroenteritis of infectious origin (A09)	37	1.8	3.4
8	Septicemia (A40-A41)	36	1.7	3.3
9	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	22	1.1	2.0
10	Chronic respiratory disease originating in the perinatal period (P27)	20	1.0	1.8
11	Diseases of the blood and blood-forming organs and certain disorders involving the immune mechanism (D50-D89)	14	0.7	*
12	Malignant neoplasms (C00-C97)	12	0.6	*
13	Anoxic brain damage, not elsewhere classified (G93.1)	11	0.5	*
13	Acute bronchitis and acute bronchiolitis (J20-J21)	11	0.5	*
15	Necrotizing enterocolitis of newborn (P77)	9	0.4	*
...	All other causes (residual)	595	28.7	54.4
Postneonates, non-Hispanic, white, female				
...	All causes	1,485	100.0	143.1
1	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	317	21.3	30.6
2	Sudden infant death syndrome (R95)	261	17.6	25.2
3	Accidents (unintentional injuries) (V01-X59)	207	13.9	20.0
4	Diseases of the circulatory system (I00-I99)	58	3.9	5.6
5	Assault (homicide) (*U01,X85-Y09)	40	2.7	3.9
6	Diarrhea and gastroenteritis of infectious origin (A09)	36	2.4	3.5
7	Septicemia (A40-A41)	32	2.2	3.1
8	Influenza and pneumonia (J09-J18)	30	2.0	2.9
9	Chronic respiratory disease originating in the perinatal period (P27)	21	1.4	2.0
10	Anoxic brain damage, not elsewhere classified (G93.1)	20	1.3	1.9
11	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	11	0.7	*
12	Acute bronchitis and acute bronchiolitis (J20-J21)	10	0.7	*
13	Volume depletion, disorders of fluid, electrolyte and acid-base balance (E86-E87)	9	0.6	*
13	Necrotizing enterocolitis of newborn (P77)	9	0.6	*
15	Malignant neoplasms (C00-C97)	8	0.5	*
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	8	0.5	*
...	All other causes (residual)	408	27.5	39.3

... Category not applicable

\* Figure does not meet standards of reliability or precision

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015

LCWK4. Infant, neonatal, and postneonatal deaths, percent of total deaths, and mortality rates for the 15 leading causes of infant death, by race and Hispanic origin, and sex: United States, 2015

[Rates are infant (under 1 year), neonatal (under 28 days), and postneonatal (28 days through 11 months) deaths per 100,000 live births in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of inconsistencies between reporting Hispanic origin on birth and death certificates and because deaths are based on race of decedent whereas live births are based on race of mother.]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race and sex	Number	Percent of total deaths	Mortality rate
Postneonates, non-Hispanic, black, both sexes				
...	All causes	2,430	100.0	412.5
1	Sudden infant death syndrome (R95)	468	19.3	79.5
2	Accidents (unintentional injuries) (V01-X59)	381	15.7	64.7
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	317	13.0	53.8
4	Diseases of the circulatory system (I00-I99)	109	4.5	18.5
5	Assault (homicide) (*U01,X85-Y09)	87	3.6	14.8
6	Diarrhea and gastroenteritis of infectious origin (A09)	80	3.3	13.6
7	Septicemia (A40-A41)	58	2.4	9.8
8	Influenza and pneumonia (J09-J18)	56	2.3	9.5
9	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	49	2.0	8.3
10	Chronic respiratory disease originating in the perinatal period (P27)	43	1.8	7.3
11	Acute bronchitis and acute bronchiolitis (J20-J21)	17	0.7	*
12	Necrotizing enterocolitis of newborn (P77)	13	0.5	*
13	Meningitis (G00,G03)	12	0.5	*
14	Diseases of the blood and blood-forming organs and certain disorders involving the immune mechanism (D50-D89)	11	0.5	*
15	Anoxic brain damage, not elsewhere classified (G93.1)	10	0.4	*
15	Renal failure and other disorders of kidney (N17-N19,N25,N27)	10	0.4	*
...	All other causes (residual)	709	29.2	120.4
Postneonates, non-Hispanic, black, male				
...	All causes	1,373	100.0	458.7
1	Sudden infant death syndrome (R95)	272	19.8	90.9
2	Accidents (unintentional injuries) (V01-X59)	210	15.3	70.2
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	173	12.6	57.8
4	Diseases of the circulatory system (I00-I99)	52	3.8	17.4
5	Diarrhea and gastroenteritis of infectious origin (A09)	46	3.4	15.4
6	Assault (homicide) (*U01,X85-Y09)	44	3.2	14.7
7	Influenza and pneumonia (J09-J18)	36	2.6	12.0
8	Septicemia (A40-A41)	33	2.4	11.0
9	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	27	2.0	9.0
10	Chronic respiratory disease originating in the perinatal period (P27)	20	1.5	6.7
11	Acute bronchitis and acute bronchiolitis (J20-J21)	10	0.7	*
12	Hernia of abdominal cavity and intestinal obstruction without hernia (K40-K46,K56)	7	0.5	*
12	Necrotizing enterocolitis of newborn (P77)	7	0.5	*
14	Diseases of the blood and blood-forming organs and certain disorders involving the immune mechanism (D50-D89)	6	0.4	*
14	Meningitis (G00,G03)	6	0.4	*
14	Anoxic brain damage, not elsewhere classified (G93.1)	6	0.4	*
14	Respiratory distress of newborn (P22)	6	0.4	*
...	All other causes (residual)	412	30.0	137.6

... Category not applicable

\* Figure does not meet standards of reliability or precision

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015

LCWK4. Infant, neonatal, and postneonatal deaths, percent of total deaths, and mortality rates for the 15 leading causes of infant death, by race and Hispanic origin, and sex: United States, 2015

[Rates are infant (under 1 year), neonatal (under 28 days), and postneonatal (28 days through 11 months) deaths per 100,000 live births in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of inconsistencies between reporting Hispanic origin on birth and death certificates and because deaths are based on race of decedent whereas live births are based on race of mother.]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race and sex	Number	Percent of total deaths	Mortality rate
Postneonates, non-Hispanic, black, female				
...	All causes	1,057	100.0	364.8
1	Sudden infant death syndrome (R95)	196	18.5	67.6
2	Accidents (unintentional injuries) (V01-X59)	171	16.2	59.0
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	144	13.6	49.7
4	Diseases of the circulatory system (I00-I99)	57	5.4	19.7
5	Assault (homicide) (*U01,X85-Y09)	43	4.1	14.8
6	Diarrhea and gastroenteritis of infectious origin (A09)	34	3.2	11.7
7	Septicemia (A40-A41)	25	2.4	8.6
8	Chronic respiratory disease originating in the perinatal period (P27)	23	2.2	7.9
9	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	22	2.1	7.6
10	Influenza and pneumonia (J09-J18)	20	1.9	6.9
11	Acute bronchitis and acute bronchiolitis (J20-J21)	7	0.7	*
11	Renal failure and other disorders of kidney (N17-N19,N25,N27)	7	0.7	*
13	Meningitis (G00,G03)	6	0.6	*
13	Necrotizing enterocolitis of newborn (P77)	6	0.6	*
15	Diseases of the blood and blood-forming organs and certain disorders involving the immune mechanism (D50-D89)	5	0.5	*
...	All other causes (residual)	291	27.5	100.4
Postneonates, non-Hispanic, American Indian or Alaska Native, both sexes				
...	All causes	141	100.0	382.9
1	Sudden infant death syndrome (R95)	40	28.4	108.6
2	Accidents (unintentional injuries) (V01-X59)	28	19.9	76.0
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	20	14.2	54.3
4	Influenza and pneumonia (J09-J18)	4	2.8	*
5	Diseases of the circulatory system (I00-I99)	3	2.1	*
5	Acute bronchitis and acute bronchiolitis (J20-J21)	3	2.1	*
7	Septicemia (A40-A41)	2	1.4	*
7	Meningitis (G00,G03)	2	1.4	*
7	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	2	1.4	*
7	Assault (homicide) (*U01,X85-Y09)	2	1.4	*
11	Diarrhea and gastroenteritis of infectious origin (A09)	1	0.7	*
11	Malignant neoplasms (C00-C97)	1	0.7	*
11	Diseases of the blood and blood-forming organs and certain disorders involving the immune mechanism (D50-D89)	1	0.7	*
11	Cystic fibrosis (E84)	1	0.7	*
11	Anoxic brain damage, not elsewhere classified (G93.1)	1	0.7	*
11	Hernia of abdominal cavity and intestinal obstruction without hernia (K40-K46,K56)	1	0.7	*
11	Renal failure and other disorders of kidney (N17-N19,N25,N27)	1	0.7	*
11	Slow fetal growth and fetal malnutrition (P05)	1	0.7	*
11	Hydrops fetalis not due to hemolytic disease (P83.2)	1	0.7	*
...	All other causes (residual)	26	18.4	70.6

... Category not applicable

\* Figure does not meet standards of reliability or precision

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015

LCWK4. Infant, neonatal, and postneonatal deaths, percent of total deaths, and mortality rates for the 15 leading causes of infant death, by race and Hispanic origin, and sex: United States, 2015

[Rates are infant (under 1 year), neonatal (under 28 days), and postneonatal (28 days through 11 months) deaths per 100,000 live births in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of inconsistencies between reporting Hispanic origin on birth and death certificates and because deaths are based on race of decedent whereas live births are based on race of mother.]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race and sex	Number	Percent of total deaths	Mortality rate
Postneonates, non-Hispanic, American Indian or Alaska Native, male				
...	All causes	77	100.0	408.2
1	Sudden infant death syndrome (R95)	25	32.5	132.5
2	Accidents (unintentional injuries) (V01-X59)	15	19.5	*
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	11	14.3	*
4	Diseases of the circulatory system (I00-I99)	2	2.6	*
4	Influenza and pneumonia (J09-J18)	2	2.6	*
4	Acute bronchitis and acute bronchiolitis (J20-J21)	2	2.6	*
7	Diarrhea and gastroenteritis of infectious origin (A09)	1	1.3	*
7	Septicemia (A40-A41)	1	1.3	*
7	Malignant neoplasms (C00-C97)	1	1.3	*
7	Diseases of the blood and blood-forming organs and certain disorders involving the immune mechanism (D50-D89)	1	1.3	*
7	Cystic fibrosis (E84)	1	1.3	*
7	Anoxic brain damage, not elsewhere classified (G93.1)	1	1.3	*
7	Hernia of abdominal cavity and intestinal obstruction without hernia (K40-K46,K56)	1	1.3	*
7	Renal failure and other disorders of kidney (N17-N19,N25,N27)	1	1.3	*
7	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	1	1.3	*
7	Assault (homicide) (*U01,X85-Y09)	1	1.3	*
...	All other causes (residual)	10	13.0	*
Postneonates, non-Hispanic, American Indian or Alaska Native, female				
...	All causes	64	100.0	356.3
1	Sudden infant death syndrome (R95)	15	23.4	*
2	Accidents (unintentional injuries) (V01-X59)	13	20.3	*
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	9	14.1	*
4	Meningitis (G00,G03)	2	3.1	*
4	Influenza and pneumonia (J09-J18)	2	3.1	*
6	Septicemia (A40-A41)	1	1.6	*
6	Diseases of the circulatory system (I00-I99)	1	1.6	*
6	Acute bronchitis and acute bronchiolitis (J20-J21)	1	1.6	*
6	Slow fetal growth and fetal malnutrition (P05)	1	1.6	*
6	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	1	1.6	*
6	Hydrops fetalis not due to hemolytic disease (P83.2)	1	1.6	*
6	Assault (homicide) (*U01,X85-Y09)	1	1.6	*
...	All other causes (residual)	16	25.0	*

... Category not applicable

\* Figure does not meet standards of reliability or precision

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015

LCWK4. Infant, neonatal, and postneonatal deaths, percent of total deaths, and mortality rates for the 15 leading causes of infant death, by race and Hispanic origin, and sex: United States, 2015

[Rates are infant (under 1 year), neonatal (under 28 days), and postneonatal (28 days through 11 months) deaths per 100,000 live births in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of inconsistencies between reporting Hispanic origin on birth and death certificates and because deaths are based on race of decedent whereas live births are based on race of mother.]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race and sex	Number	Percent of total deaths	Mortality rate
Postneonates, non-Hispanic, Asian or Pacific Islander, both sexes				
...	All causes	257	100.0	96.7
1	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	75	29.2	28.2
2	Sudden infant death syndrome (R95)	24	9.3	9.0
3	Accidents (unintentional injuries) (V01-X59)	22	8.6	8.3
4	Diseases of the circulatory system (I00-I99)	17	6.6	*
5	Diarrhea and gastroenteritis of infectious origin (A09)	13	5.1	*
6	Septicemia (A40-A41)	7	2.7	*
7	Assault (homicide) (*U01,X85-Y09)	6	2.3	*
8	Diseases of the blood and blood-forming organs and certain disorders involving the immune mechanism (D50-D89)	5	1.9	*
8	Influenza and pneumonia (J09-J18)	5	1.9	*
10	Malignant neoplasms (C00-C97)	4	1.6	*
11	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	3	1.2	*
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2	0.8	*
12	Hernia of abdominal cavity and intestinal obstruction without hernia (K40-K46,K56)	2	0.8	*
12	Renal failure and other disorders of kidney (N17-N19,N25,N27)	2	0.8	*
12	Slow fetal growth and fetal malnutrition (P05)	2	0.8	*
...	All other causes (residual)	68	26.5	25.6
Postneonates, non-Hispanic, Asian or Pacific Islander, male				
...	All causes	142	100.0	103.6
1	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	32	22.5	23.4
2	Sudden infant death syndrome (R95)	16	11.3	*
3	Accidents (unintentional injuries) (V01-X59)	14	9.9	*
4	Diseases of the circulatory system (I00-I99)	11	7.7	*
5	Diarrhea and gastroenteritis of infectious origin (A09)	8	5.6	*
6	Septicemia (A40-A41)	4	2.8	*
6	Influenza and pneumonia (J09-J18)	4	2.8	*
6	Assault (homicide) (*U01,X85-Y09)	4	2.8	*
9	Diseases of the blood and blood-forming organs and certain disorders involving the immune mechanism (D50-D89)	3	2.1	*
10	Renal failure and other disorders of kidney (N17-N19,N25,N27)	2	1.4	*
10	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	2	1.4	*
12	Malignant neoplasms (C00-C97)	1	0.7	*
12	Acute bronchitis and acute bronchiolitis (J20-J21)	1	0.7	*
12	Hernia of abdominal cavity and intestinal obstruction without hernia (K40-K46,K56)	1	0.7	*
12	Newborn affected by maternal complications of pregnancy (P01)	1	0.7	*
12	Intrauterine hypoxia and birth asphyxia (P20-P21)	1	0.7	*
12	Chronic respiratory disease originating in the perinatal period (P27)	1	0.7	*
12	Hydrops fetalis not due to hemolytic disease (P83.2)	1	0.7	*
...	All other causes (residual)	35	24.6	25.5

... Category not applicable

\* Figure does not meet standards of reliability or precision

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015

LCWK4. Infant, neonatal, and postneonatal deaths, percent of total deaths, and mortality rates for the 15 leading causes of infant death, by race and Hispanic origin, and sex: United States, 2015

[Rates are infant (under 1 year), neonatal (under 28 days), and postneonatal (28 days through 11 months) deaths per 100,000 live births in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of inconsistencies between reporting Hispanic origin on birth and death certificates and because deaths are based on race of decedent whereas live births are based on race of mother.]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race and sex	Number	Percent of total deaths	Mortality rate
Postneonates, non-Hispanic, Asian or Pacific Islander, female				
...	All causes	115	100.0	89.3
1	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	43	37.4	33.4
2	Sudden infant death syndrome (R95)	8	7.0	*
2	Accidents (unintentional injuries) (V01-X59)	8	7.0	*
4	Diseases of the circulatory system (I00-I99)	6	5.2	*
5	Diarrhea and gastroenteritis of infectious origin (A09)	5	4.3	*
6	Septicemia (A40-A41)	3	2.6	*
6	Malignant neoplasms (C00-C97)	3	2.6	*
8	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2	1.7	*
8	Diseases of the blood and blood-forming organs and certain disorders involving the immune mechanism (D50-D89)	2	1.7	*
8	Slow fetal growth and fetal malnutrition (P05)	2	1.7	*
8	Assault (homicide) (*U01,X85-Y09)	2	1.7	*
12	Anoxic brain damage, not elsewhere classified (G93.1)	1	0.9	*
12	Diseases of the ear and mastoid process (H60-H93)	1	0.9	*
12	Influenza and pneumonia (J09-J18)	1	0.9	*
12	Pneumonitis due to solids and liquids (J69)	1	0.9	*
12	Hernia of abdominal cavity and intestinal obstruction without hernia (K40-K46,K56)	1	0.9	*
12	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	1	0.9	*
12	Interstitial emphysema and related conditions originating in the perinatal period (P25)	1	0.9	*
12	Bacterial sepsis of newborn (P36)	1	0.9	*
...	All other causes (residual)	23	20.0	17.9

... Category not applicable

\* Figure does not meet standards of reliability or precision

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015