

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2014

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
All races, both sexes, all ages				
...	All causes	2,626,418	100.0	823.7
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	614,348	23.4	192.7
2	Malignant neoplasms (C00-C97)	591,699	22.5	185.6
3	Chronic lower respiratory diseases (J40-J47)	147,101	5.6	46.1
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	136,053	5.2	42.7
5	Cerebrovascular diseases (I60-I69)	133,103	5.1	41.7
6	Alzheimer's disease (G30)	93,541	3.6	29.3
7	Diabetes mellitus (E10-E14)	76,488	2.9	24.0
8	Influenza and pneumonia (J09-J18)	55,227	2.1	17.3
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	48,146	1.8	15.1
10	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	42,773	1.6	13.4
11	Septicemia (A40-A41)	38,940	1.5	12.2
12	Chronic liver disease and cirrhosis (K70,K73-K74)	38,170	1.5	12.0
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	30,221	1.2	9.5
14	Parkinson's disease (G20-G21)	26,150	1.0	8.2
15	Pneumonitis due to solids and liquids (J69)	18,792	0.7	5.9
...	All other causes (residual)	535,666	20.4	168.0
All races, both sexes, 1-4 years				
...	All causes	3,830	100.0	24.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,216	31.7	7.6
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	399	10.4	2.5
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	364	9.5	2.3
4	Malignant neoplasms (C00-C97)	321	8.4	2.0
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	149	3.9	0.9
6	Influenza and pneumonia (J09-J18)	109	2.8	0.7
7	Septicemia (A40-A41)	53	1.4	0.3
7	Chronic lower respiratory diseases (J40-J47)	53	1.4	0.3
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	38	1.0	0.2
9	Certain conditions originating in the perinatal period (P00-P96)	38	1.0	0.2
11	Cerebrovascular diseases (I60-I69)	33	0.9	0.2
12	Complications of medical and surgical care (Y40-Y84,Y88)	17	0.4	*
13	Meningitis (G00,G03)	16	0.4	*
13	Acute bronchitis and bronchiolitis (J20-J21)	16	0.4	*
15	Anemias (D50-D64)	15	0.4	*
...	All other causes (residual)	993	25.9	6.2

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2014

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2014

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
All races, both sexes, 5-9 years				
...	All causes	2,357	100.0	11.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	730	31.0	3.6
2	Malignant neoplasms (C00-C97)	436	18.5	2.1
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	192	8.1	0.9
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	123	5.2	0.6
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	69	2.9	0.3
6	Chronic lower respiratory diseases (J40-J47)	68	2.9	0.3
7	Influenza and pneumonia (J09-J18)	57	2.4	0.3
8	Cerebrovascular diseases (I60-I69)	45	1.9	0.2
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	36	1.5	0.2
10	Septicemia (A40-A41)	33	1.4	0.2
11	Certain conditions originating in the perinatal period (P00-P96)	16	0.7	*
12	Anemias (D50-D64)	14	0.6	*
13	Diabetes mellitus (E10-E14)	7	0.3	*
14	Meningitis (G00,G03)	6	0.3	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	5	0.2	*
15	Complications of medical and surgical care (Y40-Y84,Y88)	5	0.2	*
...	All other causes (residual)	515	21.8	2.5
All races, both sexes, 10-14 years				
...	All causes	2,893	100.0	14.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	750	25.9	3.6
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	425	14.7	2.1
3	Malignant neoplasms (C00-C97)	416	14.4	2.0
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	156	5.4	0.8
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	156	5.4	0.8
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	122	4.2	0.6
7	Chronic lower respiratory diseases (J40-J47)	71	2.5	0.3
8	Cerebrovascular diseases (I60-I69)	43	1.5	0.2
9	Influenza and pneumonia (J09-J18)	41	1.4	0.2
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	38	1.3	0.2
11	Septicemia (A40-A41)	30	1.0	0.1
12	Anemias (D50-D64)	20	0.7	0.1
13	Diabetes mellitus (E10-E14)	19	0.7	*
14	Certain conditions originating in the perinatal period (P00-P96)	16	0.6	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	8	0.3	*
15	Complications of medical and surgical care (Y40-Y84,Y88)	8	0.3	*
...	All other causes (residual)	574	19.8	2.8

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2014

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2014

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
All races, both sexes, 15-19 years				
...	All causes	9,586	100.0	45.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,736	39.0	17.7
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,834	19.1	8.7
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,397	14.6	6.6
4	Malignant neoplasms (C00-C97)	612	6.4	2.9
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	299	3.1	1.4
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	179	1.9	0.8
7	Influenza and pneumonia (J09-J18)	64	0.7	0.3
8	Cerebrovascular diseases (I60-I69)	58	0.6	0.3
9	Chronic lower respiratory diseases (J40-J47)	55	0.6	0.3
10	Septicemia (A40-A41)	39	0.4	0.2
11	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	38	0.4	0.2
12	Diabetes mellitus (E10-E14)	34	0.4	0.2
13	Pregnancy, childbirth and the puerperium (O00-O99)	31	0.3	0.1
14	Anemias (D50-D64)	27	0.3	0.1
15	Legal intervention (Y35,Y89.0)	24	0.3	0.1
...	All other causes (residual)	1,159	12.1	5.5
All races, both sexes, 20-24 years				
...	All causes	19,205	100.0	83.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	8,100	42.2	35.4
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	3,245	16.9	14.2
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	2,747	14.3	12.0
4	Malignant neoplasms (C00-C97)	957	5.0	4.2
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	654	3.4	2.9
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	198	1.0	0.9
7	Diabetes mellitus (E10-E14)	147	0.8	0.6
8	Influenza and pneumonia (J09-J18)	135	0.7	0.6
9	Pregnancy, childbirth and the puerperium (O00-O99)	132	0.7	0.6
10	Chronic lower respiratory diseases (J40-J47)	123	0.6	0.5
11	Cerebrovascular diseases (I60-I69)	119	0.6	0.5
12	Human immunodeficiency virus (HIV) disease (B20-B24)	109	0.6	0.5
13	Septicemia (A40-A41)	87	0.5	0.4
14	Anemias (D50-D64)	58	0.3	0.3
14	Legal intervention (Y35,Y89.0)	58	0.3	0.3
...	All other causes (residual)	2,336	12.2	10.2

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2014

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2014

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
All races, both sexes, 25-29 years				
...	All causes	21,925	100.0	99.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	8,737	39.8	39.7
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	3,283	15.0	14.9
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	2,302	10.5	10.5
4	Malignant neoplasms (C00-C97)	1,371	6.3	6.2
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,153	5.3	5.2
6	Diabetes mellitus (E10-E14)	252	1.1	1.1
7	Influenza and pneumonia (J09-J18)	231	1.1	1.1
8	Human immunodeficiency virus (HIV) disease (B20-B24)	228	1.0	1.0
8	Cerebrovascular diseases (I60-I69)	228	1.0	1.0
10	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	222	1.0	1.0
11	Chronic liver disease and cirrhosis (K70,K73-K74)	207	0.9	0.9
12	Pregnancy, childbirth and the puerperium (O00-O99)	179	0.8	0.8
13	Chronic lower respiratory diseases (J40-J47)	168	0.8	0.8
14	Septicemia (A40-A41)	143	0.7	0.7
15	Legal intervention (Y35,Y89.0)	101	0.5	0.5
...	All other causes (residual)	3,120	14.2	14.2
All races, both sexes, 30-34 years				
...	All causes	25,252	100.0	117.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	8,620	34.1	40.0
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	3,286	13.0	15.3
3	Malignant neoplasms (C00-C97)	2,253	8.9	10.5
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,188	8.7	10.2
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,857	7.4	8.6
6	Chronic liver disease and cirrhosis (K70,K73-K74)	518	2.1	2.4
7	Diabetes mellitus (E10-E14)	457	1.8	2.1
8	Human immunodeficiency virus (HIV) disease (B20-B24)	355	1.4	1.6
9	Cerebrovascular diseases (I60-I69)	351	1.4	1.6
10	Influenza and pneumonia (J09-J18)	318	1.3	1.5
11	Pregnancy, childbirth and the puerperium (O00-O99)	226	0.9	1.0
12	Septicemia (A40-A41)	216	0.9	1.0
13	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	206	0.8	1.0
14	Chronic lower respiratory diseases (J40-J47)	199	0.8	0.9
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	152	0.6	0.7
...	All other causes (residual)	4,050	16.0	18.8

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2014

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2014

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
All races, both sexes, 35-39 years				
...	All causes	29,325	100.0	147.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	7,860	26.8	39.5
2	Malignant neoplasms (C00-C97)	3,879	13.2	19.5
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,541	12.1	17.8
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	3,188	10.9	16.0
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,421	4.8	7.1
6	Chronic liver disease and cirrhosis (K70,K73-K74)	846	2.9	4.2
7	Diabetes mellitus (E10-E14)	738	2.5	3.7
8	Cerebrovascular diseases (I60-I69)	616	2.1	3.1
9	Human immunodeficiency virus (HIV) disease (B20-B24)	463	1.6	2.3
10	Influenza and pneumonia (J09-J18)	429	1.5	2.2
11	Septicemia (A40-A41)	320	1.1	1.6
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	259	0.9	1.3
13	Chronic lower respiratory diseases (J40-J47)	253	0.9	1.3
14	Pregnancy, childbirth and the puerperium (O00-O99)	211	0.7	1.1
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	188	0.6	0.9
...	All other causes (residual)	5,113	17.4	25.7
All races, both sexes, 40-44 years				
...	All causes	41,671	100.0	202.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	8,188	19.6	39.8
2	Malignant neoplasms (C00-C97)	7,388	17.7	35.9
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	6,827	16.4	33.2
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	3,518	8.4	17.1
5	Chronic liver disease and cirrhosis (K70,K73-K74)	1,736	4.2	8.4
6	Diabetes mellitus (E10-E14)	1,261	3.0	6.1
7	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,167	2.8	5.7
8	Cerebrovascular diseases (I60-I69)	1,129	2.7	5.5
9	Human immunodeficiency virus (HIV) disease (B20-B24)	711	1.7	3.5
10	Influenza and pneumonia (J09-J18)	696	1.7	3.4
11	Septicemia (A40-A41)	512	1.2	2.5
12	Chronic lower respiratory diseases (J40-J47)	501	1.2	2.4
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	414	1.0	2.0
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	295	0.7	1.4
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	221	0.5	1.1
...	All other causes (residual)	7,107	17.1	34.5

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2014

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2014

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
All races, both sexes, 45-49 years				
...	All causes	65,016	100.0	311.3
1	Malignant neoplasms (C00-C97)	14,519	22.3	69.5
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	12,182	18.7	58.3
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	9,271	14.3	44.4
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	4,112	6.3	19.7
5	Chronic liver disease and cirrhosis (K70,K73-K74)	3,094	4.8	14.8
6	Diabetes mellitus (E10-E14)	2,229	3.4	10.7
7	Cerebrovascular diseases (I60-I69)	1,933	3.0	9.3
8	Chronic lower respiratory diseases (J40-J47)	1,149	1.8	5.5
9	Influenza and pneumonia (J09-J18)	1,062	1.6	5.1
10	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,029	1.6	4.9
11	Human immunodeficiency virus (HIV) disease (B20-B24)	989	1.5	4.7
12	Septicemia (A40-A41)	873	1.3	4.2
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	719	1.1	3.4
14	Viral hepatitis (B15-B19)	507	0.8	2.4
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	489	0.8	2.3
...	All other causes (residual)	10,859	16.7	52.0
All races, both sexes, 50-54 years				
...	All causes	110,901	100.0	491.3
1	Malignant neoplasms (C00-C97)	30,315	27.3	134.3
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	22,609	20.4	100.2
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	11,339	10.2	50.2
4	Chronic liver disease and cirrhosis (K70,K73-K74)	5,533	5.0	24.5
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	4,655	4.2	20.6
6	Diabetes mellitus (E10-E14)	3,833	3.5	17.0
7	Cerebrovascular diseases (I60-I69)	3,416	3.1	15.1
8	Chronic lower respiratory diseases (J40-J47)	3,253	2.9	14.4
9	Influenza and pneumonia (J09-J18)	1,669	1.5	7.4
10	Septicemia (A40-A41)	1,641	1.5	7.3
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,313	1.2	5.8
12	Viral hepatitis (B15-B19)	1,267	1.1	5.6
13	Human immunodeficiency virus (HIV) disease (B20-B24)	1,245	1.1	5.5
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	939	0.8	4.2
15	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	914	0.8	4.0
...	All other causes (residual)	16,960	15.3	75.1

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2014

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2014

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
All races, both sexes, 55-59 years				
...	All causes	157,170	100.0	730.6
1	Malignant neoplasms (C00-C97)	49,939	31.8	232.2
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	32,987	21.0	153.3
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	10,250	6.5	47.6
4	Chronic liver disease and cirrhosis (K70,K73-K74)	6,726	4.3	31.3
5	Chronic lower respiratory diseases (J40-J47)	6,607	4.2	30.7
6	Diabetes mellitus (E10-E14)	5,742	3.7	26.7
7	Cerebrovascular diseases (I60-I69)	5,066	3.2	23.6
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	4,300	2.7	20.0
9	Septicemia (A40-A41)	2,551	1.6	11.9
10	Influenza and pneumonia (J09-J18)	2,401	1.5	11.2
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2,071	1.3	9.6
12	Viral hepatitis (B15-B19)	1,955	1.2	9.1
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,424	0.9	6.6
14	Human immunodeficiency virus (HIV) disease (B20-B24)	1,059	0.7	4.9
15	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	708	0.5	3.3
...	All other causes (residual)	23,384	14.9	108.7
All races, both sexes, 60-64 years				
...	All causes	191,638	100.0	1,032.2
1	Malignant neoplasms (C00-C97)	65,343	34.1	351.9
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	41,486	21.6	223.4
3	Chronic lower respiratory diseases (J40-J47)	9,885	5.2	53.2
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	7,780	4.1	41.9
5	Diabetes mellitus (E10-E14)	7,600	4.0	40.9
6	Cerebrovascular diseases (I60-I69)	6,661	3.5	35.9
7	Chronic liver disease and cirrhosis (K70,K73-K74)	6,066	3.2	32.7
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	3,227	1.7	17.4
9	Septicemia (A40-A41)	3,158	1.6	17.0
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2,992	1.6	16.1
11	Influenza and pneumonia (J09-J18)	2,989	1.6	16.1
12	Viral hepatitis (B15-B19)	1,975	1.0	10.6
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,938	1.0	10.4
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	964	0.5	5.2
15	Aortic aneurysm and dissection (I71)	784	0.4	4.2
...	All other causes (residual)	28,790	15.0	155.1

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2014

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2014

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
All races, both sexes, 65-69 years				
...	All causes	222,834	100.0	1,454.0
1	Malignant neoplasms (C00-C97)	77,895	35.0	508.3
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	48,142	21.6	314.1
3	Chronic lower respiratory diseases (J40-J47)	15,086	6.8	98.4
4	Diabetes mellitus (E10-E14)	9,042	4.1	59.0
5	Cerebrovascular diseases (I60-I69)	8,569	3.8	55.9
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	6,113	2.7	39.9
7	Chronic liver disease and cirrhosis (K70,K73-K74)	4,515	2.0	29.5
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	4,095	1.8	26.7
9	Septicemia (A40-A41)	3,882	1.7	25.3
10	Influenza and pneumonia (J09-J18)	3,563	1.6	23.2
11	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,306	1.0	15.0
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	2,100	0.9	13.7
13	Alzheimer's disease (G30)	1,418	0.6	9.3
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1,294	0.6	8.4
15	Pneumonitis due to solids and liquids (J69)	1,075	0.5	7.0
...	All other causes (residual)	33,739	15.1	220.2
All races, both sexes, 70-74 years				
...	All causes	248,707	100.0	2,246.1
1	Malignant neoplasms (C00-C97)	81,313	32.7	734.3
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	53,541	21.5	483.5
3	Chronic lower respiratory diseases (J40-J47)	20,531	8.3	185.4
4	Cerebrovascular diseases (I60-I69)	11,094	4.5	100.2
5	Diabetes mellitus (E10-E14)	9,162	3.7	82.7
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	5,796	2.3	52.3
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	4,965	2.0	44.8
8	Septicemia (A40-A41)	4,338	1.7	39.2
9	Influenza and pneumonia (J09-J18)	4,298	1.7	38.8
10	Alzheimer's disease (G30)	3,752	1.5	33.9
11	Chronic liver disease and cirrhosis (K70,K73-K74)	3,294	1.3	29.7
12	Parkinson's disease (G20-G21)	2,401	1.0	21.7
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	2,364	1.0	21.3
14	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,804	0.7	16.3
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1,687	0.7	15.2
...	All other causes (residual)	38,367	15.4	346.5

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2014

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2014

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
All races, both sexes, 75-79 years				
...	All causes	282,072	100.0	3,560.5
1	Malignant neoplasms (C00-C97)	79,173	28.1	999.4
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	62,999	22.3	795.2
3	Chronic lower respiratory diseases (J40-J47)	22,878	8.1	288.8
4	Cerebrovascular diseases (I60-I69)	15,253	5.4	192.5
5	Diabetes mellitus (E10-E14)	9,496	3.4	119.9
6	Alzheimer's disease (G30)	8,548	3.0	107.9
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	6,613	2.3	83.5
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	5,998	2.1	75.7
9	Influenza and pneumonia (J09-J18)	5,539	2.0	69.9
10	Septicemia (A40-A41)	4,703	1.7	59.4
11	Parkinson's disease (G20-G21)	4,376	1.6	55.2
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	2,911	1.0	36.7
13	Chronic liver disease and cirrhosis (K70,K73-K74)	2,459	0.9	31.0
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2,129	0.8	26.9
15	Pneumonitis due to solids and liquids (J69)	2,031	0.7	25.6
...	All other causes (residual)	46,966	16.7	592.8
All races, both sexes, 80-84 years				
...	All causes	342,432	100.0	5,944.6
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	83,427	24.4	1,448.3
2	Malignant neoplasms (C00-C97)	74,880	21.9	1,299.9
3	Chronic lower respiratory diseases (J40-J47)	24,880	7.3	431.9
4	Cerebrovascular diseases (I60-I69)	21,100	6.2	366.3
5	Alzheimer's disease (G30)	16,845	4.9	292.4
6	Diabetes mellitus (E10-E14)	9,911	2.9	172.1
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	8,260	2.4	143.4
8	Influenza and pneumonia (J09-J18)	7,654	2.2	132.9
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	7,499	2.2	130.2
10	Parkinson's disease (G20-G21)	6,459	1.9	112.1
11	Septicemia (A40-A41)	5,301	1.5	92.0
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	4,112	1.2	71.4
13	Pneumonitis due to solids and liquids (J69)	2,984	0.9	51.8
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2,662	0.8	46.2
15	Chronic liver disease and cirrhosis (K70,K73-K74)	1,698	0.5	29.5
...	All other causes (residual)	64,760	18.9	1,124.2

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2014

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2014

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
All races, both sexes, 85-89 years				
...	All causes	381,539	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	103,682	27.2	---
2	Malignant neoplasms (C00-C97)	60,120	15.8	---
3	Cerebrovascular diseases (I60-I69)	26,024	6.8	---
4	Alzheimer's disease (G30)	25,352	6.6	---
5	Chronic lower respiratory diseases (J40-J47)	22,710	6.0	---
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	9,826	2.6	---
7	Influenza and pneumonia (J09-J18)	9,814	2.6	---
8	Diabetes mellitus (E10-E14)	8,973	2.4	---
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	8,463	2.2	---
10	Parkinson's disease (G20-G21)	6,657	1.7	---
11	Septicemia (A40-A41)	5,556	1.5	---
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	5,414	1.4	---
13	Pneumonitis due to solids and liquids (J69)	3,893	1.0	---
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2,816	0.7	---
15	Aortic aneurysm and dissection (I71)	1,420	0.4	---
...	All other causes (residual)	80,819	21.2	---
All races, both sexes, 90-94 years				
...	All causes	300,366	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	90,664	30.2	---
2	Malignant neoplasms (C00-C97)	31,423	10.5	---
3	Alzheimer's disease (G30)	24,021	8.0	---
4	Cerebrovascular diseases (I60-I69)	21,486	7.2	---
5	Chronic lower respiratory diseases (J40-J47)	13,732	4.6	---
6	Influenza and pneumonia (J09-J18)	8,946	3.0	---
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	8,122	2.7	---
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	6,220	2.1	---
9	Diabetes mellitus (E10-E14)	5,580	1.9	---
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	4,930	1.6	---
11	Septicemia (A40-A41)	3,855	1.3	---
12	Parkinson's disease (G20-G21)	3,604	1.2	---
13	Pneumonitis due to solids and liquids (J69)	3,490	1.2	---
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1,878	0.6	---
15	Atherosclerosis (I70)	1,356	0.5	---
...	All other causes (residual)	71,059	23.7	---

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2014

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2014

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
All races, both sexes, 95-99 years				
...	All causes	118,407	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	38,308	32.4	---
2	Alzheimer's disease (G30)	10,476	8.8	---
3	Cerebrovascular diseases (I60-I69)	8,191	6.9	---
4	Malignant neoplasms (C00-C97)	8,022	6.8	---
5	Chronic lower respiratory diseases (J40-J47)	4,198	3.5	---
6	Influenza and pneumonia (J09-J18)	3,994	3.4	---
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,989	2.5	---
8	Essential hypertension and hypertensive renal disease (I10,I12,I15)	2,377	2.0	---
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2,288	1.9	---
10	Diabetes mellitus (E10-E14)	1,717	1.5	---
11	Pneumonitis due to solids and liquids (J69)	1,417	1.2	---
12	Septicemia (A40-A41)	1,257	1.1	---
13	Parkinson's disease (G20-G21)	848	0.7	---
14	Atherosclerosis (I70)	698	0.6	---
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	615	0.5	---
...	All other causes (residual)	31,012	26.2	---
All races, both sexes, 100 years and over				
...	All causes	25,914	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	8,959	34.6	---
2	Alzheimer's disease (G30)	2,192	8.5	---
3	Cerebrovascular diseases (I60-I69)	1,591	6.1	---
4	Malignant neoplasms (C00-C97)	1,059	4.1	---
5	Influenza and pneumonia (J09-J18)	1,028	4.0	---
6	Chronic lower respiratory diseases (J40-J47)	678	2.6	---
7	Essential hypertension and hypertensive renal disease (I10,I12,I15)	653	2.5	---
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	576	2.2	---
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	429	1.7	---
10	Pneumonitis due to solids and liquids (J69)	308	1.2	---
11	Diabetes mellitus (E10-E14)	280	1.1	---
12	Septicemia (A40-A41)	232	0.9	---
13	Atherosclerosis (I70)	212	0.8	---
14	Parkinson's disease (G20-G21)	107	0.4	---
15	Nutritional deficiencies (E40-E64)	91	0.4	---
...	All other causes (residual)	7,519	29.0	---

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2014

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2014

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
All races, male, all ages				
...	All causes	1,328,241	100.0	846.4
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	325,077	24.5	207.1
2	Malignant neoplasms (C00-C97)	311,296	23.4	198.4
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	85,448	6.4	54.4
4	Chronic lower respiratory diseases (J40-J47)	69,456	5.2	44.3
5	Cerebrovascular diseases (I60-I69)	55,471	4.2	35.3
6	Diabetes mellitus (E10-E14)	41,111	3.1	26.2
7	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	33,113	2.5	21.1
8	Alzheimer's disease (G30)	28,362	2.1	18.1
9	Influenza and pneumonia (J09-J18)	26,586	2.0	16.9
10	Chronic liver disease and cirrhosis (K70,K73-K74)	24,584	1.9	15.7
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	24,436	1.8	15.6
12	Septicemia (A40-A41)	18,333	1.4	11.7
13	Parkinson's disease (G20-G21)	15,681	1.2	10.0
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	12,989	1.0	8.3
15	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	12,491	0.9	8.0
...	All other causes (residual)	243,807	18.4	155.4
All races, male, 1-4 years				
...	All causes	2,172	100.0	26.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	744	34.3	9.1
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	219	10.1	2.7
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	196	9.0	2.4
4	Malignant neoplasms (C00-C97)	182	8.4	2.2
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	76	3.5	0.9
6	Influenza and pneumonia (J09-J18)	57	2.6	0.7
7	Chronic lower respiratory diseases (J40-J47)	30	1.4	0.4
8	Certain conditions originating in the perinatal period (P00-P96)	24	1.1	0.3
9	Septicemia (A40-A41)	23	1.1	0.3
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	19	0.9	*
10	Cerebrovascular diseases (I60-I69)	19	0.9	*
12	Anemias (D50-D64)	10	0.5	*
13	Meningitis (G00,G03)	7	0.3	*
13	Acute bronchitis and bronchiolitis (J20-J21)	7	0.3	*
13	Complications of medical and surgical care (Y40-Y84,Y88)	7	0.3	*
...	All other causes (residual)	552	25.4	6.8

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2014

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2014

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
All races, male, 5-9 years				
...	All causes	1,357	100.0	13.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	464	34.2	4.4
2	Malignant neoplasms (C00-C97)	242	17.8	2.3
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	89	6.6	0.8
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	61	4.5	0.6
5	Chronic lower respiratory diseases (J40-J47)	46	3.4	0.4
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	40	2.9	0.4
7	Influenza and pneumonia (J09-J18)	29	2.1	0.3
8	Cerebrovascular diseases (I60-I69)	20	1.5	0.2
9	Septicemia (A40-A41)	17	1.3	*
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	15	1.1	*
11	Certain conditions originating in the perinatal period (P00-P96)	11	0.8	*
12	Anemias (D50-D64)	6	0.4	*
13	Diabetes mellitus (E10-E14)	4	0.3	*
13	Meningitis (G00,G03)	4	0.3	*
15	Diseases of appendix (K35-K38)	3	0.2	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3	0.2	*
15	Complications of medical and surgical care (Y40-Y84,Y88)	3	0.2	*
...	All other causes (residual)	300	22.1	2.9
All races, male, 10-14 years				
...	All causes	1,771	100.0	16.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	504	28.5	4.8
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	275	15.5	2.6
3	Malignant neoplasms (C00-C97)	246	13.9	2.3
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	105	5.9	1.0
5	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	96	5.4	0.9
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	68	3.8	0.6
7	Chronic lower respiratory diseases (J40-J47)	44	2.5	0.4
8	Cerebrovascular diseases (I60-I69)	27	1.5	0.3
9	Influenza and pneumonia (J09-J18)	21	1.2	0.2
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	18	1.0	*
11	Septicemia (A40-A41)	17	1.0	*
12	Anemias (D50-D64)	11	0.6	*
12	Diabetes mellitus (E10-E14)	11	0.6	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	6	0.3	*
15	Certain conditions originating in the perinatal period (P00-P96)	5	0.3	*
...	All other causes (residual)	317	17.9	3.0

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2014

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2014

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
All races, male, 15-19 years				
...	All causes	6,828	100.0	63.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,697	39.5	25.0
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,404	20.6	13.0
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,201	17.6	11.1
4	Malignant neoplasms (C00-C97)	358	5.2	3.3
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	189	2.8	1.8
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	94	1.4	0.9
7	Influenza and pneumonia (J09-J18)	38	0.6	0.4
7	Chronic lower respiratory diseases (J40-J47)	38	0.6	0.4
9	Cerebrovascular diseases (I60-I69)	30	0.4	0.3
10	Diabetes mellitus (E10-E14)	23	0.3	0.2
11	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	21	0.3	0.2
12	Septicemia (A40-A41)	20	0.3	0.2
12	Legal intervention (Y35,Y89.0)	20	0.3	0.2
14	Aortic aneurysm and dissection (I71)	8	0.1	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	7	0.1	*
...	All other causes (residual)	680	10.0	6.3
All races, male, 20-24 years				
...	All causes	14,289	100.0	121.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	6,106	42.7	52.0
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,685	18.8	22.9
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	2,397	16.8	20.4
4	Malignant neoplasms (C00-C97)	594	4.2	5.1
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	427	3.0	3.6
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	117	0.8	1.0
7	Diabetes mellitus (E10-E14)	92	0.6	0.8
8	Influenza and pneumonia (J09-J18)	78	0.5	0.7
8	Chronic lower respiratory diseases (J40-J47)	78	0.5	0.7
10	Cerebrovascular diseases (I60-I69)	72	0.5	0.6
11	Human immunodeficiency virus (HIV) disease (B20-B24)	71	0.5	0.6
12	Legal intervention (Y35,Y89.0)	58	0.4	0.5
13	Septicemia (A40-A41)	45	0.3	0.4
14	Anemias (D50-D64)	35	0.2	0.3
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	32	0.2	0.3
...	All other causes (residual)	1,402	9.8	11.9

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2014

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2014

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
All races, male, 25-29 years				
...	All causes	15,619	100.0	139.9
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	6,482	41.5	58.1
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,631	16.8	23.6
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,948	12.5	17.5
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	787	5.0	7.1
5	Malignant neoplasms (C00-C97)	726	4.6	6.5
6	Human immunodeficiency virus (HIV) disease (B20-B24)	174	1.1	1.6
7	Diabetes mellitus (E10-E14)	151	1.0	1.4
8	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	140	0.9	1.3
9	Cerebrovascular diseases (I60-I69)	132	0.8	1.2
10	Influenza and pneumonia (J09-J18)	131	0.8	1.2
11	Chronic liver disease and cirrhosis (K70,K73-K74)	121	0.8	1.1
12	Chronic lower respiratory diseases (J40-J47)	100	0.6	0.9
13	Legal intervention (Y35,Y89.0)	96	0.6	0.9
14	Septicemia (A40-A41)	80	0.5	0.7
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	50	0.3	0.4
...	All other causes (residual)	1,870	12.0	16.8
All races, male, 30-34 years				
...	All causes	17,078	100.0	158.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	6,359	37.2	58.8
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,591	15.2	24.0
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,529	9.0	14.1
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,479	8.7	13.7
5	Malignant neoplasms (C00-C97)	1,068	6.3	9.9
6	Chronic liver disease and cirrhosis (K70,K73-K74)	316	1.9	2.9
7	Diabetes mellitus (E10-E14)	281	1.6	2.6
8	Human immunodeficiency virus (HIV) disease (B20-B24)	239	1.4	2.2
9	Cerebrovascular diseases (I60-I69)	199	1.2	1.8
10	Influenza and pneumonia (J09-J18)	157	0.9	1.5
11	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	114	0.7	1.1
12	Chronic lower respiratory diseases (J40-J47)	113	0.7	1.0
13	Septicemia (A40-A41)	105	0.6	1.0
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	79	0.5	0.7
15	Legal intervention (Y35,Y89.0)	64	0.4	0.6
...	All other causes (residual)	2,385	14.0	22.1

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2014

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2014

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
All races, male, 35-39 years				
...	All causes	18,500	100.0	186.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	5,520	29.8	55.5
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,446	13.2	24.6
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,384	12.9	24.0
4	Malignant neoplasms (C00-C97)	1,632	8.8	16.4
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,166	6.3	11.7
6	Chronic liver disease and cirrhosis (K70,K73-K74)	516	2.8	5.2
7	Diabetes mellitus (E10-E14)	446	2.4	4.5
8	Cerebrovascular diseases (I60-I69)	363	2.0	3.7
9	Human immunodeficiency virus (HIV) disease (B20-B24)	300	1.6	3.0
10	Influenza and pneumonia (J09-J18)	236	1.3	2.4
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	158	0.9	1.6
12	Septicemia (A40-A41)	146	0.8	1.5
13	Chronic lower respiratory diseases (J40-J47)	129	0.7	1.3
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	113	0.6	1.1
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	101	0.5	1.0
...	All other causes (residual)	2,844	15.4	28.6
All races, male, 40-44 years				
...	All causes	25,193	100.0	246.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	5,546	22.0	54.3
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	4,637	18.4	45.4
3	Malignant neoplasms (C00-C97)	3,197	12.7	31.3
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,589	10.3	25.3
5	Chronic liver disease and cirrhosis (K70,K73-K74)	1,101	4.4	10.8
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	896	3.6	8.8
7	Diabetes mellitus (E10-E14)	765	3.0	7.5
8	Cerebrovascular diseases (I60-I69)	638	2.5	6.2
9	Human immunodeficiency virus (HIV) disease (B20-B24)	474	1.9	4.6
10	Influenza and pneumonia (J09-J18)	380	1.5	3.7
11	Septicemia (A40-A41)	273	1.1	2.7
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	241	1.0	2.4
13	Chronic lower respiratory diseases (J40-J47)	230	0.9	2.3
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	198	0.8	1.9
15	Aortic aneurysm and dissection (I71)	157	0.6	1.5
...	All other causes (residual)	3,871	15.4	37.9

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2014

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2014

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
All races, male, 45-49 years				
...	All causes	39,281	100.0	379.6
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	8,526	21.7	82.4
2	Malignant neoplasms (C00-C97)	6,733	17.1	65.1
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	6,243	15.9	60.3
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	3,048	7.8	29.5
5	Chronic liver disease and cirrhosis (K70,K73-K74)	1,997	5.1	19.3
6	Diabetes mellitus (E10-E14)	1,411	3.6	13.6
7	Cerebrovascular diseases (I60-I69)	1,078	2.7	10.4
8	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	770	2.0	7.4
9	Human immunodeficiency virus (HIV) disease (B20-B24)	714	1.8	6.9
10	Influenza and pneumonia (J09-J18)	584	1.5	5.6
11	Chronic lower respiratory diseases (J40-J47)	526	1.3	5.1
12	Septicemia (A40-A41)	454	1.2	4.4
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	430	1.1	4.2
14	Viral hepatitis (B15-B19)	357	0.9	3.5
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	318	0.8	3.1
...	All other causes (residual)	6,092	15.5	58.9
All races, male, 50-54 years				
...	All causes	67,096	100.0	605.7
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	15,790	23.5	142.5
2	Malignant neoplasms (C00-C97)	15,320	22.8	138.3
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	7,598	11.3	68.6
4	Chronic liver disease and cirrhosis (K70,K73-K74)	3,730	5.6	33.7
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	3,372	5.0	30.4
6	Diabetes mellitus (E10-E14)	2,384	3.6	21.5
7	Cerebrovascular diseases (I60-I69)	1,917	2.9	17.3
8	Chronic lower respiratory diseases (J40-J47)	1,476	2.2	13.3
9	Human immunodeficiency virus (HIV) disease (B20-B24)	948	1.4	8.6
10	Influenza and pneumonia (J09-J18)	904	1.3	8.2
11	Septicemia (A40-A41)	889	1.3	8.0
12	Viral hepatitis (B15-B19)	856	1.3	7.7
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	801	1.2	7.2
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	695	1.0	6.3
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	602	0.9	5.4
...	All other causes (residual)	9,814	14.6	88.6

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2014

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2014

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
All races, male, 55-59 years				
...	All causes	95,992	100.0	919.1
1	Malignant neoplasms (C00-C97)	27,228	28.4	260.7
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	23,187	24.2	222.0
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	6,872	7.2	65.8
4	Chronic liver disease and cirrhosis (K70,K73-K74)	4,615	4.8	44.2
5	Diabetes mellitus (E10-E14)	3,558	3.7	34.1
6	Chronic lower respiratory diseases (J40-J47)	3,318	3.5	31.8
7	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	3,187	3.3	30.5
8	Cerebrovascular diseases (I60-I69)	2,952	3.1	28.3
9	Influenza and pneumonia (J09-J18)	1,368	1.4	13.1
10	Viral hepatitis (B15-B19)	1,360	1.4	13.0
11	Septicemia (A40-A41)	1,297	1.4	12.4
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,213	1.3	11.6
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	902	0.9	8.6
14	Human immunodeficiency virus (HIV) disease (B20-B24)	807	0.8	7.7
15	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	508	0.5	4.9
...	All other causes (residual)	13,620	14.2	130.4
All races, male, 60-64 years				
...	All causes	116,206	100.0	1,308.9
1	Malignant neoplasms (C00-C97)	36,597	31.5	412.2
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	28,608	24.6	322.2
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	5,302	4.6	59.7
4	Chronic lower respiratory diseases (J40-J47)	5,231	4.5	58.9
5	Diabetes mellitus (E10-E14)	4,638	4.0	52.2
6	Chronic liver disease and cirrhosis (K70,K73-K74)	4,256	3.7	47.9
7	Cerebrovascular diseases (I60-I69)	3,844	3.3	43.3
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,492	2.1	28.1
9	Septicemia (A40-A41)	1,710	1.5	19.3
10	Influenza and pneumonia (J09-J18)	1,708	1.5	19.2
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,705	1.5	19.2
12	Viral hepatitis (B15-B19)	1,410	1.2	15.9
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,177	1.0	13.3
14	Human immunodeficiency virus (HIV) disease (B20-B24)	582	0.5	6.6
15	Aortic aneurysm and dissection (I71)	570	0.5	6.4
...	All other causes (residual)	16,376	14.1	184.5

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2014

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2014

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
All races, male, 65-69 years				
...	All causes	129,802	100.0	1,790.6
1	Malignant neoplasms (C00-C97)	43,185	33.3	595.7
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	31,687	24.4	437.1
3	Chronic lower respiratory diseases (J40-J47)	7,828	6.0	108.0
4	Diabetes mellitus (E10-E14)	5,351	4.1	73.8
5	Cerebrovascular diseases (I60-I69)	4,702	3.6	64.9
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	4,013	3.1	55.4
7	Chronic liver disease and cirrhosis (K70,K73-K74)	2,990	2.3	41.2
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2,225	1.7	30.7
9	Septicemia (A40-A41)	2,027	1.6	28.0
10	Influenza and pneumonia (J09-J18)	1,999	1.5	27.6
11	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,788	1.4	24.7
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,232	0.9	17.0
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	785	0.6	10.8
14	Parkinson's disease (G20-G21)	682	0.5	9.4
15	Aortic aneurysm and dissection (I71)	656	0.5	9.0
...	All other causes (residual)	18,652	14.4	257.3
All races, male, 70-74 years				
...	All causes	138,846	100.0	2,722.5
1	Malignant neoplasms (C00-C97)	44,729	32.2	877.0
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	33,230	23.9	651.6
3	Chronic lower respiratory diseases (J40-J47)	10,268	7.4	201.3
4	Cerebrovascular diseases (I60-I69)	5,802	4.2	113.8
5	Diabetes mellitus (E10-E14)	5,248	3.8	102.9
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,507	2.5	68.8
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2,602	1.9	51.0
8	Influenza and pneumonia (J09-J18)	2,369	1.7	46.5
9	Septicemia (A40-A41)	2,237	1.6	43.9
10	Chronic liver disease and cirrhosis (K70,K73-K74)	2,009	1.4	39.4
11	Parkinson's disease (G20-G21)	1,589	1.1	31.2
12	Alzheimer's disease (G30)	1,587	1.1	31.1
13	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,494	1.1	29.3
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,243	0.9	24.4
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	986	0.7	19.3
...	All other causes (residual)	19,946	14.4	391.1

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2014

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2014

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
All races, male, 75-79 years				
...	All causes	149,259	100.0	4,250.5
1	Malignant neoplasms (C00-C97)	42,564	28.5	1,212.1
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	36,401	24.4	1,036.6
3	Chronic lower respiratory diseases (J40-J47)	11,181	7.5	318.4
4	Cerebrovascular diseases (I60-I69)	7,297	4.9	207.8
5	Diabetes mellitus (E10-E14)	5,098	3.4	145.2
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,766	2.5	107.2
7	Alzheimer's disease (G30)	3,406	2.3	97.0
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3,166	2.1	90.2
9	Influenza and pneumonia (J09-J18)	2,954	2.0	84.1
10	Parkinson's disease (G20-G21)	2,910	1.9	82.9
11	Septicemia (A40-A41)	2,253	1.5	64.2
12	Chronic liver disease and cirrhosis (K70,K73-K74)	1,365	0.9	38.9
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,364	0.9	38.8
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1,243	0.8	35.4
15	Pneumonitis due to solids and liquids (J69)	1,204	0.8	34.3
...	All other causes (residual)	23,087	15.5	657.5
All races, male, 80-84 years				
...	All causes	167,171	100.0	7,018.6
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	43,443	26.0	1,823.9
2	Malignant neoplasms (C00-C97)	39,204	23.5	1,646.0
3	Chronic lower respiratory diseases (J40-J47)	11,777	7.0	494.5
4	Cerebrovascular diseases (I60-I69)	8,778	5.3	368.5
5	Alzheimer's disease (G30)	6,019	3.6	252.7
6	Diabetes mellitus (E10-E14)	4,900	2.9	205.7
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	4,373	2.6	183.6
8	Parkinson's disease (G20-G21)	4,007	2.4	168.2
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3,828	2.3	160.7
10	Influenza and pneumonia (J09-J18)	3,802	2.3	159.6
11	Septicemia (A40-A41)	2,451	1.5	102.9
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,714	1.0	72.0
13	Pneumonitis due to solids and liquids (J69)	1,707	1.0	71.7
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1,471	0.9	61.8
15	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	966	0.6	40.6
...	All other causes (residual)	28,731	17.2	1,206.3

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2014

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2014

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
All races, male, 85-89 years				
...	All causes	164,609	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	46,984	28.5	---
2	Malignant neoplasms (C00-C97)	29,986	18.2	---
3	Chronic lower respiratory diseases (J40-J47)	10,083	6.1	---
4	Cerebrovascular diseases (I60-I69)	9,261	5.6	---
5	Alzheimer's disease (G30)	7,972	4.8	---
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	4,531	2.8	---
7	Influenza and pneumonia (J09-J18)	4,485	2.7	---
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	4,090	2.5	---
9	Diabetes mellitus (E10-E14)	4,065	2.5	---
10	Parkinson's disease (G20-G21)	3,837	2.3	---
11	Septicemia (A40-A41)	2,371	1.4	---
12	Pneumonitis due to solids and liquids (J69)	2,156	1.3	---
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,955	1.2	---
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1,389	0.8	---
15	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	689	0.4	---
...	All other causes (residual)	30,755	18.7	---
All races, male, 90-94 years				
...	All causes	107,387	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	34,181	31.8	---
2	Malignant neoplasms (C00-C97)	14,111	13.1	---
3	Cerebrovascular diseases (I60-I69)	6,245	5.8	---
4	Alzheimer's disease (G30)	6,230	5.8	---
5	Chronic lower respiratory diseases (J40-J47)	5,369	5.0	---
6	Influenza and pneumonia (J09-J18)	3,631	3.4	---
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,146	2.9	---
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2,778	2.6	---
9	Diabetes mellitus (E10-E14)	2,096	2.0	---
10	Parkinson's disease (G20-G21)	1,826	1.7	---
11	Pneumonitis due to solids and liquids (J69)	1,767	1.6	---
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,449	1.3	---
13	Septicemia (A40-A41)	1,378	1.3	---
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	818	0.8	---
15	Atherosclerosis (I70)	443	0.4	---
...	All other causes (residual)	21,919	20.4	---

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2014

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2014

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
All races, male, 95-99 years				
...	All causes	32,226	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	11,059	34.3	---
2	Malignant neoplasms (C00-C97)	3,046	9.5	---
3	Alzheimer's disease (G30)	1,927	6.0	---
4	Cerebrovascular diseases (I60-I69)	1,809	5.6	---
5	Chronic lower respiratory diseases (J40-J47)	1,418	4.4	---
6	Influenza and pneumonia (J09-J18)	1,306	4.1	---
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	869	2.7	---
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	848	2.6	---
9	Pneumonitis due to solids and liquids (J69)	618	1.9	---
10	Diabetes mellitus (E10-E14)	542	1.7	---
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	526	1.6	---
12	Septicemia (A40-A41)	394	1.2	---
13	Parkinson's disease (G20-G21)	356	1.1	---
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	200	0.6	---
15	Atherosclerosis (I70)	181	0.6	---
...	All other causes (residual)	7,127	22.1	---
All races, male, 100 years and over				
...	All causes	4,563	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,693	37.1	---
2	Malignant neoplasms (C00-C97)	315	6.9	---
3	Influenza and pneumonia (J09-J18)	258	5.7	---
4	Cerebrovascular diseases (I60-I69)	231	5.1	---
5	Alzheimer's disease (G30)	205	4.5	---
6	Chronic lower respiratory diseases (J40-J47)	154	3.4	---
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	128	2.8	---
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	121	2.7	---
9	Essential hypertension and hypertensive renal disease (I10,I12,I15)	118	2.6	---
10	Pneumonitis due to solids and liquids (J69)	97	2.1	---
11	Septicemia (A40-A41)	58	1.3	---
12	Diabetes mellitus (E10-E14)	41	0.9	---
13	Atherosclerosis (I70)	37	0.8	---
14	Parkinson's disease (G20-G21)	33	0.7	---
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	15	0.3	---
15	Nutritional deficiencies (E40-E64)	15	0.3	---
...	All other causes (residual)	1,044	22.9	---

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2014

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2014

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
All races, female, all ages				
...	All causes	1,298,177	100.0	801.7
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	289,271	22.3	178.6
2	Malignant neoplasms (C00-C97)	280,403	21.6	173.2
3	Chronic lower respiratory diseases (J40-J47)	77,645	6.0	48.0
4	Cerebrovascular diseases (I60-I69)	77,632	6.0	47.9
5	Alzheimer's disease (G30)	65,179	5.0	40.3
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	50,605	3.9	31.3
7	Diabetes mellitus (E10-E14)	35,377	2.7	21.8
8	Influenza and pneumonia (J09-J18)	28,641	2.2	17.7
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	23,710	1.8	14.6
10	Septicemia (A40-A41)	20,607	1.6	12.7
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	17,232	1.3	10.6
12	Chronic liver disease and cirrhosis (K70,K73-K74)	13,586	1.0	8.4
13	Parkinson's disease (G20-G21)	10,469	0.8	6.5
14	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	9,660	0.7	6.0
15	Pneumonitis due to solids and liquids (J69)	8,434	0.6	5.2
...	All other causes (residual)	289,726	22.3	178.9
All races, female, 1-4 years				
...	All causes	1,658	100.0	21.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	472	28.5	6.1
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	180	10.9	2.3
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	168	10.1	2.2
4	Malignant neoplasms (C00-C97)	139	8.4	1.8
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	73	4.4	0.9
6	Influenza and pneumonia (J09-J18)	52	3.1	0.7
7	Septicemia (A40-A41)	30	1.8	0.4
8	Chronic lower respiratory diseases (J40-J47)	23	1.4	0.3
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	19	1.1	*
10	Cerebrovascular diseases (I60-I69)	14	0.8	*
10	Certain conditions originating in the perinatal period (P00-P96)	14	0.8	*
12	Complications of medical and surgical care (Y40-Y84,Y88)	10	0.6	*
13	Meningitis (G00,G03)	9	0.5	*
13	Acute bronchitis and bronchiolitis (J20-J21)	9	0.5	*
15	Anemias (D50-D64)	5	0.3	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	5	0.3	*
...	All other causes (residual)	436	26.3	5.6

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2014

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2014

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
All races, female, 5-9 years				
...	All causes	1,000	100.0	10.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	266	26.6	2.6
2	Malignant neoplasms (C00-C97)	194	19.4	1.9
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	103	10.3	1.0
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	62	6.2	0.6
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	29	2.9	0.3
6	Influenza and pneumonia (J09-J18)	28	2.8	0.3
7	Cerebrovascular diseases (I60-I69)	25	2.5	0.2
8	Chronic lower respiratory diseases (J40-J47)	22	2.2	0.2
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	21	2.1	0.2
10	Septicemia (A40-A41)	16	1.6	*
11	Anemias (D50-D64)	8	0.8	*
12	Certain conditions originating in the perinatal period (P00-P96)	5	0.5	*
13	Diabetes mellitus (E10-E14)	3	0.3	*
14	Tuberculosis (A16-A19)	2	0.2	*
14	Meningitis (G00,G03)	2	0.2	*
14	Hernia (K40-K46)	2	0.2	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2	0.2	*
14	Complications of medical and surgical care (Y40-Y84,Y88)	2	0.2	*
...	All other causes (residual)	208	20.8	2.1
All races, female, 10-14 years				
...	All causes	1,122	100.0	11.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	246	21.9	2.4
2	Malignant neoplasms (C00-C97)	170	15.2	1.7
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	150	13.4	1.5
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	60	5.3	0.6
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	54	4.8	0.5
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	51	4.5	0.5
7	Chronic lower respiratory diseases (J40-J47)	27	2.4	0.3
8	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	20	1.8	0.2
8	Influenza and pneumonia (J09-J18)	20	1.8	0.2
10	Cerebrovascular diseases (I60-I69)	16	1.4	*
11	Septicemia (A40-A41)	13	1.2	*
12	Certain conditions originating in the perinatal period (P00-P96)	11	1.0	*
13	Anemias (D50-D64)	9	0.8	*
14	Diabetes mellitus (E10-E14)	8	0.7	*
15	Complications of medical and surgical care (Y40-Y84,Y88)	4	0.4	*
...	All other causes (residual)	263	23.4	2.6

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2014

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2014

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
All races, female, 15-19 years				
...	All causes	2,758	100.0	26.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,039	37.7	10.1
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	430	15.6	4.2
3	Malignant neoplasms (C00-C97)	254	9.2	2.5
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	196	7.1	1.9
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	110	4.0	1.1
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	85	3.1	0.8
7	Pregnancy, childbirth and the puerperium (O00-O99)	31	1.1	0.3
8	Cerebrovascular diseases (I60-I69)	28	1.0	0.3
9	Influenza and pneumonia (J09-J18)	26	0.9	0.3
10	Anemias (D50-D64)	21	0.8	0.2
11	Septicemia (A40-A41)	19	0.7	*
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	17	0.6	*
12	Chronic lower respiratory diseases (J40-J47)	17	0.6	*
14	Diabetes mellitus (E10-E14)	11	0.4	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	9	0.3	*
...	All other causes (residual)	465	16.9	4.5
All races, female, 20-24 years				
...	All causes	4,916	100.0	44.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,994	40.6	17.8
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	560	11.4	5.0
3	Malignant neoplasms (C00-C97)	363	7.4	3.2
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	350	7.1	3.1
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	227	4.6	2.0
6	Pregnancy, childbirth and the puerperium (O00-O99)	132	2.7	1.2
7	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	81	1.6	0.7
8	Influenza and pneumonia (J09-J18)	57	1.2	0.5
9	Diabetes mellitus (E10-E14)	55	1.1	0.5
10	Cerebrovascular diseases (I60-I69)	47	1.0	0.4
11	Chronic lower respiratory diseases (J40-J47)	45	0.9	0.4
12	Septicemia (A40-A41)	42	0.9	0.4
13	Human immunodeficiency virus (HIV) disease (B20-B24)	38	0.8	0.3
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	24	0.5	0.2
15	Anemias (D50-D64)	23	0.5	0.2
...	All other causes (residual)	878	17.9	7.9

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2014

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2014

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
All races, female, 25-29 years				
...	All causes	6,306	100.0	58.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,255	35.8	20.8
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	652	10.3	6.0
3	Malignant neoplasms (C00-C97)	645	10.2	6.0
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	366	5.8	3.4
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	354	5.6	3.3
6	Pregnancy, childbirth and the puerperium (O00-O99)	179	2.8	1.7
7	Diabetes mellitus (E10-E14)	101	1.6	0.9
8	Influenza and pneumonia (J09-J18)	100	1.6	0.9
9	Cerebrovascular diseases (I60-I69)	96	1.5	0.9
10	Chronic liver disease and cirrhosis (K70,K73-K74)	86	1.4	0.8
11	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	82	1.3	0.8
12	Chronic lower respiratory diseases (J40-J47)	68	1.1	0.6
13	Septicemia (A40-A41)	63	1.0	0.6
14	Human immunodeficiency virus (HIV) disease (B20-B24)	54	0.9	0.5
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	34	0.5	0.3
...	All other causes (residual)	1,171	18.6	10.8
All races, female, 30-34 years				
...	All causes	8,174	100.0	76.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,261	27.7	21.1
2	Malignant neoplasms (C00-C97)	1,185	14.5	11.1
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	709	8.7	6.6
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	695	8.5	6.5
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	328	4.0	3.1
6	Pregnancy, childbirth and the puerperium (O00-O99)	226	2.8	2.1
7	Chronic liver disease and cirrhosis (K70,K73-K74)	202	2.5	1.9
8	Diabetes mellitus (E10-E14)	176	2.2	1.6
9	Influenza and pneumonia (J09-J18)	161	2.0	1.5
10	Cerebrovascular diseases (I60-I69)	152	1.9	1.4
11	Human immunodeficiency virus (HIV) disease (B20-B24)	116	1.4	1.1
12	Septicemia (A40-A41)	111	1.4	1.0
13	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	92	1.1	0.9
14	Chronic lower respiratory diseases (J40-J47)	86	1.1	0.8
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	73	0.9	0.7
...	All other causes (residual)	1,601	19.6	14.9

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2014

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2014

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
All races, female, 35-39 years				
...	All causes	10,825	100.0	108.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,340	21.6	23.4
2	Malignant neoplasms (C00-C97)	2,247	20.8	22.5
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,157	10.7	11.6
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	742	6.9	7.4
5	Chronic liver disease and cirrhosis (K70,K73-K74)	330	3.0	3.3
6	Diabetes mellitus (E10-E14)	292	2.7	2.9
7	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	255	2.4	2.6
8	Cerebrovascular diseases (I60-I69)	253	2.3	2.5
9	Pregnancy, childbirth and the puerperium (O00-O99)	211	1.9	2.1
10	Influenza and pneumonia (J09-J18)	193	1.8	1.9
11	Septicemia (A40-A41)	174	1.6	1.7
12	Human immunodeficiency virus (HIV) disease (B20-B24)	163	1.5	1.6
13	Chronic lower respiratory diseases (J40-J47)	124	1.1	1.2
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	101	0.9	1.0
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	75	0.7	0.8
...	All other causes (residual)	2,168	20.0	21.7
All races, female, 40-44 years				
...	All causes	16,478	100.0	158.9
1	Malignant neoplasms (C00-C97)	4,191	25.4	40.4
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,642	16.0	25.5
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,190	13.3	21.1
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	929	5.6	9.0
5	Chronic liver disease and cirrhosis (K70,K73-K74)	635	3.9	6.1
6	Diabetes mellitus (E10-E14)	496	3.0	4.8
7	Cerebrovascular diseases (I60-I69)	491	3.0	4.7
8	Influenza and pneumonia (J09-J18)	316	1.9	3.0
9	Chronic lower respiratory diseases (J40-J47)	271	1.6	2.6
9	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	271	1.6	2.6
11	Septicemia (A40-A41)	239	1.5	2.3
12	Human immunodeficiency virus (HIV) disease (B20-B24)	237	1.4	2.3
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	173	1.0	1.7
14	Pregnancy, childbirth and the puerperium (O00-O99)	139	0.8	1.3
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	116	0.7	1.1
...	All other causes (residual)	3,142	19.1	30.3

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2014

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2014

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
All races, female, 45-49 years				
...	All causes	25,735	100.0	244.2
1	Malignant neoplasms (C00-C97)	7,786	30.3	73.9
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,656	14.2	34.7
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,028	11.8	28.7
4	Chronic liver disease and cirrhosis (K70,K73-K74)	1,097	4.3	10.4
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,064	4.1	10.1
6	Cerebrovascular diseases (I60-I69)	855	3.3	8.1
7	Diabetes mellitus (E10-E14)	818	3.2	7.8
8	Chronic lower respiratory diseases (J40-J47)	623	2.4	5.9
9	Influenza and pneumonia (J09-J18)	478	1.9	4.5
10	Septicemia (A40-A41)	419	1.6	4.0
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	289	1.1	2.7
12	Human immunodeficiency virus (HIV) disease (B20-B24)	275	1.1	2.6
13	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	259	1.0	2.5
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	171	0.7	1.6
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	162	0.6	1.5
...	All other causes (residual)	4,755	18.5	45.1
All races, female, 50-54 years				
...	All causes	43,805	100.0	381.1
1	Malignant neoplasms (C00-C97)	14,995	34.2	130.5
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	6,819	15.6	59.3
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,741	8.5	32.5
4	Chronic liver disease and cirrhosis (K70,K73-K74)	1,803	4.1	15.7
5	Chronic lower respiratory diseases (J40-J47)	1,777	4.1	15.5
6	Cerebrovascular diseases (I60-I69)	1,499	3.4	13.0
7	Diabetes mellitus (E10-E14)	1,449	3.3	12.6
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,283	2.9	11.2
9	Influenza and pneumonia (J09-J18)	765	1.7	6.7
10	Septicemia (A40-A41)	752	1.7	6.5
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	512	1.2	4.5
12	Viral hepatitis (B15-B19)	411	0.9	3.6
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	337	0.8	2.9
14	Human immunodeficiency virus (HIV) disease (B20-B24)	297	0.7	2.6
15	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	219	0.5	1.9
...	All other causes (residual)	7,146	16.3	62.2

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2014

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2014

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
All races, female, 55-59 years				
...	All causes	61,178	100.0	552.8
1	Malignant neoplasms (C00-C97)	22,711	37.1	205.2
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	9,800	16.0	88.5
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,378	5.5	30.5
4	Chronic lower respiratory diseases (J40-J47)	3,289	5.4	29.7
5	Diabetes mellitus (E10-E14)	2,184	3.6	19.7
6	Cerebrovascular diseases (I60-I69)	2,114	3.5	19.1
7	Chronic liver disease and cirrhosis (K70,K73-K74)	2,111	3.5	19.1
8	Septicemia (A40-A41)	1,254	2.0	11.3
9	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,113	1.8	10.1
10	Influenza and pneumonia (J09-J18)	1,033	1.7	9.3
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	858	1.4	7.8
12	Viral hepatitis (B15-B19)	595	1.0	5.4
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	522	0.9	4.7
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	259	0.4	2.3
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	259	0.4	2.3
...	All other causes (residual)	9,698	15.9	87.6
All races, female, 60-64 years				
...	All causes	75,432	100.0	778.6
1	Malignant neoplasms (C00-C97)	28,746	38.1	296.7
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	12,878	17.1	132.9
3	Chronic lower respiratory diseases (J40-J47)	4,654	6.2	48.0
4	Diabetes mellitus (E10-E14)	2,962	3.9	30.6
5	Cerebrovascular diseases (I60-I69)	2,817	3.7	29.1
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,478	3.3	25.6
7	Chronic liver disease and cirrhosis (K70,K73-K74)	1,810	2.4	18.7
8	Septicemia (A40-A41)	1,448	1.9	14.9
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,287	1.7	13.3
10	Influenza and pneumonia (J09-J18)	1,281	1.7	13.2
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	761	1.0	7.9
12	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	735	1.0	7.6
13	Viral hepatitis (B15-B19)	565	0.7	5.8
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	406	0.5	4.2
15	Alzheimer's disease (G30)	356	0.5	3.7
...	All other causes (residual)	12,248	16.2	126.4

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2014

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2014

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
All races, female, 65-69 years				
...	All causes	93,032	100.0	1,151.9
1	Malignant neoplasms (C00-C97)	34,710	37.3	429.8
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	16,455	17.7	203.7
3	Chronic lower respiratory diseases (J40-J47)	7,258	7.8	89.9
4	Cerebrovascular diseases (I60-I69)	3,867	4.2	47.9
5	Diabetes mellitus (E10-E14)	3,691	4.0	45.7
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,100	2.3	26.0
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,870	2.0	23.2
8	Septicemia (A40-A41)	1,855	2.0	23.0
9	Influenza and pneumonia (J09-J18)	1,564	1.7	19.4
10	Chronic liver disease and cirrhosis (K70,K73-K74)	1,525	1.6	18.9
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	868	0.9	10.7
12	Alzheimer's disease (G30)	799	0.9	9.9
13	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	518	0.6	6.4
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	509	0.5	6.3
15	Pneumonitis due to solids and liquids (J69)	433	0.5	5.4
...	All other causes (residual)	15,010	16.1	185.9
All races, female, 70-74 years				
...	All causes	109,861	100.0	1,839.3
1	Malignant neoplasms (C00-C97)	36,584	33.3	612.5
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	20,311	18.5	340.0
3	Chronic lower respiratory diseases (J40-J47)	10,263	9.3	171.8
4	Cerebrovascular diseases (I60-I69)	5,292	4.8	88.6
5	Diabetes mellitus (E10-E14)	3,914	3.6	65.5
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2,363	2.2	39.6
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,289	2.1	38.3
8	Alzheimer's disease (G30)	2,165	2.0	36.2
9	Septicemia (A40-A41)	2,101	1.9	35.2
10	Influenza and pneumonia (J09-J18)	1,929	1.8	32.3
11	Chronic liver disease and cirrhosis (K70,K73-K74)	1,285	1.2	21.5
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,121	1.0	18.8
13	Parkinson's disease (G20-G21)	812	0.7	13.6
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	701	0.6	11.7
15	Pneumonitis due to solids and liquids (J69)	563	0.5	9.4
...	All other causes (residual)	18,168	16.5	304.2

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2014

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2014

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
All races, female, 75-79 years				
...	All causes	132,813	100.0	3,011.1
1	Malignant neoplasms (C00-C97)	36,609	27.6	830.0
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	26,598	20.0	603.0
3	Chronic lower respiratory diseases (J40-J47)	11,697	8.8	265.2
4	Cerebrovascular diseases (I60-I69)	7,956	6.0	180.4
5	Alzheimer's disease (G30)	5,142	3.9	116.6
6	Diabetes mellitus (E10-E14)	4,398	3.3	99.7
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,847	2.1	64.5
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2,832	2.1	64.2
9	Influenza and pneumonia (J09-J18)	2,585	1.9	58.6
10	Septicemia (A40-A41)	2,450	1.8	55.5
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,547	1.2	35.1
12	Parkinson's disease (G20-G21)	1,466	1.1	33.2
13	Chronic liver disease and cirrhosis (K70,K73-K74)	1,094	0.8	24.8
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	886	0.7	20.1
15	Pneumonitis due to solids and liquids (J69)	827	0.6	18.7
...	All other causes (residual)	23,879	18.0	541.4
All races, female, 80-84 years				
...	All causes	175,261	100.0	5,187.5
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	39,984	22.8	1,183.5
2	Malignant neoplasms (C00-C97)	35,676	20.4	1,056.0
3	Chronic lower respiratory diseases (J40-J47)	13,103	7.5	387.8
4	Cerebrovascular diseases (I60-I69)	12,322	7.0	364.7
5	Alzheimer's disease (G30)	10,826	6.2	320.4
6	Diabetes mellitus (E10-E14)	5,011	2.9	148.3
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,887	2.2	115.0
8	Influenza and pneumonia (J09-J18)	3,852	2.2	114.0
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3,671	2.1	108.7
10	Septicemia (A40-A41)	2,850	1.6	84.4
11	Parkinson's disease (G20-G21)	2,452	1.4	72.6
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	2,398	1.4	71.0
13	Pneumonitis due to solids and liquids (J69)	1,277	0.7	37.8
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1,191	0.7	35.3
15	Chronic liver disease and cirrhosis (K70,K73-K74)	825	0.5	24.4
...	All other causes (residual)	35,936	20.5	1,063.7

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2014

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2014

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
All races, female, 85-89 years				
...	All causes	216,930	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	56,698	26.1	---
2	Malignant neoplasms (C00-C97)	30,134	13.9	---
3	Alzheimer's disease (G30)	17,380	8.0	---
4	Cerebrovascular diseases (I60-I69)	16,763	7.7	---
5	Chronic lower respiratory diseases (J40-J47)	12,627	5.8	---
6	Influenza and pneumonia (J09-J18)	5,329	2.5	---
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	5,295	2.4	---
8	Diabetes mellitus (E10-E14)	4,908	2.3	---
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	4,373	2.0	---
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	3,459	1.6	---
11	Septicemia (A40-A41)	3,185	1.5	---
12	Parkinson's disease (G20-G21)	2,820	1.3	---
13	Pneumonitis due to solids and liquids (J69)	1,737	0.8	---
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1,427	0.7	---
15	Enterocolitis due to <i>Clostridium difficile</i> (A04.7)	897	0.4	---
...	All other causes (residual)	49,898	23.0	---
All races, female, 90-94 years				
...	All causes	192,979	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	56,483	29.3	---
2	Alzheimer's disease (G30)	17,791	9.2	---
3	Malignant neoplasms (C00-C97)	17,312	9.0	---
4	Cerebrovascular diseases (I60-I69)	15,241	7.9	---
5	Chronic lower respiratory diseases (J40-J47)	8,363	4.3	---
6	Influenza and pneumonia (J09-J18)	5,315	2.8	---
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	4,976	2.6	---
8	Diabetes mellitus (E10-E14)	3,484	1.8	---
9	Essential hypertension and hypertensive renal disease (I10,I12,I15)	3,481	1.8	---
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3,442	1.8	---
11	Septicemia (A40-A41)	2,477	1.3	---
12	Parkinson's disease (G20-G21)	1,778	0.9	---
13	Pneumonitis due to solids and liquids (J69)	1,723	0.9	---
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1,060	0.5	---
15	Atherosclerosis (I70)	913	0.5	---
...	All other causes (residual)	49,140	25.5	---

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2014

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2014

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
All races, female, 95-99 years				
...	All causes	86,181	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	27,249	31.6	---
2	Alzheimer's disease (G30)	8,549	9.9	---
3	Cerebrovascular diseases (I60-I69)	6,382	7.4	---
4	Malignant neoplasms (C00-C97)	4,976	5.8	---
5	Chronic lower respiratory diseases (J40-J47)	2,780	3.2	---
6	Influenza and pneumonia (J09-J18)	2,688	3.1	---
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,120	2.5	---
8	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,851	2.1	---
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,440	1.7	---
10	Diabetes mellitus (E10-E14)	1,175	1.4	---
11	Septicemia (A40-A41)	863	1.0	---
12	Pneumonitis due to solids and liquids (J69)	799	0.9	---
13	Atherosclerosis (I70)	517	0.6	---
14	Parkinson's disease (G20-G21)	492	0.6	---
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	415	0.5	---
...	All other causes (residual)	23,885	27.7	---
All races, female, 100 years and over				
...	All causes	21,351	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	7,266	34.0	---
2	Alzheimer's disease (G30)	1,987	9.3	---
3	Cerebrovascular diseases (I60-I69)	1,360	6.4	---
4	Influenza and pneumonia (J09-J18)	770	3.6	---
5	Malignant neoplasms (C00-C97)	744	3.5	---
6	Essential hypertension and hypertensive renal disease (I10,I12,I15)	535	2.5	---
7	Chronic lower respiratory diseases (J40-J47)	524	2.5	---
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	448	2.1	---
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	308	1.4	---
10	Diabetes mellitus (E10-E14)	239	1.1	---
11	Pneumonitis due to solids and liquids (J69)	211	1.0	---
12	Atherosclerosis (I70)	175	0.8	---
13	Septicemia (A40-A41)	174	0.8	---
14	Nutritional deficiencies (E40-E64)	76	0.4	---
15	Parkinson's disease (G20-G21)	74	0.3	---
...	All other causes (residual)	6,460	30.3	---

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2014

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2014

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
White, both sexes, all ages				
...	All causes	2,237,880	100.0	892.9
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	524,695	23.4	209.4
2	Malignant neoplasms (C00-C97)	502,932	22.5	200.7
3	Chronic lower respiratory diseases (J40-J47)	134,541	6.0	53.7
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	117,242	5.2	46.8
5	Cerebrovascular diseases (I60-I69)	111,035	5.0	44.3
6	Alzheimer's disease (G30)	84,990	3.8	33.9
7	Diabetes mellitus (E10-E14)	59,741	2.7	23.8
8	Influenza and pneumonia (J09-J18)	47,293	2.1	18.9
9	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	38,675	1.7	15.4
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	37,976	1.7	15.2
11	Chronic liver disease and cirrhosis (K70,K73-K74)	33,508	1.5	13.4
12	Septicemia (A40-A41)	31,512	1.4	12.6
13	Parkinson's disease (G20-G21)	24,294	1.1	9.7
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	23,639	1.1	9.4
15	Pneumonitis due to solids and liquids (J69)	16,487	0.7	6.6
...	All other causes (residual)	449,320	20.1	179.3
White, both sexes, 1-4 years				
...	All causes	2,592	100.0	21.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	859	33.1	7.2
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	271	10.5	2.3
3	Malignant neoplasms (C00-C97)	250	9.6	2.1
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	200	7.7	1.7
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	84	3.2	0.7
6	Influenza and pneumonia (J09-J18)	72	2.8	0.6
7	Septicemia (A40-A41)	37	1.4	0.3
8	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	28	1.1	0.2
9	Chronic lower respiratory diseases (J40-J47)	25	1.0	0.2
10	Certain conditions originating in the perinatal period (P00-P96)	24	0.9	0.2
11	Cerebrovascular diseases (I60-I69)	20	0.8	0.2
12	Meningitis (G00,G03)	12	0.5	*
13	Complications of medical and surgical care (Y40-Y84,Y88)	10	0.4	*
14	Acute bronchitis and bronchiolitis (J20-J21)	9	0.3	*
15	Anemias (D50-D64)	8	0.3	*
...	All other causes (residual)	683	26.4	5.7

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2014

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2014

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
White, both sexes, 5-9 years				
...	All causes	1,683	100.0	10.9
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	534	31.7	3.5
2	Malignant neoplasms (C00-C97)	344	20.4	2.2
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	141	8.4	0.9
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	73	4.3	0.5
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	45	2.7	0.3
6	Influenza and pneumonia (J09-J18)	41	2.4	0.3
7	Cerebrovascular diseases (I60-I69)	32	1.9	0.2
8	Chronic lower respiratory diseases (J40-J47)	27	1.6	0.2
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	25	1.5	0.2
10	Septicemia (A40-A41)	22	1.3	0.1
11	Certain conditions originating in the perinatal period (P00-P96)	11	0.7	*
12	Anemias (D50-D64)	4	0.2	*
12	Diabetes mellitus (E10-E14)	4	0.2	*
14	Meningitis (G00,G03)	3	0.2	*
14	Pneumonitis due to solids and liquids (J69)	3	0.2	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3	0.2	*
14	Complications of medical and surgical care (Y40-Y84,Y88)	3	0.2	*
...	All other causes (residual)	368	21.9	2.4
White, both sexes, 10-14 years				
...	All causes	2,070	100.0	13.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	533	25.7	3.4
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	347	16.8	2.2
3	Malignant neoplasms (C00-C97)	322	15.6	2.1
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	113	5.5	0.7
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	87	4.2	0.6
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	71	3.4	0.5
7	Cerebrovascular diseases (I60-I69)	37	1.8	0.2
8	Influenza and pneumonia (J09-J18)	32	1.5	0.2
8	Chronic lower respiratory diseases (J40-J47)	32	1.5	0.2
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	25	1.2	0.2
11	Septicemia (A40-A41)	19	0.9	*
12	Diabetes mellitus (E10-E14)	10	0.5	*
12	Certain conditions originating in the perinatal period (P00-P96)	10	0.5	*
14	Anemias (D50-D64)	9	0.4	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	5	0.2	*
...	All other causes (residual)	418	20.2	2.7

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2014

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2014

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
White, both sexes, 15-19 years				
...	All causes	6,954	100.0	43.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,011	43.3	18.8
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,552	22.3	9.7
3	Malignant neoplasms (C00-C97)	476	6.8	3.0
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	462	6.6	2.9
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	201	2.9	1.3
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	145	2.1	0.9
7	Influenza and pneumonia (J09-J18)	53	0.8	0.3
8	Cerebrovascular diseases (I60-I69)	42	0.6	0.3
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	30	0.4	0.2
9	Chronic lower respiratory diseases (J40-J47)	30	0.4	0.2
11	Septicemia (A40-A41)	27	0.4	0.2
11	Diabetes mellitus (E10-E14)	27	0.4	0.2
13	Pregnancy, childbirth and the puerperium (O00-O99)	20	0.3	0.1
14	Legal intervention (Y35,Y89.0)	13	0.2	*
15	Anemias (D50-D64)	9	0.1	*
15	Aortic aneurysm and dissection (I71)	9	0.1	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	9	0.1	*
15	Complications of medical and surgical care (Y40-Y84,Y88)	9	0.1	*
...	All other causes (residual)	829	11.9	5.2
White, both sexes, 20-24 years				
...	All causes	13,806	100.0	80.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	6,748	48.9	39.3
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,625	19.0	15.3
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	894	6.5	5.2
4	Malignant neoplasms (C00-C97)	733	5.3	4.3
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	420	3.0	2.4
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	143	1.0	0.8
7	Influenza and pneumonia (J09-J18)	101	0.7	0.6
8	Diabetes mellitus (E10-E14)	91	0.7	0.5
9	Pregnancy, childbirth and the puerperium (O00-O99)	79	0.6	0.5
10	Cerebrovascular diseases (I60-I69)	78	0.6	0.5
11	Chronic lower respiratory diseases (J40-J47)	65	0.5	0.4
12	Septicemia (A40-A41)	62	0.4	0.4
13	Legal intervention (Y35,Y89.0)	34	0.2	0.2
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	32	0.2	0.2
15	Human immunodeficiency virus (HIV) disease (B20-B24)	29	0.2	0.2
...	All other causes (residual)	1,672	12.1	9.7

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2014

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2014

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
White, both sexes, 25-29 years				
...	All causes	16,347	100.0	98.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	7,368	45.1	44.2
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,786	17.0	16.7
3	Malignant neoplasms (C00-C97)	1,066	6.5	6.4
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	859	5.3	5.2
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	754	4.6	4.5
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	178	1.1	1.1
7	Influenza and pneumonia (J09-J18)	176	1.1	1.1
8	Chronic liver disease and cirrhosis (K70,K73-K74)	157	1.0	0.9
9	Cerebrovascular diseases (I60-I69)	152	0.9	0.9
10	Diabetes mellitus (E10-E14)	150	0.9	0.9
11	Pregnancy, childbirth and the puerperium (O00-O99)	107	0.7	0.6
12	Chronic lower respiratory diseases (J40-J47)	104	0.6	0.6
13	Septicemia (A40-A41)	100	0.6	0.6
14	Legal intervention (Y35,Y89.0)	68	0.4	0.4
15	Human immunodeficiency virus (HIV) disease (B20-B24)	61	0.4	0.4
...	All other causes (residual)	2,261	13.8	13.6
White, both sexes, 30-34 years				
...	All causes	19,061	100.0	115.9
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	7,294	38.3	44.3
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,848	14.9	17.3
3	Malignant neoplasms (C00-C97)	1,705	8.9	10.4
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,423	7.5	8.7
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	808	4.2	4.9
6	Chronic liver disease and cirrhosis (K70,K73-K74)	430	2.3	2.6
7	Diabetes mellitus (E10-E14)	285	1.5	1.7
8	Influenza and pneumonia (J09-J18)	248	1.3	1.5
9	Cerebrovascular diseases (I60-I69)	235	1.2	1.4
10	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	171	0.9	1.0
11	Septicemia (A40-A41)	158	0.8	1.0
12	Human immunodeficiency virus (HIV) disease (B20-B24)	133	0.7	0.8
13	Pregnancy, childbirth and the puerperium (O00-O99)	133	0.7	0.8
14	Chronic lower respiratory diseases (J40-J47)	126	0.7	0.8
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	93	0.5	0.6
...	All other causes (residual)	2,971	15.6	18.1

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2014

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2014

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
White, both sexes, 35-39 years				
...	All causes	21,984	100.0	144.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	6,678	30.4	43.8
2	Malignant neoplasms (C00-C97)	2,924	13.3	19.2
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,821	12.8	18.5
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,349	10.7	15.4
5	Chronic liver disease and cirrhosis (K70,K73-K74)	685	3.1	4.5
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	649	3.0	4.3
7	Diabetes mellitus (E10-E14)	491	2.2	3.2
8	Cerebrovascular diseases (I60-I69)	402	1.8	2.6
9	Influenza and pneumonia (J09-J18)	325	1.5	2.1
10	Septicemia (A40-A41)	229	1.0	1.5
11	Human immunodeficiency virus (HIV) disease (B20-B24)	184	0.8	1.2
12	Chronic lower respiratory diseases (J40-J47)	164	0.7	1.1
13	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	148	0.7	1.0
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	141	0.6	0.9
15	Pregnancy, childbirth and the puerperium (O00-O99)	133	0.6	0.9
...	All other causes (residual)	3,661	16.7	24.0
White, both sexes, 40-44 years				
...	All causes	31,690	100.0	198.9
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	6,908	21.8	43.4
2	Malignant neoplasms (C00-C97)	5,686	17.9	35.7
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	4,786	15.1	30.0
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	3,150	9.9	19.8
5	Chronic liver disease and cirrhosis (K70,K73-K74)	1,487	4.7	9.3
6	Diabetes mellitus (E10-E14)	864	2.7	5.4
7	Cerebrovascular diseases (I60-I69)	727	2.3	4.6
8	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	620	2.0	3.9
9	Influenza and pneumonia (J09-J18)	538	1.7	3.4
10	Septicemia (A40-A41)	376	1.2	2.4
11	Chronic lower respiratory diseases (J40-J47)	370	1.2	2.3
12	Human immunodeficiency virus (HIV) disease (B20-B24)	327	1.0	2.1
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	235	0.7	1.5
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	179	0.6	1.1
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	155	0.5	1.0
...	All other causes (residual)	5,282	16.7	33.1

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2014

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2014

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
White, both sexes, 45-49 years				
...	All causes	50,234	100.0	304.9
1	Malignant neoplasms (C00-C97)	11,328	22.6	68.7
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	8,845	17.6	53.7
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	7,836	15.6	47.6
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	3,802	7.6	23.1
5	Chronic liver disease and cirrhosis (K70,K73-K74)	2,708	5.4	16.4
6	Diabetes mellitus (E10-E14)	1,562	3.1	9.5
7	Cerebrovascular diseases (I60-I69)	1,283	2.6	7.8
8	Chronic lower respiratory diseases (J40-J47)	913	1.8	5.5
9	Influenza and pneumonia (J09-J18)	824	1.6	5.0
10	Septicemia (A40-A41)	629	1.3	3.8
11	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	583	1.2	3.5
12	Human immunodeficiency virus (HIV) disease (B20-B24)	458	0.9	2.8
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	443	0.9	2.7
14	Viral hepatitis (B15-B19)	437	0.9	2.7
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	285	0.6	1.7
...	All other causes (residual)	8,298	16.5	50.4
White, both sexes, 50-54 years				
...	All causes	87,263	100.0	480.8
1	Malignant neoplasms (C00-C97)	24,013	27.5	132.3
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	17,077	19.6	94.1
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	9,607	11.0	52.9
4	Chronic liver disease and cirrhosis (K70,K73-K74)	4,825	5.5	26.6
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	4,315	4.9	23.8
6	Diabetes mellitus (E10-E14)	2,804	3.2	15.4
7	Chronic lower respiratory diseases (J40-J47)	2,762	3.2	15.2
8	Cerebrovascular diseases (I60-I69)	2,290	2.6	12.6
9	Influenza and pneumonia (J09-J18)	1,302	1.5	7.2
10	Septicemia (A40-A41)	1,214	1.4	6.7
11	Viral hepatitis (B15-B19)	1,075	1.2	5.9
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	842	1.0	4.6
13	Human immunodeficiency virus (HIV) disease (B20-B24)	632	0.7	3.5
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	601	0.7	3.3
15	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	541	0.6	3.0
...	All other causes (residual)	13,363	15.3	73.6

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2014

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2014

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
White, both sexes, 55-59 years				
...	All causes	124,168	100.0	708.5
1	Malignant neoplasms (C00-C97)	39,970	32.2	228.1
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	25,325	20.4	144.5
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	8,527	6.9	48.7
4	Chronic liver disease and cirrhosis (K70,K73-K74)	5,801	4.7	33.1
5	Chronic lower respiratory diseases (J40-J47)	5,681	4.6	32.4
6	Diabetes mellitus (E10-E14)	4,166	3.4	23.8
7	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	4,051	3.3	23.1
8	Cerebrovascular diseases (I60-I69)	3,471	2.8	19.8
9	Influenza and pneumonia (J09-J18)	1,900	1.5	10.8
10	Septicemia (A40-A41)	1,882	1.5	10.7
11	Viral hepatitis (B15-B19)	1,582	1.3	9.0
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,334	1.1	7.6
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	928	0.7	5.3
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	529	0.4	3.0
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	500	0.4	2.9
...	All other causes (residual)	18,521	14.9	105.7
White, both sexes, 60-64 years				
...	All causes	153,412	100.0	998.9
1	Malignant neoplasms (C00-C97)	53,161	34.7	346.1
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	32,365	21.1	210.7
3	Chronic lower respiratory diseases (J40-J47)	8,719	5.7	56.8
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	6,485	4.2	42.2
5	Diabetes mellitus (E10-E14)	5,684	3.7	37.0
6	Chronic liver disease and cirrhosis (K70,K73-K74)	5,270	3.4	34.3
7	Cerebrovascular diseases (I60-I69)	4,705	3.1	30.6
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	3,046	2.0	19.8
9	Septicemia (A40-A41)	2,355	1.5	15.3
10	Influenza and pneumonia (J09-J18)	2,353	1.5	15.3
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,970	1.3	12.8
12	Viral hepatitis (B15-B19)	1,540	1.0	10.0
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,293	0.8	8.4
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	807	0.5	5.3
15	Aortic aneurysm and dissection (I71)	642	0.4	4.2
...	All other causes (residual)	23,017	15.0	149.9

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2014

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2014

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
White, both sexes, 65-69 years				
...	All causes	185,043	100.0	1,429.0
1	Malignant neoplasms (C00-C97)	65,541	35.4	506.2
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	39,061	21.1	301.7
3	Chronic lower respiratory diseases (J40-J47)	13,636	7.4	105.3
4	Diabetes mellitus (E10-E14)	7,078	3.8	54.7
5	Cerebrovascular diseases (I60-I69)	6,518	3.5	50.3
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	5,240	2.8	40.5
7	Chronic liver disease and cirrhosis (K70,K73-K74)	3,975	2.1	30.7
8	Septicemia (A40-A41)	3,037	1.6	23.5
9	Influenza and pneumonia (J09-J18)	2,950	1.6	22.8
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2,950	1.6	22.8
11	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,185	1.2	16.9
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,515	0.8	11.7
13	Alzheimer's disease (G30)	1,282	0.7	9.9
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1,110	0.6	8.6
15	Parkinson's disease (G20-G21)	922	0.5	7.1
...	All other causes (residual)	28,043	15.2	216.6
White, both sexes, 70-74 years				
...	All causes	211,385	100.0	2,234.9
1	Malignant neoplasms (C00-C97)	69,979	33.1	739.8
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	44,530	21.1	470.8
3	Chronic lower respiratory diseases (J40-J47)	18,891	8.9	199.7
4	Cerebrovascular diseases (I60-I69)	8,786	4.2	92.9
5	Diabetes mellitus (E10-E14)	7,148	3.4	75.6
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	5,085	2.4	53.8
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3,713	1.8	39.3
8	Influenza and pneumonia (J09-J18)	3,578	1.7	37.8
9	Septicemia (A40-A41)	3,477	1.6	36.8
10	Alzheimer's disease (G30)	3,316	1.6	35.1
11	Chronic liver disease and cirrhosis (K70,K73-K74)	2,988	1.4	31.6
12	Parkinson's disease (G20-G21)	2,211	1.0	23.4
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,743	0.8	18.4
14	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,715	0.8	18.1
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1,476	0.7	15.6
...	All other causes (residual)	32,749	15.5	346.2

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2014

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2014

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
White, both sexes, 75-79 years				
...	All causes	243,234	100.0	3,577.1
1	Malignant neoplasms (C00-C97)	68,859	28.3	1,012.7
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	53,469	22.0	786.3
3	Chronic lower respiratory diseases (J40-J47)	21,163	8.7	311.2
4	Cerebrovascular diseases (I60-I69)	12,628	5.2	185.7
5	Alzheimer's disease (G30)	7,582	3.1	111.5
6	Diabetes mellitus (E10-E14)	7,460	3.1	109.7
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	5,924	2.4	87.1
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	4,734	1.9	69.6
9	Influenza and pneumonia (J09-J18)	4,706	1.9	69.2
10	Parkinson's disease (G20-G21)	4,011	1.6	59.0
11	Septicemia (A40-A41)	3,850	1.6	56.6
12	Chronic liver disease and cirrhosis (K70,K73-K74)	2,249	0.9	33.1
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	2,205	0.9	32.4
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1,907	0.8	28.0
15	Pneumonitis due to solids and liquids (J69)	1,727	0.7	25.4
...	All other causes (residual)	40,760	16.8	599.4
White, both sexes, 80-84 years				
...	All causes	302,714	100.0	6,031.4
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	73,275	24.2	1,460.0
2	Malignant neoplasms (C00-C97)	66,169	21.9	1,318.4
3	Chronic lower respiratory diseases (J40-J47)	23,200	7.7	462.2
4	Cerebrovascular diseases (I60-I69)	18,183	6.0	362.3
5	Alzheimer's disease (G30)	15,232	5.0	303.5
6	Diabetes mellitus (E10-E14)	8,065	2.7	160.7
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	7,632	2.5	152.1
8	Influenza and pneumonia (J09-J18)	6,661	2.2	132.7
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	6,220	2.1	123.9
10	Parkinson's disease (G20-G21)	6,052	2.0	120.6
11	Septicemia (A40-A41)	4,505	1.5	89.8
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	3,320	1.1	66.1
13	Pneumonitis due to solids and liquids (J69)	2,649	0.9	52.8
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2,427	0.8	48.4
15	Chronic liver disease and cirrhosis (K70,K73-K74)	1,560	0.5	31.1
...	All other causes (residual)	57,564	19.0	1,146.9

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2014

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2014

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
White, both sexes, 85-89 years				
...	All causes	344,524	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	93,845	27.2	---
2	Malignant neoplasms (C00-C97)	53,897	15.6	---
3	Alzheimer's disease (G30)	23,254	6.7	---
4	Cerebrovascular diseases (I60-I69)	23,175	6.7	---
5	Chronic lower respiratory diseases (J40-J47)	21,257	6.2	---
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	9,225	2.7	---
7	Influenza and pneumonia (J09-J18)	8,779	2.5	---
8	Diabetes mellitus (E10-E14)	7,472	2.2	---
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	7,252	2.1	---
10	Parkinson's disease (G20-G21)	6,255	1.8	---
11	Septicemia (A40-A41)	4,805	1.4	---
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	4,594	1.3	---
13	Pneumonitis due to solids and liquids (J69)	3,539	1.0	---
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2,596	0.8	---
15	Aortic aneurysm and dissection (I71)	1,301	0.4	---
...	All other causes (residual)	73,278	21.3	---
White, both sexes, 90-94 years				
...	All causes	274,251	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	83,371	30.4	---
2	Malignant neoplasms (C00-C97)	28,394	10.4	---
3	Alzheimer's disease (G30)	22,040	8.0	---
4	Cerebrovascular diseases (I60-I69)	19,475	7.1	---
5	Chronic lower respiratory diseases (J40-J47)	12,876	4.7	---
6	Influenza and pneumonia (J09-J18)	8,038	2.9	---
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	7,688	2.8	---
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	5,495	2.0	---
9	Diabetes mellitus (E10-E14)	4,712	1.7	---
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	4,274	1.6	---
11	Septicemia (A40-A41)	3,364	1.2	---
12	Parkinson's disease (G20-G21)	3,359	1.2	---
13	Pneumonitis due to solids and liquids (J69)	3,209	1.2	---
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1,751	0.6	---
15	Atherosclerosis (I70)	1,260	0.5	---
...	All other causes (residual)	64,945	23.7	---

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2014

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2014

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
White, both sexes, 95-99 years				
...	All causes	107,768	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	35,202	32.7	---
2	Alzheimer's disease (G30)	9,562	8.9	---
3	Cerebrovascular diseases (I60-I69)	7,362	6.8	---
4	Malignant neoplasms (C00-C97)	7,170	6.7	---
5	Chronic lower respiratory diseases (J40-J47)	3,897	3.6	---
6	Influenza and pneumonia (J09-J18)	3,603	3.3	---
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,806	2.6	---
8	Essential hypertension and hypertensive renal disease (I10,I12,I15)	2,052	1.9	---
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2,029	1.9	---
10	Diabetes mellitus (E10-E14)	1,449	1.3	---
11	Pneumonitis due to solids and liquids (J69)	1,302	1.2	---
12	Septicemia (A40-A41)	1,082	1.0	---
13	Parkinson's disease (G20-G21)	796	0.7	---
14	Atherosclerosis (I70)	656	0.6	---
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	563	0.5	---
...	All other causes (residual)	28,237	26.2	---
White, both sexes, 100 years and over				
...	All causes	22,679	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	7,987	35.2	---
2	Alzheimer's disease (G30)	1,909	8.4	---
3	Cerebrovascular diseases (I60-I69)	1,367	6.0	---
4	Influenza and pneumonia (J09-J18)	909	4.0	---
5	Malignant neoplasms (C00-C97)	891	3.9	---
6	Chronic lower respiratory diseases (J40-J47)	589	2.6	---
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	538	2.4	---
8	Essential hypertension and hypertensive renal disease (I10,I12,I15)	535	2.4	---
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	356	1.6	---
10	Pneumonitis due to solids and liquids (J69)	262	1.2	---
11	Diabetes mellitus (E10-E14)	213	0.9	---
12	Atherosclerosis (I70)	189	0.8	---
13	Septicemia (A40-A41)	186	0.8	---
14	Parkinson's disease (G20-G21)	99	0.4	---
15	Nutritional deficiencies (E40-E64)	73	0.3	---
...	All other causes (residual)	6,576	29.0	---

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2014

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2014

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
White, male, all ages				
...	All causes	1,128,993	100.0	909.4
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	277,921	24.6	223.9
2	Malignant neoplasms (C00-C97)	266,137	23.6	214.4
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	72,884	6.5	58.7
4	Chronic lower respiratory diseases (J40-J47)	62,989	5.6	50.7
5	Cerebrovascular diseases (I60-I69)	45,505	4.0	36.7
6	Diabetes mellitus (E10-E14)	32,920	2.9	26.5
7	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	29,971	2.7	24.1
8	Alzheimer's disease (G30)	25,937	2.3	20.9
9	Influenza and pneumonia (J09-J18)	22,643	2.0	18.2
10	Chronic liver disease and cirrhosis (K70,K73-K74)	21,781	1.9	17.5
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	19,587	1.7	15.8
12	Septicemia (A40-A41)	14,847	1.3	12.0
13	Parkinson's disease (G20-G21)	14,594	1.3	11.8
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	9,998	0.9	8.1
15	Pneumonitis due to solids and liquids (J69)	9,119	0.8	7.3
...	All other causes (residual)	202,160	17.9	162.8
White, male, 1-4 years				
...	All causes	1,452	100.0	23.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	521	35.9	8.6
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	146	10.1	2.4
3	Malignant neoplasms (C00-C97)	142	9.8	2.3
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	105	7.2	1.7
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	42	2.9	0.7
6	Influenza and pneumonia (J09-J18)	37	2.5	0.6
7	Septicemia (A40-A41)	15	1.0	*
8	Chronic lower respiratory diseases (J40-J47)	14	1.0	*
8	Certain conditions originating in the perinatal period (P00-P96)	14	1.0	*
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	12	0.8	*
10	Cerebrovascular diseases (I60-I69)	12	0.8	*
12	Anemias (D50-D64)	4	0.3	*
12	Meningitis (G00,G03)	4	0.3	*
12	Complications of medical and surgical care (Y40-Y84,Y88)	4	0.3	*
15	Diabetes mellitus (E10-E14)	3	0.2	*
15	Nutritional deficiencies (E40-E64)	3	0.2	*
15	Acute bronchitis and bronchiolitis (J20-J21)	3	0.2	*
...	All other causes (residual)	371	25.6	6.1

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2014

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2014

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
White, male, 5-9 years				
...	All causes	965	100.0	12.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	338	35.0	4.3
2	Malignant neoplasms (C00-C97)	196	20.3	2.5
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	63	6.5	0.8
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	32	3.3	0.4
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	22	2.3	0.3
6	Influenza and pneumonia (J09-J18)	19	2.0	*
7	Chronic lower respiratory diseases (J40-J47)	16	1.7	*
8	Cerebrovascular diseases (I60-I69)	14	1.5	*
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	13	1.3	*
10	Septicemia (A40-A41)	10	1.0	*
11	Certain conditions originating in the perinatal period (P00-P96)	7	0.7	*
12	Diabetes mellitus (E10-E14)	3	0.3	*
12	Meningitis (G00,G03)	3	0.3	*
14	Anemias (D50-D64)	2	0.2	*
14	Nutritional deficiencies (E40-E64)	2	0.2	*
14	Pneumonitis due to solids and liquids (J69)	2	0.2	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2	0.2	*
14	Complications of medical and surgical care (Y40-Y84,Y88)	2	0.2	*
...	All other causes (residual)	219	22.7	2.8
White, male, 10-14 years				
...	All causes	1,266	100.0	15.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	357	28.2	4.4
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	224	17.7	2.8
3	Malignant neoplasms (C00-C97)	191	15.1	2.4
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	70	5.5	0.9
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	56	4.4	0.7
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	37	2.9	0.5
7	Cerebrovascular diseases (I60-I69)	22	1.7	0.3
8	Chronic lower respiratory diseases (J40-J47)	20	1.6	0.2
9	Influenza and pneumonia (J09-J18)	17	1.3	*
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	13	1.0	*
11	Septicemia (A40-A41)	10	0.8	*
12	Diabetes mellitus (E10-E14)	7	0.6	*
13	Anemias (D50-D64)	6	0.5	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	5	0.4	*
15	Whooping cough (A37)	2	0.2	*
15	Meningitis (G00,G03)	2	0.2	*
15	Diseases of appendix (K35-K38)	2	0.2	*
15	Hernia (K40-K46)	2	0.2	*
15	Certain conditions originating in the perinatal period (P00-P96)	2	0.2	*
15	Complications of medical and surgical care (Y40-Y84,Y88)	2	0.2	*
...	All other causes (residual)	219	17.3	2.7

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2014

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2014

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
White, male, 15-19 years				
...	All causes	4,867	100.0	59.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,172	44.6	26.5
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,209	24.8	14.7
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	368	7.6	4.5
4	Malignant neoplasms (C00-C97)	276	5.7	3.4
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	124	2.5	1.5
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	76	1.6	0.9
7	Influenza and pneumonia (J09-J18)	29	0.6	0.4
8	Cerebrovascular diseases (I60-I69)	19	0.4	*
8	Chronic lower respiratory diseases (J40-J47)	19	0.4	*
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	17	0.3	*
10	Diabetes mellitus (E10-E14)	17	0.3	*
12	Septicemia (A40-A41)	16	0.3	*
13	Legal intervention (Y35,Y89.0)	9	0.2	*
14	Aortic aneurysm and dissection (I71)	7	0.1	*
15	Complications of medical and surgical care (Y40-Y84,Y88)	6	0.1	*
...	All other causes (residual)	503	10.3	6.1
White, male, 20-24 years				
...	All causes	10,174	100.0	115.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	5,098	50.1	57.8
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,164	21.3	24.5
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	720	7.1	8.2
4	Malignant neoplasms (C00-C97)	458	4.5	5.2
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	284	2.8	3.2
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	86	0.8	1.0
7	Diabetes mellitus (E10-E14)	62	0.6	0.7
8	Influenza and pneumonia (J09-J18)	55	0.5	0.6
9	Cerebrovascular diseases (I60-I69)	49	0.5	0.6
10	Chronic lower respiratory diseases (J40-J47)	42	0.4	0.5
11	Septicemia (A40-A41)	35	0.3	0.4
12	Legal intervention (Y35,Y89.0)	34	0.3	0.4
13	Human immunodeficiency virus (HIV) disease (B20-B24)	24	0.2	0.3
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	18	0.2	*
15	Chronic liver disease and cirrhosis (K70,K73-K74)	15	0.1	*
...	All other causes (residual)	1,030	10.1	11.7

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2014

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2014

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
White, male, 25-29 years				
...	All causes	11,585	100.0	135.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	5,453	47.1	63.9
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,222	19.2	26.0
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	672	5.8	7.9
4	Malignant neoplasms (C00-C97)	577	5.0	6.8
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	522	4.5	6.1
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	119	1.0	1.4
7	Influenza and pneumonia (J09-J18)	105	0.9	1.2
8	Diabetes mellitus (E10-E14)	97	0.8	1.1
9	Cerebrovascular diseases (I60-I69)	95	0.8	1.1
10	Chronic liver disease and cirrhosis (K70,K73-K74)	91	0.8	1.1
11	Chronic lower respiratory diseases (J40-J47)	63	0.5	0.7
11	Legal intervention (Y35,Y89.0)	63	0.5	0.7
13	Septicemia (A40-A41)	56	0.5	0.7
14	Human immunodeficiency virus (HIV) disease (B20-B24)	42	0.4	0.5
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	27	0.2	0.3
...	All other causes (residual)	1,381	11.9	16.2
White, male, 30-34 years				
...	All causes	12,987	100.0	155.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	5,376	41.4	64.2
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,253	17.3	26.9
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	996	7.7	11.9
4	Malignant neoplasms (C00-C97)	848	6.5	10.1
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	590	4.5	7.0
6	Chronic liver disease and cirrhosis (K70,K73-K74)	274	2.1	3.3
7	Diabetes mellitus (E10-E14)	180	1.4	2.2
8	Cerebrovascular diseases (I60-I69)	123	0.9	1.5
8	Influenza and pneumonia (J09-J18)	123	0.9	1.5
10	Human immunodeficiency virus (HIV) disease (B20-B24)	106	0.8	1.3
11	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	92	0.7	1.1
12	Septicemia (A40-A41)	76	0.6	0.9
13	Chronic lower respiratory diseases (J40-J47)	73	0.6	0.9
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	53	0.4	0.6
15	Legal intervention (Y35,Y89.0)	43	0.3	0.5
...	All other causes (residual)	1,781	13.7	21.3

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2014

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2014

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
White, male, 35-39 years				
...	All causes	14,055	100.0	181.9
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	4,684	33.3	60.6
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,166	15.4	28.0
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,635	11.6	21.2
4	Malignant neoplasms (C00-C97)	1,269	9.0	16.4
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	493	3.5	6.4
6	Chronic liver disease and cirrhosis (K70,K73-K74)	427	3.0	5.5
7	Diabetes mellitus (E10-E14)	309	2.2	4.0
8	Cerebrovascular diseases (I60-I69)	225	1.6	2.9
9	Influenza and pneumonia (J09-J18)	182	1.3	2.4
10	Human immunodeficiency virus (HIV) disease (B20-B24)	141	1.0	1.8
11	Septicemia (A40-A41)	108	0.8	1.4
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	87	0.6	1.1
13	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	86	0.6	1.1
14	Chronic lower respiratory diseases (J40-J47)	76	0.5	1.0
15	Aortic aneurysm and dissection (I71)	59	0.4	0.8
...	All other causes (residual)	2,108	15.0	27.3
White, male, 40-44 years				
...	All causes	19,427	100.0	242.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	4,672	24.0	58.3
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,321	17.1	41.4
3	Malignant neoplasms (C00-C97)	2,518	13.0	31.4
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,320	11.9	28.9
5	Chronic liver disease and cirrhosis (K70,K73-K74)	975	5.0	12.2
6	Diabetes mellitus (E10-E14)	543	2.8	6.8
7	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	443	2.3	5.5
8	Cerebrovascular diseases (I60-I69)	408	2.1	5.1
9	Influenza and pneumonia (J09-J18)	297	1.5	3.7
10	Human immunodeficiency virus (HIV) disease (B20-B24)	249	1.3	3.1
11	Septicemia (A40-A41)	199	1.0	2.5
12	Chronic lower respiratory diseases (J40-J47)	164	0.8	2.0
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	145	0.7	1.8
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	107	0.6	1.3
15	Aortic aneurysm and dissection (I71)	100	0.5	1.2
...	All other causes (residual)	2,966	15.3	37.0

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2014

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2014

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
White, male, 45-49 years				
...	All causes	30,768	100.0	372.4
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	6,344	20.6	76.8
2	Malignant neoplasms (C00-C97)	5,354	17.4	64.8
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	5,238	17.0	63.4
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,813	9.1	34.0
5	Chronic liver disease and cirrhosis (K70,K73-K74)	1,797	5.8	21.7
6	Diabetes mellitus (E10-E14)	1,016	3.3	12.3
7	Cerebrovascular diseases (I60-I69)	732	2.4	8.9
8	Influenza and pneumonia (J09-J18)	453	1.5	5.5
9	Chronic lower respiratory diseases (J40-J47)	420	1.4	5.1
9	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	420	1.4	5.1
11	Human immunodeficiency virus (HIV) disease (B20-B24)	367	1.2	4.4
12	Septicemia (A40-A41)	333	1.1	4.0
13	Viral hepatitis (B15-B19)	310	1.0	3.8
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	257	0.8	3.1
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	187	0.6	2.3
...	All other causes (residual)	4,727	15.4	57.2
White, male, 50-54 years				
...	All causes	53,646	100.0	595.9
1	Malignant neoplasms (C00-C97)	12,308	22.9	136.7
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	12,253	22.8	136.1
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	6,431	12.0	71.4
4	Chronic liver disease and cirrhosis (K70,K73-K74)	3,292	6.1	36.6
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	3,128	5.8	34.7
6	Diabetes mellitus (E10-E14)	1,795	3.3	19.9
7	Cerebrovascular diseases (I60-I69)	1,314	2.4	14.6
8	Chronic lower respiratory diseases (J40-J47)	1,260	2.3	14.0
9	Viral hepatitis (B15-B19)	727	1.4	8.1
10	Influenza and pneumonia (J09-J18)	714	1.3	7.9
11	Septicemia (A40-A41)	666	1.2	7.4
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	551	1.0	6.1
13	Human immunodeficiency virus (HIV) disease (B20-B24)	539	1.0	6.0
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	399	0.7	4.4
15	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	393	0.7	4.4
...	All other causes (residual)	7,876	14.7	87.5

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2014

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2014

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
White, male, 55-59 years				
...	All causes	76,704	100.0	891.9
1	Malignant neoplasms (C00-C97)	21,947	28.6	255.2
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	18,240	23.8	212.1
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	5,678	7.4	66.0
4	Chronic liver disease and cirrhosis (K70,K73-K74)	4,015	5.2	46.7
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	3,004	3.9	34.9
6	Chronic lower respiratory diseases (J40-J47)	2,847	3.7	33.1
7	Diabetes mellitus (E10-E14)	2,663	3.5	31.0
8	Cerebrovascular diseases (I60-I69)	2,013	2.6	23.4
9	Viral hepatitis (B15-B19)	1,111	1.4	12.9
10	Influenza and pneumonia (J09-J18)	1,090	1.4	12.7
11	Septicemia (A40-A41)	958	1.2	11.1
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	788	1.0	9.2
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	601	0.8	7.0
14	Human immunodeficiency virus (HIV) disease (B20-B24)	387	0.5	4.5
15	Aortic aneurysm and dissection (I71)	338	0.4	3.9
...	All other causes (residual)	11,024	14.4	128.2
White, male, 60-64 years				
...	All causes	93,862	100.0	1,262.6
1	Malignant neoplasms (C00-C97)	29,928	31.9	402.6
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	22,796	24.3	306.7
3	Chronic lower respiratory diseases (J40-J47)	4,617	4.9	62.1
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	4,396	4.7	59.1
5	Chronic liver disease and cirrhosis (K70,K73-K74)	3,739	4.0	50.3
6	Diabetes mellitus (E10-E14)	3,566	3.8	48.0
7	Cerebrovascular diseases (I60-I69)	2,730	2.9	36.7
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,347	2.5	31.6
9	Influenza and pneumonia (J09-J18)	1,337	1.4	18.0
10	Septicemia (A40-A41)	1,289	1.4	17.3
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,142	1.2	15.4
12	Viral hepatitis (B15-B19)	1,104	1.2	14.9
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	817	0.9	11.0
14	Aortic aneurysm and dissection (I71)	480	0.5	6.5
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	473	0.5	6.4
...	All other causes (residual)	13,101	14.0	176.2

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2014

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2014

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
White, male, 65-69 years				
...	All causes	108,503	100.0	1,750.7
1	Malignant neoplasms (C00-C97)	36,447	33.6	588.1
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	26,236	24.2	423.3
3	Chronic lower respiratory diseases (J40-J47)	7,019	6.5	113.3
4	Diabetes mellitus (E10-E14)	4,280	3.9	69.1
5	Cerebrovascular diseases (I60-I69)	3,578	3.3	57.7
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,423	3.2	55.2
7	Chronic liver disease and cirrhosis (K70,K73-K74)	2,631	2.4	42.5
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,699	1.6	27.4
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,650	1.5	26.6
10	Influenza and pneumonia (J09-J18)	1,647	1.5	26.6
11	Septicemia (A40-A41)	1,590	1.5	25.7
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	909	0.8	14.7
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	701	0.6	11.3
14	Parkinson's disease (G20-G21)	612	0.6	9.9
15	Aortic aneurysm and dissection (I71)	585	0.5	9.4
...	All other causes (residual)	15,496	14.3	250.0
White, male, 70-74 years				
...	All causes	118,746	100.0	2,695.7
1	Malignant neoplasms (C00-C97)	38,581	32.5	875.9
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	28,128	23.7	638.6
3	Chronic lower respiratory diseases (J40-J47)	9,332	7.9	211.9
4	Cerebrovascular diseases (I60-I69)	4,621	3.9	104.9
5	Diabetes mellitus (E10-E14)	4,238	3.6	96.2
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,078	2.6	69.9
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2,006	1.7	45.5
8	Influenza and pneumonia (J09-J18)	1,941	1.6	44.1
9	Chronic liver disease and cirrhosis (K70,K73-K74)	1,816	1.5	41.2
10	Septicemia (A40-A41)	1,792	1.5	40.7
11	Parkinson's disease (G20-G21)	1,470	1.2	33.4
12	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,427	1.2	32.4
13	Alzheimer's disease (G30)	1,421	1.2	32.3
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	926	0.8	21.0
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	869	0.7	19.7
...	All other causes (residual)	17,100	14.4	388.2

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2014

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2014

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
White, male, 75-79 years				
...	All causes	129,804	100.0	4,257.6
1	Malignant neoplasms (C00-C97)	37,270	28.7	1,222.5
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	31,388	24.2	1,029.5
3	Chronic lower respiratory diseases (J40-J47)	10,286	7.9	337.4
4	Cerebrovascular diseases (I60-I69)	6,061	4.7	198.8
5	Diabetes mellitus (E10-E14)	4,115	3.2	135.0
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,370	2.6	110.5
7	Alzheimer's disease (G30)	3,033	2.3	99.5
8	Parkinson's disease (G20-G21)	2,685	2.1	88.1
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2,572	2.0	84.4
10	Influenza and pneumonia (J09-J18)	2,496	1.9	81.9
11	Septicemia (A40-A41)	1,843	1.4	60.5
12	Chronic liver disease and cirrhosis (K70,K73-K74)	1,258	1.0	41.3
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1,130	0.9	37.1
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,040	0.8	34.1
15	Pneumonitis due to solids and liquids (J69)	1,037	0.8	34.0
...	All other causes (residual)	20,220	15.6	663.2
White, male, 80-84 years				
...	All causes	149,373	100.0	7,103.0
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	38,754	25.9	1,842.8
2	Malignant neoplasms (C00-C97)	34,875	23.3	1,658.4
3	Chronic lower respiratory diseases (J40-J47)	10,936	7.3	520.0
4	Cerebrovascular diseases (I60-I69)	7,609	5.1	361.8
5	Alzheimer's disease (G30)	5,497	3.7	261.4
6	Diabetes mellitus (E10-E14)	4,173	2.8	198.4
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	4,039	2.7	192.1
8	Parkinson's disease (G20-G21)	3,771	2.5	179.3
9	Influenza and pneumonia (J09-J18)	3,351	2.2	159.3
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3,269	2.2	155.4
11	Septicemia (A40-A41)	2,093	1.4	99.5
12	Pneumonitis due to solids and liquids (J69)	1,531	1.0	72.8
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,403	0.9	66.7
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1,370	0.9	65.1
15	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	934	0.6	44.4
...	All other causes (residual)	25,768	17.3	1,225.3

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2014

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2014

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
White, male, 85-89 years				
...	All causes	150,111	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	43,182	28.8	---
2	Malignant neoplasms (C00-C97)	27,106	18.1	---
3	Chronic lower respiratory diseases (J40-J47)	9,347	6.2	---
4	Cerebrovascular diseases (I60-I69)	8,314	5.5	---
5	Alzheimer's disease (G30)	7,371	4.9	---
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	4,243	2.8	---
7	Influenza and pneumonia (J09-J18)	4,006	2.7	---
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3,611	2.4	---
9	Parkinson's disease (G20-G21)	3,602	2.4	---
10	Diabetes mellitus (E10-E14)	3,533	2.4	---
11	Septicemia (A40-A41)	2,082	1.4	---
12	Pneumonitis due to solids and liquids (J69)	1,978	1.3	---
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,693	1.1	---
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1,302	0.9	---
15	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	672	0.4	---
...	All other causes (residual)	28,069	18.7	---
White, male, 90-94 years				
...	All causes	98,836	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	31,746	32.1	---
2	Malignant neoplasms (C00-C97)	12,814	13.0	---
3	Alzheimer's disease (G30)	5,746	5.8	---
4	Cerebrovascular diseases (I60-I69)	5,701	5.8	---
5	Chronic lower respiratory diseases (J40-J47)	5,004	5.1	---
6	Influenza and pneumonia (J09-J18)	3,281	3.3	---
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,969	3.0	---
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2,503	2.5	---
9	Diabetes mellitus (E10-E14)	1,822	1.8	---
10	Parkinson's disease (G20-G21)	1,708	1.7	---
11	Pneumonitis due to solids and liquids (J69)	1,645	1.7	---
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,261	1.3	---
13	Septicemia (A40-A41)	1,224	1.2	---
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	774	0.8	---
15	Atherosclerosis (I70)	419	0.4	---
...	All other causes (residual)	20,219	20.5	---

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2014

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2014

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
White, male, 95-99 years				
...	All causes	29,475	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	10,239	34.7	---
2	Malignant neoplasms (C00-C97)	2,737	9.3	---
3	Alzheimer's disease (G30)	1,781	6.0	---
4	Cerebrovascular diseases (I60-I69)	1,623	5.5	---
5	Chronic lower respiratory diseases (J40-J47)	1,299	4.4	---
6	Influenza and pneumonia (J09-J18)	1,184	4.0	---
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	807	2.7	---
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	763	2.6	---
9	Pneumonitis due to solids and liquids (J69)	576	2.0	---
10	Diabetes mellitus (E10-E14)	464	1.6	---
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	458	1.6	---
12	Septicemia (A40-A41)	342	1.2	---
13	Parkinson's disease (G20-G21)	332	1.1	---
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	187	0.6	---
15	Atherosclerosis (I70)	172	0.6	---
...	All other causes (residual)	6,511	22.1	---
White, male, 100 years and over				
...	All causes	4,002	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,514	37.8	---
2	Malignant neoplasms (C00-C97)	264	6.6	---
3	Influenza and pneumonia (J09-J18)	222	5.5	---
4	Cerebrovascular diseases (I60-I69)	203	5.1	---
5	Alzheimer's disease (G30)	181	4.5	---
6	Chronic lower respiratory diseases (J40-J47)	122	3.0	---
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	120	3.0	---
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	100	2.5	---
9	Essential hypertension and hypertensive renal disease (I10,I12,I15)	97	2.4	---
10	Pneumonitis due to solids and liquids (J69)	84	2.1	---
11	Septicemia (A40-A41)	53	1.3	---
12	Atherosclerosis (I70)	34	0.8	---
13	Diabetes mellitus (E10-E14)	33	0.8	---
14	Parkinson's disease (G20-G21)	29	0.7	---
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	13	0.3	---
...	All other causes (residual)	933	23.3	---

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2014

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2014

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
White, female, all ages				
...	All causes	1,108,887	100.0	876.7
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	246,774	22.3	195.1
2	Malignant neoplasms (C00-C97)	236,795	21.4	187.2
3	Chronic lower respiratory diseases (J40-J47)	71,552	6.5	56.6
4	Cerebrovascular diseases (I60-I69)	65,530	5.9	51.8
5	Alzheimer's disease (G30)	59,053	5.3	46.7
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	44,358	4.0	35.1
7	Diabetes mellitus (E10-E14)	26,821	2.4	21.2
8	Influenza and pneumonia (J09-J18)	24,650	2.2	19.5
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	18,389	1.7	14.5
10	Septicemia (A40-A41)	16,665	1.5	13.2
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	13,641	1.2	10.8
12	Chronic liver disease and cirrhosis (K70,K73-K74)	11,727	1.1	9.3
13	Parkinson's disease (G20-G21)	9,700	0.9	7.7
14	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	8,704	0.8	6.9
15	Pneumonitis due to solids and liquids (J69)	7,368	0.7	5.8
...	All other causes (residual)	247,160	22.3	195.4
White, female, 1-4 years				
...	All causes	1,140	100.0	19.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	338	29.6	5.8
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	125	11.0	2.1
3	Malignant neoplasms (C00-C97)	108	9.5	1.9
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	95	8.3	1.6
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	42	3.7	0.7
6	Influenza and pneumonia (J09-J18)	35	3.1	0.6
7	Septicemia (A40-A41)	22	1.9	0.4
8	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	16	1.4	*
9	Chronic lower respiratory diseases (J40-J47)	11	1.0	*
10	Certain conditions originating in the perinatal period (P00-P96)	10	0.9	*
11	Meningitis (G00,G03)	8	0.7	*
11	Cerebrovascular diseases (I60-I69)	8	0.7	*
13	Acute bronchitis and bronchiolitis (J20-J21)	6	0.5	*
13	Complications of medical and surgical care (Y40-Y84,Y88)	6	0.5	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	5	0.4	*
...	All other causes (residual)	305	26.8	5.2

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2014

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2014

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
White, female, 5-9 years				
...	All causes	718	100.0	9.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	196	27.3	2.6
2	Malignant neoplasms (C00-C97)	148	20.6	2.0
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	78	10.9	1.0
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	41	5.7	0.5
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	23	3.2	0.3
6	Influenza and pneumonia (J09-J18)	22	3.1	0.3
7	Cerebrovascular diseases (I60-I69)	18	2.5	*
8	Septicemia (A40-A41)	12	1.7	*
8	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	12	1.7	*
10	Chronic lower respiratory diseases (J40-J47)	11	1.5	*
11	Certain conditions originating in the perinatal period (P00-P96)	4	0.6	*
12	Tuberculosis (A16-A19)	2	0.3	*
12	Anemias (D50-D64)	2	0.3	*
14	Diabetes mellitus (E10-E14)	1	0.1	*
14	Pneumonitis due to solids and liquids (J69)	1	0.1	*
14	Hernia (K40-K46)	1	0.1	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1	0.1	*
14	Complications of medical and surgical care (Y40-Y84,Y88)	1	0.1	*
...	All other causes (residual)	144	20.1	1.9
White, female, 10-14 years				
...	All causes	804	100.0	10.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	176	21.9	2.3
2	Malignant neoplasms (C00-C97)	131	16.3	1.7
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	123	15.3	1.6
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	43	5.3	0.6
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	34	4.2	0.4
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	31	3.9	0.4
7	Cerebrovascular diseases (I60-I69)	15	1.9	*
7	Influenza and pneumonia (J09-J18)	15	1.9	*
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	12	1.5	*
9	Chronic lower respiratory diseases (J40-J47)	12	1.5	*
11	Septicemia (A40-A41)	9	1.1	*
12	Certain conditions originating in the perinatal period (P00-P96)	8	1.0	*
13	Anemias (D50-D64)	3	0.4	*
13	Diabetes mellitus (E10-E14)	3	0.4	*
15	Meningitis (G00,G03)	2	0.2	*
15	Pneumonitis due to solids and liquids (J69)	2	0.2	*
15	Pregnancy, childbirth and the puerperium (O00-O99)	2	0.2	*
15	Complications of medical and surgical care (Y40-Y84,Y88)	2	0.2	*
...	All other causes (residual)	181	22.5	2.4

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2014

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2014

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
White, female, 15-19 years				
...	All causes	2,087	100.0	26.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	839	40.2	10.8
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	343	16.4	4.4
3	Malignant neoplasms (C00-C97)	200	9.6	2.6
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	94	4.5	1.2
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	77	3.7	1.0
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	69	3.3	0.9
7	Influenza and pneumonia (J09-J18)	24	1.1	0.3
8	Cerebrovascular diseases (I60-I69)	23	1.1	0.3
9	Pregnancy, childbirth and the puerperium (O00-O99)	20	1.0	0.3
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	13	0.6	*
11	Septicemia (A40-A41)	11	0.5	*
11	Chronic lower respiratory diseases (J40-J47)	11	0.5	*
13	Diabetes mellitus (E10-E14)	10	0.5	*
14	Anemias (D50-D64)	7	0.3	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	6	0.3	*
...	All other causes (residual)	340	16.3	4.4
White, female, 20-24 years				
...	All causes	3,632	100.0	43.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,650	45.4	19.8
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	461	12.7	5.5
3	Malignant neoplasms (C00-C97)	275	7.6	3.3
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	174	4.8	2.1
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	136	3.7	1.6
6	Pregnancy, childbirth and the puerperium (O00-O99)	79	2.2	0.9
7	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	57	1.6	0.7
8	Influenza and pneumonia (J09-J18)	46	1.3	0.6
9	Diabetes mellitus (E10-E14)	29	0.8	0.3
9	Cerebrovascular diseases (I60-I69)	29	0.8	0.3
11	Septicemia (A40-A41)	27	0.7	0.3
12	Chronic lower respiratory diseases (J40-J47)	23	0.6	0.3
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	16	0.4	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	14	0.4	*
15	Pneumonitis due to solids and liquids (J69)	10	0.3	*
...	All other causes (residual)	606	16.7	7.3

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2014

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2014

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
White, female, 25-29 years				
...	All causes	4,762	100.0	58.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,915	40.2	23.6
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	564	11.8	6.9
3	Malignant neoplasms (C00-C97)	489	10.3	6.0
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	232	4.9	2.9
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	187	3.9	2.3
6	Pregnancy, childbirth and the puerperium (O00-O99)	107	2.2	1.3
7	Influenza and pneumonia (J09-J18)	71	1.5	0.9
8	Chronic liver disease and cirrhosis (K70,K73-K74)	66	1.4	0.8
9	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	59	1.2	0.7
10	Cerebrovascular diseases (I60-I69)	57	1.2	0.7
11	Diabetes mellitus (E10-E14)	53	1.1	0.7
12	Septicemia (A40-A41)	44	0.9	0.5
13	Chronic lower respiratory diseases (J40-J47)	41	0.9	0.5
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	20	0.4	0.2
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	20	0.4	0.2
...	All other causes (residual)	837	17.6	10.3
White, female, 30-34 years				
...	All causes	6,074	100.0	75.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,918	31.6	23.7
2	Malignant neoplasms (C00-C97)	857	14.1	10.6
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	595	9.8	7.4
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	427	7.0	5.3
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	218	3.6	2.7
6	Chronic liver disease and cirrhosis (K70,K73-K74)	156	2.6	1.9
7	Pregnancy, childbirth and the puerperium (O00-O99)	133	2.2	1.6
8	Influenza and pneumonia (J09-J18)	125	2.1	1.5
9	Cerebrovascular diseases (I60-I69)	112	1.8	1.4
10	Diabetes mellitus (E10-E14)	105	1.7	1.3
11	Septicemia (A40-A41)	82	1.4	1.0
12	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	79	1.3	1.0
13	Chronic lower respiratory diseases (J40-J47)	53	0.9	0.7
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	40	0.7	0.5
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	29	0.5	0.4
...	All other causes (residual)	1,145	18.9	14.2

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2014

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2014

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
White, female, 35-39 years				
...	All causes	7,929	100.0	105.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,994	25.1	26.5
2	Malignant neoplasms (C00-C97)	1,655	20.9	22.0
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	714	9.0	9.5
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	655	8.3	8.7
5	Chronic liver disease and cirrhosis (K70,K73-K74)	258	3.3	3.4
6	Diabetes mellitus (E10-E14)	182	2.3	2.4
7	Cerebrovascular diseases (I60-I69)	177	2.2	2.4
8	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	156	2.0	2.1
9	Influenza and pneumonia (J09-J18)	143	1.8	1.9
10	Pregnancy, childbirth and the puerperium (O00-O99)	133	1.7	1.8
11	Septicemia (A40-A41)	121	1.5	1.6
12	Chronic lower respiratory diseases (J40-J47)	88	1.1	1.2
13	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	62	0.8	0.8
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	54	0.7	0.7
15	Human immunodeficiency virus (HIV) disease (B20-B24)	43	0.5	0.6
...	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	43	0.5	0.6
...	All other causes (residual)	1,451	18.3	19.3
White, female, 40-44 years				
...	All causes	12,263	100.0	155.0
1	Malignant neoplasms (C00-C97)	3,168	25.8	40.0
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,236	18.2	28.3
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,465	11.9	18.5
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	830	6.8	10.5
5	Chronic liver disease and cirrhosis (K70,K73-K74)	512	4.2	6.5
6	Diabetes mellitus (E10-E14)	321	2.6	4.1
7	Cerebrovascular diseases (I60-I69)	319	2.6	4.0
8	Influenza and pneumonia (J09-J18)	241	2.0	3.0
9	Chronic lower respiratory diseases (J40-J47)	206	1.7	2.6
10	Septicemia (A40-A41)	177	1.4	2.2
10	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	177	1.4	2.2
12	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	97	0.8	1.2
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	90	0.7	1.1
14	Pregnancy, childbirth and the puerperium (O00-O99)	86	0.7	1.1
15	Human immunodeficiency virus (HIV) disease (B20-B24)	78	0.6	1.0
...	All other causes (residual)	2,260	18.4	28.6

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2014

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2014

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
White, female, 45-49 years				
...	All causes	19,466	100.0	237.0
1	Malignant neoplasms (C00-C97)	5,974	30.7	72.7
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,598	13.3	31.6
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,501	12.8	30.4
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	989	5.1	12.0
5	Chronic liver disease and cirrhosis (K70,K73-K74)	911	4.7	11.1
6	Cerebrovascular diseases (I60-I69)	551	2.8	6.7
7	Diabetes mellitus (E10-E14)	546	2.8	6.6
8	Chronic lower respiratory diseases (J40-J47)	493	2.5	6.0
9	Influenza and pneumonia (J09-J18)	371	1.9	4.5
10	Septicemia (A40-A41)	296	1.5	3.6
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	186	1.0	2.3
12	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	163	0.8	2.0
13	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	143	0.7	1.7
14	Viral hepatitis (B15-B19)	127	0.7	1.5
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	98	0.5	1.2
...	All other causes (residual)	3,519	18.1	42.8
White, female, 50-54 years				
...	All causes	33,617	100.0	367.5
1	Malignant neoplasms (C00-C97)	11,705	34.8	128.0
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	4,824	14.3	52.7
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,176	9.4	34.7
4	Chronic liver disease and cirrhosis (K70,K73-K74)	1,533	4.6	16.8
5	Chronic lower respiratory diseases (J40-J47)	1,502	4.5	16.4
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,187	3.5	13.0
7	Diabetes mellitus (E10-E14)	1,009	3.0	11.0
8	Cerebrovascular diseases (I60-I69)	976	2.9	10.7
9	Influenza and pneumonia (J09-J18)	588	1.7	6.4
10	Septicemia (A40-A41)	548	1.6	6.0
11	Viral hepatitis (B15-B19)	348	1.0	3.8
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	291	0.9	3.2
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	202	0.6	2.2
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	171	0.5	1.9
15	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	148	0.4	1.6
...	All other causes (residual)	5,409	16.1	59.1

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2014

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2014

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
White, female, 55-59 years				
...	All causes	47,464	100.0	531.8
1	Malignant neoplasms (C00-C97)	18,023	38.0	201.9
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	7,085	14.9	79.4
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,849	6.0	31.9
4	Chronic lower respiratory diseases (J40-J47)	2,834	6.0	31.8
5	Chronic liver disease and cirrhosis (K70,K73-K74)	1,786	3.8	20.0
6	Diabetes mellitus (E10-E14)	1,503	3.2	16.8
7	Cerebrovascular diseases (I60-I69)	1,458	3.1	16.3
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,047	2.2	11.7
9	Septicemia (A40-A41)	924	1.9	10.4
10	Influenza and pneumonia (J09-J18)	810	1.7	9.1
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	546	1.2	6.1
12	Viral hepatitis (B15-B19)	471	1.0	5.3
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	327	0.7	3.7
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	227	0.5	2.5
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	203	0.4	2.3
...	All other causes (residual)	7,371	15.5	82.6
White, female, 60-64 years				
...	All causes	59,550	100.0	751.5
1	Malignant neoplasms (C00-C97)	23,233	39.0	293.2
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	9,569	16.1	120.8
3	Chronic lower respiratory diseases (J40-J47)	4,102	6.9	51.8
4	Diabetes mellitus (E10-E14)	2,118	3.6	26.7
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,089	3.5	26.4
6	Cerebrovascular diseases (I60-I69)	1,975	3.3	24.9
7	Chronic liver disease and cirrhosis (K70,K73-K74)	1,531	2.6	19.3
8	Septicemia (A40-A41)	1,066	1.8	13.5
9	Influenza and pneumonia (J09-J18)	1,016	1.7	12.8
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	828	1.4	10.4
11	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	699	1.2	8.8
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	476	0.8	6.0
13	Viral hepatitis (B15-B19)	436	0.7	5.5
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	334	0.6	4.2
15	Alzheimer's disease (G30)	311	0.5	3.9
...	All other causes (residual)	9,767	16.4	123.3

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2014

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2014

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
White, female, 65-69 years				
...	All causes	76,540	100.0	1,133.7
1	Malignant neoplasms (C00-C97)	29,094	38.0	430.9
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	12,825	16.8	190.0
3	Chronic lower respiratory diseases (J40-J47)	6,617	8.6	98.0
4	Cerebrovascular diseases (I60-I69)	2,940	3.8	43.5
5	Diabetes mellitus (E10-E14)	2,798	3.7	41.4
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,817	2.4	26.9
7	Septicemia (A40-A41)	1,447	1.9	21.4
8	Chronic liver disease and cirrhosis (K70,K73-K74)	1,344	1.8	19.9
9	Influenza and pneumonia (J09-J18)	1,303	1.7	19.3
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,300	1.7	19.3
11	Alzheimer's disease (G30)	723	0.9	10.7
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	606	0.8	9.0
13	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	486	0.6	7.2
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	409	0.5	6.1
15	Pneumonitis due to solids and liquids (J69)	373	0.5	5.5
...	All other causes (residual)	12,458	16.3	184.5
White, female, 70-74 years				
...	All causes	92,639	100.0	1,833.1
1	Malignant neoplasms (C00-C97)	31,398	33.9	621.3
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	16,402	17.7	324.6
3	Chronic lower respiratory diseases (J40-J47)	9,559	10.3	189.2
4	Cerebrovascular diseases (I60-I69)	4,165	4.5	82.4
5	Diabetes mellitus (E10-E14)	2,910	3.1	57.6
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,007	2.2	39.7
7	Alzheimer's disease (G30)	1,895	2.0	37.5
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,707	1.8	33.8
9	Septicemia (A40-A41)	1,685	1.8	33.3
10	Influenza and pneumonia (J09-J18)	1,637	1.8	32.4
11	Chronic liver disease and cirrhosis (K70,K73-K74)	1,172	1.3	23.2
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	817	0.9	16.2
13	Parkinson's disease (G20-G21)	741	0.8	14.7
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	607	0.7	12.0
15	Pneumonitis due to solids and liquids (J69)	464	0.5	9.2
...	All other causes (residual)	15,473	16.7	306.2

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2014

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2014

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
White, female, 75-79 years				
...	All causes	113,430	100.0	3,024.0
1	Malignant neoplasms (C00-C97)	31,589	27.8	842.1
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	22,081	19.5	588.7
3	Chronic lower respiratory diseases (J40-J47)	10,877	9.6	290.0
4	Cerebrovascular diseases (I60-I69)	6,567	5.8	175.1
5	Alzheimer's disease (G30)	4,549	4.0	121.3
6	Diabetes mellitus (E10-E14)	3,345	2.9	89.2
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,554	2.3	68.1
8	Influenza and pneumonia (J09-J18)	2,210	1.9	58.9
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2,162	1.9	57.6
10	Septicemia (A40-A41)	2,007	1.8	53.5
11	Parkinson's disease (G20-G21)	1,326	1.2	35.4
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,165	1.0	31.1
13	Chronic liver disease and cirrhosis (K70,K73-K74)	991	0.9	26.4
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	777	0.7	20.7
15	Pneumonitis due to solids and liquids (J69)	690	0.6	18.4
...	All other causes (residual)	20,540	18.1	547.6
White, female, 80-84 years				
...	All causes	153,341	100.0	5,258.6
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	34,521	22.5	1,183.8
2	Malignant neoplasms (C00-C97)	31,294	20.4	1,073.2
3	Chronic lower respiratory diseases (J40-J47)	12,264	8.0	420.6
4	Cerebrovascular diseases (I60-I69)	10,574	6.9	362.6
5	Alzheimer's disease (G30)	9,735	6.3	333.8
6	Diabetes mellitus (E10-E14)	3,892	2.5	133.5
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,593	2.3	123.2
8	Influenza and pneumonia (J09-J18)	3,310	2.2	113.5
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2,951	1.9	101.2
10	Septicemia (A40-A41)	2,412	1.6	82.7
11	Parkinson's disease (G20-G21)	2,281	1.5	78.2
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,917	1.3	65.7
13	Pneumonitis due to solids and liquids (J69)	1,118	0.7	38.3
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1,057	0.7	36.2
15	Chronic liver disease and cirrhosis (K70,K73-K74)	745	0.5	25.5
...	All other causes (residual)	31,677	20.7	1,086.3

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2014

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2014

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
White, female, 85-89 years				
...	All causes	194,413	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	50,663	26.1	---
2	Malignant neoplasms (C00-C97)	26,791	13.8	---
3	Alzheimer's disease (G30)	15,883	8.2	---
4	Cerebrovascular diseases (I60-I69)	14,861	7.6	---
5	Chronic lower respiratory diseases (J40-J47)	11,910	6.1	---
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	4,982	2.6	---
7	Influenza and pneumonia (J09-J18)	4,773	2.5	---
8	Diabetes mellitus (E10-E14)	3,939	2.0	---
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3,641	1.9	---
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	2,901	1.5	---
11	Septicemia (A40-A41)	2,723	1.4	---
12	Parkinson's disease (G20-G21)	2,653	1.4	---
13	Pneumonitis due to solids and liquids (J69)	1,561	0.8	---
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1,294	0.7	---
15	Enterocolitis due to <i>Clostridium difficile</i> (A04.7)	817	0.4	---
...	All other causes (residual)	45,021	23.2	---
White, female, 90-94 years				
...	All causes	175,415	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	51,625	29.4	---
2	Alzheimer's disease (G30)	16,294	9.3	---
3	Malignant neoplasms (C00-C97)	15,580	8.9	---
4	Cerebrovascular diseases (I60-I69)	13,774	7.9	---
5	Chronic lower respiratory diseases (J40-J47)	7,872	4.5	---
6	Influenza and pneumonia (J09-J18)	4,757	2.7	---
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	4,719	2.7	---
8	Essential hypertension and hypertensive renal disease (I10,I12,I15)	3,013	1.7	---
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2,992	1.7	---
10	Diabetes mellitus (E10-E14)	2,890	1.6	---
11	Septicemia (A40-A41)	2,140	1.2	---
12	Parkinson's disease (G20-G21)	1,651	0.9	---
13	Pneumonitis due to solids and liquids (J69)	1,564	0.9	---
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	977	0.6	---
15	Atherosclerosis (I70)	841	0.5	---
...	All other causes (residual)	44,726	25.5	---

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2014

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2014

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
White, female, 95-99 years				
...	All causes	78,293	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	24,963	31.9	---
2	Alzheimer's disease (G30)	7,781	9.9	---
3	Cerebrovascular diseases (I60-I69)	5,739	7.3	---
4	Malignant neoplasms (C00-C97)	4,433	5.7	---
5	Chronic lower respiratory diseases (J40-J47)	2,598	3.3	---
6	Influenza and pneumonia (J09-J18)	2,419	3.1	---
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,999	2.6	---
8	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,594	2.0	---
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,266	1.6	---
10	Diabetes mellitus (E10-E14)	985	1.3	---
11	Septicemia (A40-A41)	740	0.9	---
12	Pneumonitis due to solids and liquids (J69)	726	0.9	---
13	Atherosclerosis (I70)	484	0.6	---
14	Parkinson's disease (G20-G21)	464	0.6	---
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	376	0.5	---
...	All other causes (residual)	21,726	27.7	---
White, female, 100 years and over				
...	All causes	18,677	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	6,473	34.7	---
2	Alzheimer's disease (G30)	1,728	9.3	---
3	Cerebrovascular diseases (I60-I69)	1,164	6.2	---
4	Influenza and pneumonia (J09-J18)	687	3.7	---
5	Malignant neoplasms (C00-C97)	627	3.4	---
6	Chronic lower respiratory diseases (J40-J47)	467	2.5	---
7	Essential hypertension and hypertensive renal disease (I10,I12,I15)	438	2.3	---
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	418	2.2	---
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	256	1.4	---
10	Diabetes mellitus (E10-E14)	180	1.0	---
11	Pneumonitis due to solids and liquids (J69)	178	1.0	---
12	Atherosclerosis (I70)	155	0.8	---
13	Septicemia (A40-A41)	133	0.7	---
14	Parkinson's disease (G20-G21)	70	0.4	---
15	Nutritional deficiencies (E40-E64)	62	0.3	---
...	All other causes (residual)	5,641	30.2	---

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2014

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2014

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
Black, both sexes, all ages				
...	All causes	308,960	100.0	697.3
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	73,095	23.7	165.0
2	Malignant neoplasms (C00-C97)	69,090	22.4	155.9
3	Cerebrovascular diseases (I60-I69)	17,088	5.5	38.6
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	14,168	4.6	32.0
5	Diabetes mellitus (E10-E14)	13,435	4.3	30.3
6	Chronic lower respiratory diseases (J40-J47)	9,934	3.2	22.4
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	8,586	2.8	19.4
8	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	7,876	2.5	17.8
9	Alzheimer's disease (G30)	6,567	2.1	14.8
10	Septicemia (A40-A41)	6,386	2.1	14.4
11	Influenza and pneumonia (J09-J18)	5,611	1.8	12.7
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	5,399	1.7	12.2
13	Certain conditions originating in the perinatal period (P00-P96)	4,031	1.3	9.1
14	Human immunodeficiency virus (HIV) disease (B20-B24)	3,591	1.2	8.1
15	Chronic liver disease and cirrhosis (K70,K73-K74)	3,103	1.0	7.0
...	All other causes (residual)	61,000	19.7	137.7
Black, both sexes, 1-4 years				
...	All causes	1,009	100.0	37.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	291	28.8	10.7
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	151	15.0	5.6
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	104	10.3	3.8
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	55	5.5	2.0
5	Malignant neoplasms (C00-C97)	54	5.4	2.0
6	Influenza and pneumonia (J09-J18)	29	2.9	1.1
7	Chronic lower respiratory diseases (J40-J47)	23	2.3	0.8
8	Septicemia (A40-A41)	12	1.2	*
8	Certain conditions originating in the perinatal period (P00-P96)	12	1.2	*
10	Cerebrovascular diseases (I60-I69)	9	0.9	*
11	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	7	0.7	*
11	Acute bronchitis and bronchiolitis (J20-J21)	7	0.7	*
11	Complications of medical and surgical care (Y40-Y84,Y88)	7	0.7	*
14	Anemias (D50-D64)	6	0.6	*
15	Meningitis (G00,G03)	4	0.4	*
...	All other causes (residual)	238	23.6	8.8

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2014

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2014

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
Black, both sexes, 5-9 years				
...	All causes	537	100.0	15.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	154	28.7	4.5
2	Malignant neoplasms (C00-C97)	77	14.3	2.2
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	43	8.0	1.3
4	Chronic lower respiratory diseases (J40-J47)	38	7.1	1.1
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	38	7.1	1.1
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	19	3.5	*
7	Influenza and pneumonia (J09-J18)	12	2.2	*
8	Cerebrovascular diseases (I60-I69)	10	1.9	*
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	9	1.7	*
9	Anemias (D50-D64)	9	1.7	*
11	Septicemia (A40-A41)	8	1.5	*
12	Certain conditions originating in the perinatal period (P00-P96)	5	0.9	*
13	Meningitis (G00,G03)	3	0.6	*
14	Diabetes mellitus (E10-E14)	2	0.4	*
14	Acute bronchitis and bronchiolitis (J20-J21)	2	0.4	*
14	Hernia (K40-K46)	2	0.4	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2	0.4	*
14	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2	0.4	*
14	Complications of medical and surgical care (Y40-Y84,Y88)	2	0.4	*
...	All other causes (residual)	100	18.6	2.9
Black, both sexes, 10-14 years				
...	All causes	667	100.0	19.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	182	27.3	5.4
2	Malignant neoplasms (C00-C97)	69	10.3	2.0
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	62	9.3	1.8
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	56	8.4	1.7
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	42	6.3	1.2
6	Chronic lower respiratory diseases (J40-J47)	37	5.5	1.1
7	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	34	5.1	1.0
8	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	12	1.8	*
9	Anemias (D50-D64)	9	1.3	*
9	Diabetes mellitus (E10-E14)	9	1.3	*
11	Septicemia (A40-A41)	8	1.2	*
12	Cerebrovascular diseases (I60-I69)	5	0.7	*
12	Influenza and pneumonia (J09-J18)	5	0.7	*
12	Certain conditions originating in the perinatal period (P00-P96)	5	0.7	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3	0.4	*
15	Complications of medical and surgical care (Y40-Y84,Y88)	3	0.4	*
...	All other causes (residual)	126	18.9	3.7

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2014

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2014

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
Black, both sexes, 15-19 years				
...	All causes	2,165	100.0	62.3
1	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	901	41.6	25.9
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	548	25.3	15.8
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	157	7.3	4.5
4	Malignant neoplasms (C00-C97)	101	4.7	2.9
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	86	4.0	2.5
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	27	1.2	0.8
7	Chronic lower respiratory diseases (J40-J47)	23	1.1	0.7
8	Anemias (D50-D64)	17	0.8	*
9	Septicemia (A40-A41)	12	0.6	*
9	Cerebrovascular diseases (I60-I69)	12	0.6	*
11	Pregnancy, childbirth and the puerperium (O00-O99)	11	0.5	*
12	Influenza and pneumonia (J09-J18)	9	0.4	*
13	Legal intervention (Y35,Y89.0)	8	0.4	*
14	Human immunodeficiency virus (HIV) disease (B20-B24)	7	0.3	*
14	Diabetes mellitus (E10-E14)	7	0.3	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	7	0.3	*
...	All other causes (residual)	232	10.7	6.7
Black, both sexes, 20-24 years				
...	All causes	4,437	100.0	114.1
1	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,770	39.9	45.5
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	975	22.0	25.1
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	387	8.7	10.0
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	198	4.5	5.1
5	Malignant neoplasms (C00-C97)	171	3.9	4.4
6	Human immunodeficiency virus (HIV) disease (B20-B24)	79	1.8	2.0
7	Diabetes mellitus (E10-E14)	53	1.2	1.4
8	Chronic lower respiratory diseases (J40-J47)	49	1.1	1.3
9	Anemias (D50-D64)	48	1.1	1.2
10	Pregnancy, childbirth and the puerperium (O00-O99)	46	1.0	1.2
11	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	40	0.9	1.0
12	Cerebrovascular diseases (I60-I69)	37	0.8	1.0
13	Influenza and pneumonia (J09-J18)	29	0.7	0.7
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	22	0.5	0.6
15	Legal intervention (Y35,Y89.0)	21	0.5	0.5
...	All other causes (residual)	512	11.5	13.2

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2014

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2014

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
Black, both sexes, 25-29 years				
...	All causes	4,513	100.0	135.1
1	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,377	30.5	41.2
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	969	21.5	29.0
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	340	7.5	10.2
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	308	6.8	9.2
5	Malignant neoplasms (C00-C97)	220	4.9	6.6
6	Human immunodeficiency virus (HIV) disease (B20-B24)	162	3.6	4.8
7	Diabetes mellitus (E10-E14)	89	2.0	2.7
8	Pregnancy, childbirth and the puerperium (O00-O99)	62	1.4	1.9
9	Cerebrovascular diseases (I60-I69)	61	1.4	1.8
10	Chronic lower respiratory diseases (J40-J47)	60	1.3	1.8
11	Anemias (D50-D64)	52	1.2	1.6
12	Influenza and pneumonia (J09-J18)	43	1.0	1.3
13	Septicemia (A40-A41)	36	0.8	1.1
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	33	0.7	1.0
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	33	0.7	1.0
...	All other causes (residual)	668	14.8	20.0
Black, both sexes, 30-34 years				
...	All causes	5,036	100.0	163.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	996	19.8	32.3
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	983	19.5	31.8
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	670	13.3	21.7
4	Malignant neoplasms (C00-C97)	409	8.1	13.2
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	259	5.1	8.4
6	Human immunodeficiency virus (HIV) disease (B20-B24)	214	4.2	6.9
7	Diabetes mellitus (E10-E14)	150	3.0	4.9
8	Cerebrovascular diseases (I60-I69)	94	1.9	3.0
9	Pregnancy, childbirth and the puerperium (O00-O99)	81	1.6	2.6
10	Anemias (D50-D64)	75	1.5	2.4
11	Chronic lower respiratory diseases (J40-J47)	66	1.3	2.1
12	Influenza and pneumonia (J09-J18)	56	1.1	1.8
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	51	1.0	1.7
14	Septicemia (A40-A41)	45	0.9	1.5
15	Chronic liver disease and cirrhosis (K70,K73-K74)	35	0.7	1.1
...	All other causes (residual)	852	16.9	27.6

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2014

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2014

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
Black, both sexes, 35-39 years				
...	All causes	5,959	100.0	211.7
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,026	17.2	36.4
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	903	15.2	32.1
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	720	12.1	25.6
4	Malignant neoplasms (C00-C97)	697	11.7	24.8
5	Human immunodeficiency virus (HIV) disease (B20-B24)	275	4.6	9.8
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	234	3.9	8.3
7	Diabetes mellitus (E10-E14)	205	3.4	7.3
8	Cerebrovascular diseases (I60-I69)	179	3.0	6.4
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	101	1.7	3.6
10	Influenza and pneumonia (J09-J18)	84	1.4	3.0
11	Chronic lower respiratory diseases (J40-J47)	78	1.3	2.8
12	Septicemia (A40-A41)	74	1.2	2.6
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	73	1.2	2.6
14	Anemias (D50-D64)	66	1.1	2.3
15	Pregnancy, childbirth and the puerperium (O00-O99)	61	1.0	2.2
...	All other causes (residual)	1,183	19.9	42.0
Black, both sexes, 40-44 years				
...	All causes	8,038	100.0	284.0
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,749	21.8	61.8
2	Malignant neoplasms (C00-C97)	1,279	15.9	45.2
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	954	11.9	33.7
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	499	6.2	17.6
5	Human immunodeficiency virus (HIV) disease (B20-B24)	365	4.5	12.9
6	Diabetes mellitus (E10-E14)	342	4.3	12.1
7	Cerebrovascular diseases (I60-I69)	337	4.2	11.9
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	207	2.6	7.3
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	164	2.0	5.8
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	124	1.5	4.4
10	Chronic liver disease and cirrhosis (K70,K73-K74)	124	1.5	4.4
12	Influenza and pneumonia (J09-J18)	117	1.5	4.1
13	Septicemia (A40-A41)	115	1.4	4.1
14	Chronic lower respiratory diseases (J40-J47)	110	1.4	3.9
15	Anemias (D50-D64)	77	1.0	2.7
...	All other causes (residual)	1,475	18.4	52.1

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2014

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2014

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
Black, both sexes, 45-49 years				
...	All causes	12,101	100.0	431.4
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,875	23.8	102.5
2	Malignant neoplasms (C00-C97)	2,458	20.3	87.6
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,122	9.3	40.0
4	Diabetes mellitus (E10-E14)	570	4.7	20.3
5	Cerebrovascular diseases (I60-I69)	522	4.3	18.6
6	Human immunodeficiency virus (HIV) disease (B20-B24)	507	4.2	18.1
7	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	398	3.3	14.2
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	233	1.9	8.3
9	Chronic liver disease and cirrhosis (K70,K73-K74)	229	1.9	8.2
10	Septicemia (A40-A41)	207	1.7	7.4
10	Chronic lower respiratory diseases (J40-J47)	207	1.7	7.4
12	Influenza and pneumonia (J09-J18)	203	1.7	7.2
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	187	1.5	6.7
14	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	179	1.5	6.4
15	Aortic aneurysm and dissection (I71)	73	0.6	2.6
...	All other causes (residual)	2,131	17.6	76.0
Black, both sexes, 50-54 years				
...	All causes	19,714	100.0	678.6
1	Malignant neoplasms (C00-C97)	5,034	25.5	173.3
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	4,812	24.4	165.6
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,410	7.2	48.5
4	Cerebrovascular diseases (I60-I69)	951	4.8	32.7
5	Diabetes mellitus (E10-E14)	869	4.4	29.9
6	Human immunodeficiency virus (HIV) disease (B20-B24)	593	3.0	20.4
7	Chronic liver disease and cirrhosis (K70,K73-K74)	495	2.5	17.0
8	Chronic lower respiratory diseases (J40-J47)	444	2.3	15.3
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	427	2.2	14.7
10	Septicemia (A40-A41)	386	2.0	13.3
11	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	327	1.7	11.3
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	298	1.5	10.3
13	Influenza and pneumonia (J09-J18)	297	1.5	10.2
14	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	196	1.0	6.7
15	Viral hepatitis (B15-B19)	145	0.7	5.0
...	All other causes (residual)	3,030	15.4	104.3

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2014

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2014

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
Black, both sexes, 55-59 years				
...	All causes	27,864	100.0	1,054.3
1	Malignant neoplasms (C00-C97)	8,090	29.0	306.1
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	6,755	24.2	255.6
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,433	5.1	54.2
4	Cerebrovascular diseases (I60-I69)	1,331	4.8	50.4
5	Diabetes mellitus (E10-E14)	1,325	4.8	50.1
6	Chronic lower respiratory diseases (J40-J47)	813	2.9	30.8
7	Chronic liver disease and cirrhosis (K70,K73-K74)	683	2.5	25.8
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	656	2.4	24.8
9	Septicemia (A40-A41)	609	2.2	23.0
10	Human immunodeficiency virus (HIV) disease (B20-B24)	586	2.1	22.2
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	449	1.6	17.0
12	Influenza and pneumonia (J09-J18)	414	1.5	15.7
13	Viral hepatitis (B15-B19)	298	1.1	11.3
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	219	0.8	8.3
15	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	144	0.5	5.4
...	All other causes (residual)	4,059	14.6	153.6
Black, both sexes, 60-64 years				
...	All causes	32,108	100.0	1,531.1
1	Malignant neoplasms (C00-C97)	9,950	31.0	474.5
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	7,883	24.6	375.9
3	Cerebrovascular diseases (I60-I69)	1,632	5.1	77.8
4	Diabetes mellitus (E10-E14)	1,575	4.9	75.1
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,030	3.2	49.1
6	Chronic lower respiratory diseases (J40-J47)	1,019	3.2	48.6
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	927	2.9	44.2
8	Septicemia (A40-A41)	714	2.2	34.0
9	Chronic liver disease and cirrhosis (K70,K73-K74)	629	2.0	30.0
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	569	1.8	27.1
11	Influenza and pneumonia (J09-J18)	522	1.6	24.9
12	Human immunodeficiency virus (HIV) disease (B20-B24)	410	1.3	19.6
13	Viral hepatitis (B15-B19)	353	1.1	16.8
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	139	0.4	6.6
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	129	0.4	6.2
...	All other causes (residual)	4,627	14.4	220.6

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2014

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2014

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
Black, both sexes, 65-69 years				
...	All causes	30,985	100.0	2,031.8
1	Malignant neoplasms (C00-C97)	9,853	31.8	646.1
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	7,668	24.7	502.8
3	Cerebrovascular diseases (I60-I69)	1,675	5.4	109.8
4	Diabetes mellitus (E10-E14)	1,581	5.1	103.7
5	Chronic lower respiratory diseases (J40-J47)	1,237	4.0	81.1
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,008	3.3	66.1
7	Septicemia (A40-A41)	731	2.4	47.9
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	674	2.2	44.2
9	Essential hypertension and hypertensive renal disease (I10,I12,I15)	506	1.6	33.2
10	Influenza and pneumonia (J09-J18)	476	1.5	31.2
11	Chronic liver disease and cirrhosis (K70,K73-K74)	409	1.3	26.8
12	Human immunodeficiency virus (HIV) disease (B20-B24)	215	0.7	14.1
13	Viral hepatitis (B15-B19)	165	0.5	10.8
14	Pneumonitis due to solids and liquids (J69)	152	0.5	10.0
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	151	0.5	9.9
...	All other causes (residual)	4,484	14.5	294.0
Black, both sexes, 70-74 years				
...	All causes	29,970	100.0	2,897.0
1	Malignant neoplasms (C00-C97)	8,847	29.5	855.2
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	7,462	24.9	721.3
3	Cerebrovascular diseases (I60-I69)	1,836	6.1	177.5
4	Diabetes mellitus (E10-E14)	1,632	5.4	157.8
5	Chronic lower respiratory diseases (J40-J47)	1,352	4.5	130.7
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,074	3.6	103.8
7	Septicemia (A40-A41)	750	2.5	72.5
8	Influenza and pneumonia (J09-J18)	536	1.8	51.8
9	Essential hypertension and hypertensive renal disease (I10,I12,I15)	518	1.7	50.1
10	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	508	1.7	49.1
11	Alzheimer's disease (G30)	372	1.2	36.0
12	Pneumonitis due to solids and liquids (J69)	208	0.7	20.1
13	Chronic liver disease and cirrhosis (K70,K73-K74)	188	0.6	18.2
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	160	0.5	15.5
15	Parkinson's disease (G20-G21)	120	0.4	11.6
...	All other causes (residual)	4,407	14.7	426.0

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2014

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2014

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
Black, both sexes, 75-79 years				
...	All causes	30,364	100.0	4,217.1
1	Malignant neoplasms (C00-C97)	7,814	25.7	1,085.2
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	7,686	25.3	1,067.5
3	Cerebrovascular diseases (I60-I69)	2,020	6.7	280.5
4	Diabetes mellitus (E10-E14)	1,624	5.3	225.5
5	Chronic lower respiratory diseases (J40-J47)	1,332	4.4	185.0
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,042	3.4	144.7
7	Alzheimer's disease (G30)	803	2.6	111.5
8	Septicemia (A40-A41)	727	2.4	101.0
9	Influenza and pneumonia (J09-J18)	589	1.9	81.8
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	579	1.9	80.4
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	475	1.6	66.0
12	Pneumonitis due to solids and liquids (J69)	237	0.8	32.9
13	Parkinson's disease (G20-G21)	230	0.8	31.9
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	159	0.5	22.1
15	Chronic liver disease and cirrhosis (K70,K73-K74)	124	0.4	17.2
...	All other causes (residual)	4,923	16.2	683.7
Black, both sexes, 80-84 years				
...	All causes	30,064	100.0	6,305.1
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	7,889	26.2	1,654.5
2	Malignant neoplasms (C00-C97)	6,462	21.5	1,355.2
3	Cerebrovascular diseases (I60-I69)	2,163	7.2	453.6
4	Diabetes mellitus (E10-E14)	1,388	4.6	291.1
5	Alzheimer's disease (G30)	1,270	4.2	266.3
6	Chronic lower respiratory diseases (J40-J47)	1,226	4.1	257.1
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,041	3.5	218.3
8	Septicemia (A40-A41)	660	2.2	138.4
9	Essential hypertension and hypertensive renal disease (I10,I12,I15)	639	2.1	134.0
10	Influenza and pneumonia (J09-J18)	627	2.1	131.5
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	386	1.3	81.0
12	Parkinson's disease (G20-G21)	259	0.9	54.3
13	Pneumonitis due to solids and liquids (J69)	249	0.8	52.2
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	155	0.5	32.5
15	Aortic aneurysm and dissection (I71)	91	0.3	19.1
...	All other causes (residual)	5,559	18.5	1,165.8

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2014

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2014

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
Black, both sexes, 85-89 years				
...	All causes	27,254	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	7,345	27.0	---
2	Malignant neoplasms (C00-C97)	4,523	16.6	---
3	Cerebrovascular diseases (I60-I69)	2,005	7.4	---
4	Alzheimer's disease (G30)	1,575	5.8	---
5	Diabetes mellitus (E10-E14)	1,125	4.1	---
6	Chronic lower respiratory diseases (J40-J47)	1,008	3.7	---
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	956	3.5	---
8	Influenza and pneumonia (J09-J18)	665	2.4	---
9	Septicemia (A40-A41)	613	2.2	---
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	610	2.2	---
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	346	1.3	---
12	Pneumonitis due to solids and liquids (J69)	239	0.9	---
13	Parkinson's disease (G20-G21)	238	0.9	---
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	150	0.6	---
15	Nutritional deficiencies (E40-E64)	86	0.3	---
...	All other causes (residual)	5,770	21.2	---
Black, both sexes, 90-94 years				
...	All causes	18,794	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	5,347	28.5	---
2	Malignant neoplasms (C00-C97)	2,201	11.7	---
3	Alzheimer's disease (G30)	1,430	7.6	---
4	Cerebrovascular diseases (I60-I69)	1,411	7.5	---
5	Diabetes mellitus (E10-E14)	643	3.4	---
6	Chronic lower respiratory diseases (J40-J47)	554	2.9	---
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	550	2.9	---
8	Influenza and pneumonia (J09-J18)	522	2.8	---
9	Essential hypertension and hypertensive renal disease (I10,I12,I15)	475	2.5	---
10	Septicemia (A40-A41)	420	2.2	---
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	250	1.3	---
12	Pneumonitis due to solids and liquids (J69)	174	0.9	---
13	Parkinson's disease (G20-G21)	143	0.8	---
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	82	0.4	---
15	Atherosclerosis (I70)	69	0.4	---
...	All other causes (residual)	4,523	24.1	---

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2014

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2014

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
Black, both sexes, 95-99 years				
...	All causes	7,746	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,297	29.7	---
2	Alzheimer's disease (G30)	660	8.5	---
3	Malignant neoplasms (C00-C97)	644	8.3	---
4	Cerebrovascular diseases (I60-I69)	596	7.7	---
5	Essential hypertension and hypertensive renal disease (I10,I12,I15)	227	2.9	---
6	Influenza and pneumonia (J09-J18)	225	2.9	---
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	204	2.6	---
8	Diabetes mellitus (E10-E14)	193	2.5	---
9	Chronic lower respiratory diseases (J40-J47)	186	2.4	---
10	Septicemia (A40-A41)	147	1.9	---
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	115	1.5	---
12	Pneumonitis due to solids and liquids (J69)	78	1.0	---
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	41	0.5	---
14	Parkinson's disease (G20-G21)	35	0.5	---
15	Nutritional deficiencies (E40-E64)	33	0.4	---
...	All other causes (residual)	2,065	26.7	---
Black, both sexes, 100 years and over				
...	All causes	2,533	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	763	30.1	---
2	Alzheimer's disease (G30)	231	9.1	---
3	Cerebrovascular diseases (I60-I69)	178	7.0	---
4	Malignant neoplasms (C00-C97)	126	5.0	---
5	Essential hypertension and hypertensive renal disease (I10,I12,I15)	90	3.6	---
6	Influenza and pneumonia (J09-J18)	77	3.0	---
7	Chronic lower respiratory diseases (J40-J47)	64	2.5	---
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	61	2.4	---
9	Diabetes mellitus (E10-E14)	51	2.0	---
10	Septicemia (A40-A41)	39	1.5	---
11	Pneumonitis due to solids and liquids (J69)	30	1.2	---
12	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	23	0.9	---
13	Atherosclerosis (I70)	19	0.8	---
14	Nutritional deficiencies (E40-E64)	12	0.5	---
15	Anemias (D50-D64)	9	0.4	---
...	All other causes (residual)	760	30.0	---

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2014

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2014

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
Black, male, all ages				
...	All causes	157,733	100.0	742.6
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	37,962	24.1	178.7
2	Malignant neoplasms (C00-C97)	35,061	22.2	165.1
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	9,567	6.1	45.0
4	Cerebrovascular diseases (I60-I69)	7,747	4.9	36.5
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	6,798	4.3	32.0
6	Diabetes mellitus (E10-E14)	6,452	4.1	30.4
7	Chronic lower respiratory diseases (J40-J47)	5,035	3.2	23.7
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	4,034	2.6	19.0
9	Septicemia (A40-A41)	2,969	1.9	14.0
10	Influenza and pneumonia (J09-J18)	2,736	1.7	12.9
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	2,481	1.6	11.7
12	Human immunodeficiency virus (HIV) disease (B20-B24)	2,373	1.5	11.2
13	Certain conditions originating in the perinatal period (P00-P96)	2,266	1.4	10.7
14	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,946	1.2	9.2
15	Chronic liver disease and cirrhosis (K70,K73-K74)	1,940	1.2	9.1
...	All other causes (residual)	28,366	18.0	133.5
Black, male, 1-4 years				
...	All causes	583	100.0	42.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	179	30.7	12.9
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	81	13.9	5.9
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	59	10.1	4.3
4	Malignant neoplasms (C00-C97)	28	4.8	2.0
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	26	4.5	1.9
6	Influenza and pneumonia (J09-J18)	19	3.3	*
7	Chronic lower respiratory diseases (J40-J47)	11	1.9	*
8	Certain conditions originating in the perinatal period (P00-P96)	9	1.5	*
9	Septicemia (A40-A41)	7	1.2	*
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	6	1.0	*
11	Anemias (D50-D64)	5	0.9	*
11	Cerebrovascular diseases (I60-I69)	5	0.9	*
13	Acute bronchitis and bronchiolitis (J20-J21)	4	0.7	*
14	Meningitis (G00,G03)	3	0.5	*
14	Complications of medical and surgical care (Y40-Y84,Y88)	3	0.5	*
...	All other causes (residual)	138	23.7	10.0

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2014

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2014

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
Black, male, 5-9 years				
...	All causes	314	100.0	18.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	102	32.5	5.8
2	Malignant neoplasms (C00-C97)	37	11.8	2.1
3	Chronic lower respiratory diseases (J40-J47)	28	8.9	1.6
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	25	8.0	1.4
5	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	21	6.7	1.2
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	14	4.5	*
7	Influenza and pneumonia (J09-J18)	8	2.5	*
8	Septicemia (A40-A41)	5	1.6	*
9	Anemias (D50-D64)	4	1.3	*
9	Certain conditions originating in the perinatal period (P00-P96)	4	1.3	*
11	Cerebrovascular diseases (I60-I69)	3	1.0	*
12	Acute bronchitis and bronchiolitis (J20-J21)	2	0.6	*
13	Diabetes mellitus (E10-E14)	1	0.3	*
13	Meningitis (G00,G03)	1	0.3	*
13	Diseases of appendix (K35-K38)	1	0.3	*
13	Hernia (K40-K46)	1	0.3	*
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1	0.3	*
13	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1	0.3	*
13	Complications of medical and surgical care (Y40-Y84,Y88)	1	0.3	*
...	All other causes (residual)	54	17.2	3.1
Black, male, 10-14 years				
...	All causes	411	100.0	24.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	123	29.9	7.2
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	45	10.9	2.6
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	41	10.0	2.4
4	Malignant neoplasms (C00-C97)	40	9.7	2.3
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	26	6.3	1.5
6	Chronic lower respiratory diseases (J40-J47)	23	5.6	1.3
7	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	21	5.1	1.2
8	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	5	1.2	*
9	Septicemia (A40-A41)	4	1.0	*
9	Anemias (D50-D64)	4	1.0	*
9	Diabetes mellitus (E10-E14)	4	1.0	*
9	Cerebrovascular diseases (I60-I69)	4	1.0	*
13	Certain conditions originating in the perinatal period (P00-P96)	3	0.7	*
14	Influenza and pneumonia (J09-J18)	2	0.5	*
14	Diseases of appendix (K35-K38)	2	0.5	*
...	All other causes (residual)	64	15.6	3.7

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2014

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2014

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
Black, male, 15-19 years				
...	All causes	1,668	100.0	94.4
1	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	806	48.3	45.6
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	407	24.4	23.0
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	119	7.1	6.7
4	Malignant neoplasms (C00-C97)	63	3.8	3.6
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	59	3.5	3.3
6	Chronic lower respiratory diseases (J40-J47)	17	1.0	*
7	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	15	0.9	*
8	Cerebrovascular diseases (I60-I69)	9	0.5	*
9	Influenza and pneumonia (J09-J18)	8	0.5	*
9	Legal intervention (Y35,Y89.0)	8	0.5	*
11	Diabetes mellitus (E10-E14)	6	0.4	*
12	Human immunodeficiency virus (HIV) disease (B20-B24)	5	0.3	*
13	Septicemia (A40-A41)	4	0.2	*
13	Anemias (D50-D64)	4	0.2	*
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	4	0.2	*
...	All other causes (residual)	134	8.0	7.6
Black, male, 20-24 years				
...	All causes	3,390	100.0	172.3
1	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,610	47.5	81.8
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	728	21.5	37.0
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	329	9.7	16.7
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	113	3.3	5.7
5	Malignant neoplasms (C00-C97)	98	2.9	5.0
6	Human immunodeficiency virus (HIV) disease (B20-B24)	47	1.4	2.4
7	Chronic lower respiratory diseases (J40-J47)	30	0.9	1.5
8	Anemias (D50-D64)	29	0.9	1.5
9	Diabetes mellitus (E10-E14)	27	0.8	1.4
10	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	23	0.7	1.2
11	Legal intervention (Y35,Y89.0)	21	0.6	1.1
12	Cerebrovascular diseases (I60-I69)	20	0.6	1.0
13	Influenza and pneumonia (J09-J18)	19	0.6	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	13	0.4	*
15	Septicemia (A40-A41)	8	0.2	*
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	8	0.2	*
...	All other causes (residual)	267	7.9	13.6

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2014

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2014

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
Black, male, 25-29 years				
...	All causes	3,292	100.0	200.4
1	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,224	37.2	74.5
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	733	22.3	44.6
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	264	8.0	16.1
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	219	6.7	13.3
5	Human immunodeficiency virus (HIV) disease (B20-B24)	127	3.9	7.7
6	Malignant neoplasms (C00-C97)	100	3.0	6.1
7	Diabetes mellitus (E10-E14)	46	1.4	2.8
8	Chronic lower respiratory diseases (J40-J47)	36	1.1	2.2
9	Anemias (D50-D64)	31	0.9	1.9
10	Cerebrovascular diseases (I60-I69)	30	0.9	1.8
11	Legal intervention (Y35,Y89.0)	29	0.9	1.8
12	Septicemia (A40-A41)	21	0.6	1.3
13	Influenza and pneumonia (J09-J18)	20	0.6	1.2
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	20	0.6	1.2
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	15	0.5	*
...	All other causes (residual)	377	11.5	22.9
Black, male, 30-34 years				
...	All causes	3,330	100.0	225.1
1	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	883	26.5	59.7
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	741	22.3	50.1
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	410	12.3	27.7
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	212	6.4	14.3
5	Malignant neoplasms (C00-C97)	158	4.7	10.7
6	Human immunodeficiency virus (HIV) disease (B20-B24)	126	3.8	8.5
7	Diabetes mellitus (E10-E14)	84	2.5	5.7
8	Cerebrovascular diseases (I60-I69)	61	1.8	4.1
9	Anemias (D50-D64)	49	1.5	3.3
10	Chronic lower respiratory diseases (J40-J47)	38	1.1	2.6
11	Influenza and pneumonia (J09-J18)	28	0.8	1.9
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	23	0.7	1.6
13	Septicemia (A40-A41)	22	0.7	1.5
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	22	0.7	1.5
15	Aortic aneurysm and dissection (I71)	19	0.6	*
...	All other causes (residual)	454	13.6	30.7

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2014

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2014

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
Black, male, 35-39 years				
...	All causes	3,629	100.0	272.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	643	17.7	48.3
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	636	17.5	47.8
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	626	17.2	47.0
4	Malignant neoplasms (C00-C97)	265	7.3	19.9
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	193	5.3	14.5
6	Human immunodeficiency virus (HIV) disease (B20-B24)	155	4.3	11.6
7	Diabetes mellitus (E10-E14)	113	3.1	8.5
8	Cerebrovascular diseases (I60-I69)	110	3.0	8.3
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	60	1.7	4.5
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	50	1.4	3.8
11	Chronic lower respiratory diseases (J40-J47)	46	1.3	3.5
12	Influenza and pneumonia (J09-J18)	41	1.1	3.1
13	Anemias (D50-D64)	33	0.9	2.5
14	Septicemia (A40-A41)	30	0.8	2.3
14	Chronic liver disease and cirrhosis (K70,K73-K74)	30	0.8	2.3
...	All other causes (residual)	598	16.5	44.9
Black, male, 40-44 years				
...	All causes	4,581	100.0	344.4
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,091	23.8	82.0
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	651	14.2	48.9
3	Malignant neoplasms (C00-C97)	501	10.9	37.7
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	418	9.1	31.4
5	Human immunodeficiency virus (HIV) disease (B20-B24)	211	4.6	15.9
6	Cerebrovascular diseases (I60-I69)	194	4.2	14.6
7	Diabetes mellitus (E10-E14)	188	4.1	14.1
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	159	3.5	12.0
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	90	2.0	6.8
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	79	1.7	5.9
11	Septicemia (A40-A41)	63	1.4	4.7
12	Influenza and pneumonia (J09-J18)	62	1.4	4.7
13	Chronic liver disease and cirrhosis (K70,K73-K74)	61	1.3	4.6
14	Chronic lower respiratory diseases (J40-J47)	55	1.2	4.1
15	Aortic aneurysm and dissection (I71)	49	1.1	3.7
...	All other causes (residual)	709	15.5	53.3

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2014

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2014

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
Black, male, 45-49 years				
...	All causes	6,892	100.0	521.8
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,823	26.5	138.0
2	Malignant neoplasms (C00-C97)	1,039	15.1	78.7
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	778	11.3	58.9
4	Diabetes mellitus (E10-E14)	336	4.9	25.4
5	Human immunodeficiency virus (HIV) disease (B20-B24)	327	4.7	24.8
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	324	4.7	24.5
7	Cerebrovascular diseases (I60-I69)	285	4.1	21.6
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	143	2.1	10.8
9	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	137	2.0	10.4
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	119	1.7	9.0
10	Chronic liver disease and cirrhosis (K70,K73-K74)	119	1.7	9.0
12	Influenza and pneumonia (J09-J18)	113	1.6	8.6
13	Septicemia (A40-A41)	105	1.5	7.9
14	Chronic lower respiratory diseases (J40-J47)	88	1.3	6.7
15	Aortic aneurysm and dissection (I71)	56	0.8	4.2
...	All other causes (residual)	1,100	16.0	83.3
Black, male, 50-54 years				
...	All causes	11,119	100.0	817.5
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,989	26.9	219.8
2	Malignant neoplasms (C00-C97)	2,413	21.7	177.4
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	954	8.6	70.1
4	Cerebrovascular diseases (I60-I69)	509	4.6	37.4
5	Diabetes mellitus (E10-E14)	488	4.4	35.9
6	Human immunodeficiency virus (HIV) disease (B20-B24)	395	3.6	29.0
7	Chronic liver disease and cirrhosis (K70,K73-K74)	311	2.8	22.9
8	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	267	2.4	19.6
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	222	2.0	16.3
10	Septicemia (A40-A41)	198	1.8	14.6
11	Chronic lower respiratory diseases (J40-J47)	192	1.7	14.1
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	176	1.6	12.9
13	Influenza and pneumonia (J09-J18)	151	1.4	11.1
14	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	147	1.3	10.8
15	Viral hepatitis (B15-B19)	93	0.8	6.8
...	All other causes (residual)	1,614	14.5	118.7

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2014

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2014

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
Black, male, 55-59 years				
...	All causes	16,177	100.0	1,324.1
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	4,275	26.4	349.9
2	Malignant neoplasms (C00-C97)	4,263	26.4	348.9
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,005	6.2	82.3
4	Cerebrovascular diseases (I60-I69)	788	4.9	64.5
5	Diabetes mellitus (E10-E14)	738	4.6	60.4
6	Chronic liver disease and cirrhosis (K70,K73-K74)	442	2.7	36.2
7	Chronic lower respiratory diseases (J40-J47)	411	2.5	33.6
8	Human immunodeficiency virus (HIV) disease (B20-B24)	405	2.5	33.1
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	380	2.3	31.1
10	Septicemia (A40-A41)	314	1.9	25.7
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	272	1.7	22.3
12	Influenza and pneumonia (J09-J18)	220	1.4	18.0
13	Viral hepatitis (B15-B19)	197	1.2	16.1
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	178	1.1	14.6
15	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	107	0.7	8.8
...	All other causes (residual)	2,182	13.5	178.6
Black, male, 60-64 years				
...	All causes	18,666	100.0	1,984.9
1	Malignant neoplasms (C00-C97)	5,478	29.3	582.5
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	4,924	26.4	523.6
3	Cerebrovascular diseases (I60-I69)	928	5.0	98.7
4	Diabetes mellitus (E10-E14)	863	4.6	91.8
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	721	3.9	76.7
6	Chronic lower respiratory diseases (J40-J47)	528	2.8	56.1
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	509	2.7	54.1
8	Chronic liver disease and cirrhosis (K70,K73-K74)	421	2.3	44.8
9	Septicemia (A40-A41)	370	2.0	39.3
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	317	1.7	33.7
11	Influenza and pneumonia (J09-J18)	308	1.7	32.8
12	Human immunodeficiency virus (HIV) disease (B20-B24)	298	1.6	31.7
13	Viral hepatitis (B15-B19)	247	1.3	26.3
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	110	0.6	11.7
15	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	87	0.5	9.3
...	All other causes (residual)	2,557	13.7	271.9

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2014

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2014

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
Black, male, 65-69 years				
...	All causes	17,440	100.0	2,617.7
1	Malignant neoplasms (C00-C97)	5,367	30.8	805.6
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	4,581	26.3	687.6
3	Cerebrovascular diseases (I60-I69)	919	5.3	137.9
4	Diabetes mellitus (E10-E14)	862	4.9	129.4
5	Chronic lower respiratory diseases (J40-J47)	688	3.9	103.3
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	500	2.9	75.0
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	463	2.7	69.5
8	Septicemia (A40-A41)	374	2.1	56.1
9	Chronic liver disease and cirrhosis (K70,K73-K74)	290	1.7	43.5
10	Influenza and pneumonia (J09-J18)	281	1.6	42.2
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	279	1.6	41.9
12	Human immunodeficiency virus (HIV) disease (B20-B24)	156	0.9	23.4
13	Viral hepatitis (B15-B19)	115	0.7	17.3
14	Pneumonitis due to solids and liquids (J69)	100	0.6	15.0
15	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	81	0.5	12.2
...	All other causes (residual)	2,384	13.7	357.8
Black, male, 70-74 years				
...	All causes	16,059	100.0	3,708.6
1	Malignant neoplasms (C00-C97)	4,810	30.0	1,110.8
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	4,154	25.9	959.3
3	Cerebrovascular diseases (I60-I69)	948	5.9	218.9
4	Diabetes mellitus (E10-E14)	794	4.9	183.4
5	Chronic lower respiratory diseases (J40-J47)	776	4.8	179.2
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	504	3.1	116.4
7	Septicemia (A40-A41)	382	2.4	88.2
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	321	2.0	74.1
9	Influenza and pneumonia (J09-J18)	316	2.0	73.0
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	270	1.7	62.4
11	Alzheimer's disease (G30)	138	0.9	31.9
12	Pneumonitis due to solids and liquids (J69)	130	0.8	30.0
13	Chronic liver disease and cirrhosis (K70,K73-K74)	129	0.8	29.8
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	82	0.5	18.9
15	Parkinson's disease (G20-G21)	77	0.5	17.8
...	All other causes (residual)	2,228	13.9	514.5

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2014

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2014

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
Black, male, 75-79 years				
...	All causes	14,965	100.0	5,248.3
1	Malignant neoplasms (C00-C97)	3,969	26.5	1,391.9
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,961	26.5	1,389.1
3	Cerebrovascular diseases (I60-I69)	942	6.3	330.4
4	Diabetes mellitus (E10-E14)	779	5.2	273.2
5	Chronic lower respiratory diseases (J40-J47)	690	4.6	242.0
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	475	3.2	166.6
7	Septicemia (A40-A41)	341	2.3	119.6
8	Influenza and pneumonia (J09-J18)	319	2.1	111.9
9	Alzheimer's disease (G30)	312	2.1	109.4
10	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	273	1.8	95.7
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	267	1.8	93.6
12	Parkinson's disease (G20-G21)	140	0.9	49.1
13	Pneumonitis due to solids and liquids (J69)	132	0.9	46.3
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	77	0.5	27.0
15	Chronic liver disease and cirrhosis (K70,K73-K74)	66	0.4	23.1
...	All other causes (residual)	2,222	14.8	779.3
Black, male, 80-84 years				
...	All causes	13,196	100.0	7,713.2
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,573	27.1	2,088.5
2	Malignant neoplasms (C00-C97)	3,191	24.2	1,865.2
3	Cerebrovascular diseases (I60-I69)	852	6.5	498.0
4	Chronic lower respiratory diseases (J40-J47)	590	4.5	344.9
5	Diabetes mellitus (E10-E14)	512	3.9	299.3
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	433	3.3	253.1
7	Alzheimer's disease (G30)	410	3.1	239.6
8	Septicemia (A40-A41)	299	2.3	174.8
9	Influenza and pneumonia (J09-J18)	269	2.0	157.2
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	244	1.8	142.6
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	207	1.6	121.0
12	Parkinson's disease (G20-G21)	153	1.2	89.4
13	Pneumonitis due to solids and liquids (J69)	124	0.9	72.5
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	58	0.4	33.9
15	Chronic liver disease and cirrhosis (K70,K73-K74)	33	0.3	19.3
...	All other causes (residual)	2,248	17.0	1,314.0

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2014

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2014

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
Black, male, 85-89 years				
...	All causes	10,227	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,688	26.3	---
2	Malignant neoplasms (C00-C97)	2,059	20.1	---
3	Cerebrovascular diseases (I60-I69)	661	6.5	---
4	Chronic lower respiratory diseases (J40-J47)	476	4.7	---
5	Alzheimer's disease (G30)	438	4.3	---
6	Diabetes mellitus (E10-E14)	371	3.6	---
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	364	3.6	---
8	Influenza and pneumonia (J09-J18)	290	2.8	---
9	Septicemia (A40-A41)	225	2.2	---
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	187	1.8	---
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	164	1.6	---
12	Parkinson's disease (G20-G21)	132	1.3	---
13	Pneumonitis due to solids and liquids (J69)	109	1.1	---
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	53	0.5	---
15	Nutritional deficiencies (E40-E64)	34	0.3	---
...	All other causes (residual)	1,976	19.3	---
Black, male, 90-94 years				
...	All causes	5,688	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,659	29.2	---
2	Malignant neoplasms (C00-C97)	923	16.2	---
3	Cerebrovascular diseases (I60-I69)	334	5.9	---
4	Alzheimer's disease (G30)	326	5.7	---
5	Chronic lower respiratory diseases (J40-J47)	218	3.8	---
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	194	3.4	---
7	Diabetes mellitus (E10-E14)	182	3.2	---
8	Influenza and pneumonia (J09-J18)	165	2.9	---
9	Essential hypertension and hypertensive renal disease (I10,I12,I15)	125	2.2	---
10	Septicemia (A40-A41)	124	2.2	---
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	97	1.7	---
12	Parkinson's disease (G20-G21)	70	1.2	---
13	Pneumonitis due to solids and liquids (J69)	68	1.2	---
14	Nutritional deficiencies (E40-E64)	26	0.5	---
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	25	0.4	---
...	All other causes (residual)	1,152	20.3	---

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2014

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2014

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
Black, male, 95-99 years				
...	All causes	1,809	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	563	31.1	---
2	Malignant neoplasms (C00-C97)	222	12.3	---
3	Cerebrovascular diseases (I60-I69)	116	6.4	---
4	Alzheimer's disease (G30)	97	5.4	---
5	Chronic lower respiratory diseases (J40-J47)	66	3.6	---
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	65	3.6	---
7	Influenza and pneumonia (J09-J18)	52	2.9	---
8	Diabetes mellitus (E10-E14)	51	2.8	---
9	Essential hypertension and hypertensive renal disease (I10,I12,I15)	44	2.4	---
10	Septicemia (A40-A41)	42	2.3	---
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	37	2.0	---
12	Pneumonitis due to solids and liquids (J69)	23	1.3	---
13	Parkinson's disease (G20-G21)	13	0.7	---
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	9	0.5	---
15	Nutritional deficiencies (E40-E64)	8	0.4	---
...	All other causes (residual)	401	22.2	---
Black, male, 100 years and over				
...	All causes	376	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	119	31.6	---
2	Malignant neoplasms (C00-C97)	35	9.3	---
3	Chronic lower respiratory diseases (J40-J47)	22	5.9	---
4	Influenza and pneumonia (J09-J18)	17	4.5	---
4	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	17	4.5	---
6	Alzheimer's disease (G30)	16	4.3	---
6	Essential hypertension and hypertensive renal disease (I10,I12,I15)	16	4.3	---
6	Cerebrovascular diseases (I60-I69)	16	4.3	---
9	Pneumonitis due to solids and liquids (J69)	8	2.1	---
10	Diabetes mellitus (E10-E14)	5	1.3	---
11	Septicemia (A40-A41)	4	1.1	---
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	4	1.1	---
13	Atherosclerosis (I70)	3	0.8	---
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2	0.5	---
14	Nutritional deficiencies (E40-E64)	2	0.5	---
...	All other causes (residual)	90	23.9	---

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2014

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2014

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
Black, female, all ages				
...	All causes	151,227	100.0	655.5
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	35,133	23.2	152.3
2	Malignant neoplasms (C00-C97)	34,029	22.5	147.5
3	Cerebrovascular diseases (I60-I69)	9,341	6.2	40.5
4	Diabetes mellitus (E10-E14)	6,983	4.6	30.3
5	Chronic lower respiratory diseases (J40-J47)	4,899	3.2	21.2
6	Alzheimer's disease (G30)	4,730	3.1	20.5
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	4,601	3.0	19.9
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	4,552	3.0	19.7
9	Septicemia (A40-A41)	3,417	2.3	14.8
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	2,918	1.9	12.6
11	Influenza and pneumonia (J09-J18)	2,875	1.9	12.5
12	Certain conditions originating in the perinatal period (P00-P96)	1,765	1.2	7.7
13	Human immunodeficiency virus (HIV) disease (B20-B24)	1,218	0.8	5.3
14	Chronic liver disease and cirrhosis (K70,K73-K74)	1,163	0.8	5.0
15	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,078	0.7	4.7
...	All other causes (residual)	32,525	21.5	141.0
Black, female, 1-4 years				
...	All causes	426	100.0	31.9
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	112	26.3	8.4
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	70	16.4	5.2
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	45	10.6	3.4
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	29	6.8	2.2
5	Malignant neoplasms (C00-C97)	26	6.1	1.9
6	Chronic lower respiratory diseases (J40-J47)	12	2.8	*
7	Influenza and pneumonia (J09-J18)	10	2.3	*
8	Septicemia (A40-A41)	5	1.2	*
9	Cerebrovascular diseases (I60-I69)	4	0.9	*
9	Complications of medical and surgical care (Y40-Y84,Y88)	4	0.9	*
11	Acute bronchitis and bronchiolitis (J20-J21)	3	0.7	*
11	Certain conditions originating in the perinatal period (P00-P96)	3	0.7	*
13	Salmonella infections (A01-A02)	1	0.2	*
13	Meningococcal infection (A39)	1	0.2	*
13	Malaria (B50-B54)	1	0.2	*
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1	0.2	*
13	Anemias (D50-D64)	1	0.2	*
13	Nutritional deficiencies (E40-E64)	1	0.2	*
13	Meningitis (G00,G03)	1	0.2	*
13	Pneumonitis due to solids and liquids (J69)	1	0.2	*
...	All other causes (residual)	95	22.3	7.1

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2014

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2014

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
Black, female, 5-9 years				
...	All causes	223	100.0	13.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	52	23.3	3.1
2	Malignant neoplasms (C00-C97)	40	17.9	2.4
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	18	8.1	*
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	17	7.6	*
5	Chronic lower respiratory diseases (J40-J47)	10	4.5	*
6	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	9	4.0	*
7	Cerebrovascular diseases (I60-I69)	7	3.1	*
8	Anemias (D50-D64)	5	2.2	*
8	Diseases of heart (I00-I09,I11,I13,I20-I51)	5	2.2	*
10	Influenza and pneumonia (J09-J18)	4	1.8	*
11	Septicemia (A40-A41)	3	1.3	*
12	Meningitis (G00,G03)	2	0.9	*
13	Diabetes mellitus (E10-E14)	1	0.4	*
13	Hernia (K40-K46)	1	0.4	*
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1	0.4	*
13	Certain conditions originating in the perinatal period (P00-P96)	1	0.4	*
13	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1	0.4	*
13	Complications of medical and surgical care (Y40-Y84,Y88)	1	0.4	*
...	All other causes (residual)	45	20.2	2.7
Black, female, 10-14 years				
...	All causes	256	100.0	15.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	59	23.0	3.6
2	Malignant neoplasms (C00-C97)	29	11.3	1.7
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	17	6.6	*
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	16	6.3	*
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	15	5.9	*
6	Chronic lower respiratory diseases (J40-J47)	14	5.5	*
7	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	13	5.1	*
8	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	7	2.7	*
9	Anemias (D50-D64)	5	2.0	*
9	Diabetes mellitus (E10-E14)	5	2.0	*
11	Septicemia (A40-A41)	4	1.6	*
12	Influenza and pneumonia (J09-J18)	3	1.2	*
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2	0.8	*
13	Certain conditions originating in the perinatal period (P00-P96)	2	0.8	*
13	Complications of medical and surgical care (Y40-Y84,Y88)	2	0.8	*
...	All other causes (residual)	63	24.6	3.8

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2014

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2014

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
Black, female, 15-19 years				
...	All causes	497	100.0	29.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	141	28.4	8.2
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	95	19.1	5.6
3	Malignant neoplasms (C00-C97)	38	7.6	2.2
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	38	7.6	2.2
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	27	5.4	1.6
6	Anemias (D50-D64)	13	2.6	*
7	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	12	2.4	*
8	Pregnancy, childbirth and the puerperium (O00-O99)	11	2.2	*
9	Septicemia (A40-A41)	8	1.6	*
10	Chronic lower respiratory diseases (J40-J47)	6	1.2	*
11	Cerebrovascular diseases (I60-I69)	3	0.6	*
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3	0.6	*
13	Human immunodeficiency virus (HIV) disease (B20-B24)	2	0.4	*
13	Complications of medical and surgical care (Y40-Y84,Y88)	2	0.4	*
15	Diabetes mellitus (E10-E14)	1	0.2	*
15	Influenza and pneumonia (J09-J18)	1	0.2	*
15	Diseases of appendix (K35-K38)	1	0.2	*
15	Infections of kidney (N10-N12,N13.6,N15.1)	1	0.2	*
15	Certain conditions originating in the perinatal period (P00-P96)	1	0.2	*
...	All other causes (residual)	93	18.7	5.4
Black, female, 20-24 years				
...	All causes	1,047	100.0	54.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	247	23.6	12.9
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	160	15.3	8.3
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	85	8.1	4.4
4	Malignant neoplasms (C00-C97)	73	7.0	3.8
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	58	5.5	3.0
6	Pregnancy, childbirth and the puerperium (O00-O99)	46	4.4	2.4
7	Human immunodeficiency virus (HIV) disease (B20-B24)	32	3.1	1.7
8	Diabetes mellitus (E10-E14)	26	2.5	1.4
9	Anemias (D50-D64)	19	1.8	*
9	Chronic lower respiratory diseases (J40-J47)	19	1.8	*
11	Cerebrovascular diseases (I60-I69)	17	1.6	*
11	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	17	1.6	*
13	Septicemia (A40-A41)	12	1.1	*
14	Influenza and pneumonia (J09-J18)	10	1.0	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	9	0.9	*
...	All other causes (residual)	217	20.7	11.3

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2014

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2014

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
Black, female, 25-29 years				
...	All causes	1,221	100.0	71.9
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	236	19.3	13.9
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	153	12.5	9.0
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	121	9.9	7.1
4	Malignant neoplasms (C00-C97)	120	9.8	7.1
5	Pregnancy, childbirth and the puerperium (O00-O99)	62	5.1	3.7
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	44	3.6	2.6
7	Diabetes mellitus (E10-E14)	43	3.5	2.5
8	Human immunodeficiency virus (HIV) disease (B20-B24)	35	2.9	2.1
9	Cerebrovascular diseases (I60-I69)	31	2.5	1.8
10	Chronic lower respiratory diseases (J40-J47)	24	2.0	1.4
11	Influenza and pneumonia (J09-J18)	23	1.9	1.4
12	Anemias (D50-D64)	21	1.7	1.2
13	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	18	1.5	*
14	Septicemia (A40-A41)	15	1.2	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	13	1.1	*
...	All other causes (residual)	262	21.5	15.4
Black, female, 30-34 years				
...	All causes	1,706	100.0	106.1
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	260	15.2	16.2
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	255	14.9	15.9
3	Malignant neoplasms (C00-C97)	251	14.7	15.6
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	100	5.9	6.2
5	Human immunodeficiency virus (HIV) disease (B20-B24)	88	5.2	5.5
6	Pregnancy, childbirth and the puerperium (O00-O99)	81	4.7	5.0
7	Diabetes mellitus (E10-E14)	66	3.9	4.1
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	47	2.8	2.9
9	Cerebrovascular diseases (I60-I69)	33	1.9	2.1
10	Influenza and pneumonia (J09-J18)	28	1.6	1.7
10	Chronic lower respiratory diseases (J40-J47)	28	1.6	1.7
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	28	1.6	1.7
13	Anemias (D50-D64)	26	1.5	1.6
14	Septicemia (A40-A41)	23	1.3	1.4
14	Chronic liver disease and cirrhosis (K70,K73-K74)	23	1.3	1.4
...	All other causes (residual)	369	21.6	23.0

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2014

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2014

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
Black, female, 35-39 years				
...	All causes	2,330	100.0	157.0
1	Malignant neoplasms (C00-C97)	432	18.5	29.1
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	400	17.2	27.0
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	260	11.2	17.5
4	Human immunodeficiency virus (HIV) disease (B20-B24)	120	5.2	8.1
5	Diabetes mellitus (E10-E14)	92	3.9	6.2
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	84	3.6	5.7
7	Cerebrovascular diseases (I60-I69)	69	3.0	4.6
8	Pregnancy, childbirth and the puerperium (O00-O99)	61	2.6	4.1
9	Septicemia (A40-A41)	44	1.9	3.0
10	Influenza and pneumonia (J09-J18)	43	1.8	2.9
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	41	1.8	2.8
11	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	41	1.8	2.8
13	Anemias (D50-D64)	33	1.4	2.2
14	Chronic lower respiratory diseases (J40-J47)	32	1.4	2.2
15	Chronic liver disease and cirrhosis (K70,K73-K74)	30	1.3	2.0
...	All other causes (residual)	548	23.5	36.9
Black, female, 40-44 years				
...	All causes	3,457	100.0	230.5
1	Malignant neoplasms (C00-C97)	778	22.5	51.9
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	658	19.0	43.9
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	303	8.8	20.2
4	Human immunodeficiency virus (HIV) disease (B20-B24)	154	4.5	10.3
4	Diabetes mellitus (E10-E14)	154	4.5	10.3
6	Cerebrovascular diseases (I60-I69)	143	4.1	9.5
7	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	81	2.3	5.4
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	74	2.1	4.9
9	Chronic liver disease and cirrhosis (K70,K73-K74)	63	1.8	4.2
10	Influenza and pneumonia (J09-J18)	55	1.6	3.7
10	Chronic lower respiratory diseases (J40-J47)	55	1.6	3.7
12	Septicemia (A40-A41)	52	1.5	3.5
13	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	48	1.4	3.2
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	45	1.3	3.0
15	Pregnancy, childbirth and the puerperium (O00-O99)	42	1.2	2.8
...	All other causes (residual)	752	21.8	50.1

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2014

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2014

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
Black, female, 45-49 years				
...	All causes	5,209	100.0	351.0
1	Malignant neoplasms (C00-C97)	1,419	27.2	95.6
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,052	20.2	70.9
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	344	6.6	23.2
4	Cerebrovascular diseases (I60-I69)	237	4.5	16.0
5	Diabetes mellitus (E10-E14)	234	4.5	15.8
6	Human immunodeficiency virus (HIV) disease (B20-B24)	180	3.5	12.1
7	Chronic lower respiratory diseases (J40-J47)	119	2.3	8.0
8	Chronic liver disease and cirrhosis (K70,K73-K74)	110	2.1	7.4
9	Septicemia (A40-A41)	102	2.0	6.9
10	Influenza and pneumonia (J09-J18)	90	1.7	6.1
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	90	1.7	6.1
12	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	74	1.4	5.0
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	68	1.3	4.6
14	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	42	0.8	2.8
15	Pregnancy, childbirth and the puerperium (O00-O99)	36	0.7	2.4
...	All other causes (residual)	1,012	19.4	68.2
Black, female, 50-54 years				
...	All causes	8,595	100.0	556.3
1	Malignant neoplasms (C00-C97)	2,621	30.5	169.7
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,823	21.2	118.0
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	456	5.3	29.5
4	Cerebrovascular diseases (I60-I69)	442	5.1	28.6
5	Diabetes mellitus (E10-E14)	381	4.4	24.7
6	Chronic lower respiratory diseases (J40-J47)	252	2.9	16.3
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	205	2.4	13.3
8	Human immunodeficiency virus (HIV) disease (B20-B24)	198	2.3	12.8
9	Septicemia (A40-A41)	188	2.2	12.2
10	Chronic liver disease and cirrhosis (K70,K73-K74)	184	2.1	11.9
11	Influenza and pneumonia (J09-J18)	146	1.7	9.5
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	122	1.4	7.9
13	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	60	0.7	3.9
14	Viral hepatitis (B15-B19)	52	0.6	3.4
15	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	49	0.6	3.2
...	All other causes (residual)	1,416	16.5	91.7

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2014

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2014

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
Black, female, 55-59 years				
...	All causes	11,687	100.0	822.3
1	Malignant neoplasms (C00-C97)	3,827	32.7	269.3
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,480	21.2	174.5
3	Diabetes mellitus (E10-E14)	587	5.0	41.3
4	Cerebrovascular diseases (I60-I69)	543	4.6	38.2
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	428	3.7	30.1
6	Chronic lower respiratory diseases (J40-J47)	402	3.4	28.3
7	Septicemia (A40-A41)	295	2.5	20.8
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	276	2.4	19.4
9	Chronic liver disease and cirrhosis (K70,K73-K74)	241	2.1	17.0
10	Influenza and pneumonia (J09-J18)	194	1.7	13.7
11	Human immunodeficiency virus (HIV) disease (B20-B24)	181	1.5	12.7
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	177	1.5	12.5
13	Viral hepatitis (B15-B19)	101	0.9	7.1
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	44	0.4	3.1
15	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	41	0.4	2.9
...	All other causes (residual)	1,870	16.0	131.6
Black, female, 60-64 years				
...	All causes	13,442	100.0	1,162.2
1	Malignant neoplasms (C00-C97)	4,472	33.3	386.6
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,959	22.0	255.8
3	Diabetes mellitus (E10-E14)	712	5.3	61.6
4	Cerebrovascular diseases (I60-I69)	704	5.2	60.9
5	Chronic lower respiratory diseases (J40-J47)	491	3.7	42.5
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	418	3.1	36.1
7	Septicemia (A40-A41)	344	2.6	29.7
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	309	2.3	26.7
9	Essential hypertension and hypertensive renal disease (I10,I12,I15)	252	1.9	21.8
10	Influenza and pneumonia (J09-J18)	214	1.6	18.5
11	Chronic liver disease and cirrhosis (K70,K73-K74)	208	1.5	18.0
12	Human immunodeficiency virus (HIV) disease (B20-B24)	112	0.8	9.7
13	Viral hepatitis (B15-B19)	106	0.8	9.2
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	57	0.4	4.9
15	Aortic aneurysm and dissection (I71)	42	0.3	3.6
15	Pneumonitis due to solids and liquids (J69)	42	0.3	3.6
...	All other causes (residual)	2,000	14.9	172.9

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2014

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2014

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
Black, female, 65-69 years				
...	All causes	13,545	100.0	1,577.3
1	Malignant neoplasms (C00-C97)	4,486	33.1	522.4
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,087	22.8	359.5
3	Cerebrovascular diseases (I60-I69)	756	5.6	88.0
4	Diabetes mellitus (E10-E14)	719	5.3	83.7
5	Chronic lower respiratory diseases (J40-J47)	549	4.1	63.9
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	508	3.8	59.2
7	Septicemia (A40-A41)	357	2.6	41.6
8	Essential hypertension and hypertensive renal disease (I10,I12,I15)	227	1.7	26.4
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	211	1.6	24.6
10	Influenza and pneumonia (J09-J18)	195	1.4	22.7
11	Chronic liver disease and cirrhosis (K70,K73-K74)	119	0.9	13.9
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	82	0.6	9.5
13	Alzheimer's disease (G30)	64	0.5	7.5
14	Human immunodeficiency virus (HIV) disease (B20-B24)	59	0.4	6.9
15	Pneumonitis due to solids and liquids (J69)	52	0.4	6.1
...	All other causes (residual)	2,074	15.3	241.5
Black, female, 70-74 years				
...	All causes	13,911	100.0	2,312.7
1	Malignant neoplasms (C00-C97)	4,037	29.0	671.2
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,308	23.8	550.0
3	Cerebrovascular diseases (I60-I69)	888	6.4	147.6
4	Diabetes mellitus (E10-E14)	838	6.0	139.3
5	Chronic lower respiratory diseases (J40-J47)	576	4.1	95.8
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	570	4.1	94.8
7	Septicemia (A40-A41)	368	2.6	61.2
8	Essential hypertension and hypertensive renal disease (I10,I12,I15)	248	1.8	41.2
9	Alzheimer's disease (G30)	234	1.7	38.9
10	Influenza and pneumonia (J09-J18)	220	1.6	36.6
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	187	1.3	31.1
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	78	0.6	13.0
12	Pneumonitis due to solids and liquids (J69)	78	0.6	13.0
14	Chronic liver disease and cirrhosis (K70,K73-K74)	59	0.4	9.8
15	Aortic aneurysm and dissection (I71)	51	0.4	8.5
...	All other causes (residual)	2,171	15.6	360.9

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2014

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2014

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
Black, female, 75-79 years				
...	All causes	15,399	100.0	3,541.0
1	Malignant neoplasms (C00-C97)	3,845	25.0	884.2
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,725	24.2	856.6
3	Cerebrovascular diseases (I60-I69)	1,078	7.0	247.9
4	Diabetes mellitus (E10-E14)	845	5.5	194.3
5	Chronic lower respiratory diseases (J40-J47)	642	4.2	147.6
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	567	3.7	130.4
7	Alzheimer's disease (G30)	491	3.2	112.9
8	Septicemia (A40-A41)	386	2.5	88.8
9	Essential hypertension and hypertensive renal disease (I10,I12,I15)	312	2.0	71.7
10	Influenza and pneumonia (J09-J18)	270	1.8	62.1
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	202	1.3	46.4
12	Pneumonitis due to solids and liquids (J69)	105	0.7	24.1
13	Parkinson's disease (G20-G21)	90	0.6	20.7
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	82	0.5	18.9
15	Chronic liver disease and cirrhosis (K70,K73-K74)	58	0.4	13.3
...	All other causes (residual)	2,701	17.5	621.1
Black, female, 80-84 years				
...	All causes	16,868	100.0	5,517.2
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	4,316	25.6	1,411.7
2	Malignant neoplasms (C00-C97)	3,271	19.4	1,069.9
3	Cerebrovascular diseases (I60-I69)	1,311	7.8	428.8
4	Diabetes mellitus (E10-E14)	876	5.2	286.5
5	Alzheimer's disease (G30)	860	5.1	281.3
6	Chronic lower respiratory diseases (J40-J47)	636	3.8	208.0
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	608	3.6	198.9
8	Essential hypertension and hypertensive renal disease (I10,I12,I15)	395	2.3	129.2
9	Septicemia (A40-A41)	361	2.1	118.1
10	Influenza and pneumonia (J09-J18)	358	2.1	117.1
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	179	1.1	58.5
12	Pneumonitis due to solids and liquids (J69)	125	0.7	40.9
13	Parkinson's disease (G20-G21)	106	0.6	34.7
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	97	0.6	31.7
15	Aortic aneurysm and dissection (I71)	59	0.3	19.3
...	All other causes (residual)	3,310	19.6	1,082.6

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2014

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2014

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
Black, female, 85-89 years				
...	All causes	17,027	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	4,657	27.4	---
2	Malignant neoplasms (C00-C97)	2,464	14.5	---
3	Cerebrovascular diseases (I60-I69)	1,344	7.9	---
4	Alzheimer's disease (G30)	1,137	6.7	---
5	Diabetes mellitus (E10-E14)	754	4.4	---
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	592	3.5	---
7	Chronic lower respiratory diseases (J40-J47)	532	3.1	---
8	Essential hypertension and hypertensive renal disease (I10,I12,I15)	423	2.5	---
9	Septicemia (A40-A41)	388	2.3	---
10	Influenza and pneumonia (J09-J18)	375	2.2	---
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	182	1.1	---
12	Pneumonitis due to solids and liquids (J69)	130	0.8	---
13	Parkinson's disease (G20-G21)	106	0.6	---
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	97	0.6	---
15	Enterocolitis due to <i>Clostridium difficile</i> (A04.7)	61	0.4	---
...	All other causes (residual)	3,785	22.2	---
Black, female, 90-94 years				
...	All causes	13,106	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,688	28.1	---
2	Malignant neoplasms (C00-C97)	1,278	9.8	---
3	Alzheimer's disease (G30)	1,104	8.4	---
4	Cerebrovascular diseases (I60-I69)	1,077	8.2	---
5	Diabetes mellitus (E10-E14)	461	3.5	---
6	Influenza and pneumonia (J09-J18)	357	2.7	---
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	356	2.7	---
8	Essential hypertension and hypertensive renal disease (I10,I12,I15)	350	2.7	---
9	Chronic lower respiratory diseases (J40-J47)	336	2.6	---
10	Septicemia (A40-A41)	296	2.3	---
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	153	1.2	---
12	Pneumonitis due to solids and liquids (J69)	106	0.8	---
13	Parkinson's disease (G20-G21)	73	0.6	---
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	57	0.4	---
15	Atherosclerosis (I70)	50	0.4	---
...	All other causes (residual)	3,364	25.7	---

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2014

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2014

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
Black, female, 95-99 years				
...	All causes	5,937	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,734	29.2	---
2	Alzheimer's disease (G30)	563	9.5	---
3	Cerebrovascular diseases (I60-I69)	480	8.1	---
4	Malignant neoplasms (C00-C97)	422	7.1	---
5	Essential hypertension and hypertensive renal disease (I10,I12,I15)	183	3.1	---
6	Influenza and pneumonia (J09-J18)	173	2.9	---
7	Diabetes mellitus (E10-E14)	142	2.4	---
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	139	2.3	---
9	Chronic lower respiratory diseases (J40-J47)	120	2.0	---
10	Septicemia (A40-A41)	105	1.8	---
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	78	1.3	---
12	Pneumonitis due to solids and liquids (J69)	55	0.9	---
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	32	0.5	---
14	Nutritional deficiencies (E40-E64)	25	0.4	---
15	Atherosclerosis (I70)	24	0.4	---
...	All other causes (residual)	1,662	28.0	---
Black, female, 100 years and over				
...	All causes	2,157	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	644	29.9	---
2	Alzheimer's disease (G30)	215	10.0	---
3	Cerebrovascular diseases (I60-I69)	162	7.5	---
4	Malignant neoplasms (C00-C97)	91	4.2	---
5	Essential hypertension and hypertensive renal disease (I10,I12,I15)	74	3.4	---
6	Influenza and pneumonia (J09-J18)	60	2.8	---
7	Diabetes mellitus (E10-E14)	46	2.1	---
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	44	2.0	---
9	Chronic lower respiratory diseases (J40-J47)	42	1.9	---
10	Septicemia (A40-A41)	35	1.6	---
11	Pneumonitis due to solids and liquids (J69)	22	1.0	---
12	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	19	0.9	---
13	Atherosclerosis (I70)	16	0.7	---
14	Nutritional deficiencies (E40-E64)	10	0.5	---
15	Anemias (D50-D64)	8	0.4	---
...	All other causes (residual)	669	31.0	---

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2014

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2014

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, both sexes, all ages				
...	All causes	18,008	100.0	398.5
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,288	18.3	72.8
2	Malignant neoplasms (C00-C97)	3,153	17.5	69.8
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,997	11.1	44.2
4	Chronic liver disease and cirrhosis (K70,K73-K74)	951	5.3	21.0
5	Diabetes mellitus (E10-E14)	945	5.2	20.9
6	Chronic lower respiratory diseases (J40-J47)	788	4.4	17.4
7	Cerebrovascular diseases (I60-I69)	649	3.6	14.4
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	489	2.7	10.8
9	Influenza and pneumonia (J09-J18)	412	2.3	9.1
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	338	1.9	7.5
11	Alzheimer's disease (G30)	304	1.7	6.7
12	Septicemia (A40-A41)	289	1.6	6.4
13	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	263	1.5	5.8
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	176	1.0	3.9
15	Certain conditions originating in the perinatal period (P00-P96)	142	0.8	3.1
...	All other causes (residual)	3,824	21.2	84.6
American Indian, both sexes, 1-4 years				
...	All causes	95	100.0	30.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	36	37.9	11.7
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	8	8.4	*
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	6	6.3	*
4	Malignant neoplasms (C00-C97)	5	5.3	*
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	4	4.2	*
6	Septicemia (A40-A41)	2	2.1	*
6	Chronic lower respiratory diseases (J40-J47)	2	2.1	*
6	Certain conditions originating in the perinatal period (P00-P96)	2	2.1	*
9	Meningococcal infection (A39)	1	1.1	*
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1	1.1	*
9	Cerebrovascular diseases (I60-I69)	1	1.1	*
9	Influenza and pneumonia (J09-J18)	1	1.1	*
...	All other causes (residual)	26	27.4	8.4

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2014

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2014

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, both sexes, 5-9 years				
...	All causes	42	100.0	10.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	21	50.0	5.4
2	Malignant neoplasms (C00-C97)	3	7.1	*
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	3	7.1	*
4	Influenza and pneumonia (J09-J18)	2	4.8	*
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	2	4.8	*
6	Diabetes mellitus (E10-E14)	1	2.4	*
6	Cerebrovascular diseases (I60-I69)	1	2.4	*
6	Diseases of appendix (K35-K38)	1	2.4	*
...	All other causes (residual)	8	19.0	*
American Indian, both sexes, 10-14 years				
...	All causes	47	100.0	12.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	16	34.0	*
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	12	25.5	*
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	5	10.6	*
4	Malignant neoplasms (C00-C97)	4	8.5	*
5	Influenza and pneumonia (J09-J18)	2	4.3	*
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	1	2.1	*
6	Chronic lower respiratory diseases (J40-J47)	1	2.1	*
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	1	2.1	*
6	Complications of medical and surgical care (Y40-Y84,Y88)	1	2.1	*
...	All other causes (residual)	4	8.5	*
American Indian, both sexes, 15-19 years				
...	All causes	185	100.0	49.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	87	47.0	23.4
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	58	31.4	15.6
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	16	8.6	*
4	Malignant neoplasms (C00-C97)	4	2.2	*
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	3	1.6	*
5	Legal intervention (Y35,Y89.0)	3	1.6	*
7	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1	0.5	*
7	Influenza and pneumonia (J09-J18)	1	0.5	*
...	All other causes (residual)	12	6.5	*

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2014

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2014

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, both sexes, 20-24 years				
...	All causes	401	100.0	100.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	195	48.6	49.0
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	71	17.7	17.8
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	40	10.0	10.0
4	Malignant neoplasms (C00-C97)	10	2.5	*
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	10	2.5	*
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	8	2.0	*
7	Chronic lower respiratory diseases (J40-J47)	5	1.2	*
8	Pregnancy, childbirth and the puerperium (O00-O99)	4	1.0	*
9	Influenza and pneumonia (J09-J18)	3	0.7	*
9	Chronic liver disease and cirrhosis (K70,K73-K74)	3	0.7	*
11	Septicemia (A40-A41)	2	0.5	*
12	Diabetes mellitus (E10-E14)	1	0.2	*
12	Meningitis (G00,G03)	1	0.2	*
12	Cerebrovascular diseases (I60-I69)	1	0.2	*
12	Aortic aneurysm and dissection (I71)	1	0.2	*
12	Pneumonitis due to solids and liquids (J69)	1	0.2	*
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1	0.2	*
...	All other causes (residual)	44	11.0	11.0
American Indian, both sexes, 25-29 years				
...	All causes	460	100.0	128.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	201	43.7	56.1
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	74	16.1	20.7
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	37	8.0	10.3
4	Chronic liver disease and cirrhosis (K70,K73-K74)	28	6.1	7.8
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	14	3.0	*
6	Malignant neoplasms (C00-C97)	10	2.2	*
7	Diabetes mellitus (E10-E14)	7	1.5	*
8	Influenza and pneumonia (J09-J18)	5	1.1	*
9	Cerebrovascular diseases (I60-I69)	4	0.9	*
9	Pregnancy, childbirth and the puerperium (O00-O99)	4	0.9	*
11	Septicemia (A40-A41)	2	0.4	*
11	Human immunodeficiency virus (HIV) disease (B20-B24)	2	0.4	*
11	Pneumonitis due to solids and liquids (J69)	2	0.4	*
11	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	2	0.4	*
11	Legal intervention (Y35,Y89.0)	2	0.4	*
...	All other causes (residual)	66	14.3	18.4

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2014

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2014

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, both sexes, 30-34 years				
...	All causes	483	100.0	142.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	179	37.1	52.8
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	59	12.2	17.4
3	Chronic liver disease and cirrhosis (K70,K73-K74)	45	9.3	13.3
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	36	7.5	10.6
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	35	7.2	10.3
6	Malignant neoplasms (C00-C97)	14	2.9	*
7	Diabetes mellitus (E10-E14)	9	1.9	*
8	Cerebrovascular diseases (I60-I69)	6	1.2	*
8	Influenza and pneumonia (J09-J18)	6	1.2	*
10	Septicemia (A40-A41)	5	1.0	*
11	Legal intervention (Y35,Y89.0)	3	0.6	*
12	Aortic aneurysm and dissection (I71)	2	0.4	*
12	Chronic lower respiratory diseases (J40-J47)	2	0.4	*
14	Viral hepatitis (B15-B19)	1	0.2	*
14	Human immunodeficiency virus (HIV) disease (B20-B24)	1	0.2	*
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1	0.2	*
14	Pneumonitis due to solids and liquids (J69)	1	0.2	*
14	Peptic ulcer (K25-K28)	1	0.2	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1	0.2	*
14	Infections of kidney (N10-N12,N13.6,N15.1)	1	0.2	*
14	Pregnancy, childbirth and the puerperium (O00-O99)	1	0.2	*
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	1	0.2	*
14	Enterocolitis due to <i>Clostridium difficile</i> (A04.7)	1	0.2	*
...	All other causes (residual)	72	14.9	21.2
American Indian, both sexes, 35-39 years				
...	All causes	583	100.0	192.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	171	29.3	56.3
2	Chronic liver disease and cirrhosis (K70,K73-K74)	80	13.7	26.4
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	55	9.4	18.1
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	46	7.9	15.2
5	Malignant neoplasms (C00-C97)	38	6.5	12.5
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	26	4.5	8.6
7	Diabetes mellitus (E10-E14)	23	3.9	7.6
8	Influenza and pneumonia (J09-J18)	12	2.1	*
9	Septicemia (A40-A41)	11	1.9	*
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	6	1.0	*
11	Cerebrovascular diseases (I60-I69)	5	0.9	*
11	Pregnancy, childbirth and the puerperium (O00-O99)	5	0.9	*
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	4	0.7	*
14	Viral hepatitis (B15-B19)	3	0.5	*
15	Chronic lower respiratory diseases (J40-J47)	2	0.3	*
15	Legal intervention (Y35,Y89.0)	2	0.3	*
...	All other causes (residual)	94	16.1	31.0

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2014

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2014

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, both sexes, 40-44 years				
...	All causes	737	100.0	254.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	183	24.8	63.2
2	Chronic liver disease and cirrhosis (K70,K73-K74)	95	12.9	32.8
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	90	12.2	31.1
4	Malignant neoplasms (C00-C97)	59	8.0	20.4
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	56	7.6	19.3
6	Diabetes mellitus (E10-E14)	30	4.1	10.4
7	Influenza and pneumonia (J09-J18)	20	2.7	6.9
8	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	18	2.4	*
9	Septicemia (A40-A41)	14	1.9	*
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	7	0.9	*
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	6	0.8	*
11	Chronic lower respiratory diseases (J40-J47)	6	0.8	*
13	Viral hepatitis (B15-B19)	5	0.7	*
13	Human immunodeficiency virus (HIV) disease (B20-B24)	5	0.7	*
13	Cerebrovascular diseases (I60-I69)	5	0.7	*
...	All other causes (residual)	138	18.7	47.6
American Indian, both sexes, 45-49 years				
...	All causes	988	100.0	359.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	190	19.2	69.0
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	145	14.7	52.7
3	Chronic liver disease and cirrhosis (K70,K73-K74)	120	12.1	43.6
4	Malignant neoplasms (C00-C97)	105	10.6	38.2
5	Diabetes mellitus (E10-E14)	54	5.5	19.6
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	30	3.0	10.9
7	Cerebrovascular diseases (I60-I69)	27	2.7	9.8
8	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	23	2.3	8.4
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	20	2.0	7.3
10	Septicemia (A40-A41)	16	1.6	*
10	Influenza and pneumonia (J09-J18)	16	1.6	*
12	Viral hepatitis (B15-B19)	13	1.3	*
12	Human immunodeficiency virus (HIV) disease (B20-B24)	13	1.3	*
14	Chronic lower respiratory diseases (J40-J47)	12	1.2	*
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	5	0.5	*
...	All other causes (residual)	199	20.1	72.3

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2014

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2014

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, both sexes, 50-54 years				
...	All causes	1,388	100.0	502.7
1	Malignant neoplasms (C00-C97)	242	17.4	87.6
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	231	16.6	83.7
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	186	13.4	67.4
4	Chronic liver disease and cirrhosis (K70,K73-K74)	160	11.5	57.9
5	Diabetes mellitus (E10-E14)	71	5.1	25.7
6	Influenza and pneumonia (J09-J18)	38	2.7	13.8
7	Chronic lower respiratory diseases (J40-J47)	35	2.5	12.7
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	31	2.2	11.2
9	Cerebrovascular diseases (I60-I69)	27	1.9	9.8
10	Septicemia (A40-A41)	24	1.7	8.7
11	Viral hepatitis (B15-B19)	21	1.5	7.6
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	20	1.4	7.2
13	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	18	1.3	*
14	Human immunodeficiency virus (HIV) disease (B20-B24)	11	0.8	*
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	8	0.6	*
...	All other causes (residual)	265	19.1	96.0
American Indian, both sexes, 55-59 years				
...	All causes	1,585	100.0	665.4
1	Malignant neoplasms (C00-C97)	352	22.2	147.8
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	276	17.4	115.9
3	Chronic liver disease and cirrhosis (K70,K73-K74)	156	9.8	65.5
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	135	8.5	56.7
5	Diabetes mellitus (E10-E14)	99	6.2	41.6
6	Chronic lower respiratory diseases (J40-J47)	58	3.7	24.3
7	Cerebrovascular diseases (I60-I69)	46	2.9	19.3
8	Viral hepatitis (B15-B19)	34	2.1	14.3
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	28	1.8	11.8
10	Septicemia (A40-A41)	25	1.6	10.5
11	Influenza and pneumonia (J09-J18)	24	1.5	10.1
12	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	19	1.2	*
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	17	1.1	*
14	Human immunodeficiency virus (HIV) disease (B20-B24)	10	0.6	*
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	10	0.6	*
...	All other causes (residual)	296	18.7	124.3

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2014

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2014

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, both sexes, 60-64 years				
...	All causes	1,717	100.0	949.3
1	Malignant neoplasms (C00-C97)	397	23.1	219.5
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	358	20.9	197.9
3	Diabetes mellitus (E10-E14)	137	8.0	75.7
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	115	6.7	63.6
5	Chronic liver disease and cirrhosis (K70,K73-K74)	104	6.1	57.5
6	Chronic lower respiratory diseases (J40-J47)	76	4.4	42.0
7	Cerebrovascular diseases (I60-I69)	61	3.6	33.7
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	36	2.1	19.9
9	Influenza and pneumonia (J09-J18)	33	1.9	18.2
10	Septicemia (A40-A41)	31	1.8	17.1
11	Viral hepatitis (B15-B19)	24	1.4	13.3
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	19	1.1	*
13	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	11	0.6	*
14	Pneumonitis due to solids and liquids (J69)	10	0.6	*
15	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	7	0.4	*
...	All other causes (residual)	298	17.4	164.8
American Indian, both sexes, 65-69 years				
...	All causes	1,723	100.0	1,310.6
1	Malignant neoplasms (C00-C97)	477	27.7	362.8
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	382	22.2	290.6
3	Diabetes mellitus (E10-E14)	129	7.5	98.1
4	Chronic lower respiratory diseases (J40-J47)	104	6.0	79.1
5	Chronic liver disease and cirrhosis (K70,K73-K74)	75	4.4	57.0
6	Cerebrovascular diseases (I60-I69)	65	3.8	49.4
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	65	3.8	49.4
8	Septicemia (A40-A41)	33	1.9	25.1
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	33	1.9	25.1
10	Influenza and pneumonia (J09-J18)	31	1.8	23.6
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	18	1.0	*
12	Viral hepatitis (B15-B19)	11	0.6	*
13	Enterocolitis due to <i>Clostridium difficile</i> (A04.7)	9	0.5	*
14	Aortic aneurysm and dissection (I71)	7	0.4	*
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	6	0.3	*
15	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	6	0.3	*
...	All other causes (residual)	272	15.8	206.9

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2014

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2014

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, both sexes, 70-74 years				
...	All causes	1,707	100.0	2,012.4
1	Malignant neoplasms (C00-C97)	465	27.2	548.2
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	378	22.1	445.6
3	Chronic lower respiratory diseases (J40-J47)	128	7.5	150.9
4	Diabetes mellitus (E10-E14)	112	6.6	132.0
5	Cerebrovascular diseases (I60-I69)	71	4.2	83.7
6	Chronic liver disease and cirrhosis (K70,K73-K74)	46	2.7	54.2
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	46	2.7	54.2
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	41	2.4	48.3
9	Influenza and pneumonia (J09-J18)	37	2.2	43.6
10	Septicemia (A40-A41)	28	1.6	33.0
11	Alzheimer's disease (G30)	22	1.3	25.9
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	18	1.1	*
13	Pneumonitis due to solids and liquids (J69)	15	0.9	*
14	Parkinson's disease (G20-G21)	12	0.7	*
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	9	0.5	*
...	All other causes (residual)	279	16.3	328.9
American Indian, both sexes, 75-79 years				
...	All causes	1,673	100.0	3,065.6
1	Malignant neoplasms (C00-C97)	399	23.8	731.1
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	349	20.9	639.5
3	Chronic lower respiratory diseases (J40-J47)	145	8.7	265.7
4	Diabetes mellitus (E10-E14)	101	6.0	185.1
5	Cerebrovascular diseases (I60-I69)	90	5.4	164.9
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	50	3.0	91.6
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	45	2.7	82.5
8	Influenza and pneumonia (J09-J18)	40	2.4	73.3
9	Septicemia (A40-A41)	36	2.2	66.0
10	Alzheimer's disease (G30)	34	2.0	62.3
11	Parkinson's disease (G20-G21)	20	1.2	36.6
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	19	1.1	*
12	Chronic liver disease and cirrhosis (K70,K73-K74)	19	1.1	*
14	Pneumonitis due to solids and liquids (J69)	18	1.1	*
15	Aortic aneurysm and dissection (I71)	9	0.5	*
...	All other causes (residual)	299	17.9	547.9

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2014

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2014

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, both sexes, 80-84 years				
...	All causes	1,476	100.0	4,480.1
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	346	23.4	1,050.2
2	Malignant neoplasms (C00-C97)	287	19.4	871.1
3	Diabetes mellitus (E10-E14)	79	5.4	239.8
4	Chronic lower respiratory diseases (J40-J47)	76	5.1	230.7
5	Cerebrovascular diseases (I60-I69)	74	5.0	224.6
6	Alzheimer's disease (G30)	70	4.7	212.5
7	Influenza and pneumonia (J09-J18)	49	3.3	148.7
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	42	2.8	127.5
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	36	2.4	109.3
10	Septicemia (A40-A41)	23	1.6	69.8
11	Parkinson's disease (G20-G21)	19	1.3	*
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	19	1.3	*
13	Pneumonitis due to solids and liquids (J69)	14	0.9	*
14	Enterocolitis due to <i>Clostridium difficile</i> (A04.7)	12	0.8	*
15	Aortic aneurysm and dissection (I71)	11	0.7	*
...	All other causes (residual)	319	21.6	968.3
American Indian, both sexes, 85-89 years				
...	All causes	1,226	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	316	25.8	---
2	Malignant neoplasms (C00-C97)	189	15.4	---
3	Alzheimer's disease (G30)	78	6.4	---
3	Cerebrovascular diseases (I60-I69)	78	6.4	---
5	Chronic lower respiratory diseases (J40-J47)	74	6.0	---
6	Diabetes mellitus (E10-E14)	51	4.2	---
7	Influenza and pneumonia (J09-J18)	42	3.4	---
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	34	2.8	---
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	31	2.5	---
10	Septicemia (A40-A41)	19	1.5	---
11	Parkinson's disease (G20-G21)	15	1.2	---
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	15	1.2	---
13	Pneumonitis due to solids and liquids (J69)	12	1.0	---
14	Chronic liver disease and cirrhosis (K70,K73-K74)	8	0.7	---
15	Nutritional deficiencies (E40-E64)	7	0.6	---
15	Atherosclerosis (I70)	7	0.6	---
...	All other causes (residual)	250	20.4	---

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2014

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2014

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, both sexes, 90-94 years				
...	All causes	800	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	201	25.1	---
2	Malignant neoplasms (C00-C97)	77	9.6	---
3	Alzheimer's disease (G30)	63	7.9	---
4	Cerebrovascular diseases (I60-I69)	59	7.4	---
5	Chronic lower respiratory diseases (J40-J47)	45	5.6	---
6	Diabetes mellitus (E10-E14)	37	4.6	---
7	Influenza and pneumonia (J09-J18)	27	3.4	---
8	Essential hypertension and hypertensive renal disease (I10,I12,I15)	19	2.4	---
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	19	2.4	---
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	16	2.0	---
11	Parkinson's disease (G20-G21)	11	1.4	---
12	Septicemia (A40-A41)	10	1.3	---
13	Pneumonitis due to solids and liquids (J69)	6	0.8	---
14	Nutritional deficiencies (E40-E64)	5	0.6	---
15	Anemias (D50-D64)	4	0.5	---
15	Hernia (K40-K46)	4	0.5	---
...	All other causes (residual)	197	24.6	---
American Indian, both sexes, 95-99 years				
...	All causes	261	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	72	27.6	---
2	Alzheimer's disease (G30)	24	9.2	---
3	Cerebrovascular diseases (I60-I69)	22	8.4	---
4	Influenza and pneumonia (J09-J18)	17	6.5	---
5	Chronic lower respiratory diseases (J40-J47)	13	5.0	---
6	Malignant neoplasms (C00-C97)	11	4.2	---
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	9	3.4	---
8	Essential hypertension and hypertensive renal disease (I10,I12,I15)	8	3.1	---
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	5	1.9	---
10	Septicemia (A40-A41)	4	1.5	---
11	Diabetes mellitus (E10-E14)	2	0.8	---
11	Enterocolitis due to <i>Clostridium difficile</i> (A04.7)	2	0.8	---
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1	0.4	---
13	Nutritional deficiencies (E40-E64)	1	0.4	---
13	Parkinson's disease (G20-G21)	1	0.4	---
13	Pneumonitis due to solids and liquids (J69)	1	0.4	---
13	Cholelithiasis and other disorders of gallbladder (K80-K82)	1	0.4	---
13	Hyperplasia of prostate (N40)	1	0.4	---
13	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	1	0.4	---
...	All other causes (residual)	65	24.9	---

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2014

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2014

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, both sexes, 100 years and over				
...	All causes	71	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	20	28.2	---
2	Malignant neoplasms (C00-C97)	5	7.0	---
3	Essential hypertension and hypertensive renal disease (I10,I12,I15)	4	5.6	---
3	Cerebrovascular diseases (I60-I69)	4	5.6	---
5	Alzheimer's disease (G30)	3	4.2	---
5	Pneumonitis due to solids and liquids (J69)	3	4.2	---
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3	4.2	---
8	Diabetes mellitus (E10-E14)	2	2.8	---
8	Nutritional deficiencies (E40-E64)	2	2.8	---
8	Chronic lower respiratory diseases (J40-J47)	2	2.8	---
11	Septicemia (A40-A41)	1	1.4	---
11	Anemias (D50-D64)	1	1.4	---
11	Influenza and pneumonia (J09-J18)	1	1.4	---
11	Enterocolitis due to <i>Clostridium difficile</i> (A04.7)	1	1.4	---
...	All other causes (residual)	19	26.8	---
American Indian, male, all ages				
...	All causes	9,829	100.0	433.2
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,915	19.5	84.4
2	Malignant neoplasms (C00-C97)	1,727	17.6	76.1
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,332	13.6	58.7
4	Diabetes mellitus (E10-E14)	504	5.1	22.2
5	Chronic liver disease and cirrhosis (K70,K73-K74)	481	4.9	21.2
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	363	3.7	16.0
7	Chronic lower respiratory diseases (J40-J47)	360	3.7	15.9
8	Cerebrovascular diseases (I60-I69)	296	3.0	13.0
9	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	208	2.1	9.2
10	Influenza and pneumonia (J09-J18)	195	2.0	8.6
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	163	1.7	7.2
12	Septicemia (A40-A41)	123	1.3	5.4
13	Viral hepatitis (B15-B19)	86	0.9	3.8
14	Alzheimer's disease (G30)	80	0.8	3.5
14	Certain conditions originating in the perinatal period (P00-P96)	80	0.8	3.5
...	All other causes (residual)	1,916	19.5	84.4

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2014

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2014

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, male, 1-4 years				
...	All causes	64	100.0	41.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	24	37.5	15.4
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	8	12.5	*
3	Malignant neoplasms (C00-C97)	5	7.8	*
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	3	4.7	*
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	3	4.7	*
6	Chronic lower respiratory diseases (J40-J47)	2	3.1	*
7	Meningococcal infection (A39)	1	1.6	*
7	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1	1.6	*
7	Cerebrovascular diseases (I60-I69)	1	1.6	*
7	Certain conditions originating in the perinatal period (P00-P96)	1	1.6	*
...	All other causes (residual)	15	23.4	*
American Indian, male, 5-9 years				
...	All causes	25	100.0	12.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	13	52.0	*
2	Malignant neoplasms (C00-C97)	2	8.0	*
3	Cerebrovascular diseases (I60-I69)	1	4.0	*
3	Influenza and pneumonia (J09-J18)	1	4.0	*
3	Diseases of appendix (K35-K38)	1	4.0	*
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	1	4.0	*
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1	4.0	*
...	All other causes (residual)	5	20.0	*
American Indian, male, 10-14 years				
...	All causes	23	100.0	12.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	10	43.5	*
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	4	17.4	*
3	Malignant neoplasms (C00-C97)	2	8.7	*
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	2	8.7	*
5	Chronic lower respiratory diseases (J40-J47)	1	4.3	*
5	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	1	4.3	*
5	Complications of medical and surgical care (Y40-Y84,Y88)	1	4.3	*
...	All other causes (residual)	2	8.7	*

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2014

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2014

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, male, 15-19 years				
...	All causes	121	100.0	63.9
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	54	44.6	28.5
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	37	30.6	19.5
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	14	11.6	*
4	Malignant neoplasms (C00-C97)	3	2.5	*
4	Legal intervention (Y35,Y89.0)	3	2.5	*
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	1	0.8	*
6	Influenza and pneumonia (J09-J18)	1	0.8	*
...	All other causes (residual)	8	6.6	*
American Indian, male, 20-24 years				
...	All causes	289	100.0	140.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	136	47.1	66.1
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	56	19.4	27.2
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	34	11.8	16.5
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	9	3.1	*
5	Malignant neoplasms (C00-C97)	7	2.4	*
6	Influenza and pneumonia (J09-J18)	3	1.0	*
6	Chronic lower respiratory diseases (J40-J47)	3	1.0	*
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	3	1.0	*
9	Chronic liver disease and cirrhosis (K70,K73-K74)	2	0.7	*
10	Diabetes mellitus (E10-E14)	1	0.3	*
10	Meningitis (G00,G03)	1	0.3	*
10	Cerebrovascular diseases (I60-I69)	1	0.3	*
10	Aortic aneurysm and dissection (I71)	1	0.3	*
...	All other causes (residual)	32	11.1	15.6
American Indian, male, 25-29 years				
...	All causes	322	100.0	172.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	143	44.4	76.5
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	60	18.6	32.1
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	26	8.1	13.9
4	Chronic liver disease and cirrhosis (K70,K73-K74)	15	4.7	*
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	11	3.4	*
6	Malignant neoplasms (C00-C97)	10	3.1	*
7	Diabetes mellitus (E10-E14)	3	0.9	*
7	Cerebrovascular diseases (I60-I69)	3	0.9	*
9	Human immunodeficiency virus (HIV) disease (B20-B24)	2	0.6	*
9	Influenza and pneumonia (J09-J18)	2	0.6	*
9	Pneumonitis due to solids and liquids (J69)	2	0.6	*
9	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	2	0.6	*
9	Legal intervention (Y35,Y89.0)	2	0.6	*
14	Anemias (D50-D64)	1	0.3	*
...	All other causes (residual)	40	12.4	21.4

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2014

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2014

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, male, 30-34 years				
...	All causes	326	100.0	186.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	129	39.6	73.8
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	40	12.3	22.9
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	30	9.2	17.2
4	Chronic liver disease and cirrhosis (K70,K73-K74)	26	8.0	14.9
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	24	7.4	13.7
6	Malignant neoplasms (C00-C97)	7	2.1	*
6	Diabetes mellitus (E10-E14)	7	2.1	*
8	Influenza and pneumonia (J09-J18)	4	1.2	*
9	Septicemia (A40-A41)	3	0.9	*
9	Cerebrovascular diseases (I60-I69)	3	0.9	*
9	Legal intervention (Y35,Y89.0)	3	0.9	*
12	Viral hepatitis (B15-B19)	1	0.3	*
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1	0.3	*
12	Aortic aneurysm and dissection (I71)	1	0.3	*
12	Pneumonitis due to solids and liquids (J69)	1	0.3	*
12	Peptic ulcer (K25-K28)	1	0.3	*
12	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	1	0.3	*
12	Enterocolitis due to <i>Clostridium difficile</i> (A04.7)	1	0.3	*
...	All other causes (residual)	43	13.2	24.6
American Indian, male, 35-39 years				
...	All causes	351	100.0	226.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	117	33.3	75.3
2	Chronic liver disease and cirrhosis (K70,K73-K74)	42	12.0	27.0
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	34	9.7	21.9
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	33	9.4	21.2
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	21	6.0	13.5
6	Malignant neoplasms (C00-C97)	17	4.8	*
7	Diabetes mellitus (E10-E14)	15	4.3	*
8	Influenza and pneumonia (J09-J18)	8	2.3	*
9	Septicemia (A40-A41)	5	1.4	*
9	Cerebrovascular diseases (I60-I69)	5	1.4	*
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	5	1.4	*
12	Viral hepatitis (B15-B19)	2	0.6	*
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	2	0.6	*
12	Legal intervention (Y35,Y89.0)	2	0.6	*
15	Meningitis (G00,G03)	1	0.3	*
...	All other causes (residual)	42	12.0	27.0

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2014

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2014

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, male, 40-44 years				
...	All causes	448	100.0	305.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	116	25.9	79.0
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	65	14.5	44.3
3	Chronic liver disease and cirrhosis (K70,K73-K74)	42	9.4	28.6
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	41	9.2	27.9
5	Malignant neoplasms (C00-C97)	29	6.5	19.8
6	Diabetes mellitus (E10-E14)	17	3.8	*
7	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	13	2.9	*
8	Septicemia (A40-A41)	8	1.8	*
8	Influenza and pneumonia (J09-J18)	8	1.8	*
10	Human immunodeficiency virus (HIV) disease (B20-B24)	4	0.9	*
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	3	0.7	*
11	Chronic lower respiratory diseases (J40-J47)	3	0.7	*
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3	0.7	*
11	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	3	0.7	*
15	Viral hepatitis (B15-B19)	2	0.4	*
15	Cerebrovascular diseases (I60-I69)	2	0.4	*
15	Complications of medical and surgical care (Y40-Y84,Y88)	2	0.4	*
...	All other causes (residual)	87	19.4	59.3
American Indian, male, 45-49 years				
...	All causes	578	100.0	417.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	132	22.8	95.4
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	100	17.3	72.3
3	Chronic liver disease and cirrhosis (K70,K73-K74)	54	9.3	39.0
4	Malignant neoplasms (C00-C97)	51	8.8	36.9
5	Diabetes mellitus (E10-E14)	29	5.0	21.0
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	25	4.3	18.1
7	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	16	2.8	*
8	Viral hepatitis (B15-B19)	12	2.1	*
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	12	2.1	*
10	Human immunodeficiency virus (HIV) disease (B20-B24)	10	1.7	*
10	Cerebrovascular diseases (I60-I69)	10	1.7	*
12	Chronic lower respiratory diseases (J40-J47)	7	1.2	*
13	Septicemia (A40-A41)	6	1.0	*
13	Influenza and pneumonia (J09-J18)	6	1.0	*
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	3	0.5	*
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	3	0.5	*
...	All other causes (residual)	102	17.6	73.7

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2014

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2014

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, male, 50-54 years				
...	All causes	816	100.0	600.9
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	163	20.0	120.0
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	129	15.8	95.0
3	Malignant neoplasms (C00-C97)	110	13.5	81.0
4	Chronic liver disease and cirrhosis (K70,K73-K74)	83	10.2	61.1
5	Diabetes mellitus (E10-E14)	44	5.4	32.4
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	24	2.9	17.7
7	Influenza and pneumonia (J09-J18)	22	2.7	16.2
8	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	16	2.0	*
9	Viral hepatitis (B15-B19)	15	1.8	*
10	Chronic lower respiratory diseases (J40-J47)	14	1.7	*
11	Septicemia (A40-A41)	13	1.6	*
11	Cerebrovascular diseases (I60-I69)	13	1.6	*
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	12	1.5	*
14	Human immunodeficiency virus (HIV) disease (B20-B24)	6	0.7	*
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	6	0.7	*
...	All other causes (residual)	146	17.9	107.5
American Indian, male, 55-59 years				
...	All causes	948	100.0	824.8
1	Malignant neoplasms (C00-C97)	216	22.8	187.9
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	187	19.7	162.7
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	88	9.3	76.6
4	Chronic liver disease and cirrhosis (K70,K73-K74)	86	9.1	74.8
5	Diabetes mellitus (E10-E14)	60	6.3	52.2
6	Cerebrovascular diseases (I60-I69)	27	2.8	23.5
7	Chronic lower respiratory diseases (J40-J47)	24	2.5	20.9
8	Viral hepatitis (B15-B19)	23	2.4	20.0
9	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	14	1.5	*
10	Influenza and pneumonia (J09-J18)	13	1.4	*
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	12	1.3	*
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	12	1.3	*
13	Septicemia (A40-A41)	9	0.9	*
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	8	0.8	*
15	Human immunodeficiency virus (HIV) disease (B20-B24)	6	0.6	*
...	All other causes (residual)	163	17.2	141.8

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2014

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2014

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, male, 60-64 years				
...	All causes	1,033	100.0	1,197.8
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	250	24.2	289.9
2	Malignant neoplasms (C00-C97)	229	22.2	265.5
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	83	8.0	96.2
4	Diabetes mellitus (E10-E14)	78	7.6	90.4
5	Chronic liver disease and cirrhosis (K70,K73-K74)	54	5.2	62.6
6	Chronic lower respiratory diseases (J40-J47)	39	3.8	45.2
7	Cerebrovascular diseases (I60-I69)	33	3.2	38.3
8	Viral hepatitis (B15-B19)	20	1.9	23.2
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	18	1.7	*
10	Septicemia (A40-A41)	16	1.5	*
10	Influenza and pneumonia (J09-J18)	16	1.5	*
12	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	9	0.9	*
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	6	0.6	*
13	Pneumonitis due to solids and liquids (J69)	6	0.6	*
15	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	5	0.5	*
...	All other causes (residual)	171	16.6	198.3
American Indian, male, 65-69 years				
...	All causes	968	100.0	1,547.8
1	Malignant neoplasms (C00-C97)	270	27.9	431.7
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	230	23.8	367.8
3	Diabetes mellitus (E10-E14)	69	7.1	110.3
4	Chronic lower respiratory diseases (J40-J47)	52	5.4	83.1
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	42	4.3	67.2
6	Cerebrovascular diseases (I60-I69)	37	3.8	59.2
6	Chronic liver disease and cirrhosis (K70,K73-K74)	37	3.8	59.2
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	18	1.9	*
9	Septicemia (A40-A41)	17	1.8	*
10	Influenza and pneumonia (J09-J18)	12	1.2	*
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	9	0.9	*
12	Viral hepatitis (B15-B19)	7	0.7	*
13	Aortic aneurysm and dissection (I71)	5	0.5	*
13	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	5	0.5	*
13	Enterocolitis due to <i>Clostridium difficile</i> (A04.7)	5	0.5	*
...	All other causes (residual)	153	15.8	244.6

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2014

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2014

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, male, 70-74 years				
...	All causes	895	100.0	2,280.1
1	Malignant neoplasms (C00-C97)	260	29.1	662.4
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	222	24.8	565.6
3	Diabetes mellitus (E10-E14)	59	6.6	150.3
4	Chronic lower respiratory diseases (J40-J47)	56	6.3	142.7
5	Cerebrovascular diseases (I60-I69)	39	4.4	99.4
6	Chronic liver disease and cirrhosis (K70,K73-K74)	23	2.6	58.6
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	22	2.5	56.0
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	20	2.2	51.0
9	Influenza and pneumonia (J09-J18)	19	2.1	*
10	Pneumonitis due to solids and liquids (J69)	10	1.1	*
11	Alzheimer's disease (G30)	9	1.0	*
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	9	1.0	*
13	Septicemia (A40-A41)	8	0.9	*
13	Parkinson's disease (G20-G21)	8	0.9	*
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	6	0.7	*
15	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	6	0.7	*
...	All other causes (residual)	119	13.3	303.2
American Indian, male, 75-79 years				
...	All causes	858	100.0	3,583.5
1	Malignant neoplasms (C00-C97)	228	26.6	952.3
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	202	23.5	843.7
3	Chronic lower respiratory diseases (J40-J47)	69	8.0	288.2
4	Diabetes mellitus (E10-E14)	46	5.4	192.1
5	Cerebrovascular diseases (I60-I69)	41	4.8	171.2
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	22	2.6	91.9
7	Influenza and pneumonia (J09-J18)	20	2.3	83.5
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	19	2.2	*
9	Septicemia (A40-A41)	13	1.5	*
10	Parkinson's disease (G20-G21)	10	1.2	*
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	9	1.0	*
12	Alzheimer's disease (G30)	8	0.9	*
12	Pneumonitis due to solids and liquids (J69)	8	0.9	*
12	Chronic liver disease and cirrhosis (K70,K73-K74)	8	0.9	*
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	4	0.5	*
15	Peptic ulcer (K25-K28)	4	0.5	*
15	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	4	0.5	*
15	Enterocolitis due to <i>Clostridium difficile</i> (A04.7)	4	0.5	*
...	All other causes (residual)	139	16.2	580.5

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2014

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2014

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, male, 80-84 years				
...	All causes	671	100.0	5,019.1
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	163	24.3	1,219.2
2	Malignant neoplasms (C00-C97)	151	22.5	1,129.5
3	Diabetes mellitus (E10-E14)	39	5.8	291.7
4	Chronic lower respiratory diseases (J40-J47)	35	5.2	261.8
5	Cerebrovascular diseases (I60-I69)	31	4.6	231.9
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	25	3.7	187.0
7	Alzheimer's disease (G30)	24	3.6	179.5
8	Influenza and pneumonia (J09-J18)	22	3.3	164.6
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	19	2.8	*
10	Parkinson's disease (G20-G21)	9	1.3	*
10	Pneumonitis due to solids and liquids (J69)	9	1.3	*
12	Septicemia (A40-A41)	8	1.2	*
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	7	1.0	*
14	Aortic aneurysm and dissection (I71)	4	0.6	*
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	3	0.4	*
15	Nutritional deficiencies (E40-E64)	3	0.4	*
15	Chronic liver disease and cirrhosis (K70,K73-K74)	3	0.4	*
15	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	3	0.4	*
15	Enterocolitis due to <i>Clostridium difficile</i> (A04.7)	3	0.4	*
...	All other causes (residual)	110	16.4	822.8
American Indian, male, 85-89 years				
...	All causes	530	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	147	27.7	---
2	Malignant neoplasms (C00-C97)	93	17.5	---
3	Chronic lower respiratory diseases (J40-J47)	36	6.8	---
4	Alzheimer's disease (G30)	26	4.9	---
5	Cerebrovascular diseases (I60-I69)	25	4.7	---
6	Diabetes mellitus (E10-E14)	21	4.0	---
7	Influenza and pneumonia (J09-J18)	19	3.6	---
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	18	3.4	---
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	14	2.6	---
10	Septicemia (A40-A41)	12	2.3	---
11	Parkinson's disease (G20-G21)	11	2.1	---
12	Pneumonitis due to solids and liquids (J69)	8	1.5	---
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	6	1.1	---
14	Atherosclerosis (I70)	5	0.9	---
14	Chronic liver disease and cirrhosis (K70,K73-K74)	5	0.9	---
...	All other causes (residual)	84	15.8	---

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2014

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2014

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, male, 90-94 years				
...	All causes	280	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	79	28.2	---
2	Malignant neoplasms (C00-C97)	33	11.8	---
3	Cerebrovascular diseases (I60-I69)	18	6.4	---
4	Diabetes mellitus (E10-E14)	16	5.7	---
5	Chronic lower respiratory diseases (J40-J47)	15	5.4	---
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	13	4.6	---
7	Alzheimer's disease (G30)	11	3.9	---
8	Influenza and pneumonia (J09-J18)	9	3.2	---
9	Parkinson's disease (G20-G21)	6	2.1	---
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	6	2.1	---
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	5	1.8	---
12	Septicemia (A40-A41)	2	0.7	---
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2	0.7	---
12	Nutritional deficiencies (E40-E64)	2	0.7	---
12	Pneumoconioses and chemical effects (J60-J66,J68)	2	0.7	---
12	Enterocolitis due to <i>Clostridium difficile</i> (A04.7)	2	0.7	---
...	All other causes (residual)	59	21.1	---
American Indian, male, 95-99 years				
...	All causes	69	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	19	27.5	---
2	Influenza and pneumonia (J09-J18)	8	11.6	---
3	Cerebrovascular diseases (I60-I69)	5	7.2	---
4	Chronic lower respiratory diseases (J40-J47)	4	5.8	---
5	Malignant neoplasms (C00-C97)	3	4.3	---
6	Septicemia (A40-A41)	2	2.9	---
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2	2.9	---
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2	2.9	---
6	Enterocolitis due to <i>Clostridium difficile</i> (A04.7)	2	2.9	---
10	Alzheimer's disease (G30)	1	1.4	---
10	Hyperplasia of prostate (N40)	1	1.4	---
10	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	1	1.4	---
...	All other causes (residual)	19	27.5	---

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2014

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2014

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, male, 100 years and over				
...	All causes	12	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	5	41.7	---
2	Malignant neoplasms (C00-C97)	1	8.3	---
2	Nutritional deficiencies (E40-E64)	1	8.3	---
2	Pneumonitis due to solids and liquids (J69)	1	8.3	---
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1	8.3	---
...	All other causes (residual)	3	25.0	---
American Indian, female, all ages				
...	All causes	8,179	100.0	363.5
1	Malignant neoplasms (C00-C97)	1,426	17.4	63.4
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,373	16.8	61.0
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	665	8.1	29.6
4	Chronic liver disease and cirrhosis (K70,K73-K74)	470	5.7	20.9
5	Diabetes mellitus (E10-E14)	441	5.4	19.6
6	Chronic lower respiratory diseases (J40-J47)	428	5.2	19.0
7	Cerebrovascular diseases (I60-I69)	353	4.3	15.7
8	Alzheimer's disease (G30)	224	2.7	10.0
9	Influenza and pneumonia (J09-J18)	217	2.7	9.6
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	175	2.1	7.8
11	Septicemia (A40-A41)	166	2.0	7.4
12	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	126	1.5	5.6
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	101	1.2	4.5
14	Certain conditions originating in the perinatal period (P00-P96)	62	0.8	2.8
15	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	55	0.7	2.4
...	All other causes (residual)	1,897	23.2	84.3
American Indian, female, 1-4 years				
...	All causes	31	100.0	20.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	12	38.7	*
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	3	9.7	*
3	Septicemia (A40-A41)	2	6.5	*
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	1	3.2	*
4	Influenza and pneumonia (J09-J18)	1	3.2	*
4	Certain conditions originating in the perinatal period (P00-P96)	1	3.2	*
...	All other causes (residual)	11	35.5	*

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2014

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2014

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, female, 5-9 years				
...	All causes	17	100.0	*
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	8	47.1	*
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	2	11.8	*
3	Malignant neoplasms (C00-C97)	1	5.9	*
3	Diabetes mellitus (E10-E14)	1	5.9	*
3	Influenza and pneumonia (J09-J18)	1	5.9	*
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1	5.9	*
...	All other causes (residual)	3	17.6	*
American Indian, female, 10-14 years				
...	All causes	24	100.0	12.9
1	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	8	33.3	*
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	6	25.0	*
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	3	12.5	*
4	Malignant neoplasms (C00-C97)	2	8.3	*
4	Influenza and pneumonia (J09-J18)	2	8.3	*
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	1	4.2	*
...	All other causes (residual)	2	8.3	*
American Indian, female, 15-19 years				
...	All causes	64	100.0	35.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	33	51.6	18.0
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	21	32.8	11.5
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	2	3.1	*
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	2	3.1	*
5	Malignant neoplasms (C00-C97)	1	1.6	*
5	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1	1.6	*
...	All other causes (residual)	4	6.3	*
American Indian, female, 20-24 years				
...	All causes	112	100.0	58.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	59	52.7	30.6
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	15	13.4	*
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	6	5.4	*
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	5	4.5	*
5	Pregnancy, childbirth and the puerperium (O00-O99)	4	3.6	*
6	Malignant neoplasms (C00-C97)	3	2.7	*
7	Septicemia (A40-A41)	2	1.8	*
7	Chronic lower respiratory diseases (J40-J47)	2	1.8	*
9	Diseases of heart (I00-I09,I11,I13,I20-I51)	1	0.9	*
9	Pneumonitis due to solids and liquids (J69)	1	0.9	*
9	Chronic liver disease and cirrhosis (K70,K73-K74)	1	0.9	*
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1	0.9	*
...	All other causes (residual)	12	10.7	*

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2014

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2014

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, female, 25-29 years				
...	All causes	138	100.0	80.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	58	42.0	33.9
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	14	10.1	*
3	Chronic liver disease and cirrhosis (K70,K73-K74)	13	9.4	*
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	11	8.0	*
5	Diabetes mellitus (E10-E14)	4	2.9	*
5	Pregnancy, childbirth and the puerperium (O00-O99)	4	2.9	*
7	Diseases of heart (I00-I09,I11,I13,I20-I51)	3	2.2	*
7	Influenza and pneumonia (J09-J18)	3	2.2	*
9	Septicemia (A40-A41)	2	1.4	*
10	Viral hepatitis (B15-B19)	1	0.7	*
10	Cerebrovascular diseases (I60-I69)	1	0.7	*
10	Chronic lower respiratory diseases (J40-J47)	1	0.7	*
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1	0.7	*
...	All other causes (residual)	22	15.9	12.9
American Indian, female, 30-34 years				
...	All causes	157	100.0	95.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	50	31.8	30.5
2	Chronic liver disease and cirrhosis (K70,K73-K74)	19	12.1	*
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	19	12.1	*
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	12	7.6	*
5	Malignant neoplasms (C00-C97)	7	4.5	*
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	5	3.2	*
7	Cerebrovascular diseases (I60-I69)	3	1.9	*
8	Septicemia (A40-A41)	2	1.3	*
8	Diabetes mellitus (E10-E14)	2	1.3	*
8	Influenza and pneumonia (J09-J18)	2	1.3	*
8	Chronic lower respiratory diseases (J40-J47)	2	1.3	*
12	Human immunodeficiency virus (HIV) disease (B20-B24)	1	0.6	*
12	Aortic aneurysm and dissection (I71)	1	0.6	*
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1	0.6	*
12	Infections of kidney (N10-N12,N13.6,N15.1)	1	0.6	*
12	Pregnancy, childbirth and the puerperium (O00-O99)	1	0.6	*
...	All other causes (residual)	29	18.5	17.7

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2014

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2014

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, female, 35-39 years				
...	All causes	232	100.0	156.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	54	23.3	36.4
2	Chronic liver disease and cirrhosis (K70,K73-K74)	38	16.4	25.6
3	Malignant neoplasms (C00-C97)	21	9.1	14.2
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	21	9.1	14.2
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	13	5.6	*
6	Diabetes mellitus (E10-E14)	8	3.4	*
7	Septicemia (A40-A41)	6	2.6	*
8	Pregnancy, childbirth and the puerperium (O00-O99)	5	2.2	*
8	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	5	2.2	*
10	Influenza and pneumonia (J09-J18)	4	1.7	*
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	2	0.9	*
11	Chronic lower respiratory diseases (J40-J47)	2	0.9	*
13	Viral hepatitis (B15-B19)	1	0.4	*
13	Pneumonitis due to solids and liquids (J69)	1	0.4	*
13	Cholelithiasis and other disorders of gallbladder (K80-K82)	1	0.4	*
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1	0.4	*
13	Infections of kidney (N10-N12,N13.6,N15.1)	1	0.4	*
...	All other causes (residual)	48	20.7	32.4
American Indian, female, 40-44 years				
...	All causes	289	100.0	202.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	67	23.2	46.9
2	Chronic liver disease and cirrhosis (K70,K73-K74)	53	18.3	37.1
3	Malignant neoplasms (C00-C97)	30	10.4	21.0
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	25	8.7	17.5
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	15	5.2	*
6	Diabetes mellitus (E10-E14)	13	4.5	*
7	Influenza and pneumonia (J09-J18)	12	4.2	*
8	Septicemia (A40-A41)	6	2.1	*
9	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	5	1.7	*
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	4	1.4	*
11	Viral hepatitis (B15-B19)	3	1.0	*
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	3	1.0	*
11	Cerebrovascular diseases (I60-I69)	3	1.0	*
11	Chronic lower respiratory diseases (J40-J47)	3	1.0	*
15	Pneumonitis due to solids and liquids (J69)	2	0.7	*
...	All other causes (residual)	45	15.6	31.5

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2014

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2014

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, female, 45-49 years				
...	All causes	410	100.0	299.6
1	Chronic liver disease and cirrhosis (K70,K73-K74)	66	16.1	48.2
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	58	14.1	42.4
3	Malignant neoplasms (C00-C97)	54	13.2	39.5
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	45	11.0	32.9
5	Diabetes mellitus (E10-E14)	25	6.1	18.3
6	Cerebrovascular diseases (I60-I69)	17	4.1	*
7	Septicemia (A40-A41)	10	2.4	*
7	Influenza and pneumonia (J09-J18)	10	2.4	*
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	8	2.0	*
10	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	7	1.7	*
11	Chronic lower respiratory diseases (J40-J47)	5	1.2	*
11	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	5	1.2	*
13	Human immunodeficiency virus (HIV) disease (B20-B24)	3	0.7	*
13	Enterocolitis due to <i>Clostridium difficile</i> (A04.7)	3	0.7	*
15	Tuberculosis (A16-A19)	2	0.5	*
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2	0.5	*
15	Pregnancy, childbirth and the puerperium (O00-O99)	2	0.5	*
15	Complications of medical and surgical care (Y40-Y84,Y88)	2	0.5	*
...	All other causes (residual)	86	21.0	62.8
American Indian, female, 50-54 years				
...	All causes	572	100.0	407.7
1	Malignant neoplasms (C00-C97)	132	23.1	94.1
2	Chronic liver disease and cirrhosis (K70,K73-K74)	77	13.5	54.9
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	68	11.9	48.5
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	57	10.0	40.6
5	Diabetes mellitus (E10-E14)	27	4.7	19.2
6	Chronic lower respiratory diseases (J40-J47)	21	3.7	15.0
7	Influenza and pneumonia (J09-J18)	16	2.8	*
8	Cerebrovascular diseases (I60-I69)	14	2.4	*
9	Septicemia (A40-A41)	11	1.9	*
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	8	1.4	*
11	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	7	1.2	*
12	Viral hepatitis (B15-B19)	6	1.0	*
13	Human immunodeficiency virus (HIV) disease (B20-B24)	5	0.9	*
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	3	0.5	*
15	Pneumonitis due to solids and liquids (J69)	2	0.3	*
15	Peptic ulcer (K25-K28)	2	0.3	*
15	Infections of kidney (N10-N12,N13.6,N15.1)	2	0.3	*
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	2	0.3	*
15	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	2	0.3	*
15	Enterocolitis due to <i>Clostridium difficile</i> (A04.7)	2	0.3	*
...	All other causes (residual)	108	18.9	77.0

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2014

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2014

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, female, 55-59 years				
...	All causes	637	100.0	516.8
1	Malignant neoplasms (C00-C97)	136	21.4	110.3
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	89	14.0	72.2
3	Chronic liver disease and cirrhosis (K70,K73-K74)	70	11.0	56.8
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	47	7.4	38.1
5	Diabetes mellitus (E10-E14)	39	6.1	31.6
6	Chronic lower respiratory diseases (J40-J47)	34	5.3	27.6
7	Cerebrovascular diseases (I60-I69)	19	3.0	*
8	Septicemia (A40-A41)	16	2.5	*
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	16	2.5	*
10	Viral hepatitis (B15-B19)	11	1.7	*
10	Influenza and pneumonia (J09-J18)	11	1.7	*
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	5	0.8	*
12	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	5	0.8	*
14	Human immunodeficiency virus (HIV) disease (B20-B24)	4	0.6	*
15	Peptic ulcer (K25-K28)	3	0.5	*
15	Complications of medical and surgical care (Y40-Y84,Y88)	3	0.5	*
...	All other causes (residual)	129	20.3	104.6
American Indian, female, 60-64 years				
...	All causes	684	100.0	722.7
1	Malignant neoplasms (C00-C97)	168	24.6	177.5
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	108	15.8	114.1
3	Diabetes mellitus (E10-E14)	59	8.6	62.3
4	Chronic liver disease and cirrhosis (K70,K73-K74)	50	7.3	52.8
5	Chronic lower respiratory diseases (J40-J47)	37	5.4	39.1
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	32	4.7	33.8
7	Cerebrovascular diseases (I60-I69)	28	4.1	29.6
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	18	2.6	*
9	Influenza and pneumonia (J09-J18)	17	2.5	*
10	Septicemia (A40-A41)	15	2.2	*
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	13	1.9	*
12	Viral hepatitis (B15-B19)	4	0.6	*
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	4	0.6	*
12	Alzheimer's disease (G30)	4	0.6	*
12	Pneumonitis due to solids and liquids (J69)	4	0.6	*
...	All other causes (residual)	123	18.0	130.0

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2014

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2014

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, female, 65-69 years				
...	All causes	755	100.0	1,095.4
1	Malignant neoplasms (C00-C97)	207	27.4	300.3
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	152	20.1	220.5
3	Diabetes mellitus (E10-E14)	60	7.9	87.1
4	Chronic lower respiratory diseases (J40-J47)	52	6.9	75.4
5	Chronic liver disease and cirrhosis (K70,K73-K74)	38	5.0	55.1
6	Cerebrovascular diseases (I60-I69)	28	3.7	40.6
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	23	3.0	33.4
8	Influenza and pneumonia (J09-J18)	19	2.5	*
9	Septicemia (A40-A41)	16	2.1	*
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	15	2.0	*
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	9	1.2	*
12	Viral hepatitis (B15-B19)	4	0.5	*
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	4	0.5	*
12	Alzheimer's disease (G30)	4	0.5	*
12	Enterocolitis due to <i>Clostridium difficile</i> (A04.7)	4	0.5	*
...	All other causes (residual)	120	15.9	174.1
American Indian, female, 70-74 years				
...	All causes	812	100.0	1,781.7
1	Malignant neoplasms (C00-C97)	205	25.2	449.8
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	156	19.2	342.3
3	Chronic lower respiratory diseases (J40-J47)	72	8.9	158.0
4	Diabetes mellitus (E10-E14)	53	6.5	116.3
5	Cerebrovascular diseases (I60-I69)	32	3.9	70.2
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	24	3.0	52.7
7	Chronic liver disease and cirrhosis (K70,K73-K74)	23	2.8	50.5
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	21	2.6	46.1
9	Septicemia (A40-A41)	20	2.5	43.9
10	Influenza and pneumonia (J09-J18)	18	2.2	*
11	Alzheimer's disease (G30)	13	1.6	*
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	9	1.1	*
13	Pneumonitis due to solids and liquids (J69)	5	0.6	*
14	Parkinson's disease (G20-G21)	4	0.5	*
14	Enterocolitis due to <i>Clostridium difficile</i> (A04.7)	4	0.5	*
...	All other causes (residual)	153	18.8	335.7

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2014

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2014

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, female, 75-79 years				
...	All causes	815	100.0	2,660.7
1	Malignant neoplasms (C00-C97)	171	21.0	558.3
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	147	18.0	479.9
3	Chronic lower respiratory diseases (J40-J47)	76	9.3	248.1
4	Diabetes mellitus (E10-E14)	55	6.7	179.6
5	Cerebrovascular diseases (I60-I69)	49	6.0	160.0
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	31	3.8	101.2
7	Alzheimer's disease (G30)	26	3.2	84.9
8	Septicemia (A40-A41)	23	2.8	75.1
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	23	2.8	75.1
10	Influenza and pneumonia (J09-J18)	20	2.5	65.3
11	Chronic liver disease and cirrhosis (K70,K73-K74)	11	1.3	*
12	Parkinson's disease (G20-G21)	10	1.2	*
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	10	1.2	*
12	Pneumonitis due to solids and liquids (J69)	10	1.2	*
15	Aortic aneurysm and dissection (I71)	6	0.7	*
...	All other causes (residual)	147	18.0	479.9
American Indian, female, 80-84 years				
...	All causes	805	100.0	4,112.0
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	183	22.7	934.8
2	Malignant neoplasms (C00-C97)	136	16.9	694.7
3	Alzheimer's disease (G30)	46	5.7	235.0
4	Cerebrovascular diseases (I60-I69)	43	5.3	219.6
5	Chronic lower respiratory diseases (J40-J47)	41	5.1	209.4
6	Diabetes mellitus (E10-E14)	40	5.0	204.3
7	Influenza and pneumonia (J09-J18)	27	3.4	137.9
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	17	2.1	*
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	17	2.1	*
10	Septicemia (A40-A41)	15	1.9	*
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	12	1.5	*
12	Parkinson's disease (G20-G21)	10	1.2	*
13	Enterocolitis due to <i>Clostridium difficile</i> (A04.7)	9	1.1	*
14	Aortic aneurysm and dissection (I71)	7	0.9	*
14	Chronic liver disease and cirrhosis (K70,K73-K74)	7	0.9	*
...	All other causes (residual)	195	24.2	996.1

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2014

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2014

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, female, 85-89 years				
...	All causes	696	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	169	24.3	---
2	Malignant neoplasms (C00-C97)	96	13.8	---
3	Cerebrovascular diseases (I60-I69)	53	7.6	---
4	Alzheimer's disease (G30)	52	7.5	---
5	Chronic lower respiratory diseases (J40-J47)	38	5.5	---
6	Diabetes mellitus (E10-E14)	30	4.3	---
7	Influenza and pneumonia (J09-J18)	23	3.3	---
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	17	2.4	---
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	16	2.3	---
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	9	1.3	---
11	Septicemia (A40-A41)	7	1.0	---
12	Nutritional deficiencies (E40-E64)	6	0.9	---
13	Parkinson's disease (G20-G21)	4	0.6	---
13	Pneumonitis due to solids and liquids (J69)	4	0.6	---
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	3	0.4	---
15	Hernia (K40-K46)	3	0.4	---
15	Chronic liver disease and cirrhosis (K70,K73-K74)	3	0.4	---
...	All other causes (residual)	163	23.4	---
American Indian, female, 90-94 years				
...	All causes	520	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	122	23.5	---
2	Alzheimer's disease (G30)	52	10.0	---
3	Malignant neoplasms (C00-C97)	44	8.5	---
4	Cerebrovascular diseases (I60-I69)	41	7.9	---
5	Chronic lower respiratory diseases (J40-J47)	30	5.8	---
6	Diabetes mellitus (E10-E14)	21	4.0	---
7	Influenza and pneumonia (J09-J18)	18	3.5	---
8	Essential hypertension and hypertensive renal disease (I10,I12,I15)	14	2.7	---
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	10	1.9	---
10	Septicemia (A40-A41)	8	1.5	---
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	6	1.2	---
12	Parkinson's disease (G20-G21)	5	1.0	---
12	Pneumonitis due to solids and liquids (J69)	5	1.0	---
14	Hernia (K40-K46)	4	0.8	---
15	Anemias (D50-D64)	3	0.6	---
15	Nutritional deficiencies (E40-E64)	3	0.6	---
...	All other causes (residual)	134	25.8	---

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2014

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2014

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, female, 95-99 years				
...	All causes	192	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	53	27.6	---
2	Alzheimer's disease (G30)	23	12.0	---
3	Cerebrovascular diseases (I60-I69)	17	8.9	---
4	Influenza and pneumonia (J09-J18)	9	4.7	---
4	Chronic lower respiratory diseases (J40-J47)	9	4.7	---
6	Malignant neoplasms (C00-C97)	8	4.2	---
6	Essential hypertension and hypertensive renal disease (I10,I12,I15)	8	4.2	---
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	7	3.6	---
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3	1.6	---
10	Septicemia (A40-A41)	2	1.0	---
10	Diabetes mellitus (E10-E14)	2	1.0	---
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1	0.5	---
12	Nutritional deficiencies (E40-E64)	1	0.5	---
12	Parkinson's disease (G20-G21)	1	0.5	---
12	Pneumonitis due to solids and liquids (J69)	1	0.5	---
12	Cholelithiasis and other disorders of gallbladder (K80-K82)	1	0.5	---
...	All other causes (residual)	46	24.0	---
American Indian, female, 100 years and over				
...	All causes	59	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	15	25.4	---
2	Malignant neoplasms (C00-C97)	4	6.8	---
2	Essential hypertension and hypertensive renal disease (I10,I12,I15)	4	6.8	---
2	Cerebrovascular diseases (I60-I69)	4	6.8	---
5	Alzheimer's disease (G30)	3	5.1	---
6	Diabetes mellitus (E10-E14)	2	3.4	---
6	Chronic lower respiratory diseases (J40-J47)	2	3.4	---
6	Pneumonitis due to solids and liquids (J69)	2	3.4	---
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2	3.4	---
10	Septicemia (A40-A41)	1	1.7	---
10	Anemias (D50-D64)	1	1.7	---
10	Nutritional deficiencies (E40-E64)	1	1.7	---
10	Influenza and pneumonia (J09-J18)	1	1.7	---
10	Enterocolitis due to <i>Clostridium difficile</i> (A04.7)	1	1.7	---
...	All other causes (residual)	16	27.1	---

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2014

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2014

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, both sexes, all ages				
...	All causes	61,570	100.0	317.4
1	Malignant neoplasms (C00-C97)	16,524	26.8	85.2
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	13,270	21.6	68.4
3	Cerebrovascular diseases (I60-I69)	4,331	7.0	22.3
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,646	4.3	13.6
5	Diabetes mellitus (E10-E14)	2,367	3.8	12.2
6	Influenza and pneumonia (J09-J18)	1,911	3.1	9.9
7	Chronic lower respiratory diseases (J40-J47)	1,838	3.0	9.5
8	Alzheimer's disease (G30)	1,680	2.7	8.7
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,246	2.0	6.4
10	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,188	1.9	6.1
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,007	1.6	5.2
12	Septicemia (A40-A41)	753	1.2	3.9
13	Parkinson's disease (G20-G21)	611	1.0	3.1
14	Chronic liver disease and cirrhosis (K70,K73-K74)	608	1.0	3.1
15	Certain conditions originating in the perinatal period (P00-P96)	508	0.8	2.6
...	All other causes (residual)	11,082	18.0	57.1
Asian or Pacific Islander, both sexes, 1-4 years				
...	All causes	134	100.0	13.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	30	22.4	3.0
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	18	13.4	*
3	Malignant neoplasms (C00-C97)	12	9.0	*
4	Influenza and pneumonia (J09-J18)	7	5.2	*
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	6	4.5	*
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	5	3.7	*
7	Cerebrovascular diseases (I60-I69)	3	2.2	*
7	Chronic lower respiratory diseases (J40-J47)	3	2.2	*
9	Septicemia (A40-A41)	2	1.5	*
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2	1.5	*
11	Anemias (D50-D64)	1	0.7	*
11	Hernia (K40-K46)	1	0.7	*
...	All other causes (residual)	44	32.8	4.4

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2014

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2014

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, both sexes, 5-9 years				
...	All causes	95	100.0	7.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	21	22.1	1.7
2	Malignant neoplasms (C00-C97)	12	12.6	*
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	10	10.5	*
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	5	5.3	*
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	5	5.3	*
6	Septicemia (A40-A41)	3	3.2	*
6	Chronic lower respiratory diseases (J40-J47)	3	3.2	*
8	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2	2.1	*
8	Cerebrovascular diseases (I60-I69)	2	2.1	*
8	Influenza and pneumonia (J09-J18)	2	2.1	*
11	Anemias (D50-D64)	1	1.1	*
11	Nutritional deficiencies (E40-E64)	1	1.1	*
11	Atherosclerosis (I70)	1	1.1	*
11	Diseases of appendix (K35-K38)	1	1.1	*
11	Enterocolitis due to <i>Clostridium difficile</i> (A04.7)	1	1.1	*
...	All other causes (residual)	25	26.3	2.0
Asian or Pacific Islander, both sexes, 10-14 years				
...	All causes	109	100.0	8.8
1	Malignant neoplasms (C00-C97)	21	19.3	1.7
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	19	17.4	*
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	10	9.2	*
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	8	7.3	*
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	8	7.3	*
6	Septicemia (A40-A41)	3	2.8	*
7	Anemias (D50-D64)	2	1.8	*
7	Influenza and pneumonia (J09-J18)	2	1.8	*
7	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	2	1.8	*
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1	0.9	*
10	Cerebrovascular diseases (I60-I69)	1	0.9	*
10	Chronic lower respiratory diseases (J40-J47)	1	0.9	*
10	Peptic ulcer (K25-K28)	1	0.9	*
10	Certain conditions originating in the perinatal period (P00-P96)	1	0.9	*
...	All other causes (residual)	29	26.6	2.4

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2014

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2014

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, both sexes, 15-19 years				
...	All causes	282	100.0	23.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	90	31.9	7.4
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	67	23.8	5.5
3	Malignant neoplasms (C00-C97)	31	11.0	2.5
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	18	6.4	*
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	9	3.2	*
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	7	2.5	*
7	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	5	1.8	*
8	Cerebrovascular diseases (I60-I69)	4	1.4	*
9	Chronic lower respiratory diseases (J40-J47)	2	0.7	*
10	Anemias (D50-D64)	1	0.4	*
10	Influenza and pneumonia (J09-J18)	1	0.4	*
10	Chronic liver disease and cirrhosis (K70,K73-K74)	1	0.4	*
10	Certain conditions originating in the perinatal period (P00-P96)	1	0.4	*
...	All other causes (residual)	45	16.0	3.7
Asian or Pacific Islander, both sexes, 20-24 years				
...	All causes	561	100.0	38.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	182	32.4	12.4
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	162	28.9	11.1
3	Malignant neoplasms (C00-C97)	43	7.7	2.9
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	43	7.7	2.9
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	26	4.6	1.8
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	7	1.2	*
7	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	5	0.9	*
8	Chronic lower respiratory diseases (J40-J47)	4	0.7	*
9	Septicemia (A40-A41)	3	0.5	*
9	Cerebrovascular diseases (I60-I69)	3	0.5	*
9	Pneumonitis due to solids and liquids (J69)	3	0.5	*
9	Pregnancy, childbirth and the puerperium (O00-O99)	3	0.5	*
9	Legal intervention (Y35,Y89.0)	3	0.5	*
14	Diabetes mellitus (E10-E14)	2	0.4	*
14	Meningitis (G00,G03)	2	0.4	*
14	Influenza and pneumonia (J09-J18)	2	0.4	*
...	All other causes (residual)	68	12.1	4.7

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2014

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2014

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, both sexes, 25-29 years				
...	All causes	605	100.0	37.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	199	32.9	12.2
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	115	19.0	7.1
3	Malignant neoplasms (C00-C97)	75	12.4	4.6
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	45	7.4	2.8
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	29	4.8	1.8
6	Cerebrovascular diseases (I60-I69)	11	1.8	*
7	Chronic liver disease and cirrhosis (K70,K73-K74)	9	1.5	*
7	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	9	1.5	*
9	Influenza and pneumonia (J09-J18)	7	1.2	*
10	Diabetes mellitus (E10-E14)	6	1.0	*
10	Pregnancy, childbirth and the puerperium (O00-O99)	6	1.0	*
12	Septicemia (A40-A41)	5	0.8	*
13	Human immunodeficiency virus (HIV) disease (B20-B24)	3	0.5	*
13	Aortic aneurysm and dissection (I71)	3	0.5	*
13	Chronic lower respiratory diseases (J40-J47)	3	0.5	*
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3	0.5	*
13	Complications of medical and surgical care (Y40-Y84,Y88)	3	0.5	*
...	All other causes (residual)	74	12.2	4.5
Asian or Pacific Islander, both sexes, 30-34 years				
...	All causes	672	100.0	40.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	151	22.5	9.1
2	Malignant neoplasms (C00-C97)	125	18.6	7.6
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	120	17.9	7.3
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	59	8.8	3.6
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	31	4.6	1.9
6	Cerebrovascular diseases (I60-I69)	16	2.4	*
7	Diabetes mellitus (E10-E14)	13	1.9	*
8	Pregnancy, childbirth and the puerperium (O00-O99)	11	1.6	*
9	Septicemia (A40-A41)	8	1.2	*
9	Influenza and pneumonia (J09-J18)	8	1.2	*
9	Chronic liver disease and cirrhosis (K70,K73-K74)	8	1.2	*
12	Human immunodeficiency virus (HIV) disease (B20-B24)	7	1.0	*
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	7	1.0	*
14	Viral hepatitis (B15-B19)	5	0.7	*
14	Chronic lower respiratory diseases (J40-J47)	5	0.7	*
...	All other causes (residual)	98	14.6	5.9

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2014

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2014

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, both sexes, 35-39 years				
...	All causes	799	100.0	51.7
1	Malignant neoplasms (C00-C97)	220	27.5	14.2
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	111	13.9	7.2
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	108	13.5	7.0
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	87	10.9	5.6
5	Cerebrovascular diseases (I60-I69)	30	3.8	1.9
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	26	3.3	1.7
7	Chronic liver disease and cirrhosis (K70,K73-K74)	21	2.6	1.4
8	Diabetes mellitus (E10-E14)	19	2.4	*
9	Pregnancy, childbirth and the puerperium (O00-O99)	12	1.5	*
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	11	1.4	*
11	Aortic aneurysm and dissection (I71)	9	1.1	*
11	Chronic lower respiratory diseases (J40-J47)	9	1.1	*
13	Influenza and pneumonia (J09-J18)	8	1.0	*
14	Viral hepatitis (B15-B19)	7	0.9	*
15	Septicemia (A40-A41)	6	0.8	*
...	All other causes (residual)	115	14.4	7.4
Asian or Pacific Islander, both sexes, 40-44 years				
...	All causes	1,206	100.0	78.4
1	Malignant neoplasms (C00-C97)	364	30.2	23.7
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	202	16.7	13.1
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	143	11.9	9.3
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	105	8.7	6.8
5	Cerebrovascular diseases (I60-I69)	60	5.0	3.9
6	Chronic liver disease and cirrhosis (K70,K73-K74)	30	2.5	2.0
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	30	2.5	2.0
8	Diabetes mellitus (E10-E14)	25	2.1	1.6
9	Influenza and pneumonia (J09-J18)	21	1.7	1.4
10	Chronic lower respiratory diseases (J40-J47)	15	1.2	*
11	Human immunodeficiency virus (HIV) disease (B20-B24)	14	1.2	*
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	10	0.8	*
12	Pregnancy, childbirth and the puerperium (O00-O99)	10	0.8	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	8	0.7	*
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	8	0.7	*
...	All other causes (residual)	161	13.3	10.5

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2014

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2014

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, both sexes, 45-49 years				
...	All causes	1,693	100.0	127.2
1	Malignant neoplasms (C00-C97)	628	37.1	47.2
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	317	18.7	23.8
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	123	7.3	9.2
4	Cerebrovascular diseases (I60-I69)	101	6.0	7.6
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	101	6.0	7.6
6	Diabetes mellitus (E10-E14)	43	2.5	3.2
7	Chronic liver disease and cirrhosis (K70,K73-K74)	37	2.2	2.8
8	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	25	1.5	1.9
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	23	1.4	1.7
10	Septicemia (A40-A41)	21	1.2	1.6
11	Influenza and pneumonia (J09-J18)	19	1.1	*
12	Chronic lower respiratory diseases (J40-J47)	17	1.0	*
13	Viral hepatitis (B15-B19)	14	0.8	*
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	14	0.8	*
13	Aortic aneurysm and dissection (I71)	14	0.8	*
...	All other causes (residual)	196	11.6	14.7
Asian or Pacific Islander, both sexes, 50-54 years				
...	All causes	2,536	100.0	204.5
1	Malignant neoplasms (C00-C97)	1,026	40.5	82.7
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	489	19.3	39.4
3	Cerebrovascular diseases (I60-I69)	148	5.8	11.9
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	136	5.4	11.0
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	113	4.5	9.1
6	Diabetes mellitus (E10-E14)	89	3.5	7.2
7	Chronic liver disease and cirrhosis (K70,K73-K74)	53	2.1	4.3
8	Essential hypertension and hypertensive renal disease (I10,I12,I15)	34	1.3	2.7
9	Influenza and pneumonia (J09-J18)	32	1.3	2.6
10	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	28	1.1	2.3
11	Viral hepatitis (B15-B19)	26	1.0	2.1
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	24	0.9	1.9
13	Septicemia (A40-A41)	17	0.7	*
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	14	0.6	*
15	Aortic aneurysm and dissection (I71)	13	0.5	*
...	All other causes (residual)	294	11.6	23.7

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2014

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2014

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, both sexes, 55-59 years				
...	All causes	3,553	100.0	321.4
1	Malignant neoplasms (C00-C97)	1,527	43.0	138.1
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	631	17.8	57.1
3	Cerebrovascular diseases (I60-I69)	218	6.1	19.7
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	155	4.4	14.0
5	Diabetes mellitus (E10-E14)	152	4.3	13.8
6	Chronic liver disease and cirrhosis (K70,K73-K74)	86	2.4	7.8
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	86	2.4	7.8
8	Influenza and pneumonia (J09-J18)	63	1.8	5.7
9	Chronic lower respiratory diseases (J40-J47)	55	1.5	5.0
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	53	1.5	4.8
11	Viral hepatitis (B15-B19)	41	1.2	3.7
12	Septicemia (A40-A41)	35	1.0	3.2
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	30	0.8	2.7
14	Aortic aneurysm and dissection (I71)	28	0.8	2.5
15	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	27	0.8	2.4
...	All other causes (residual)	366	10.3	33.1
Asian or Pacific Islander, both sexes, 60-64 years				
...	All causes	4,401	100.0	473.0
1	Malignant neoplasms (C00-C97)	1,835	41.7	197.2
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	880	20.0	94.6
3	Cerebrovascular diseases (I60-I69)	263	6.0	28.3
4	Diabetes mellitus (E10-E14)	204	4.6	21.9
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	150	3.4	16.1
6	Influenza and pneumonia (J09-J18)	81	1.8	8.7
7	Chronic lower respiratory diseases (J40-J47)	71	1.6	7.6
8	Chronic liver disease and cirrhosis (K70,K73-K74)	63	1.4	6.8
9	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	62	1.4	6.7
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	59	1.3	6.3
11	Septicemia (A40-A41)	58	1.3	6.2
11	Viral hepatitis (B15-B19)	58	1.3	6.2
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	57	1.3	6.1
14	Aortic aneurysm and dissection (I71)	30	0.7	3.2
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	22	0.5	2.4
...	All other causes (residual)	508	11.5	54.6

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2014

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2014

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, both sexes, 65-69 years				
...	All causes	5,083	100.0	706.1
1	Malignant neoplasms (C00-C97)	2,024	39.8	281.2
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,031	20.3	143.2
3	Cerebrovascular diseases (I60-I69)	311	6.1	43.2
4	Diabetes mellitus (E10-E14)	254	5.0	35.3
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	134	2.6	18.6
6	Chronic lower respiratory diseases (J40-J47)	109	2.1	15.1
7	Influenza and pneumonia (J09-J18)	106	2.1	14.7
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	104	2.0	14.4
9	Septicemia (A40-A41)	81	1.6	11.3
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	61	1.2	8.5
11	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	58	1.1	8.1
12	Chronic liver disease and cirrhosis (K70,K73-K74)	56	1.1	7.8
13	Viral hepatitis (B15-B19)	44	0.9	6.1
14	Parkinson's disease (G20-G21)	34	0.7	4.7
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	27	0.5	3.8
15	Aortic aneurysm and dissection (I71)	27	0.5	3.8
...	All other causes (residual)	622	12.2	86.4
Asian or Pacific Islander, both sexes, 70-74 years				
...	All causes	5,645	100.0	1,140.1
1	Malignant neoplasms (C00-C97)	2,022	35.8	408.4
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,171	20.7	236.5
3	Cerebrovascular diseases (I60-I69)	401	7.1	81.0
4	Diabetes mellitus (E10-E14)	270	4.8	54.5
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	162	2.9	32.7
6	Chronic lower respiratory diseases (J40-J47)	160	2.8	32.3
7	Influenza and pneumonia (J09-J18)	147	2.6	29.7
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	132	2.3	26.7
9	Essential hypertension and hypertensive renal disease (I10,I12,I15)	85	1.5	17.2
10	Septicemia (A40-A41)	83	1.5	16.8
11	Chronic liver disease and cirrhosis (K70,K73-K74)	72	1.3	14.5
12	Parkinson's disease (G20-G21)	58	1.0	11.7
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	42	0.7	8.5
13	Alzheimer's disease (G30)	42	0.7	8.5
15	Pneumonitis due to solids and liquids (J69)	36	0.6	7.3
...	All other causes (residual)	762	13.5	153.9

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2014

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2014

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, both sexes, 75-79 years				
...	All causes	6,801	100.0	1,954.6
1	Malignant neoplasms (C00-C97)	2,101	30.9	603.8
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,495	22.0	429.7
3	Cerebrovascular diseases (I60-I69)	515	7.6	148.0
4	Diabetes mellitus (E10-E14)	311	4.6	89.4
5	Chronic lower respiratory diseases (J40-J47)	238	3.5	68.4
6	Influenza and pneumonia (J09-J18)	204	3.0	58.6
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	172	2.5	49.4
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	169	2.5	48.6
9	Alzheimer's disease (G30)	129	1.9	37.1
10	Parkinson's disease (G20-G21)	115	1.7	33.1
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	108	1.6	31.0
12	Septicemia (A40-A41)	90	1.3	25.9
13	Chronic liver disease and cirrhosis (K70,K73-K74)	67	1.0	19.3
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	55	0.8	15.8
15	Pneumonitis due to solids and liquids (J69)	49	0.7	14.1
...	All other causes (residual)	983	14.5	282.5
Asian or Pacific Islander, both sexes, 80-84 years				
...	All causes	8,178	100.0	3,530.5
1	Malignant neoplasms (C00-C97)	1,962	24.0	847.0
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,917	23.4	827.6
3	Cerebrovascular diseases (I60-I69)	680	8.3	293.6
4	Diabetes mellitus (E10-E14)	379	4.6	163.6
5	Chronic lower respiratory diseases (J40-J47)	378	4.6	163.2
6	Influenza and pneumonia (J09-J18)	317	3.9	136.9
7	Alzheimer's disease (G30)	273	3.3	117.9
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	202	2.5	87.2
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	200	2.4	86.3
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	134	1.6	57.8
11	Parkinson's disease (G20-G21)	129	1.6	55.7
12	Septicemia (A40-A41)	113	1.4	48.8
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	72	0.9	31.1
13	Pneumonitis due to solids and liquids (J69)	72	0.9	31.1
15	Chronic liver disease and cirrhosis (K70,K73-K74)	63	0.8	27.2
...	All other causes (residual)	1,287	15.7	555.6

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2014

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2014

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, both sexes, 85-89 years				
...	All causes	8,535	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,176	25.5	---
2	Malignant neoplasms (C00-C97)	1,511	17.7	---
3	Cerebrovascular diseases (I60-I69)	766	9.0	---
4	Alzheimer's disease (G30)	445	5.2	---
5	Chronic lower respiratory diseases (J40-J47)	371	4.3	---
6	Influenza and pneumonia (J09-J18)	328	3.8	---
7	Diabetes mellitus (E10-E14)	325	3.8	---
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	224	2.6	---
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	221	2.6	---
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	195	2.3	---
11	Parkinson's disease (G20-G21)	149	1.7	---
12	Septicemia (A40-A41)	119	1.4	---
13	Pneumonitis due to solids and liquids (J69)	103	1.2	---
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	65	0.8	---
15	Aortic aneurysm and dissection (I71)	49	0.6	---
...	All other causes (residual)	1,488	17.4	---
Asian or Pacific Islander, both sexes, 90-94 years				
...	All causes	6,521	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,745	26.8	---
2	Malignant neoplasms (C00-C97)	751	11.5	---
3	Cerebrovascular diseases (I60-I69)	541	8.3	---
4	Alzheimer's disease (G30)	488	7.5	---
5	Influenza and pneumonia (J09-J18)	359	5.5	---
6	Chronic lower respiratory diseases (J40-J47)	257	3.9	---
7	Diabetes mellitus (E10-E14)	188	2.9	---
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	165	2.5	---
9	Essential hypertension and hypertensive renal disease (I10,I12,I15)	162	2.5	---
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	159	2.4	---
11	Pneumonitis due to solids and liquids (J69)	101	1.5	---
12	Parkinson's disease (G20-G21)	91	1.4	---
13	Septicemia (A40-A41)	61	0.9	---
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	43	0.7	---
15	Atherosclerosis (I70)	26	0.4	---
15	Aortic aneurysm and dissection (I71)	26	0.4	---
...	All other causes (residual)	1,358	20.8	---

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2014

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2014

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, both sexes, 95-99 years				
...	All causes	2,632	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	737	28.0	---
2	Alzheimer's disease (G30)	230	8.7	---
3	Cerebrovascular diseases (I60-I69)	211	8.0	---
4	Malignant neoplasms (C00-C97)	197	7.5	---
5	Influenza and pneumonia (J09-J18)	149	5.7	---
6	Chronic lower respiratory diseases (J40-J47)	102	3.9	---
7	Essential hypertension and hypertensive renal disease (I10,I12,I15)	90	3.4	---
8	Diabetes mellitus (E10-E14)	73	2.8	---
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	59	2.2	---
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	50	1.9	---
11	Pneumonitis due to solids and liquids (J69)	36	1.4	---
12	Septicemia (A40-A41)	24	0.9	---
13	Parkinson's disease (G20-G21)	16	0.6	---
14	Atherosclerosis (I70)	13	0.5	---
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	10	0.4	---
...	All other causes (residual)	635	24.1	---
Asian or Pacific Islander, both sexes, 100 years and over				
...	All causes	631	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	189	30.0	---
2	Alzheimer's disease (G30)	49	7.8	---
3	Cerebrovascular diseases (I60-I69)	42	6.7	---
4	Influenza and pneumonia (J09-J18)	41	6.5	---
5	Malignant neoplasms (C00-C97)	37	5.9	---
6	Essential hypertension and hypertensive renal disease (I10,I12,I15)	24	3.8	---
7	Chronic lower respiratory diseases (J40-J47)	23	3.6	---
8	Diabetes mellitus (E10-E14)	14	2.2	---
9	Pneumonitis due to solids and liquids (J69)	13	2.1	---
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	12	1.9	---
10	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	12	1.9	---
12	Septicemia (A40-A41)	6	1.0	---
13	Nutritional deficiencies (E40-E64)	4	0.6	---
13	Parkinson's disease (G20-G21)	4	0.6	---
13	Atherosclerosis (I70)	4	0.6	---
...	All other causes (residual)	157	24.9	---

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2014

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2014

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, male, all ages				
...	All causes	31,686	100.0	341.3
1	Malignant neoplasms (C00-C97)	8,371	26.4	90.2
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	7,279	23.0	78.4
3	Cerebrovascular diseases (I60-I69)	1,923	6.1	20.7
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,665	5.3	17.9
5	Diabetes mellitus (E10-E14)	1,235	3.9	13.3
6	Chronic lower respiratory diseases (J40-J47)	1,072	3.4	11.5
7	Influenza and pneumonia (J09-J18)	1,012	3.2	10.9
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	833	2.6	9.0
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	652	2.1	7.0
10	Alzheimer's disease (G30)	508	1.6	5.5
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	435	1.4	4.7
12	Septicemia (A40-A41)	394	1.2	4.2
13	Chronic liver disease and cirrhosis (K70,K73-K74)	382	1.2	4.1
14	Parkinson's disease (G20-G21)	362	1.1	3.9
15	Certain conditions originating in the perinatal period (P00-P96)	281	0.9	3.0
...	All other causes (residual)	5,282	16.7	56.9
Asian or Pacific Islander, male, 1-4 years				
...	All causes	73	100.0	14.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	20	27.4	3.9
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	11	15.1	*
3	Malignant neoplasms (C00-C97)	7	9.6	*
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	5	6.8	*
5	Chronic lower respiratory diseases (J40-J47)	3	4.1	*
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	2	2.7	*
7	Septicemia (A40-A41)	1	1.4	*
7	Anemias (D50-D64)	1	1.4	*
7	Cerebrovascular diseases (I60-I69)	1	1.4	*
7	Influenza and pneumonia (J09-J18)	1	1.4	*
...	All other causes (residual)	21	28.8	4.1
Asian or Pacific Islander, male, 5-9 years				
...	All causes	53	100.0	8.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	11	20.8	*
2	Malignant neoplasms (C00-C97)	7	13.2	*
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	4	7.5	*
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	4	7.5	*
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	3	5.7	*
6	Septicemia (A40-A41)	2	3.8	*
6	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2	3.8	*
6	Cerebrovascular diseases (I60-I69)	2	3.8	*
6	Chronic lower respiratory diseases (J40-J47)	2	3.8	*
10	Atherosclerosis (I70)	1	1.9	*
10	Influenza and pneumonia (J09-J18)	1	1.9	*
...	All other causes (residual)	14	26.4	*

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2014

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2014

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, male, 10-14 years				
...	All causes	71	100.0	11.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	14	19.7	*
2	Malignant neoplasms (C00-C97)	13	18.3	*
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	6	8.5	*
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	5	7.0	*
5	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	4	5.6	*
6	Septicemia (A40-A41)	3	4.2	*
7	Influenza and pneumonia (J09-J18)	2	2.8	*
7	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	2	2.8	*
9	Anemias (D50-D64)	1	1.4	*
9	Cerebrovascular diseases (I60-I69)	1	1.4	*
9	Peptic ulcer (K25-K28)	1	1.4	*
...	All other causes (residual)	19	26.8	*
Asian or Pacific Islander, male, 15-19 years				
...	All causes	172	100.0	27.9
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	64	37.2	10.4
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	39	22.7	6.3
3	Malignant neoplasms (C00-C97)	16	9.3	*
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	13	7.6	*
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	5	2.9	*
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	3	1.7	*
7	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2	1.2	*
7	Cerebrovascular diseases (I60-I69)	2	1.2	*
7	Chronic lower respiratory diseases (J40-J47)	2	1.2	*
10	Chronic liver disease and cirrhosis (K70,K73-K74)	1	0.6	*
10	Certain conditions originating in the perinatal period (P00-P96)	1	0.6	*
...	All other causes (residual)	24	14.0	3.9

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2014

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2014

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, male, 20-24 years				
...	All causes	436	100.0	58.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	144	33.0	19.4
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	136	31.2	18.3
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	33	7.6	4.4
4	Malignant neoplasms (C00-C97)	31	7.1	4.2
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	21	4.8	2.8
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	5	1.1	*
7	Chronic lower respiratory diseases (J40-J47)	3	0.7	*
7	Pneumonitis due to solids and liquids (J69)	3	0.7	*
7	Legal intervention (Y35,Y89.0)	3	0.7	*
10	Septicemia (A40-A41)	2	0.5	*
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2	0.5	*
10	Diabetes mellitus (E10-E14)	2	0.5	*
10	Cerebrovascular diseases (I60-I69)	2	0.5	*
14	Parkinson's disease (G20-G21)	1	0.2	*
14	Influenza and pneumonia (J09-J18)	1	0.2	*
14	Chronic liver disease and cirrhosis (K70,K73-K74)	1	0.2	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1	0.2	*
...	All other causes (residual)	45	10.3	6.1
Asian or Pacific Islander, male, 25-29 years				
...	All causes	420	100.0	52.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	153	36.4	19.2
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	85	20.2	10.7
3	Malignant neoplasms (C00-C97)	39	9.3	4.9
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	35	8.3	4.4
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	26	6.2	3.3
6	Chronic liver disease and cirrhosis (K70,K73-K74)	7	1.7	*
7	Diabetes mellitus (E10-E14)	5	1.2	*
8	Cerebrovascular diseases (I60-I69)	4	1.0	*
8	Influenza and pneumonia (J09-J18)	4	1.0	*
8	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	4	1.0	*
11	Septicemia (A40-A41)	3	0.7	*
11	Human immunodeficiency virus (HIV) disease (B20-B24)	3	0.7	*
11	Aortic aneurysm and dissection (I71)	3	0.7	*
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3	0.7	*
15	Legal intervention (Y35,Y89.0)	2	0.5	*
15	Complications of medical and surgical care (Y40-Y84,Y88)	2	0.5	*
...	All other causes (residual)	42	10.0	5.3

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2014

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2014

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, male, 30-34 years				
...	All causes	435	100.0	55.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	113	26.0	14.4
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	86	19.8	11.0
3	Malignant neoplasms (C00-C97)	55	12.6	7.0
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	49	11.3	6.3
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	26	6.0	3.3
6	Cerebrovascular diseases (I60-I69)	12	2.8	*
7	Diabetes mellitus (E10-E14)	10	2.3	*
8	Human immunodeficiency virus (HIV) disease (B20-B24)	7	1.6	*
9	Viral hepatitis (B15-B19)	5	1.1	*
10	Septicemia (A40-A41)	4	0.9	*
10	Chronic liver disease and cirrhosis (K70,K73-K74)	4	0.9	*
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3	0.7	*
12	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	3	0.7	*
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2	0.5	*
14	Aortic aneurysm and dissection (I71)	2	0.5	*
14	Influenza and pneumonia (J09-J18)	2	0.5	*
14	Chronic lower respiratory diseases (J40-J47)	2	0.5	*
14	Legal intervention (Y35,Y89.0)	2	0.5	*
...	All other causes (residual)	48	11.0	6.1
Asian or Pacific Islander, male, 35-39 years				
...	All causes	465	100.0	64.0
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	89	19.1	12.2
2	Malignant neoplasms (C00-C97)	81	17.4	11.1
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	76	16.3	10.5
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	54	11.6	7.4
5	Cerebrovascular diseases (I60-I69)	23	4.9	3.2
6	Chronic liver disease and cirrhosis (K70,K73-K74)	17	3.7	*
7	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	16	3.4	*
8	Diabetes mellitus (E10-E14)	9	1.9	*
9	Aortic aneurysm and dissection (I71)	8	1.7	*
10	Chronic lower respiratory diseases (J40-J47)	7	1.5	*
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	6	1.3	*
12	Influenza and pneumonia (J09-J18)	5	1.1	*
13	Viral hepatitis (B15-B19)	4	0.9	*
13	Human immunodeficiency virus (HIV) disease (B20-B24)	4	0.9	*
15	Septicemia (A40-A41)	3	0.6	*
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	3	0.6	*
...	All other causes (residual)	60	12.9	8.3

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2014

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2014

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, male, 40-44 years				
...	All causes	737	100.0	102.1
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	160	21.7	22.2
2	Malignant neoplasms (C00-C97)	149	20.2	20.6
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	107	14.5	14.8
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	69	9.4	9.6
5	Cerebrovascular diseases (I60-I69)	34	4.6	4.7
6	Chronic liver disease and cirrhosis (K70,K73-K74)	23	3.1	3.2
7	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	22	3.0	3.0
8	Diabetes mellitus (E10-E14)	17	2.3	*
9	Influenza and pneumonia (J09-J18)	13	1.8	*
10	Human immunodeficiency virus (HIV) disease (B20-B24)	10	1.4	*
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	9	1.2	*
12	Chronic lower respiratory diseases (J40-J47)	8	1.1	*
13	Aortic aneurysm and dissection (I71)	7	0.9	*
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	6	0.8	*
15	Viral hepatitis (B15-B19)	4	0.5	*
...	All other causes (residual)	99	13.4	13.7
Asian or Pacific Islander, male, 45-49 years				
...	All causes	1,043	100.0	166.7
1	Malignant neoplasms (C00-C97)	289	27.7	46.2
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	259	24.8	41.4
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	95	9.1	15.2
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	73	7.0	11.7
5	Cerebrovascular diseases (I60-I69)	51	4.9	8.2
6	Diabetes mellitus (E10-E14)	30	2.9	4.8
7	Chronic liver disease and cirrhosis (K70,K73-K74)	27	2.6	4.3
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	18	1.7	*
9	Influenza and pneumonia (J09-J18)	12	1.2	*
10	Chronic lower respiratory diseases (J40-J47)	11	1.1	*
11	Septicemia (A40-A41)	10	1.0	*
11	Human immunodeficiency virus (HIV) disease (B20-B24)	10	1.0	*
11	Aortic aneurysm and dissection (I71)	10	1.0	*
11	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	10	1.0	*
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	9	0.9	*
...	All other causes (residual)	129	12.4	20.6

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2014

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2014

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, male, 50-54 years				
...	All causes	1,515	100.0	261.5
1	Malignant neoplasms (C00-C97)	489	32.3	84.4
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	385	25.4	66.5
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	84	5.5	14.5
4	Cerebrovascular diseases (I60-I69)	81	5.3	14.0
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	73	4.8	12.6
6	Diabetes mellitus (E10-E14)	57	3.8	9.8
7	Chronic liver disease and cirrhosis (K70,K73-K74)	44	2.9	7.6
8	Essential hypertension and hypertensive renal disease (I10,I12,I15)	24	1.6	4.1
9	Viral hepatitis (B15-B19)	21	1.4	3.6
10	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	19	1.3	*
11	Influenza and pneumonia (J09-J18)	17	1.1	*
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	16	1.1	*
13	Septicemia (A40-A41)	12	0.8	*
14	Aortic aneurysm and dissection (I71)	11	0.7	*
15	Chronic lower respiratory diseases (J40-J47)	10	0.7	*
...	All other causes (residual)	172	11.4	29.7
Asian or Pacific Islander, male, 55-59 years				
...	All causes	2,163	100.0	426.4
1	Malignant neoplasms (C00-C97)	802	37.1	158.1
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	485	22.4	95.6
3	Cerebrovascular diseases (I60-I69)	124	5.7	24.4
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	101	4.7	19.9
5	Diabetes mellitus (E10-E14)	97	4.5	19.1
6	Chronic liver disease and cirrhosis (K70,K73-K74)	72	3.3	14.2
7	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	62	2.9	12.2
8	Influenza and pneumonia (J09-J18)	45	2.1	8.9
9	Chronic lower respiratory diseases (J40-J47)	36	1.7	7.1
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	33	1.5	6.5
11	Viral hepatitis (B15-B19)	29	1.3	5.7
12	Aortic aneurysm and dissection (I71)	21	1.0	4.1
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	17	0.8	*
13	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	17	0.8	*
15	Septicemia (A40-A41)	16	0.7	*
...	All other causes (residual)	206	9.5	40.6

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2014

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2014

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, male, 60-64 years				
...	All causes	2,645	100.0	633.5
1	Malignant neoplasms (C00-C97)	962	36.4	230.4
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	638	24.1	152.8
3	Cerebrovascular diseases (I60-I69)	153	5.8	36.6
4	Diabetes mellitus (E10-E14)	131	5.0	31.4
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	102	3.9	24.4
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	49	1.9	11.7
7	Influenza and pneumonia (J09-J18)	47	1.8	11.3
7	Chronic lower respiratory diseases (J40-J47)	47	1.8	11.3
9	Chronic liver disease and cirrhosis (K70,K73-K74)	42	1.6	10.1
10	Viral hepatitis (B15-B19)	39	1.5	9.3
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	37	1.4	8.9
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	36	1.4	8.6
13	Septicemia (A40-A41)	35	1.3	8.4
14	Aortic aneurysm and dissection (I71)	20	0.8	4.8
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	11	0.4	*
...	All other causes (residual)	296	11.2	70.9
Asian or Pacific Islander, male, 65-69 years				
...	All causes	2,891	100.0	896.2
1	Malignant neoplasms (C00-C97)	1,101	38.1	341.3
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	640	22.1	198.4
3	Cerebrovascular diseases (I60-I69)	168	5.8	52.1
4	Diabetes mellitus (E10-E14)	140	4.8	43.4
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	85	2.9	26.3
6	Chronic lower respiratory diseases (J40-J47)	69	2.4	21.4
7	Influenza and pneumonia (J09-J18)	59	2.0	18.3
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	57	2.0	17.7
9	Septicemia (A40-A41)	46	1.6	14.3
10	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	41	1.4	12.7
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	35	1.2	10.8
12	Chronic liver disease and cirrhosis (K70,K73-K74)	32	1.1	9.9
13	Parkinson's disease (G20-G21)	22	0.8	6.8
14	Viral hepatitis (B15-B19)	20	0.7	6.2
15	Pneumonitis due to solids and liquids (J69)	14	0.5	*
...	All other causes (residual)	362	12.5	112.2

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2014

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2014

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, male, 70-74 years				
...	All causes	3,146	100.0	1,412.6
1	Malignant neoplasms (C00-C97)	1,078	34.3	484.0
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	726	23.1	326.0
3	Cerebrovascular diseases (I60-I69)	194	6.2	87.1
4	Diabetes mellitus (E10-E14)	157	5.0	70.5
5	Chronic lower respiratory diseases (J40-J47)	104	3.3	46.7
6	Influenza and pneumonia (J09-J18)	93	3.0	41.8
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	88	2.8	39.5
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	70	2.2	31.4
9	Septicemia (A40-A41)	55	1.7	24.7
10	Chronic liver disease and cirrhosis (K70,K73-K74)	41	1.3	18.4
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	38	1.2	17.1
12	Parkinson's disease (G20-G21)	34	1.1	15.3
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	29	0.9	13.0
14	Aortic aneurysm and dissection (I71)	21	0.7	9.4
15	Pneumonitis due to solids and liquids (J69)	20	0.6	9.0
...	All other causes (residual)	398	12.7	178.7
Asian or Pacific Islander, male, 75-79 years				
...	All causes	3,632	100.0	2,362.4
1	Malignant neoplasms (C00-C97)	1,097	30.2	713.5
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	850	23.4	552.9
3	Cerebrovascular diseases (I60-I69)	253	7.0	164.6
4	Diabetes mellitus (E10-E14)	158	4.4	102.8
5	Chronic lower respiratory diseases (J40-J47)	136	3.7	88.5
6	Influenza and pneumonia (J09-J18)	119	3.3	77.4
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	101	2.8	65.7
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	100	2.8	65.0
9	Parkinson's disease (G20-G21)	75	2.1	48.8
10	Septicemia (A40-A41)	56	1.5	36.4
11	Alzheimer's disease (G30)	53	1.5	34.5
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	48	1.3	31.2
13	Chronic liver disease and cirrhosis (K70,K73-K74)	33	0.9	21.5
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	32	0.9	20.8
15	Aortic aneurysm and dissection (I71)	28	0.8	18.2
...	All other causes (residual)	493	13.6	320.7

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2014

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2014

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, male, 80-84 years				
...	All causes	3,931	100.0	4,164.2
1	Malignant neoplasms (C00-C97)	987	25.1	1,045.6
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	953	24.2	1,009.5
3	Cerebrovascular diseases (I60-I69)	286	7.3	303.0
4	Chronic lower respiratory diseases (J40-J47)	216	5.5	228.8
5	Diabetes mellitus (E10-E14)	176	4.5	186.4
6	Influenza and pneumonia (J09-J18)	160	4.1	169.5
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	107	2.7	113.3
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	102	2.6	108.1
9	Alzheimer's disease (G30)	88	2.2	93.2
10	Parkinson's disease (G20-G21)	74	1.9	78.4
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	60	1.5	63.6
12	Septicemia (A40-A41)	51	1.3	54.0
13	Pneumonitis due to solids and liquids (J69)	43	1.1	45.6
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	40	1.0	42.4
15	Aortic aneurysm and dissection (I71)	27	0.7	28.6
...	All other causes (residual)	561	14.3	594.3
Asian or Pacific Islander, male, 85-89 years				
...	All causes	3,741	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	967	25.8	---
2	Malignant neoplasms (C00-C97)	728	19.5	---
3	Cerebrovascular diseases (I60-I69)	261	7.0	---
4	Chronic lower respiratory diseases (J40-J47)	224	6.0	---
5	Influenza and pneumonia (J09-J18)	170	4.5	---
6	Diabetes mellitus (E10-E14)	140	3.7	---
7	Alzheimer's disease (G30)	137	3.7	---
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	106	2.8	---
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	101	2.7	---
10	Parkinson's disease (G20-G21)	92	2.5	---
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	69	1.8	---
12	Pneumonitis due to solids and liquids (J69)	61	1.6	---
13	Septicemia (A40-A41)	52	1.4	---
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	32	0.9	---
15	Aortic aneurysm and dissection (I71)	17	0.5	---
...	All other causes (residual)	584	15.6	---

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2014

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2014

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, male, 90-94 years				
...	All causes	2,583	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	697	27.0	---
2	Malignant neoplasms (C00-C97)	341	13.2	---
3	Cerebrovascular diseases (I60-I69)	192	7.4	---
4	Influenza and pneumonia (J09-J18)	176	6.8	---
5	Alzheimer's disease (G30)	147	5.7	---
6	Chronic lower respiratory diseases (J40-J47)	132	5.1	---
7	Diabetes mellitus (E10-E14)	76	2.9	---
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	75	2.9	---
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	67	2.6	---
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	58	2.2	---
11	Pneumonitis due to solids and liquids (J69)	53	2.1	---
12	Parkinson's disease (G20-G21)	42	1.6	---
13	Septicemia (A40-A41)	28	1.1	---
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	17	0.7	---
15	Tuberculosis (A16-A19)	9	0.3	---
15	Cholelithiasis and other disorders of gallbladder (K80-K82)	9	0.3	---
...	All other causes (residual)	464	18.0	---
Asian or Pacific Islander, male, 95-99 years				
...	All causes	873	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	238	27.3	---
2	Malignant neoplasms (C00-C97)	84	9.6	---
3	Cerebrovascular diseases (I60-I69)	65	7.4	---
4	Influenza and pneumonia (J09-J18)	62	7.1	---
5	Chronic lower respiratory diseases (J40-J47)	49	5.6	---
6	Alzheimer's disease (G30)	48	5.5	---
7	Diabetes mellitus (E10-E14)	27	3.1	---
8	Essential hypertension and hypertensive renal disease (I10,I12,I15)	24	2.7	---
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	23	2.6	---
10	Pneumonitis due to solids and liquids (J69)	19	2.2	---
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	18	2.1	---
12	Parkinson's disease (G20-G21)	11	1.3	---
13	Septicemia (A40-A41)	8	0.9	---
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	4	0.5	---
14	Atherosclerosis (I70)	4	0.5	---
14	Chronic liver disease and cirrhosis (K70,K73-K74)	4	0.5	---
...	All other causes (residual)	185	21.2	---

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2014

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2014

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, male, 100 years and over				
...	All causes	173	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	55	31.8	---
2	Influenza and pneumonia (J09-J18)	19	11.0	---
3	Malignant neoplasms (C00-C97)	15	8.7	---
4	Cerebrovascular diseases (I60-I69)	12	6.9	---
5	Chronic lower respiratory diseases (J40-J47)	10	5.8	---
6	Alzheimer's disease (G30)	8	4.6	---
7	Essential hypertension and hypertensive renal disease (I10,I12,I15)	5	2.9	---
8	Pneumonitis due to solids and liquids (J69)	4	2.3	---
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	4	2.3	---
10	Diabetes mellitus (E10-E14)	3	1.7	---
10	Parkinson's disease (G20-G21)	3	1.7	---
10	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3	1.7	---
13	Septicemia (A40-A41)	1	0.6	---
13	Anemias (D50-D64)	1	0.6	---
13	Nutritional deficiencies (E40-E64)	1	0.6	---
13	Peptic ulcer (K25-K28)	1	0.6	---
13	Diseases of appendix (K35-K38)	1	0.6	---
13	Hernia (K40-K46)	1	0.6	---
13	Hyperplasia of prostate (N40)	1	0.6	---
13	Complications of medical and surgical care (Y40-Y84,Y88)	1	0.6	---
13	Enterocolitis due to <i>Clostridium difficile</i> (A04.7)	1	0.6	---
...	All other causes (residual)	23	13.3	---
Asian or Pacific Islander, female, all ages				
...	All causes	29,884	100.0	295.5
1	Malignant neoplasms (C00-C97)	8,153	27.3	80.6
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	5,991	20.0	59.2
3	Cerebrovascular diseases (I60-I69)	2,408	8.1	23.8
4	Alzheimer's disease (G30)	1,172	3.9	11.6
5	Diabetes mellitus (E10-E14)	1,132	3.8	11.2
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	981	3.3	9.7
7	Influenza and pneumonia (J09-J18)	899	3.0	8.9
8	Chronic lower respiratory diseases (J40-J47)	766	2.6	7.6
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	594	2.0	5.9
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	572	1.9	5.7
11	Septicemia (A40-A41)	359	1.2	3.5
12	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	355	1.2	3.5
13	Parkinson's disease (G20-G21)	249	0.8	2.5
14	Certain conditions originating in the perinatal period (P00-P96)	227	0.8	2.2
15	Chronic liver disease and cirrhosis (K70,K73-K74)	226	0.8	2.2
...	All other causes (residual)	5,800	19.4	57.3

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2014

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2014

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, female, 1-4 years				
...	All causes	61	100.0	12.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	10	16.4	*
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	7	11.5	*
3	Influenza and pneumonia (J09-J18)	6	9.8	*
4	Malignant neoplasms (C00-C97)	5	8.2	*
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	3	4.9	*
6	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2	3.3	*
6	Cerebrovascular diseases (I60-I69)	2	3.3	*
8	Septicemia (A40-A41)	1	1.6	*
8	Diseases of heart (I00-I09,I11,I13,I20-I51)	1	1.6	*
8	Hernia (K40-K46)	1	1.6	*
...	All other causes (residual)	23	37.7	4.7
Asian or Pacific Islander, female, 5-9 years				
...	All causes	42	100.0	6.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	10	23.8	*
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	6	14.3	*
3	Malignant neoplasms (C00-C97)	5	11.9	*
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	2	4.8	*
5	Septicemia (A40-A41)	1	2.4	*
5	Anemias (D50-D64)	1	2.4	*
5	Nutritional deficiencies (E40-E64)	1	2.4	*
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	1	2.4	*
5	Influenza and pneumonia (J09-J18)	1	2.4	*
5	Chronic lower respiratory diseases (J40-J47)	1	2.4	*
5	Diseases of appendix (K35-K38)	1	2.4	*
5	Enterocolitis due to <i>Clostridium difficile</i> (A04.7)	1	2.4	*
...	All other causes (residual)	11	26.2	*
Asian or Pacific Islander, female, 10-14 years				
...	All causes	38	100.0	6.2
1	Malignant neoplasms (C00-C97)	8	21.1	*
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	5	13.2	*
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	4	10.5	*
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	4	10.5	*
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	3	7.9	*
6	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1	2.6	*
6	Anemias (D50-D64)	1	2.6	*
6	Chronic lower respiratory diseases (J40-J47)	1	2.6	*
6	Certain conditions originating in the perinatal period (P00-P96)	1	2.6	*
...	All other causes (residual)	10	26.3	*

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2014

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2014

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, female, 15-19 years				
...	All causes	110	100.0	18.3
1	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	28	25.5	4.7
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	26	23.6	4.3
3	Malignant neoplasms (C00-C97)	15	13.6	*
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	5	4.5	*
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	4	3.6	*
5	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	4	3.6	*
7	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	3	2.7	*
8	Cerebrovascular diseases (I60-I69)	2	1.8	*
9	Anemias (D50-D64)	1	0.9	*
9	Influenza and pneumonia (J09-J18)	1	0.9	*
...	All other causes (residual)	21	19.1	3.5
Asian or Pacific Islander, female, 20-24 years				
...	All causes	125	100.0	17.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	38	30.4	5.3
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	26	20.8	3.6
3	Malignant neoplasms (C00-C97)	12	9.6	*
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	10	8.0	*
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	5	4.0	*
6	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	3	2.4	*
6	Pregnancy, childbirth and the puerperium (O00-O99)	3	2.4	*
8	Meningitis (G00,G03)	2	1.6	*
8	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	2	1.6	*
10	Septicemia (A40-A41)	1	0.8	*
10	Human immunodeficiency virus (HIV) disease (B20-B24)	1	0.8	*
10	Cerebrovascular diseases (I60-I69)	1	0.8	*
10	Influenza and pneumonia (J09-J18)	1	0.8	*
10	Chronic lower respiratory diseases (J40-J47)	1	0.8	*
...	All other causes (residual)	19	15.2	*

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2014

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2014

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, female, 25-29 years				
...	All causes	185	100.0	22.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	46	24.9	5.5
2	Malignant neoplasms (C00-C97)	36	19.5	4.3
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	30	16.2	3.6
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	10	5.4	*
5	Cerebrovascular diseases (I60-I69)	7	3.8	*
6	Pregnancy, childbirth and the puerperium (O00-O99)	6	3.2	*
7	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	5	2.7	*
8	Influenza and pneumonia (J09-J18)	3	1.6	*
8	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	3	1.6	*
10	Septicemia (A40-A41)	2	1.1	*
10	Chronic lower respiratory diseases (J40-J47)	2	1.1	*
10	Chronic liver disease and cirrhosis (K70,K73-K74)	2	1.1	*
13	Tuberculosis (A16-A19)	1	0.5	*
13	Anemias (D50-D64)	1	0.5	*
13	Diabetes mellitus (E10-E14)	1	0.5	*
13	Meningitis (G00,G03)	1	0.5	*
13	Complications of medical and surgical care (Y40-Y84,Y88)	1	0.5	*
...	All other causes (residual)	28	15.1	3.4
Asian or Pacific Islander, female, 30-34 years				
...	All causes	237	100.0	27.3
1	Malignant neoplasms (C00-C97)	70	29.5	8.1
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	38	16.0	4.4
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	34	14.3	3.9
4	Pregnancy, childbirth and the puerperium (O00-O99)	11	4.6	*
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	10	4.2	*
6	Influenza and pneumonia (J09-J18)	6	2.5	*
7	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	5	2.1	*
8	Septicemia (A40-A41)	4	1.7	*
8	Cerebrovascular diseases (I60-I69)	4	1.7	*
8	Chronic liver disease and cirrhosis (K70,K73-K74)	4	1.7	*
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	4	1.7	*
12	Diabetes mellitus (E10-E14)	3	1.3	*
12	Chronic lower respiratory diseases (J40-J47)	3	1.3	*
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2	0.8	*
15	Nutritional deficiencies (E40-E64)	1	0.4	*
15	Aortic aneurysm and dissection (I71)	1	0.4	*
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	1	0.4	*
...	All other causes (residual)	36	15.2	4.1

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2014

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2014

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, female, 35-39 years				
...	All causes	334	100.0	40.8
1	Malignant neoplasms (C00-C97)	139	41.6	17.0
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	33	9.9	4.0
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	32	9.6	3.9
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	22	6.6	2.7
5	Pregnancy, childbirth and the puerperium (O00-O99)	12	3.6	*
6	Diabetes mellitus (E10-E14)	10	3.0	*
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	10	3.0	*
8	Cerebrovascular diseases (I60-I69)	7	2.1	*
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	5	1.5	*
10	Chronic liver disease and cirrhosis (K70,K73-K74)	4	1.2	*
11	Septicemia (A40-A41)	3	0.9	*
11	Viral hepatitis (B15-B19)	3	0.9	*
11	Influenza and pneumonia (J09-J18)	3	0.9	*
11	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	3	0.9	*
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2	0.6	*
15	Anemias (D50-D64)	2	0.6	*
15	Chronic lower respiratory diseases (J40-J47)	2	0.6	*
...	All other causes (residual)	42	12.6	5.1
Asian or Pacific Islander, female, 40-44 years				
...	All causes	469	100.0	57.5
1	Malignant neoplasms (C00-C97)	215	45.8	26.4
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	42	9.0	5.1
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	36	7.7	4.4
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	36	7.7	4.4
5	Cerebrovascular diseases (I60-I69)	26	5.5	3.2
6	Pregnancy, childbirth and the puerperium (O00-O99)	10	2.1	*
7	Diabetes mellitus (E10-E14)	8	1.7	*
7	Influenza and pneumonia (J09-J18)	8	1.7	*
7	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	8	1.7	*
10	Chronic lower respiratory diseases (J40-J47)	7	1.5	*
10	Chronic liver disease and cirrhosis (K70,K73-K74)	7	1.5	*
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	5	1.1	*
13	Septicemia (A40-A41)	4	0.9	*
13	Human immunodeficiency virus (HIV) disease (B20-B24)	4	0.9	*
15	Viral hepatitis (B15-B19)	3	0.6	*
...	All other causes (residual)	50	10.7	6.1

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2014

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2014

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, female, 45-49 years				
...	All causes	650	100.0	92.2
1	Malignant neoplasms (C00-C97)	339	52.2	48.1
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	58	8.9	8.2
3	Cerebrovascular diseases (I60-I69)	50	7.7	7.1
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	28	4.3	4.0
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	28	4.3	4.0
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	15	2.3	*
7	Diabetes mellitus (E10-E14)	13	2.0	*
8	Septicemia (A40-A41)	11	1.7	*
9	Chronic liver disease and cirrhosis (K70,K73-K74)	10	1.5	*
10	Influenza and pneumonia (J09-J18)	7	1.1	*
11	Viral hepatitis (B15-B19)	6	0.9	*
11	Chronic lower respiratory diseases (J40-J47)	6	0.9	*
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	5	0.8	*
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	5	0.8	*
15	Aortic aneurysm and dissection (I71)	4	0.6	*
...	All other causes (residual)	65	10.0	9.2
Asian or Pacific Islander, female, 50-54 years				
...	All causes	1,021	100.0	154.5
1	Malignant neoplasms (C00-C97)	537	52.6	81.2
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	104	10.2	15.7
3	Cerebrovascular diseases (I60-I69)	67	6.6	10.1
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	52	5.1	7.9
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	40	3.9	6.1
6	Diabetes mellitus (E10-E14)	32	3.1	4.8
7	Influenza and pneumonia (J09-J18)	15	1.5	*
8	Essential hypertension and hypertensive renal disease (I10,I12,I15)	10	1.0	*
9	Chronic liver disease and cirrhosis (K70,K73-K74)	9	0.9	*
9	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	9	0.9	*
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	8	0.8	*
12	Septicemia (A40-A41)	5	0.5	*
12	Viral hepatitis (B15-B19)	5	0.5	*
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	5	0.5	*
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	4	0.4	*
...	All other causes (residual)	119	11.7	18.0

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2014

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2014

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, female, 55-59 years				
...	All causes	1,390	100.0	232.4
1	Malignant neoplasms (C00-C97)	725	52.2	121.2
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	146	10.5	24.4
3	Cerebrovascular diseases (I60-I69)	94	6.8	15.7
4	Diabetes mellitus (E10-E14)	55	4.0	9.2
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	54	3.9	9.0
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	24	1.7	4.0
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	20	1.4	3.3
8	Septicemia (A40-A41)	19	1.4	*
8	Chronic lower respiratory diseases (J40-J47)	19	1.4	*
10	Influenza and pneumonia (J09-J18)	18	1.3	*
11	Chronic liver disease and cirrhosis (K70,K73-K74)	14	1.0	*
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	13	0.9	*
13	Viral hepatitis (B15-B19)	12	0.9	*
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	10	0.7	*
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	10	0.7	*
...	All other causes (residual)	157	11.3	26.2
Asian or Pacific Islander, female, 60-64 years				
...	All causes	1,756	100.0	342.4
1	Malignant neoplasms (C00-C97)	873	49.7	170.2
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	242	13.8	47.2
3	Cerebrovascular diseases (I60-I69)	110	6.3	21.4
4	Diabetes mellitus (E10-E14)	73	4.2	14.2
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	48	2.7	9.4
6	Influenza and pneumonia (J09-J18)	34	1.9	6.6
7	Chronic lower respiratory diseases (J40-J47)	24	1.4	4.7
8	Septicemia (A40-A41)	23	1.3	4.5
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	23	1.3	4.5
10	Chronic liver disease and cirrhosis (K70,K73-K74)	21	1.2	4.1
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	20	1.1	3.9
12	Viral hepatitis (B15-B19)	19	1.1	*
13	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	13	0.7	*
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	11	0.6	*
15	Aortic aneurysm and dissection (I71)	10	0.6	*
...	All other causes (residual)	212	12.1	41.3

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2014

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2014

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, female, 65-69 years				
...	All causes	2,192	100.0	551.8
1	Malignant neoplasms (C00-C97)	923	42.1	232.3
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	391	17.8	98.4
3	Cerebrovascular diseases (I60-I69)	143	6.5	36.0
4	Diabetes mellitus (E10-E14)	114	5.2	28.7
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	49	2.2	12.3
6	Influenza and pneumonia (J09-J18)	47	2.1	11.8
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	47	2.1	11.8
8	Chronic lower respiratory diseases (J40-J47)	40	1.8	10.1
9	Septicemia (A40-A41)	35	1.6	8.8
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	26	1.2	6.5
11	Viral hepatitis (B15-B19)	24	1.1	6.0
11	Chronic liver disease and cirrhosis (K70,K73-K74)	24	1.1	6.0
13	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	17	0.8	*
14	Aortic aneurysm and dissection (I71)	15	0.7	*
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	14	0.6	*
...	All other causes (residual)	283	12.9	71.2
Asian or Pacific Islander, female, 70-74 years				
...	All causes	2,499	100.0	917.3
1	Malignant neoplasms (C00-C97)	944	37.8	346.5
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	445	17.8	163.4
3	Cerebrovascular diseases (I60-I69)	207	8.3	76.0
4	Diabetes mellitus (E10-E14)	113	4.5	41.5
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	74	3.0	27.2
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	62	2.5	22.8
7	Chronic lower respiratory diseases (J40-J47)	56	2.2	20.6
8	Influenza and pneumonia (J09-J18)	54	2.2	19.8
9	Essential hypertension and hypertensive renal disease (I10,I12,I15)	47	1.9	17.3
10	Chronic liver disease and cirrhosis (K70,K73-K74)	31	1.2	11.4
11	Septicemia (A40-A41)	28	1.1	10.3
12	Parkinson's disease (G20-G21)	24	1.0	8.8
13	Alzheimer's disease (G30)	23	0.9	8.4
14	Pneumonitis due to solids and liquids (J69)	16	0.6	*
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	13	0.5	*
15	Aortic aneurysm and dissection (I71)	13	0.5	*
15	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	13	0.5	*
...	All other causes (residual)	336	13.4	123.3

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2014

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2014

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, female, 75-79 years				
...	All causes	3,169	100.0	1,631.7
1	Malignant neoplasms (C00-C97)	1,004	31.7	517.0
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	645	20.4	332.1
3	Cerebrovascular diseases (I60-I69)	262	8.3	134.9
4	Diabetes mellitus (E10-E14)	153	4.8	78.8
5	Chronic lower respiratory diseases (J40-J47)	102	3.2	52.5
6	Influenza and pneumonia (J09-J18)	85	2.7	43.8
7	Alzheimer's disease (G30)	76	2.4	39.1
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	72	2.3	37.1
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	68	2.1	35.0
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	60	1.9	30.9
11	Parkinson's disease (G20-G21)	40	1.3	20.6
12	Septicemia (A40-A41)	34	1.1	17.5
12	Chronic liver disease and cirrhosis (K70,K73-K74)	34	1.1	17.5
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	23	0.7	11.8
15	Pneumonitis due to solids and liquids (J69)	22	0.7	11.3
...	All other causes (residual)	489	15.4	251.8
Asian or Pacific Islander, female, 80-84 years				
...	All causes	4,247	100.0	3,094.6
1	Malignant neoplasms (C00-C97)	975	23.0	710.4
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	964	22.7	702.4
3	Cerebrovascular diseases (I60-I69)	394	9.3	287.1
4	Diabetes mellitus (E10-E14)	203	4.8	147.9
5	Alzheimer's disease (G30)	185	4.4	134.8
6	Chronic lower respiratory diseases (J40-J47)	162	3.8	118.0
7	Influenza and pneumonia (J09-J18)	157	3.7	114.4
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	98	2.3	71.4
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	95	2.2	69.2
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	74	1.7	53.9
11	Septicemia (A40-A41)	62	1.5	45.2
12	Parkinson's disease (G20-G21)	55	1.3	40.1
13	Chronic liver disease and cirrhosis (K70,K73-K74)	41	1.0	29.9
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	32	0.8	23.3
15	Pneumonitis due to solids and liquids (J69)	29	0.7	21.1
...	All other causes (residual)	721	17.0	525.4

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2014

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2014

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, female, 85-89 years				
...	All causes	4,794	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,209	25.2	---
2	Malignant neoplasms (C00-C97)	783	16.3	---
3	Cerebrovascular diseases (I60-I69)	505	10.5	---
4	Alzheimer's disease (G30)	308	6.4	---
5	Diabetes mellitus (E10-E14)	185	3.9	---
6	Influenza and pneumonia (J09-J18)	158	3.3	---
7	Chronic lower respiratory diseases (J40-J47)	147	3.1	---
8	Essential hypertension and hypertensive renal disease (I10,I12,I15)	126	2.6	---
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	123	2.6	---
10	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	115	2.4	---
11	Septicemia (A40-A41)	67	1.4	---
12	Parkinson's disease (G20-G21)	57	1.2	---
13	Pneumonitis due to solids and liquids (J69)	42	0.9	---
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	33	0.7	---
15	Aortic aneurysm and dissection (I71)	32	0.7	---
...	All other causes (residual)	904	18.9	---
Asian or Pacific Islander, female, 90-94 years				
...	All causes	3,938	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,048	26.6	---
2	Malignant neoplasms (C00-C97)	410	10.4	---
3	Cerebrovascular diseases (I60-I69)	349	8.9	---
4	Alzheimer's disease (G30)	341	8.7	---
5	Influenza and pneumonia (J09-J18)	183	4.6	---
6	Chronic lower respiratory diseases (J40-J47)	125	3.2	---
7	Diabetes mellitus (E10-E14)	112	2.8	---
8	Essential hypertension and hypertensive renal disease (I10,I12,I15)	104	2.6	---
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	98	2.5	---
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	84	2.1	---
11	Parkinson's disease (G20-G21)	49	1.2	---
12	Pneumonitis due to solids and liquids (J69)	48	1.2	---
13	Septicemia (A40-A41)	33	0.8	---
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	26	0.7	---
15	Aortic aneurysm and dissection (I71)	23	0.6	---
...	All other causes (residual)	905	23.0	---

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2014

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2014

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, female, 95-99 years				
...	All causes	1,759	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	499	28.4	---
2	Alzheimer's disease (G30)	182	10.3	---
3	Cerebrovascular diseases (I60-I69)	146	8.3	---
4	Malignant neoplasms (C00-C97)	113	6.4	---
5	Influenza and pneumonia (J09-J18)	87	4.9	---
6	Essential hypertension and hypertensive renal disease (I10,I12,I15)	66	3.8	---
7	Chronic lower respiratory diseases (J40-J47)	53	3.0	---
8	Diabetes mellitus (E10-E14)	46	2.6	---
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	36	2.0	---
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	32	1.8	---
11	Pneumonitis due to solids and liquids (J69)	17	1.0	---
12	Septicemia (A40-A41)	16	0.9	---
13	Atherosclerosis (I70)	9	0.5	---
14	Cholelithiasis and other disorders of gallbladder (K80-K82)	7	0.4	---
14	Enterocolitis due to <i>Clostridium difficile</i> (A04.7)	7	0.4	---
...	All other causes (residual)	443	25.2	---
Asian or Pacific Islander, female, 100 years and over				
...	All causes	458	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	134	29.3	---
2	Alzheimer's disease (G30)	41	9.0	---
3	Cerebrovascular diseases (I60-I69)	30	6.6	---
4	Malignant neoplasms (C00-C97)	22	4.8	---
4	Influenza and pneumonia (J09-J18)	22	4.8	---
6	Essential hypertension and hypertensive renal disease (I10,I12,I15)	19	4.1	---
7	Chronic lower respiratory diseases (J40-J47)	13	2.8	---
8	Diabetes mellitus (E10-E14)	11	2.4	---
9	Pneumonitis due to solids and liquids (J69)	9	2.0	---
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	9	2.0	---
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	8	1.7	---
12	Septicemia (A40-A41)	5	1.1	---
13	Atherosclerosis (I70)	4	0.9	---
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	3	0.7	---
14	Nutritional deficiencies (E40-E64)	3	0.7	---
...	All other causes (residual)	125	27.3	---

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2014